# Higgs signal in Chargino-Neutralino production at the LHC

## Diptimoy Ghosh<sup>a</sup>, Monoranjan Guchait<sup>b</sup>, Dipan Sengupta<sup>b</sup>

 $^{a}$ Department of Theoretical Physics, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai, India

E-mail: diptimoyghosh@theory.tifr.res.in

<sup>b</sup> Department of High Energy Physics, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai, India

E-mail: guchait@tifr.res.in, dipan@tifr.res.in

Abstract. We have analyzed the prospect of detecting a Higgs signal in mSUGRA/cMSSM based Supersymmetric (SUSY) model via chargino-neutralino( $\tilde{\chi}_1^{\pm}\tilde{\chi}_2^0$ ) production at 8 TeV and 14 TeV LHC energies. The signal is studied in the  $\ell + b\bar{b} + p_T$  channel following the decays,  $\tilde{\chi}_1^{\pm} \rightarrow W^{\pm}\tilde{\chi}_1^0$ ,  $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 h$  and  $h \rightarrow b\bar{b}$ . In this analysis reconstruction of the Higgs mass out of two b-jets plays a very crucial role in determining the signal to background ratio. We follow two techniques to reconstruct the Higgs mass: (A) adding momenta of two identified b-jets, (B) jet substructure technique. In addition, imposing a certain set of selection cuts we observe that the significance is better for the method (B). We find that a signal can be observed for the Higgs mass ~ 125 GeV with an integrated luminosity 100 fb<sup>-1</sup> for both 8 TeV and 14 TeV LHC energies.

#### 1. Introduction

The quest for the Higgs boson is one of the high priority programme of the LHC experiment. Recently, both ATLAS and CMS collaborations have published preliminary results on Higgs searches accumulating data of integrated luminosity about  $5\text{fb}^{-1}$  each at 8 TeV LHC energy. They have constrained the light Higgs mass within the range of 122 - 131 GeV at 95% C.L. which is [1] consistent with the prediction based on electro-weak precision measurements [2]. However, interestingly both the groups have also reported the discovery of a standard model(SM) like Higgs boson with mass ~ 125 GeV [1]. Further investigations are required to confirm that it is indeed the Higgs boson predicted by the SM.

In this paper we investigate the implications of Higgs searches mentioned above assuming that the observed boson is indeed the Higgs boson. As we know, many models beyond the SM also predict the existence of a Higgs particle. For instance, the Minimal Supersymmetric Standard Model(MSSM) contains five Higgs bosons, two CP even Higgs h and H, one CP odd Higgs A and two charged Higgs  $H^{\pm}$ . At the tree level, the masses of all the Higgs particles in the MSSM can be predicted in terms of the two parameters, the CP-odd Higgs mass  $m_A$  and the ratio  $\tan \beta$  of the vacuum expectation values of the two Higgs doublets. The mass of the lightest  $\text{Higgs}(m_h)$  is bounded by  $m_h \leq m_Z$  [3] at the tree level, but loop corrections enhance this limit to  $m_h \leq 140 \text{ GeV}[3]$ . Notice that this theoretical upper limit is consistent with the present limits set on the Higgs mass by LHC experiments. Note that, in the decoupling regime  $m_A \gg m_Z$ , the lightest Higgs becomes SM like. Evidently, there is a correlation between  $m_h$  and other sparticle masses(and hence other model parameters) because of loop effects. The dependence of  $m_h$  on model parameters including recent Higgs mass constraints are discussed by a number of authors in the framework of constrained MSSM(cMSSM) or the minimal supergravity(mSUGRA) [4, 5, 6] model and also other variations of SUSY models [5].

The mSUGRA model is described by four parameters,  $m_0, m_{1/2}, A_0$  (defined at the GUT scale),  $\tan\beta$  (defined at the Electroweak scale) and a sign, the sign of  $\mu$ . Here  $m_0$  is the universal soft mass of the scalars,  $m_{1/2}$  is the unified gaugino mass,  $A_0$  is the universal trilinear coupling and  $\mu$  is the supersymmetric Higgs(ino) mass parameter. The lightest Higgs mass is highly sensitive to  $m_0, A_0$  and  $\tan \beta$ , as the square of the third generation squark mass matrix which contributes dominantly to the loop correction is controlled by these parameters. A detailed scan of parameter space shows that the current constraints on the Higgs mass from LHC experiments is compatible with certain regions of parameter space in mSUGRA/cMSSM. For example, for low  $m_0 (\leq 4 \text{ TeV})$ case, to achieve  $m_h \sim 125$  GeV, a high value of  $A_0$  is required, whereas for high  $m_0 \sim$ 4 TeV, one needs a moderate value of  $A_0[5, 6]$ . As a consequence, the parameter space in mSUGRA allowed by Higgs mass constraints predict the masses of sfermions (squarks and sleptons) to be of multi-TeV range. However, the mass of top squark $(\tilde{t}_1)$  remains comparatively lighter because of mixing effects and is likely to be accessible within the LHC energy range along with gauginos (charginos and neutralinos) and gluinos [6]. It is worth mentioning here that from the negative results in direct searches at LHC, both ATLAS and CMS collaborations have excluded a region in the  $m_0 - m_{1/2}$  plane imposing a limit,  $m_{\tilde{g}} \gtrsim 1.2$  TeV for  $m_{\tilde{q}} \sim m_{\tilde{g}}$  case, and  $m_{\tilde{g}} \gtrsim 800$  GeV for  $m_{\tilde{q}} >> m_{\tilde{g}}$  scenario[7].

In this paper, we explore the detectability of Higgs signal in SUSY cascade decay chain which may enable us to confirm the existence of a SUSY Higgs. With this motivation we investigate the Higgs signal in chargino( $\tilde{\chi}_1^{\pm}$ ) and second lightest neutralino( $\tilde{\chi}_2^0$ ) pair production following the dominant decays,  $\tilde{\chi}_1^{\pm} \rightarrow \tilde{\chi}_1^0 W^{\pm}$  and  $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 h$ . Higgs signal in SUSY cascade decays has been studied previously in detail by the authors of Ref.[8]. It is well known that in hadron colliders, strongly interacting colored sparticles,  $\tilde{g}$  and  $\tilde{q}$  are produced copiously. The current exclusions by the LHC experiments from SUSY and Higgs searches in mSUGRA favor high  $\tilde{q}$  and  $\tilde{g}$  masses( $m_{\tilde{g}}, m_{\tilde{q}} \sim 1 \text{ TeV}$ ). For these ranges of  $\tilde{g}$  and  $\tilde{q}$  masses, the  $\tilde{g}$  pair production is expected to dominate over the  $\tilde{q}$  production. Eventually, the Higgs boson may arise in  $\tilde{g}$  cascade decay chains involving heavy flavors, i.e.,  $\tilde{g} \rightarrow tb \tilde{\chi}_1^{\pm}, t\bar{t} \tilde{\chi}_1^0, t\bar{t} \tilde{\chi}_2^0$  and  $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 h$ . We checked that the probability of finding Higgs events via  $\tilde{g}$  pair production and its subsequent cascade decays is  $\sim 1-3\%$ . Moreover, with the increase of  $\tilde{g}$  and  $\tilde{q}$  masses,

Higgs signal in Chargino-Neutralino production at the LHC

	$m_{1/2}$	$\mu$	$m_h$	$m_{ ilde{g}}$	$m_{\tilde{q}}$	$m_{\tilde{t}_1}$	$m_{ ilde{\chi}_1^0}$	$m_{ ilde{\chi}^0_2}$	$m_{\tilde{\chi}_1^{\pm}}$
P1	300	1541	122.4	865	3000	1305	133	265	265
P2	380	1660	122.8	1046	3060	1335	168	332	332
$\mathbf{P3}$	450	1653	123.2	1200	3096	1370	198	390	390

**Table 1.** Masses of some of the sparticles for three benchmark points. In all the cases  $m_0 = 3000$ ,  $\tan \beta = 30$  and  $A_0 = -4500$ . All mass units are in GeV.

strong production cross sections drop significantly (~ few fb) and electro-weak gaugino pair production takes over. In view of this fact, we consider  $\tilde{\chi}_1^{\pm} \tilde{\chi}_2^0$  pair production to study the Higgs signal instead of the  $\tilde{g}$  pair production. The detection of Higgs signal in this channel has not been studied before for 8 TeV LHC energy. It is to be noted that the  $\tilde{\chi}_1^{\pm} \tilde{\chi}_2^0$  production is regarded to be a promising SUSY discovery channel through the clean trilepton signal. Recently, this channel has also received a lot of attention to probe SUSY signal [9, 10] at the LHC due to the higher limits on  $\tilde{g}$  and  $\tilde{q}$  masses [7]. Similar analysis has also been performed for LHC in [11] for 14 TeV energy. The Higgs production via  $\tilde{t}_1$  production and its subsequent decays has also been discussed in [12].

In mSUGRA, at the GUT scale masses of all the gauginos are given by  $m_{1/2}$  and at the electro-weak scale they are related as  $M_2 \simeq M_3/3 \sim m_{\tilde{g}}/3$  and  $M_1 \simeq M_2/2$  because of renormalization group evolution(RGE). Here  $M_1, M_2$  and  $M_3$  are the U(1), SU(2) and SU(3) gaugino mass parameters respectively. To get a reasonable branching ratio for the decay  $\tilde{\chi}_2^0 \to \tilde{\chi}_1^0 h$  we select parameter space where  $|\mu|$ , the Higgsino mass parameter is very large leading to  $\tilde{\chi}_1^{\pm}, \tilde{\chi}_2^0$  and  $\tilde{\chi}_1^0$  states gaugino dominated. Therefore, for very high values of  $|\mu|$  (i.e.,  $|\mu| >> M_2, M_1$ ),  $m_{\tilde{\chi}_1^{\pm}}, m_{\tilde{\chi}_2^0} \sim M_2 \sim M_{\tilde{g}}/3$ . Hence, in view of the current limit on  $\tilde{g}$  mass,  $m_{\tilde{\chi}_1^{\pm}}$  and  $m_{\tilde{\chi}_2^0}$  are expected to be around 250 GeV or more. For the sake of presentation of our results we select three benchmark points(P1-P3) corresponding to progressively higher values of gaugino masses which are presented in Table 1. For this region of parameter space,  $\tilde{\chi}_2^0 \to \tilde{\chi}_1^0 h$  branching ratio(BR) is more than 80% and the  $h \to b\bar{b}$  BR is about 70%.

In the next sections we discuss our simulation strategy for signal and backgrounds and then present our results. Finally we summarize our study in the last section.

#### 2. Signal and Background

We investigate the Higgs signal in SUSY cascade decay,  $pp \to \tilde{\chi}_1^{\pm} \tilde{\chi}_2^0 \to (\tilde{\chi}_1^0 W^{\pm})(\tilde{\chi}_1^0 h)$ , leading to a final state with a hard lepton $(e, \mu)$  from W decay and two b-jets from Higgs decay and a large  $p_T$  due to the presence of  $\tilde{\chi}_1^0$  and  $\nu$ , but without any additional jet. The identical final state may also come from t $\bar{t}$ , Wb $\bar{b}$ , Zb $\bar{b}$ , WZ, Wh, Zh, tb, tbW processes. Recall that the  $\tilde{\chi}_1^{\pm} \tilde{\chi}_2^0$  pair production cross sections(C.S.) for our considered parameter space are about 30-175 fb (LO) for 8 TeV in contrast to background cross sections which vary from few picobarn(pb) to more than 100 pb. Thus a huge suppression of background events is required to achieve a reasonable sensitivity, which is a challenging task. The added advantage is that the invariant mass constructed out of two b jets is expected to show a peak at the Higgs mass, which can be exploited to identify the signal region. Therefore, a good reconstruction of Higgs mass out of two b jets is one of the crucial issue to be studied in this analysis. In this paper we report about the simulation of signal and backgrounds adopting two methods for Higgs reconstruction. In the first method(Method A) we identify two b-jets out of all jets in the events and obtain the Higgs mass by calculating their invariant mass. In the second method(Method B) reconstruction of Higgs mass is performed by using the jet substructures which will be discussed later. In this paper we present our results for both cases, method A and B.

In our simulation, events are generated using PYTHIA[14] for the signal and  $t\bar{t}$ , WZ, Wh, Zh backgrounds whereas ALPGEN[15] interfaced with PYTHIA has been used for the generation of tb, tbW,  $Wb\bar{b}$  and  $Zb\bar{b}$  backgrounds. We adopt MLM matching [16] to avoid double counting while performing parton showering after matrix element calculations in ALPGEN. We use FastJet for jet reconstruction using built-in anti- $k_T$  algorithm with  $\Delta R=0.5$  [17] in method A, whereas Cambridge-Aachen [18] algorithm is used for method B. We use CTEQ6L parton distribution function while calculating cross sections [19]. SuSpect interfaced with SUSYHIT is used to calculate SUSY mass spectrum and corresponding branching ratios [20].

We observe that use of Higgs mass reconstruction alone is not enough to eliminate backgrounds substantially. A certain set of selection cuts described below are necessary to reject backgrounds.

• Lepton : Leptons (e and  $\mu$ ) are selected with  $p_T^{\ell} \ge 20$  GeV and  $|\eta| \le 2.5$ . Isolation of leptons are ensured by estimating the total transverse energy  $p_T^{AC} \le 20\%$  of  $p_T^{\ell}$ , where  $p_T^{AC}$  is the scalar sum of transverse energies of jets close to leptons satisfying  $\Delta R(\ell, j) \le 0.2$ . We veto events if there exists a second lepton with a loose criteria of  $p_T^{\ell} \ge 10$  GeV, primarily to suppress top background.

• Jets: Jets are selected using FastJet [17] with a  $p_T \ge 50$  GeV and  $|\eta| \le 3(|\eta| < 2.5$  for method B).

• b-Jets: b like jets are identified by performing a matching of jets with b quarks assuming a matching cone  $\Delta R(b, j) = 0.5$ . In addition, we require that the matched b jet transverse momentum should have at least 80% of the b quark transverse momentum. A proper method of b-tagging using displaced vertex is beyond the scope of this analysis. Finally, we multiply by a b-tagging efficiency( $\epsilon_b$ ) of 70% [21] for each b-tagging i.e  $\epsilon_b^2 = 0.5$  for two b-tagged jets while estimating total event rates.

•  $p_T$ : Missing transverse momentum is calculated out of all visible stable particles. The  $p_T$  in the signal arises due to the massive  $\tilde{\chi}_1^0$  and  $\nu$  as well, whereas in background events, it solely arises from  $\nu$  in W decay. Nevertheless, the hardness of  $p_T$  in signal is not significantly different than the large  $t\bar{t}$  background making it very difficult to distinguish the signal from the background.

•  $R_T^{b\bar{b}}$ : We define a very robust variable which is extremely efficient in eliminating backgrounds by huge fraction as discussed in a previous analysis [22]. It is defined as  $R_T^{b\bar{b}} = \frac{p_T^{b_1} + p_T^{b_2}}{H_T}$ , where the numerator is the scalar sum of  $p_T$  of the two b-jets and  $H_T$  is

the scalar sum of  $p_T$  of all jets passing our pre-selection criteria. We define this variable keeping in mind that in the signal process no hard jets are expected except two b-jets from Higgs decay. Of course, few jets may arise from initial and final state radiations, but the number of such jets with  $p_T \geq 50$  GeV is expected to be low. Hence  $R_T^{b\bar{b}}$  turns out to be ~ 1 for signal. In backgrounds, particularly in top pair production there are additional hard jets arising due to the hadronic decay of W(since we are giving a veto on the second lepton) resulting in  $R_T^{b\bar{b}} < 1$ . Thus a judicious choice of a upper cut on  $R_T^{b\bar{b}}$  suppresses backgrounds enormously without affecting the signal much.

•  $\phi^{b\bar{b}}$ : The azimuthal angle  $\phi^{b\bar{b}}$  is defined as the angle between two b-jets in the transverse plane. In the signal process the angle is expected to be small, but in backgrounds, for example in top pair production they are in general widely separated. It has to be emphasized here that with the increase in  $\tilde{\chi}_2^0$  mass, the  $h \to b\bar{b}$  system gets more and more boosted and hence the b jets become more and more collinear which is an ideal situation for the jet substructure analysis described in the following section. We find that a reasonable cut on  $\phi^{b\bar{b}}(\phi^{b\bar{b}} \leq 2)$  suppresses the background considerably.

•  $m_T(\ell, \not{p}_T)$ : The transverse mass is defined as  $m_T = \sqrt{2p_T^\ell \not{p}_T (1 - \cos\phi(\ell, \not{p}_T))}$ , where  $\phi(\ell, \not{p}_T)$  is the azimuthal angle between the lepton and  $\not{p}_T$  direction. The value of  $m_T(\ell, \not{p}_T)$  is expected to be restricted by W mass if both leptons and  $\not{p}_T$  originate from W decay, which is the case for backgrounds, particularly for  $t\bar{t}$  and  $Wb\bar{b}$  channels. Therefore, a reasonable cut on  $m_T(\ell, \not{p}_T)$  is found to be extremely effective to reduce the background level.

•  $m_{b\bar{b}}$ : As mentioned above the invariant mass of two b-jets is very useful in isolating the signal region. In method A, this reconstruction is straight forward and is performed using two b-jets momenta obtained by matching b-jets with b-quarks. However, in method B, we use jet substructures to find b-jets inside a "fat-jet" from the Higgs decay. The use of jet substructure for the reconstruction of hadronic decays of boosted W, Z, Higgs boson and top quark has received considerable attention in recent years and the available literature is steadily increasing [23]. In our present study this method was motivated following the work of Ref. [24] where the authors reconstructed the Higgs mass using jet substructures to increase the signal sensitivity. The efficiency of jet substructure technique depends on the boost factor of the decayed object. A highly boosted system ensures that decay products are well collimated and appear as a "fatjet". However, in the scenario of interest to us Higgs is moderately boosted as its  $p_T$ depends on  $\Delta m = m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$ . In our analysis we first cluster all the stable final state particles into a "fat jets" using the C/A algorithm [18] with R = 1.2 as implemented in Fastjet [17]. We select "fat jets" with  $p_T \geq 100$  GeV and  $|\eta| < 2.5$  and then perform jet substructure analysis. There are various methods of finding jet substructures [23]. We use the mass drop(MD) method |24| (coded in the FastJet package |17|) in our analysis optimizing the two input parameters,  $\mu = 0.4$  and  $y_{cut} = 0.1$ . In the simulation we use PYTHIA event generator by setting Tune  $Z2^*$  parameters described in Ref. [25] for underlying event modeling. In Figure.1, we show the reconstructed Higgs mass following method A(blue) and B(red) corresponding to parameters P2. This figure

Higgs signal in Chargino-Neutralino production at the LHC

Process	$\sigma(\mathrm{pb})$	$N_{EV}$	$1\ell$	$R_T^{b\bar{b}}$	$m_{b\bar{b}}$	$p_T$	$\phi_{bb}$	c.s.
			2b-jets	$\geq 0.7$	110 - 130	$\geq \! 175$	$\leq 2$	(fb)
P1	0.175	0.1M	1392	1162	723	92	76	0.065
P2	0.065	$0.1 \mathrm{M}$	1767	1478	933	217	178	0.06
P3	0.03	0.1M	2142	1774	1122	424	391	0.055
Wh	0.58	50K	702	594	394	8	2	0.01
Zh	0.3	50K	210	162	51	1	1	0.003
$W b ar{b}$	3	619685	26841	24513	2269	8	3	0.014
Z b ar b	5.1	378098	3863	2937	269	< 1	< 1	< 1
$t\bar{t}$								
5-100	48.2	4M	207335	94337	10145	< 1	< 1	< 1
100-200	36.3	2M	158450	50967	1205	8	1	0.01
200-500	9.5	1M	134238	22473	116	42	4	0.02

**Table 2.** Event summary for signal and backgrounds(method A) for 8 TeV after each set of cuts described in the text. The  $t\bar{t}$  events are simulated for different  $\hat{p}_T$  bins as shown. Efficiency for tagging two b jets is multiplied in the last column. Note that for entries with < 1 the event yield in our case is 0; however since we have simulated finite number of events we denote them as < 1. The energy units are in GeV.

clearly demonstrates the usefulness of the jet substructure technique for Higgs mass reconstruction. In case of method A, some of the soft jets are incidentally passing the matching criteria resulting in a spread towards the lower side, whereas in the jet substructure method this type of contamination is avoided by the filtering procedure described in [24].

#### 3. Results

#### Method A:

In this section we discuss the simulation strategy of signal and backgrounds by reconstructing the Higgs mass out of two identified b-jets obtained by matching techniques as discussed above. In order to eliminate SM backgrounds additional cuts are applied with the following requirements:

- $R_T^{b\bar{b}} \ge 0.7$ ,
- $m_{b\bar{b}} = 110-130 \text{ GeV},$
- $p_T \ge 175 \text{ GeV},$
- $\phi_{b\bar{b}} \leq 2$ .

In Table 2 we present event summaries of signal for three benchmark points shown in Table 1, along with backgrounds after applying these set of cuts. The second and third column present the raw leading order(LO) cross section and number of events simulated respectively. In the fourth column, we present the number of events requiring one single hard lepton along with two identified b-jets and veto the second lepton as

Higgs signal in Chargino-Neutralino production at the LHC

Process	$\sigma(\mathrm{pb})$	$N_{EV}$	$1\ell$	$R_T^{b\bar{b}}$	$m_{bar{b}}$	$p_T$	$\phi_{bb}$	c.s.
			2b-jets	$\geq 0.7$	110-130	$\geq 175$	$\leq 2$	(fb)
P1	0.502	0.1M	3867	2500	1213	89	73	0.18
P2	0.202	$0.1 \mathrm{M}$	4391	2756	1381	273	229	0.23
P3	0.104	$0.1 \mathrm{M}$	4517	2824	1431	373	323	0.17
Wh	1.26	0.1M	3002	1639	750	21	15	0.09
Zh	0.69	$0.1 {\rm M}$	799	280	85	1	1	0.004
$W b ar{b}$	4.5	362018	57764	47883	44160	3948	9	0.055
Z b ar b	7.2	406110	442	380	322	< 1	< 1	< 1
$t\bar{t}$								
5-100	188	10M	1163903	188856	16910	7	4	0.04
100-200	156	10M	1202319	82970	4367	70	32	0.25
200-500	48.5	$1\mathrm{M}$	133840	2020	252	61	32	0.8

Table 3. Same as Table 2 but for 14 TeV. The same conventions as in Table 2 are used.

well. Although we simulated all possible SM backgrounds including QCD, tb, tbW, but we present results only for non-negligible contributing channels. It clearly demonstrates that the  $R_T^{b\bar{b}}$  cut is very effective in reducing backgrounds by an enormous amount, but except for channels, like  $Wb\bar{b}$  and Wh. Selection of events in the Higgs mass window between 110-130 GeV is also useful to remove backgrounds keeping almost more than 50% of signal events. Finally, a very strong  $p_T$  cut is used to eliminate remaining backgrounds, but at the cost of a sizable signal cross section; nevertheless we retain a good number of signal events. After all cuts, we find the total background cross section is about 0.057 fb with dominant contribution from  $t\bar{t}$ , whereas signal cross sections are in the range 0.065-0.055 fb. In both cases we use LO production cross sections. However, if we use NLO cross sections by multiplying K-factors which is ~ 1.5 for signal [26] and about 1.6 for  $t\bar{t}$  [27], then assuming a luminosity 100 fb<sup>-1</sup>, one can expect S/ $\sqrt{B}$  about 3.5 for these mass ranges of  $\tilde{\chi}_1^{\pm}$  and  $\tilde{\chi}_2^0$ .

In Table 3 , we present results for 14 TeV energy corresponding to the same set of benchmark points along with the SM background. We observe that the signal efficiency remains fairly the same as 8 TeV with the enhancement occurring only due to the increase in cross section. The top background however increases significantly due to a presence of a stronger missing momentum and more reconstruction of Higgs mass from the  $b\bar{b}$  system. The total background cross section at 14 TeV turns out to be 1.23 fb as compared to the signal cross sections which are between 0.18 fb and 0.25 fb. As a consequence it becomes difficult to observe a signal with low luminosity options in this approach. However for an integrated luminosity of  $1000 \text{fb}^{-1}$  it may be possible to observe a signal in this method at a  $5\sigma$  level.



Figure 1. The reconstructed Higgs mass for method A (blue) and method B (red) for  $\sqrt{s} = 8$  TeV. The y-axis is normalized to unity.

Process	C.S.	$N_{EV}$	$m_{b\bar{b}}$	Lepton	$m_T(\ell, p_T)$	$R_T^{b\bar{b}}$	$p_T$	C.S.
	(pb)			$\geq 20$	$\geq 90$	$\geq 0.9$	$\geq 125$	(fb)
P1	0.175	$25\mathrm{K}$	179	55	24	19	12	0.042
P2	0.065	10K	168	42	23	18	12	0.04
P3	0.03	10K	273	75	44	36	31	0.045
Wh	0.58	0.1M	871	239	12	11	< 1	< 1
$\mathrm{Zh}$	0.3	0.2M	1698	37	< 1	< 1	< 1	< 1
$W b \bar{b}$	3	619671	191	111	10	8	< 1	< 1
Z b ar b	5.1	378086	81	13	< 1	< 1	< 1	< 1
$t\bar{t}$								
5-100	48.2	5M	1669	454	38	1	< 1	< 1
100-200	36.3	4M	1583	440	42	3	1	0.005
200-500	9.5	$1\mathrm{M}$	315	98	9	2	< 1	< 1

**Table 4.** Event summary for signal and backgrounds(method B) for 8 TeV after each set of cuts described in the text. The same conventions as in Table 2 are used.

<u>Method B</u>: In this method we apply jet substructure technique in reconstructing mass of Higgs within the mass window between 117 - 128 GeV and with additional cuts as before to control background events,

- $m_T(\ell, \not p_T) \ge 90 \text{ GeV } \ddagger,$
- $R_T^{b\bar{b}} \ge 0.9$ ,
- $p_T \ge 125 \text{ GeV}(150 \text{ GeV for } 14 \text{ TeV}).$

‡ Note that we have taken the finite width effects of W boson into account in our simulation. This results in a tail in the  $m_T(\ell, \not p_T)$  distribution in processes like Wh and  $t\bar{t}$ . Thus forced us to opt for a higher value for  $m_T(\ell, \not p_T)$  selection cut.

After the Higgs mass reconstruction the remaining stable particles are used to find jets with C/A algorithm with  $\Delta R = 0.5$ ,  $p_T \geq 50$  GeV,  $|\eta| \leq 2.5$ . The Table 4 displays the robustness of  $R_T^{b\bar{b}}$  cut along with  $m_T(\ell, p_T)$  leading to a suppression of backgrounds to a negligible level without affecting signal significantly.

Process	C.S.	$N_{EV}$	$m_{b\bar{b}}$	Lepton	$m_T(\ell, p_T)$	$R_T^{b\bar{b}}$	$p_T$	C.S.
	(pb)			$\geq 20$	$\geq 90$	$\geq 0.9$	$\geq 150$	(fb)
P1	504	25K	242	55	23	16	5	0.05
P2	204	25K	461	113	55	43	26	0.1
P3	104	25K	713	197	116	67	46	0.095
Wh	1.3	0.1M	946	289	17	11	4	0.026
Zh	704	$0.1 \mathrm{M}$	866	13	1	< 1	< 1	< 1
$W b ar{b}$	5.5	431062	159	92	8	7	< 1	< 1
$Zbar{b}$	7.2	571166	150	< 1	< 1	< 1	< 1	< 1
$t\bar{t}$								
5-100	190	10M	4178	1016	121	12	< 1	< 1
100-200	158	$1\mathrm{M}$	4463	1181	137	11	2	0.01
200-500	49	$0.25 \mathrm{M}$	867	296	25	7	1	0.02

Table 5. Same as Table 4 but for for 14 TeV. The same conventions as in Table 2 are used.

Notice that after cuts signal cross sections remain the same for all cases although production cross sections decrease with the increase of gaugino masses, which is compensated by the increase of acceptance efficiencies. The total background cross section turn out to be 0.007 fb, an order of magnitude less than the method A whereas signal cross sections are of the same level. Assuming 100 fb<sup>-1</sup> luminosity, one can expect signal to background ratio  $S/\sqrt{B} \sim 7$  using NLO cross sections as before. It implies that probing the Higgs signal in this channel is promising with 8 TeV LHC energy and high luminosity options. In both cases signal sensitivity is low because of the tiny production cross section in comparison with the backgrounds.

For 14 TeV energy, as presented in Table 5, we find that the results are not significantly different for method B. Comparing Table 4 and Table 5 we observe a better reconstruction of the Higgs mass because of the enhanced boost of the  $b\bar{b}$  system at 14 TeV energy. However this gain is diluted due to an increase in  $\not{p}_T$  cut compared to 8 TeV to suppress the backgrounds. It has to be noted that at 14 TeV we receive a finite background contribution from Wh process due to an increase in  $\not{p}_T$ . We find that after all cuts the total background cross section is 0.05 fb while the signal cross sections vary between 0.05 fb to 0.1 fb . It is therefore possible to discover a signal for this type of parameter space at the 5  $\sigma$  level at ~ 100  $fb^{-1}$  luminosity.

### 4. Summary

We investigate the discovery potential of a Higgs signal in  $\tilde{\chi}_1^{\pm} \tilde{\chi}_2^0$  production and its subsequent decay channels at 8 TeV and 14 TeV LHC energy. This study is performed in the context of mSUGRA model taking into account the current Higgs mass constraints predicted by recent measurements by CMS and ATLAS experiments. We simulate signal events in the final state with a single hard lepton and  $p_T$  along with a reconstructed Higgs mass. The Higgs mass reconstruction is performed following two ways, first by identifying b-jets using matching technique (method A) and secondly by using the method of jet substructures (method B). We present results for both cases and find that the latter method is more promising than the former one. Incidentally, for low luminosity ( $\mathcal{L} \sim 20 f b^{-1}$ ), which is the projected luminosity for 8 TeV LHC run, the signal cross section is too low to be observed and hence we require high luminosity. For instance, we expect a significance of about ~ 7 for 250-400 GeV masses of  $\tilde{\chi}_1^{\pm}$  and  $\tilde{\chi}_2^0$  and  $m_h \sim 125$  GeV with 100 fb<sup>-1</sup> luminosity by using jet substructure method for 8 TeV. We also performed the analysis for 14 TeV energy and found identical results. The observations made in this paper therefore suggest that the jet substructure method works better for both 8 TeV and 14 TeV LHC energy. It is observed that results do not change significantly for other values of  $\tan \beta$ . In order to increase the sensitivity of Higgs signal one needs to devise more effective selection cuts to isolate tiny signal events out of the huge backgrounds. If Higgs is discovered it is worthwhile to study this channel to identify the model framework. The signal acceptance efficiency is dependent on  $\Delta m$ , which is sensitive to different models. Therefore, our conclusions are model specific and expected to be different in the case of other SUSY models, particularly in models where mass relations among gauginos follow a different pattern. It is also worth investigating the feasibility of detecting supersymmetric Higgs in the  $h \to \tau^+ \tau^-$  channel which will be presented in a future work [28].

#### References

- S. Chatrchyan *et al*, The CMS Collaboration, arXiv:1207.7235[hep-ex]; G. Aad *et al*, The ATLAS Collaboration, arXiv:1207.7214.
- [2] K. Nakamura et al. (Particle Data Group), J. Phys. G 37(2010), 075021.
- [3] A.Djouadi, Phys. Rept. 459, 1(2008), arXiv[hep-ph/0503173] and references therein.
- [4] H. Baer, V.Barger, A. Mustafayev Phys.Rev. D85 075010(2012), arXiv:1112.3017.
- [5] A. Arbey, M. Battaglia, A. Djouadi, F. Mahmoudi, J. Quevillon Phys.Lett, B 708 (2012) 162-169, arXiv:1112.3028.
- [6] S. Akula, B. Altunkaynak, D. Feldman, P. Nath, G. Peim, arXiv:1112.3645.
- [7] The CMS collaboration, CMS-PAS-SUS-:12-005(2012); G. Aad et al. ATLAS Collaboration arXiv:1206.1760 [hep-ex].
- [8] H. Baer, M. Bisset, X. Tata, J. Woodside, Phys. Rev D46(1992) 303; H. Baer, M. Bisset, C. Kao, X. Tata, Phys.Rev. D50(1994)316; A. Datta, A. Djouadi, M. Guchait, Y. Mambrini, Phys.Rev. D65 (2002) 015007; A. Datta, A. Djouadi, M. Guchait, F. Moortgat Nucl.Phys.B681(2004)31; S. Gori, P. Schwaller, C. E. M. Wagner, Phys.Rev. D83(2011), 115022.
- [9] S. Mrenna, arXiv:1110.4078.

- [10] H. Baer, V. Barger, S. Kraml, A. Lessa, W. Sreethawong, X. Tata, arXiv:1201.5382.
- [11] H. Baer, V. Barger, A. Lessa, W. Sreethawong, X. Tata, arXiv:1201.2949; P. Byakti and D. Ghosh, arXiv:1204.0415.
- [12] S. Heinemeyer, F. v. d. Pahlen, H. Rzehak, C. Schappacher, arXiv:1201.6305.
- [13] A. Arbey, F. Mahmoudi, Comput. Phys. Commun. 182 (2011) 1582.
- [14] T. Sjostrand, S. Mrenna, P.Z. Skands, J. High Energy Phys. 05 (2006) 026, [arXiv:hep-ph/0603175].
- [15] M.L. Mangano, M. Moretti, F. Piccinini, R. Pittau, A.D. Polosa, JHEP 0307 (2003) 001.
- [16] S. Hoche, F. Kraus, N. Lavesson, L. Lonnbald, and M. Mangano, arXiv:hep-ph/0602031.
- [17] M. Cacciari, G.P. Salam, and G. Soyez, Phys. Lett. B 641, 57 (2006), [arXiv:hep-ph/0512210].
- [18] Y.L. Dokshitzer, G.D. Leder, S. Moretti and B.R. Webber, JHEP 9708 (1997) 001, [arXiv:hep-ph/9707323].
- [19] H.L. Lai et al. (CTEQ Collaboration), Eur. Phys. J. C 12, 375 (2000).
- [20] A. Djouadi, M.M. Muhlleitner, M. Spira Acta Phys.Polon.B 38:635-644,2007.
- [21] CMS collaboration Report NO.CMS-PAS-BTV-11-001.
- M. Guchait, D. Sengupta Phys.Rev.D84(2011) 055010 [arXiv:1102.4785]; D. Ghosh, M. Guchait,
  S. Raychaudhuri and D. Sengupta, arXiv:1205.2283; R.M. Chatterjee, M. Guchait and
  D. Sengupta, arXiv:1206.5770.
- [23] A Abdesallam et.al. Eur.Phys.J.C71,1661(2011).
- [24] J. M. Butterworth, A. R. Davison, M. Rubin, G. P. Salam Phys.Rev.Lett.100:242001,2008.
- [25] A. Knutsson, talk given in MB & UE meeting, CERN, 17th June, 2011.
- [26] W. Beenakker, R. Hopker, M. Spira, and P. Zerwas, Nucl. Phys. **B492**, 51 (1997).
- [27] N. Kidonakis, arXiv:1109.3231.
- [28] D. Ghosh, M. Guchait, D.Sengupta, in preparation.