# HOLOMORPHIC QUILLEN DETERMINANT LINE BUNDLES ON INTEGRAL COMPACT KÄHLER MANIFOLDS

RUKMINI DEY AND VARGHESE MATHAI

Dedicated to the memory of Daniel G. Quillen

Abstract. We show that any compact Kahler manifold with integral Kahler form, parametrizes a natural holomorphic family of Cauchy-Riemann operators on the Riemann sphere such that the Quillen determinant line bundle of this family is isomorphic to a sufficiently high tensor power of the holomorphic line bundle determined by the integral Kahler form. We also establish a symplectic version of the result. We conjecture that an equivariant version of our result is true.

### 1. INTRODUCTION

In geometric quantization, given a type  $(1, 1)$  integral form  $\xi$  on a compact Kähler manifold, there is a holomorphic line bundle  $\mathcal L$  on the manifold with connection and curvature equal to  $\xi$ . For positive integral (1, 1) forms, we prove the following partial refinement.

<span id="page-0-0"></span>**Theorem 1.** Any compact Kähler manifold M with integral Kähler form  $\omega$ , parametrizes a natural holomorphic family of Cauchy-Riemann operators  $\{\overline{\partial}_z : z \in M\}$  on  $\mathbb{C}P^1$  such that the Quillen determinant line bundle  $\det(\overline{\partial}) \cong \mathcal{L}^{\otimes k}$  as holomorphic line bundles, where  $\mathcal L$  is the holomorphic line bundle determined by  $\omega$ , for some sufficiently large k.

Here we recall that in the seminal paper [\[9\]](#page-9-0), Quillen constructed a holomorphic line bundle  $\mathfrak{L}$ , termed the determinant line bundle, associated to any holomorphic family of Cauchy-Riemann operators over a compact Riemann surface. Quillen also constructed a hermitian metric on  $\mathfrak{L}$ , and calculated its curvature. His construction is referred to in the statement of Theorem [1.](#page-0-0)

The strategy of the proof of Theorem [1](#page-0-0) goes briefly as follows. We first establish the theorem for complex projective spaces  $\mathbb{C}P^N$  for all integers N and for  $k = 1$ . This is achieved by viewing  $\mathbb{C}P^N$  as the moduli space of N-vortices on  $\mathbb{C}P^1$ . The position of each N-vortex z determines a Cauchy-Riemann operator  $\overline{\partial}_z$  on  $\mathbb{C}P^1$ , and the Quillen determinant line bundle, denoted det( $\overline{\partial}$ ), is

<sup>2010</sup> Mathematics Subject Classification. 58J52, 14D21, 53D50, 32Q15, 53D30.

Key words and phrases. holomorphic Quillen determinant line bundles, vortex moduli space, Kodaira embedding theorem, Gromov's embedding theorem.

Acknowledgments. The first author thanks the SYM project grant for support and the second author thanks the Australian Research Council for support and the Harishchandra Research Institute for hospitality during the period when this work was completed. Both authors thank M.S. Narasimhan for pointing out the error in the statement of Kodaira's embedding theorem in the previous version of the paper. We also thank the referees for suggestions leading to improvements in the paper.The first author would like to thank Professor Leon Takhtajan for introducing her to the Quillen determinant bundle during her thesis work. V.M. was fortunate to have been a Ph.D. student of Daniel Quillen at MIT.

isomorphic as a holomorphic line bundle to the hyperplane bundle  $\mathcal{O}(1)$  on  $\mathbb{C}P^N$ , see §[2.](#page-1-0) The next step is to apply the Kodaira embedding theorem to establish the theorem in general, see §[3.](#page-7-0) In §[4,](#page-7-1) we prove a symplectic analogue of our theorem, established using Gromov's embedding theorem. In §[5,](#page-8-0) we conjecture that an equivariant version of our result is true.

Our use of the vortex moduli space seems to be novel: the moduli space is usually a subject of investigations for its own sake, rather than a tool for applications. As an application of our theorem, recall that one can relate the holomorphic sections of the hyperplane bundle  $\mathcal{O}(1)$  on  $\mathbb{C}P^k$ to coherent states in physics, cf. [\[11,](#page-9-1) [13\]](#page-9-2). Our use of the Kodaira embedding theorem shows that the same is true for holomorphic sections of  $\mathcal{L}^{\otimes k}$  for some sufficiently large k, and by our theorem, the same is also true for  $\det(\overline{\partial})$ , which is a topic that we plan to explore.

## <span id="page-1-0"></span>2. Moduli space of N-vortices on a sphere and determinant line bundle

2.1. Vortex equations. The vortex equations are as follows. Let  $\Sigma$  be a compact Riemann surface (in our case the sphere) and let  $\omega = \frac{1}{2}$  $\frac{1}{2}h^2 dz \wedge d\overline{z}$  be the purely imaginary volume form on it, (i.e. h is real). Let A be a unitary connection on a principal  $U(1)$  bundle P i.e. A is a purely imaginary valued one form i.e.  $A = A^{(1,0)} + A^{(0,1)}$  such that  $A^{(1,0)} = -A^{(0,1)}$ . Let L be a complex line bundle associated to  $P$  by the defining representation. We denote by the same  $A$  the unitary connection on L. Let  $\Psi$  be a section of L, i.e.  $\Psi \in \Gamma(\Sigma, L)$  and  $\Psi$  be a section of its dual, L. There is a Hermitian metric H on L, i.e. the inner product  $\langle \Psi_1, \Psi_2 \rangle_{H} = \Psi_1 H \overline{\Psi}_2$  is a smooth function on Σ. (Here *H* is real).

The pair  $(A, \Psi)$  will be said to satisfy the vortex equations if

(1) 
$$
F(A) = \frac{1}{2}(1 - |\Psi|_H^2)\omega,
$$
  
(2)  $\bar{\partial}_A \Psi = 0,$ 

where  $F(A)$  is the curvature of the connection A and  $d_A = \partial_A + \overline{\partial}_A$  is the decomposition of the covariant derivative operator into  $(1,0)$  and  $(0,1)$  pieces. Let S be the space of solutions to  $(1)$ and (2). There is a gauge group G acting on the space of  $(A, \Psi)$  which leaves the equations invariant. We take the group G to be abelian and locally it looks like Maps( $\Sigma$ ,  $U(1)$ ). If g is an U(1) gauge transformation then  $(A_1, \Psi_1)$  and  $(A_2, \Psi_2)$  are gauge equivalent if  $A_2 = g^{-1}dg + A_1$ and  $\Psi_2 = g^{-1}\Psi_1$ . Taking the quotient by the gauge group of S gives the moduli space of solutions to these equations and is denoted by M.

 $F(A) = 4\pi N$  where N is an integer called the vortex number.

Σ There is a theorem by Taubes (cf. [\[14\]](#page-9-3)) and Bradlow [\[2\]](#page-8-1), which says that the moduli space of vortices is parametrised by the zeroes of  $\Psi$ . Thus the moduli space is the symmetric product, Sym<sup>N</sup>( $\Sigma$ ). When the Riemann surface is  $\mathbb{C}P^1$ , the moduli space is complex projective space,  $\text{Sym}^N(\mathbb{C}P^1) = \mathbb{C}P^N.$ 

2.2. The metric and symplectic form. Let  $A$  be the space of all unitary connections on L and  $\Gamma(\Sigma, L)$  be sections of L.

There is a natural norm on A and  $\Gamma(\Sigma, L)$ , as follows:

Z

Let  $A = A_0 + \alpha$  where  $\alpha$  is a unitary one form on the Riemann surface and  $A_0$  is fixed. Then we define  $||A||^2 = \int_{\Sigma} \alpha \wedge \alpha$  and  $||\Psi||^2 = \int_{\Sigma} |\Psi|_H^2 d\mu$ . (The Hodge star acting on an imaginary valued 1 form is spelt out below).

In general, if  $\tau$  is a k- form on a manifold  $\Sigma$ , the norm of  $\tau$  is given by  $\int_{\Sigma} * \tau \wedge \tau$ ,  $*$  being the Hodge star.

With these norms we consider the Sobolev space  $H_3(\mathcal{A})$  to be  $\{A \in L^2(\mathcal{A}) \mid ||A||^2 + ||F(A)||^2 < \infty\}$  $\infty$ , { (since  $d(F(A)) = 0$  on a Riemann surface), where  $F(A)$  is defined, for the time being, in the distributional sense. Let  $H_3(\Gamma(\Sigma, L))$  to be  $\{\Psi \in L^2(\Gamma(\Sigma, L)) \mid ||\Psi||^2 + ||\nabla \Psi||^2 + ||\nabla^2 \Psi||^2 < \infty\}$ where the covariant derivative  $\nabla$  is taken w.r.t a fixed connection, say  $A_0$ , and is defined (for the time being) in the distributional sense.

On a Riemann surface  $(A, \Psi) \in H_3$  implies (by Sobolev lemma) that A and  $\Psi$  are actually  $C^1$  $(5, page 86).$ 

Recall that  $H_3(\Gamma(\Sigma, L))$  and  $H_3(\mathcal{A})$  are Hilbert spaces, a fact that will be useful in Marsden-Weinstein reduction.

Let  $C = H_3(\mathcal{A}) \times H_3(\Gamma(\Sigma, L))$  be the affine space on which equations (1) and (2) are imposed. Then the tangent space at  $p = (A, \Psi)$  is  $T_p C = H_3(\Omega^1(\Sigma, i\mathbb{R})) \times H_3(\Gamma(\Sigma, L))$ 

Once again,  $(\alpha, \beta)$  in  $H_3$  implies they are actually  $C^1$  on a Riemann surface. Let  $p = (A, \Psi) \in \mathcal{C}, X = (\alpha_1, \beta), Y = (\alpha_2, \eta) \in T_p\mathcal{C}$  i.e.  $\alpha_i = \alpha_i^{(0,1)} + \alpha_i^{(1,0)}$  $i^{(1,0)}$  such that  $\alpha_i^{(0,1)} = -\alpha_i^{(1,0)}$  $i^{(1,0)}$ ,  $i = 1, 2$ .

On C one can define a metric

$$
\mathcal{G}(X,Y) = \int_{\Sigma} * \alpha_1 \wedge \alpha_2 + i \int_{\Sigma} \text{Re} \langle \beta, \eta \rangle_H \, \omega
$$

and an almost complex structure  $\mathcal{I} =$ " ∗ 0  $0 \quad i$ 1 :  $T_p \mathcal{C} \to T_p \mathcal{C}$  where  $* : \Omega^1 \to \Omega^1$  is the Hodge star operator on  $\Sigma$  such that  $*\alpha^{1,0} = -i\alpha^{1,0}$  and  $*\alpha^{0,1} = i\alpha^{0,1}$  (i.e. it makes  $A^{0,1}$  the holomorphic coordinate on  $A$ ).

It is easy to check that G is positive definite. In fact, if  $\alpha_1 = \alpha^{(1,0)} + \alpha^{(0,1)} = adz - \bar{a}d\bar{z} = i(A_1)$  $dx + A_2 dy$ ) is an imaginary valued 1-form,  $*\alpha_1 = -i(adz + \bar{a}d\bar{z})$  and

$$
\mathcal{G}(X,X) = \int_{\Sigma} 4|a|^2 dx \wedge dy + \int_{\Sigma} |\beta|_H^2 h^2 dx \wedge dy
$$

where  $\omega = \frac{1}{2}$  $\frac{1}{2}h^2 dz \wedge d\bar{z} = -ih^2 dx \wedge dy.$ 

We define

$$
\Omega(X, Y) = -\frac{1}{2} \int_{\Sigma} \alpha_1 \wedge \alpha_2 + \frac{i}{2} \int_{\Sigma} \text{Re} < i\beta, \eta > H \omega
$$

$$
= -\frac{1}{2} \int_{\Sigma} \alpha_1 \wedge \alpha_2 - \frac{1}{4} \int_{\Sigma} (\beta H \bar{\eta} - \bar{\beta} H \eta) \omega
$$

such that  $\mathcal{G}(\mathcal{I}X, Y) = 2\Omega(X, Y)$ . It is closed, since it is constant. In fact,  $\Omega$  is the real Kähler form on the affine space, [\[17\]](#page-9-4) page 93. It is positive, since it comes from a positive definite metric.

**Lemma 1.**  $\Omega$  is a symplectic form on the vortex moduli space.

*Proof.* Let  $\mathcal{C}' = \{ (A, \Psi) \in \mathcal{C} | \bar{\partial}_A \Psi = 0 \}$  be the submanifold of  $\mathcal C$  defined by the second equation.

a) We first show that  $\mathcal{C}'$  is a symplectic submanifold with the same symplectic form. By linearizing the equation  $\bar{\partial}_A \Psi = 0$  we get  $\Psi \alpha_1^{0,1} = -\bar{\partial}_A \beta$  so that if  $X = (\alpha_1, \beta) \in T_p \mathcal{C}'$  then  $\alpha_1^{0,1} = -\frac{\bar{\partial}_A \beta}{\Psi}$  $\frac{dA}{dP}$ , when  $\Psi \neq 0.$ 

(Note that:

(i)  $\bar{\partial}_A\beta$  takes values in  $H_3(\Omega^{0,1}(\Sigma)) \otimes H_3(\Gamma(L,\Sigma))$  and hence  $\frac{\bar{\partial}_A\beta}{\Psi}$  is a  $(0,1)$  form.

(ii)  $\alpha_1$  is  $C^1$  since it is in  $H_3(\Omega^1(\Sigma, i\mathbb{R}))$  and we are on a Riemann surface.

(iii) We show that  $\Psi$  (which is  $C^1$ ) is zero only on a set of measure zero. Suppose  $\Psi$  is zero on a set of non-zero measure, say on an open set U. Then, since  $A^{0,1} = \overline{\partial} \log \Psi$  and is not finite on U,  $||A||^2$  will not be finite, a contradiction.)

Suppose that,

$$
\Omega(X,Y) = -\frac{1}{2} \int_{\Sigma} \alpha_1 \wedge \alpha_2 - \frac{1}{4} \int_{\Sigma} (\beta H \bar{\eta} - \bar{\beta} H \eta) \omega = 0
$$

for all  $Y = (\alpha_2, \eta) \in T_P C'$ .

Then, we show that  $X = (\alpha_1, \beta) \in T_P C'$  is zero. Recall that  $\alpha_1^{0,1} = -\frac{\bar{\partial}_A \beta}{\Psi}$  $\frac{A\beta}{\Psi}$ , a.e. Take  $\alpha_2 = \alpha_1$  and  $\eta = i\beta$ . Note that this satisfies the linearized equation  $\Psi \alpha_2^{0,1} = -\bar{\partial}_A \eta$  and hence  $(\alpha_2, \eta) \in T_P \mathcal{C}'$ . From the expression of the symplectic form, we get

$$
||\alpha_1||^2 + ||\beta||^2 = 0.
$$

Therefore,  $\alpha_1 = 0$  a.e. and  $\beta = 0$  a.e.

Since  $\alpha_1$  and  $\beta$  are  $C^1$ , therefore,  $\alpha_1$  and  $\beta$  are zero on the complement of a set of measure zero implies that they are identically zero.

b) In this part we mostly follow [\[3\]](#page-8-3) with slight modification.

Let  $\zeta \in \Omega(\Sigma, i\mathbb{R})$  be the Lie algebra of the gauge group (the gauge group element being  $g = e^{\zeta}$ ); note that  $\zeta$  is purely imaginary. It generates a vector field  $X_{\zeta}$  on  $\mathcal{C}'$  as follows :

$$
X_{\zeta}(A,\Psi)=(d\zeta,-\zeta\Psi)\in T_p\mathcal{C}'
$$

where  $p = (A, \Psi) \in \mathcal{C}'$ .

(Note that  $(d\zeta, -\zeta\Psi) \in T_p\mathcal{C}'$  follows from the fact that  $\bar{\partial}_A\Psi = 0$  is invariant under the gauge transformation and can be checked easily too).

We show next that  $X_{\zeta}$  is Hamiltonian. Let us define  $H_{\zeta}: \mathcal{C}' \to \mathbb{C}$  as follows:

 $H_{\zeta}(A,\Psi) = \frac{1}{2} [\int_{\Sigma} \zeta \cdot (F(A) - \frac{1}{2}]$  $\frac{1}{2}(1 - |\Psi|_H^2)\omega$  to be the Hamiltonian for the gauge group action. Then for  $X = (\alpha, \eta) \in T_p \mathcal{C}'$ ,

$$
dH_{\zeta}(X) = \frac{1}{2} \int_{\Sigma} \zeta d\alpha + \frac{1}{4} \int_{M} \zeta (\Psi H \bar{\eta} + \bar{\Psi} H \eta) \omega
$$
  
= 
$$
-\frac{1}{2} \int_{\Sigma} (d\zeta) \wedge \alpha - \frac{1}{4} \int_{\Sigma} [(-\zeta \Psi) H \bar{\eta} - \overline{(-\zeta \Psi)} H \eta] \omega
$$
  
= 
$$
\Omega(X_{\zeta}, X),
$$

where we use that  $\bar{\zeta} = -\zeta$ .

Thus we can define the moment map  $\mu: \mathcal{C}' \to \Omega^2(\Sigma, i\mathbb{R}) = \mathcal{G}^*$  (the dual of the Lie algebra of the gauge group) to be

$$
\mu(A, \Psi) \doteq \frac{1}{2} [F(A) - \frac{1}{2} (1 - |\Psi|_H^2) \omega].
$$

Thus equation (1) is  $\mu = 0$  and the form descends as a symplectic form to  $\mu^{-1}(0)/\mathcal{G}$ . This follows from Marsden and Weinstein, [\[8\]](#page-8-4), who proved the symplectic reduction even for infinite-dimensional case.

The moduli space of vortices,  $Sym^N(\Sigma)$  is a smooth Kähler manifold with the Manton-Nasir Kähler form  $\omega_{MN}$ , (which is equivalent to our metric).

This metric is given by  $([7], eqn (2.14)).$ 

This is the same as the descendent of our metric:

$$
\mathcal{G}(X,X) = \int_{\Sigma} 4|a|^2 dx \wedge dy + \int_{\Sigma} |\beta|_H^2 h^2 dx \wedge dy
$$

$$
= \int_{\Sigma} (\dot{A}_1 \dot{A}_1 + \dot{A}_2 \dot{A}_2) dx \wedge dy + \int_{\Sigma} |\beta|_H^2 h^2 dx \wedge dy
$$

The complex structure defined by us is exactly the usual complex structure of the vortex moduli space i.e. on  $\text{Sym}^N(\Sigma)$ . This can be proved using Ruback's argument mentioned in the appendix of [\[12\]](#page-9-5), or in the mathscinet review of [\[7\]](#page-8-5). (Recall it takes  $\alpha^{0,1}$  to  $i\alpha^{0,1}$  and  $\alpha^{1,0}$  to  $-i\alpha^{1,0}$ , and  $\beta$ to  $i\beta$  which is the case in Ruback's argument).

Now we restrict our attention to the case when the Riemann surface is  $\mathbb{C}P<sup>1</sup>$  of radius R. The Manton-Nasir Kähler form on the moduli space of N vortices on  $\mathbb{C}P^1$  of radius R is given by, [\[7\]](#page-8-5), [\[10\]](#page-9-6),

$$
\omega_{MN} = \frac{i}{2} \sum_{r,s=1}^{N} \left( \frac{4R^2 \delta_{rs}}{(1+|z_r|^2)^2} + 2 \frac{\partial \bar{b_s}}{\partial z_r} \right) dz_r \wedge d\bar{z_s}
$$

This real 2-form is exactly the symplectic reduction of  $\Omega$  on the affine space C of  $(A, \Psi)$  to the vortex moduli space, when the number of vortices is N and the Riemann surface is  $\mathbb{C}P^1$  of radius R. Thus  $\Omega = \omega_{MN}$ . (This statement is true even for a compact Riemann surface of genus  $q > 0$ .)

We study this form on the moduli space of  $N$  vortices on the sphere of radius  $R$ . In the following we refer to Romao, [\[10\]](#page-9-6), where  $\omega_{MN}$  is denoted by  $\omega_{sam}$ . If  $\frac{1}{2\pi}\Omega$  is integral, then  $\frac{1}{2\pi}[\Omega] = \ell[\omega_{FS}]$ , for some  $\ell \in \mathbb{Z}$  since  $H^2(\mathbb{C}P^N,\mathbb{Z})$  is generated by  $[\omega_{FS}]$  (the cohomology class of the Fubini-Study Kähler form.)

By [\[10\]](#page-9-6), we can fine tune the volume of the sphere such that  $\ell = 1$ . Namely, we take  $R^2 = \frac{1}{2} + N$ and  $\kappa = 2k, k \in \mathbb{Z}$ . This satisfies the constraints that  $\kappa$  and  $\kappa(R^2 - N)$  is an integer.

Then  $\left[\frac{1}{2\pi}\Omega\right] = \left[\frac{1}{2\pi}\omega_{MN}\right] = \left[\omega_{FS}\right]$ . (One can also derive the formula  $\frac{1}{2\pi}[\omega_{MN}] = 2(R^2 - N)[\omega_{FS}]$ from Manton and Nasir,  $[7]$   $((3.20),$  pg 600).

Note that the Bradlow criterion for existence of solution holds for this choice of  $R^2$ , since area of the sphere is greater than  $4\pi N$ , [\[10\]](#page-9-6) equation (18), [\[2\]](#page-8-1).

2.3. Quillen Determinant Line bundle and the hyperplane bundle on  $\mathbb{C}P^N$ . We denote the Quillen bundle  $\mathcal{P} = \det(\bar{\partial}_A)$  which is well defined on  $\mathcal{C} = \mathcal{A} \times \Gamma(L)$  (over every  $(A, \Psi)$ ) the fiber is that of det $(\bar{\partial} + A^{0,1})$ ). In [\[4\]](#page-8-6), following Biswas and Raghavendra's work on stable triples, [\[1\]](#page-8-7), we had given  $P$  a modified Quillen metric, namely, we multiply the Quillen metric  $e^{-\zeta'_A(0)}$  by the

factor  $e^{-\frac{i}{4\pi}\int_M |\Psi|^2_H\omega}$ , where recall  $\zeta_A(s)$  is the zeta-function corresponding to the Laplacian of the  $\bar{\partial}$  +  $A^{0,1}$  operator. Here we calculate the curvature for this modified metric on the affine space since the details are not given in [\[4\]](#page-8-6). The factor  $e^{-\zeta'_A(0)}$  contributes  $\frac{i}{2\pi}$  $\left(-\frac{1}{2}\right)$ 2 Z Σ  $\alpha_1 \wedge \alpha_2$  to the curvature, and the factor  $e^{-\frac{i}{4\pi}\int_M |\Psi|^2_H\omega}$  contributes  $\frac{i}{2\pi}$  $\left(-\frac{1}{4}\right)$ 4 Z Σ  $(\beta H \bar{\eta} - \bar{\beta} H \eta)\omega$  to the curvature.

**Lemma 2.** The curvature  $\Omega_{det}$  of  $P$  with the modified Quillen metric is indeed  $\frac{i}{2\pi}\Omega$  on the affine space C.  $\Omega$  is the real  $(1,1)$  given above which is positive.

*Proof.* Quillen, [\[9\]](#page-9-0), constructs the deteminant line bundle on the affine space  $\mathcal{A}^{0,1}$ .

We consider the space of unitary connections, i.e  $\overline{A^{0,1}} = -A^{1,0}$ . The complex structure on C defined by

 $\mathcal{I}$  = " ∗ 0  $0 \quad i$ 1 :  $T_p C \to T_p C$  makes  $(A^{0,1}, \Psi)$  the holomorphic variables on C. This is because  $*\alpha^{0,1} = i\alpha^{0,1}.$ 

Thus the holomorhic coordinate w in [\[9\]](#page-9-0) corresponds to  $A^{0,1}$  in our notation.

By Quillen's computation, the curvature two form is  $\bar{\partial}\partial log||\sigma||^2$  where  $log||\sigma||^2 = -\zeta_A'(0)$  where  $\zeta_A$  is the zeta-function corresponding to the Laplacian of  $\bar{\partial}_A$ .

We first note that  $\bar{\partial}\partial log||\sigma||^2 = \partial \bar{\partial}[-log||\sigma||^2]$  since the order of the derivatives is switched. The Quillen curvature form is  $\bar{\partial}\partial log||\sigma||^2 = \partial \bar{\partial}[-log||\sigma||^2] = \partial \bar{\partial}\zeta'_{A}(0)$ . Now

$$
\frac{\partial^2 \zeta'_A(0)}{\partial w \partial \bar{w}} dw \wedge d\bar{w} = \frac{i}{2\pi} \int_{\Sigma} \overline{\delta A^{0,1}} \wedge \delta A^{0,1},
$$

that is,

$$
\begin{aligned}\n(\frac{\partial^2 \zeta'(0)}{\partial w \partial \bar{w}} dw \wedge d\bar{w}) (\alpha_1^{0,1}, \alpha_2^{0,1}) &= \frac{i}{2\pi} \int_{\Sigma} \frac{1}{2} [\overline{\alpha_1^{0,1}} \wedge \alpha_2^{0,1} - \overline{\alpha_2^{0,1}} \wedge \alpha_1^{0,1}] \\
&= \frac{i}{2\pi} \int_{\Sigma} \frac{-1}{2} [\alpha_1 \wedge \alpha_2]\n\end{aligned}
$$

where recall we have used the fact that  $\alpha_i^{0,1} = -\alpha_i^{1,0}$  This precisely corresponds to  $i/2\pi$  times the first term in our Kähler form.

Next, the term  $e^{-\frac{i}{4\pi}\int_M |\Psi|^2_H\omega}$  contributes to the curvature as:

$$
-\frac{i}{4\pi}\tau = -\frac{i}{4\pi} \int_{\Sigma} <\delta\Psi \wedge \delta\bar{\Psi} >_{H} \omega
$$

where  $-\frac{i}{4}$  $\frac{i}{4\pi}\tau$  is a two-form on the affine space  $\Gamma(\Sigma, L)$ . Details of this calculation can be found in [\[4\]](#page-8-6). Here  $\tau(\beta, \eta) = \frac{1}{2} \int_{\Sigma} (\beta H \bar{\eta} - \bar{\eta} H \beta) \omega$ .

Thus the contribution is

$$
-\frac{i}{4\pi}\tau(\beta,\eta) = \frac{i}{2\pi}\left[-\frac{1}{4}\int_M(\beta H\bar{\eta}-\bar{\eta}H\beta)\omega\right],
$$

which is precisely corresponds to  $i/2\pi$  times the second term in our Kähler form.

To define it on  $\mathcal{C}/\mathcal{G}$ , we repeat the argument in [\[3\]](#page-8-3), [\[4\]](#page-8-6) with  $B = 0$ .

]

**Lemma 3.** P descends to a well defined line bundle (denoted by the same symbol) on  $C/G$ .

*Proof.* Let  $D = \bar{\partial} + A^{(0,1)}$  and let  $D_g = g(\bar{\partial} + A^{(0,1)})g^{-1}$  (the gauge transformed operator), then the gauge transformed Laplacian of D, namely  $\Delta_g = g \Delta g^{-1}$ . Thus there is an isomorphism of eigenspaces, namely,  $s \rightarrow gs$ .

Let  $K^a(\Delta)$  be the direct sum of eigenspaces of the operator  $\Delta$  of eigenvalues  $\langle a, a \rangle$  over the open subset  $U^a = \{A^{(0,1)} | a \notin \text{Spec}\Delta\}$  of the affine space C. The determinant line bundle is defined using the exact sequence

$$
0 \to \text{Ker} D \to K^a(\Delta) \to D(K^a(\Delta)) \to \text{Coker} D \to 0
$$

Thus one identifies

$$
\wedge^{\rm top}({\rm Ker} D)^*\otimes \wedge^{\rm top}({\rm Coker} D)
$$

with  $\wedge^{top}(K^a(\Delta))^* \otimes \wedge^{top}(D(K^a(\Delta)))$  (see [\[9\]](#page-9-0), for more details) and there is an isomorphism of the fibers as  $D \to D_q$ . Therefore one can identify

$$
\wedge^{\text{top}}(K^a(\Delta))^* \otimes \wedge^{\text{top}}(D(K^a(\Delta))) \equiv \wedge^{\text{top}}(K^a(\Delta_g))^* \otimes \wedge^{\text{top}}(D(K^a(\Delta_g))).
$$

Let  $U_g^a = g \cdot U^a$  where  $U_g^a$  is the open set formed out of gauge transformation of  $U^a$ , namely,  $U_g^a = \{A_g, |a \notin Spec(g\Delta g^{-1})\}$ . But  $A_g \in U_g^a$  imples  $a \notin Spec(\Delta)$  and thus  $A_g \in U^a$ . Hence  $U^a \subset U_g^a$ . Similarly,  $U_g^a \subset U^a$ . Thus  $U^a = U_g^a$ .

On  $U^a$  one defines the equivalence class of the fiber

$$
\wedge^{\text{top}}(K^a(\Delta))^* \otimes \wedge^{\text{top}}(D(K^a(\Delta))) \equiv \wedge^{\text{top}}(K^a(\Delta_g))^* \otimes \wedge^{\text{top}}(D(K^a(\Delta_g))).
$$

 $\Box$ 

When the form  $\frac{1}{2\pi}\Omega$  is integral, this construction holds for a compact Riemann surface of genus g. Thus  $P$  descends to the quotient of the affine space  $C$  modulo the gauge group and further restricts to a line bundle on the vortex moduli space. We consider this in the case when the Riemann surface is  $\mathbb{C}P^1$  of radius R where  $R^2 = \frac{1}{2} + N$ . We denote this line bundle again by P and its curvature by  $\Omega_{det} = \frac{i}{2\pi} \Omega$  such that  $\Omega$  coincides with the usual K ähler form on the vortex moduli space, namely  $\omega_{MN}$ , which is cohomologous to  $2\pi\omega_{FS}$ . The Chern class of  $\mathcal P$  is  $\frac{1}{2\pi}[\Omega]$  which is integral and  $\Omega$  is positive.

The first Chern class of  $P$ , namely  $\frac{1}{2\pi}[\Omega]$ , is equal to the first Chern-class of the hyperplane line bundle on  $\mathbb{C}P^N$ , namely  $[\omega_{FS}]$ .

Since the Picard variety of  $\mathbb{C}P^N$  is trivial, the hyperplane bundle is uniquely defined as a holomorphic line bundle by its first Chern class and so is the determinant bundle. Since the first Chern classes agree, they are the equivalent as holomorphic line bundles.

Thus we have the following proposition (where we denote  $\mathcal{P} = \det(\overline{\partial})$ ).

<span id="page-6-0"></span>**Proposition 4.** The determinant line bundle  $\mathcal{P} = \det(\overline{\partial})$  of the family of Cauchy-Riemann operators on  $\mathbb{C}P^1$ , as a line bundle on  $\mathbb{C}P^N$  is equivalent, as holomorphic line bundles, to the hyperplane line bundle  $\mathcal{O}(1)$  on  $\mathbb{C}P^N$ , if the radius R of  $\mathbb{C}P^1$  satisfies  $R^2 = \frac{1}{2} + N$ .

#### 3. Kodaira embedding theorem and application

<span id="page-7-0"></span>Here we recall the Kodaira embedding theorem (cf. [\[5,](#page-8-2) [16\]](#page-9-7)) in a form that is suitable for our application.

<span id="page-7-2"></span>**Theorem 2** (Kodaira embedding theorem). Let M be a compact Kähler manifold with integral Kähler form  $\omega$ . Let  $\mathcal{L} \to M$  be the line bundle with connection  $\nabla$  and curvature  $\omega$ . Then there is a positive integer  $k_0$  (which we will assume is minimal) such that for all  $k \geq k_0$  the natural map

$$
\phi_k: M \hookrightarrow \mathbb{P}(H^0(M, \mathcal{O}(\mathcal{L}^{\otimes k}))^*)
$$

is a holomorphic embedding. In particular, one has the equality of cohomology classes  $[\phi_k^*(\Omega_k)] =$  $k[\omega]$ , where  $\Omega_k$  denotes the Kähler form of the Fubini-Study metric on  $\mathbb{P}(H^0(M,\mathcal{O}(\mathcal{L}^{\otimes k}))^*)$ . In  $\textit{fact}, \ \phi_k^*(\mathcal{O}(1)) \cong \mathcal{L}^{\otimes k}$  as holomorphic line bundles on M, where  $\mathcal{O}(1)$  is the hyperplane bundle  $\mathbb{P}(H^0(M,\mathcal{O}(\mathcal{L}^{\otimes k}))^*).$ 

Here  $H^0(M, \mathcal{O}(\mathcal{L}^{\otimes k}))$  denotes the finite dimensional vector space of all holomorphic sections of the line bundle  $\mathcal{L}^{\otimes k}$ , which has a natural  $L^2$ -metric using the Kähler structure on M, which in turn induces the Fubini-Study Kähler form  $\Omega_k$  on  $\mathbb{P}(H^0(M,\mathcal{O}(\mathcal{L}^{\otimes k}))^*)$ .

For instance, for a compact Riemann surface of genus greater than 2, the minimal choice of  $k_0$ is equal to 3.

Proof. Here we outline a few more details re the last half of the Kodaira embedding theorem above, which may not appear in the standard versions of the theorem in reference books (cf. [\[5,](#page-8-2) [16\]](#page-9-7)). The vector space of all holomorphic sections  $H^0(M, \mathcal{O}(\mathcal{L}^{\otimes k})$  of  $\mathcal{L}^{\otimes k}$  has an inner product induced by the Kähler volume form,

$$
\langle \sigma_1, \sigma_2 \rangle = \int_M h^{\otimes k}(\sigma_1(x), \sigma_2(x)) \frac{\omega^n}{n!}
$$

where  $n = \dim_{\mathbb{C}}(M)$  and  $h^{\otimes k}$  is the induced hermitian inner product on  $\mathcal{L}^{\otimes k}$ . This in turn determines a Fubini-Study Kähler form  $\Omega_k$  on  $\mathbb{P}(H^0(M,\mathcal{O}(\mathcal{L}^{\otimes k}))^*)$ , which is the first Chern class of the hyperplane bundle  $\mathcal{O}(1)$ , satisfying  $[\phi_k^*(\Omega_k)] = k[\omega] \in H^{1,1}(M) \cap H^2(M;\mathbb{Z})$ . Therefore  $\phi_k^*(\mathcal{O}(1)) \cong \mathcal{L}^{\otimes k}$  as holomorphic line bundles on M.

Proof of Theorem [1.](#page-0-0) By Proposition [4,](#page-6-0) we know that Theorem [1](#page-0-0) is true for complex projective space  $\mathbb{C}P^N$  for all positive integers N. By the Kodaira embedding theorem, Theorem [2,](#page-7-2) and with the  $N = \dim H^0(M, \mathcal{O}(\mathcal{L}^{\otimes k_0})) - 1$  $N = \dim H^0(M, \mathcal{O}(\mathcal{L}^{\otimes k_0})) - 1$ , Theorem 1 is true in the general case.

## 4. Symplectic variant and application

<span id="page-7-1"></span>The symplectic variant of the Kodaira embedding theorem is due to Gromov and we quote a version in [\[15\]](#page-9-8), Remark following Theorem B.

<span id="page-7-3"></span>**Theorem 3** (Gromov's embedding theorem). Let M be a compact symplectic manifold with integral symplectic form  $\omega$ . Let  $\mathcal{L} \to M$  be the line bundle with connection  $\nabla$  and curvature  $\omega$ . Then there is a positive integer  $k_0$  (which we will assume is minimal) such that for all  $k \geq k_0$  there is a symplectic embedding

$$
\phi_k: M \hookrightarrow \mathbb{C}P^k.
$$

In particular,  $\phi_k^*(\Omega_k) = \omega$ , where  $\Omega_k$  denotes the symplectic form of the Fubini-Study metric on  $\mathbb{C}P^{k}.$ 

Our next result is the symplectic analogue of Theorem [1.](#page-0-0)

<span id="page-8-8"></span>**Theorem 4.** Any compact symplectic manifold M with integral symplectic form  $\omega$ , parametrizes a natural smooth family of Cauchy-Riemann operators  $\{\overline{\partial}_z : z \in M\}$  on  $\mathbb{C}P^1$  such that the determinant line bundle det $(\overline{\partial}) \cong \mathcal{L}$  as complex line bundles, where  $\mathcal{L}$  is the prequantum line bundle determined by  $\omega$ .

<span id="page-8-0"></span>*Proof.* By Proposition [4,](#page-6-0) we know that Theorem [4](#page-8-8) is true for complex projective space  $\mathbb{C}P^N$  for all positive integers N. By Gromov's embedding theorem, Theorem [3,](#page-7-3) and with the minimal choice of  $k_0$  as before, Theorem [1](#page-0-0) is true in the general case.

### 5. Conjecture

We conjecture that equivariant versions of our results are true. More precisely,

**Conjecture.** Let M be a compact Kähler G-manifold M with integral Kähler form  $\omega$ , where G is a compact Lie group. That is,  $\omega$  is a G-invariant Kähler form on the G-manifold M and all structures are G-invariant. Let  $\mathcal L$  denote the holomorphic G-line bundle determined by  $\omega$ . Then M parametrizes a natural holomorphic G-equivariant family of Cauchy-Riemann operators  ${\{\overline{\partial}_z : z \in M\}}$  on a compact Kähler G-manifold Z such that the Quillen determinant line bundle  $\det(\overline{\partial}) \cong \mathcal{L}^{\otimes k}$  as holomorphic G-line bundles, for some sufficiently large k.

There is an analogous conjecture in the equivariant symplectic case. If the conjectures are valid, then they would apply to coadjoint integral maximal orbits and thus to representation theory. It would also be relevant to the popularly known principle, "quantization commutes with reduction" of Guillemin and Sternberg, cf. [\[6\]](#page-8-9)

#### **REFERENCES**

- <span id="page-8-7"></span><span id="page-8-1"></span>[1] I. Biswas and N. Raghavendra, The determinant bundle on the moduli space of stable triples over a curve. Proc. Indian Acad. Sc. Mat. Sci. 112, no. 3, 367-382, (2002).
- <span id="page-8-3"></span>[2] S. Bradlow, Vortices in holomorphic line bundles over closed K¨ahler manifolds. Commun.Math.Phys. 135, 1-17 (1990).
- <span id="page-8-6"></span>[3] R. Dey, Geometric prequantization of the moduli space of the vortex equations on a Riemann surface; J. of Math. Phys., vol. 47, issue 10, (2006), page 103501-103508; [math-ph/0605025.](http://arxiv.org/abs/math-ph/0605025)
- [4] R. Dey, Erratum: Geometric prequantization of the moduli space of the vortex equations on a Riemann surface. J. of Math. Phys. 50, 119901 (2009).
- <span id="page-8-9"></span><span id="page-8-2"></span>[5] P. Griffiths and J. Harris, Principles of algebraic geometry. Pure and Applied Mathematics. Wiley-Interscience [John Wiley & Sons], New York, 1978.
- <span id="page-8-5"></span>[6] V. Guillemin and S. Sternberg, Geometric quantization and multiplicities of group representations, Invent. Math. 67 (1982), no. 3, 515–538.
- <span id="page-8-4"></span>[7] N.S. Manton and S.M. Nasir, Volume of vortex moduli spaces. Comm. Math. Phys. 199, no. 3, 591–604 (1998).
- [8] J.M. Marsden and A. Weinstein, Reduction of symplectic manifolds with symmetry. Rep. Math. Phys, 5, no.1, 121-130 (1974).
- <span id="page-9-6"></span><span id="page-9-0"></span>[9] D.G. Quillen, Determinants of Cauchy-Riemann operators on Riemann surfaces. Functional Anal. Appl. 19 (1985), no. 1, 31-34.
- <span id="page-9-1"></span>[10] N. M. Romao, Quantum Chern-Simons vortices on a sphere. J.Math. Phys., 42, no. 8, 3445-3469 (2001).
- <span id="page-9-5"></span>[11] J. H. Rawnsley, Coherent states and Kähler manifolds, Quart. J. Math. Oxford (2) 28 (1977), 403-415.
- <span id="page-9-2"></span>[12] T.M. Samols, Vortex scattering, Commun. Math. Phy. 145, 149-179 (1992).
- <span id="page-9-3"></span>[13] M. Spera, On Kählerian coherent states, Geometry, Integrability and Quantization, Sept , 1999, Coral Press.
- <span id="page-9-8"></span>[14] A. Jaffe and C. Taubes, Vortices and monopoles. Structure of static gauge theories. Progress in Physics, 2. Birkhuser, Boston, Mass., 1980.
- <span id="page-9-7"></span>[15] D. Tischler, Closed 2-forms and an embedding theorem for symplectic manifolds. J. Differential Geometry, 12 (1977), no. 2, 229-235.
- <span id="page-9-4"></span>[16] R.O. Wells Jr. Differential analysis on complex manifolds. Prentice-Hall Series in Modern Analysis. Prentice-Hall, Inc., Englewood Cliffs, N.J., 1973.
- [17] N.M.J. Woodhouse. Geometric Quantization, Second Edition, Clarendon Press, Oxford, 1994.

Department of Mathematics, Harishchandra Research Institute, Allahabad, India E-mail address: rkmn@hri.res.in

Department of Pure Mathematics, University of Adelaide, Adelaide 5005, Australia  $E\text{-}mail\ address: \text{mathai}.\text{varghese@adelaide.edu.au}$