

On the FitzHugh - Nagumo Model

M. DE ANGELIS, P. RENNO *

Abstract

The initial value problem \mathcal{P}_0 , in all of the space, for the spatio-temporal FitzHugh - Nagumo equations is analyzed. When the reaction kinetics of the model can be outlined by means of piecewise linear approximations, then the solution of \mathcal{P}_0 is explicitly obtained. For periodic initial data are possible damped travelling waves and their speed of propagation is evaluated. The results imply applications also to the non linear case.

Keywords: *Parabolic systems. Biological models. Laplace transforms*

1 Introduction

One of the reaction diffusion systems which models various important biological phenomena is given by

$$(1.1) \quad \begin{cases} \frac{\partial u}{\partial t} = \varepsilon \frac{\partial^2 u}{\partial x^2} - v + f(u) \\ \frac{\partial v}{\partial t} = bu - \beta v \end{cases}$$

where the appropriate class of functions $f(u)$ depends on the reaction kinetics of the model [1] - [4]. In the theory of nerve membranes, for example, the

*Faculty of Engineering, Department of Mathematics and Applications, Naples, Italy.
e-mail modeange@unina.it

system (1.1) is related to the spatio-temporal FitzHugh - Nagumo equations (FHN) with

$$(1.2) \quad f(u) = u(a - u)(u - 1) \quad (0 < a < 1)$$

and where $u(x, t)$ models the transmembrane voltage of a nerve axon at distance x and time t , $v(x, t)$ is an auxiliary variable that acts as recovery variable. Further, the diffusion coefficient ε and the parameters b, β are all non negative [5],[9].

In addition to the propagation of nerve action potentials, (1.1) - (1.2) govern other several biological and biochemical phenomena; the list of references is very long and the variety of analytical aspects examined is wide. [10] - [15]. Further, we must remark travelling pulses and periodic wavetrains obtained by means of piecewise linear approximations of $f(u)$ as

$$(1.3) \quad f(u) = \eta(u - a) - u \quad (0 < a < 1)$$

where η denotes the unit- step function.[16]- [21].

Typical boundary value problems related to the linear case (1.3) can be explicitly solved. Aim of this paper is the analysis of the initial value problem \mathcal{P}_0 in all of the space; the fundamental solution and the explicit solution of \mathcal{P}_0 are determined when (1.3) holds. More, in the non linear case of the FHN model given by (1.1)-(1.2), the problem \mathcal{P}_0 is reduced to appropriate integral equations whose kernels are functions characterized by basic properties. All this implies existence and uniqueness properties, together with a priori estimates.

2 Statement of the problem and results

Both the non linear source (1.2) and the linear approximation (1.3) involve a linear term $-ku$ with $k = a$ for (1.2), and $k = 1$ for (1.3). As consequence, the system (1.1) becomes

$$(2.1) \quad \begin{cases} u_t - \varepsilon u_{xx} + k u + v = \varphi(u) \\ v_t + \beta v - b u = 0 \end{cases}, \quad (x, t) \in \mathfrak{R}$$

where $\varphi(u) = u^2(a + 1 - u)$ when f is given by (1.2), while, in the linear case, $\varphi(u)$ is equal to the constant $\bar{\eta}$ that holds zero or one.

The initial- value problem \mathcal{P}_0 related to (2.1) with $\varphi(u) = \bar{\eta}$ is analyzed in the set

$$(2.2) \quad \Omega_T = \{(x, t) : x \in \mathfrak{R}, \quad 0 < t \leq T\},$$

with the conditions

$$(2.3) \quad u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), \quad x \in \mathfrak{R}.$$

When the linear approximation of f holds, then the explicit solution of the problem \mathcal{P}_0 can be obtained by means of functional transforms (Fourier with respect to x and Laplace as for t). If one puts formally

$$(2.4) \quad \hat{u}(x, s) = \int_0^\infty e^{-st} u(x, t) dt, \quad \hat{v}(x, s) = \int_0^\infty e^{-st} v(x, t) dt,$$

from (2.1) (2.3) one deduces:

$$(2.5) \quad \hat{u} = \int_{\mathfrak{R}} [u_0(\xi) + \frac{\bar{\eta}}{s}] \hat{K}_0[(x - \xi, s)] d\xi - \int_{\mathfrak{R}} v_0(\xi) \hat{K}_1(x - \xi, s) d\xi$$

$$(2.6) \quad \hat{v} = b \int_{\mathfrak{R}} [u_0(\xi) + \frac{\bar{\eta}}{s}] \hat{K}_1(x - \xi, s) d\xi - b \int_{\mathfrak{R}} v_0(\xi) \hat{K}_2(x - \xi, s) d\xi + \frac{v_0(x)}{s + \beta},$$

where

$$(2.7) \quad \hat{K}_n(x, s) = \frac{e^{-\frac{|x|}{\sqrt{\varepsilon}} \sigma}}{2 \sqrt{\varepsilon} \sigma (s + \beta)^n} \quad (n = 0, 1, 2)$$

with $\sigma^2 = s + k + \frac{b}{s + \beta}$.

Now, let us consider the fundamental solution

$$(2.8) \quad g(x, y) = \frac{1}{2 \sqrt{\pi \varepsilon y}} e^{-ky - \frac{x^2}{4\varepsilon y}}$$

of the heat equation $\varepsilon g_{xx} - g_y - k g = 0$ and, for $n = 0, 1, 2$, let

$$(2.9) \quad G_n(x, t) = \int_0^t e^{-\beta(t-y)} g(x, y) \left(\frac{t-y}{by}\right)^{\frac{n-1}{2}} J_{n-1}\left(2\sqrt{b(t-y)y}\right) dy,$$

where $J_n(z)$ denotes the Bessel function of first kind. Then, if one puts

$$(2.10) \quad K_0 = g(x, t) + G_0(x, t), \quad K_i = G_i(x, t) \quad (i = 1, 2),$$

the following theorems hold.

Theorem 2.1- *In the half-plane $\Re s > \max(-k, -\beta)$, the Laplace integrals of $K_n(x, t)$ ($n = 0, 1, 2$) converge absolutely for all $|x| > 0$, and one has $\mathcal{L}_t K_n(x, t) = \hat{K}_n(x, s)$.*

Theorem 2.2- *The functions K_0, K_1, K_2 are $C^\infty(\Omega_T)$ solutions of the integro differential equation :*

$$(2.11) \quad z_t - \varepsilon z_{xx} + k z + b \int_0^t e^{-\beta(t-\tau)} z(x, \tau) d\tau = 0$$

and have the same basic properties of the fundamental solution (2.8) of the heat operator. Further, for $i = 0$ and $i = 1$, it results:

$$(2.12) \quad K_i(x, t) = (\partial_t + \beta) K_{i+1}, \quad \lim_{t \downarrow 0} K_{i+1} = 0 \quad (i = 0, 1)$$

while $\lim_{t \rightarrow 0} K_0(x, t) = 0$ only for $|x| > 0$.

These properties assure the convergence of the convolutions

$$(2.13) \quad K_n * \psi = \int_{\mathfrak{R}} K_n(x - \xi, t) \psi(\xi) d\xi \quad (n = 0, 1, 2)$$

for all the functions that satisfy a *growth condition* of the form

$$(2.14) \quad |\psi(x)| < c_1 \exp [c_2 |x|^{\alpha+1}], \quad 0 < \alpha < 1$$

with c_1 and c_2 positive constants.

Further, let consider the following functions

$$(2.15) \quad N_1(t) = \frac{\bar{\eta}\beta}{b + \beta} \left[1 - e^{-\frac{1+\beta}{2}t} \cos(\gamma t) \right] + \frac{\bar{\eta}(2b + \beta - \beta^2)}{2\gamma(b + \beta)} e^{-\frac{1+\beta}{2}t} \sin(\gamma t)$$

$$(2.16) \quad N_2(t) = \frac{\bar{\eta}}{b + \beta} \left[1 - e^{-\frac{1+\beta}{2}t} \cos(\gamma t) \right] - \frac{\bar{\eta}(1 + \beta)}{2\gamma(b + \beta)} e^{-\frac{1+\beta}{2}t} \sin(\gamma t),$$

where $\gamma = \left[b - (\beta - 1)^2/4 \right]^{1/2}$. Then, by (2.5)-(2.6) and the foregoing statements, the explicit solution of the linear problem \mathcal{P}_0 is given by

$$(2.17) \quad \begin{cases} u = N_1(t) + u_0 * K_0 - v_0 * K_1 \\ v = N_2(t) + u_0 * K_1 - v_0 * K_2 + v_0(x) e^{-\beta t}, \end{cases}$$

and the following conclusion is deduced.

Theorem 2.3 - *When the data (u_0, v_0) are continuous functions that satisfy the growth condition (2.14), then the formulae (2.17) represent the unique solution of the problem \mathcal{P}_0 in the class of solutions compatible with (2.14).*

3 Travelling waves and a priori estimates

A first example of applications is related to the linear case and concerns the analysis of travelling waves. By the explicit solution (2.17), for instance, when $\bar{\eta} = 0$, $v_0 = 0$, $u_0 = A \cos(wx)$, then one obtains:

$$(3.1) \quad u = (\partial_t + \beta)w, \quad v = bw$$

where

$$(3.2) \quad w = \frac{A}{2\alpha} e^{-\frac{1+\beta+\varepsilon w^2}{2}t} [\sin(wx + \alpha t) - \sin(wx - \alpha t)]$$

with $\alpha = [b - (\frac{1+\varepsilon w^2 - \beta}{2})^2]^{1/2}$. So, when $b > (\frac{1+\varepsilon w^2 - \beta}{2})^2$, there exist damped travelling waves with speed equal to α/w .

Moreover, in the non linear case, by (2.5),(2.6) one deduces the following integral equations

$$(3.3) \left\{ \begin{array}{l} u = u_0 * K_0 - v_0 * K_1 + \int_0^t \varphi(u) * K_0 d\tau \\ v = bu_0 * K_1 - bv_0 * K_2 + v_0(x) e^{-\beta t} + b \int_0^t \varphi(u) * K_1 d\tau \end{array} \right.$$

that imply a priori estimates. In fact, in the class of bounded solutions, if one puts

$$(3.4) \quad \|u_0\| = \sup_{x \in \mathfrak{R}} |u_0(x)|, \quad \|v_0\| = \sup_{x \in \mathfrak{R}} |v_0(x)|, \quad \|\varphi\| = \sup_{u \in \mathfrak{R}} |\varphi(u)|$$

by means of the basic properties of the kernels K_0, K_1, K_2 , one obtains:

$$(3.5) \quad \begin{cases} |u| < c_1 \|\varphi\| + c_2 (\|u_0\| t + \|v_0\|) E(t) \\ |v| < c_1 \|\varphi\| + c_2 (\|u_0\| + \|v_0\| t) E(t) \end{cases}$$

where

$$(3.6) \quad E(t) = \frac{e^{-kt} - e^{-\beta t}}{\beta - k}$$

and where the constants c_1, c_2 depend on the parameters b, k, β . Worthy of remark is the fact that the estimates (3.5) hold for all t , also when $T \rightarrow \infty$.

References

- [1] Izhikevich E.M. *Dynamical Systems in Neuroscience: The Geometry of Excitability and Bursting*. The MIT press. England (2007)
- [2] J.D. Murray, *Mathematical Biology. II. Spatial models and biomedical applications*, Springer-Verlag, N.Y 2003
- [3] J.D. Murray, *Mathematical Biology. I. An Introduction*, Springer-Verlag, N.Y 2002
- [4] J. P. Keener - J. Sneyd *Mathematical Physiology* Springer-Verlag, N.Y 1998
- [5] Krupa, M; Sandstede, B; Szmolyan, P *Fast and slow waves in the FitzHugh-Nagumo equation*. J. Differential Equations 133 (1997), no. 1, 49–97.
- [6] L.Glass, M.E. Josephson *Resetting and annihilation of reentrant abnormally rapid heartbeat* Phys. Rev. Lett 75,10 (1995) 2059-2062
- [7] J Rinzel *Models in neurobiology* Nonlinear Phenomena in Physics and Biology. Edit by R. H. Enns, B. L. Jones, R. M. Miura, and S.nd S. Rangnekar. D. Reidel Publishing Company, Dordrecht-Holland. NATO Advanced Study Institutes Series. Volume B75, (1981), p.345

- [8] Nagumo, J., Animoto, S., Yoshizawa, S *An active pulse transmission line simulating nerve axon* (1962), Proc. Inst. Radio Engineers, 50, 2061-2070.
- [9] FitzHugh R. *Impulses and physiological states in theoretical models of nerve membrane*. Biophysical journal 1. (1961), 445- 466.
- [10] Wang, Jiang; Zhang, Ting; Deng, Bin *Synchronization of FitzHugh-Nagumo neurons in external electrical stimulation via nonlinear control*. Chaos Solitons Fractals 31 (2007), no. 1, 30–38.
- [11] J.G. Alford, G. Auchmuty *Rotating wave solutions of the FitzHugh-Nagumo equations*. J. Math. Biol. 53(2006) 797-819
- [12] Bini D., Cherubini C., Filippi S. *Heat transfer in FitzHugh-Nagumo models*. Physical Review E. 74, 041905 (2006)
- [13] Bini, D.; Cherubini, C.; Filippi, S. *Viscoelastic FitzHugh-Nagumo models*. Phys. Rev. E (3) 72 (2005), no. 4, 041905-01 12
- [14] Rinzel, J; Terman, D *Propagation phenomena in a bistable reaction-diffusion system*. SIAM J. Appl. Math. 42 (1982), no. 5, 1111–1137.
- [15] J. Nagumo, S. Yoshizawa, Animoto, S. *Bistable Transmission Lines* IEEE Transactions on Circuits and Systems, 12, Issue 3, (1965) Page(s): 400 - 412
- [16] McKean, H. P., Jr. *Nagumo's equation*. Advances in Math. 4 1970 209–223 (1970).
- [17] J Rinzel and J B Keller. *Traveling wave solutions of a nerve conduction equation*. Biophysical Journal, 13: 1313-1337, 1973.
- [18] Rinzel J. *Spatial stability of traveling wave solutions of a nerve conduction equation*. Biophys J. 1975 15 975-988.
- [19] JA Feroe *Temporal stability of solitary impulse solutions of a nerve equation* Biophys. J. 1978 21 103-110.
- [20] J. A. Feroe, *Existence of travelling wave trains in nerve axon equations*, SIAM J. Appl. Math. 46 (1986), 1079-1097.
- [21] A. Tonnelier, *The McKean's caricature of the FitzHugh-Nagumo model. I : The space-clamped system* , SIAM Journal on Applied Mathematics 63, pp. 459-484 (2002)