

ON A PARTICULAR CASE OF THE DIRICHLET'S THEOREM AND THE MIDY'S PROPERTY

JOHN H. CASTILLO, GILBERTO GARCÍA-PULGARÍN,
AND JUAN MIGUEL VELÁSQUEZ-SOTO

ABSTRACT. We give new characterizations of the Midy's property and using these results we obtain a new proof of a special case of the Dirichlet's theorem about primes in arithmetic progression.

1. PRELIMINARIES

Let b be a positive integer greater than 1, b will denote the base of numeration, N a positive integer relatively prime to b , i.e $(N, b) = 1$, $|b|_N$ the order of b in the multiplicative group \mathbb{U}_N of positive integers less than N and relatively primes to N , and $x \in \mathbb{U}_N$. It is well known that when we write the fraction $\frac{x}{N}$ in base b , it is periodic. By period we mean the smallest repeating sequence of digits in base b in such expansion, it is easy to see that $|b|_N$ is the length of the period of the fractions $\frac{x}{N}$ (see Exercise 2.5.9 in [Nat00]). Let d, k be positive integers with $|b|_N = dk$, $d > 1$ and $\frac{x}{N} = 0.\overline{a_1 a_2 \cdots a_{|b|_N}}$ where the bar indicate the period and a_i 's are digits in base b . We separate the period $a_1 a_2 \cdots a_{|b|_N}$ in d blocks of length k and let

$$A_j = [a_{(j-1)k+1} a_{(j-1)k+2} \cdots a_{jk}]_b$$

be the number represented in base b by the j -th block and $S_d(x) = \sum_{j=1}^d A_j$.

If for all $x \in \mathbb{U}_N$, the sum $S_d(x)$ is a multiple of $b^k - 1$ we say that N has the Midy's property for b and d . It is named after E. Midy (1836), to read historical aspects about this property see [Lew07] and its references.

We denote with $\mathcal{M}_b(N)$ the set of positive integers d such that N has the Midy's property for b and d and we will call it the Midy's set of N to base b . As usual, let $\nu_p(N)$ be the greatest exponent of p in the prime factorization of N .

For example 13 has the Midy's property to the base 10 and $d = 3$, because $|13|_{10} = 6$, $1/13 = 0.\overline{076923}$ and $07 + 69 + 23 = 99$. Also, 75 has the Midy's property to the base 8 and $d = 4$, since $|75|_8 = 20$, $1/75 = [0.\overline{00664720155164033235}]_8$ and $[00664]_8 + [72015]_8 + [51640]_8 + [33235]_8 = 2 * (8^5 - 1)$. But 75 does not have the Midy's property to 8 and 5. Actually, we can see that $\mathcal{M}_{10}(13) = \{2, 3, 6\}$ and $\mathcal{M}_8(75) = \{4, 20\}$.

2000 *Mathematics Subject Classification.* 11A05, 11A07, 11A15, 11A63, 16U60.

In [GPG09] is given the following characterization of Midy's property.

Theorem 1. *If N is a positive integer and $|b|_N = kd$, then $d \in \mathcal{M}_b(N)$ if and only if $\nu_p(N) \leq \nu_p(d)$ for all prime divisor p of $(b^k - 1, N)$.*

The next theorem is a different way to write Theorem 1.

Theorem 2. *Let N be a positive integer and d a divisor of $|b|_N$. The following statements are equivalent*

- (1) $d \in \mathcal{M}_b(N)$
- (2) *For each prime divisor p of N such that $\nu_p(N) > \nu_p(d)$, there exists a prime q divisor of $|b|_N$ that satisfies $\nu_q(|b|_p) > \nu_q(|b|_N) - \nu_q(d)$.*

In [CGPVS11] the authors prove the following theorem.

Theorem 3. *Let d_1, d_2 be divisors of $|b|_N$ and assume that $d_1 \mid d_2$ and $d_1 \in \mathcal{M}_b(N)$, then $d_2 \in \mathcal{M}_b(N)$.*

The following result has a big influence on our work, it is Theorem 3.6 in [Nat00].

Theorem 4. *Let p be an odd prime not divisor of b , $m = \nu_p(b^{|b|_p} - 1)$ and let t a positive integer, then*

$$|b|_{p^t} = \begin{cases} |b|_p & \text{if } t \leq m, \\ p^{t-m} |b|_p & \text{if } t > m. \end{cases}$$

The Dirichlet's theorem about primes in arithmetic progression states that there are infinitely many primes in any arithmetic progression of initial term a and common difference l , with $(a, l) = 1$. It was conjectured by Carl Friedrich Gauss and it was first proved in 1826 by Peter Gustav Lejeune Dirichlet. The theorem which establishes that there are infinitely many primes is a particular case of the Dirichlet's theorem, taking $a = 1$ and $l = 1$.

A related result, to the Dirichlet's theorem, proved by B. Green and T. Tao in 2004, guarantees that we can find lists of primes in arithmetic progression of arbitrary length; see [GT04].

The biggest list of primes in arithmetic progression has 26 terms, it was founded in April 12, 2010 by Benot Perichon and PrimeGrid. It is formed by the numbers $43142746595714191 + 23681770 \cdot 23\#\cdot n$, for $0 \leq n \leq 25$ where $a\#$ denotes the primorial of a , it is the product of all primes less or equal to a ; see [And12].

2. OTHER CHARACTERIZATIONS OF MIDY'S PROPERTY.

In this section, we will study some consequences of Theorem 2.

Theorem 5. *Let N be a positive integer and $|b|_N = kd$. If, for all prime divisor p of N , we have $\nu_p(N) > \nu_p(d)$, then the followings statements are equivalent*

- (1) $(b^k - 1, N) = 1$
- (2) $d \in \mathcal{M}_b(N)$
- (3) *For each prime divisor p of N , there exists a prime q divisor of d such that $\nu_q(|b|_p) > \nu_q(|b|_N) - \nu_q(d)$.*

Proof. The equivalence between (2) and (3) is immediate from Theorem 2.

By Theorem 1 we get that (1) implies (2). Now we prove that (2) implies (1). Suppose that $d \in \mathcal{M}_b(N)$ and let $g = (b^k - 1, N)$. If there exists a prime divisor p of g , from Theorem 1, we have $0 < \nu_p(N) \leq \nu_p(d)$ and this is impossible because $\nu_p(d) < \nu_p(N)$. Therefore $(b^k - 1, N) = 1$. \square

We now will study Theorem 4 of [Lew07], which is attributed by its author to M. Jenkins (1867). Let p_1, p_2, \dots, p_t be different primes such that $d \in \mathcal{M}_b(p_i)$ for each i and let h_1, h_2, \dots, h_t be positive integers, when does $N = p_1^{h_1} p_2^{h_2} \dots p_t^{h_t}$ have the Midy's property for b and d ? The Jenkins' Theorem gives the answer to this question and the same is independent from the h_i 's.

By simplicity we go to study the case when $t = 3$, but the argument is true for any t . Let p_1, p_2, p_3 be different primes such that $d \in \mathcal{M}_b(p_i)$, $|b|_{p_i} = dk_i$, $m_i = \nu_{p_i}(b^{|b|_{p_i}} - 1)$ for $i = 1, 2, 3$ then

$$|b|_N = d \left[p_1^{h_1 - m_1} k_1, p_2^{h_2 - m_2} k_2, p_3^{h_3 - m_3} k_3 \right] = dk.$$

We have to check up the prime divisors of $(b^k - 1, N)$, to determine when $d \in \mathcal{M}_b(N)$. As d is a divisor of $|b|_{p_1}$ thus $d \leq p_1 - 1$ and if, say, $p_1 \mid (b^k - 1, N)$ we get that $h_1 = \nu_{p_1}(N) > 0 = \nu_{p_1}(d)$ and so N does not have the Midy's property for b and d . In consequence $d \in \mathcal{M}_b(N)$ if and only if $(b^k - 1, N) = 1$. It is clear that $(b^k - 1, N) = 1$ is equivalent to say that for each i , $|b|_{p_i} \nmid k$. We will see when this fact is verified. Let $d = \prod_{i=1}^s q_i^{r_i}$ be the prime decomposition of d , for each $i = 1, 2, 3$ take $c_i = \nu_d(k_i)$ and

$$\begin{aligned} p^{h_1 - m_1} k_1 &= \left(d^{c_1} \prod_{i=1}^s q_i^{\alpha_i^{(1)}} \right) y_1 \\ p^{h_2 - m_2} k_2 &= \left(d^{c_2} \prod_{i=1}^s q_i^{\alpha_i^{(2)}} \right) y_2 \\ p^{h_3 - m_3} k_3 &= \left(d^{c_3} \prod_{i=1}^s q_i^{\alpha_i^{(3)}} \right) y_3 \end{aligned}$$

where $(q_i, y_j) = 1$ for $i = 1, 2, \dots, s$ and $j = 1, 2, 3$. Then

$$\begin{aligned} k &= \left[p^{h_1-m_1} k_1, p^{h_2-m_2} k_2, p^{h_3-m_3} k_3 \right] \\ &= \left[d^{c_1} \prod_{i=1}^s q_i^{\alpha_i^{(1)}}, d^{c_2} \prod_{i=1}^s q_i^{\alpha_i^{(2)}}, d^{c_3} \prod_{i=1}^s q_i^{\alpha_i^{(3)}} \right] [y_1, y_2, y_3]. \end{aligned}$$

We want, as was said before, that k not be divisible for any $|b|_{p_i}$, $i = 1, 2, 3$. Now if, say, $k = |b|_{p_1} l$ with l integer, then $k = |b|_{p_1} l = (k_1 d) l$ therefore $\frac{k}{k_1}$ is a multiple of d , and as the y_i 's are relatively primes with d this is equivalent to

$$\frac{\left[d^{c_1} \prod_{i=1}^s q_i^{\alpha_i^{(1)}}, d^{c_2} \prod_{i=1}^s q_i^{\alpha_i^{(2)}}, d^{c_3} \prod_{i=1}^s q_i^{\alpha_i^{(3)}} \right]}{d^{c_1} \prod_{i=1}^s q_i^{\alpha_i^{(1)}}} \equiv 0 \pmod{d}$$

From the above analysis follows the next theorem.

Theorem 6 (Jenkins' Theorem). *Let p_1, p_2, \dots, p_t be different primes such that $d \in \mathcal{M}_b(p_i)$ for each i , let h_1, h_2, \dots, h_t be positive integers and $N = p_1^{h_1} p_2^{h_2} \dots p_t^{h_t}$. With the notations introduced above we obtain that $d \in \mathcal{M}_b(N)$ if and only if, for each $j = 1, 2, \dots, t$ is satisfied*

$$\frac{\left[d^{c_1} \prod_{i=1}^s q_i^{\alpha_i^{(1)}}, d^{c_2} \prod_{i=1}^s q_i^{\alpha_i^{(2)}}, \dots, d^{c_t} \prod_{i=1}^s q_i^{\alpha_i^{(t)}} \right]}{d^{c_t} \prod_{i=1}^s q_i^{\alpha_i^{(j)}}} \not\equiv 0 \pmod{d}$$

Our next result has a similar flavor of the Jenkins' Theorem, although its statement and proof are simpler.

Theorem 7. *Let N, q, v be integers with q prime and $v > 0$. Then $q^v \in \mathcal{M}_b(N)$ if and only if $N = q^n p_1^{h_1} p_2^{h_2} \dots p_t^{h_t}$ where n is a non-negative integer, p_i 's are different primes and h_i 's are non-negatives integers not all zero, verifying $0 \leq n \leq v$, $\nu_q(|b|_{p_i}) > 0$ and*

$$\max_{1 \leq i \leq t} \left\{ n - m, \nu_q(|b|_{p_i}) \right\} - v < \min_{1 \leq i \leq t} \left\{ \nu_q(|b|_{p_i}) \right\}$$

where $m = \nu_q(b^{|b|_q} - 1)$.

Proof. We write $|b|_N = q^t k$, with $(q, k) = 1$ and $t \geq v$. Let us denote $g = (b^{kq^{t-v}} - 1, N)$. Suppose that $q^v \in \mathcal{M}_b(N)$. By Theorem 1 we know that g can not be divisible by other prime different from q and that $\nu_q(N) \leq v$. Let $p \neq q$ be a prime divisor of N . Because p not divides g , we have $|b|_p \nmid kq^{t-v}$ and thus $\nu_q(|b|_p) > t - v \geq 0$ and it is easy to see that $t = \max_{1 \leq i \leq t} \left\{ n - m, \nu_q(|b|_{p_i}) \right\}$. Therefore for all prime divisor p of N we get that $\nu_q(|b|_p) > \max_{1 \leq i \leq t} \left\{ n - m, \nu_q(|b|_{p_i}) \right\} - v$.

Conversely, from the hypothesis the unique prime divisor of N that could be a divisor of g is q and thus Theorem 1 implies that $q^v \in \mathcal{M}_b(N)$. \square

As a special case, if in the above theorem we do $v = 1$ and since for any p prime divisor of N we have $\nu_q(|b|_N) \geq \nu_q(|b|_p)$ and thus we obtain the next corollary.

Corollary 8. *Let N be a positive integer and let q be a prime divisor of $|b|_N$, then $q \in \mathcal{M}_b(N)$ if and only if*

- (1) *If $(N, q) = 1$, then $\nu_q(|b|_p) = \nu_q(|b|_N)$ for all p prime divisor of N .*
- (2) *If $(N, q) > 1$, then q^2 not divides N and $\nu_q(|b|_p) = \nu_q(|b|_N)$ for all p prime divisor of N different from q .*

Note that, from the last theorem, the smallest number N such that $q^v \in \mathcal{M}_b(N)$ has to be a prime P which satisfies $\nu_q(|b|_P) > 0$ and $q^v \mid |b|_P$. We obtain the below corollary, recalling that $|b|_P$ divides $P - 1$.

Corollary 9. *If q is a prime and v is a positive integer, then the smallest integer N such that $q^v \in \mathcal{M}_b(N)$ is a prime congruent with $1 \pmod{q^v}$.*

As a consequence of the above corollary we have a particular case of the Dirichlet's Theorem about primes in arithmetic progressions.

Corollary 10. *If q is a prime and v is a positive integer, there are infinitely many primes which are congruent to 1 modulo q^v .*

Proof. By the last corollary there exists a prime P_1 congruent with 1 modulo q^v and which satisfies $q^v \in \mathcal{M}_b(P_1)$. Take t_1 an integer such that $q^{t_1 v} > P_1$. Once again, we can find a prime P_2 congruent with 1 modulo $q^{t_1 v}$ such that $q^{t_1 v} \in \mathcal{M}_b(P_2)$ and so on. \square

ACKNOWLEDGEMENTS

The authors are members of the research group: Álgebra, Teoría de Números y Aplicaciones, ERM. J.H. Castillo was partially supported by CAPES, CNPq from Brazil and Universidad de Nariño from Colombia. J.M. Velásquez-Soto was partially supported by CONICET from Argentina and Universidad del Valle from Colombia.

REFERENCES

- [And12] Jens Kruse Andersen, *Primes in arithmetic progression records*, retrieved 05 March 2012, URL <http://users.cybercity.dk/~ds1522332/math/aprecords.htm> (2012).
- [CGPVS11] John H. Castillo, Gilberto García-Pulgarín, and Juan Miguel Velásquez-Soto, *Structure of associated sets to Midy's Property*, accepted to publication in *Matemáticas: Enseñanza Universitaria*, arXiv:1110.3308v2 [math.NT] (2011).
- [GPG09] Gilberto García-Pulgarín and Hernán Giraldo, *Characterizations of Midy's property*, *Integers* **9** (2009), A18, 191–197. MR MR2506150
- [GT04] Ben Green and Terence Tao, *The primes contain arbitrarily long arithmetic progressions*, arXiv:math/0404188v6 [math.NT] (2004).

- [Lew07] Joseph Lewittes, *Midy's theorem for periodic decimals*, *Integers* **7** (2007), A2, 11 pp. (electronic). MR MR2282184 (2008c:11004)
- [Nat00] Melvyn B. Nathanson, *Elementary methods in number theory*, Graduate Texts in Mathematics, vol. 195, Springer-Verlag, New York, 2000. MR 1732941 (2001j:11001)

JOHN H. CASTILLO, DEPARTAMENTO DE MATEMÁTICAS Y ESTADÍSTICA, UNIVERSIDAD DE NARIÑO, SAN JUAN DE PASTO-COLOMBIA
E-mail address: `jhcastillo@gmail.com`

GILBERTO GARCÍA-PULGARÍN, UNIVERSIDAD DE ANTIOQUIA, MEDELLÍN-COLOMBIA
E-mail address: `gigarcia@ciencias.udea.edu.co`

JUAN MIGUEL VELÁSQUEZ-SOTO, DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DEL VALLE, CALI-COLOMBIA
E-mail address: `jumiveso@univalle.edu.co`