

Projective symplectic geometry on regular subspaces; Grassmann spaces over symplectic copolar spaces

M. Prażmowska, K. Prażmowski, M. Żynel

November 23, 2018

Abstract

We construct Grassmann spaces associated with the incidence geometry of regular and tangential subspaces of a symplectic copolar space, show that the underlying metric projective space can be recovered in terms of the corresponding adjacencies on so distinguished family of k -subspaces ($2k + 1 \neq$ geometrical dimension of the space), and thus we prove that bijections which preserve the adjacency are determined by automorphisms of the underlying space.

Mathematics Subject Classification (2000): 51A50, 51F20.

Key words: symplectic copolar space, Grassmann space, adjacency.

Introduction

It is a classical result that the projective geometry can be recovered from its associated Grassmann space (cf. [1], [21], [16]) and adjacency preserving bijections of projective k -subspaces ($2k + 1 \neq$ the dimension of the space) are determined by collineations (cf. e.g. [9]). How to transfer these notions to metric projective geometry to obtain reasonable results?

Several distinct ways lead to problems investigated in this paper. Let us consider a metric projective geometry i.e. a projective space \mathfrak{P} equipped with a nondegenerate polarity \varkappa . In a more specific sense, let us consider a vector space \mathbb{V} (which represents \mathfrak{P}) equipped with a nondegenerate bilinear reflexive form ξ (which determines \varkappa and the conjugacy \perp). We also assume that the coordinate field of \mathbb{V} has characteristic $\neq 2$. If $\text{ind}(\xi) > 1$ then the geometry of (\mathfrak{P}, \perp) can be expressed in terms of the associated polar space, whose points are the isotropic (singular) points of \mathfrak{P} and whose lines are the isotropic lines (cf. [22], [3]). This geometry can be also expressed in terms of the adjacency: binary collinearity of points. This result can be extended to isotropic subspaces of higher dimensions, and on isotropic subspaces Grassmann spaces can be defined quite naturally (cf. [17]). Analogous results remain valid for affine polar spaces ("polar spaces" associated with metric affine geometries, cf. [5], [18]). But within (\mathfrak{P}, \perp) isotropic subspaces are "most" degenerate. "Least" degenerate and "opposite" to isotropic are regular (radical-free) subspaces, which are more suitable to develop the geometry, especially when reflections are considered (cf. [20]).

If $\text{ind}(\xi) \neq 0$ and ξ is symmetric then the structure of regular points and regular lines is equivalent to the underlying metric projective space. One can also extend

this result to regular subspaces of higher dimensions defining associated Grassmann spaces of regular subspaces. The underlying metric projective geometry can be expressed in terms of such Grassmannians (cf. [19]). However, if ξ is symplectic (anti-symmetric or skew-symmetric in other words) then no point is regular and the above, so elegant approach fails. Instead, one can consider symplectic copolar space (cf. [11], [15], also called hyperbolic symplectic space, cf. [10]) with the isotropic points and the regular lines. How to extend this approach to higher dimensions?

Secondly, if ξ is symmetric then quite interesting geometry arises when we consider the structure of regular points and so called tangential lines (lines, which contain exactly one nonregular point, cf. [7]). If ξ is symplectic, then such a structure has no sense, but instead, we can consider the structure with regular lines and tangential planes. Here, a tangential subspace is defined as a subspace, whose radical is a point.

All of that suggests that extending all the machinery of adjacency and Grassmann spaces to symplectic copolar spaces one should investigate "best possible": regular subspaces of even (linearly computed) dimensions and tangential subspaces of odd dimensions. And indeed, as we prove in this paper, classical results concerning geometry of Grassmannians remain valid: the underlying metric projective geometry can be expressed in terms of Grassmann spaces of such a family of subspaces, and an analogue of Chow Theorem (cf. [4], [9]) holds i.e. bijections which preserve the adjacency are determined by automorphisms of the underlying metric projective space.

1 Notions, results

Let \mathbb{V} be a vector space with the field of scalars of $\text{char} \neq 2$, let a nondegenerate symplectic form ξ be defined on \mathbb{V} , and let $n = \dim(\mathbb{V})$. Then $n = 2m$, where $m = \text{ind}(\xi)$.

Let $\text{Sub}(\mathbb{V})$ ($\text{Sub}_k(\mathbb{V})$, resp.) stand for all the (all the k -dimensional) subspaces of \mathbb{V} . For any $\mathcal{H} \subset \text{Sub}(\mathbb{V})$ and integer k we write $\mathcal{H}_k = \mathcal{H} \cap \text{Sub}_k(\mathbb{V})$.

The structure

$$\mathfrak{P} := \mathbf{P}_1(\mathbb{V}) := \langle \text{Sub}_1(\mathbb{V}), \text{Sub}_2(\mathbb{V}), \subset \rangle$$

is the projective space over \mathbb{V} . In what follows we shall refer mostly to linear dimension, so a point of \mathfrak{P} has dimension 1, a line has dimension 2 and so on. The orthogonality \perp determined by ξ is defined by the condition

$$U_1 \perp U_2 \iff \xi(U_1, U_2) = 0, \text{ for } U_1, U_2 \in \text{Sub}(\mathbb{V}).$$

For $u, v \in V$ and $U \in \text{Sub}(\mathbb{V})$ we write $u \perp U$ if $\langle u \rangle \perp U$ and $u \perp v$ if $\langle u \rangle \perp \langle v \rangle$ i.e. if $\xi(u, v) = 0$. Set $U^\perp := \{u \in V : U \perp u\}$ and then $\text{Rad}(U) := U \cap U^\perp$. Write $\text{rdim}(U) = \dim(\text{Rad}(U))$.

The structure (\mathfrak{P}, \perp) is called a *symplectic (metric-)projective space*.

Let $U \in \text{Sub}(\mathbb{V})$. The subspace U is *isotropic* (the terms *totally isotropic* and *singular* are also used) if $U \perp U$; if U is isotropic then $\dim(U) \leq m$. The subspace U is *regular* iff $\text{Rad}(U)$ is the zero-subspace of \mathbb{V} . In other words, U is regular if the form $\xi \upharpoonright U$ is nondegenerate. Let \mathbf{Q} stand for the class of isotropic subspaces of \mathbb{V} and let \mathbf{R} stand for the class of regular subspaces of \mathbb{V} . Since ξ is symplectic,

$\mathbf{R}_k \neq \emptyset$ yields $2 \mid k$. In particular, $\mathbf{R}_1 = \emptyset$ and $\mathbf{Q}_1 = \text{Sub}_1(\mathbb{V})$. Moreover, $\text{Sub}_2(\mathbb{V}) = \mathbf{Q}_2 \cup \mathbf{R}_2$; i.e. a line of \mathfrak{P} is either regular or isotropic.

A subspace U of \mathbb{V} will be called *tangential* (cf. [7]) if $\text{Rad}(U)$ is a projective point i.e. if $\text{rdim}(U) = 1$. Let \mathbf{T} be the class of the tangential subspaces of \mathbb{V} . It is evident that $\mathbf{T}_k \neq \emptyset$ yields $2 \nmid k$.

The following evident observation is worth to note

FACT 1.1. *Let $U \in \text{Sub}(\mathbb{V})$. The following conditions are equivalent:*

- (i) $U \in \mathbf{T}$;
- (ii) there are $U_0 \in \mathbf{R}$ and a point $p \subset U_0^\perp$ such that $U = U_0 + p$.

FACT 1.2 ([19, Lemma 1.2, Cor. 1.3]). *Let $Y \in \text{Sub}(\mathbb{V})$. If $Y \supset X \in \mathbf{R}$ ($Y \subset X \in \mathbf{R}$) then $\text{rdim}(Y) \leq \text{codim}_Y(X)$ ($\text{rdim}(Y) \leq \text{codim}_X(Y)$), resp.).*

Consequently, if Y contains a regular hyperplane or Y is a hyperplane of a regular subspace then Y is tangential.

Let us consider the incidence geometry $\Upsilon = ((\mathbf{T-R})_k: k = 0, \dots, n)$, where

$$(\mathbf{T-R})_k = \begin{cases} \mathbf{R}_k & \text{when } 2 \mid k \\ \mathbf{T}_k & \text{when } 2 \nmid k \end{cases}.$$

In a more concise way we can simply write

$$(1) \quad (\mathbf{T-R}) = \{U \in \text{Sub}(\mathbb{V}) : \text{rdim}(U) \leq 1\}.$$

Our incidence geometry Υ is a quasi Curtis-Phan-Tits as introduced in [2, Sec. 3] (cf. also [8], [12], [13]). The family $(\mathbf{T-R})_k$ is the set of all objects of type k in this geometry.

Note that the family $(\mathbf{T-R})$ remains invariant under the map

$$\varkappa: \text{Sub}(\mathbb{V}) \ni U \longmapsto U^\perp.$$

We adopt the following convention.

- Let $H \in \text{Sub}_{k-1}(\mathbb{V})$, $B \in \text{Sub}_{k+1}(\mathbb{V})$.

A *projective top* is a set of the form $\text{T}^\times(B) = \{U \in \text{Sub}_k(\mathbb{V}) : U \subset B\}$ and a *projective star* is a set of the form $\text{S}^\times(H) = \{U \in \text{Sub}_k(\mathbb{V}) : H \subset U\}$. For $H \subset B$ a *projective pencil* is a set of the form $\mathbf{p}^\times(H, B) = \text{T}^\times(B) \cap \text{S}^\times(H)$. The class of projective pencils will be denoted by $\mathcal{P}_k^\times = \mathcal{P}_k^\times(\mathbb{V})$.

- Let $H \in (\mathbf{T-R})_{k-1}$, $B \in (\mathbf{T-R})_{k+1}$.

A $(\mathbf{T-R})$ -*top* is a set of the form $\text{T}(B) = \{U \in (\mathbf{T-R})_k : U \subset B\}$ and a $(\mathbf{T-R})$ -*star* is a set of the form $\text{S}(H) = \{U \in (\mathbf{T-R})_k : H \subset U\}$. Clearly, $\text{T}(B) \subset \text{T}^\times(B)$ and $\text{S}(H) \subset \text{S}^\times(H)$. For $H \subset B$ a $(\mathbf{T-R})$ -*pencil* is a set of the form $\mathbf{p}(H, B) = \text{T}(B) \cap \text{S}(H)$. The class of *nonempty* $(\mathbf{T-R})$ -pencils will be denoted by $\mathcal{P}_k = \mathcal{P}_k((\mathbf{T-R}))$.

From 1.2 we immediately get

FACT 1.3.

- (i) *Let $H \in \mathbf{R}_{k-1}$, $H \subset B \in \mathbf{R}_{k+1}$. Then $\mathbf{p}^\times(H, B) \subset \mathbf{T}$.*

(ii) Let $H \in \text{Sub}_{k-1}(\mathbb{V})$, $H \subset B \in \text{Sub}_{k+1}(\mathbb{V})$, and $\mathbf{p}^\alpha(H, B) \cap \mathbf{R}_k \neq \emptyset$. Then $H, B \in \mathbf{T}$.

In view of 1.3, following a standard way (cf. [19], [6]) one can construct the *Grassmann space* (or the *space of pencils* in other words)

$$\mathbf{P}_k(\mathbf{T-R}) = \langle (\mathbf{T-R})_k, \mathcal{P}_k((\mathbf{T-R})) \rangle$$

for every integer $1 \leq k \leq n-1$. In particular (cf. 1.1), each regular subspace and each tangential subspace is in a pencil, and each pencil, if nonempty, has at least two elements. Therefore, $\mathbf{P}_k(\mathbf{T-R})$ is a partial linear space. The map \varkappa yields an isomorphism of $\mathbf{P}_k(\mathbf{T-R})$ and $\mathbf{P}_{n-k}(\mathbf{T-R})$. Let us also point out that our Grassmann space $\mathbf{P}_k(\mathbf{T-R})$ can be viewed as a specific shadow space of the Curtis-Phan-Tits geometry Υ (cf. [6, Sec. 4]).

In view of 1.3, there is no way to define a space of pencils associated with regular subspaces alone, i.e. with the incidence geometry $(\mathbf{R}_k: 1 \leq k \leq n-1)$. That is why we need the geometry with both regular *and* tangential subspaces in symplectic case.

Clearly, the notion of a tangential subspace makes sense in arbitrary metric projective geometry determined by a reflexive nondegenerate form (cf. [7]). In the case of a symmetric form ξ one can consider incidence geometries with tangential subspaces of arbitrary dimension only, or with tangential *and* isotropic, or with tangential *and* regular; in each case a reasonable incidence geometry arises and sensible Grassmann spaces can be investigated. Many of the lemmas which we prove in the paper remain valid for a symmetric form ξ . However, we do not intend to develop the general theory of Grassmannians of regular and tangential subspaces. In essence, our intention is to show how to enrich the class of regular subspaces of a *symplectic* geometry, as easily as possible, to be able to construct a reasonable Grassmann space over it and obtain analogous results as in the case of symmetric form and regular subspaces alone.

For $U, W \in (\mathbf{T-R})_k$ we write $U \sim W$ when U, W are collinear in $\mathbf{P}_k(\mathbf{T-R})$. The goal of this paper is to prove

THEOREM 1.4. *Let $k \neq n-k$. Then the underlying projective symplectic geometry (\mathfrak{P}, \perp) can be defined in terms of the adjacency \sim on $(\mathbf{T-R})_k$; consequently, (\mathfrak{P}, \perp) can be defined in terms of the geometry of its Grassmann space $\mathbf{P}_k(\mathbf{T-R})$. In particular, when $2 \mid k$, then (\mathfrak{P}, \perp) can be defined in terms of the adjacency \sim on its regular k -subspaces $(\mathbf{T-R})_k = \mathbf{R}_k$.*

For this purpose we first prove

PROPOSITION 1.5. *Let $k \neq n-k$. If $k > 1$ then $\mathbf{P}_{k-1}(\mathbf{T-R})$ can be defined in terms of the binary adjacency relation \sim on $(\mathbf{T-R})_k$ and consequently, it can be defined in terms of $\mathbf{P}_k(\mathbf{T-R})$. If $k < n-1$ then $\mathbf{P}_{k+1}(\mathbf{T-R})$ can be defined in terms of the binary adjacency relation \sim on $(\mathbf{T-R})_k$ so, it can be defined in terms of $\mathbf{P}_k(\mathbf{T-R})$.*

In case $k = 1$, the Grassmann space $\mathbf{P}_1(\mathbf{T-R})$ is isomorphic to the structure $\mathfrak{R} = \langle (\mathbf{T-R})_1, (\mathbf{T-R})_2, \subset \rangle$, which is a *copolar space* embedded in \mathfrak{P} (cf. [11], [8], [15], and [14]); it is simply the line-complement of the corresponding polar space $(\text{Sub}_1(\mathbb{V}), \mathbf{Q}_2, \subset)$. Informally, we can say that the incidence geometry (1) is defined

over subspaces of \mathfrak{R} and write $\mathbf{P}_k(\mathbf{T-R}) = \mathbf{P}_{k-1}(\mathfrak{R})$. This approach, however, makes it hard to characterize tangential and regular subspaces in the language of pure incidence structure \mathfrak{R} .

The validity of Theorem 1.4 in case $k = 1$ is a consequence of elementary properties of polar spaces (cf. (28) in Sec. 2). Therefore, by induction, Theorem 1.4 follows from 1.5. As a corollary to 1.4 we obtain

COROLLARY 1.6 (A variant of Chow Theorem). *Let $k \neq n - k$. The three classes of maps of $(\mathbf{T-R})_k$:*

- the automorphisms of $\mathbf{P}_k(\mathbf{T-R})$,
- the bijections which preserve the adjacency \approx (in both directions),
- the collineations of \mathfrak{P} preserving the conjugacy \perp and acting on $(\mathbf{T-R})_k$.

all coincide.

If $k = n - k$ then a bijection of $(\mathbf{T-R})_k$ which preserves \approx is either determined by an automorphism of (\mathfrak{P}, \perp) or it is a composition of the duality \varkappa and the map determined by an automorphism of (\mathfrak{P}, \perp) .

Frequently (cf. e.g. [17], [18]), dealing with incidence geometry like (1), two more adjacencies are considered: for $U, W \in (\mathbf{T-R})_k$ we write $U \sim_- W$ when $U \cap W \in (\mathbf{T-R})_{k-1}$, and $U \sim^+ W$ when $U \neq W$ and $U, W \subset B$ for some $B \in (\mathbf{T-R})_{k+1}$. Clearly, $U \approx W$ iff $U \sim_- W$ and $U \sim^+ W$. If $2 \mid k$ then \approx , \sim_- , and \sim^+ coincide on $(\mathbf{T-R})_k$. (cf. (12) in Sec. 2). However, if $2 \nmid k$ then the relations \approx, \sim_-, \sim^+ on $(\mathbf{T-R})_k$ are pairwise distinct. To complete the results, we prove also the following.

THEOREM 1.7. *Let $2 \nmid k$. and $1 \leq k \leq n - 1$. For $k \neq 1$, the underlying projective symplectic geometry can be defined in terms of \sim_- on \mathbf{T}_k , and for $k \neq n - 1$ in terms of \sim^+ on \mathbf{T}_k . Consequently, a bijection of \mathbf{T}_k which preserves in both directions the relation \sim_- or preserves in both directions the relation \sim^+ is determined by an automorphism of the underlying projective symplectic space.*

After a quite technical analysis of geometry of Grassmannians $\mathbf{P}_k(\mathbf{T-R})$ in the first two subsections of Sec. 2, we give complete proofs of Proposition 1.5, Theorems 1.4, 1.7, and Corollary 1.6 in Subs. 2.6. Some results, e.g. (4), (5), (13), (27), and (29), seem to be interesting also on their own right.

2 Technical details and proof of Theorems

2.1 Triangles and planes of a symplectic copolar space

In the remainder of this paper we shall need some more information on triangles of symplectic copolar spaces. We gather these in the following list. Most of them are folklore, but some are given with a short proof for completeness sake.

FACT 2.1.

- (i) *Let a_1, a_2 be distinct projective points. Then a_1, a_2 are collinear in \mathfrak{R} iff $a_1 \not\perp a_2$.*

(ii) A projective plane π contains a triangle of \mathfrak{R} iff $\text{rdim}(\pi) = 1$ i.e. iff $\pi \in \mathbf{T}$.

(iii) If $\text{rdim}(\pi) = 1$ then each projective line on π is either isotropic and passes through $p = \text{Rad}(\pi)$, or it is regular and misses p .

Let $\pi \in \mathbf{T}_3$ (i.e. let π be a tangential plane). We write $\pi^\infty = \text{Rad}(\pi)$ and $[\pi] = \pi \setminus \{\pi^\infty\}$.

(iv) Let a plane $\pi \in \mathbf{T}$ contain a triangle Δ with regular sides. Then the set $\overline{\Delta}$ of points on lines of \mathfrak{R} which cross all the sides of Δ coincides with $[\pi]$. This can be read as follows: a triangle spans a plane of \mathfrak{R} , which is a dual affine plane, cf. [8], [10], and [14].

(v) Let planes $\pi_1, \pi_2 \in \mathbf{T}$ have a regular line L in common. There is a tetrahedron with regular edges, whose one edge is L , one face is in π_1 , and other in π_2 .

PROOF. Let $p_i = \text{Rad}(\pi_i)$ for $i = 1, 2$. Then $p_1, p_2 \notin L$, since otherwise (say, $p_1 \in L$), we would get $p_1 \in L \perp p_1$, so $p_1 \in \text{Rad}(L)$. Take any $a_1 \in \pi_1 \setminus (L \cup \{p_1\})$. Suppose that $a_1 \perp (\pi_2 \setminus (L \cup \{p_2\}))$. Then $a_1 \perp \pi_2$ and thus $a_1 \perp L$, which gives $a_1 \perp L + a_1 = \pi_1$ and $a_1 = p_1$. Consequently, there is $a_2 \in \pi_2 \setminus (L \cup \{p_2\})$ with $a_1 \not\perp a_2$ and then a_1, a_2 are on a regular line. One can easily find $a_3, a_4 \in L$ such that $a_1, a_2 \not\perp a_3, a_4$. \square

Write $\pi_1 \wedge \pi_2$ iff there is a tetrahedron as in (v)

(vi) Let planes $\pi_1, \pi_2 \in \mathbf{T}$ have an isotropic line L in common. Let $a_3, a_4 \in L$, $a_3, a_4 \neq \text{Rad}(\pi_1), \text{Rad}(\pi_2)$, and $a_3 \neq a_4$. Then there are $a_i \in \pi_i \setminus L$ such that $a_1 \not\perp a_2$ and $a_1, a_2 \not\perp a_3, a_4$, so a_1, a_2, a_3 and a_1, a_2, a_4 span two planes $\pi'_1, \pi'_2 \in \mathbf{T}$ with common regular line $M = a_1 + a_2$. By the above, $\pi'_1 \wedge \pi'_2$.

PROOF. From assumption, $L \perp L$. Write $p_i = \text{Rad}(\pi_i)$. Then $p_1, p_2 \in L$. Let $a_1 \in \pi_1 \setminus L$ be arbitrary. Then $p_1 \perp a_1$; if $a_1 \perp a_3$ or $a_1 \perp a_4$ then $a_1 \perp L$ and thus $a_1 \perp \pi_1$. As above, we find $a_2 \in \pi_2 \setminus L$ with $a_1 \not\perp a_2$ and we are through. \square

Write $\pi_1 \wedge \pi_2$ iff there are planes π'_1, π'_2 as in (vi).

In view of (v) and (vi), we obtain

(vii) Let π_1, π_2 be two tangential planes. If $|\pi_1 \cap \pi_2| \geq 2$ then $\pi_1 \wedge \pi_2$ or $\pi_1 \wedge \pi_2$.

(viii) Any two planes in \mathbf{T} can be joined by a sequence $(\pi_i: i = 0, \dots, t)$ of tangential planes such that $|\pi_{i-1} \cap \pi_i| \geq 2$ for $i = 1, \dots, t$.

PROOF. By (i), any two points a', a'' can be joined in \mathfrak{R} by a path (sometimes also called a polygonal path, which by the way is of length ≤ 2 here). Let a_0, \dots, a_t be such a path in \mathfrak{R} that joins a', a'' and let L_i be the line through a_{i-1}, a_i for $i = 1, \dots, t$. By (iv), any two consecutive sides L_i, L_{i+1} lie in a plane $\pi_i \in \mathbf{T}$, $i = 1, \dots, t-1$. Let $\pi', \pi'' \in \mathbf{T}_2$; take regular lines L' in π' and L'' in π'' and points a' on L' , a'' on L'' . Taking $\pi_0 = L' + L_1$ if L' is not on π_1 and $\pi_0 = \pi_1$ otherwise, and similarly, $\pi_t = L_t + L''$ or $\pi_t = \pi_{t-1}$ we obtain a desired sequence of planes. \square

(ix) For each point p of \mathfrak{X} there is a tangential plane π such that $p \in [\pi]$.

PROOF. First, we take any point q such that $q \not\sim p$ and let L be a line through p, q . Next, let π be a plane that contains L . Then L is regular, so π is tangential. By (iii), L misses $\text{Rad}(\pi)$ and thus $p \neq \text{Rad}(\pi)$. \square

Let us write $\bar{\wedge}$ for the transitive closure of the relation $\wedge \cup \mathfrak{A}$. By (vii), (viii), and (ix) we obtain

(x) Let Δ be any triangle of \mathfrak{X} . Then the set of points of \mathfrak{X} is the union

$$(2) \quad \tilde{\Delta} = \bigcup \left\{ \overline{\Delta'} : \overline{\Delta'} \bar{\wedge} \overline{\Delta}, \Delta' \text{ is a triangle} \right\}.$$

Let a_1, a_2 be points of \mathfrak{X} . Write $a_1 \sim a_2$ when a_1, a_2 are collinear in \mathfrak{X} . In view of (i), the orthogonality of points is definable in \mathfrak{X} : $a_1 \perp a_2$ iff $a_1 = a_2$ or $a_1 \not\sim a_2$.

(xi) Let a_1, a_2, b_1, b_2 be four points of \mathfrak{X} such that $a_1 \sim a_2$, $b_1 \not\sim b_2$, and $a_1, a_2 \sim b_1, b_2$ (cf. (vi)). Then there are points c_1, c_2 such that $c_1 \not\sim c_2$ and $a_1, a_2, b_1, b_2 \sim c_1, c_2$.

PROOF. Let L be the line through a_1, a_2 . Suppose, first, that b_1, b_2 are on a plane π through L . Then π is tangential; let $p = \text{Rad}(\pi)$, so the line M through b_1, b_2 is isotropic and it passes through p . Clearly, $b_1, b_2 \neq p$, as the lines $\overline{b_1, a_i}$ and $\overline{b_2, a_i}$ are not isotropic. We take any line $M_0 \neq M$ through p that misses a_1, a_2 . Any two points c_1, c_2 on M_0 that are distinct from p and not lie on L satisfy our requirements.

If there is no plane through L which contains b_1, b_2 then the lines L and M are skew and they span a projective 3-space Γ (linearly, $\dim(\Gamma) = 4$); as Γ contains L , $\text{rdim}(\Gamma) \leq 2$, and as $\Gamma/\text{Rad}(\Gamma)$ is nondegenerate $2 \mid (4 - \text{rdim}(\Gamma))$. Consequently, there are two cases to consider. Firstly, if $\text{Rad}(\Gamma)$ is a line K then isotropic lines in Γ are exactly the projective lines which cross K ; in particular, M crosses K in a point p . Let $\pi_i = K + a_i$ for $i = 1, 2$; then M is neither contained in π_1 nor in π_2 . Let π be a plane that contains M and does not contain K ; then $\pi \cap K = p$. Let M_0 be other line in π that passes through p and is not contained in π_1, π_2 . For each $i = 1, 2$ there is at most one point on M_0 which is not collinear with a_i (as otherwise, M_0 is contained in π_i). Thus one can find on M_0 two points c_1, c_2 distinct from p and collinear with a_1, a_2 . From construction, $b_1, b_2 \sim c_1, c_2$ and $b_1 \not\sim c_2$. Secondly, let Γ be regular; then $\text{ind}(\xi \upharpoonright \Gamma) \leq 2$, so each plane in Γ is tangential. Let π_1 be the plane spanned by a_1, a_2, b_2 , M be a line through $p = \text{Rad}(\pi_1)$ contained in π_1 and missing a_1, a_2, b_2 , π_2 be the plane through M, b_1 , and $q = \text{Rad}(\pi_2)$. Since $M \subset \pi_2$ is isotropic, by (iii), $q \subset M$. Take c_1, c_2 on M distinct from p, q ; then $c_1 \not\sim c_2$. By (iii), $c_1, c_2 \sim a_1, a_2, b_1, b_2$. \square

(xii) Let a_1, a_2, a_3 be a triangle in \mathfrak{X} . There are points b_1, b_2, b_3 such that $a_1, a_2, a_3 \sim b_1, b_2, b_3$, $b_1 \not\sim b_2, b_3$, and $b_2 \sim b_3$.

PROOF. Let π_1 be the plane spanned by a_1, a_2, a_3 . Then $\pi_1 \in \mathbf{T}_3$ and one can find $\Gamma \in \mathbf{R}_4$ with $\pi \subset \Gamma$. Let us restrict to Γ considered as a 3-dimensional symplectic projective space. As above, each plane contained in Γ is tangential. One can complete in Γ given triangle to a tetrahedron a_1, a_2, a_3, b_3 . Let us take on π_1 a line L through $p = \text{Rad}(\pi_1)$ which misses a_1, a_2, a_3 ; let π_2 be the plane through L, b_3 , and let $q = \text{Rad}(\pi_2)$. π_2 contains an isotropic line L , so L contains q . Suppose that $p = q$; then $p \perp \pi_1, \pi_2$ gives $p \perp \Gamma$, a contradiction. Set $b_1 := q$. Then $b_1 \sim a_1, a_2, a_3$. Let b_2 be a point on L distinct from p, q . By (iii), considering π_1 we get $b_2 \sim a_1, a_2, a_3, b_1 \not\sim b_2$, and considering π_2 we get $b_2 \sim b_3, b_3 \not\sim b_1$. \square

Now we are going to reconstruct the Grassmann space $\mathbf{P}_k(\mathbf{T-R})$ in terms of binary adjacency \approx . This involves the family of maximal cliques of \approx , i.e. stars and tops, and we shall show that they are definable and distinguishable in the first place. Our reasoning depends on whether k is even or odd.

2.2 Case 2 $\not\sim k$

Let $H \in \mathbf{R}_{k-1}, H \subset B \in \mathbf{R}_{k+1}$.

Let $U \in \mathbf{S}^\infty(H)$ i.e. let $H \subset U \in \text{Sub}_k(\mathbb{V})$. From 1.2, $\text{rdim}(U) = 1$. Consequently,

$$\mathbf{S}(H) = \mathbf{S}^\infty(H).$$

Let $q = \text{Rad}(U)$, then $q \perp U$; in particular, $q \perp H$. Since $H \in \mathbf{R}$, $q \notin H$ and thus $U = H \oplus q$ with $q \in H^\perp$. Consequently, an element of $\mathbf{S}(H)$ is any subspace U of the form as above i.e.

$$\mathbf{S}(H) = \left\{ H + q : q \text{ is a point, } q \subset H^\perp \right\}.$$

Thus the elements of $\mathbf{S}(H)$ can be identified with the elements of $\text{Sub}_1(H^\perp)$ i.e. with the points of the nondegenerate metric projective space defined over H^\perp .

Analogous remarks concern the geometry of $\mathbf{T}(B) = \mathbf{T}^\infty(B)$. In particular, we can write

$$\mathbf{T}(B) = \left\{ B \cap q^\perp : q \text{ is a point on } B \right\}.$$

From the above we get:

(3) if $H \subset B$ then $\mathbf{S}(H) \cap \mathbf{T}(B) = \mathbf{p}^\infty(H, B)$, so it contains at least two elements.

One can note that

$$\mathbf{p}(H, B) = \{ H \oplus \langle u \rangle : u \in P, u \neq \theta \}, \text{ where } P = H^\perp \cap B, \text{ and } P \in \mathbf{R}_2.$$

The form ξ restricted to H^\perp is symplectic, so the points of the metric projective space $\mathbf{S}(H)$ are simply the points of the corresponding projective space $\mathbf{P}_1(H^\perp)$; the distinction lies in the adopted family of lines. These lines correspond to suitable pencils i.e. to $B \in \mathbf{R}_{k+1}$ with $H \subset B$. We see that the B above correspond to regular 2-subspaces of H^\perp and thus

(4) the geometry of the restriction of $\mathbf{P}_k(\mathbf{T-R})$ to $\mathbf{S}(H)$ is an $(n-k)$ -dimensional symplectic copolar space.

With analogous reasoning we obtain that

(5) the restriction of $\mathbf{P}_k(\mathbf{T-R})$ to $\mathbf{T}(B)$ is a k -dimensional symplectic copolar space.

Let Δ be a triangle in $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$. Then, by common projective geometry, its vertices are either in a top $\mathbf{T}(B) =: \mathcal{X}$ or in a star $\mathbf{S}(H) =: \mathcal{X}$. Clearly, $\overline{\Delta}$, as defined in 2.1(iv) with \mathfrak{R} replaced by $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$, is a plane in \mathcal{X} . Note that if Δ' is a different triangle in $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$ and $\overline{\Delta'} \wedge \overline{\Delta}$ or $\overline{\Delta'} \wedge \overline{\Delta}$ (cf. 2.1), then Δ' lies in \mathcal{X} as well. From the above, (4), (5), and 2.1(x), we get that

$$\widetilde{\Delta} = \mathcal{X},$$

$\widetilde{\Delta}$ being the set defined in $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$ by formula (2). Finally, we get that

- (6) *the family of stars and tops is definable in terms of geometry of $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$; by (4) and (5), if $k \neq n - k$, stars and tops are intrinsically distinguishable.*

Let U_1, U_2 be distinct points of $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$ such that $U_1 \approx U_2$, and let $U_1, U_2 \in g = \mathbf{p}(H, B) \in \mathcal{P}_k(\mathbf{T}\text{-}\mathbf{R})$. Next, let W_1, W_2 satisfy

$$W_1, W_2 \approx U_1, U_2 \quad \text{and} \quad W_1 \not\approx W_2.$$

Consider the set

$$[U_1, U_2, W_1, W_2]_{\approx} := \{U : U \approx U_1, U_2, W_1, W_2\}.$$

From common projective geometry, as \approx -neighbor subspaces are adjacent in the projective Grassmannian over \mathfrak{F} , there are three possibilities to consider

- $W_1, W_2 \in \mathbf{S}(H)$. By 2.1(xi) and (4), there are $U, W \in [U_1, U_2, W_1, W_2]_{\approx}$ such that $U \not\approx W$.
- $W_1, W_2 \in \mathbf{T}(B)$. Analogously, there are $U, W \in [U_1, U_2, W_1, W_2]_{\approx}$ such that $U \not\approx W$.
- $W_1 \in \mathbf{T}(B) \setminus \mathbf{S}(H)$ and $W_2 \in \mathbf{S}(H) \setminus \mathbf{T}(B)$ or $W_2 \in \mathbf{T}(B) \setminus \mathbf{S}(H)$ and $W_1 \in \mathbf{S}(H) \setminus \mathbf{T}(B)$. In that case $[U_1, U_2, W_1, W_2]_{\approx} \subset \mathbf{p}(H, B)$ and thus $U \approx W$ for all $U, W \in [U_1, U_2, W_1, W_2]_{\approx}$.

Let $U_3 \in g$ be arbitrary; clearly, one can find $W_1 \in \mathbf{T}(B) \setminus \mathbf{S}(H)$, $W_2 \in \mathbf{S}(H) \setminus \mathbf{T}(B)$ such that $U_1, U_2, U_3 \approx W_1, W_2$. Write \mathbf{L} for the ternary collinearity relation of $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$. In view of the above analysis, the formula

$$(7) \quad \mathbf{L}(U_1, U_2, U_3) \iff U_1 \approx U_2 \wedge (\exists W_1, W_2) \left[W_1, W_2 \approx U_1, U_2 \wedge W_1 \not\approx W_2 \wedge \right. \\ \left. (\forall U, W) [U_1, U_2, W_1, W_2 \approx U, W \implies U \approx W] \wedge U_3 \approx U_1, U_2, W_1, W_2 \right]$$

defines the relation \mathbf{L} in terms of the adjacency \approx for distinct U_1, U_2 . Finally, we conclude that

- (8) *$\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$ is definable in terms of the adjacency \approx .*

2.3 Case 2 | k

Let $H \in \mathbf{T}_{k-1}$, $B \in \mathbf{T}_{k+1}$. Set $p = \text{Rad}(H)$ and $q = \text{Rad}(B)$.

Note that $U \in \text{Sub}_k(B)$ is regular iff U is a linear complement of q in B i.e. iff $B = q \oplus U$. This can be written as

$$(9) \quad \mathbf{T}(B) = \{U \in \mathbf{R}_k : U \subset B\} = \{U \in \text{Sub}_k(B) : q \not\subset U\}.$$

It is easy to note that

$$(10) \quad \mathbf{S}(H) = \{U \in \mathbf{R}_k : H \subset U\} = \{H + \langle u \rangle : u \notin p^\perp\}.$$

Indeed, let $U = H \oplus \langle u \rangle$ with $u \neq \theta$. Suppose that $p \perp u$; then $p \perp U$ and thus $U \notin \mathbf{R}$. Conversely, let $w \in \text{Rad}(U)$, $w \neq \theta$. Then $w \in U$, $w \perp U$, so $w \perp p$. If $w \notin H$ then $p \perp H + \langle w \rangle = U$ and thus $p \perp u$. If $w \in H$ then $w \perp U \supset H$ gives $w = p$; we get $p \perp U$ and thus $p \perp u$.

Assume that $H \subset B$. There are two cases to consider

$q \in H$: Then $q \perp B \supset H$ gives $p = q$. For arbitrary $U \in \mathbf{p}^\times(H, B)$ we have then $p \in \text{Rad}(U)$ and thus $\mathbf{p}(H, B) = \emptyset$.

$q \notin H$: Then $p \neq q$. Set $L := p + q$ and $U_0 := H + q = H + L$. Note that $L \perp L$, so $L \in \mathbf{Q}_2$; moreover, $L \perp H$. From this we obtain $\text{Rad}(U_0) = L$ (in particular, $\text{rdim}(U_0) = 2$). We have $\mathbf{p}(H, B) = \mathbf{p}^\times(H, B) \setminus \{U_0\}$. Indeed, let $U \in \mathbf{p}^\times(H, B)$ and $U \neq U_0$. Then $q \notin U$ and, by (9), $U \in \mathbf{T}(B)$ and thus $U \in \mathbf{p}(H, B)$.

From the above we have the following:

(11) *if $H \subset B$ then $\mathbf{S}(H) \cap \mathbf{T}(B)$ is either empty or it contains at least two elements.*

Let $U_1, U_2 \in \mathbf{R}_k$. Assume that $U_1 \cap U_2 \in \text{Sub}_{k-1}(\mathbb{V})$. Then $U_1 + U_2 \in \text{Sub}_{k+1}(\mathbb{V})$; in view of 1.2, $U_1 \cap U_2, U_1 + U_2 \in \mathbf{T}$ and thus $U_1, U_2 \in \mathbf{p}(U_1 \cap U_2, U_1 + U_2)$. Therefore,

(12) *the binary collinearity in $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$ coincides with the projective adjacency on \mathbf{R}_k .*

Next, note that

(13) *the geometry of restriction of $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$ to a top and to a star is a k -dimensional and a $(n - k)$ -dimensional resp., affine geometry.*

Indeed,

– The family $\mathcal{X} := \mathbf{S}^\times(H)$ has the natural structure of a projective space. The set $\mathbf{S}(H)$ is obtained by removing the segment $[H, p^\perp]_k = \{U \in \wp_k(\mathbb{V}) : H \subset U \subset p^\perp\}$ from \mathcal{X} (cf. (10)). Computing dimensions we see that $[H, p^\perp]_k$ is a hyperplane in \mathcal{X} , which justifies our claim. The induced affine geometry has dimension $n - k$.

– By (9), $\mathbf{T}(B)$ consists of the hyperplanes in the projective space $\mathbf{T}^\times(B)$ which omit a point q ; this procedure results in a k -dimensional affine space.

Let us note a straightforward consequence of (13) that

(14) *stars and tops are distinguishable in terms of \approx provided $k \neq n - k$.*

The sets $\mathbf{S}(H)$ and $\mathbf{T}(B)$ both are strong subspaces of $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$; in particular, they are cliques of the binary collinearity relation \approx . Moreover, they are exactly the maximal cliques of \approx . This yields that

(15) *the family of stars and tops is definable in terms of the adjacency \approx on $(\mathbf{T}\text{-}\mathbf{R})_k$.*

An intersection of two maximal \approx -cliques has at least two elements iff one of these cliques is a star and another one is a top, and then their intersection is a pencil i.e. a line of $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$. This justifies that

(16) *the structure $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$ is definable in terms of the adjacency \approx .*

2.4 Adjacencies \sim_- and \sim^+ on \mathbf{T}

As we already stated, if $2 \mid k$, then all the three adjacency relations \sim , \sim_- , and \sim^+ coincide on $(\mathbf{T}\text{-}\mathbf{R})_k = \mathbf{R}_k$. In case k is odd we need more elaboration to get the Grassmann space $\mathbf{P}_k(\mathbf{T}\text{-}\mathbf{R})$ defined in terms of \sim_- , as well as in terms of \sim^+ .

Let $2 \nmid k$ and $1 < k < n - 1$. We begin our analysis with \sim_- on $(\mathbf{T}\text{-}\mathbf{R})_k = \mathbf{T}_k$. Note that if $U \sim_- W$ then U, W are adjacent in the projective Grassmannian over \mathfrak{P} .

Let U_1, U_2, U_3 be a \sim_- -clique i.e. let them be pairwise distinct and $U_i \sim_- U_j$ for distinct $i, j = 1, 2, 3$. Consider the set

$$[U_1, U_2, U_3]_{\sim_-} = \{U : U \sim_- U_1, U_2, U_3\}.$$

By known properties of projective Grassmannians the following possibilities arise.

- (i) U_1, U_2, U_3 are in a projective pencil $g = \mathbf{p}^\infty(H, B)$; then $H \in \mathbf{R}_{k-1}$.
- (ii) (i) fails, but $U_1, U_2, U_3 \in \mathbf{S}^\infty(H)$ for some H ; then $H \in \mathbf{R}_{k-1}$. We say that U_1, U_2, U_3 yield a \mathcal{S} -triangle.
- (iii) (i) fails, but $U_1, U_2, U_3 \in \mathbf{T}^\infty(B)$ for some B . Since B contains a regular subspace $U_1 \cap U_2$ by 1.2 we get $\text{rdim}(B) \leq 2$ and thus two cases are possible:
 - (a) $B \in \mathbf{R}_{k+1}$, we say that U_1, U_2, U_3 yield a \mathcal{T} -triangle; or
 - (b) $\text{Rad}(B)$ is a line L , we say that U_1, U_2, U_3 yield a \mathcal{T}^* -triangle.

In case (ii) the set $[U_1, U_2, U_3]_{\sim_-}$ is a \sim_- -clique. Indeed, if $U, W \in [U_1, U_2, U_3]_{\sim_-}$ are distinct then $U, W \in \mathbf{S}(H)$.

In case (iii) the set $[U_1, U_2, U_3]_{\sim_-}$ is not a \sim_- -clique. Consider (b) first. A $(k-1)$ -subspace H of B is regular iff H misses L . On the other hand, a k -subspace U is tangential iff it crosses L in a point (which turns out to be its radical). Write $p_i = L \cap U_i$, then p_1, p_2, p_3 are pairwise distinct, as the intersections $U_i \cap U_j$ are regular. Take any point p on L distinct from p_1, p_2, p_3 and two W_1, W_2 through p which do not contain L ; then $W_1, W_2 \sim_- U_1, U_2, U_3$ (the intersections $W_1 \cap U_i, W_2 \cap U_i$ miss L) but $W_1 \not\sim_- W_2$. Note that if $W_1, W_3 \in [U_1, U_2, U_3]_{\sim_-}$ and $W_1 \not\sim_- W_3$ then $p \in W_3$ and thus $W_2 \not\sim_- W_3$. Hence the relation $\not\sim_-$ is transitive on $[U_1, U_2, U_3]_{\sim_-}$.

Let us pass to case (a). Then U_1, U_2, U_3 can be considered as points of a suitable symplectic copolar space $\mathbf{T}(B) = \mathbf{T}^\infty(B)$ and we can use known properties of symplectic projective geometry. So, let π be the plane spanned in $\mathbf{T}(B)$ by U_1, U_2, U_3 . To justify that $[U_1, U_2, U_3]_{\sim_-}$ is not a \sim_- -clique it suffices to take on $\mathbf{T}(B)$ any line through the radical of π and missing U_1, U_2, U_3 , and distinct points U, W on this line. On the other hand by 2.1(xii), one can find $W_1, W_2, W_3 \in \mathbf{T}(B)$ such that $U_1, U_2, U_3 \sim_- W_1, W_2, W_3$, $W_2 \sim_- W_3$, and $W_1 \not\sim_- W_2, W_3$. This justifies that the relation $\not\sim_-$ is not transitive on $[U_1, U_2, U_3]_{\sim_-}$.

In case (i), clearly, the set $[U_1, U_2, U_3]_{\sim_-}$ is not a \sim_- -clique. To see this it suffices to find $U \in \mathbf{S}(H) \setminus g$ and $W \in \mathbf{T}^\infty(B) \setminus g$ with $U_1, U_2, U_3 \sim_- U, W$. Moreover, from properties of the copolar space $\mathbf{S}(H)$, one can find in $\mathbf{S}(H)$ a \sim_- -clique W_1, W_2, W_3 which does not fall into case (i) such that $U_1, U_2, U_3 \in [W_1, W_2, W_3]_{\sim_-}$. Moreover, if $B \in \mathbf{R}_{k+1}$ then an analogous triple can be found in $\mathbf{T}(B)$.

In view of the above analysis, a \sim_- -clique U_1, U_2, U_3 forms a \mathcal{S} -triangle iff $[U_1, U_2, U_3]_{\sim_-}$ is a \sim_- -clique, so

(17) *the class of \mathcal{S} -triangles can be distinguished in terms of \sim_- .*

Now, \mathcal{T} -triangles/ \mathcal{T}^* -triangles can be characterized as \sim_- -cliques U_1, U_2, U_3 which are, firstly, not contained in any $[W_1, W_2, W_3]_{\sim_-}$ for a \mathcal{S} -triangle W_1, W_2, W_3 and, secondly, such that $\mathcal{X} = [U_1, U_2, U_3]_{\sim_-}$ is not a \sim_- -clique and $\not\sim_-$ is not transitive/is transitive on \mathcal{X} . Consequently,

(18) *the class of \mathcal{T} -triangles and the class of \mathcal{T}^* -triangles can be distinguished in terms of \sim_- .*

Let \mathbf{L} be the ternary collinearity relation of $\mathbf{P}_k(\mathbf{T-R})$. Similarly as in (7) we can write

$$(19) \quad \mathbf{L}(U_1, U_2, U_3) \iff (\exists U'_1, U'_2, U'_3, U''_1, U''_2, U''_3) [U'_1, U'_2, U'_3 \text{ is a } \mathcal{S}\text{-triangle} \wedge U''_1, U''_2, U''_3 \text{ is a } \mathcal{T}\text{-triangle} \wedge U_1, U_2, U_3 \in [U'_1, U'_2, U'_3]_{\sim_-} \cap [U''_1, U''_2, U''_3]_{\sim_-}].$$

By (17), (18), and (19) we obtain that

(20) *the structure $\mathbf{P}_k(\mathbf{T-R})$ can be defined in terms of the adjacency \sim_- .*

Clearly, \varkappa maps the relation \sim_- on $(\mathbf{T-R})_k$ onto \sim^+ on $(\mathbf{T-R})_{n-k}$. Thus, (20) with k replaced by $n - k$ yields that

(21) *the structure $\mathbf{P}_k(\mathbf{T-R})$ can be defined in terms of the adjacency \sim^+ .*

Finally, (20) and (21) enable us to conclude with the following.

(22) *The structure $\mathbf{P}_k(\mathbf{T-R})$ can be defined in terms of both \sim_- and \sim^+ on \mathbf{T}_k . Consequently, \approx can be defined in terms of \sim_- and in terms of \sim^+ .*

2.5 Connectedness

We use standard methods to show that automorphisms of binary adjacency preserve, or exchange, the two families of its maximal cliques. These methods rely on the fact that the adjacency in question is connected.

For a relation ρ on $(\mathbf{T-R})_k$ we say that $U, W \in (\mathbf{T-R})_k$ are ρ -connected if there is a sequence $U = U_0, \dots, U_t = W$ such that $U_{i-1} \rho U_i$ or $U_{i-1} = U_i$ for $i = 1, \dots, t$.

By 2.1(i) we easily get that

(23) *the binary collinearity of points of a symplectic copolar space is connected.*

Let $k \leq n - 1$, $H \in (\mathbf{T-R})_{k-1}$, and $U, W \in S(H)$. If $2 \mid k$ then by (12), $U \approx W$. If $2 \nmid k$ then, by (4) and (23), the points U, W of $S(H)$ can be joined in $S(H)$ by a polygonal path. This gives the following.

(24) *Let $U \sim_- W$. Then U, W are \sim -connected.*

In a consequence, we get the following.

(25) *Let $U, W \in (\mathbf{T-R})_k$ and $U \sim_- W$. Then there are \sim_- -connected $B', B'' \in (\mathbf{T-R})_{k+1}$ such that $U \subset B'$ and $W \subset B''$.*

Now, it is easy to prove that

(26) *the conclusions of (24) and of (25) hold for any \sim_- -connected $U, W \in (\mathbf{T-R})_k$.*

Let $U, W \in (\mathbf{T-R})_k$. Consider two flags $U_1 \subset \dots \subset U_k = U$ and $W_1 \subset \dots \subset W_k = W$ such that $U_i, W_i \in (\mathbf{T-R})_i$. Note that if $U_i \subset B', W_i \subset B''$ and $B', B'' \in (\mathbf{T-R})_{i+1}$ then $U_{i+1} \sim_- B'$ and $W_{i+1} \sim_- B''$, so if B', B'' are \sim_- -connected then U_{i+1}, W_{i+1} are \sim_- -connected as well. Starting from (23) and applying, consecutively, (26) and the observation above to $U_i, W_i, i = 1, \dots, k$, by induction, we get that U, W are \sim -connected. This proves that

(27) *the relation \sim on $(\mathbf{T-R})_k$ is connected.*

2.6 Proofs of the results

Now we are able to complete the proofs of our theorems from Sec. 1 by gathering together the facts proved above. The reasoning is more or less typical for Chow type theorems and its crucial step consists in proving that the adjacency structure $\langle (\mathbf{T-R})_{k-1}, \sim \rangle$ (as well as $\langle (\mathbf{T-R})_{k+1}, \sim \rangle$) is definable in $\langle (\mathbf{T-R})_k, \sim \rangle$.

Let a_1, a_2 be points of \mathfrak{A} such that $a_1 \not\sim a_2$. By known properties of symplectic polar spaces and 2.1(i), the set $\{p: (\forall q) [a_1, a_2 \not\sim q \implies q \not\sim p]\}$ is the projective line through a_1, a_2 . Thus the class of lines of \mathfrak{P} is definable in \mathfrak{A} . Finally,

(28) *the metric projective geometry (\mathfrak{P}, \perp) is definable in \mathfrak{A} .*

Fix k with $1 < k < n - 1$. Recall that, due to (8) and (16),

(29) *the two structures $\mathfrak{A}_k := \mathbf{P}_k(\mathbf{T-R})$ and $\mathfrak{B}_k := \langle (\mathbf{T-R})_k, \sim \rangle$ are mutually definable and consequently $\text{Aut}(\mathfrak{A}_k) = \text{Aut}(\mathfrak{B}_k)$.*

Let \mathcal{S}_k be the family of all the stars and \mathcal{T}_k be the family of all the tops in \mathfrak{A}_k . By (6), (8), and (15), in both cases $2 \mid k$ and $2 \nmid k$ the class $\mathcal{S}_k \cup \mathcal{T}_k$ is definable in \mathfrak{B}_k and thus it remains invariant under automorphisms of \mathfrak{B}_k .

LEMMA 2.2. *Assume that stars and tops in $\mathbf{P}_k(\mathbf{T-R})$ are distinguishable.*

(i) *The families $(\mathbf{T-R})_{k-1}$ and $(\mathbf{T-R})_{k+1}$ are definable in \mathfrak{B}_k .*

(ii) *If $k > 1$ then $\mathbf{P}_{k-1}(\mathbf{T-R})$ can be defined in terms of the binary adjacency relation \approx on $(\mathbf{T-R})_k$ and consequently, it can be defined in terms of $\mathbf{P}_k(\mathbf{T-R})$. If $k < n - 1$ then $\mathbf{P}_{k+1}(\mathbf{T-R})$ can be defined in terms of the binary adjacency relation \approx on $(\mathbf{T-R})_k$ so, it can be defined in terms of $\mathbf{P}_k(\mathbf{T-R})$.*

(iii) *Each automorphism F of \mathfrak{A}_k determines an automorphism F^+ of \mathfrak{A}_{k+1} and an automorphism F^- of \mathfrak{A}_{k-1} such that $U \in \text{S}(H)$ iff $F(U) \in \text{S}(F^-(H))$ and $U \in \text{T}(B)$ iff $F(U) \in \text{T}(F^+(B))$ for all $H \in (\mathbf{T-R})_{k-1}$, $U \in (\mathbf{T-R})_k$, and $B \in (\mathbf{T-R})_{k+1}$.*

PROOF. (i) It is enough to see that we can identify $H \in (\mathbf{T-R})_{k-1}$ with $\text{S}(H)$ and $B \in (\mathbf{T-R})_{k+1}$ with $\text{T}(B)$.

(ii) Let $H_1, H_2 \in (\mathbf{T-R})_{k-1}$, and $B_1, B_2 \in (\mathbf{T-R})_{k+1}$. Clearly, $H_1 \sim^+ H_2$ iff $\text{S}(H_1) \cap \text{S}(H_2) \neq \emptyset$, and dually $B_1 \sim_- B_2$ iff $\text{T}(B_1) \cap \text{T}(B_2) \neq \emptyset$. By (i) the maps $\text{S}(\cdot)$ and $\text{T}(\cdot)$ are definable, so up to them, the relations: \sim^+ on $(\mathbf{T-R})_{k-1}$ and \sim_- on $(\mathbf{T-R})_{k+1}$ are definable in \mathfrak{A}_k . This together with (29) when $2 \mid (k - 1)$, and together with (22) when $2 \nmid (k - 1)$, proves our statement.

(iii) By (i) we have bijections F^+ of $(\mathbf{T-R})_{k+1}$ and F^- of $(\mathbf{T-R})_{k-1}$ such that $\text{T}(F^+(B)) = F(\text{T}(B))$ and $\text{S}(F^-(H)) = F(\text{S}(H))$. Now by (ii) these maps are automorphisms as required. \square

PROOF OF PROPOSITION 1.5. The proposition follows directly from Lemma 2.2(ii), (6), and (14). \square

PROOF OF THEOREM 1.4. Assume that $k \neq n - k$. Then by (6), (14) stars and tops are definable and distinguishable in \mathfrak{R}_k . Hence, we can define \mathfrak{R}_{k-1} in \mathfrak{R}_k by Lemma 2.2(ii). A top of \mathfrak{R}_{k-1} has form $\{H \in (\mathbf{T-R})_{k-1} : U \in S(H)\}$ for a point U of \mathfrak{R}_k , so these tops are determined by the points of \mathfrak{R}_k . Analogously, the stars of \mathfrak{R}_{k+1} are determined by the points of \mathfrak{R}_k . Therefore, the stars and the tops of \mathfrak{R}_{k-1} (of \mathfrak{R}_{k+1} , resp.) can be distinguished. Now, from Lemma 2.2(ii) by induction on k we infer that \mathfrak{R}_1 , which is \mathfrak{R} up to an isomorphism, can be defined in \mathfrak{R}_k . In view of (28) we are through. \square

PROOF OF COROLLARY 1.6. Let $F \in \text{Aut}(\mathfrak{B}_k)$. In view of (29) it suffices to show that F is induced by a collineation of \mathfrak{P} preserving \perp and acting on $(\mathbf{T-R})_k$.

In case $k \neq n - k$ the proof runs by induction on k via Lemma 2.2(iii). So, assume that $k = m$. Note that if $\mathcal{X}', \mathcal{X}'' \in \mathcal{S}_k$, $|\mathcal{X}' \cap \mathcal{X}''| = 1$, and $F(\mathcal{X}') \in \mathcal{S}_k$ then $F(\mathcal{X}'') \in \mathcal{S}_k$ as well. Indeed, if there were $F(\mathcal{X}'') \in \mathcal{T}_k$ then $|F(\mathcal{X}') \cap F(\mathcal{X}'')| = 1$ would contradict (3) and (11). Suppose that $F(\mathcal{X}) \in \mathcal{S}_k$ for some $\mathcal{X} = S(H) \in \mathcal{S}_k$. By (27), for each $H' \in (\mathbf{T-R})_{k-1}$ there is a sequence $H = H_0 \sim^+ \dots \sim^+ H_t = H'$ and then $F(S(H')) \in \mathcal{S}_k$, so F preserves \mathcal{S}_k . As in the proof of Lemma 2.2(iii), we consider F^+ and F^- to justify that F is determined by a collineation of \mathfrak{R} . If $F(\mathcal{X}_0) \notin \mathcal{S}_k$ for some $\mathcal{X}_0 \in \mathcal{S}_k$ then $F(\mathcal{X}) \notin \mathcal{S}_k$ for all $\mathcal{X} \in \mathcal{S}_k$, and thus $F(\mathcal{X}) \in \mathcal{T}_k$ for all $\mathcal{X} \in \mathcal{S}_k$. Let $G = F \circ \varkappa$; then $G \in \text{Aut}(\mathfrak{B}_k)$ preserves the families \mathcal{S}_k and \mathcal{T}_k and thus it is determined by a collineation. This completes the proof. \square

PROOF OF THEOREM 1.7. We have $\sim^+ = \sim$ for $k = 1$ and $\sim_- = \sim$ for $k = n - 1$. Consequently, for $k \in \{1, n - 1\}$, we are done by Theorem 1.4. Let $2 \nmid k$ and $1 < k < n - 1$. By (17) and (18), \mathcal{S}_k and \mathcal{T}_k can be distinguished in terms of \sim_- ; applying the duality \varkappa we see that the stars and the tops are distinguishable in terms of \sim^+ as well. As above, from (22) we get that \mathfrak{R}_{k-1} can be defined both in terms of \sim_- and in terms of \sim^+ on $(\mathbf{T-R})_k$. Finally, this observation together with Theorem 1.4 and Corollary 1.6 completes the proof. \square

3 Comments

We do not pay special attention to geometry of the Grassmannians introduced in the paper. Some comments which can be derived immediately from facts established in Sec. 2 are in order, though.

(A) If $2 \mid k$ then $\mathbf{P}_k(\mathbf{T-R})$ is a connected partial linear Γ -space, whose strong (or linear in other words) subspaces are affine spaces. This resembles an affine polar space. And indeed, there are connections. The Grassmannian of regular lines in a projective 3-space endowed with a symplectic polarity is an affine polar space. In general, however, the axiom (3.1.iii) of [5] fails here (and only this one from the list (3.1) fails). Moreover, our Grassmannians have maximal strong subspaces of two distinct dimensions allowed, while maximal strong subspaces of a polar space all have the same dimension. And here, each line is the intersection of exactly two maximal strong subspaces which contain it.

(B) If we consider the Grassmannian $\mathbf{P}_k(\mathbf{T-R})$ with $2 \mid k$ as a part of the whole incidence geometry (1) then we see that a plane in a strong subspace \mathcal{X} is determined either by an element of $\mathbf{R}_{k\pm 2}$ or by $X \in \text{Sub}_{k\pm 2}(\mathbb{V})$ with $\text{rdim}(X) = 2$. Clearly, there is no way to distinguish these two types within the affine space \mathcal{X} . Let us consider an “*affine geometry*” of the form

$$\mathfrak{A} = \langle \text{points of } \mathcal{X}, \text{lines of } \mathcal{X}, \Pi \rangle,$$

where Π are the regular planes in the above meaning. For any 3-subspace Γ of \mathfrak{A} which contains a plane in Π the “dual” structure

$$\langle \{\pi \in \Pi: \pi \subset \Gamma\}, \{L: L \subset \Gamma, L \text{ is a line of } \mathfrak{A}\}, \supset \rangle$$

is an affine 3-space.

(C) Let $2 \nmid k$. Then $\mathbf{P}_k(\mathbf{T-R})$ is a connected partial linear space which satisfies the following variant of the Δ -axiom (cf. [11], [15]):

*a point not on a line L is collinear with
none, exactly one, or all except one point on L .*

Let \mathcal{M} be the family of the maximal subspaces of $\mathbf{P}_k(\mathbf{T-R})$ which are (up to an isomorphism) symplectic copolar spaces. The family \mathcal{M} covers the point set of our Grassmannian in such a way that any two elements of \mathcal{M} intersect in a point, in a line, or are disjoint, each line has exactly two extensions to a subspace in \mathcal{M} , and each clique of collinearity is contained in an element of \mathcal{M} . Very nice characterizations of the geometry on elements of \mathcal{M} can be found, e.g. in [15] and [8].

We conjecture that starting with the properties (A)–(C) one can obtain characterizations of respective Grassmannians in the style of [21].

References

- [1] BICHARA A., TALLINI G., *On a characterization of Grassmann space representing the h -dimensional subspaces in a projective space*, Ann. Discrete Math. **18** (1983), 113–132.
- [2] BLOK R. J., HOFFMAN C., *A quasi Curtis-Tits-Phan theorem for the symplectic group*, J. Algebra **319** (2008), no. 11, 4662–4691.
- [3] BUEKENHOUT F., SHULT E., *On the foundations of polar geometry*, Geom. Dedicata **3** (1974), 155–170.
- [4] CHOW W.-L., *On the geometry of algebraic homogeneous spaces*, Ann. of Math. **50** (1949), 32–67.
- [5] COHEN A. M., SHULT E. E., *Affine polar spaces*, Geom. Dedicata **35** (1990), 43–76.
- [6] COHEN M. A., *Point-line spaces related to buildings*, In *Handbook of incidence geometry*, F. Buekenhout, Ed. North-Holland, Amsterdam, 1995, pp. 647–737.
- [7] CUYPERS H., PASSINI A., *Locally polar geometries with affine planes*, European J. Combin. **13** (1992), no. 1, 39–57.
- [8] CUYPERS H., *Symplectic geometries, transvection groups, and modules*, J. Combin. Theory Ser. A **65** (1994), no. 1, 39–59.
- [9] DIEUDONNÉ J., *La géométrie des groupes classiques*, Springer-Verlag, Berlin, 1971.

- [10] GRAMLICH R., *On the hyperbolic symplectic geometry*, J. Combin. Theory Ser. A **105** (2004), 97–110.
- [11] HALL J. I., *Classifying copolar spaces and graphs*, Quart. J. Math. Oxford (2) **33** (1982), 421–449.
- [12] HALL J. I., *Graphs, geometry, 3-transpositions, and symplectic F_2 -transvection groups*, Proc. London Math. Soc. (3) **58** (1989), no. 1, 89–111.
- [13] HALL J. I., *The hyperbolic lines of finite symplectic spaces*, J. Comb. Theory Ser. A **47** (1988), 284–298.
- [14] LEFÈVRE-PERCSY C., *Geometries with dual affine planes and symplectic quadrics*, Linear Algebra Appl. **42** (1982), 31–37.
- [15] LEFÈVRE-PERCSY C., *Copolar spaces fully embedded in projective spaces*, Ann. Discrete Math. **18** (1983), 553–566.
- [16] PAMBUCCIAN V., *Elementary axiomatizations of projective space and of its associated Grassmann space*, Note Mat. **24** (2004/2005), no. 1, 129–141.
- [17] PANKOV M., PRAŻMOWSKI K., ŻYNEL M., *Geometry of polar Grassmann spaces*, Demonstratio Math. **39** (2006), no. 3, 625–637.
- [18] PRAŻMOWSKA M., PRAŻMOWSKI K., ŻYNEL M., *Affine polar spaces, their Grassmannians, and adjacencies*, Math. Pannonica **20** (2009), no. 1, 37–59.
- [19] PRAŻMOWSKA M., PRAŻMOWSKI K., ŻYNEL M., *Grassmann spaces of regular subspaces*. In review.
- [20] SCHRÖDER E., *Symmetrie als fundamentales Prinzip der Geometrie: Entwicklungen der Spiegelungsgeometrie im vorigen Jahrhundert*, Mitt. Math. Ges. Hamburg **20** (2001), 55–70.
- [21] TALLINI G., *Partial line spaces and algebraic varieties*, Symp. Math., **28** (1986), 203–217.
- [22] VELDKAMP F. D., *Polar geometry I-IV*, Indag. Math. **21** (1959), 512–551.

Authors' address:

Małgorzata Prażmowska, Krzysztof Prażmowski, Mariusz Żynel

Institute of Mathematics, University of Białystok

ul. Akademicka 2, 15-267 Białystok, Poland

malgpraz@math.uwb.edu.pl, krzypraz@math.uwb.edu.pl, mariusz@math.uwb.edu.pl