

IP-RIGIDITY AND EIGENVALUE GROUPS

JON. AARONSON, MARYAM HOSSEINI & MARIUSZ LEMAŃCZYK

ABSTRACT. We examine the class of increasing sequences of natural numbers which are IP-rigidity sequences for some weakly mixing probability preserving transformation. This property is closely related to the uncountability of the eigenvalue group of a corresponding non-singular transformation. We give examples, including a super-lacunary sequence which is not IP-rigid.

§0 INTRODUCTION

Let (X, \mathcal{B}, m) be a standard, continuous, probability space (that is, (X, \mathcal{B}) is a Polish space equipped with its Borel sets and a non-atomic $m \in \mathcal{P}(X)$ (the collection of probability measures on (X, \mathcal{B}))).

We'll denote by $\text{MPT} = \text{MPT}(X, \mathcal{B}, m)$ the collection of invertible, probability preserving transformations of (X, \mathcal{B}, m) . This is a Polish space when equipped with the *coarse topology* with basic neighborhoods of form

$$U(T_0, f_1, \dots, f_N; \epsilon) := \{T \in \text{MPT} : \|f_j \circ T^s - f_j \circ T_0^s\|_{L^2(m)} < \epsilon \ \forall \ 1 \leq j \leq N, \ s = \pm 1\}$$

where $T_0 \in \text{MPT}(X, \mathcal{B}, m)$ and $f_1, \dots, f_N \in L^2(m)$.

Equipped with the coarse topology, $\text{MPT}(X, \mathcal{B}, m)$ is a topological group under composition. It is embedded into the Polish, topological group $\mathcal{U}(L^2(m))$ of unitary operators on $L^2(m)$ equipped with the strong operator topology by the *Koopman representation* $U_T f := f \circ T$ ($T \in \text{MPT}(X, \mathcal{B}, m)$, $f \in L^2(m)$). Accounts of the spectral theory of unitary operators can be found in [KT] and [N2].

Date: May 24th 2012.

2010 Mathematics Subject Classification. 37A05, (37A30, 37A40).

Key words and phrases. Probability preserving transformation, spectral type, rigidity, IP-convergence, Dirichlet measure, eigenvalue group.

Aaronson and Hosseini would like to thank Nicolaus Copernicus University, and Aaronson the University of Surrey, for hospitality provided when this paper was done. Aaronson's research was partially supported by Israel Science Foundation grant No. 1114/08.

Recurrence and rigidity.

A sequence $q \in \mathbb{N}^{\mathbb{N}}(\uparrow) := \{q \in \mathbb{N}^{\mathbb{N}} : q_n < q_{n+1} \ \forall n \geq 1\}$ is called a sequence of *recurrence* for $T \in \text{MPT}(X, \mathcal{B}, m)$ if $\limsup_{n \rightarrow \infty} \mu(A \cap T^{-q_n} A) > 0$ for each $A \in \mathcal{B}$ of positive measure.

Rigidity is a stronger version of recurrence.

An sequence $q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is called a *rigidity sequence* for $T \in \text{MPT}(X, \mathcal{B}, m)$ if $\mu(T^{q_n} A \Delta A) \rightarrow 0$ for each $A \in \mathcal{B}$; equivalently

$$(\times) \quad T^{q_n} \xrightarrow[n \rightarrow \infty]{\text{MPT}} \text{Id.}$$

Using spectral theory one sees that (\times) is equivalent to the **restricted spectral type** σ_T of T (i.e. $U_T|_{L^2(m)_0}$) having the *Dirichlet property* along q , that is

$$(\dagger) \quad \chi_{q_n} \xrightarrow[n \rightarrow \infty]{L^2(\mathbb{T}, \sigma_T)} 1.$$

where $\chi_k(t) := e^{2\pi i k t}$.

Rigidity sequences for non-trivial transformations must be sparse. In particular, unless $T \in \text{MPT}$ is purely periodic any rigidity sequence for T has at most finite intersection with each of its translates whence has zero Banach density.

Additional properties of rigidity sequences are studied in [BJLR] & [EG] including the rigidity properties of **lacunary** and **super-lacunary** sequences, a sequence $q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ being called *lacunary* if $\frac{q_{n+1}}{q_n} \geq \lambda > 1 \ \forall n \geq 1$ and *super-lacunary* if $\frac{q_{n+1}}{q_n} \xrightarrow[n \rightarrow \infty]{} \infty$.

Rigid factors, mild mixing and IP sets. Let $T \in \text{MPT}(X, \mathcal{B}, m)$ and let $q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$. It is well known that the collection of sets

$$(\mathcal{R}) \quad \mathcal{R}(q) := \{A \in \mathcal{B} : m(A \Delta T^{q_n} A) \xrightarrow[n \rightarrow \infty]{} 0\}$$

is a T -invariant, σ -algebra. It corresponds to the maximal factor of T which is rigid along q . The transformation $T \in \text{MPT}(X, \mathcal{B}, m)$ is called *mildly mixing* if it has no non-trivial, rigid factor along any $q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ (as in [FW]).

Since the spectral type σ_S of a factor S of T is absolutely continuous with respect to σ_T , it is evident that T has some non-trivial rigid factor if and only if \exists a *Dirichlet measure* $\mu \ll \sigma_T$, (that is, one satisfying (\dagger) along some $q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$).

An *IP-set* is a collection of “finite sum sets” of form

$$\text{FS}(q) := \{q(F) : F \in \mathcal{F}\}$$

where $q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ and for

$$F \in \mathcal{F} := \{\text{finite, nonempty subsets of } \mathbb{N}\}, \quad q(F) := \sum_{j \in F} q_j.$$

This notion appears in combinatorics, ultrafilter theory, topological dynamics (see [Fu] and [HS]) and also in ergodic theory.

As shown in [Fu], $T \in \text{MPT}(X, \mathcal{B}, m)$ is mildly mixing if and only if $\exists K \subset \mathbb{N}$ which intersects with every finite sum set so that

$$m(A \cap T^{-n} B) \xrightarrow[n \rightarrow \infty, n \in K]{} m(A)m(B) \quad \forall A, B \in \mathcal{B};$$

equivalently (see [HMP1]),

T is not mildly mixing if and only if $\exists q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ so that

$$\inf_{n \in \text{FS}(q)} |\widehat{\mu}(n)| > 0.$$

The considerations involved give rise to the notion of

IP convergence.

Let $q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$. We'll say that a sequence $a : \mathbb{N} \rightarrow Z$ (a metric space) *converges IP to* $L \in Z$ *along* $\text{FS}(q)$

$$\text{written } a(n) \xrightarrow[n \rightarrow \infty]{\text{FS}(q)} L, \text{ in } Z \text{ if}$$

$$a(q(F)) \xrightarrow[F \in \mathcal{F}, \min F \rightarrow \infty]{Z} L.$$

This paper is about

IP-rigidity.

We'll say that $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is

an *IP-rigidity sequence for* T and that T is *IP-rigid along* b if

$$T^n \xrightarrow[n \rightarrow \infty]{\text{FS}(b)} \text{Id in MPT.}$$

Let

$$\text{IPRWM} := \{b \in \mathbb{N}^{\mathbb{N}}(\uparrow) : \exists T \in \text{MPT}(X, \mathcal{B}, m), \text{ weakly mixing \& IP-rigid along } b\}.$$

Any rigid transformation is IP-rigid on some subsequence (see [Fu]). On the other hand if a transformation is IP-rigid on $q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$, then it is rigid along much thicker subsequences (see §5).

Similarly to (\mathfrak{A}), for $q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$, the collection

$$\mathcal{R}_{\text{IP}}(q) := \{A \in \mathcal{B} : m(A \Delta T^n A) \xrightarrow[n \rightarrow \infty]{\text{FS}(q)} 0\}$$

is a T -invariant, σ -algebra. It corresponds to the maximal factor of T which is IP-rigid along q . As above, T has a non-trivial factor, IP-rigid along q if and only if $\lim_{N \rightarrow \infty} \inf_{F \in \mathcal{F}, \min F \geq N} |\widehat{\sigma}_T(q(F))| > 0$.

The existence of IP-Dirichlet measures along $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is related to the groups

$$G_p(b) := \{t \in \mathbb{T} : \sum_{n=1}^{\infty} \|b_n t\|^p < \infty\} \quad (0 < p < \infty) \ \&$$

$$G_{\infty}(b) := \{t \in \mathbb{T} : \|b_n t\| \xrightarrow{n \rightarrow \infty} 0\}$$

where for $x \in \mathbb{R}$, $\|x\| := \min_{n \in \mathbb{Z}} |x - n|$.

These groups are discussed in [AN] and [HMP2].

Results.

Proposition 1 *Suppose that $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$, then*

$$|G_1(b)| > \aleph_0 \implies b \in \text{IPRWM}.$$

Proposition 1 (which is folklore) can be proved using Propositions 1.1 and 1.2 (below).

Theorem 2 *If $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$, then*

$$b \in \text{IPRWM} \implies |G_2(b)| > \aleph_0.$$

Theorem 2 also provides an answer to a question in [BJLR]:

☺ *if $b \in \text{IPRWM}$, then some irrational rotation is rigid along b because if $|G_2(b)| > \aleph_0$ then $\exists \alpha \in G_2(b) \setminus \mathbb{Q}$. It follows that rotation of \mathbb{T} by α is rigid along b . \checkmark*

The converse of theorem 2 holds for **arithmetic** sequences and **Erdos-Taylor** sequences for different reasons.

A sequence $q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is called *arithmetic* if it is either

- *multiplicative* in the sense that $q_n | q_{n+1} \ \forall n \geq 1$; or it is the
- *principal denominator sequence* of some $\alpha \in \mathbb{T} \setminus \mathbb{Q}$, being defined by $q_0 = 1$, $q_1 = a_1$, $q_{n+1} := a_{n+1}q_n + q_{n-1}$ where $\alpha = [0; a_1, a_2, \dots]$ is the continued fraction expansion of α .

Proposition 3 *Let $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$, be arithmetic. The following statements are equivalent.*

- (a) $\overline{\lim}_{n \rightarrow \infty} \frac{b_{n+1}}{b_n} = \infty$. (b) $|G_1(b)| > \aleph_0$.
- (c) $b \in \text{IPRWM}$. (d) $|G_2(b)| > \aleph_0$.

The *Erdos-Taylor sequence associated to* $(a_1, a_2, \dots) \in \mathbb{N}^{\mathbb{N}}$ is $b = (b_1, b_2, \dots) \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ defined by

$$b_1 := 1. \quad b_{n+1} := a_n b_n + 1.$$

Erdos-Taylor sequences were introduced in [ET] and are considered to be “extremely non-arithmetic”.

Proposition 4 *If $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$, is an Erdos-Taylor sequence, then*

$$(i) \sum_{n \geq 1} \left(\frac{b_n}{b_{n+1}} \right)^2 < \infty \iff (ii) b \in \text{IPRWM} \iff (iii) |G_2(b)| > \aleph_0.$$

We’ll see that there are super-lacunary Erdos-Taylor sequences b & $q \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ satisfying $|G_2(b)| > \aleph_0$ & $G_1(b) = \{0\}$ and $G_2(q) = \{0\}$.

Eigenvalue Groups and theorem 2.

Groups of form G_2 appear as eigenvalue groups (see [AN]). Eigenvalue groups and rigidity are related as follows:

An ergodic probability preserving transformation S is not mildly mixing (i.e. has a rigid factor) if and only if there is a conservative, ergodic non-singular transformation T so that $S \times T$ is not ergodic (see [FW]). By the ergodic multiplier theorem of M. Keane (see e.g. §2.7 of [A]), this situation is characterized by $\sigma_S(e(T)) > 0$ where σ_S is the restricted spectral type of S and $e(T)$ is the group of eigenvalues of T .

We prove Theorem 2 in §4 by considering a dyadic cocycle (see below) associated to $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ over the dyadic adding machine. The eigenvalue group of the Mackey range (as in p. 76-77 in [Z]) of this cocycle is $G_2(b)$. In case b is a *growth sequence* as in [A2], that is $b(n) > \sum_{1 \leq k < n} b(k)$, then the Mackey range preserves a σ -finite measure and is isomorphic to the appropriate dyadic tower over the dyadic adding machine (defined in [A2]).

Organization of the paper.

In §1 we establish the basic results on Dirichlet sets and measures and begin to consider membership of IPRWM.

In §2 we consider the class of arithmetic sequences, and prove Proposition 3.

In §3 we prove proposition 4 for Erdos-Taylor sequences and give our main examples.

The proofs of propositions 3 & 4 both use Theorem 2 which is established in §4. In §5 we make some quantitative remarks on the growth of rigid sequences for transformations IP-rigid along some (particular) $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$.

§1 DIRICHLET SETS AND MEASURES

Dirichlet sets. A *Dirichlet set* is a subset $\Gamma \subset \mathbb{T}$ of form

$$\Gamma(b) = \{t \in \mathbb{T} : \chi_{b_n}(t) \xrightarrow[n \rightarrow \infty]{} 1\}$$

where $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ & $\chi_n(t) := e^{2\pi i n t}$.

An *IP-Dirichlet set* is a subset $\Gamma \subset \mathbb{T}$ of form

$$\Gamma(\text{FS}(b)) = \{t \in \mathbb{T} : \chi_n(t) \xrightarrow[n \rightarrow \infty]{\text{FS}(b)} 1\}$$

where $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$. Here, we have

Proposition 1.1 For $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$,

$$\Gamma(\text{FS}(b)) = G_1(b).$$

Proof sketch of \subseteq

It suffices to show that for $t \in \mathbb{R}$,

$$\|nt\| \xrightarrow[n \rightarrow \infty]{\text{FS}(b)} 0 \Rightarrow \sum_{n \geq 1} \|b_n t\| < \infty.$$

For $x \in \mathbb{R}$, let $\lfloor x \rfloor$ be the nearest integer to x (if there are two, take the lesser one), and let

$$\langle x \rangle := x - \lfloor x \rfloor, \text{ then } |\langle x \rangle| = \|x\| \leq \frac{1}{2}.$$

Fix $t \in \mathbb{R}$ so that $\|nt\| \xrightarrow[n \rightarrow \infty]{\text{FS}(b)} 0$, let $K > 0$ be so that

$$\|b(F)t\| < \frac{1}{16} \quad \forall F \in \mathcal{F}, \min F \geq K.$$

If $F, G \subset [K, \infty) \cap \mathbb{N}$ are disjoint finite sets, then

$$\|b(F)t\|, \|b(G)t\|, \|b(F \cup G)t\| < \frac{1}{16}.$$

Since $\langle b(F \cup G)t \rangle - \langle b(F)t \rangle - \langle b(G)t \rangle \in \mathbb{Z}$, this forces $\langle b(F \cup G)t \rangle = \langle b(F)t \rangle + \langle b(G)t \rangle$.

It follows that

$$\sum_{n \geq K, \langle b_n t \rangle \geq 0} \|b_n t\| \leq \frac{1}{16}, \quad \sum_{n \geq K, \langle b_n t \rangle < 0} \|b_n t\| \leq \frac{1}{16} \quad \& \quad \sum_{n \in \mathbb{N}, n \geq K} \|b_n t\| < \frac{1}{8}. \quad \square.$$

Dirichlet measures. A probability measure $\mu \in \mathcal{P}(\mathbb{T})$ is called

- a *Dirichlet* measure if

$$\|\chi_{b_n} - 1\|_{L^2(\mu)} \xrightarrow[n \rightarrow \infty]{} 0$$

for some $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ in which case μ is called *Dirichlet along b* and

- an *IP Dirichlet* measure if

$$\|\chi_n - 1\|_{L^2(\mu)} \xrightarrow[n \rightarrow \infty]{\text{FS}(b)} 0$$

for some $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ in which case μ is called *IP-Dirichlet along b* .

Evidently:

$$\chi_{n_k} \xrightarrow[k \rightarrow \infty]{L^2(\mu)} 1 \text{ if and only if } \widehat{\mu}(n_k) \xrightarrow[k \rightarrow \infty]{} \mu(\mathbb{T}),$$

if μ is IP-Dirichlet along b , then so is any $\nu \ll \mu$,

if μ is IP-Dirichlet, then $\exists b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ so that $\sum_{n \geq 1} \|\chi_{b_n} - 1\|_{L^2(\mu)} < \infty$, whence $\mu(G_1(b)) = 1$ and μ is IP-rigid along b .

By Proposition 1.1, a totally atomic measure $\mu \in \mathcal{P}(\mathbb{T})$ is IP-Dirichlet along b if and only if $\mu(G_1(b)) = 1$. Examples in §4 (below) show that this is false for continuous measures $\mu \in \mathcal{P}(\mathbb{T})$.

Proposition 1.2

$$\text{IPRWM} = \{b \in \mathbb{N}^{\mathbb{N}}(\uparrow) : \exists \mu \in \mathcal{P}(\mathbb{T}) \text{ continuous \& IP-Dirichlet along } b\}.$$

Proof of \subseteq Suppose that $b \in \text{IPRWM}$ and that (X, \mathcal{B}, m, T) is a weakly mixing, probability preserving transformation so that $T^n \xrightarrow[n \rightarrow \infty]{\text{FS}(b)} \text{Id}$. Fix $f \in L^2(m)$, $\int_X f dm = 0$, $\int_X |f|^2 dm = 1$. The spectral measure of f : $\mu \in \mathcal{P}(\mathbb{T})$ is continuous and IP-Dirichlet along b .

Proof of \supseteq Suppose that $\mu_0 \in \mathcal{P}(\mathbb{T})$ is continuous and IP-Dirichlet along b . Let μ be the symmetrization of μ_0 (also continuous and IP-Dirichlet along b) and let (X, \mathcal{B}, m, T) be the shift of the Gaussian process with spectral measure μ . The spectral type of (X, \mathcal{B}, m, T) is $\sigma_T = \sum_{n \geq 0} \mu^{n*}$ where μ^{n*} denotes the n -fold convolution of μ with itself (see e.g. [CFS]). Each μ^{K*} is continuous (whence T is weakly mixing) and IP-Dirichlet along b (since $\widehat{\mu^{K*}}(n) = \widehat{\mu}(n)^K \xrightarrow[n \rightarrow \infty]{\text{FS}(b)} 1$). Every $\nu \ll \sigma_T$ is also IP-Dirichlet along b and T is IP-rigid along b . Thus $b \in \text{IPRWM}$. \square

It follows from proposition 1.2 that if $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is an IP rigidity sequence for some $T \in \text{MPT}$ not of discrete spectrum (i.e. σ_T is not totally atomic) then $b \in \text{IPRWM}$.

We complete this section with a “mixed” multiplicative-finite sum condition for membership in IPRWM.

Proposition 1.3 *Suppose that $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ and that $\exists S \subset \mathbb{N}$ infinite, so that $\sum_{n \in S} \frac{b_n}{b_{n+1}} < \infty$, and $b_n | b_{n+1}$ for $n \notin S$, then $b \in \text{IPRWM}$.*

If $\mathbb{N} \setminus S$ is finite, then $|G_1(b)| > \aleph_0$ by Theorem 5 in [ET], whence $b \in \text{IPRWM}$ by proposition 1.

Proof Assume (without loss of generality) that $b_1 = 1$. We construct a weakly mixing $T \in \text{MPT}(X, \mathcal{B}, m)$ which is IP-rigid along b by cutting and stacking as in Ch. 7 of [N1].

To this end, we construct a nested sequence of Rokhlin towers $(\tau_n)_{n \geq 1}$ of intervals where τ_n has height b_n .

Let τ_1 be $[0, 1]$. To construct τ_{n+1} from τ_n :

- if $n \notin S$ & $b_{n+1} = a_n b_n$, $a_n \in \mathbb{N}$, $a_n \geq 2$ then we cut τ_n into a_n columns and stack.
- If $n \in S$ & $b_{n+1} = a_n b_n + r_n$, $a_n, r_n \in \mathbb{N}$, $1 \leq r_n < b_n$, we cut τ_n into a_n columns and put one spacer interval above the $\lfloor \frac{a_n}{2} \rfloor$ 'th column from the left and $r_n - 1$ spacer intervals above the last column in the right and then we stack.

The tower τ_n is a stack of b_n intervals of length $\prod_{j=1}^{n-1} \frac{1}{a_j}$ called *levels* of τ_n .

It follows from §7.31 in [N1] that the transformation T constructed preserves a finite measure m . A standard argument as in the proof of Proposition 3.10 of [BJLR] shows that T is weakly mixing.

Next, we show that if A is a union of levels in some τ_K , then

$$\sum_{n=1}^{\infty} m(A \Delta T^{b_n} A) < \infty.$$

To see this, we note first that A is also a union of levels in every τ_n ($n \geq K$). Fix $n \geq K$ and write

$$S \cap [K, \infty) = \{s_1 < s_2 < s_3 < \dots\}.$$

Since $\frac{1}{a_{s_\ell+1}} \leq \frac{b_{s_\ell}}{b_{s_\ell+1}} \leq \frac{1}{a_{s_\ell}}$ our assumptions imply $\sum_{\ell=1}^{\infty} \frac{1}{a_{s_\ell}} < \infty$.

To estimate $m(A \Delta T^{-b_n} A)$ for $n \in (s_{\ell-1}, s_\ell]$ we consider the appearance of the tower τ_n as “ τ_n -stalks” inside $\tau_{s_{\ell+1}}$.

A τ_n -stalk in $\tau_{s_\ell+1}$ is a union $\mathfrak{s} = \bigcup_{k=0}^{b_n-1} T^k B$ of levels of $\tau_{s_\ell+1}$ where B is contained in the base of τ_n .

Let $\mathfrak{s} \in \mathcal{B}$ be a τ_n -stalk in $\tau_{s_\ell+1}$, then

$$m(\mathfrak{s} \cap A) = \prod_{j=n}^{s_\ell} \frac{1}{a_j} \cdot m(A).$$

By construction, $\tau_{s_\ell+1}$ consists entirely of τ_n -stalks and spacer stalks added in $\tau_{s_\ell+1}$. Thus, all points of A except those contained in the two τ_n -stalks preceding the spacer stalks added in $\tau_{s_\ell+1}$, return to A at time b_n , so

$$m(A\Delta T^{-b_n} A) = 2m(A \setminus T^{-b_n} A) = 4 \prod_{j=n}^{s_\ell} \frac{1}{a_j} \cdot m(A) \leq \frac{4}{2^{s_\ell-n}} \cdot \frac{1}{a_{s_\ell}} \cdot m(A).$$

Thus, writing $s_0 := K - 1$, we have

$$\begin{aligned} \sum_{n=K}^{\infty} m(A\Delta T^{-b_n} A) &= \sum_{\ell=1}^{\infty} \sum_{n \in (s_{\ell-1}, s_\ell]} m(T^{b_n}(A)\Delta A) \\ &\leq \sum_{\ell=1}^{\infty} \sum_{n \in (s_{\ell-1}, s_\ell]} \frac{4}{2^{s_\ell-n}} \cdot \frac{1}{a_{s_\ell}} \cdot m(A) \\ &\leq 4m(A) \sum_{\ell=1}^{\infty} \frac{1}{a_{s_\ell}} \\ &< \infty. \end{aligned}$$

It follows from this that for A a union of levels in some tower τ_n and $F = \{n_1 < n_2 < \dots < n_k\} \in \mathcal{F}$,

$$\begin{aligned} m(A\Delta T^{b(F)} A) &= m(A\Delta T^{\sum_{j=1}^k b_{n_j}} A) \\ &\leq m(A\Delta T^{b_1} A) + m(T^{b_1} A\Delta T^{\sum_{j=1}^k b_{n_j}} A) \\ &= m(A\Delta T^{b_1} A) + m(A\Delta T^{\sum_{j=2}^k b_{n_j}} A) \\ &\leq \\ &\vdots \\ &\leq \sum_{j=1}^k m(A\Delta T^{b_j} A) \\ &\leq \sum_{n=\min F}^{\infty} m(T^{b_n}(A)\Delta A) \\ &\xrightarrow{F \in \mathcal{F}, \min F \rightarrow \infty} 0. \end{aligned}$$

The collection of measurable sets \mathcal{C} satisfying this last convergence is a σ -algebra and

$$\mathcal{C} \supset \sigma\left(\bigcup_{n \geq 1} \{\text{unions of levels in } \tau_n\}\right) = \mathcal{B}.$$

Thus T is IP-rigid along b . \square

Remark 1.4. The proof of Proposition 1.3 establishes the following proposition:

Suppose that $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$, $p > 0$ and that $\exists S \subset \mathbb{N}$ infinite, so that $\sum_{n \in S} (\frac{b_n}{b_{n+1}})^p < \infty$, and $b_n | b_{n+1}$ for $n \notin S$, then $\exists T \in \text{MPT}(X, \mathcal{B}, m)$ weakly mixing and a dense collection $\mathcal{A} \subset \mathcal{B}$ so that

$$(\heartsuit(p)) \quad \sum_{n \geq 1} m(A \Delta T^{b_n} A)^p < \infty \quad \forall A \in \mathcal{A}.$$

As above, $\heartsuit(1) \implies$ IP-rigidity along b whence $\sigma_T(G_2(b)^c) = 0$ by Theorem 2. Using the spectral theorem, one sees that $\heartsuit(\frac{1}{2}) \implies \sigma_T(G_1(b)^c) = 0$.

§2 ARITHMETIC SEQUENCES

In this section, we prove proposition 3. The implications (b) \implies (c) \implies (d) \implies (a) follow from proposition 1, theorem 2 and theorem 16 in [E] (respectively). None of these uses arithmeticity.

We turn to the remaining implication (a) \implies (b).

Lemma 2.1

If $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is multiplicative and $\sup_{n \geq 1} \frac{b_{n+1}}{b_n} = \infty$, then $|G_1(b)| > \aleph_0$.

C.f. theorem 3 in [ET].

Proof Suppose that $b_{n+1} = a_{n+1}b_n$, where $a_n \geq 2 \forall n \geq 1$.

Since $\sup_{n \geq 1} a_n = \infty$, \exists a subsequence $(n_k)_{k \geq 1}$ such that $a_{n_{k+1}}/a_{n_k} \geq 3 \forall k \geq 1$. Define $t : \Omega := \{0, 1\}^{\mathbb{N}} \rightarrow [0, 1]$ by

$$t(\omega) := \sum_{k=1}^{\infty} \frac{\omega_k}{b_{n_k}}.$$

Since

$$\sum_{k=L+1}^{\infty} \frac{\omega_k}{b_{n_k}} \leq \frac{1}{b_{n_L}} \sum_{j=1}^{\infty} \frac{1}{a_{n_{L+1}} a_{n_{L+2}} \cdots a_{n_j}} \leq \frac{1}{b_{n_L}} \sum_{j=1}^{\infty} \frac{1}{3^j} < \frac{1}{b_{n_L}},$$

we have that $t : \Omega := \{0, 1\}^{\mathbb{N}} \rightarrow [0, 1]$ is strictly increasing (with respect to lexicographic order on Ω), whence injective and $|t(\Omega)| > \aleph_0$.

It suffices to show that $t(\Omega) \subset G_1(b)$.

To see this, fix $\omega \in \Omega$. For $N \geq 1$, we have that

$$b_N t(\omega) = \sum_{k=1}^{\infty} \frac{b_N \omega_k}{b_{n_k}} = \sum_{k \geq 1, n_k \geq N} \frac{\omega_k}{a_{N+1} a_{N+2} \cdots a_{n_k}} \pmod{1},$$

whence

$$\|b_N t(\omega)\| = \sum_{k \geq 1, n_k \geq N} \frac{\omega_k}{a_{N+1} a_{N+2} \cdots a_{n_k}} \leq \sum_{k \geq 1, n_k \geq N} \frac{1}{a_{N+1} a_{N+2} \cdots a_{n_k}} =: \Delta_N.$$

For $n_{K-1} < N \leq n_K$,

$$\Delta_N = \sum_{k \geq K} \frac{1}{a_{N+1} a_{N+2} \cdots a_{n_k}} \leq \sum_{k \geq K} \frac{1}{a_{N+1} a_{N+2} \cdots a_{n_K} \cdot a_{n_k}} \leq \frac{1}{2^{n_K - N} a_{n_K}}.$$

Thus

$$\begin{aligned} \sum_{N=n_1}^{\infty} \|b_N t(\omega)\| &\leq \sum_{N=n_1}^{\infty} \Delta_N = \sum_{k=2}^{\infty} \sum_{N=n_{k-1}+1}^{n_k} \sum_{\nu=k}^{\infty} \frac{1}{a_{N+1} a_{N+2} \cdots a_{n_\nu}} \\ &\leq \sum_{k=2}^{\infty} \sum_{N=n_{k-1}+1}^{n_k} \frac{1}{2^{n_k - N} a_{n_k}} \leq \sum_{k=2}^{\infty} \frac{2}{a_{n_k}} \leq 4 \end{aligned}$$

and $t(\omega) \in G_1(b)$. \square

Lemma 2.2

Let $q = q(\alpha) \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ be the principal denominator sequence of $\alpha \in (0, 1) \setminus \mathbb{Q}$.

If $\sup_{n \geq 1} \frac{q_{n+1}}{q_n} = \infty$, then $|G_1(q)| > \aleph_0$.

Proof As shown in [IN],

for any $t \in [0, 1]$ there is a unique sequence $(\omega_n)_{n \geq 1} \in \prod_{n \geq 1} \{0, 1, \dots, a_n\}$ such that

- $\omega_k \leq a_k$, $\omega_k = a_k \Rightarrow \omega_{k+1} = 0$ and
- $t = \sum_{n=1}^{\infty} \omega_n \langle q_n \alpha \rangle \pmod{1}$.

Since $\sup_{n \geq 1} a_n = \sup_{n \geq 1} \frac{q_{n+1}}{q_n} = \infty$, we can choose a sub-sequence a_{n_k} such that $\sum_{k \geq 1} 1/a_{n_k} < \infty$ and define

$$t : \Omega = \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{T} \text{ by } t(\omega) := \sum_{k=1}^{\infty} \omega_k \langle q_{n_k-1} \alpha \rangle \pmod{1}.$$

By the above, $t : \Omega \rightarrow \mathbb{T}$ is injective. It follows that $|t(\Omega)| > \aleph_0$ and it suffices to show that $t(\Omega) \subset G_1(q)$.

We claim that $\sup_{\omega \in \Omega} \sum_{n=1}^{\infty} \|q_n t(\omega)\| < \infty$.

Fix $K \geq 1$ and consider $n_K - 1 \leq N \leq n_{K+1} - 1$. Then

$$\begin{aligned} \|q_N t(\omega)\| &\leq \sum_{k=1}^{\infty} \|\omega_{n_k-1} q_N q_{n_k-1} \alpha\| \\ &\leq \sum_{k=1}^{\infty} \|q_N q_{n_k-1} \alpha\| \\ &\leq \sum_{k=1}^K \frac{q_{n_k-1}}{q_{N+1}} + \sum_{k=K+1}^{\infty} \frac{q_N}{q_{n_k}}. \end{aligned}$$

Using the fact that $\frac{q_{j+n}}{q_j} \geq \sqrt{2}^{n-2} \forall j, n \geq 1$ we have for some absolute constant C ,

$$\frac{q_{n_k-1}}{q_{N+1}} = \frac{q_{n_k-1}}{q_{n_{K-1}}} \cdot \frac{q_{n_{K-1}}}{q_{n_K}} \cdot \frac{q_{n_K}}{q_{N+1}} \leq \frac{C}{\sqrt{2}^{n_K-n_k}} \cdot \frac{1}{a_{n_K}} \cdot \frac{C}{\sqrt{2}^{N-n_K}} \quad \text{for } k \leq K$$

and

$$\frac{q_N}{q_{n_k}} = \frac{q_N}{q_{n_{K+1}-1}} \cdot \frac{q_{n_{K+1}-1}}{q_{n_{K+1}}} \cdot \frac{q_{n_{K+1}}}{q_{n_k}} \leq \frac{C}{\sqrt{2}^{n_{K+1}-N}} \cdot \frac{1}{a_{n_{K+1}}} \cdot \frac{C}{\sqrt{2}^{n_k-n_{K+1}}} \quad \text{for } k > K.$$

Therefore,

$$\begin{aligned} \sum_{N=n_{K-1}}^{n_{K+1}-2} \|q_N t(\omega)\| &\leq \sum_{N=n_{K-1}}^{n_{K+1}-2} \left(\sum_{k=1}^K \frac{q_{n_k-1}}{q_{N+1}} + \sum_{k=K+1}^{\infty} \frac{q_N}{q_{n_k}} \right) \\ &\leq C^2 \sum_{N=n_{K-1}}^{n_{K+1}-2} \left(\sum_{k=1}^K \frac{1}{a_{n_K} \sqrt{2}^{N-n_k}} + \sum_{k=K+1}^{\infty} \frac{1}{a_{n_{K+1}} \sqrt{2}^{n_k-N}} \right) \\ &\leq C^3 \sum_{N=n_{K-1}}^{n_{K+1}-2} \left(\frac{1}{a_{n_K} \sqrt{2}^{N-n_K}} + \frac{1}{a_{n_{K+1}} \sqrt{2}^{n_{K+1}-N}} \right) \\ &\leq C^4 \left(\frac{1}{a_{n_K}} + \frac{1}{a_{n_{K+1}}} \right) \end{aligned}$$

and

$$\sum_{N=1}^{\infty} \|q_N t(\omega)\| \leq C^4 \sum_{K=1}^{\infty} \left(\frac{1}{a_{n_K}} + \frac{1}{a_{n_{K+1}}} \right) < \infty. \quad \checkmark$$

Hence $q \in \text{IPRWM}$. \checkmark The proof of Proposition 3 is now complete. \checkmark

§3 SUPER-LACUNARY SEQUENCES

Suppose that $b = (b_1, b_2, \dots) \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is super-lacunary, i.e. $\frac{b_{n+1}}{b_n} \xrightarrow{n \rightarrow \infty} \infty$.

As in theorem 17 in [E], we fix $N \geq 1$ with $\frac{b_{n+1}}{b_n} > 10 \forall n \geq N$ and let

$$E := \bigcap_{n \geq N} E_n$$

where

$$E_n := \{t \in [0, 1] : \|b_n t\| \leq \frac{4b_n}{b_{n+1}}\}.$$

Now

$$E_n \supseteq \bigcup_{k=1}^{b_n-1} I_{k,n}$$

where

$$I_{k,n} := \left[\frac{k}{b_n} - \frac{4}{b_{n+1}}, \frac{k}{b_n} + \frac{4}{b_{n+1}} \right].$$

For each $n \geq N$, the intervals $\{I_{k,n} : 1 \leq k < b_n\}$ are disjoint, and each interval $I_{k,n}$ contains at least five disjoint intervals of form $I_{k',n+1}$. It follows that E contains a Cantor set and $|E| > \aleph_0$. Thus, (c.f of theorem 5 in [ET])

Proposition 3.1 *Suppose that $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ & $p > 0$, then*

$$\sum_{n \geq 1} \left(\frac{b_n}{b_{n+1}} \right)^p < \infty \implies |G_p(b)| > \aleph_0.$$

Proof Let $N \geq 1$ & E be as above, then for $t \in E$,

$$\sum_{n \geq 1} \|b_n t\|^p \leq N + 4^p \sum_{n \geq N} \left(\frac{b_n}{b_{n+1}} \right)^p < \infty. \quad \square$$

Proposition 3.2 *Suppose that $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ and that*

$$\sum_{n \geq 1} \left(\frac{b_n}{b_{n+1}} \right)^2 < \infty, \quad \text{then } b \in \text{IPRWM}.$$

Proof By Proposition 1.2, it suffices to construct a continuous probability in \mathbb{T} which is IP-Dirichlet along b .

To this end, let $N \geq 1$, E and $\{I_{k,n} : 1 \leq k < b_n\}$ be as above. As above, for each $n \geq N$, the intervals $\{I_{k,n} : 1 \leq k < b_n\}$ are disjoint, and each interval $I_{k,n}$ contains at least five disjoint intervals of form $I_{k',n+1}$.

Thus we may choose

$$I_n(\omega) = I_{k_n(\omega),n} \quad (n \geq 1, \omega \in \{0, 1\}^n)$$

so that

$$I_{n+1}(\omega, \epsilon) \subset I_n(\omega) \quad \forall n \geq 1, \omega \in \{0, 1\}^n \text{ & } \epsilon = 0, 1$$

and

$$X_{n+1}(\omega, 0) < X_n(\omega) < X_{n+1}(\omega, 1) \text{ where } X_n(\omega) := \frac{k_n(\omega)}{b_n}.$$

Next, for $\omega \in \Omega := \{0, 1\}^{\mathbb{N}}$,

$$X_n(\omega_1, \dots, \omega_n) \xrightarrow{n \rightarrow \infty} X(\omega) \text{ where } \bigcap_{n \geq 1} I_n(\omega_1, \dots, \omega_n) = \{X(\omega)\}$$

and

$$\begin{aligned} b_n X(\omega) &= b_n X_n(\omega) + b_n (X_{n+1}(\omega) - X_n(\omega)) + b_n (X(\omega) - X_{n+1}(\omega)) \\ &= k_n(\omega) + \xi_n(\omega) + \theta_n(\omega) \end{aligned}$$

where

$$\begin{aligned} \xi_n(\omega) &:= b_n (X_{n+1}(\omega) - X_n(\omega)) \text{ \&} \\ \theta_n(\omega) &:= b_n (X(\omega) - X_{n+1}(\omega)). \end{aligned}$$

Note that $|\theta_n(\omega)| \leq \mathcal{E}_n := \frac{4b_n}{b_{n+2}}$ and that by assumption, $\sum_{n \geq 1} \mathcal{E}_n < \infty$.

For $n \geq 1$, $\omega \in \{0, 1\}^n$, $\exists ! p_{n,\omega} : \{0, 1\} \rightarrow (0, 1)$ so that

$$p_{n,\omega}(0) + p_{n,\omega}(1) = 1 \text{ \&} X_{n+1}(\omega, 0)p_{n,\omega}(0) + X_{n+1}(\omega, 1)p_{n,\omega}(1) = X_n(\omega).$$

Define $P : \{\text{cylinders}\} \rightarrow (0, 1)$ by

$$P([a_1, a_2, \dots, a_n]) := \frac{1}{2} \prod_{k=1}^{n-1} p_{k,(a_1, a_2, \dots, a_k)}(a_k).$$

It follows that P is additive and by standard extension theory $\exists \mathbb{P} \in \mathcal{P}(\Omega)$ extending P .

Denoting expectation with respect to \mathbb{P} by \mathbb{E} and writing $\omega = (\omega_1, \omega_2, \dots, \omega_n) \in \{0, 1\}^n$, we have

$$\mathbb{E}(X_{n+1} \|\omega_1, \omega_2, \dots, \omega_n) = X_{n+1}(\omega, 0)p_{n,\omega}(0) + X_{n+1}(\omega, 1)p_{n,\omega}(1) = X_n(\omega),$$

whence

$$\mathbb{E}(\xi_n \|\omega_1, \omega_2, \dots, \omega_n) = b_n (\mathbb{E}(X_{n+1} \|\omega_1, \omega_2, \dots, \omega_n) - X_n(\omega)) = 0$$

and $\mathbb{E}(\xi_n) = 0$.

For $n, k \geq 1$,

$$\begin{aligned} \mathbb{E}(\xi_n \xi_{n+k}) &= b_n b_{n+k} \mathbb{E}(\mathbb{E}(\xi_n \xi_{n+k} \|\omega_1, \omega_2, \dots, \omega_{n+k})) \\ &= b_n b_{n+k} \mathbb{E}((\xi_n(\omega) \mathbb{E} \xi_{n+k} \|\omega_1, \omega_2, \dots, \omega_{n+k})) = 0 \quad \& \end{aligned}$$

$$\mathbb{E}(\xi_n^2) =: \Delta_n \leq \frac{16b_n^2}{b_{n+1}^2}.$$

By assumption $\sum_{n \geq 1} \Delta_n < \infty$, so $\sum_{n \in K} \xi_n$ converges in $L^2(\mathbb{P})$ for every $K \subset \mathbb{N}$ and

$$\mathbb{E}\left(\left(\sum_{n \in K} \xi_n\right)^2\right) = \sum_{n \in K} \mathbb{E}(\xi_n^2) = \sum_{n \in K} \Delta_n.$$

The measure $\mu := \mathbb{P} \circ X^{-1} \in \mathcal{P}(\mathbb{T})$ is continuous. We claim that it is IP-Dirichlet along b .

To check this, let $F \subset \mathbb{N} \cap [K, \infty)$ be finite and write $\Xi_F := \sum_{n \in F} \xi_n$, then

$$\begin{aligned} \|\chi_{b(F)} - 1\|_{L^2(\mu)} &\leq \left\| \left\langle \sum_{N \in F} b_N t \right\rangle \right\|_{L^2(\mu)} \\ &= \left\| \left\langle \sum_{N \in F} (\xi_N + \theta_N) \right\rangle \right\|_{L^2(\mathbb{P})} \\ &\leq \|\langle \Xi_F \rangle\|_{L^2(\mathbb{P})} + \sum_{N \in F} \mathcal{E}_N. \end{aligned}$$

Next,

$$\begin{aligned} \|\langle \Xi_F \rangle\|_{L^2(\mathbb{P})}^2 &= \mathbb{E}(\langle \Xi_F \rangle^2) \\ &= \mathbb{E}(1_{[|\Xi_F| \leq \frac{1}{2}]}) \langle \Xi_F \rangle^2 + \mathbb{E}(1_{[|\Xi_F| > \frac{1}{2}]}) \langle \Xi_F \rangle^2 \\ &\leq \mathbb{E}(\Xi_F^2) + \frac{1}{4} \mathbb{P}[|\Xi_F| > \frac{1}{2}] \quad \because \langle \Xi_F \rangle^2 \leq \frac{1}{4}, \\ &\leq 2\mathbb{E}(\Xi_F^2) \quad \text{by Tchebychev's inequality} \\ &= 2 \sum_{N \in F} \mathbb{E}(\xi_N^2) \leq 2 \sum_{N=K}^{\infty} \Delta_N. \end{aligned}$$

Thus,

$$\begin{aligned} \|\chi_{n(F)} - 1\|_{L^2(\mu)} &\leq \|\langle \Xi_F \rangle\|_{L^2(\mathbb{P})} + \sum_{N=K}^{\infty} \mathcal{E}_N \\ &\leq \sqrt{2 \sum_{N=K}^{\infty} \Delta_N} + \sum_{N=K}^{\infty} \mathcal{E}_N \\ &\xrightarrow{K \rightarrow \infty} 0 \end{aligned}$$

proving that μ is IP-Dirichlet along b . \square

Remark. The converses to propositions 3.1 & 3.2 are false. It is easy to construct $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ multiplicative, super-lacunary so that $\sum_{n \geq 1} (\frac{b_n}{b_{n+1}})^p = \infty \quad \forall p > 0$. By proposition 3, $|G_1(b)| > \aleph_0$ & $T \in \text{IPRWM}$.

Erdos-Taylor sequences & proposition 4.

We begin with a strong converse to Proposition 3.1 for Erdos-Taylor sequences:

Proposition 3.3 *Suppose that $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is an Erdos-Taylor sequence and let $p > 0$, then*

$$\sum_{n \geq 1} \left(\frac{b_n}{b_{n+1}} \right)^p = \infty \quad \implies \quad G_p(b) = \{0\}.$$

This was stated in [ET] for $p = 1$ and b the Erdos-Taylor sequence associated to $(2, 3, \dots)$. See also Théorème 2 in [P].

Proof Let $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ be the Erdos-Taylor sequence associated to $(a_1, a_2, \dots) \in \mathbb{N}^{\mathbb{N}}$ and let $t \in \mathbb{R} \setminus \mathbb{Z}$, then

$$\|b_n t\| < \frac{\|t\|}{2a_n} \implies \|b_{n+1} t\| > \frac{\|t\|}{2}.$$

If $p > 0$ and $\sum_{n \geq 1} (\frac{b_n}{b_{n+1}})^p = \infty$, then for $t \in \mathbb{R} \setminus \mathbb{Z}$,

- either $\|b_n t\| \geq \frac{\|t\|}{2a_n}$ eventually and $\sum_{n \geq 1} \|b_n t\|^p = \infty$, or
 - $\|b_{n+1} t\| > \frac{\|t\|}{2}$ infinitely often and $\sum_{n \geq 1} \|b_n t\|^p = \infty$.
- Either way, $t \notin G_p(b)$. \square

Proof of Proposition 4 The implications (i) \implies (ii) \implies (iii) \implies (i) follow from proposition 3.2, theorem 2 and proposition 3.3 (respectively). \square

Examples.

- If $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is the Erdos-Taylor sequence associated to $(2, 3, \dots)$, then

$$(a) \quad b \in \text{IPRWM} \quad \& \quad (b) \quad G_1(b) = \{0\}.$$

- If $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is the Erdos-Taylor sequence associated to (a_2, a_3, \dots) where $a_n \rightarrow \infty$ and $\sum_{n \geq 1} \frac{1}{a_n^2} = \infty$ (e.g. $a_n := \lfloor \sqrt{n} \rfloor$), then b is superlacunary, $G_2(b) = \{0\}$ and b is not a sequence of IP-rigidity for any probability preserving transformation other than the identity.

§4 PROOF OF THEOREM 2

We prove Theorem 2 using dyadic cocycles over the dyadic odometer.

Let $\Omega := \{0, 1\}^{\mathbb{N}}$, and let $P \in \mathcal{P}(\Omega)$ be symmetric product measure: $P = \prod(\frac{1}{2}, \frac{1}{2})$, and let $\tau : \Omega \rightarrow \Omega$ be the dyadic odometer defined by

$$\tau(1, \dots, 1, 0, \omega_{\ell+1}, \dots) = (0, \dots, 0, 1, \omega_{\ell+1}, \dots)$$

where $\ell = \ell(\omega) := \min \{n \geq 1 : \omega_n = 0\}$.

The *dyadic cocycle* $\varphi : \Omega \rightarrow \mathbb{Z}$ associated to $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is defined by

$$\varphi(\omega) := b_{\ell(\omega)} - \sum_{k=1}^{\ell(\omega)-1} b_k$$

and its *skew product*

$$\tau_\varphi : \Omega \times \mathbb{Z} \rightarrow \Omega \times \mathbb{Z} \text{ is defined by } \tau_\varphi(x, n) = (\tau(x), n + \varphi(x)).$$

Define $Q : \Omega \times \mathbb{Z} \rightarrow \Omega \times \mathbb{Z}$ by $Q(x, n) := (x, n + 1)$ and fix $p \in \mathcal{P}(\Omega \times \mathbb{Z})$, $p \sim P \times \#$ where $\#$ is counting measure on \mathbb{Z} .

There is (see [Z], pp. 76–77) an ergodic, non-singular transformation $(\mathfrak{X}, \mathcal{B}(\mathfrak{X}), \mathfrak{q}, \mathfrak{T})$ of a standard probability space and a map $\pi : \Omega \times \mathbb{Z} \rightarrow \mathfrak{X}$ so that

$$p \circ \pi^{-1} = \mathfrak{q}, \quad \pi^{-1}\mathcal{B}(\mathfrak{X}) = \{A \in \mathcal{B}(\Omega \times \mathbb{Z}) : \tau_\varphi A = A\} \quad \& \quad \pi \circ Q = \mathfrak{T} \circ \pi.$$

The ergodic, non-singular transformation $(\mathfrak{X}, \mathcal{B}(\mathfrak{X}), \mathfrak{q}, \mathfrak{T})$ is called the *Mackey range of (τ, φ)* .

In case b is a growth sequence, equivalently $\varphi : \Omega \rightarrow \mathbb{N}$, there is a σ -finite, invariant T -invariant measure $\mathfrak{m} \sim \mathfrak{q}$ with respect to which the Mackey range $(\mathfrak{X}, \mathcal{B}(\mathfrak{X}), \mathfrak{m}, \mathfrak{T})$ is isomorphic to the tower over $(\Omega, \mathcal{B}(\Omega), P, \tau)$ with height function φ (aka the *dyadic tower with growth sequence b* in [A2]).

The collection of eigenvalues of the Mackey range is

$$e(\mathfrak{T}) := \{t \in \mathbb{T} : \exists F \in L^\infty(\mathfrak{q}), F \neq 0, F \circ \mathfrak{T} = e^{2\pi i t} F\}$$

and it follows from the definitions that

$$e(\mathfrak{T}) = \mathcal{T}(\tau, \varphi) := \{s \in \mathbb{T} : \exists f \in L^\infty(P), f \neq 0, f \circ \tau = e^{2\pi i s \varphi} f\}.$$

It is shown in §2 of [AN] (see also Theorem 2.6.3 of [A1]) that

$$(\clubsuit) \quad \mathcal{T}(\tau, \varphi) = G_2(b).$$

Although formally, (\clubsuit) was only stated for growth sequences in [AN] and [A1], the proofs do not use this condition and apply to arbitrary $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$.

Consider the Polish group $\mathfrak{B}(\mu) := \{f \in L^2(\mu) : |f| \equiv 1\}$ equipped with $L^2(\mu)$ -distance.

Lemma 4.1

If the probability $\mu \in \mathcal{P}(\mathbb{T})$ is IP Dirichlet along b , then $\exists \mathcal{X} : \Omega \rightarrow \mathfrak{B}(\mu)$ continuous so that

$$(\spadesuit) \quad \sup_{\omega \in \Omega} \|\chi_{b(K(\omega) \cap [1, n])} - \mathcal{X}(\omega)\|_{L^2(\mu)} \xrightarrow{n \rightarrow \infty} 0$$

where $K(\omega) := \{n \geq 1 : \omega_n = 1\}$.

Proof

Suppose that $\mu \in \mathcal{P}(\mathbb{T})$ is IP Dirichlet along b . Fix $\omega \in \Omega$. We claim that the sequence $n \mapsto \chi_{b(K(\omega) \cap [1, n])}$ is Cauchy in $L^2(\mu)$. To see this, let

$$\mathcal{E}_n := \sup_{F \in \mathcal{F}, \min F \geq n} \|\chi_{b(F)} - 1\|_{L^2(\mu)}$$

then by assumption $\mathcal{E}_n \xrightarrow{n \rightarrow \infty} 0$. Evidently

$$\|\chi_{b(K(\omega) \cap [1, n])} - \chi_{b(K(\omega) \cap [1, n+k])}\|_{L^2(\mu)} = \|\chi_{b(K(\omega) \cap [n+1, n+k])} - 1\|_{L^2(\mu)} \leq \mathcal{E}_n$$

whence $\exists \mathcal{X} : \Omega \rightarrow \mathfrak{B}(\mu)$ so that

$$\chi_{b(K(\omega) \cap [1, n])} \xrightarrow[n \rightarrow \infty]{L^2(\mu)} \mathcal{X}(\omega) \text{ uniformly in } \omega \in \Omega.$$

For $\omega \in \Omega$,

$$\|\chi_{b(K(\omega) \cap [1, n])} - \mathcal{X}(\omega)\|_{L^2(\mu)} \xleftarrow[k \rightarrow \infty]{} \|\chi_{b(K(\omega) \cap [1, n])} - \chi_{b(K(\omega) \cap [1, n+k])}\|_{L^2(\mu)} \leq 2\mathcal{E}_n$$

proving (\heartsuit) . Clearly, for each $n \geq 1$, $\omega \mapsto \chi_{b(K(\omega) \cap [1, n])}$ is continuous ($\Omega \rightarrow \mathfrak{B}(\mu)$) and so continuity of $\mathcal{X} : \Omega \rightarrow \mathfrak{B}(\mu)$ follows from the uniformity of the convergence. \checkmark

Note that the converse of Lemma 4.1 is also true.

Completion of the proof

Now suppose that $\mu \in \mathcal{P}(\mathbb{T})$ is IP-Dirichlet along b . By Lemma 4.1, $\exists \mathcal{X} : \Omega \rightarrow \mathfrak{B}(\mu)$ satisfying (\heartsuit) .

We claim that

$$(\clubsuit) \quad \mathcal{X}(\tau\omega) = \chi_{\varphi(\omega)} \mathcal{X}(\omega).$$

To see this, note that $\chi_{b(K(\omega) \cap [1, n])}(t) = \prod_{k=1}^n \chi_{\omega_k b_k}(t) =: \mathcal{X}_n(\omega, t)$. For $n > \ell(\omega)$,

$$\frac{\mathcal{X}_n(\tau\omega)}{\mathcal{X}_n(\omega)} = \frac{\chi_{b(K(\tau\omega) \cap [1, n])}}{\chi_{b(K(\omega) \cap [1, n])}} = \prod_{k=1}^n \frac{\chi_{(\tau\omega)_k b_k}}{\chi_{\omega_k b_k}} = \chi_{\varphi(\omega)}.$$

Since $\frac{\mathcal{X}_n(\tau\omega)}{\mathcal{X}_n(\omega)} \xrightarrow[n \rightarrow \infty]{\mathfrak{B}(\mu)} \frac{\mathcal{X}(\tau\omega)}{\mathcal{X}(\omega)}$, this proves (\clubsuit) .

By (\heartsuit) , $\exists n_J \rightarrow \infty$ so that

$$\sum_{J \geq 1} \|\mathcal{X}_{n_J} - \mathcal{X}\|_{L^2(P \times \mu)} < \infty \leq \sum_{J \geq 1} \sup_{\omega \in \Omega} \|\chi_{b(K(\omega) \cap [1, n_J])} - \mathcal{X}(\omega)\|_{L^2(\mu)} < \infty$$

(where $\mathcal{X}(\omega, t) := \mathcal{X}(\omega)(t)$). Hence $\mathcal{X}_{n_J} \rightarrow \mathcal{X}$ $P \times \mu$ -a.e. and by Fubini's theorem, $\exists \Lambda \in \mathcal{B}(\mathbb{T})$, $\mu(\Lambda) = 1$ so that

$$\prod_{k=1}^{n_J} \chi_{\omega_k b_k}(t) \xrightarrow[J \rightarrow \infty]{} \mathfrak{X}_t(\omega) = \mathcal{X}(\omega)(t) \quad \forall t \in \Lambda \text{ \& } P\text{-a.e. } \omega \in \Omega.$$

By (\clubsuit) , for $t \in \Lambda$,

$$\mathfrak{X}_t \circ \tau = e^{2\pi i t \varphi} \mathfrak{X}_t \quad P\text{-a.e.}$$

and $t \in \mathcal{T}(\tau, \varphi)$. By (\clubsuit) , $t \in G_2(b)$. \square

The following example shows that the converse to Theorem 2 is false.

Example 4.2 $\exists b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ & $\mu \in \mathcal{P}(\mathbb{T})$ non-atomic, not IP-Dirichlet along b but so that $\mu(G_2(b)) = 1$.

Construction:

Define $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ by $b_n := \prod_{k=1}^n a_k$ with $a_k := k + 1$.

Consider the mapping $t: \Omega := \prod_{k \geq 1} \{0, 1\} \rightarrow [0, 1]$ defined by

$$t(\omega) = t(\omega_1, \omega_2, \dots) := \sum_{n \geq 1} \frac{\omega_n}{b_n}.$$

This is injective and Borel measurable, so $t(\Omega)$ is an uncountable, Borel set in $[0, 1]$. We claim that

$$(\heartsuit) \quad t(\Omega) \subset G_2(b).$$

Proof of (\heartsuit)

For $\omega \in \Omega$ and $N \geq 1$, we have that

$$b_N t(\omega) = \frac{\omega_{N+1}}{a_{N+1}} + \frac{1}{a_{N+1}} \sum_{k \geq 2} \frac{\omega_{N+k}}{a_{N+2} \cdots a_{N+k}} \pmod{1}.$$

Now

$$\left| \sum_{k \geq 2} \frac{\omega_{N+k}}{a_{N+2} \cdots a_{N+k}} \right| \leq \sum_{k \geq 2} \frac{1}{a_{N+2} \cdots a_{N+k}} < \frac{1}{a_{N+1}}.$$

Thus, we have

$$(\spadesuit) \quad \langle b_N t(\omega) \rangle = \frac{\omega_{N+1}}{a_{N+1}} + \theta_N(\omega) \quad \text{where } \theta_N \leq \frac{1}{a_{N+1}^2},$$

whence $|\langle b_N t(\omega) \rangle|^2 \leq \frac{2}{N^2}$ and

$$\sum_{k=1}^{\infty} |\langle b_k t(\omega) \rangle|^2 \leq \frac{\pi}{3}. \quad \square \quad (\heartsuit)$$

Now define $P \in \mathcal{P}(\Omega)$ by $P := \prod_{k \geq 1} (\frac{1}{2}\delta_0 + \frac{1}{2}\delta_1)$ and set $\mu := P \circ t^{-1}$, then $\mu \in \mathcal{P}(G_2(b))$. We now show that μ is not IP-Dirichlet along b .

Fix $1 < \lambda < e^{\frac{1}{3}}$, then by (\spadesuit) ,

$$\left| \sum_{\lambda^N < k < \lambda^{N+1}} \langle b_N t(\omega) \rangle - \sum_{\lambda^N < k < \lambda^{N+1}} \frac{\omega_{N+1}}{a_{N+1}} \right| \leq \sum_{\lambda^N < k < \lambda^{N+1}} \frac{2}{k^2} \xrightarrow{N \rightarrow \infty} 0$$

and setting $s_n := \sum_{j=1}^n \omega_j$, $\kappa_n := \lceil \lambda^n \rceil$, $\ell_n := \lfloor \lambda^{n+1} \rfloor$, we have

$$\begin{aligned} \sum_{\lambda^N < k < \lambda^{N+1}} \frac{\omega_k}{a_k} &= \sum_{\kappa_N \leq k \leq \ell_N} \frac{s_k - s_{k-1}}{k+1} \\ &= \sum_{\kappa_N \leq k \leq \ell_N} \frac{s_k}{k+1} - \sum_{\kappa_{N-1} \leq k \leq \ell_{N-1}} \frac{s_k}{k+2} \\ &= \frac{s_{\ell_N}}{\ell_N + 1} - \frac{s_{\kappa_N}}{\kappa_N + 2} + \sum_{\kappa_N \leq k \leq \ell_{N-1}} \frac{s_k}{(k+1)(k+2)}. \end{aligned}$$

By the SLLN, we have

$$\frac{s_N}{N} \xrightarrow[N \rightarrow \infty]{} \frac{1}{2} \quad \text{a.s.}$$

Writing $\Gamma_N \approx \Delta_N$ as $N \rightarrow \infty$ to mean $\Gamma_N \approx \Delta_N \xrightarrow[N \rightarrow \infty]{} 0$, we have for P -a.e. $\omega \in \Omega$, as $N \rightarrow \infty$:

$$\begin{aligned} \sum_{\lambda^N < k < \lambda^{N+1}} \langle b_N t(\omega) \rangle &\approx \sum_{\lambda^N < k < \lambda^{N+1}} \frac{\omega_k}{a_k} \\ &= \frac{s_{\ell_N}}{\ell_N + 1} - \frac{s_{\kappa_N}}{\kappa_N + 2} + \sum_{\kappa_N \leq k \leq \ell_{N-1}} \frac{s_k}{(k+1)(k+2)} \\ &\approx \sum_{\kappa_N \leq k \leq \ell_{N-1}} \frac{1}{2k} \\ &\longrightarrow \log \lambda. \end{aligned}$$

Thus

$$\begin{aligned} \sup_{F \in \mathcal{F}, \min F > \lambda^N} \|\chi_{b(F)} - 1\|_{L^2(\mu)}^2 &\geq \mathbb{E}(|\exp[2\pi i \sum_{\lambda^N < k < \lambda^{N+1}} b_N t(\omega)] - 1|^2) \\ &\geq 4\mathbb{E}(|\sum_{\lambda^N < k < \lambda^{N+1}} \langle b_N t(\omega) \rangle|^2) \\ &\longrightarrow 2 \log \lambda. \quad \square \end{aligned}$$

§5 REMARKS ON THE THICKNESS OF RIGIDITY SEQUENCES

Remark 5.1. Rigidity sequences for weakly mixing transformations can be arbitrarily "large" within the limitation of density zero.

It follows from the definitions that for $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ a growth sequence,

$$|\text{FS}(b) \cap [1, n]| \asymp 2^{c(n)}$$

where $c(n) = \min \{k \geq 1 : b_k \geq n\}$ and if $T \in \text{MPT}(X, \mathcal{B}, m)$ is IP-rigid along b , one might expect a rigid sequence at least of this thickness.

Indeed, in this case, by Theorem 2, $\sigma_T(e(\mathfrak{T})^c) = 0$ where \mathfrak{T} is the dyadic tower with growth sequence b (since by [AN] $e(\mathfrak{T}) = G_2(b)$). By theorem 4 in [A2] $\exists L \subset \mathbb{N}$ with

$$T^n \xrightarrow[n \rightarrow \infty, n \in L]{\text{MPT}} \text{Id} \quad \& \quad \frac{|L \cap [1, n]|}{2^{c(n)}} \xrightarrow[n \rightarrow \infty]{} \infty.$$

By the Corollary in [A2], $\forall a(n) > 0$, $\frac{a(n)}{n} \xrightarrow[n \rightarrow \infty]{} 0$, there is a weakly mixing $T \in \text{MPT}(X, \mathcal{B}, m)$ and $L \subset \mathbb{N}$ such that

$$T^n \xrightarrow[n \rightarrow \infty, n \in L]{\text{MPT}} \text{Id} \quad \& \quad \frac{|L \cap [1, n]|}{a(n)} \xrightarrow[n \rightarrow \infty]{} \infty.$$

For more on this phenomenon, see §3 in [BJLR].

Remark 5.2. Let $b \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ be a growth sequence and suppose that $\text{H-dim}(G_1(b)) > \alpha$ (where $\alpha \in (0, 1)$ and H-dim denotes Hausdorff dimension).

We claim that $\exists T \in \text{MPT}$ weakly mixing & IP-rigid along b with the property that for any sequence $L \subset \mathbb{N}$ along which T is rigid:

$$(\star) \quad \sum_{n=1}^{\infty} \frac{|L \cap [1, n]|}{n^{2-\alpha}} < \infty.$$

Note that it follows from this that $\sum_{n=1}^{\infty} \frac{2^{c(n)}}{n^{2-\alpha}} < \infty$.

Proof of (\star)

As in the proof of theorem 1 of [A3], it follows from Frostman's theorem ([Fr], see also [KS]) that $\exists \mu \in \mathcal{P}(G_1(b))$ so that

$$\sum_{n=1}^{\infty} \frac{|\widehat{\mu}(n)|}{n^{1-\alpha}} < \infty.$$

Let $T \in \text{MPT}(X, \mathcal{B}, m)$ be the associated Gaussian automorphism. By Proposition 1, T is IP-rigid along b .

Suppose that T is rigid along $L \subset \mathbb{N}$. In particular $\exists N \geq 1$ so that $|\widehat{\mu}(n)| \geq \frac{1}{2} \quad \forall n \in L, n > N$.

It follows that

$$\begin{aligned}
\sum_{n=1}^{\infty} \frac{|L \cap [1, n]|}{n^{2-\alpha}} &= \sum_{n=1}^{\infty} \sum_{k=1}^n 1_L(k) \frac{1}{n^{2-\alpha}} \\
&= \sum_{k=1}^{\infty} 1_L(k) \sum_{n=k}^{\infty} \frac{1}{n^{2-\alpha}} \\
&\leq \frac{1}{1-\alpha} \sum_{k=1}^{\infty} 1_L(k) \frac{1}{k^{1-\alpha}} \\
&\leq \frac{N}{1-\alpha} + \frac{2}{1-\alpha} \sum_{k=N+1}^{\infty} \frac{|\widehat{\mu}(k)|}{k^{1-\alpha}} \\
&< \infty. \quad \heartsuit(\heartsuit)
\end{aligned}$$

Remark 5.3. The condition (\heartsuit) is sharp. In §5 of [A3], $\forall \alpha \in (0, 1)$ a growth sequence $b^{(\alpha)} \in \mathbb{N}^{\mathbb{N}}(\uparrow)$ is exhibited with

$$\text{H-dim}(G_1(b^{(\alpha)})) = \alpha \ \& \ 2^{c^{(\alpha)}(n)} \gg n^{1-\alpha}$$

whence if $T \in \text{MPT}(X, \mathcal{B}, m)$ is so that $\sigma_T(G_2(b)) = 1$, then (again by theorem 4 in [A2]) $\exists L \subset \mathbb{N}$ rigid for T with $\frac{|L \cap [1, n]|}{n^{1-\alpha}} \rightarrow \infty$ and therefore

$$\sum_{n=1}^{\infty} \frac{|L \cap [1, n]|}{n^{2-\alpha}} = \infty.$$

REFERENCES

- [A1] J. Aaronson, *An introduction to infinite ergodic theory*; Mathematical Surveys and Monographs, 50. American Mathematical Society Providence, RI(1997)
- [A2] J. Aaronson, *Rational ergodicity, bounded rational ergodicity and some continuous measures on the circle*. A collection of invited papers on ergodic theory. Israel J. Math. 33 (1979), no. 3-4, 181–197 (1980).
- [A3] J. Aaronson, *The eigenvalues of non-singular transformations*. Israel Journal of Math. **45**, 297-312 (1983).
- [AN] J. Aaronson, M.G. Nadkarni, *L_{∞} eigenvalues and L_2 spectra of non-singular transformations*, Proc. London Math. Soc. 930 55 91987), 538-570.
- [BJLR] V. Bergelson, A. del Junco, M. Lemańczyk, J. Rosenblatt, *Rigidity and non-recurrence along sequences*, submitted (arxiv.org/pdf/1103.0905).
- [CFS] I.P. Cornfeld, S.V. Fomin, Y.G. Sinai, *Ergodic Theory*, Springer-Verlag, New York, 1982.
- [E] H. G. Eggleston, *Sets of fractional dimensions which occur in some problems of number theory*, Proceedings of the London Mathematics Soc (2) 54 (1952) 42-93.
- [EG] T. Eisner, S. Grivaux, *Hilbertian Jamison sequences and rigid dynamical systems*, J. Functional Analysis **261** (2011), 2013-2052.
- [ET] P. Erdős and S. J. Taylor, *On the set of points of convergence of a lacunary trigonometric series and the equidistribution properties of related sequences*, Proceedings of the London Mathematics Society (3) 7 (1957) 598-615.

- [Fr] O. Frostman, *Potentiel d'équilibre et capacité des ensembles*, Thesis, Lund, 1935
- [Fu] H. Furstenberg, *Recurrence in Ergodic Theory and Combinatorial Number Theory*, Princeton University Press, Princeton, New Jersey, 1981.
- [FW] H. Furstenberg, B. Weiss, *The finite multipliers of infinite ergodic transformations*, in “The structure of attractors in dynamical systems” (Proc. Conf., North Dakota State Univ., Fargo, N.D., 1977), pp. 127–132, *Lecture Notes in Math.* **668**, Springer, Berlin, 1978.
- [HS] N. Hindman, D. Strauss, *Algebra in the Stone-Čech compactification*, De Gruyter, 1998
- [HMP1] B. Host, J.-F. Mela and F. Parreau. *Analyse harmonique des mesures*. Asterisque **135–136**, SMF, (1986).
- [HMP2] B. Host, J.-F. Mela and F. Parreau. *Nonsingular transformations and spectral analysis of measures*. Bull. Soc. Math. France **119** (1991), no. 1, 33–90.
- [IN] Ito, Sh. ; Nakada, H. *Approximations of real numbers by the sequence $\{n\alpha\}$ and their metrical theory*. Acta Math. Hungar. **52** (1988), no. 1-2, 91–100.
- [KS] J. P. Kahane and R. Salem, *Ensembles parfaites et séries trigonométriques*, Hermann, Paris, 1963.
- [KT] A. Katok and J.-P. Thouvenot, *Spectral properties and combinatorial constructions in ergodic theory*, Handbook of Dynamical Systems. Vol. 1B, 649–743, Elsevier, Amsterdam, 2006.
- [N1] Nadkarni, M. G. *Basic ergodic theory*. Hindustan Book Agency, New Delhi, 1998.
- [N2] M.G. Nadkarni, *Spectral Theory of Dynamical Systems*, Hindustan Book Agency, New Delhi, 1998.
- [P] F. Parreau, *Ergodicité et pureté des produits de Riesz*, Ann. Inst. Fourier, **40** (1990) 391–405.
- [Z] R.J. Zimmer, *Ergodic theory and semisimple groups*, Monographs in Mathematics, vol 81, Birkhäuser Verlag, Basel, 1984.

(Aaronson) SCHOOL OF MATH. SCIENCES, TEL AVIV UNIVERSITY, 69978
TEL AVIV, ISRAEL.

E-mail address: aaro@tau.ac.il

(Hosseini) FACULTY OF MATHEMATICAL SCIENCE, UNIVERSITY OF GUILAN,
RASHT, IRAN

E-mail address: hoseini@guilan.ac.ir

(Lemańczyk) FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, NICO-
LAUS COPERNICUS UNIVERSITY, UL. CHOPINA 12/18, 87-100 TORUŃ, POLAND

E-mail address: mlem@mat.uni.torun.pl