Grassmannians of lines defined in the geometry of a pseudo-polarity

K. Prażmowski and M. Żynel

June 21, 2018

Abstract

The regular point-line geometry with respect to a pseudo-polarity is introduced. It is weaker than the underlying metric-projective geometry. The automorphism group of this geometry is determined. This geometry can be also expressed as the geometry of regular lines and planes.

Mathematics Subject Classification (2010): 51A50, 51F20, 51A45. Keywords: pseudo-polarity, regular subspaces, Grassmannian.

Introduction

Projective geometry and affine geometry over fields with even characteristic, though have some "strange" properties do not differ so strongly from geometries over arbitrary field. In particular, most of standard methods used to characterize geometry of the Grassmann spaces associated with them can be applied here.

The situation changes when we pass to the orthogonal geometry. Let \mathfrak{P} be a projective space coordinatized by a field \mathfrak{F} and let \perp be a projective conjugacy (projective polarity) defined on \mathfrak{P} ; then \mathfrak{P} equipped with \perp is referred to as a *metric-projective space*.

Some standard derivatives are associated with the space (\mathfrak{P}, \perp) . The first is a polar space, which consists of selfconjugate (absolute) points and isotropic (singular or absolute) lines (cf. [11]). One of fundamental results which states that the underlying metric-projective space can be reconstructed in terms of the associated polar space, is valid in all cases except exactly one: when $\operatorname{char}(\mathfrak{F}) = 2$ and \perp is a *pseudo-polarity*. In [13, Subsection 2.4.18] a polarity \perp is said to be a pseudopolarity if the set **H** of selfconjugate points with respect to \perp is a proper (possibly empty) subspace of \mathfrak{P} . In this paper we additionally assume that **H** is a hyperplane (this means that \mathfrak{F} is perfect, see also [5, Subsection 2.1.5]). Pseudo-polarities are related to pseudo-quadrics and pseudo-quadratic forms (cf. [2, Section 10.2], [12, Chapter 6]). So, in the aforementioned point the geometry of a pseudo-polarity is exceptional, though the geometry of its polar space is, generally, known.

In a metric projective geometry an important role is played by the family of regular subspaces i.e. subspaces with the trivial radical.

If the conjugacy \perp is not symplectic then the structure of regular points and regular lines is equivalent to the underlying metric geometry (cf. [10]). If \perp is

symplectic then one cannot have a space with regular points and lines. If \perp is a pseudo-polarity then, admittedly char(\mathfrak{F}) = 2, but (\mathfrak{P}, \perp) does contain regular points and lines. They yield a new geometry a subject of our paper. This geometry is stronger than the affine geometry \mathfrak{A} obtained by deleting nonregular points i.e. a projective hyperplane **H** from \mathfrak{P} , but *weaker* than the underlying metric-projective geometry in that one can reconstruct the space \mathfrak{P} in terms of regular points and lines, (cf. 3.13), but it is impossible to reconstruct the polarity \perp in these terms (cf. 3.15). In this point the geometry of a pseudo-polarity is essentially exceptional. The geometry of regular points and lines satisfies the dual of Γ -axiom (cf. [3]), a relative of the Δ -axiom (cf. [4], [6]). However, our geometry is neither one of copolar spaces, nor a regular point-line geometry of a polarity (cf. [9]). Therefore, results of these theories cannot be directly copied into our work.

In our paper we also discuss more properties of the *regular geometry* of a pseudopolarity, in particular, we determine the automorphism group of this geometry (cf. 3.22, 3.24).

The regular geometry of a pseudo-polarity, considered on the varieties of the regular subspaces, presents some more oddities: the incidence system of the regular subspaces contains isolated objects; there are also flags of regular objects whose end-object may be contained in (or contains) no regular successor (or predecessor respectively). In geometry of incidence structures isolated objects generally are trash: nothing can be defined in terms of them and they are unredefinable in general, as each automorphism of the structure can arbitrarily permute them. So, it is a good reason to remove them from considerations. Since there are many combinations of isolated objects we get various structures of Grassmannians over regular subspaces. Without entering into details of the general theory of k-Grassmannians we show how our apparatus works in case of structures over regular lines. It turns out (cf. 3.20, 4.1, 5.14) that all they are definitionally equivalent to the regular point-line geometry. This, in particular, let us determine the automorphisms. We are convinced that analogous results can be achieved for arbitrary k.

1 Basic notions

Let \perp be a projective polarity in a finite dimensional metric projective space (\mathfrak{P}, \perp) coordinatized by a vector space \mathbb{V} over a perfect field \mathfrak{F} with even characteristic. In a suitable coordinate system the form ξ which determines \perp has its matrix of one of the following forms (cf. [5, Subsection 2.1.5]). Let us set

$$O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad \nabla = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \text{and} \quad \nabla' = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}.$$

It is seen that $\det(\nabla), \det(\nabla') \neq 0$. Then the matrix of ξ is one of the following

$$\begin{bmatrix} 1 & 0 & \cdots & \cdots & 0 \\ 0 & \nabla & 0 & \cdots & 0 \\ \vdots & 0 & \nabla & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & 0 \\ 0 & 0 & \cdots & 0 & \nabla \end{bmatrix} \begin{bmatrix} \nabla' & 0 & \cdots & 0 \\ 0 & \nabla & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & \nabla \end{bmatrix} \begin{bmatrix} \nabla & 0 & \cdots & 0 \\ 0 & \nabla & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & \nabla \end{bmatrix}$$

Type (1) Type (2) Type (3)

For a subspace U we write $U^{\perp} = \{p : p \perp U\}$, $\operatorname{Rad}(U) = U \cap U^{\perp}$, and $\operatorname{rdim}(U) = \operatorname{dim}(\operatorname{Rad}(U))$. A subspace U is called *regular* when $\operatorname{Rad}(U) = \emptyset$. The dimension $\operatorname{dim}(U)$ of a subspace U coincides with the dimension of the vector subspace W of \mathbb{V} such that $U = \{\langle u \rangle : u \in W, u \neq \theta\}$. In particular, a point has dimension 1 and a line has dimension 2. Frequently, a point p will be identified with the one-element subspace $\{p\}$ which consists of p; consequently, we frequently write $U \cap U' = p$ instead of $U \cap U' = \{p\}$. We write \mathcal{R}_k for the class of regular k-dimensional subspaces of (\mathfrak{P}, \perp) . In the sequel we are interested in the incidence structure (cf. [3])

$$\mathcal{R} := (\mathcal{R}_k \colon 0 \le k \le \dim(\mathfrak{P}) + 1),$$

which is more closely investigated for indices 1, 2 and 2, 3. Various geometries arise when we delete isolated objects in such structures.

1.1 Grassmannians

Grassmann spaces frequently appear in the literature, just to mention [3], [8]. The most general definition could be probably as follows. Let X be a nonempty set and let \mathcal{P} be a family of subsets of X. Assume that there is a dimension function dim : $\mathcal{P} \to \{0, \ldots, n\}$ such that $I = \langle \mathcal{P}, \subset, \dim \rangle$ is an incidence geometry. Write \mathcal{P}_k for the set of all $U \in \mathcal{P}$ with dim(U) = k. For $H \in \mathcal{P}_{k-1}$ and $B \in \mathcal{P}_{k+1}$ with $H \subset B$ the set

$$\mathbf{p}(H,B) := \{ U \in \mathcal{P}_k \colon H \subset U \subset B \}$$
(4)

is called a *pencil*; \mathcal{G}_k stands for the family of all such pencils. If 0 < k < n, then a *k*-th Grassmann space over I is a point-line geometry

$$\mathbf{P}_k(I) := \langle \mathfrak{P}_k, \mathfrak{G}_k \rangle.$$

This is the most common understanding of a Grassmann space. We used to call it a *space of pencils* for its specific lineset and to distinguish it from a closely related point-line geometry consisting of \mathcal{P}_k as points and \mathcal{P}_{k+1} as lines, namely

$$\mathbf{G}_k(I) := \langle \mathfrak{P}_k, \mathfrak{P}_{k+1}, \subset
angle$$

which we call a k-th Grassmannian over I (cf. [7, Section 1.1.2]).

In our settings, we write \mathcal{H}_k for the set of k-dimensional subspaces of \mathfrak{P} . Then a projective pencil is a set (4) with $\mathcal{P} = \mathcal{H}$ and a regular pencil is a set (4) with $\mathcal{P} = \mathcal{R}$.

1.2 Methodological issues

Most of the results of our paper consists in various definabilities/undefinabilities of particular structures, derived from the incidence geometry \mathcal{R} , in terms of other structures of the same type. Clearly, in a structure like this various notions can be introduced and some notions can originate in the underlying projective and metric projective geometry. But proving our results we must be strict: in each particular case we must be sure that the respective definition can be expressed entirely in terms of the language with names only for primitive notions of the structure in which the second one is defined. The safest way to ensure that is to write down respective definitions as "formal formulas" in a formal language, and we do follow this convention. Our structures are, primarily, point-line geometries and the only primitive notion used to characterize their geometry is the *incidence* relation. We use the symbol |to name the incidence relation between points and lines. Note that $| = \in$ in $\mathbf{P}_k(I)$ while $| = \subset$ in $\mathbf{G}_k(I)$.

By a *triangle* in a point-line geometry we mean three points, called *vertices*, and three lines, called *sides*, where every vertex is incident to exactly two sides (or dually, every side is incident to exactly two vertices).

1.3 \perp is symplectic

The form is of type (3). Consequently, $\dim(\mathbb{V}) = 2m$ for some integer m. In this case each point is selfconjugate; the point set of the respective polar space is the point set of \mathfrak{P} .

1.4 \perp is a pseudo-polarity

The form is of type (1) or (2). Then the set of selfconjugate points under \perp is a subspace **H** of \mathfrak{P} . We assume that **H** is a hyperplane; then the pole **b** of **H** is a point. The set \mathcal{R}_1 of regular points is the hyperplane complement: the complement of **H**. One can imagine the geometry of regular points and regular lines as a fragment of an affine geometry.

The restriction $\perp_{\mathbf{H}}$ of the conjugacy \perp to \mathbf{H} determines on \mathbf{H} a (possibly degenerate) symplectic polarity. If $\operatorname{Rad}(\mathbf{H}) = \emptyset$, this polarity is nondegenerate. In general, $\dim(\operatorname{Rad}(\mathbf{H})) \leq 1$.

Again, two types of geometry may occur.

1.4.1 **b** lies on **H**

In this case $\operatorname{Rad}(\mathbf{H}) = \mathbf{b}$, so the polarity \perp restricted to \mathbf{H} is a degenerate symplectic polarity. Then $n = \dim(\mathbb{V}) = 2k$ for some integer k and the form ξ has form (2); it may be written as

 $\xi([x_0, x_1, \dots, x_{2k-1}], [y_0, y_1, \dots, y_{2k-1}]) = x_0 y_0 + \sum_{i=0}^{k-1} (x_{2i} y_{2i+1} + x_{2i+1} y_{2i})$ (cf. [5]). The hyperplane **H** and its pole **b** are characterized by the equations **H**: $x_0 = 0$, **b** = [0, 1, 0, ..., 0].

1.4.2 b does not lie on H

Then $n = \dim(\mathbb{V}) = 2k + 1$ and the form ξ has form (1); it may be written as

 $\xi([x_0, x_1, \dots, x_{2k}], [y_0, y_1, \dots, y_{2k}]) = x_0 y_0 + \sum_{i=1}^k (x_{2i-1}y_{2i} + x_{2i}y_{2i-1})$ (cf. [5]). The hyperplane **H** and its pole **b** are characterized by the equations

$$\mathbf{H}: x_0 = 0, \qquad \mathbf{b} = [1, 0, \dots, 0].$$

Note that for $x \in \mathbf{H}$ we have $x^{\perp} = \mathbf{b} + (x^{\perp} \cap \mathbf{H})$. Therefore, to determine x^{\perp} it suffices to know the restriction $\perp_{\mathbf{H}}$ of the conjugacy \perp to \mathbf{H} . On the other hand, this restriction determines on \mathbf{H} a symplectic polarity and this polarity is nondegenerate as $\operatorname{Rad}(\mathbf{H}) = \emptyset$.

2 Results, general

Let \perp , **H**, and **b** be like in Section 1.4. Two families of specific regular subspaces arise:

$$\mathcal{A}_k = \{ A \in \mathcal{R}_k \colon A \not\subset \mathbf{H} \}, \qquad \qquad \mathcal{A}_k^{\circ} = \{ A \in \mathcal{R}_k \colon \mathbf{b} \notin A \}.$$

If A is a subspace of \mathfrak{P} we write $A^{\infty} := A \cap \mathsf{H}$ for the set of selfconjugate points of A. Note that $\operatorname{Rad}(A)$ is always a (possibly empty) set of selfconjugate points, i.e. $\operatorname{Rad}(A) \subset A^{\infty} \subset \mathsf{H}$.

We begin with a simple but very significant fact.

FACT 2.1 ([9, Cor. 1.3]). If a subspace A contains a regular hyperplane or A is a hyperplane of a regular subspace, then $\operatorname{rdim}(A) \leq 1$.

FACT 2.2. If A is a subspace not contained in **H**, then $\dim(A \cap (A^{\infty})^{\perp}) \geq 1$.

PROOF. Set $k := \dim(A)$, $m := \dim(A \cap (A^{\infty})^{\perp})$. Then $\dim(A^{\infty}) = k - 1$ and $n \ge \dim(A + (A^{\infty})^{\perp}) = k + (n - (k - 1)) - m = n + 1 - m$, which yields our claim.

The subspace $A \cap (A^{\infty})^{\perp}$ will be denoted by $\operatorname{Hrd}(A)$ and will play an essential role (slightly similar to the role of the radical $\operatorname{Rad}(A)$ of A). Fact 2.2 says that $\operatorname{Hrd}(A)$ is at least a point for every subspace A not contained in \mathbf{H} .

PROPOSITION 2.3. Let A be a subspace of \mathfrak{P} not contained in H. The following conditions are equivalent:

- (i) the subspace A is regular;
- (ii) the subspace Hrd(A) is a point (cf. 2.2);
- (iii) $\operatorname{rdim}(A^{\infty}) \leq 1$ and
 - (a) either the subspace A^{∞} is regular in the (possibly degenerated) symplectic projective geometry induced on **H**,
 - (b) or $\operatorname{Rad}(A^{\infty})$ is a point p and $A \not\subset p^{\perp}$.

PROOF. From 2.2, Hrd(A) is at least a point.

(i) \implies (iii): Since A^{∞} is a hyperplane in A, by 2.1, $\operatorname{rdim}(A^{\infty}) \leq 1$. If $\operatorname{rdim}(A^{\infty}) = 0$, then A^{∞} is regular. If $\operatorname{rdim}(A^{\infty}) = 1$, then $\operatorname{Rad}(A^{\infty})$ is a point, say p. The point p could be the only point in $\operatorname{Rad}(A)$. As A is regular, we have $p \notin \operatorname{Rad}(A)$ i.e. $p \not\perp A$ or, in other words, $A \not\subset p^{\perp}$.

(iii) \Longrightarrow (ii): Let A^{∞} be regular. Suppose that $\operatorname{Hrd}(A)$ contains a line K. Then $K \subset A$. As A^{∞} is a hyperplane in A so, K and A^{∞} share a point q. Note that $q \in A^{\infty} \cap (A^{\infty})^{\perp} = \operatorname{Rad}(A^{\infty})$ which is impossible. Consequently, $\operatorname{Hrd}(A)$ is a point.

Now, let $\operatorname{Rad}(A^{\infty})$ be a point p and $A \not\subset p^{\perp}$. Then $p \in \operatorname{Hrd}(A)$. Suppose that there is $q \in \operatorname{Hrd}(A)$, $q \neq p$. If $q \in A^{\infty}$, then $q \in \operatorname{Rad}(A^{\infty})$, which is impossible. So $q \in A \setminus A^{\infty}$. On the other hand $p \in A^{\infty} \cap (A^{\infty})^{\perp}$ and $q \in (A^{\infty})^{\perp}$, hence $p \perp A^{\infty} + q = A$ which contradicts our assumption that $A \not\subset p^{\perp}$. So, $\operatorname{Hrd}(A)$ is a point.

(ii) \implies (i): Let $\operatorname{Hrd}(A)$ be a point p and suppose that $q \in \operatorname{Rad}(A)$. Then $q \in A$ and $q \in A^{\perp} \subset (A^{\infty})^{\perp}$, so q = p. Hence $p \perp A$ and $p \in A^{\infty}$, which gives $p \perp A + (A^{\infty})^{\perp}$. Note that $\dim(A + (A^{\infty})^{\perp}) = n$, so $p \perp V$ a contradiction. \Box

NOTE. If $p = \operatorname{Rad}(A^{\infty})$, then $p = \operatorname{Hrd}(A)$.

NOTE. Let A be a regular subspace not contained in \mathbf{H} and $k = \dim(A)$. Then the restriction \perp_A of \perp to A is a pseudo-polarity and $\operatorname{Hrd}(A)$ is the pole of A^{∞} within A. Thus

 $if \ 2 \mid k, \ then \ \mathrm{Hrd}(A) \in A^{\infty}, \qquad and \qquad if \ 2 \nmid k, \ then \ \mathrm{Hrd}(A) \notin A^{\infty}.$

Recall that the induced geometry on **H** is symplectic. Therefore, A^{∞} may be regular iff $2 \mid \dim(A^{\infty})$ i.e. iff $2 \nmid \dim(A)$. This gives

COROLLARY 2.4. Let A be a subspace not contained in \mathbf{H} with $k = \dim(A)$. If $2 \mid k$ then A is regular iff $\operatorname{Rad}(A^{\infty})$ is a point p and $A \setminus \mathbf{H}$ misses p^{\perp} . If $2 \nmid k$ then A is regular iff A^{∞} is regular.

As particular instances of 2.3 we obtain a series of criterions of regularity.

LEMMA 2.5. A line L not contained in **H** is regular iff it is not contained in the hyperplane $(L^{\infty})^{\perp}$.

COROLLARY 2.6. Let p be a regular point. A line L through p is regular iff it misses $p^{\perp} \cap \mathbf{H}$.

PROOF. Let $q := L \cap H$. By 2.5 L is regular iff $p \notin q^{\perp}$ i.e. iff $q \notin p^{\perp}$, as required. \Box

Note that every nonregular line on \mathbf{H} is totally isotropic.

LEMMA 2.7. If $\mathbf{b} \in \mathbf{H}$, then each affine line with direction \mathbf{b} is regular and no line on \mathbf{H} through \mathbf{b} is regular. If $\mathbf{b} \notin \mathbf{H}$, then no affine line through \mathbf{b} is regular.

PROOF. Let $\mathbf{b} \in \mathbf{H}$. In view of 2.5 it suffices to note that $\mathbf{b}^{\perp} = \mathbf{H}$ and affine line are those not contained in \mathbf{H} .

Let $\mathbf{b} \notin \mathbf{H}$. Then for each $p \in \mathbf{H}$ we have $p \in \operatorname{Rad}(\overline{p, \mathbf{b}})$ so, the line $\overline{p, \mathbf{b}}$ is not regular.

A direct consequence of 2.3 and 2.4 is

LEMMA 2.8. Let A be a plane not contained in **H**. Clearly, A^{∞} is a line. The following conditions are equivalent:

- (i) the plane A is regular;
- (ii) the subspace Hrd(A) is a point (cf. 2.2);
- (iii) the line A^{∞} is regular in the (possibly degenerated) symplectic projective geometry induced on **H**.

LEMMA 2.9. Let A be a plane contained in **H**. Then either

(i) $\operatorname{Rad}(A)$ is a point and all non-regular lines on A form a pencil through that point, or

- (ii) $\operatorname{Rad}(A)$ is a line and no line on A is regular.
- (iii) $\operatorname{Rad}(A)$ is the entire plane A, i.e. A is totally isotropic.

PROOF. It is clear that $1 \leq \operatorname{rdim}(A)$ as $\dim(A) = 3$ and $A \subseteq \mathbf{H}$.

In case $p := \operatorname{Rad}(A)$ is a point then $A \subseteq p^{\perp}$, so p is the radical of every line on A through p. Note that if q is the radical of some line on A not through p, then $A \subseteq q^{\perp}$ and we would have $\overline{p, q} = \operatorname{Rad}(A)$ which is impossible.

Now, if $L := \operatorname{Rad}(A)$ is a line, then every line of A crosses L and thus is non-regular.

Remark. Let A be a plane not contained in \mathbf{H} . If Hrd(A) is a point p not on \mathbf{H} then A is regular, but no line through p contained in A is regular.

Further we assume that:

lines of \mathfrak{P} are of size at least 6,

which means that the ground field of \mathfrak{P} is not GF(2) and not GF(4). Most of our reasonings remain true for GF(4) and those few which fail will be indicated.

3 Grassmannians of regular points and lines

3.1 Regular point-line geometry

In what follows we shall pay attention to the Grassmannian of regular points and lines of the pseudo-polarity \perp , namely

$$\mathfrak{G}_1 := \mathbf{G}_1(\mathfrak{R}) = \langle \mathfrak{R}_1, \mathfrak{R}_2, \subset \rangle.$$

Observe, first, that the set \mathcal{R}_1 is simply the point-complement of the hyperplane \mathbf{H} . Let $\mathfrak{A} = \langle \mathcal{R}_1, \mathcal{L} \rangle$ be the affine space obtained from \mathfrak{P} by deleting the hyperplane \mathbf{H} ; then $\mathcal{A}_2 \subset \mathcal{L}$.

FACT 3.1. If $\mathbf{b} \in \mathbb{R}_1$, then \mathbf{b} is an isolated point in \mathfrak{G}_1 . If $L \in \mathbb{R}_2$ and $L \subset \mathbf{H}$, then L is an isolated line in \mathfrak{G}_1 .

PROOF. The first statement restates 2.7, while the other is trivial.

Let \mathfrak{B}_1 be the structure obtained from \mathfrak{G}_1 by deleting its isolated points and lines. Then

$$\mathfrak{B}_1 = \langle \mathcal{A}_1^{\circ}, \mathcal{A}_2, \in \rangle$$

Note that \mathfrak{B}_1 is a substructure of \mathfrak{A} . This structure, primarily, and related structures will be investigated in this section.

A plane A of \mathfrak{P} not contained in \mathbf{H} is a plane of \mathfrak{A} and will be referred to as an affine plane; A^{∞} is the set of its improper points. Similarly, a line L of \mathfrak{P} not contained in \mathbf{H} is a line of \mathfrak{A} and will be referred to as an affine line; L^{∞} is its improper point.

So, let A be an affine plane and set $L := A^{\infty}$.

FACT 3.2. Through each $q \in L$ there passes a nonregular affine line M contained in A (in every direction in A there is a nonregular line contained in A).

PROOF. It suffices to consider $A \cap q^{\perp}$, which is at least a line.

FACT 3.3. If A contains two parallel nonregular lines then A is nonregular as well.

PROOF. Let $M_1 \parallel M_2$ be nonregular, $M_1, M_2 \subset A$. Take $q = M_1^{\infty}$. Then $q \perp M_1, M_2$, so $q \perp M_1 + M_2 = A$ i.e. $q \in \text{Rad}(A)$.

FACT 3.4. If A contains a triangle with all its sides nonregular then $\operatorname{Rad}(A) = A^{\infty}$ and no affine line contained in A is regular.

PROOF. Let a_1, a_2, a_3 be the vertices of a required triangle and $q_i = \overline{a_j, a_k}^{\infty}$ for $\{i, j, k\} = \{1, 2, 3\}$. Then $a_i \perp q_j, q_k$, so $a_i \perp A^{\infty}$ for each *i*. Thus $A \perp A^{\infty}$.

FACT 3.5. The following possibilities may occur.

- L is nonregular: Then L is isotropic i.e. $L \perp L$, A is nonregular, and we have two cases.
 - $\operatorname{Rad}(A)$ is the line L: Then $L \perp A$ and A contains no regular line. In this case $\operatorname{Hrd}(A) = A$.
 - $\operatorname{Rad}(A)$ is a point q: Then $q \in L$. An affine line K on A is nonregular iff $K^{\infty} = q$. In this case $\operatorname{Hrd}(A) = L$.
- L is regular: Then A is regular and Hrd(A) is an affine point p on A. An affine line K on A is nonregular iff $p \in K$.

PROOF. The claim is nearly evident. It only remains to prove the above characterization of nonregular lines on A. Let K be an affine line on A and $x = K^{\infty}$.

Let $A \perp L$; then $L \ni x \perp A \supset K$ and thus K is not regular.

Let $\operatorname{Rad}(A) = q$ and $L \perp L$; clearly, $q \in L$. Let $q \in K$; as above, $q \perp A \supset K$ and thus $q \perp K$. Let $q \notin K$ and suppose that K is not regular. Then $x \neq q$ and $x \perp L + K = A$, which gives $A \perp q + x = L$. The obtained contradiction yields that K is regular.

Let A be regular. From definition, $p \perp L$ and thus $p \perp x$; with $x \perp x$ from $p \in K$ we obtain $x \perp K$, so K is not regular. Assume that K misses p and suppose that K is not regular. Then K and $\overline{x, p}$ are two parallel nonregular lines on A and thus A is not regular. This contradiction yields that K must be regular.

As a consequence we get

FACT 3.6. Let A be an affine plane with $\operatorname{rdim}(A) \leq 1$. In view of 3.5 the nonregular lines on A form a pencil. Its vertex will be denoted by $\mathbf{q}(A)$. This pencil is

proper and $\mathbf{q}(A)$ is an affine point when $\operatorname{rdim}(A) = 0$, or

parallel and $\mathbf{q}(A)$ is a point on A^{∞} when $\operatorname{rdim}(A) = 1$.

LEMMA 3.7. Let M_1, M_2 be two parallel regular lines in \mathcal{L} and let A be the affine plane that contains them. Then either A is regular or Rad(A) is a point. In both cases A contains a pair K_1, K_2 of regular lines which intersect in an affine point such that K_i crosses M_j in an affine point for all i, j.

PROOF. Let a_1 be any point on M_1 and K_0 be the unique nonregular line through a_1 (in above notation, either $K_0 = \overline{a_1, q}$ or $K_0 = \overline{a_1, p}$, resp.). Let $y = M_2 \cap K_0$ and a_2 be an affine point on M_2 distinct from y; take $K_1 = \overline{a_1, a_2}$. Let $b \in K_1$ be an affine point distinct from a_1, a_2 and $K_2 = \overline{b, y}$.

COROLLARY 3.8. The formula

$$M_{1} \parallel M_{2} \iff (\exists K_{1}, K_{2})(\exists p, a_{1}, a_{2}, b_{1}, b_{2}) [K_{1} \neq K_{2} \land p \mid K_{1}, K_{2}$$

$$a_{1} \mid K_{1}, M_{1} \land a_{2} \mid K_{1}, M_{2} \land b_{1} \mid K_{2}, M_{1} \land b_{2} \mid K_{2}, M_{2} \land p \not \mid M_{1}, M_{2}]$$

$$\land \neg \exists a[a \mid M_{1}, M_{2}] \quad (5)$$

defines the parallelism of regular lines in terms of the geometry of \mathfrak{B}_1 .

LEMMA 3.9. Let A be an affine plane with $\operatorname{rdim}(A) \leq 1$. Then A contains a triangle Δ with regular sides and, moreover, through each affine point on A distinct from $\mathbf{q}(A)$ there passes a regular line which crosses the sides of Δ in at least two affine points.

PROOF. In view of 3.6 the nonregular lines on A form a pencil with the vertex $\mathbf{q}(A)$: a proper one if $\mathbf{q}(A)$ is an affine point p or a parallel one when $\mathbf{q}(A)$ is a point qon A^{∞} . Thus the existence of a required triangle Δ is evident. Let a_1, a_2, a_3 be the vertices of Δ and x be an arbitrary affine point on A. The only lines through x that may not cross appropriately the sides of Δ are the following $\overline{x}, a_1 \parallel \overline{a_2}, a_3$ $\overline{x}, a_2 \parallel \overline{a_1, a_3}, \overline{x}, \overline{a_3} \parallel \overline{a_1, a_2}, \text{ and } \overline{x}, \mathbf{q}(A)$. (Note that from the Fano axiom valid in \mathfrak{A} , the lines through a_i parallel to $\overline{a_j, a_l}, \{i, j, l\} = \{1, 2, 3\}$ have indeed a common point x.) From assumptions, there are at least 5 lines through x contained in A and thus there exists a line required as well.

In \mathfrak{B}_1 for a triangle Δ with the sides L_1, L_2, L_3 we define

$$\pi(\Delta) := \{ x \colon (\exists K) (\exists a, b) [x, a, b \mid K \land a \neq b \land ((a \mid L_1 \land b \mid L_2) \lor (a \mid L_2 \land b \mid L_3) \lor (a \mid L_1 \land b \mid L_3))] \}.$$
(6)

Let \mathcal{P} be the set of planes of \mathfrak{A} , and let \mathcal{P}_i be the set of planes in \mathcal{P} with $\operatorname{rdim} = i$. Then $\mathcal{P}_{01} := \mathcal{P}_0 \cup \mathcal{P}_1$ is the set of planes with $\operatorname{rdim} \leq 1$. For $A \in \mathcal{P}_{01}$ write $[A] := A \setminus \{\mathbf{q}(A)\}$.

COROLLARY 3.10. If $A \in \mathcal{P}_1$, then [A] = A. If $A \in \mathcal{P}_0$, then [A] is an affine plane with one point deleted. Moreover, we have

$$\{[A]: A \in \mathcal{P}_{01}\} = \{\pi(\Delta): \Delta \text{ is a triangle in } \mathfrak{B}_1\}.$$
(7)

LEMMA 3.11. Let *L* be a nonregular affine line through an affine point *p* in \mathfrak{A} . Then $q := L^{\infty} \neq \mathbf{b}$. Let $A \in \mathcal{P}_{01}$ contain *L* and $M := A^{\infty}$, so A = L + M with $q \in M \subset \mathbf{H}$. Then one of the following holds.

(i) $M \cap q^{\perp} = q \ (M \not\subset q^{\perp})$. In this case M is regular, so A is regular as well. Clearly, $\mathbf{q}(A) \in L$, so $M \subset \mathbf{q}(A)^{\perp}$. To have $p \neq \mathbf{q}(A)$ we need $M \not\subset p^{\perp}$.

(ii) $M \subset q^{\perp}$. In this case either $M \not\perp L$ or **b** is an affine point on L.

PROOF. Case (i) is evident. If $q \in M \subset q^{\perp}$, then $M \perp M$. To have $M \neq \operatorname{Rad}(A)$ we must have $x \not\perp M$ for each $x \in A \setminus M$; this is equivalent to $L \not\perp M$. Assume that $L \perp M$. Then $M = \operatorname{Rad}(A)$ and $L \perp q^{\perp} \cap \mathbf{H}$. Comparing dimensions we get $L^{\perp} = q^{\perp} \cap \mathbf{H}$ which gives $L = q + \mathbf{b}$.

LEMMA 3.12. Let L be a nonregular line and a_1, a_2, a_3 be affine points on L in \mathfrak{A} . If $\mathbf{b} \neq a_1, a_2, a_3$, then there are distinct $A_1, A_2 \in \mathcal{P}_{01}$ with $a_1, a_2, a_3 \in [A_1] \cap [A_2]$.

PROOF. In view of 3.11 it suffices to find two lines $M_1, M_2 \subset \mathbf{H}$ through $q = L^{\infty}$ such that $M_1, M_2 \not\subset q^{\perp}, a_1^{\perp}, a_2^{\perp}, a_3^{\perp}$. The required lines exist as we have at least 6 lines in a projective pencil.

THEOREM 3.13. The affine space \mathfrak{A} is definable in terms of \mathfrak{B}_1 and, consequently, \mathfrak{P} is definable in terms of \mathfrak{B}_1 as well.

PROOF. Let \mathbf{L}_0 be the ternary collinearity relation on the set of affine points distinct from **b** on nonregular lines. By 3.12 this relation is definable in terms of the geometry of \mathfrak{B}_1 . On the other hand \mathbf{L}_0 is a ternary equivalence relation in the sense of [1, §4.5]. So, let \mathcal{L}_0 be the family of equivalence classes of \mathbf{L}_0 . If **b** is an improper point we are through, as \mathcal{L}_0 consists of the affine nonregular lines. So, assume that **b** is an affine point. Then \mathcal{L}_0 consists of the nonregular lines not through **b** and the sets $L \setminus \{\mathbf{b}\}$ where L is nonregular through **b**. For every triangle Δ of \mathfrak{B}_1 we set

$$\pi'(\Delta) = \bigcup \{ L \in \mathcal{A}_2 \cup \mathcal{L}_0 \colon |\pi(\Delta) \cap L| \ge 2 \}.$$

Let $L \in \mathcal{L}_0$ and L' be the nonregular line of \mathfrak{A} that contains L. Either $\mathbf{b} \in L'$ and then every plane containing L also contains \mathbf{b} , or $\mathbf{b} \notin L'$ and then there is exactly one plane $L + \mathbf{b}$ containing L and \mathbf{b} . Therefore, if no plane through a line L is an affine plane, then $L \in \mathcal{L}_0$ and we set $L' := L \cup \{\mathbf{b}\}$; otherwise we set L' := L. After that $\mathcal{L}'_0 := \{L' : L \in \mathcal{L}_0\}$ is the set of all nonregular lines of \mathfrak{A} and $\mathcal{L} = \mathcal{A}_2 \cup \mathcal{L}'_0$. \Box

For a set X of affine points we write \overline{X} for the least subspace of \mathfrak{P} which contains X.

LEMMA 3.14. Let $q \in H$. Then the set

$$[q] := \{a: a \text{ is a point of } \mathfrak{A}, \ \overline{a,q} \notin \mathcal{A}_2\}$$

$$(8)$$

is the set of affine points on q^{\perp} , and thus $\overline{[q]} = q^{\perp}$.

Similarly, if a is a point of \mathfrak{A} then the set $[a] = \{K^{\perp} : a \mid K, K \in \mathcal{L} \setminus \mathcal{A}_2\}$ coincides with the set $a^{\perp} \cap \mathbf{H}$; but not with the set a^{\perp} , unhappily.

THEOREM 3.15. The metric projective space (\mathfrak{P}, \perp) is not definable in terms of the geometry of \mathfrak{B}_1 .

PROOF. Let W be the subspace of \mathbb{V} with $\mathbf{H} = \{\langle h \rangle : h \in W, h \neq \theta\}$, and ξ be the form defined on \mathbb{V} which determines the conjugacy \bot . Write $\mathbf{b} = \langle e_0 \rangle$ for a vector e_0 . There are two cases to consider.

b \notin **H**: Let ξ_0 be the restriction of ξ to W. Then ξ_0 is a nondegenerate symplectic form. Write $\varepsilon = \xi(e_0, e_0)$. Let e_1, \ldots, e_n be a basis of **H**; then the family $\mathcal{E} = (e_i : i = 0, \ldots, n)$ is a basis of \mathbb{V} . We have $e_0 \perp h$ for every $h \in \mathbf{H}$ and thus the formula defining the form ξ is the following

$$\xi(h_1 + \alpha_1 e_0, h_2 + \alpha_2 e_0) = \xi_0(h_1, h_2) + \alpha_1 \alpha_2 \varepsilon, \tag{9}$$

where $h_1, h_2 \in W$ and α_1, α_2 are scalars of the coordinate field. Note that, conversely, for every nondegenerate symplectic form ξ_0 defined on W and every scalar

 $\varepsilon \neq 0$ the formula (9) defines a nondegenerate bilinear form $\xi = \xi_{\varepsilon}$. Indeed, if M is the matrix of ξ_0 in the given basis then

$$M_{\varepsilon} = \begin{bmatrix} \varepsilon & 0 & \cdots & 0 \\ 0 & & \\ \vdots & M & \\ 0 & & \end{bmatrix}$$

is the matrix of ξ_{ε} and $\det(M_{\varepsilon}) \neq 0$. In particular, for $h, h_1 \in W$ and a scalar α we have

$$\xi_{\varepsilon}(h_1, h + \alpha e_0) = \xi_0(h_1, h)$$
 and $\xi_{\varepsilon}(h + \alpha e, h + \alpha e_0) = \alpha^2 \varepsilon_1$

Let us write \perp_{ε} for the conjugacy determined by ξ_{ε} and $\mathfrak{B}_{\varepsilon}$ for the induced structure of regular points and lines wrt. the conjugacy \perp_{ε} . Let $\varepsilon_1, \varepsilon_2$ be any two nonzero scalars. Then the following holds

$$q' \perp_{\varepsilon_1} q'' \iff q' \perp_{\varepsilon_2} q'', \tag{10}$$

$$q' \perp_{\varepsilon_1} a \iff q' \perp_{\varepsilon_2} a, \tag{11}$$

for all $q', q'' \in \mathbf{H}$, $a \notin \mathbf{H}$. From (9), (10) and (11) we derive that the set of points selfconjugate under ξ_{ε_i} is \mathbf{H} , and a line of \mathfrak{A} is regular under \perp_{ε_1} iff it is regular under \perp_{ε_2} . This yields $\mathfrak{B}_{\varepsilon_1} = \mathfrak{B}_{\varepsilon_2}$.

Let us take any $h_1, h_2 \in W$ with $\xi_0(h_1, h_2) \neq 0$. Set $\varepsilon_1 = \xi_0(h_1, h_2)$, $a_i = \langle h_i + e_0 \rangle$ for i = 1, 2, and let ε_2 be a nonzero scalar $\neq \varepsilon_1$. From (9) we directly compute that $a_1 \perp_{\varepsilon_1} a_2$ and $a_1 \not\perp_{\varepsilon_2} a_2$. This yields that \perp_{ε_i} cannot be defined in terms of $\mathfrak{B}_{\varepsilon_i}$.

b \in **H**: Let $\omega \notin W$. The form $\xi \upharpoonright W$ is a degenerate symplectic form. Set $Y = \omega^{\perp} \cap W$, then $e_0, \omega \notin Y$. Let e_1, \ldots, e_{n-1} be a basis of Y; then (e_0, \ldots, e_{n-1}) is a basis of W and $(\omega, e_0, \ldots, e_{n-1})$ is a basis of \mathbb{V} . Let $y \in Y$, then

$$\xi(y,\omega) = 0, \qquad \xi(y,e_0) = 0, \qquad \text{and} \qquad \xi(e_0,e_0) = 0.$$

Take any two vectors $y_i + \alpha_i e_0 + \beta_i \omega$, $(y_i \in Y, i = 1, 2)$ of \mathbb{V} . We have

$$\xi(y_1 + \alpha_1 e_0 + \beta_1 \omega, y_2 + \alpha_2 e_0 + \beta_2 \omega) = \xi_0(y_1, y_2) + (\alpha_1 \beta_2 + \alpha_2 \beta_1)\lambda + \beta_1 \beta_2 \mu, \quad (12)$$

where $\lambda = \xi(e_0, \omega) \neq 0$, $\mu = \xi(\omega, \omega) \neq 0$, and ξ_0 is the restriction of ξ to Y.

For any scalars $\lambda, \mu \neq 0$ let $\xi_{\mu,\lambda}$ be a bilinear form defined on \mathbb{V} by formula (12). Let M be the matrix of ξ_0 in the given basis. Note that ξ_0 is a nondegenerate symplectic form. Then

$$M_{\mu,\lambda} = \begin{bmatrix} \mu & \lambda & 0 & \cdots & 0 \\ \lambda & 0 & 0 & \cdots & 0 \\ 0 & 0 & & & \\ \vdots & \vdots & & M \\ 0 & 0 & & & \end{bmatrix}$$

is the matrix of $\xi_{\mu,\lambda}$ in our basis. Clearly, $\det(M_{\mu\lambda}) \neq 0$, so $\xi_{\mu,\lambda}$ is nondegenerate. Let $\perp_{\mu,\lambda}$ be the conjugacy determined by the form $\xi_{\mu,\lambda}$ and $\mathfrak{B}_{\mu,\lambda}$ be the induced structure of regular points and lines. Since $\xi_{\mu,\lambda}(\omega,\omega) \neq 0$, the form $\xi_{\mu,\lambda}$ is not symplectic. Easy computation gives that $\xi_{\mu,\lambda}(y + \alpha e_0, y + \alpha e_0) = 0$ for each $y \in Y$ and each scalar α . Consequently, **H** is the set of points selfconjugate under $\perp_{\mu,\lambda}$. From (12) we compute

$$\begin{aligned} \xi_{\mu,\lambda}(y + \alpha e_0, y_1 + \alpha_1 e_0) &= \xi_0(y, y_1) \text{ and} \\ \xi_{\mu,\lambda}(y_1 + \alpha_1 e_0, y + \alpha e_0 + \beta \omega) &= \xi_0(y_1, y) + \lambda \alpha_1 \beta \end{aligned}$$

for all $y, y_1 \in Y$ and scalars α, α_1, β . This yields that for any nonzero scalars $\lambda, \lambda_1, \mu, \mu_1$ and $q', q'' \in \mathbf{H}, a \notin \mathbf{H}$ we have

$$q' \perp_{\mu,\lambda} q'' \iff q' \perp_{\mu_1,\lambda_1} q'', \tag{13}$$

$$q' \perp_{\mu,\lambda} a \iff q' \perp_{\mu_1,\lambda} a.$$
 (14)

Let us take any $y_1, y_2 \in Y$ with $\xi_0(y_1, y_2) \neq 0$ and let $\mu_1 = \xi_0(y_1, y_2)$. Write $a_i = \langle y_i + \omega \rangle$. Finally, let $\mu_2 \neq 0, \mu_1$ be a scalar. Then $\mathfrak{B}_{\mu_1,\lambda} = \mathfrak{B}_{\mu_2,\lambda}, a_1 \perp_{\mu_1,\lambda} a_2$, and $a_1 \not\perp_{\mu_2,\lambda} a_2$. Consequently, $\perp_{\mu_i,\lambda}$ cannot be defined in $\mathfrak{B}_{\mu_i,\lambda}$.

Since in any case, the conjugacy \perp cannot be defined in terms of the geometry of \mathfrak{B}_1 , our proof is complete.

Gathering together 3.13 and 3.14 we conclude with

COROLLARY 3.16. The structure of the form

 $\langle \text{points of } \mathfrak{A}, \text{lines of } \mathfrak{A}, \text{points of } \mathfrak{P}, \text{lines of } \mathfrak{P}, \mathsf{H} \ (= \text{the horizon of } \mathfrak{A}), \perp_{\mathsf{H}}, \\ \perp \cap (\mathfrak{R}_1 \times \mathsf{H}) \rangle$

is definable in terms of the structure \mathfrak{B}_1 .

Finally, taking into account 2.4 from 3.16 we obtain

PROPOSITION 3.17. For each integer k the family \mathcal{R}_k is definable in terms of the structure \mathfrak{B}_1 . Consequently, the family

 $\{A: A \text{ is a subspace of } \mathfrak{A}, \overline{A} \in \mathfrak{R}_k\}$

is definable in \mathfrak{B}_1 as well.

Then from 3.15 we conclude with

COROLLARY 3.18. Let $1 \leq k < \dim(\mathbb{V})$ be an integer. The underlying metric projective space (\mathfrak{P}, \bot) cannot be defined neither in terms of the structure $\mathbf{G}_k(\mathfrak{R}) = \langle \mathfrak{R}_k, \mathfrak{R}_{k+1}, \subset \rangle$ nor in terms of the structure $\mathbf{P}_k(\mathfrak{R}) = \langle \mathfrak{R}_k, \mathfrak{G}_k(\mathfrak{R}) \rangle$, where $\mathfrak{G}_k(\mathfrak{R})$ is the family of regular k-pencils.

In view of 3.18, the two geometries: of regular subspaces of (\mathfrak{P}, \perp) and of the metric projective space (\mathfrak{P}, \perp) , are distinct.

3.2 More regular point-line geometry

Now, let us have a look at the incidence structure $\langle \mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_3 \rangle$. From 2.7 and 2.8 we see that

FACT 3.19. The lines through **b** are either nonregular, when $\mathbf{b} \notin \mathbf{H}$, or isolated as there are no regular planes containing such lines when $\mathbf{b} \in \mathbf{H}$.

This is the reason to investigate a new lineset

 $\mathcal{L}_{\mathrm{r}} := \mathcal{A}_2 \cap \mathcal{A}_2^{\circ}$

of regular affine lines not through **b**. This lineset gives rise to a new geometry

 $\mathfrak{C}_1 := \langle \mathcal{A}_1^{\circ}, \mathcal{L}_r, \subset \rangle,$

which is a substructure of \mathfrak{B}_1 (and of \mathfrak{G}_1). This slight difference between \mathfrak{B}_1 and \mathfrak{C}_1 has no impact on the validity of 3.9, 3.10 and 3.13 with \mathfrak{B}_1 replaced by \mathfrak{C}_1 . The respective proofs for \mathfrak{C}_1 become a bit more complex but are based on the same ideas. Actually we can state even more:

THEOREM 3.20. The structures \mathfrak{B}_1 and \mathfrak{C}_1 are mutually definable. Consequently, the affine space \mathfrak{A} , and thus the projective space \mathfrak{P} , is definable in terms of \mathfrak{C}_1 .

3.3 Automorphism group of regular point-line geometry

In view of 3.13, $\operatorname{Aut}(\mathfrak{B}_1)$ is a subgroup of $\operatorname{Aut}(\mathfrak{A})$. Let $f \in \operatorname{Aut}(\mathfrak{A})$ and let f^{∞} be its action on the horizon \mathbf{H} of \mathfrak{A} .

If $f \in Aut(\mathfrak{B}_1)$ then f^{∞} must be an automorphism of the induced metric projective symplectic geometry on \mathbf{H} .

Moreover, in view of 3.14, f must preserve the family of hyperplanes $\{[q]: q \in \mathbf{H}\}$. The following is simple, though quite useful.

LEMMA 3.21. Let $f \in Aut(\mathfrak{A})$. The following conditions are equivalent.

- (i) $f \in \operatorname{Aut}(\mathfrak{B}_1)$
- (ii) $f^{\infty} \in \operatorname{Aut}(\langle \mathbf{H}, \bot_{\mathbf{H}} \rangle)$ and (f, f^{∞}) preserves $\bot \cap (\mathfrak{R}_1 \times \mathbf{H})$.

PROOF. Immediate by 3.16.

PROPOSITION 3.22. Let W be the subspace of \mathbb{V} with $\mathbf{H} = \{\langle u \rangle : u \in W, u \neq \theta\}$ and let ξ_0 be the restriction of ξ to W. Then, clearly, ξ_0 determines $\perp_{\mathbf{H}}$. Assume that $\mathbf{b} \notin \mathbf{H}$. Then

$$\operatorname{Aut}(\mathfrak{B}_1) = \{ \varphi \in \Gamma L(W) \colon \varphi \text{ preserves } \bot_{\mathsf{H}} \}.$$
(15)

PROOF. Note that \mathfrak{A} can be presented as the affine space $\mathbf{A}(W)$ over W. **b** is the unique point of \mathfrak{A} such that each line through it is nonregular and thus **b** remains invariant under automorphisms of \mathfrak{B}_1 . One can coordinatize W so as **b** is the origin of the coordinate system and thus each automorphism φ of \mathfrak{B}_1 is a semilinear bijection of W. Since **H** is the horizon of \mathfrak{A} , from 3.14 we get that φ preserves $\perp_{\mathbf{H}}$. A direct computation based on (9) justifies that if $\varphi \in \Gamma L(W)$ preserves the conjugacy defined on **H** by the symplectic form ξ_0 then φ preserves the class of regular lines. This closes our proof.

The technique used in the proof of 3.15 enables us to formulate a more elementary definition of the structure \mathfrak{B}_1 .

PROPOSITION 3.23. Let W be the subspace of \mathbb{V} with $\mathbf{H} = \{\langle u \rangle : u \in W, u \neq \theta\}$ and let ξ_0 be the restriction of ξ to W. Then, clearly, ξ_0 determines $\perp_{\mathbf{H}}$. Moreover, one can represent \mathfrak{B}_1 as a line reduct of the affine space $\mathfrak{A} = \mathbf{A}(W)$ and \mathbf{H} is the horizon of \mathfrak{A} . Assume that $\mathbf{b} \notin \mathbf{H}$; consequently, \mathbf{b} is a point of \mathfrak{B}_1 .

(i) Let \mathcal{L}_* be the class of lines in the symplectic polar space determined by the conjugacy $\perp_{\mathbf{H}}$ on \mathbf{H} , let a be a point of \mathfrak{A} , and \mathcal{P}^a_* be the set of planes of the form a + L with $L \in \mathcal{L}_*$. Finally, write $\mathcal{L}^a_* = \{L \in \mathcal{L} : L \subset A \text{ for some } A \in \mathcal{P}^a_*\}$. Then $\mathcal{A}_2 = \mathcal{L} \setminus \mathcal{L}^{\mathbf{b}}_*$. Consequently, for an arbitrary affine point a we have $\mathfrak{B}_1 \cong \langle W, \mathcal{L} \setminus \mathcal{L}^a_* \rangle$.

(ii) Let $u, v \in W$. Then u, v lie on a nonregular line iff $\xi_0(u, v) = 0$.

Now, we pass to the case $\mathbf{b} \in \mathbf{H}$. Let us adopt the coordinate system as in subs. 1.4.1 and let ξ_0 be the restriction of ξ to W.

PROPOSITION 3.24. The following conditions are equivalent.

(i) $f \in \operatorname{Aut}(\mathfrak{B}_1)$

(ii) There are $\varphi \in \Gamma L(W)$ and a vector $\omega \in W$ such that $f(x) = \varphi(x) + \omega$ for each $x \in W$, φ preserves $\bot_{\mathbf{H}}$, and the following holds

$$\xi_0(x,y) = \pi_1(y) \implies \xi_0(\varphi(x),\varphi(y)) + \xi_0(\omega,\varphi(y)) = \pi_1(\varphi(y)),$$

for all $x, y \in W$, where π_1 is the projection on the 1st coordinate.

PROOF. It is clear that each automorphism f of \mathfrak{B}_1 is a composition of a semilinear map φ and a translation on a vector ω . In the projective coordinates we can write $f([1,x]) = [1,\varphi(x) + \omega]$ and $f([0,y]) = [0,\varphi(y)]$. The map f of such a form is an automorphism of \mathfrak{B}_1 iff it preserves \perp_{H} and it preserves regular lines. From 3.21, f preserves regular lines iff it preserves suitable restriction of the polarity. To complete the proof it suffices to note that

$$\xi([1,x],[0,y]) = \pi_1(y) + \xi_0(x,y)$$

for all $x, y \in W$.

Suppose that $\varphi \in GL(W)$ and $\xi_0(\varphi(x), \varphi(y)) = c\xi_0(x, y)$ for some $c \neq 0$ and all $x, y \in W$ (then, clearly, φ preserves $\perp_{\mathbf{H}}$). The conditions of 3.24 yield $\xi_0(\omega, \varphi(y)) = \pi_1(cy + \varphi(y))$ for each $y \in W$. In particular, if $\varphi = \mathrm{id}$ we obtain $\omega \perp W$ and thus $\omega \parallel \mathbf{b}$. If φ is a homothety $x \mapsto \alpha x$ with $\alpha \neq 0$ then $c = \alpha^2$ and the condition of 3.24 is read as $\xi_0(\omega, y) = (\alpha + 1)\pi_1(y)$ for all $y \in W$; this yields $\omega \perp \{y: \pi_1(y) = 0\}$.

4 Grassmannians of regular secunda and hyperplanes

Note that $\operatorname{Rad}(U) = \operatorname{Rad}(U^{\perp})$. This yields that

Remark 1. The mapping \perp is a correlation in \mathfrak{P} which maps regular subspaces to regular subspaces.

Now set $n := \dim(\mathbb{V}) = \dim(\mathfrak{P}) + 1$. In this case $\mathcal{R}_k^{\perp} = \mathcal{R}_{n-k}$ and, clearly, $\mathcal{A}_k^{\perp} = \mathcal{A}_{n-k}^{\circ}$ for each $k, 1 \leq k \leq n$. Consider the Grassmannian of regular secunda and hyperplanes

$$\mathfrak{G}_{n-2} := \mathbf{G}_{n-2}(\mathfrak{R}) = \langle \mathfrak{R}_{n-2}, \mathfrak{R}_{n-1}, \subset \rangle.$$

It can be easily seen that $\mathfrak{G}_1 \cong \mathfrak{G}_{n-2}^{\perp}$. So, based on 3.13 we can state that the dual of the affine space \mathfrak{A} can be defined in terms of \mathfrak{G}_{n-2} . In the dual of \mathfrak{A} we can define \mathfrak{P} as well as in \mathfrak{A} . These observation can be summarized in the following

THEOREM 4.1. The structure \mathfrak{B}_1 can be reconstructed in terms of \mathfrak{G}_{n-2} .

The case where n = 4, i.e. $\dim(\mathfrak{P}) = 3$, seems quite interesting. Then **H** is a plane and $\mathbf{b} \in \mathbf{H}$. In this case the Grassmannian \mathfrak{G}_1 of regular points and lines is dual isomorphic to the Grassmannian \mathfrak{G}_2 of regular lines and planes, i.e. $\mathfrak{G}_1 \cong \mathfrak{G}_2^{\perp}$. Since $\mathcal{A}_2^{\perp} = \mathcal{A}_2^{\circ}$, we have $\mathfrak{B}_1^{\perp} \cong \langle \mathcal{A}_2^{\circ}, \mathcal{R}_3, \subset \rangle$. The latter structure is studied in the next section for arbitrary n.

5 Grassmannians of regular lines and planes

It is a quite complex, though more or less routine, job to define \mathfrak{B}_1 in terms of $\mathbf{P}_k(\mathfrak{R})$ or $\mathbf{G}_k(\mathfrak{R})$. In this section we shall discuss one particular case of this problem where k = 2. It seems, however, that the techniques used here can be applied generally.

Consider the Grassmannian of regular lines and planes

$$\mathfrak{G}_2 := \mathbf{G}_2(\mathfrak{R}) = \langle \mathfrak{R}_2, \mathfrak{R}_3, \subset \rangle.$$

In view of 3.17 the structure \mathfrak{G}_2 is definable in \mathfrak{B}_1 . Note by 3.19 that there are isolated points in \mathfrak{G}_2 iff $\mathbf{b} \in \mathbf{H}$. When we get rid of these isolated points we get a new structure

$$\mathfrak{B}_2:=\langle \mathcal{A}_2^{\circ}, \mathfrak{R}_3, \subset
angle$$

of the regular lines not through **b** and regular planes, a substructure of \mathfrak{G}_2 . Note that \mathfrak{B}_2 is the dual of \mathfrak{B}_1 when dim $(\mathfrak{P}) = 3$.

As we already know the incidence structure $\langle \mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_3 \rangle$ of regular points, lines and planes contains isolated objects: the point **b**, when **b** \notin **H**, regular lines on **H** and regular lines through **b** when **b** \in **H**. So, now we introduce the structure

$$\mathfrak{C}_2 := \langle \mathcal{L}_r, \mathfrak{R}_3, \subset \rangle$$

of the regular affine lines not through **b** and regular planes, a substructure of \mathfrak{B}_2 . Note that, when dim(\mathfrak{P}) = 3, we have $\mathcal{L}_r^{\perp} = \mathcal{L}_r$ and thus \mathfrak{C}_2 is the dual of \mathfrak{C}_1 .

Note an evident (cf. 2.9)

Remark 2. There is no regular plane on **H**.

This means that the points and lines of \mathfrak{C}_2 are respectively lines and planes of the affine space $\mathfrak{A} = \langle \mathcal{R}_1, \mathcal{L} \rangle$ obtained from \mathfrak{P} by deleting the hyperplane H.

Recall a known fact

FACT 5.1. In the structure $\mathbf{G}_2(\mathfrak{P}) = \langle \mathfrak{H}_2, \mathfrak{H}_3, \subset \rangle$ consider a triangle with the vertices L_0, L_1, L_2 and the sides A_0, A_1, A_2 labelled so as $L_i \not\subset A_i$ for i = 0, 1, 2. Then the lines L_i have a common point p. Moreover

$$\{L \in \mathcal{H}_2 \colon L \subset A_0 \land \exists A \in \mathcal{H}_3 [L, L_0 \subset A]\} = \{L \in \mathcal{H}_2 \colon p \in L \subset A_0\} = =: \mathbf{p}(p, A_0) \in \mathcal{G}_2(\mathfrak{P}), \quad (16)$$

where $\mathfrak{G}_2(\mathfrak{P})$ is the set of projective planar pencils of lines of \mathfrak{P} . Consequently, the relation

defined for arbitrary lines L_1, L_2, L_3 coincides with the collinearity relation in the space $\mathbf{P}_2(\mathfrak{P})$ of pencils of lines.

The following is evident by 3.6 and inspection of possible cases.

LEMMA 5.2. Let $A \in \mathbb{R}_3$ and p be a point on A. Then $\mathbf{q}(A)$ is an affine point and

$$\mathbf{p}(p,A) \cap \mathcal{A}_{2}^{\circ} = \begin{cases} \emptyset, & \text{when } p = \mathbf{q}(A), \\ \mathbf{p}(p,A) \setminus \{\overline{p, \mathbf{q}(A)}\}, & \text{when } p \neq \mathbf{q}(A). \end{cases}$$
$$\mathbf{p}(p,A) \cap \mathcal{L}_{r} = \begin{cases} \emptyset, & \text{when } p = \mathbf{q}(A), \\ \mathbf{p}(p,A) \setminus \{\overline{p, \mathbf{q}(A)}\}, & \text{when } p \neq \mathbf{q}(A) \text{ and } p \notin A^{\infty}, \\ \mathbf{p}(p,A) \setminus \{\overline{p, \mathbf{q}(A)}, A^{\infty}\}, & \text{when } p \in A^{\infty}, \end{cases}$$

Consequently, if $p \notin A^{\infty}$ (or equivalently $p \notin \mathbf{H}$), then $\mathbf{p}(p, A) \cap \mathcal{A}_2 = \mathbf{p}(p, A) \cap \mathcal{R}_2$. Moreover, $\mathbf{p}(p, A) \cap \mathcal{R}_2 = \mathbf{p}(p, A) \cap \mathcal{A}_2^{\circ}$ and $\mathbf{p}(p, A) \cap \mathcal{A}_2 = \mathbf{p}(p, A) \cap \mathcal{L}_r$ as $\mathbf{b} \notin A^{\infty}$.

LEMMA 5.3. Let the relation $\mathbf{L}_{\mathfrak{C}_2}$ be defined by the formula (17) with \mathfrak{H}_2 replaced by \mathcal{L}_r , \mathfrak{H}_3 replaced by \mathfrak{R}_3 , and let $L_1, L_2, L_3 \in \mathcal{L}_r$. Then the relation $\mathbf{L}_{\mathfrak{C}_2}(L_1, L_2, L_3)$ holds iff $L_1, L_2, L_3 \in \mathbf{p}(p, A)$ for some point p and some $A \in \mathfrak{R}_3$.

PROOF. \Rightarrow : Straightforward by 5.1.

 \Leftarrow : Let $A \in \mathcal{R}_3$, p be a point on A, and $L_1, L_2, L_3 \in \mathcal{L}_r$ be three lines on A through p. Set $M := A^{\infty}$; then $M \in \mathcal{R}_2$. To close the proof we need to find a line $L_0 \in \mathcal{L}_r$ through p but not on A such that the planes $A_1 = L_0 + L_1$, $A_2 = L_0 + L_2$ and $A_3 = L_0 + L_3$ are all regular. There are two cases to consider.

(i) p is an affine point i.e. $p \notin M$.

Note that $M \cap M^{\perp} = \emptyset$. Write $q_i := M \cap L_i$ for i = 1, 2, 3. Since the L_i are regular, $q_i \notin p^{\perp}$ and thus $p \not\perp M$. Therefore, the intersection $M \cap p^{\perp}$ is a single point $q_0 \neq q_1, q_2, q_3$.

Take $x \in M$ with $x \neq q_i$ for i = 0, ..., 3 and $y \in M^{\perp} \cap \mathbf{H} + x$ with $y \neq x$ and $y \notin M^{\perp}$. Then $y \neq \mathbf{b}$ and $y \in x + m = \overline{x, m}$ for some (uniquely determined by y) point $m \in M^{\perp}$. We have $y \notin M$, since otherwise $M = \overline{x, y} \ni m$, so $m \in \text{Rad}(M)$.

Suppose that $y \in q_i^{\perp}$ for some i = 1, 2, 3. Evidently, $q_i \perp M^{\perp}$ and $q_i \perp q_i$; comparing dimensions we get $q_i^{\perp} = q_i + M^{\perp}$. From $y \in q_i^{\perp}$ we get that $y \in q_i + m_i$ for some $m_i \in M^{\perp}$. Since M and M^{\perp} are skew, we get $x = q_i$, which is impossible.

Suppose that $y \in p^{\perp}$ and write $K = y + q_0$; then $K \subset p^{\perp}$. For each i = 1, 2, 3 the line K crosses $q_i + M^{\perp}$ in a point y_i and $y_i \neq y$. Note: $y_i \perp p$. There are $m_i \in M^{\perp}$ such that $y_i \in q_i + m_i$; it is seen that $m_i = m$, as otherwise the lines M and $\overline{m, m_i} \subset M^{\perp}$ are contained in the plane M + K and thus they have a common

point. Consider another point y' on m + x. If there were $y' \in p^{\perp}$ we would obtain another point y'_1 on $m + q_1$ with $p \perp y'_1$. This leads, contradictory, to $q_i \perp p$. Without loss of generality we can assume that $y \notin p^{\perp}$.

Take $L_0 := \overline{p, y}$; then $L_0 \in \mathcal{L}_r$ as p is an affine point. Take $A_i := L_0 + L_i$ for i = 1, 2, 3. Then $A_i^{\infty} = \overline{q_i, y}$. Since $q_i \not\perp y$, the lines A_i^{∞} are regular and thus $A_i \in \mathcal{R}_3$.

(ii) $p \in M, M \neq L_1, L_2, L_3$ i.e. all the L_i are affine lines.

By assumption, M is regular. Let D be a plane on \mathbf{H} containing the line M. By 2.9 the radical of D is a point, say u. The line $L := \overline{p, u}$ is nonregular. Now, in the three space A + D we take a line L_0 through p that is not contained in any of the planes $L_i + L$ and is not contained in p^{\perp} . This is doable thanks to assumption that there are at least 5 lines distinct from M through p on every plane containing M. Note that L_0 is regular by 2.5.

Let $A_i := L_i + L_0$. By 2.8 each of the planes A_i is regular as it meets D in a line distinct from L thus, in a regular one.

This completes the reasoning.

COROLLARY 5.4. Let the relation $\mathbf{L}_{\mathfrak{B}_2}$ be defined by the formula (17) with \mathfrak{H}_2 , \mathfrak{H}_3 replaced by \mathcal{A}_2° , \mathfrak{R}_3 respectively and let $L_1, L_2, L_3 \in \mathcal{A}_2^{\circ}$. The relation $\mathbf{L}_{\mathfrak{B}_2}(L_1, L_2, L_3)$ holds iff $L_1, L_2, L_3 \in \mathbf{p}(p, A)$ for some regular point p and some $A \in \mathfrak{R}_3$.

PROOF. \Rightarrow : Clear by 5.1.

 \Leftarrow : Let $A \in \mathcal{R}_3$, p be a point on A, and $L_1, L_2, L_3 \in \mathcal{A}_2^\circ$ be three lines on A through p. Set $M := A^{\infty}$; then $M \in \mathcal{A}_2^\circ$. We have three cases

- (i) p is an affine point i.e. $p \notin M$,
- (ii) $p \in M, M \neq L_1, L_2, L_3$ i.e. all the L_i are affine lines,
- (iii) $p \in M = L_i$ for some $i \in \{1, 2, 3\}$.

Cases (i) and (ii) follow directly by 5.3 (though in case (ii) there is a simple independent proof here that do not require 6 lines in a projective pencil). In the remaining case (iii) without loss of generality we take i = 1. Let D be a plane through Mcontained in **H**. By 2.9, since D contains a regular line M, $\operatorname{Rad}(D)$ is a point q. If there was p = q we would have $p \perp D$, which gives, contradictory, $p \in \operatorname{Rad}(L_1)$. The unique nonregular line through p on A is $K_1 := \overline{p, q(A)}$ and the unique nonregular line through p on D is $K_2 := \overline{p, q}$. Write $A_0 := K_1 + K_2$; since $p \perp K_1, K_2$ we have $p \perp A_0$. Set Y := D + A; then $Y \in \mathcal{H}_4$. Note that $\operatorname{rdim}(Y) \leq 1$; if $\operatorname{Rad}(Y) \neq \emptyset$ then $\operatorname{Rad}(Y) = q$. Since, either $Y \subset p^{\perp}$, which gives, contradictory, $p \perp L_1$, or $Y \cap p^{\perp}$ is a plane, we obtain $Y \cap p^{\perp} = A_0$.

Take $M_3 \subset D$ with $p \in M_3 \neq K_2, M$; then M_3 is regular. For $A_3 = L_3 + M_3$ we have $A_3 \in \mathcal{R}_3$, because $A_3^{\infty} = M_3$. From $p \in A_3, A_0 \subset Y$ we get that $K_0 := A_3 \cap A_0$ is a line.

Consider the plane $B = L_2 + K_2$; then B and A_3 have a common line K_4 . Let L_0 be a line in $\mathbf{p}(p, A_3)$ distinct from L_3, K_4, K_0, M_3 . Then $L_0 \not\subset A_0$ and thus L_0 is regular. Set $A_2 = L_0 + L_2$; then $p \in A_2^{\infty}$ and $A_2^{\infty} = A_2 \cap \mathbf{H} = A_2 \cap D \neq K_2$, which gives that A_2 is regular. Finally, we take $A_1 = L_0 + L_1$. Then $A_1^{\infty} = M$ and thus $A_1 \in \mathcal{R}_3$.

The family of equivalence classes of the relation $\mathbf{L}_{\mathfrak{C}_2}$ is the set

 $\{\mathbf{p}_{\mathbf{r}}^{\mathrm{af}}(p,A): p \in A \in \mathcal{R}_3\} \setminus \{\emptyset\}, \text{ where } \mathbf{p}_{\mathbf{r}}^{\mathrm{af}}(p,A) = \{L \in \mathcal{L}_{\mathbf{r}}: p \in L \subset A\}.$

So, we get the space of affine regular pencils $\mathbf{A}_2(\mathcal{R})$ with regular affine lines not through **b**, i.e. elements of \mathcal{L}_r , as points and affine regular pencils $\mathbf{p}_r^{\mathrm{af}}(p, A)$ as lines. Note that among pencils $\mathbf{p}_r^{\mathrm{af}}(p, A)$ we have proper pencils, those with $p \notin \mathbf{H}$, and parallel pencils, those with $p \in \mathbf{H}$.

The family of equivalence classes of the relation $\mathbf{L}_{\mathfrak{B}_2}$, is the set

 $\{\mathbf{p}_{\mathbf{r}}(p,A) \colon p \in A \in \mathfrak{R}_3\} \setminus \{\emptyset\}, \text{ where } \mathbf{p}_{\mathbf{r}}(p,A) = \{L \in \mathcal{A}_2^\circ \colon p \in L \subset A\},\$

and we have the space of regular pencils $\mathbf{P}_2(\mathcal{R})$ with regular lines not through **b** as points and regular pencils $\mathbf{p}_r(p, A)$ as lines. The space of affine regular pencils $\mathbf{A}_2(\mathcal{R})$ is a substructure of the space of regular pencils $\mathbf{P}_2(\mathcal{R})$ in the sense that points of $\mathbf{A}_2(\mathcal{R})$ are points of $\mathbf{P}_2(\mathcal{R})$ and, in view of 5.2, lines of $\mathbf{P}_2(\mathcal{R})$ are a bit "richer".

Loosely speaking, we have proved that the family of regular pencils is definable in both \mathfrak{C}_2 and in \mathfrak{B}_2 (cf. [9]).

Recall that 2.7 and 2.8 together say the following.

FACT 5.5. Let $\mathbf{b} \in \mathbf{H}$. If A is a plane not contained in \mathbf{H} and $\mathbf{b} \in A$, then A is not regular (comp. 2). Consequently, no regular pencil exists that contains a line through $\mathbf{b} \in \mathbf{H}$.

This means that there are no pencils, neither in $\mathbf{A}_2(\mathcal{R})$ nor in $\mathbf{P}_2(\mathcal{R})$, with a vertex **b** when $\mathbf{b} \in \mathbf{H}$, though there are regular affine lines through **b**.

For points L_1, L_2, L_3 of a point-line geometry X we write

$$\Delta_X(L_1, L_2, L_3) \iff L_1, L_2, L_3$$
 are the vertices of a triangle in X. (18)

From common projective geometry (cf. 5.1) it follows that if $\Delta_X(L_1, L_2, L_3)$ holds and X is \mathfrak{G}_2 , \mathfrak{B}_2 or \mathfrak{C}_2 and $p \in L_1, L_2$, then $p \in L_3$ as well.

We are going to identify points of \mathfrak{B}_1 with stars of lines in \mathfrak{B}_2 as well as with stars of lines in \mathfrak{C}_2 . The star of regular lines through a point p is the set

$$S_{\mathbf{r}}(p) = \{ L \in \mathcal{R}_2 \colon p \in L \}$$

and the star of regular affine lines through a point p is

$$S_{\mathbf{r}}^{\mathrm{af}}(p) = \{ L \in \mathcal{A}_2 \colon p \in L \}.$$

Note that

$$S_{\mathbf{r}}(p) = \begin{cases} S_{\mathbf{r}}^{\mathrm{af}}(p), & p \notin \mathbf{H}, \\ S_{\mathbf{r}}^{\mathrm{af}}(p) \cup \{L \in \mathcal{R}_{2} \colon p \in L \subset \mathbf{H}\}, & p \in \mathbf{H}. \end{cases}$$

To express the notion of a star of lines purely in terms of the geometry of \mathfrak{B}_2 or \mathfrak{C}_2 for a given pencil $\mathcal{Q} = \mathbf{p}_r(a, A) \neq \emptyset$ (and respectively for $\mathcal{Q} = \mathbf{p}_r^{\mathrm{af}}(a, A) \neq \emptyset$) we write

$$\begin{split} \mathbf{S}_{\Delta}(\mathcal{Q}) &:= \{ L \in \mathcal{R}_{2} \colon (\exists \ L_{1}, L_{2} \in \mathcal{Q}) \ \mathbf{\Delta}(L, L_{1}, L_{2}) \}, \\ \mathbf{S}_{\mathrm{L}}(\mathcal{Q}) &:= \{ L \in \mathcal{R}_{2} \colon (\exists \ L', L'' \in \mathbf{S}_{\Delta}(\mathcal{Q})) \ [L' \neq L'' \land \mathbf{L}(L, L', L'')] \}, \\ \mathbf{S}(\mathcal{Q}) &:= \mathcal{Q} \cup \mathbf{S}_{\Delta}(\mathcal{Q}) \cup \mathbf{S}_{\mathrm{L}}(\mathcal{Q}), \end{split}$$

where $\Delta = \Delta_{\mathfrak{B}_2}$ and $\mathbf{L} = \mathbf{L}_{\mathfrak{B}_2}$ (or respectively $\Delta = \Delta_{\mathfrak{C}_2}$ and $\mathbf{L} = \mathbf{L}_{\mathfrak{C}_2}$). The set $S_{\Delta}(\mathcal{Q})$ contains those lines L with $L^{\infty} \not\perp A^{\infty}$ while we need $S_{\mathrm{L}}(\mathcal{Q})$ for the other lines L with $L^{\infty} \perp A^{\infty}$. It will become more apparent later in 5.6 and 5.7.

Let ~ be the binary collinearity in \mathfrak{G}_2 , i.e. for $L_1, L_2 \in \mathfrak{R}_2$ we write

$$L_1 \sim L_2$$
 iff there is $A \in \mathcal{R}_3$ with $L_1, L_2 \subset A$, and
 $L_1 \not\sim L_2$ when $L_1 \sim L_2$ does not hold.

Following 2.7 recall that

$$S_{r}^{af}(\mathbf{b}) = S_{r}(\mathbf{b}) \begin{cases} = \emptyset \text{ if } \mathbf{b} \notin \mathbf{H}; & \text{there are } L_{1}, L_{2} \in S_{r}(\mathbf{b}) \text{ with } L_{1} \sim L_{2}, L_{1} \neq L_{2}, \\ \neq \emptyset \text{ if } \mathbf{b} \in \mathbf{H}; & L_{1} \not\sim L_{2} \text{ for all } L_{1}, L_{2} \in S_{r}(\mathbf{b}). \end{cases}$$

LEMMA 5.6. Let $\mathcal{Q} = \mathbf{p}_r(p, A) \neq \emptyset$ with $p \in A \in \mathcal{R}_3$ and $p \notin \mathbf{H}$, and let L be a regular line not in \mathcal{Q} . Write $q := L^{\infty}$ and $M := A^{\infty}$.

If $p \in L$ and $q \neq \mathbf{b}$, then one of the following holds:

(i) $q \notin M^{\perp}$. Then there are distinct $L_1, L_2 \in \mathcal{Q}$ such that $\Delta_{\mathfrak{B}_2}(L, L_1, L_2)$.

(ii) $q \in M^{\perp}$. Then there are distinct regular lines L', L'' through q such that $\mathbf{L}_{\mathfrak{B}_2}(L, L', L'')$ and $L'^{\infty}, L''^{\infty} \notin M^{\perp}$, so for both L' and L'' the condition (i) holds.

Conversely, if a line L satisfies (i) or (ii) then $p \in L$.

PROOF. Note that M is a regular line and $x_1 := p^{\perp} \cap M$ is a point, since otherwise $p \perp M$ and then no line in $\mathbf{p}(p, A)$ is regular. The line L is regular so, $p \not\perp q$.

Let $q \notin M^{\perp}$. Then $x_2 := M \cap q^{\perp}$ is a point, for if not then $q \perp M$, so $q \in M^{\perp}$. Take distinct $y_1, y_2 \in M$ such that $y_1, y_2 \neq x_1, x_2$ and set $L_i = \overline{p, y_i}$ for i = 1, 2. From construction, $y_1, y_2 \not\perp p, q$. Then $L_1, L_2 \in \mathbf{p}_r(a, A)$ by 2.5. In view of 2.9 applied for the plane M + q the lines $M_i = \overline{q, y_i}, i = 1, 2$ are regular. So, the plane $p + M_i$ is regular for i = 1, 2, which gives $\Delta_{\mathfrak{B}_2}(L, L_1, L_2)$. This completes the proof of (i).

Let $q \in M^{\perp}$. The nonregular lines contained in **H** through q are all contained in the hyperplane $q^{\perp} \cap \mathbf{H}$ of **H** (the assumption $q \neq \mathbf{b}$ turns out to be essential here). It is impossible to decompose **H** into the union of three proper subspaces $q^{\perp} \cap \mathbf{H}, p^{\perp} \cap \mathbf{H}, \text{ and } M^{\perp} \cap \mathbf{H}$, so there is a point $q' \in \mathbf{H}$ with $q' \notin q^{\perp}, M^{\perp}, p^{\perp}$. Then the line $K := \overline{q, q'} \subset \mathbf{H}$ is regular and $p \not\perp K$. Let $z := K \cap p^{\perp}$ and $q'' \in M$, $q'' \neq q, q', z$. Then $q'' \not\perp p$ and thus the lines $L' := \overline{p, q'}$ and $L'' := \overline{p, q''}$ are regular. Write B = K + p; then $B \in \mathcal{R}_3$ and, evidently, $L, L', L'' \in \mathbf{p_r}(p, B)$. Moreover, $q'' \notin M^{\perp}$, as M is regular by 2.8. Since $L'^{\infty} = q'$ and $L''^{\infty} = q''$, the proof in case (ii) is complete.

Now, let L be an arbitrary regular line. From 5.1 we get that (i) implies $p \in L$. In case when (ii) holds we get $p \in L', L''$, which directly gives $p \in L$.

Since $p \notin \mathbf{H}$ in 5.6, then in view of 5.2 we have

COROLLARY 5.7. Let $\mathcal{Q} = \mathbf{p}_{r}^{\text{af}}(p, A) \neq \emptyset$ with $p \in A \in \mathcal{R}_{3}$ and $p \notin \mathbf{H}$, and let L be a regular affine line not in \mathcal{Q} . Write $q := L^{\infty}$ and $M := A^{\infty}$.

If $p \in L$ and $q \neq \mathbf{b}$, then one of the following holds:

(i) $q \notin M^{\perp}$. Then there are distinct $L_1, L_2 \in \mathcal{Q}$ such that $\Delta_{\mathfrak{C}_2}(L, L_1, L_2)$.

(ii) $q \in M^{\perp}$. Then there are distinct regular affine lines L', L'' through p such that $\mathbf{L}_{\mathfrak{C}_2}(L, L', L'')$ and $L'^{\infty}, L''^{\infty} \notin M^{\perp}$, so for both L' and L'' the condition (i) holds.

Conversely, if a line L satisfies (i) or (ii) then $p \in L$.

LEMMA 5.8. Let $\mathcal{Q} = \mathbf{p}_{\mathbf{r}}^{\mathrm{af}}(p, A) \neq \emptyset$ with $p \in A \in \mathbb{R}_3$, $p \in \mathbf{H}$ and let $L \notin \mathcal{Q}$ be a regular affine line. Then $p \in L$ iff there are distinct $L_1, L_2 \in \mathcal{Q}$ such that $\Delta_{\mathfrak{C}_2}(L, L_1, L_2)$ holds.

PROOF. By 2.8 and our assumptions the line $M := A^{\infty}$ is regular. Hence $p \neq \mathbf{b}$. By 5.2 the projective pencil $\mathbf{p}(p, A)$ contains exactly two lines that are not in $\mathbf{p}_{r}^{af}(p, A)$, namely M and $K_{0} := \overline{p}, \mathbf{q}(A)$.

⇒: We have $L \not\subset \mathbf{H}$ as L is affine. Since A + L is a projective 3-space, $D := (A + L) \cap \mathbf{H}$ is a plane. Note that $M \subset D$ so, by 2.9 $q := \operatorname{Rad}(D)$ is a point such that $q \notin M$. Set $K_1 = \overline{p,q}$. Consider the line $K_2 := (K_0 + L) \cap D$. Take two lines M_1, M_2 on D through p with $M_1, M_2 \neq M, K_1, K_2$. So, M_i is regular, and by 2.8 the plane $L + M_i$ is regular for i = 1, 2. Now take $L_i := (L + M_i) \cap A, i = 1, 2$. Observe that $L_1, L_2 \neq K_0$. Suppose that L_i is nonregular. Then by 2.5 we have $p \perp L_i$ which is impossible as the plane $A = L_1 + L_2$ is regular. Clearly, $L_1, L_2 \in \mathcal{Q}$ and $\Delta_{\mathfrak{C}_2}(L, L_1, L_2)$.

 \Leftarrow : A direct consequence of 5.1.

COROLLARY 5.9. Let $\mathcal{Q} = \mathbf{p}_r(p, A) \neq \emptyset$ with $p \in A \in \mathcal{R}_3$, $p \in \mathbf{H}$, and let $L \notin \mathcal{Q}$ be a regular line. Then $p \in L$ iff there are distinct $L_1, L_2 \in \mathcal{Q}$ such that $\Delta_{\mathfrak{B}_2}(L, L_1, L_2)$ holds.

PROOF. \Rightarrow : There are two cases to consider here: (i) $L \subset H$ and (ii) $L \not\subset H$.

In case (i) it suffices to take any two affine lines $L_1, L_2 \in \mathcal{Q}$ so, we have two new regular planes $L + L_1$, $L + L_2$ by 2.8 and $A = L_1 + L_2$ which means that $\Delta_{\mathfrak{B}_2}(L, L_1, L_2)$ is valid.

The case (ii) follows from 5.8.

 \Leftarrow : A direct consequence of 5.1.

Note that when $p \in \mathbf{H}$, then in $S(\mathcal{Q})$ there is no line L with $L^{\infty} \perp A^{\infty}$ as such a line is nonregular. Thus $S_L(\mathcal{Q}) = \emptyset$ and consequently $S(\mathcal{Q}) = \mathcal{Q} \cup S_{\Delta}(\mathcal{Q})$. Taking this into account and summing up 5.6 together with 5.9 and 5.7 together with 5.8 we get

PROPOSITION 5.10. Let $p \in A \in \mathbb{R}_3$. If $p \notin H$ and $\mathbf{b} \notin H$, or $p \in H$ and $p \neq \mathbf{b}$, then

 $S(\mathbf{p}_{r}(p, A)) = S_{r}(p)$ and $S(\mathbf{p}_{r}^{af}(p, A)) = S_{r}^{af}(p).$

If $p \notin \mathbf{H}$ and $\mathbf{b} \in \mathbf{H}$, then

$$S(\mathbf{p}_{r}(p,A)) = S(\mathbf{p}_{r}^{af}(p,A)) = S_{r}(p) \setminus \{\overline{p, \mathbf{b}}\} = S_{r}^{af}(p) \setminus \{\overline{p, \mathbf{b}}\}.$$

Roughly speaking, all the points of \mathfrak{P} except **b** can be reconstructed in terms of \mathfrak{B}_2 as well as in terms of \mathfrak{C}_2 . Actually, this is expectable since no plane through improper **b** is regular (comp. 2 and 5.5).

Now we will try to distinguish regular and nonregular points of \mathfrak{P} in terms of \mathfrak{B}_2 and \mathfrak{C}_2 . To do that we need a convenient characterization of binary collinearity

(adjacency) in these two structures. Basically two regular affine lines are adjacent iff they are coplanar and the plane is regular. Note that if a regular line from \mathbf{H} and a regular affine line are coplanar then the plane is always regular, while regular lines on \mathbf{H} are never adjacent by 2. The next lemma sheds yet more light on (non)adjacency of regular lines.

LEMMA 5.11. Let L_1, L_2 be distinct regular lines through a point p. Then $L_1 \not\sim L_2$ iff one of the following holds

- (i) $p \notin \mathbf{H}$ and $L_1^{\infty} \perp L_2^{\infty}$;
- (ii) $p \in \mathbf{H}$ and
 - a) either $L_1, L_2 \subset \mathbf{H}$,
 - b) or $L_1, L_2 \not\subset \mathbf{H}$ and $p \perp (L_1 + L_2)^{\infty}$

PROOF. Clearly, $A := L_1 + L_2$ is a plane. Therefore, $L_1 \not\sim L_2$ iff A is not regular.

In case $p \notin \mathbf{H}$ the plane A is affine and, in view of 2.8, it is nonregular iff $A^{\infty} = \overline{L_1^{\infty}, L_2^{\infty}}$ is nonregular i.e. iff $L_1^{\infty} \perp L_2^{\infty}$.

In case $p \in \mathbf{H}$ there are three possibilities. If $L_1, L_2 \subset \mathbf{H}$, then $A \subset \mathbf{H}$, so A is nonregular by 2. If $L_1 \subset \mathbf{H}$ and $L_2 \not\subset \mathbf{H}$, then $A^{\infty} = L_1$, so A is regular by 2.8. Finally, if $L_1, L_2 \not\subset \mathbf{H}$, then A is nonregular iff $A^{\infty} \perp A^{\infty}$, which is equivalent to $p \perp A^{\infty}$.

LEMMA 5.12. Assume that dim(\mathbb{V}) > 4. Let $\mathcal{Q} = \mathbf{p}_{r}(p, A)$ or $\mathcal{Q} = \mathbf{p}_{r}^{af}(p, A)$ with $\mathcal{Q} \neq \emptyset$ and $A \in \mathbb{R}_{3}$. The following conditions are equivalent:

- (i) $p \in \mathbf{H}$;
- (ii) for all $L, L_1, L_2 \in S(\mathcal{Q})$ from $L \not\sim L_1, L_2$ it follows that $L_1 \not\sim L_2$.

PROOF. Let $L \in S(Q)$. Then $p \in L$ by 5.10. In view of 5.11 it suffices to consider the following cases.

(i) Let $p \notin \mathbf{H}$. Set $q := L^{\infty}$. Take $q_1 \in q^{\perp} \cap \mathbf{H}$ with $q_1 \notin p^{\perp}$ and $q_2 \in q^{\perp} \cap \mathbf{H}$ with $q_2 \notin p^{\perp}, q_1^{\perp}$. The later is doable when dim $(\mathbb{V}) > 4$. So, we have $q_i \neq \mathbf{b}, i = 1, 2$ for if not we would have $q_1 \perp q_2$, a contradiction. Set $L_i := p + q_i$ for i = 1, 2. Then $L \not\sim L_1, L_2 \in \mathcal{S}(\mathcal{Q})$ and $L_1 \sim L_2$.

(ii) Let $p \in \mathbf{H}$, $L \not\subset \mathbf{H}$, and $L \not\sim L_1, L_2 \in \mathcal{S}(\mathcal{Q})$. From 5.11 we get $L_1, L_2 \not\subset \mathbf{H}$. **H**. Set $A_i := L + L_i$ and $M_i := A_i^{\infty}$. From assumption, the M_i are nonregular lines on **H** and therefore $p \perp M_i$; this yields $p \perp M_1 + M_2$. If $M_1 = M_2$ then $(L_1 + L_2)^{\infty} = M_1$, so $L_1 \not\sim L_2$. Assume that $M_1 \neq M_2$. Then there is a line $M_0 := (L_1 + L_2) \cap (M_1 + M_2)$. Clearly, $M_0 = (L_1 + L_2)^{\infty}$ and $p \in M_0 \perp p$, so M_0 is nonregular. Consequently, $L_1 \not\sim L_2$.

(iii) Let $p \in \mathbf{H}$, $L \subset \mathbf{H}$, and $L \not\sim L_1, L_2 \in \mathcal{S}(\mathcal{Q})$. From 5.11 we get $L_1, L_2 \subset \mathbf{H}$, and then $L_1 \not\sim L_2$ follows.

This closes our proof.

LEMMA 5.13. If dim(\mathbb{V}) = 4, then condition (ii) in 5.12 is always valid.

PROOF. Let \mathcal{Q} be as assumed in 5.12 and let $L, L_1, L_2 \in \mathcal{S}(\mathcal{Q}) = \mathcal{S}_r(p)$ with $L \not\sim L_1, L_2$. Set $q := L^{\infty}, q_i := L^{\infty}_i$ for i = 1, 2. By 5.11, $q \perp q_1, q_2$ which, in view of 2.8, means that planes $L + L_1, L + L_2$ are nonregular.

If $p = \mathbf{b}$, then the plane $L_1 + L_2$ is nonregular as no plane through **b** is regular, hence $L_1 \not\sim L_2$.

Now assume that $p \notin \mathbf{H}$. According to the definition of $S(\mathcal{Q})$ there is always a regular plane A' such that $L \subset A'$. Hence $q \neq \mathbf{b}$ by 5.5. In 3-space \mathfrak{P} the subspace $K := q^{\perp} \cap \mathbf{H}$ is a line on \mathbf{H} . Note that $q_1, q_2 \in K$ and also $q \in K$ as $q \in \mathbf{H}$ and thus $q \in q^{\perp}$. This means that K is isotropic so, $q_1 \perp q_2$. From 5.11 we get $L_1 \not\sim L_2$.

Finally, if $\mathbf{b} \neq p \in \mathbf{H}$, then we have three cases:

(i) Let $L_1, L_2 \subset \mathbf{H}$. Clearly $L_1 \not\sim L_2$.

(ii) Let $L_1 \subset \mathbf{H}$ and $L_2 \not\subset \mathbf{H}$. If $L \subset \mathbf{H}$, then the plane $L + L_2$ is regular by 2.8 which contradicts our assumptions that $L \not\sim L_2$. If $L \not\subset \mathbf{H}$, then the plane $L + L_1$ is regular, again a contradiction.

(iii) Let $L_1, L_2 \not\subset \mathbf{H}$. If $L \subset \mathbf{H}$, then both planes $L + L_1, L + L_2$ are regular which is impossible by our assumptions that $L \not\sim L_1, L_2$. If $L \not\subset \mathbf{H}$, then we have two distinct isotropic lines $M_i := (L + L_i) \cap \mathbf{H}, i = 1, 2$ on \mathbf{H} . Since $p \in M_1, M_2$ and $M_1 \cap M_2 = \mathbf{b}$, a contradiction and the proof is complete.

As it has been shown in 5.13 we need to treat the case dim(\mathbb{V}) = 4 separately. Recall that $\mathfrak{B}_2 \cong \mathfrak{B}_1^{\perp}$ and $\mathfrak{C}_2 \cong \mathfrak{C}_1^{\perp}$. So, we can state that the dual of \mathfrak{A} is definable in \mathfrak{B}_2 and \mathfrak{C}_2 by 3.20 and 3.13. Hence \mathfrak{P} is definable in \mathfrak{B}_2 . Moreover, the horizon $\langle \mathbf{H}, \perp_{\mathbf{H}} \rangle$ is also definable, so is \mathfrak{B}_1 .

In 5.10 we have defined points of \mathfrak{P} and thanks to 5.12 we are able to distinguish regular and nonregular points, all strictly in languages of \mathfrak{B}_2 and of \mathfrak{C}_2 . So, we have $\langle \mathsf{H}, \bot_{\mathsf{H}} \rangle$ reconstructed. Gathering all together we get

THEOREM 5.14. The structure \mathfrak{B}_1 can be defined in terms of both \mathfrak{B}_2 and \mathfrak{C}_2 .

References

- BACHMANN, F., Aufbau der Geometrie aus dem Spiegelungsbegriff, 2. Auflage, Springer-Verlag, Berlin, 1973.
- [2] BUEKENHOUT J., COHEN A., *Diagram Geometry*, (to appear).
- [3] COHEN M. A. Point-line spaces related to buildings. In Handbook of incidence geometry, F. Buekenhout, Ed. North-Holland, Amsterdam, 1995, pp. 647–737.
- [4] HALL, J. I., Classifying copolar spaces and graphs, Quart. J. Math. Oxford (2), 33 (1982), 421–449.
- [5] HIRSCHFELD J. W. W. P., Projective Geometries over Finite Fields, Clarendon Press, Oxford, 1998.
- [6] LEFÈVRE-PERCSY C., Copolar spaces fully embedded in projective spaces, Ann. Discrete Math. 18 (1983), 553–566.
- [7] PANKOV M., Grassmannians of classical buildings, Algebra and Discrete Mathematics Vol. 2, World Scientific, New Jersey, 2010.
- [8] PANKOV M., PRA£MOWSKI K., ŔYNEL M., Geometry of polar Grassmann spaces, Demonstratio Math. 39 (2006), no. 3, 625–637.

- [9] PRAŻMOWSKA M., PRAŻMOWSKI K., ŻYNEL M., Grassmann spaces of regular subspaces, J. Geom. 97 (2010), 99-123.
- [10] PRAŻMOWSKI K., ŻYNEL M., Orthogonality of subspaces in metric-projective geometry, Adv. Geom. 11 (2011), no. 1, 103-116.
- [11] TITS, J. Buildings of spherical type and finite BN-pairs, vol. 386 of Lecture Notes in Mathematics. Springer, Berlin, 1974.
- [12] UEBERBERG J., Foundations of Incidence Geometry: Projective and Polar Spaces, Springer-Verlag, Berlin, 2011.
- [13] VAN MALDEGHEM H., Generalized Polygons, Birkhäuser, Basel, 1998.

Authors' address: Krzysztof Prażmowski, Mariusz Żynel Institute of Mathematics, University of Białystok ul. Akademicka 2, 15-267 Białystok, Poland krzypraz@math.uwb.edu.pl, mariusz@math.uwb.edu.pl