## ON MULTIPARTITE HAJNAL-SZEMERÉDI THEOREMS

JIE HAN AND YI ZHAO

ABSTRACT. Let G be a k-partite graph with  $n$  vertices in parts such that each vertex is adjacent to at least  $\delta^*(G)$  vertices in each of the other parts. Magyar and Martin [\[20\]](#page-13-0) proved that for  $k=3$ , if  $\delta^*(G) \geq \frac{2}{3}n + 1$  and n is sufficiently large, then G contains a K<sub>3</sub>-factor (a spanning subgraph consisting of n vertex-disjoint copies of  $K_3$ ). Martin and Szemerédi [\[21\]](#page-13-1) proved that G contains a K<sub>4</sub>-factor when  $\delta^*(G) \geq \frac{3}{4}n$  and n is sufficiently large. Both results were proved using the Regularity Lemma. In this paper we give a proof of these two results by the absorbing method. Our absorbing lemma actually works for all  $k \geq 3$  and may be utilized to prove a general and tight multipartite Hajnal-Szemerédi theorem.

#### 1. INTRODUCTION

Let H be a graph on h vertices, and let G be a graph on n vertices. Packing (or tiling) problems in extremal graph theory are investigations of conditions under which  $G$  must contain many vertex disjoint copies of  $H$  (as subgraphs), where minimum degree conditions are studied the most. An H-matching of  $G$  is a subgraph of  $G$  which consists of vertex-disjoint copies of  $H$ . A perfect  $H$ matching, or H-factor, of G is an H-matching consisting of  $|n/h|$  copies of H. Let  $K_k$  denote the complete graph on  $k$  vertices. The celebrated theorem of Hajnal and Szemerédi  $[6]$  says that every n-vertex graph G with  $\delta(G) \geq (k-1)n/k$  contains a  $K_k$ -factor (see [\[11\]](#page-12-1) for another proof).

Using the Regularity Lemma of Szemerédi [\[25\]](#page-13-2), researchers have generalized this theorem for packing arbitrary  $H$  [\[1,](#page-12-2) [15,](#page-12-3) [24,](#page-13-3) [16\]](#page-12-4). Results and methods for packing problems can be found in the survey of Kühn and Osthus [\[17\]](#page-12-5).

In this paper we consider multipartite packing, which restricts G to be a k-partite graph for  $k \geq 2$ . A k-partite graph is called balanced if its partition sets have the same size. Given a k-partite graph G, it is natural to consider the minimum partite degree  $\delta^*(G)$ , the minimum degree from a vertex in one partition set to any other partition set. When  $k = 2$ ,  $\delta^*(G)$  is simply  $\delta(G)$ . In most of the rest of this paper, the minimum degree condition stands for the minimum partite degree for short.

Let  $\mathcal{G}_k(n)$  denote the family of balanced k-partite graphs with n vertices in each of its partition sets. It is easy to see (e.g. using the König-Hall Theorem) that every bipartite graph  $G \in \mathcal{G}_2(n)$ with  $\delta^*(G) \geq n/2$  contains a 1-factor. Fischer [\[5\]](#page-12-6) conjectured that if  $G \in \mathcal{G}_k(n)$  satisfies

<span id="page-0-0"></span>
$$
\delta^*(G) \ge \frac{k-1}{k}n,\tag{1}
$$

then G contains a  $K_k$ -factor and proved the existence of an *almost*  $K_k$ -factor for  $k = 3, 4$ . Magyar and Martin [\[20\]](#page-13-0) noticed that the condition [\(1\)](#page-0-0) is not sufficient for odd  $k$  and instead proved the following theorem for  $k = 3$ . (They actually showed that when n is divisible by 3, there is only one graph in  $G_3(n)$ , denoted by  $\Gamma_3(n/3)$ , that satisfies [\(1\)](#page-0-0) but fails to contain a  $K_3$ -factor, and adding any new edge to  $\Gamma_3(n/3)$  results in a  $K_3$ -factor.)

Date: November 7, 2018.

<sup>1991</sup> Mathematics Subject Classification. Primary 05C35, 05C70.

Key words and phrases. graph packing, Hajnal-Szemerédi, absorbing method.

The second author was supported NSA grants H98230-10-1-0165 and H98230-12-1-0283.

<span id="page-1-0"></span>**Theorem 1** ([\[20\]](#page-13-0)). There exists an integer  $n_0$  such that If  $n \ge n_0$  and  $G \in \mathcal{G}_3(n)$  satisfies  $\delta^*(G) \ge$  $2n/3+1$ , then G contains a K<sub>3</sub>-factor.

On the other hand, Martin and Szemerédi [\[21\]](#page-13-1) proved the original conjecture holds for  $k = 4$ .

<span id="page-1-1"></span>**Theorem 2** ([\[21\]](#page-13-1)). There exists an integer  $n_0$  such that if  $n \ge n_0$  and  $G \in \mathcal{G}_4(n)$  satisfies  $\delta^*(G) \ge$  $3n/4$ , then G contains a K<sub>4</sub>-factor.

Recently Keevash and Mycroft [\[9\]](#page-12-7) and independently Lo and Markström [\[19\]](#page-12-8) proved that Fischer's conjecture is asymptotically true, namely,  $\delta^*(G) \ge \frac{k-1}{k}n + o(n)$  guarantees a  $K_k$ -factor for all  $k \ge 3$ . Very recently, Keevash and Mycroft [\[10\]](#page-12-9) improved this to an exact result.

In this paper we give a new proof of Theorems [1](#page-1-0) and [2](#page-1-1) by the absorbing method. Our approach is similar to that of [\[19\]](#page-12-8) (in contrast, a geometric approach was employed in [\[9\]](#page-12-7)). However, in order to prove exact results by the absorbing lemma, one needs only assume  $\delta^*(G) \geq (1 - 1/k)n$ , instead of  $\delta^*(G) \ge (1 - 1/k + \alpha)n$  for some  $\alpha > 0$  as in [\[19\]](#page-12-8). In fact, our absorbing lemma uses an even weaker assumption  $\delta^*(G) \ge (1 - 1/k - \alpha)n$  and has a more complicated absorbing structure.

The absorbing method, initiated by Rödl, Ruciński, and Szemerédi [\[23\]](#page-13-4), has been shown to be effective handling extremal problems in graphs and hypergraphs. One example is the re-proof of Posa's conjecture by Levitt, Sárközy, and Szemerédi [\[18\]](#page-12-10), while the original proof of Komlós, Sárközy, and Szemerédi [\[13\]](#page-12-11) used the Regularity Lemma. Our paper is another example of replacing the regularity method with the absorbing method. Compared with the threshold  $n_0$  in Theorems [1](#page-1-0) and [2](#page-1-1) derived from the Regularity Lemma, the value of our  $n_0$  is much smaller.

Before presenting our proof, let us first recall the approach used in  $[20, 21]$  $[20, 21]$ . Given a k-partite graph  $G \in \mathcal{G}_k(n)$  with parts  $V_1, \ldots, V_k$ , the authors said that G is  $\Delta$ -extremal if each  $V_i$  contains a subset  $A_i$  of size  $\lfloor n/k \rfloor$  such that the density  $d(A_i, A_j) \leq \Delta$  for all  $i \neq j$ . Using standard but involved graph theoretic arguments, they solved the extremal case for  $k = 3,4$  [\[20,](#page-13-0) Theorem 3.1], [\[21,](#page-13-1) Theorem 2.1].

<span id="page-1-4"></span>**Theorem 3.** Let  $k = 3, 4$ . There exists  $\Delta$  and  $n_0$  such that the following holds. Let  $n \geq n_0$  and  $G \in \mathcal{G}_k(n)$  be a k-partite graph satisfying  $\delta^*(G) \ge (2/3)n + 1$  when  $k = 3$  and [\(1\)](#page-0-0) when  $k = 4$ . If G is  $\Delta$ -extremal, then G contains a  $K_k$ -factor.

To handle the non-extremal case, they proved the following lemma([\[20,](#page-13-0) Lemma 2.2] and [\[21,](#page-13-1) Lemma 2.2]).

<span id="page-1-2"></span>**Lemma 4** (Almost Covering Lemma). Let  $k = 3, 4$ . Given  $\Delta > 0$ , there exists  $\alpha > 0$  such that for every graph  $G \in \mathcal{G}_k(n)$  with  $\delta^*(G) \geq (1 - 1/k)n - \alpha n$  either G contains an almost  $K_k$ -factor that leaves at most  $C = C(k)$  vertices uncovered or G is  $\Delta$ -extremal.

To improve the almost  $K_k$ -factor obtained from Lemma [4,](#page-1-2) they used the Regularity Lemma and Blow-up Lemma [\[14\]](#page-12-12). Here is where we need our absorbing lemma whose proof is given in Section 2. Our lemma actually gives a more detailed structure than what is needed for the extremal case when G does not satisfy the absorbing property.

We need some definitions. Given positive integers k and r, let  $\Theta_{k \times r}$  denote the graph with vertices  $a_{ij}, i = 1, \ldots, k, j = 1, \ldots, r$ , and  $a_{ij}$  is adjacent to  $a_{i'j'}$  if and only if  $i \neq i'$  and  $j \neq j'$ . In addition, given a positive integer t, the graph  $\Theta_{k \times r}(t)$  denotes the blow-up of  $\Theta_{k \times r}$ , obtained by replacing vertices  $a_{ij}$  with sets  $A_{ij}$  of size t, and edges  $a_{ij}a_{i'j'}$  with complete bipartite graphs between  $A_{ij}$ and  $A_{i'j'}$ . Given  $\epsilon, \Delta > 0$  and  $t \ge 1$  (not necessarily an integer), we say that a k-partite graph G is  $(\epsilon, \Delta)$ -approximate to  $\Theta_{k \times r}(t)$  if each of its partition sets  $V_i$  can be partitioned into  $\bigcup_{i=1}^r V_{ij}$  such that  $||V_{ij}|-t| \leq \epsilon t$  for all  $i, j$  and  $d(V_{ij}, V_{i'j}) \leq \Delta$  whenever  $i \neq i'$ .<sup>[1](#page-1-3)</sup>

<span id="page-1-3"></span><sup>&</sup>lt;sup>1</sup>Here we follow the definition of  $(\epsilon, \Delta)$ -approximation in [\[20,](#page-13-0) [21\]](#page-13-1). It seems natural to require that  $d(V_{ij}, V_{i'j'}) \ge$ 1 – ∆ whenever  $i \neq i'$  and  $j \neq j'$  as well. However, this follows from  $d(V_{ij}, V_{i'j}) \leq \Delta (i \neq i')$  when  $\delta^*(G) \geq (1-1/r)rt$ .

<span id="page-2-0"></span>**Lemma 5** (Absorbing Lemma). Given  $k \geq 3$  and  $\Delta > 0$ , there exists  $\alpha = \alpha(k, \Delta) > 0$  and an integer  $n_1 > 0$  such that the following holds. Let  $n \geq n_1$  and  $G \in \mathcal{G}_k(n)$  be a k-partite graph on  $V_1 \cup \cdots \cup V_k$  such that  $\delta^*(G) \geq (1 - 1/k)n - \alpha n$ . Then one of the following cases holds.

- (1) G contains a  $K_k$ -matching M of size  $|M| \leq 2(k-1)\alpha^{4k-2}n$  in G such that for every  $W \subset V \backslash V(M)$  with  $|W \cap V_1| = \cdots = |W \cap V_k| \leq \alpha^{8k-6} n/4$ , there exists a  $K_k$ -matching covering exactly the vertices in  $V(M) \cup W$ .
- (2) We may remove some edges from G so that the resulting graph G' satisfies  $\delta^*(G') \geq (1 1/k$ )n –  $\alpha n$  and is  $(\Delta/6, \Delta/2)$ -approximate to  $\Theta_{k \times k}(\frac{n}{k})$ .

The  $K_k$ -matching M in Lemma [5](#page-2-0) has the so-called *absorbing* property: it can absorb any balanced set with a much smaller size.

*Proof of Theorems [1](#page-1-0) and [2.](#page-1-1)* Let  $k = 3, 4$  $k = 3, 4$  $k = 3, 4$ . Let  $\alpha \ll \Delta$ , where  $\Delta$  is given by Theorem 3 and  $\alpha$  satisfies both Lemmas [4](#page-1-2) and [5.](#page-2-0) Suppose that n is sufficiently large. Let  $G \in \mathcal{G}_k(n)$  be a k-partite graph satisfying  $\delta^*(G) \ge (2/3)n + 1$  when  $k = 3$  and [\(1\)](#page-0-0) when  $k = 4$ . By Lemma [5,](#page-2-0) either G contains a subgraph which is  $(\Delta/6, \Delta/2)$ -approximate to  $\Theta_{k \times k}(\frac{n}{k})$  or G contains an absorbing  $K_k$ -matching M. In the former case, for  $i = 1, ..., k$ , we add or remove at most  $\frac{\Delta n}{6k}$  vertices from  $V_{i1}$  to obtain a set  $A_i \subset V_i$  of size  $\lfloor n/k \rfloor$ . For  $i \neq i'$ , we have

$$
e(A_i, A_{i'}) \le e(V_{i1}, V_{i'1}) + \frac{\Delta n}{6k}(|A_i| + |A_{i'}|)
$$
  
\n
$$
\le \frac{\Delta}{2}|V_{i1}||V_{i'1}| + 2\frac{\Delta n}{6k} \left\lfloor \frac{n}{k} \right\rfloor
$$
  
\n
$$
\le \frac{\Delta}{2} \left(1 + \frac{\Delta}{6}\right)^2 \left(\frac{n}{k}\right)^2 + \frac{\Delta n}{3k} \left\lfloor \frac{n}{k} \right\rfloor
$$
  
\n
$$
\le \Delta \left\lfloor \frac{n}{k} \right\rfloor \left\lfloor \frac{n}{k} \right\rfloor,
$$

which implies that  $d(A_i, A_{i'}) \leq \Delta$ . Thus G is  $\Delta$ -extremal. By Theorem [3,](#page-1-4) G contains a  $K_k$ -factor. In the latter case, G contains a  $K_k$ -matching M is of size  $|M| \leq 2(k-1)\alpha^{4k-2}n$  such that for every  $W \subset V \backslash V(M)$  with  $|W \cap V_1| = \cdots = |W \cap V_k| \leq \alpha^{8k-6} n/4$ , there exists a  $K_k$ -matching on  $V(M) \cup W$ . Let  $G' = G \setminus V(M)$  be the induced subgraph of G on  $V(G) \setminus V(M)$ , and  $n' = |V(G')|$ . Clearly  $G'$  is balanced. As  $\alpha \ll 1$ , we have

$$
\delta^*(G') \ge \delta^*(G) - |M| \ge \left(1 - \frac{1}{k}\right)n - 2(k-1)\alpha^{4k-2}n \ge \left(1 - \frac{1}{k} - \alpha\right)n'.
$$

By Lemma [4,](#page-1-2) G' contains a  $K_k$ -matching M' such that  $|V(G') \setminus V(M')| \leq C$ . Let  $W = V(G') \setminus V(G)$  $V(M')$ . Clearly  $|W \cap V_1| = \cdots = |W \cap V_k|$ . Since  $C/k \le \alpha^{8k-6}n/4$  for sufficiently large n, by the absorbing property of M, there is a  $K_k$ -matching  $M''$  on  $V(M) \cup W$ . This gives the desired  $K_k$ -factor  $M' \cup M''$  of G.  $K_k$ -factor  $M' \cup M''$  of  $G$ .

## Remarks.

- Since our Lemma [5](#page-2-0) works for all  $k \geq 3$ , it has the potential of proving a general multipartite Hajnal-Szemerédi theorem. To do it, one only needs to prove Theorem [3](#page-1-4) and Lemma [4](#page-1-2) for  $k > 5$ .
- Since our Lemma [5](#page-2-0) gives a detailed structure of  $G$  when  $G$  does not have desired absorbing  $K_k$ -matching, it has the potential of simplifying the proof of the extremal case. Indeed, if one can refine Lemma [4](#page-1-2) such that it concludes that G either contains an almost  $K_k$ -factor or it is approximate to  $\Theta_{k\times k}(\frac{n}{k})$  and other extremal graphs, then in Theorem [3](#page-1-4) we may assume that  $G$  is actually approximate to these extremal graphs.
- Using the Regularity Lemma, researchers have obtained results on packing arbitrary graphs in k-partite graphs, see [\[26,](#page-13-5) [8,](#page-12-13) [3,](#page-12-14) [2\]](#page-12-15) for  $k = 2$  and [\[22\]](#page-13-6) for  $k = 3$ . With the help of the recent

#### 4 JIE HAN AND YI ZHAO

result of Keevash–Mycroft  $[9]$  and Lo-Markström  $[19]$ , it seems not very difficult to extend these results to the  $k > 4$  case (though exact results may be much harder). However, it seems difficult to replace the regularity method by the absorbing method for these problems.

## 2. Proof of the Absorbing Lemma

In this section we prove the Absorbing Lemma (Lemma [5\)](#page-2-0). We first introduce the concepts of reachability.

**Definition 6.** In a graph G, a vertex x is reachable from another vertex y by a set  $S \subseteq V(G)$  if both  $G[x \cup S]$  and  $G[y \cup S]$  contain  $K_k$ -factors. In this case, we say S connects x and y.

The following lemma plays a key role in constructing absorbing structures. We postpone its proof to the end of the section.

<span id="page-3-0"></span>**Lemma 7** (Reachability Lemma). Given  $k \geq 3$  and  $\Delta > 0$ , there exists  $\alpha = \alpha(k, \Delta) > 0$  and an integer  $n_2 > 0$  such that the following holds. Let  $n \geq n_2$  and  $G \in \mathcal{G}_k(n)$  be a k-partite graph on  $V_1 \cup \cdots \cup V_k$  such that  $\delta^*(G) \geq (1-1/k)n - \alpha n$ . Then one of the following cases holds.

- (1) For any x and y in  $V_i$ ,  $i \in [k]$ , x is reachable from y by either at least  $\alpha^3 n^{k-1}$   $(k-1)$ -sets or at least  $\alpha^3 n^{2k-1}$   $(2k-1)$ -sets in G.
- (2) We may remove some edges from G so that the resulting graph G' satisfies  $\delta^*(G') \geq (1 1/k$ )n –  $\alpha n$  and is  $(\Delta/6, \Delta/2)$ -approximate to  $\Theta_{k \times k}(\frac{n}{k})$ .

With the aid of Lemma [7,](#page-3-0) the proof of Lemma [5](#page-2-0) becomes standard counting and probabilistic arguments, as shown in [\[7\]](#page-12-16).

Proof of Lemma [5.](#page-2-0) We assume that G does not satisfy the second property stated in the lemma.

Given a crossing k-tuple  $T = (v_1, \dots, v_k)$ , with  $v_i \in V_i$ , for  $i = 1, \dots, k$ , we call a set A an absorbing set for T if both  $G[A]$  and  $G[A \cup T]$  contain  $K_k$ -factors. Let  $\mathcal{L}(T)$  denote the family of all  $2k(k-1)$ -sets that absorb T (the reason why our absorbing sets are of size  $2k(k-1)$  can be seen from the proof of Claim [8](#page-3-1) below).

<span id="page-3-1"></span>**Claim 8.** For every crossing k-tuple T, we have  $|\mathcal{L}(T)| > \alpha^{4k-3} n^{2k(k-1)}$ .

*Proof.* Fix a crossing k-tuple T. First we try to find a copy of  $K_k$  containing  $v_1$  and avoiding  $v_2$ ,  $\dots, v_k$ . By the minimum degree condition, there are at least

$$
\prod_{i=2}^{k} \left( n - 1 - (i - 1) \left( \frac{1}{k} + \alpha \right) n \right) \ge \prod_{i=2}^{k} \left( n - (i - 1) \frac{n}{k} - ((k - 1)\alpha n + 1) \right)
$$

such copies of  $K_k$ . When  $n \geq 3k^2$  and  $\frac{1}{\alpha} \geq 3k^2$ , we have  $(k-1)\alpha n + 1 \leq n/(3k)$  and thus the number above is at least

$$
\prod_{i=2}^{k} \left( n - (i-1)\frac{n}{k} - \frac{n}{3k} \right) \ge \left( \frac{n}{k} \right)^{k-1}, \text{ when } k \ge 3.
$$

Fix such a copy of  $K_k$  on  $\{v_1, u_2, u_3, \cdots, u_k\}$ . Consider  $u_2$  and  $v_2$ . By Lemma [7](#page-3-0) and the assumption that G does not satisfy the second property of the lemma, we can find at least  $\alpha^3 n^{k-1}$  $(k-1)$ -sets or  $\alpha^3 n^{2k-1}$   $(2k-1)$ -sets to connect  $u_2$  and  $v_2$ . If S is a  $(k-1)$ -set that connects  $u_2$ and  $v_2$ , then  $S \cup K$  also connects  $u_2$  and  $v_2$  for any k-set K such that  $G[K] \cong K_k$  and  $K \cap S = \emptyset$ . There are at least

$$
(n-2)\prod_{i=2}^{k} \left(n-1-(i-1)\left(\frac{1}{k}+\alpha\right)n\right) \ge \frac{n}{2} \left(\frac{n}{k}\right)^{k-1}
$$

copies of  $K_k$  in G avoiding  $u_2$ ,  $v_2$  and S. If there are at least  $\alpha^3 n^{k-1}$  (k - 1)-sets that connect  $u_2$ and  $v_2$ , then at least

$$
\alpha^3 n^{k-1} \cdot \frac{n}{2} \left(\frac{n}{k}\right)^{k-1} \frac{1}{\binom{2k-1}{k-1}} \ge 2\alpha^4 n^{2k-1}
$$

 $(2k-1)$ -sets connect  $u_2$  and  $v_2$  because a  $(2k-1)$ -set can be counted at most  $\binom{2k-1}{k-1}$  times. Since  $2\alpha^4 < \alpha^3$ , we can assume that there are always at least  $2\alpha^4 n^{2k-1} (2k-1)$ -sets connecting  $u_2$  and v<sub>2</sub>. We inductively choose disjoint  $(2k-1)$ -sets that connects v<sub>i</sub> and u<sub>i</sub> for  $i = 2, \ldots, k$ . For each i, we must avoid T,  $u_2, \ldots, u_k$ , and  $i-2$  previously selected  $(2k-1)$ -sets. Hence there are at least  $2\alpha^4 n^{2k-1} - (2k-1)(i-1)n^{2k-2} > \alpha^4 n^{2k-1}$  choices of such  $(2k-1)$ -sets for each  $i \ge 2$ . Putting all these together, and using the assumption that  $\alpha$  is sufficiently small, we have

$$
|\mathcal{L}(T)| \ge \left(\frac{n}{k}\right)^{k-1} \cdot (\alpha^4 n^{2k-1})^{k-1} > \alpha^{4k-3} n^{2k(k-1)}.
$$

Every set  $S \in \mathcal{L}(T)$  is balanced because  $G[S]$  contains a  $K_k$ -factor and thus  $|S \cap V_1| = \cdots =$  $|S \cap V_k| = 2(k-1)$ . Note that there are  $\binom{n}{2(k-1)}^k$  balanced  $2k(k-1)$ -sets in G. Let F be the random family of  $2k(k-1)$ -sets obtained by selecting each balanced  $2k(k-1)$ -set from  $V(G)$  independently with probability  $p := \alpha^{4k-3} n^{1-2k(k-1)}$ . Then by Chernoff's bound, since n is sufficiently large, with probability  $1 - o(1)$ , the family F satisfies the following properties:

$$
|\mathcal{F}| \le 2\mathbb{E}(|\mathcal{F}|) \le 2p \binom{n}{2(k-1)}^k \le \alpha^{4k-2}n,\tag{2}
$$

$$
|\mathcal{L}(T) \cap \mathcal{F}| \ge \frac{1}{2} \mathbb{E}(|\mathcal{L}(T) \cap \mathcal{F}|) \ge \frac{1}{2} p|\mathcal{L}(T)| \ge \frac{\alpha^{8k - 6}n}{2} \text{ for every crossing } k \text{-tuple } T. \tag{3}
$$

Let Y be the number of intersecting pairs of members of F. Since each fixed balanced  $2k(k-1)$ -set intersects at most  $2k(k-1)\binom{n-1}{2(k-1)-1}\binom{n}{2(k-1)}^{k-1}$  other balanced  $2k(k-1)$ -sets in G,

$$
\mathbb{E}(Y) \le p^2 \binom{n}{2(k-1)}^k 2k(k-1) \binom{n-1}{2k-3} \binom{n}{2(k-1)}^{k-1} \le \frac{1}{8} \alpha^{8k-6} n.
$$

By Markov's bound, with probability at least  $\frac{1}{2}$ ,  $Y \le \alpha^{8k-6}n/4$ . Therefore, we can find a family  $\mathcal F$ satisfying [\(2\)](#page-4-0), [\(3\)](#page-4-1) and having at most  $\alpha^{8k-6}n/4$  intersecting pairs. Remove one set from each of the intersecting pairs and the sets that have no  $K_k$ -factor from  $\mathcal{F}$ , we get a subfamily  $\mathcal{F}'$  consisting of pairwise disjoint absorbing  $2k(k-1)$ -sets which satisfies  $|\mathcal{F}'| \leq |\mathcal{F}| \leq \alpha^{4k-2}n$  and for all crossing T,

$$
|\mathcal{L}(T) \cap \mathcal{F}'| \geq \frac{\alpha^{8k-6}n}{2} - \frac{\alpha^{8k-6}n}{4} \geq \frac{\alpha^{8k-6}n}{4}.
$$

Since  $\mathcal{F}'$  consists of disjoint absorbing sets and each absorbing set is covered by a  $K_k$ -matching,  $V(F')$  is covered by some  $K_k$ -matching M. Since  $|\mathcal{F}'| \leq \alpha^{4k-2}n$ , we have  $|M| \leq 2k(k-1)\alpha^{4k-2}n/k =$  $2(k-1)\alpha^{4k-2}n$ . Now consider a balanced set  $W \subseteq V(G) \backslash V(\mathcal{F}')$  such that  $|W \cap V_1| = \cdots = |W \cap V_k| \le$  $\alpha^{8k-6}n/4$ . Arbitrarily partition W into at most  $\alpha^{8k-6}n/4$  crossing k-tuples. We absorb each of the k-tuples with a different  $2k(k-1)$ -set from  $\mathcal{L}(T) \cap \mathcal{F}'$ . As a result,  $V(\mathcal{F}') \cup W$  is covered by a  $K_k$ -matching, as desired.

<span id="page-4-1"></span><span id="page-4-0"></span> $\Box$ 

The rest of the paper is devoted to proving Lemma [7.](#page-3-0) First we prove a useful lemma. A weaker version of it appears in [\[21,](#page-13-1) Proposition 1.4] with a brief proof sketch.

<span id="page-4-2"></span>**Lemma 9.** Let  $k \geq 2$  be an integer,  $t \geq 1$  and  $\epsilon \ll 1$ . Let H be a k-partite graph on  $V_1 \cup \cdots \cup V_k$ such that  $|V_i| \geq (k-1)(1-\epsilon)t$  for all i and each vertex is nonadjacent to at most  $(1+\epsilon)t$  vertices in each of the other color classes. Then either H contains at least  $\epsilon^2 t^k$  copies of  $K_k$ , or H is  $(16k^4\epsilon^{1/2^{k-2}}, 16k^4\epsilon^{1/2^{k-2}})$ -approximate to  $\Theta_{k\times (k-1)}(t)$ .

*Proof.* First we derive an upper bound for  $|V_i|, i \in [k]$ . Suppose for example, that  $|V_k| \ge (k-1)(1+1)$  $\epsilon$ )t +  $\epsilon t$ . Then if we greedily construct copies of  $K_k$  while choosing the last vertex from  $V_k$ , by the minimum degree condition and  $\epsilon \ll 1$ , there are at least

$$
|V_1| \cdot (|V_2| - (1 + \epsilon)t) \cdots (|V_{k-1}| - (k-2)(1+\epsilon)t) \cdot (|V_k| - (k-1)(1+\epsilon)t)
$$
  
\n
$$
\geq (k-1)(1-\epsilon)t \cdot (k-2-k\epsilon)t \cdots (1-(2k-3)\epsilon)t \cdot \epsilon t
$$
  
\n
$$
\geq (k-1-\frac{1}{2})(k-2-\frac{1}{2}) \cdots (1-\frac{1}{2})\epsilon t^k \geq \frac{\epsilon}{2}t^k
$$

copies of  $K_k$  in  $H$ , so we are done. We thus assume that for all  $i$ ,

<span id="page-5-0"></span>
$$
|V_i| \le (k-1)(1+\epsilon)t + \epsilon t < (k-1)(1+2\epsilon)t. \tag{4}
$$

Now we proceed by induction on k. The base case is  $k = 2$ . If H has at least  $\epsilon^2 t^2$  edges, then we are done. Otherwise  $e(H) < \epsilon^2 t^2$ . Using the lower bound for  $|V_i|$ , we obtain that

$$
d(V_1, V_2) < \frac{\epsilon^2 t^2}{|V_1| \cdot |V_2|} \le \frac{\epsilon^2}{(1 - \epsilon)^2} < \epsilon.
$$

Hence H is  $(2\epsilon, \epsilon)$ -approximate to  $\Theta_{2\times1}(t)$ . When  $k=2$ ,  $16k^4\epsilon^{1/2^{k-2}}=256\epsilon$ , so we are done.

Now assume that  $k \geq 3$  and the conclusion holds for  $k-1$ . Let H be a k-partite graph satisfying the assumptions and assume that H contains less than  $\epsilon^2 t^k$  copies of  $K_k$ .

For simplicity, write  $N_i(v) = N(v) \cap V_i$  for any vertex v. Let  $V'_1 \subset V_1$  be the vertices which are in at least  $\epsilon t^{k-1}$  copies of  $K_k$  in H, and let  $\tilde{V}_1 = V_1 \setminus V'_1$ . Note that  $|V'_1| < \epsilon t$  otherwise we get at least  $\epsilon^2 t^k$  copies of  $K_k$  in H. Fix  $v_0 \in \tilde{V}_1$ . For  $2 \leq i \leq k$ , by the minimum degree condition and  $k \geq 3$ ,

$$
|N_i(v_0)| \ge (k-1)(1-\epsilon)t - (1+\epsilon)t = (k-2)\left(1 - \frac{k}{k-2}\epsilon\right)t \ge (k-2)(1-3\epsilon)t.
$$

On the other hand, following the same arguments as we used for [\(4\)](#page-5-0), we derive that

<span id="page-5-1"></span>
$$
|N_i(v_0)| \le (k-2)(1+2\epsilon t). \tag{5}
$$

The minimum degree condition implies that a vertex in  $N(v_0)$  misses at most  $(1+\epsilon)t$  vertices in each  $N_i(v_0)$ . We now apply induction with  $k-1$ , t and  $3\epsilon$  on  $H[N(v_0)]$ . Because of the definition of  $V'_1$ , we conclude that  $N(v_0)$  is  $(\epsilon', \epsilon')$ -approximate to  $\Theta_{(k-1)\times (k-2)}(t)$ , where

$$
\epsilon' := 16(k-1)^4 (3\epsilon)^{1/2^{k-3}}
$$

This means that we can partition  $N_i(v_0)$  into  $A_{i1} \cup \cdots A_{i(k-2)}$  for  $2 \leq i \leq k$  such that

$$
\forall 2 \le i \le k, 1 \le j \le k-2, \quad (1 - \epsilon')t \le |A_{ij}| \le (1 + \epsilon')t \quad \text{and} \tag{6}
$$

<span id="page-5-3"></span><span id="page-5-2"></span>.

$$
\forall 2 \leq i < i' \leq k, \ 1 \leq j \leq k-2, \quad d(A_{ij}, A_{i'j}) \leq \epsilon'.\tag{7}
$$

Furthermore, let  $A_{i(k-1)} := V_i \setminus N(v_0)$  for  $i = 2, \dots, k$ . By [\(5\)](#page-5-1) and the minimum degree condition, we get that

<span id="page-5-4"></span>
$$
(1 - (3k - 5)\epsilon)t \le |A_{i(k-1)}| \le (1 + \epsilon)t,\tag{8}
$$

for  $i = 2, \cdots, k$ .

Let  $A_{ij}^c = V_i \backslash A_{ij}$  denote the complement of  $A_{ij}$ . Let  $\bar{e}(A, B) = |A||B| - e(A, B)$  denote the number of non-edges between two disjoint sets A and B, and  $\bar{d}(A, B) = \bar{e}(A, B)/(|A||B|) = 1 - d(A, B)$ . Given two disjoint sets A and B (with density close to one) and  $\alpha > 0$ , we call a vertex  $a \in A$  is  $\alpha$ -typical to B if deg<sub>B</sub> $(a) \geq (1 - \alpha)|B|$ .

<span id="page-5-5"></span>Claim 10. Let  $2 \le i \neq i' \le k$ ,  $1 \le j \neq j' \le k - 1$ . (1)  $d(A_{ij}, A_{i'j'}) \geq 1 - 3\epsilon'$  and  $d(A_{ij}, A_{i'j}^c) \geq 1 - 3\epsilon'.$  (2) All but at most  $\sqrt{3\epsilon'}$  vertices in  $A_{ij}$  are  $\sqrt{3\epsilon'}$ -typical to  $A_{i'j'}$ ; at most  $\sqrt{3\epsilon'}$  vertices in  $A_{ij}$ are  $\sqrt{3\epsilon'}$ -typical to  $A_{i'j}^c$ .

*Proof.* (1). Since  $A_{i'j}^c = \bigcup_{j' \neq j} A_{i'j'}$ , the second assertion  $d(A_{ij}, A_{i'j}^c) \geq 1 - 3\epsilon'$  immediately follows from the first assertion  $d(\tilde{A}_{ij}, \tilde{A}_{i'j'}) \geq 1-3\epsilon'$ . Thus it suffices to show that  $d(A_{ij}, A_{i'j'}) \geq 1-3\epsilon'$ , or equivalently that  $\bar{d}(A_{ij}, \dot{A}_{i'j'}) \leq 3\epsilon'.$ 

Assume  $j \ge 2$ . By [\(7\)](#page-5-2), we have  $e(A_{ij}, A_{i'j}) \le \epsilon' |A_{ij}| |A_{i'j}|$ . So  $\bar{e}(A_{ij}, A_{i'j}) \ge (1 - \epsilon') |A_{ij}| |A_{i'j}|$ . By the minimum degree condition and [\(6\)](#page-5-3),

$$
\bar{e}(A_{ij}, A_{i'j}^c) \le [(1+\epsilon)t - (1-\epsilon')|A_{i'j}|||A_{ij}|
$$
  
\n
$$
\le [(1+\epsilon)t - (1-\epsilon')(1-\epsilon')t]|A_{ij}|
$$
  
\n
$$
< (\epsilon + 2\epsilon')t|A_{ij}|,
$$

which implies that  $\bar{e}((A_{ij}, A_{i'j'}) \leq (\epsilon + 2\epsilon')t|A_{ij}|$  for any  $j' \neq j$  and  $1 \leq j' \leq k - 1$ . By [\(6\)](#page-5-3) and [\(8\)](#page-5-4), we have  $|A_{i'j'}| \geq (1 - \epsilon')t$ . Hence

$$
\bar{d}(A_{ij}, A_{i'j'}) \leq (\epsilon + 2\epsilon') \frac{t}{|A_{i'j'}|} \leq (\epsilon + 2\epsilon') \frac{t}{(1 - \epsilon')t} \leq 3\epsilon',
$$

where the last inequality holds because  $\epsilon \ll \epsilon' \ll 1$ .

(2) Given two disjoint sets A and B, if  $d(A, B) \leq \alpha$  for some  $\alpha > 0$ , then at most  $\sqrt{\alpha}|A|$  vertices  $a \in A$  satisfy  $\deg_B(a) < (1 - \sqrt{\alpha})|B|$ . Hence Part (2) immediately follows from Part (1).

We need a lower bound for the number of copies of  $K_k$  in a dense k-partite graph.

<span id="page-6-0"></span>**Proposition 11.** Let G be a k-partite graph with vertex class  $V_1, \dots, V_k$ . Suppose for every two vertex classes, the pairwise density  $d(V_i, V_j) \geq 1 - \alpha$  for some  $\alpha \leq (k+1)^{-4}$ , then there are at least  $\frac{1}{2} \prod_i |V_i|$  copies of  $K_k$  in G.

*Proof.* Given two disjoint sets  $V_i$  and  $V_j$ , if  $\bar{d}(V_i, V_j) \leq \alpha$  for some  $\alpha > 0$ , then at most  $\sqrt{\alpha}|V_i|$ vertices  $v \in V_i$  satisfy  $\deg_{V_j}(v) < (1 - \sqrt{\alpha})|V_j|$ . Thus, by choosing typical vertices greedily and the assumption  $\alpha \leq (k+1)^{-4}$ , there are at least

$$
(1 - \sqrt{\alpha})|V_1|(1 - 2\sqrt{\alpha})|V_2|\cdots(1 - k\sqrt{\alpha})|V_k| > (1 - (1 + \cdots + k)\sqrt{\alpha})\prod_i |V_i| > \frac{1}{2}\prod_i |V_i|
$$

copies of  $K_k$  in  $G$ .

Let  $\epsilon'' = 2k\sqrt{\epsilon'}$ . Now we want to study the structure of  $\tilde{V}_1$ .

<span id="page-6-1"></span>**Claim 12.** Given  $v \in \tilde{V}_1$  and  $2 \le i \le k$ , there exists  $j \in [k-1]$ , such that  $|N_{A_{ij}}(v)| < \epsilon''$ t.

*Proof.* Suppose instead, that there exist  $v \in V_1$  and some  $2 \leq i_0 \leq k$ , such that  $|N_{A_{i_0j}}(v)| \geq \epsilon'' t$  for all  $j \in [k-1]$ . By the minimum degree condition, for each  $2 \le i \le k$ , there is at most one  $j \in [k-1]$ . such that  $|N_{A_{ij}}(v)| < t/3$ . Therefore we can greedily choose  $k-2$  distinct  $j_i$  for  $i \neq i_0$ , such that  $|N_{A_{ij_i}}(v)| \ge t/3$ . Let  $j_{i_0}$  be the the (unique) unused index. Note that

$$
\forall i \neq i_0, \quad \frac{|A_{ij_i}|}{|N_{A_{ij_i}}(v)|} \le \frac{(1+\epsilon')t}{t/3} < 4, \quad \text{and} \quad \frac{|A_{i_0j_{i_0}}|}{|N_{A_{i_0j_{i_0}}}(v)|} \le \frac{(1+\epsilon')t}{\epsilon''t} < \frac{2}{\epsilon''}
$$

So for any  $i \neq i'$ , by Claim [10](#page-5-5) and the definition of  $\epsilon''$ , we have

<span id="page-6-2"></span>
$$
\bar{d}(N_{A_{ij_i}}(v), N_{A_{i'j_{i'}}}(v)) \le \frac{3\epsilon'|A_{ij_i}||A_{i'j_{i'}}|}{|N_{A_{ij_i}}(v)||N_{A_{i'j_{i'}}}(v)|} \le 3\epsilon' \cdot 4 \cdot \frac{2}{\epsilon''} = \frac{6}{k^2}\epsilon''.
$$
\n(9)

$$
\Box
$$

Since  $\epsilon \ll \epsilon'' \ll 1$ , by Proposition [11,](#page-6-0) there are at least

$$
\frac{1}{2} \prod_{i} N_{A_{ij_i}}(v) \ge \frac{1}{2} \cdot \epsilon'' t \left(\frac{t}{3}\right)^{k-2} = \frac{\epsilon''}{2 \cdot 3^{k-2}} t^{k-1} > \epsilon t^{k-1}
$$

copies of  $K_{k-1}$  in  $N(v)$ , contradicting the assumption  $v \in V_1$ .

Note that if  $\deg_{A_{ij}}(v) < \epsilon''t$ , at least  $|A_{ij}| - \epsilon''t$  vertices of  $A_{ij}$  are not in  $N(v)$ . By the minimum degree condition, [\(6\)](#page-5-3) and [\(8\)](#page-5-4), it follows that

<span id="page-7-0"></span>
$$
|A_{ij}^c \setminus N(v)| \le (1+\epsilon)t - (|A_{ij}| - \epsilon''t) \le (1+\epsilon)t - (1-\epsilon')t + \epsilon''t \le 2\epsilon''t.
$$
 (10)

Fix a vertex  $v \in \tilde{V}_1$ . Given  $2 \leq i \leq k$ , let  $\ell_i$  denote the (unique) index such that  $|N_{A_{i\ell_i}}(v)| < \epsilon'' t$ (the existence of  $\ell_i$  follows from Claim [12\)](#page-6-1).

# <span id="page-7-1"></span>Claim 13. We have  $\ell_2 = \ell_3 = \cdots = \ell_k$ .

*Proof.* Otherwise, say  $\ell_2 \neq \ell_3$ , then we set  $j_2 = \ell_3$  and for  $3 \leq i \leq k$ , greedily choose distinct  $j_k, j_{k-1}, \ldots, j_3 \in [k-1] \setminus \{\ell_3\}$  such that  $j_i \neq \ell_i$  (this is possible as  $j_3$  is chosen at last). Let us bound the number of copies of  $K_{k-1}$  in  $\bigcup_{i=2}^{k} N_{A_{ij_i}}(v)$ . By [10,](#page-7-0) we get  $|N_{A_{ij_i}}(v)| \geq |A_{ij_i}| - 2\epsilon''t \geq t/2$  for all *i*. As in [\(9\)](#page-6-2), for any  $i \neq i'$ , we derive that  $\bar{d}(N_{A_{ij_i}}(v), N_{A_{i'j_i'}}(v)) \leq 3\epsilon'' \cdot 4 \cdot 4 = 48\epsilon''$ . Since  $\epsilon'' \ll 1$ , by Proposition [11,](#page-6-0) we get at least  $\frac{1}{2} \left(\frac{t}{2}\right)^{k-1} > \epsilon t^{k-1}$  copies of  $K_{k-1}$  in  $N(v)$ , a contradiction.  $\square$ 

We define  $A_{1j} := \{v \in \tilde{V}_1 : |N_{A_{2j}}(v)| < \epsilon''t\}$  for  $j \in [k-1]$ . By Claims [12](#page-6-1) and [13,](#page-7-1) this yields a partition of  $\tilde{V}_1 = \bigcup_{j=1}^{k-1} A_{1j}$  such that

<span id="page-7-2"></span>
$$
d(A_{1j}, A_{ij}) < \frac{\epsilon''t|A_{1j}|}{|A_{1j}||A_{ij}|} \le \frac{\epsilon''t}{(1 - \epsilon')t} < (1 + 2\epsilon')\epsilon'' \quad \text{for} \quad i \ge 2 \quad \text{and} \quad j \ge 1. \tag{11}
$$

By [\(6\)](#page-5-3), [\(8\)](#page-5-4) and [\(10\)](#page-7-0), as  $(3k-5)\epsilon \leq \epsilon'$ , we have

$$
\bar{d}(A_{1j}, A_{ij'}) < \frac{|A_{1j}| 2\epsilon''t}{|A_{1j}| |A_{ij'}|} \le \frac{2\epsilon''t}{(1 - \epsilon')t} < 3\epsilon'' \quad \text{for } i \ge 2 \text{ and } j \ne j'.
$$
\n(12)

We claim  $|A_{1j}| \leq (1+\epsilon)t + (1+2\epsilon')\epsilon''|A_{1j}|$  for all j. Otherwise, by the minimum degree condition, we have  $\deg_{A_{1j}}(v) > (1+2\epsilon')\epsilon''|A_{1j}|$  for all  $v \in A_{ij}$ , and consequently  $d(A_{1j}, A_{ij}) > (1+2\epsilon')\epsilon''$ , contradicting [\(11\)](#page-7-2). We thus conclude that

<span id="page-7-3"></span>
$$
|A_{1j}| \le \frac{1+\epsilon}{1 - (1+2\epsilon')\epsilon''} t < (1+2\epsilon'')t. \tag{13}
$$

Since  $|V'_1| \leq \epsilon t$ , we have  $|\bigcup_{j=1}^{k-1} A_{1j}| = |V_1 \setminus V'_1| \geq |V_1| - \epsilon t$ . Using [\(13\)](#page-7-3), we now obtain a lower bound for  $|A_{1j}|, j \in [k-1]$ :

<span id="page-7-4"></span>
$$
|A_{1j}| \ge (k-1)(1-\epsilon)t - (k-2)(1+2\epsilon'')t - \epsilon t \ge (1-2k\epsilon'')t.
$$
 (14)

It remains to show that for  $2 \leq i \neq i' \leq k$ ,  $d(A_{i(k-1)}, A_{i'(k-1)})$  is small. Write  $N(v_1 \cdots v_m)$  $\bigcap_{1\leq i\leq m} N(v_i).$ 

<span id="page-7-5"></span>Claim 14.  $d(A_{i(k-1)}, A_{i'(k-1)}) \leq 6\epsilon''$  for  $2 \leq i, i' \leq k$ .

*Proof.* Suppose to the contrary, that say  $d(A_{(k-1)(k-1)}, A_{k(k-1)}) > 6\epsilon''$ . We first select  $k-2$  sets  $A_{ij}$  with  $1 \le i \le k-2$  and  $1 \le j \le k-2$  such that no two of them are on the same row or column – there are  $(k-2)!$  choices. Fix one of them, say  $A_{11}, A_{22}, \cdots, A_{(k-2)(k-2)}$ . We construct copies of  $K_{k-2}$  in  $A_{11} \cup A_{22} \cup \cdots \cup A_{(k-2)(k-2)}$  as follows. Pick arbitrary  $v_1 \in A_{11}$ . For  $2 \le i \le k-2$ , we select  $v_i \in N_{A_{ii}}(v_1 \cdots v_{i-1})$  such that  $v_i$  is  $\sqrt{3\epsilon'}$ -typical to  $A_{(k-1)(k-1)}$ ,  $A_{k(k-1)}$  and all  $A_{jj}$ ,  $i < j \le k-2$ . By Claim [10](#page-5-5) and [\(10\)](#page-7-0), there are at least  $(1 - (k-2)\sqrt{3\epsilon'})|A_{ii}| - 2\epsilon''t \ge t/2$  choices for each  $v_i$ . After selecting  $v_1, \ldots, v_{k-2}$ , we select adjacent vertices  $v_{k-1} \in A_{(k-1)(k-1)}$  and  $v_k \in A_{k(k-1)}$  such

$$
\frac{1}{1}.
$$

that  $v_{k-1}, v_k \in N(v_1 \cdots v_{k-2})$ . For  $j \in \{k-1, k\}$ , we know that  $N(v_1)$  misses at most  $2\epsilon''t$  vertices in  $A_{j(k-1)}$ , and at most  $(k-3)\sqrt{3\epsilon'}A_{j(k-1)}$  vertices of  $A_{j(k-1)}$  are not in  $N(v_2\cdots v_{k-2})$ . Since  $d(A_{(k-1)1}, A_{k1}) > 6\epsilon''$  and  $\epsilon'' = 2k\sqrt{\epsilon'},$  there are at least

$$
6\epsilon''|A_{(k-1)(k-1)}||A_{k(k-1)}| - 2\epsilon''t(|A_{(k-1)(k-1)}| + |A_{k(k-1)}|) - 2(k-3)\sqrt{3\epsilon'}|A_{(k-1)(k-1)}||A_{k(k-1)}|
$$
  
\n
$$
\geq (6\epsilon'' - 4\epsilon'' - 4(k-3)\sqrt{\epsilon'})|A_{(k-1)(k-1)}||A_{k(k-1)}|
$$
  
\n
$$
= 12\sqrt{\epsilon'}|A_{(k-1)(k-1)}||A_{k(k-1)}| \geq 6\sqrt{\epsilon'}t^2
$$

such pairs  $v_{k-1}, v_k$ . Together with the choices of  $v_1, \dots, v_{k-2}$ , we obtain at least  $(k-2)!(\frac{t}{2})^{k-2} 6\sqrt{\epsilon'}t^2 >$  $\epsilon t^k$  copies of  $K_k$ , a contradiction.

In summary, by [\(6\)](#page-5-3), [\(8\)](#page-5-4), [\(13\)](#page-7-3) and [\(14\)](#page-7-4), we have  $(1-2k\epsilon'')t \leq |A_{ij}| \leq (1+2\epsilon'')t$  for all i and j. In order to make  $\bigcup_{j=1}^{k-1} A_{1j}$  a partition of  $V_1$ , we move the vertices of  $V'_1$  to  $A_{11}$ . Since  $|V'_1| < \epsilon t$ , we still have  $||A_{ij} - t| \leq 2k\epsilon''t$  after moving these vertices. On the other hand, by [\(7\)](#page-5-2), [\(11\)](#page-7-2), and Claim [14,](#page-7-5) we have  $d(A_{ij}, A_{i'j}) \leq 6\epsilon'' \leq 2k\epsilon''$  for  $i \neq i'$  and all j (we now have  $d(A_{11}, A_{i1}) \leq 2\epsilon''$  for all  $i \geq 2$ because  $|A_{11}|$  becomes slightly larger). Therefore H is  $(2k\epsilon'', 2k\epsilon'')$ -approximate to  $\Theta_{k\times (k-1)}(t)$ . By the definitions of  $\epsilon''$  and  $\epsilon'$ ,

$$
2k\epsilon'' = 4k^2\sqrt{\epsilon'} = 4k^2\sqrt{16(k-1)^4(3\epsilon)^{1/2^{k-3}}} \le 16k^4\epsilon^{1/2^{k-2}},
$$

where the last inequality is equivalent to  $(\frac{k-1}{k})^2 3^{1/2^{k-2}} \le 1$  or  $3^{1/2^{k-1}} \le \frac{k}{k-1}$ , which holds because  $3 \leq 1 + \frac{2^{k-1}}{k-1} \leq (1 + \frac{1}{k-1})^{2^{k-1}}$  for  $k \geq 2$ . This completes the proof of Lemma [9.](#page-4-2)

We are ready to prove Lemma [7.](#page-3-0)

*Proof of Lemma [7.](#page-3-0)* First assume that  $G \in \mathcal{G}_3(n)$  is minimal, namely, G satisfies the minimum partite degree condition but removing any edge of G will destroy this condition. Note that this assumption is only needed by Claim [20.](#page-11-0)

Given  $0 < \Delta \leq 1$ , let

<span id="page-8-0"></span>
$$
\alpha = \frac{1}{2k} \left( \frac{\Delta}{24k(k-1)\sqrt{2k}} \right)^{2^{k-1}}.
$$
\n(15)

Without loss of generality, assume that  $x, y \in V_1$  and y is not reachable by  $\alpha^3 n^{k-1}$   $(k-1)$ -sets or  $\alpha^3 n^{2k-1}$  (2k – 1)-sets from x.

For  $2 \leq i \leq k$ , define

$$
A_{i1} = V_i \cap (N(x) \setminus N(y)), \quad A_{ik} = V_i \cap (N(y) \setminus N(x)),
$$
  
\n
$$
B_i = V_i \cap (N(x) \cap N(y)), \quad A_{i0} = V_i \setminus (N(x) \cup N(y)).
$$

Let  $B = \bigcup_{i \geq 2} B_i$ . If there are at least  $\alpha^3 n^{k-1}$  copies of  $K_{k-1}$  in B, then x is reachable from y by at least  $\alpha^3 n^{k-1}$  (k – 1)-sets. We thus assume there are less than  $\alpha^3 n^{k-1}$  copies of  $K_{k-1}$  in B.

Clearly, for  $i \geq 2$ ,  $A_{i1}$ ,  $A_{ik}$ ,  $B_i$  and  $A_{i0}$  are pairwise disjoint. The following claim bounds the sizes of  $A_{ik}$ ,  $B_i$  and  $A_{i0}$ .

Claim 15. (1)  $(1 - k^2 \alpha) \frac{n}{k} < |A_{i1}|, |A_{ik}| \le (1 + k \alpha) \frac{n}{k}$ ,  $(2)$   $(k-2-2k\alpha)\frac{n}{k} \leq |B_i| < (k-2+k(k-1)\alpha)\frac{n}{k},$ (3)  $|A_{i0}| < (k+1)\alpha n$ .

*Proof.* For  $v \in V$ , and  $i \in [k]$ , write  $N_i(v) := N(v) \cap V_i$ . By the minimum degree condition, we have  $|A_{i1}|, |A_{ik}| \leq (1/k + \alpha)n$ . Since  $N_i(x) = A_{i1} \cup B_i$ , it follows that

<span id="page-9-0"></span>
$$
|B_i| \ge \left(\frac{k-1}{k} - \alpha\right)n - \left(\frac{1}{k} + \alpha\right)n. \tag{16}
$$

If some  $B_i$ , say  $B_k$ , has at least  $(\frac{k-2}{k} + (k-1)\alpha)n$  vertices, then there are at least  $\prod_{i=2}^k |B_i|$  –  $(i-2)(\frac{1}{k}+\alpha)$  n copies of  $K_{k-1}$  in B. By [\(16\)](#page-9-0) and  $|B_k| \geq (\frac{k-2}{k}+(k-1)\alpha)n$ , this is at least

$$
\alpha n \cdot \prod_{i=2}^{k-1} \left( \frac{k-1}{k} - \alpha \right) n - (i-1) \left( \frac{1}{k} + \alpha \right) n
$$
  
= 
$$
\alpha n \cdot \prod_{i=2}^{k-1} \left( \frac{k-i}{k} - i\alpha \right) n
$$
  

$$
\geq \alpha n \cdot \prod_{i=2}^{k-1} \left( \frac{k-i-\frac{1}{2}}{k} \right) n \quad \text{because } 2k^2 \alpha \leq 1,
$$
  

$$
\geq \alpha n \cdot \frac{1}{2} \left( \frac{n}{k} \right)^{k-2}
$$
  

$$
\geq \alpha^2 n^{k-1} \quad \text{because } 2k^{k-2} \alpha \leq 1.
$$

This is a contradiction.

We may thus assume that  $|B_i| < (\frac{k-2}{k} + (k-1)\alpha)n$  for  $2 \le i \le k$ , as required for Part (2). As  $N_i(x) = A_{i1} \cup B_i$ , it follows that

$$
|A_{i1}| > (\frac{k-1}{k} - \alpha)n - (\frac{k-2}{k} + (k-1)\alpha)n = (\frac{1}{k} - k\alpha)n.
$$

The same holds for  $|A_{ik}|$  thus Part (1) follows. Finally

$$
|A_{i0}| = |V_i| - |N_i(x)| - |A_{ik}| < n - \left(\frac{k-1}{k} - \alpha\right)n - \left(\frac{1}{k} - k\alpha\right)n = (k+1)\alpha n,
$$
\nas required for Part (3).

Let  $t = n/k$  and  $\epsilon = 2k\alpha$ . By the minimum degree condition, every vertex  $u \in B$  is nonadjacent to at most  $(1 + k\alpha)n/k < (1 + \epsilon)t$  vertices in other color classes of B. By Claim 15,  $|B_i| \ge$  $(k-2-2k\alpha)\frac{n}{k} = (k-2-\epsilon)t \ge (k-2)(1-\epsilon)t$ . Thus  $G[B]$  is a  $(k-1)$ -partite graph that satisfies the assumptions of Lemma [9.](#page-4-2) We assumed that B contains less than  $\alpha^3 n^{k-1} < \epsilon^2 t^{k-1}$  copies of  $K_{k-1}$ , so by Lemma [9,](#page-4-2) B is  $(\alpha', \alpha')$ -approximate to  $\Theta_{(k-1)\times (k-2)}(\frac{n}{k})$ , where

$$
\alpha' := 16(k-1)^4 (2k\alpha)^{1/2^{k-3}}.
$$

This means that we can partition  $B_i$ ,  $2 \le i \le k$ , into  $A_{i2} \cup \cdots A_{i(k-1)}$  such that  $(1 - \alpha')\frac{n}{k} \le |A_{ij}| \le k$  $(1+\alpha')\frac{n}{k}$  for  $2 \leq j \leq k-1$  and

<span id="page-9-3"></span>
$$
\forall 2 \le i < i' \le k, 2 \le j \le k - 1, \quad d(A_{ij}, A_{i'j}) \le \alpha'.\tag{17}
$$

Together with Claim 15 Part (1), we obtain that (using  $k^2 \alpha \le \alpha'$ )

<span id="page-9-1"></span>
$$
\forall 2 \le i \le k, 1 \le j \le k, \quad (1 - \alpha')^{\frac{n}{k}} \le |A_{ij}| \le (1 + \alpha')^{\frac{n}{k}}.
$$
 (18)

Let  $A_{ij}^c = V_i \setminus A_{ij}$  denote the complement of  $A_{ij}$ . The following claim is an analog of Claim [10,](#page-5-5) and its proof is almost the same – after we replace  $(1+\epsilon)t$  with  $(1+k\alpha)n/k$  and  $\epsilon'$  with  $\alpha'$  (and we use  $\alpha \ll \alpha'$ ). We thus omit the proof.

<span id="page-9-2"></span>Claim 16. Let  $2 \le i \neq i' \le k$ ,  $1 \le j \neq j' \le k$ , and  $\{j, j'\} \neq \{1, k\}$ .

- (1)  $d(A_{ij}, A_{i'j'}) \ge 1 3\alpha'$  and  $d(A_{ij}, A_{i'j}^c) \ge 1 3\alpha'$ .
- (2) All but at most  $\sqrt{3\alpha'}$  vertices in  $A_{ij}$  are  $\sqrt{3\alpha'}$ -typical to  $A_{i'j'}$ ; at most  $\sqrt{3\alpha'}$  vertices in  $A_{ij}$ are  $\sqrt{3\alpha'}$ -typical to  $A_{i'j}^c$ .

Now let us study the structure of  $V_1$ . Let  $\alpha'' = 2k\sqrt{\alpha'}$ . Recall that  $N(xv) = N(x) \cap N(v)$ . Let  $V_1'$ be the set of the vertices  $v \in V_1$  such that there are at least  $\alpha n^{k-1}$  copies of  $K_{k-1}$  in each of  $N(xv)$ and  $N(yx)$ . We claim that  $|V'_1| < 2\alpha n$ . Otherwise, since a  $(k-1)$ -set intersects at most  $(k-1)n^{k-2}$ other  $(k-1)$ -sets, there are at least

$$
2\alpha n\cdot\alpha n^{k-1}(\alpha n^{k-1}-(k-1)n^{k-2})>\alpha^3n^{2k-1}
$$

copies of  $(2k-1)$ -sets connecting x and y, a contradiction.

Let  $\tilde{V}_1 := V_1 \setminus V'_1$ . The following claim is an analog of Claim [12](#page-6-1) for Lemma [9](#page-4-2) and can be proved similarly. The only difference between their proofs is that here we find at least  $\alpha n^{k-1}$  copies of  $K_{k-1}$ in each of  $N(xv)$  and  $N(yv)$ , which contradicts the definition of  $\tilde{V}_1$ .

<span id="page-10-0"></span>**Claim 17.** Given 
$$
v \in \tilde{V}_1
$$
 and  $2 \le i \le k$ , there exists  $j \in [k]$  such that  $|N_{A_{ij}}(v)| < \alpha''t$ .

Fix an vertex  $v \in V_1$ . Claim [17](#page-10-0) implies that for each  $2 \leq i \leq k$ , there exists  $\ell_i$  such that  $|N_{A_{i\ell_i}}(v)| < \alpha'' t$ . Our next claim is an analog of Claim [13](#page-7-1) for Lemma [9](#page-4-2) and can be proved similarly. **Claim 18.** We have  $\ell_2 = \ell_3 = \cdots = \ell_k$ .

<span id="page-10-1"></span>We now define  $A_{1j} := \{v \in \tilde{V}_1 : |N_{A_{2j}}(v)| < \alpha'' t\}$  for  $j \in [k]$ . By Claims [17](#page-10-0) and [18,](#page-10-1) this yields a partition of  $\tilde{V}_1 = \bigcup_{j=1}^k A_{1j}$  such that

<span id="page-10-3"></span>
$$
d(A_{1j}, A_{ij}) < \frac{\alpha''t|A_{1j}|}{|A_{1j}||A_{ij}|} \le \frac{\alpha''t}{(1-\alpha')t} < (1+2\alpha')\alpha'' \text{ for } i \ge 2 \text{ and } j \ge 1.
$$
 (19)

For  $v \in A_{1j}$ , we have  $|N_{A_{ij}}(v)| < \alpha'' t$  for  $i \geq 2$ . By the minimum degree condition and [\(18\)](#page-9-1),

<span id="page-10-2"></span>
$$
|A_{ij}^c \setminus N(v)| \le (\frac{1}{k} + \alpha)n - (|A_{ij}| - \alpha''t) < 2\alpha''t. \tag{20}
$$

By  $(18)$  and  $(20)$ , we derive that

$$
\bar{d}(A_{1j}, A_{ij'}) < \frac{|A_{1j}| \cdot 2\alpha''t}{|A_{1j}||A_{ij'}|} \le \frac{2\alpha''t}{(1 - \alpha')t} < 3\alpha'' \text{ for } i \ge 2 \text{ and } j \ne j'.
$$
\n(21)

We claim that  $|A_{1j}| \leq (1+\alpha)t + (1+2\alpha')\alpha''|A_{1j}|$  for all j. Otherwise, by the minimum degree condition, we have  $\deg_{A_{1j}}(v) > (1+2\alpha')\alpha''|A_{1j}|$  for all  $v \in A_{ij}$ , and consequently  $d(A_{1j}, A_{ij}) >$  $(1+2\alpha')\alpha''$ , contradicting [\(19\)](#page-10-3). We thus conclude that

<span id="page-10-4"></span>
$$
|A_{1j}| \le \frac{1+\alpha}{1 - (1 + 2\alpha')\alpha''}t < (1 + 2\alpha'')\frac{n}{k}.\tag{22}
$$

Since  $|V'_1| \leq 2\alpha n$ , we have  $|\bigcup_{j=1}^k A_{1j}| = |V_1 \setminus V'_1| \geq |V_1| - 2\alpha n$ . Using [\(22\)](#page-10-4), we now obtain a lower bound for  $|A_{1j}|, j \in [k]$ .

<span id="page-10-6"></span>
$$
|A_{1j}| \ge n - (k-1)(1 + 2\alpha'')\frac{n}{k} - 2\alpha n \ge (1 - 2k\alpha'')\frac{n}{k}.
$$
 (23)

It remains to show that  $d(A_{i1}, A_{i'1})$  and  $d(A_{ik}, A_{i'k})$ ,  $2 \leq i, i' \leq k$ , are small. First we show that if both densities are reasonably large then there are too many reachable  $(2k - 1)$ -sets from x to y. The proof resembles the one of Claim [14.](#page-7-5)

<span id="page-10-5"></span>Claim 19. For  $2 \leq i \neq i' \leq k$ , either  $d(A_{i1}, A_{i'1}) \leq 6\alpha''$  or  $d(A_{ik}, A_{i'k}) \leq 6\alpha''$ .

*Proof.* Suppose instead, that say  $d(A_{(k-1)1}, A_{k1}), d(A_{(k-1)k}, A_{kk}) > 6\alpha''$ . Fix a vertex  $v_1$  in  $A_{1j}$ , for some  $2 \le j \le k-1$ , say  $v_1 \in A_{12}$ . We construct two vertex disjoint copies of  $K_{k-1}$  in  $N(xv_1)$  and  $N(yv_1)$  as follows. We first select  $k-3$  sets  $A_{ij}$  with  $2 \leq i \leq k-2$  and  $3 \leq j \leq k-1$  such that no two of them are on the same row or column – there are  $(k-3)!$  choices. Fix one of them, say  $A_{23}, \cdots, A_{(k-2)(k-1)}$ . For  $2 \leq i \leq k-2$ , we select  $v_i \in N_{A_{i(i+1)}}(v_1 \cdots v_{i-1})$  that is  $\sqrt{3\alpha'}$ -typical to  $A_{(k-1)1}$ ,  $A_{k1}$  and  $A_{i(i+1)}$ ,  $i < j \leq k-2$ . By Claim [16](#page-9-2) and [\(20\)](#page-10-2), there are at least

$$
(1 - (k - 2)\sqrt{3\alpha'})|A_{i(i+1)}| - (k\alpha + \alpha' + \alpha'')\frac{n}{k} \ge \frac{n}{2k}
$$

such  $v_i$ . After selecting  $v_2, \ldots, v_{k-2}$ , we select two adjacent vertices  $v_{k-1} \in A_{(k-1)1}$  and  $v_k \in A_{k1}$ such that  $v_{k-1}$  and  $v_k$  are in  $N(v_1 \cdots v_{k-2})$ . For  $j = k-1, k$ , we know that  $N(v_1)$  misses at most  $(k\alpha + \alpha' + \alpha'')n/k$  vertices in  $A_{j1}$  and at most  $(k-3)\sqrt{3\alpha'}|A_{j1}|$  vertices of  $A_{j1}$  are not in  $N(v_2 \cdots v_{k-2})$ . Since  $d(A_{(k-1)1}, A_{k1}) > 6\alpha''$ , there are at least

$$
6\alpha''|A_{(k-1)1}||A_{k1}| - (k\alpha + \alpha' + \alpha'')\frac{n}{k}(|A_{(k-1)1}| + |A_{k1}|)
$$

$$
-2(k-3)\sqrt{3\alpha'}|A_{(k-1)1}||A_{k1}| \ge 6\sqrt{\alpha'}\left(\frac{n}{k}\right)^2
$$

such pairs  $v_{k-1}, v_k$ . Hence  $N(xv_1)$  contains at least

$$
(k-3)!\left(\frac{n}{2k}\right)^{k-3}6\sqrt{\alpha'}\left(\frac{n}{k}\right)^2 \ge \sqrt{\alpha'}\left(\frac{n}{k}\right)^{k-1} \ge \sqrt{\alpha}n^{k-1}
$$

copies of  $K_{k-1}$ . Let C be such a copy of  $K_{k-1}$ . Then we follow the same procedure and construct a copy of  $K_{k-1}$  on  $N(yv_1) \setminus C$ . After fixing  $k-3$  sets  $A_{ij}$  with  $2 \le i \le k-2$  and  $3 \le j \le k-1$ such that no two of them are on the same row or column, there are still at least  $\frac{n}{2k}$  such  $v_i$  for  $2 \leq i \leq k-2$ . Then, as  $d(A_{ik}, A_{i'k}) > 6\alpha''$ , there are at least  $6\sqrt{\alpha'} \left(\frac{n}{k}\right)^2$  choices of  $v_{k-1} \in A_{(k-1)k}$ and  $v_k \in A_{kk}$  such that  $v_{k-1}$  and  $v_k$  are in  $N(v_1 \cdots v_{k-2})$ . This gives at least  $\sqrt{\alpha} n^{k-1}$  copies of  $K_{k-1}$  in  $N(yv_1)$ . Then, since there are at least  $|V_1| - |A_{11}| - |A_{1k}| \ge \alpha n$  choices of  $v_1$ , totally there are at least  $\alpha n(\sqrt{\alpha}n^{k-1})^2 = \alpha^2 n^{2k-1}$  reachable  $(2k-1)$ -sets from x to y, a contradiction.

Next we show that if any of  $d(A_{i1}, A_{i'1})$  or  $d(A_{ik}, A_{i'k})$ ,  $2 \leq i, i' \leq k$ , is sufficiently large, then we can remove edges from  $G$  such that the resulting graph still satisfies the minimum degree condition, which contradicts the assumption that  $G$  is minimal.

<span id="page-11-0"></span>Claim 20. For  $2 \le i \ne i' \le k$ ,  $d(A_{i1}, A_{i'1}), d(A_{ik}, A_{i'k}) \le 6k\sqrt{\alpha''}$ .

*Proof.* Without loss of generality, assume that  $d(A_{2k}, A_{3k}) > 6k\sqrt{\alpha''}$ . By Claim [19,](#page-10-5) we have  $d(A_{21}, A_{31}) < 6\alpha''$ . Combining this with [\(17\)](#page-9-3), we have  $d(A_{2j}, A_{3j}) < 6\alpha''$  for all  $j \in [k-1]$ . Now fix  $j \in [k-1]$ . The number of non-edges between  $A_{2j}$  and  $A_{3j}$  satisfies  $\bar{e}(A_{2j}, A_{3j}) > (1-6\alpha'')|A_{2j}||A_{3j}|$ . By the minimum degree condition and [\(18\)](#page-9-1),

$$
\bar{e}(A_{2k},A_{3j}) < (1+ \ k\alpha)\frac{n}{k} |A_{3j}| - (1- 6 \alpha'')|A_{2j}||A_{3j}| \leq 7 \alpha'' \frac{n}{k} |A_{3j}|.
$$

Using [\(18\)](#page-9-1) again, we obtain that

$$
d(A_{2k},A_{3j}) \geq 1 - \frac{7\alpha''\frac{n}{k}|A_{3j}|}{|A_{2k}||A_{3j}|} \geq 1 - 8\alpha''.
$$

This implies that  $d(A_{2k}, A_{3k}^c) \geq 1-8\alpha''$ . Similarly we derive that  $d(A_{3k}, A_{2k}^c) \geq 1-8\alpha''$ . For  $i = 2, 3$ , define  $A_{ik}^T$  as the set of the vertices in  $A_{ik}$  that are  $\sqrt{8\alpha''}$ -typical to  $A_{(5-i)k}^c$ . Thus  $|A_{ik} \setminus A_{ik}^T| \le$  $\sqrt{8\alpha''}|A_{ik}|.$ 

Let 
$$
A_{ik}^{T_1} = \{v \in A_{ik}^T : \deg_{A_{(5-i)k}}(v) \le \sqrt{8\alpha''}|A_{jk}^c|\}
$$
 and  $A_{ik}^{T_2} = A_{ik}^T \setminus A_{ik}^{T_1}$ . For  $u \in A_{2k}^{T_2}$ , we have  
\n
$$
\deg_{V_3}(u) = \deg_{A_{3k}^c}(u) + \deg_{A_{3k}}(u) > (1 - \sqrt{8\alpha''})|A_{3k}^c| + \sqrt{8\alpha''}|A_{3k}^c| = |A_{3k}^c|.
$$

Since  $|A_{3k}^c| \ge \deg_{V_3}(x)$  and  $|A_{3k}^c|$  is an integer, we conclude that  $\deg_{V_3}(u) \ge \deg_{V_3}(x) + 1$ . Similarly we can derive that  $\deg_{V_2}(v) \ge \deg_{V_2}(x) + 1$  for every  $v \in A_{3k}^{T_2}$ . If there is an edge uv joining some  $u \in A_{2k}^{T_2}$  and some  $v \in A_{3k}^{T_2}$ , then we can delete this edge and the resulting graph still satisfies the minimum degree condition. Therefore we may assume that there is no edge between  $A_{2k}^{T_2}$  and  $A_{3k}^{T_2}$ . Then

$$
e(A_{2k}, A_{3k}) \le e(A_{2k} \setminus A_{2k}^T, A_{3k}) + e(A_{2k}, A_{3k} \setminus A_{3k}^T) + e(A_{2k}^T, A_{3k}^T) + e(A_{2k}^T, A_{3k}^T)
$$
  
\n
$$
\le 2\sqrt{8\alpha''}|A_{2k}||A_{3k}| + |A_{2k}^T|\sqrt{8\alpha''}|A_{3k}^c| + |A_{3k}^T|\sqrt{8\alpha''}|A_{2k}^c|
$$
  
\n
$$
\le \sqrt{8\alpha''}(2|A_{2k}||A_{3k}| + |A_{2k}||A_{3k}^c| + |A_{3k}||A_{2k}^c|)
$$
  
\n
$$
= \sqrt{8\alpha''}(|A_{2k}||V_3| + |A_{3k}||V_2|)
$$
  
\n
$$
\le 3\sqrt{\alpha''} \cdot 2k|A_{2k}||A_{3k}| \text{ by (18)}.
$$
  
\n
$$
\Rightarrow d(A_{2k}, A_{3k}) \le 6k\sqrt{\alpha''}.
$$

Therefore  $d(A_{2k}, A_{3k}) \leq 6k\sqrt{\alpha''}$ 

In summary, by [\(18\)](#page-9-1), [\(22\)](#page-10-4) and [\(23\)](#page-10-6), we have  $(1 - 2k\alpha'')\frac{n}{k} \le |A_{ij}| \le (1 + 2\alpha'')\frac{n}{k}$  for all i and j. In order to make  $\bigcup_{j=1}^k A_{ij}$  a partition of  $V_i$ , we move the vertices of  $V'_1$  to  $A_{11}$  and the vertices of  $A_{i0}$  to  $A_{i2}$  for  $2 \leq i \leq \tilde{k}$ . Since  $|V_1'| < 2\alpha n$  and  $|A_{i0}| \leq (k+1)\alpha n$ , we have  $||A_{ij}|- \frac{n}{k}| \leq 2k\alpha'' \frac{n}{k}$  after moving these vertices. On the other hand, by [\(17\)](#page-9-3), [\(19\)](#page-10-3), and Claim [20,](#page-11-0) we have  $d(A_{ij}, A_{i'j}) \leq 6k\sqrt{\alpha''}$  for  $i \neq i'$  and all j. (In fact, for  $i \geq 2$ , we now have  $d(A_{11}, A_{i1}) \leq 2\alpha''$  as we added at most  $2\alpha n$ vertices to  $A_{11}$ . For  $i' > i \geq 2$ , we now have  $d(A_{12}, A_{i2}) \leq 2\alpha''$  and  $d(A_{i2}, A_{i2}) \leq 2\alpha'$  as we moved at most  $(k + 1)\alpha n$  vertices to  $A_{i2}$ .) Therefore after deleting edges, G is  $(2k\alpha'', 6k\sqrt{\alpha''})$ approximate to  $\Theta_{k\times k}(n/k)$ . By [\(15\)](#page-8-0), and the definitions of  $\alpha''$  and  $\alpha'$ , G is ( $\Delta/6$ ,  $\Delta/2$ )-approximate to  $\Theta_{k\times k}(n/k)$ .

#### **ACKNOWLEDGMENTS**

The authors thank two referees for their valuable comments that improved the presentation and shortened the proofs of Lemmas [7](#page-3-0) and [9.](#page-4-2)

#### **REFERENCES**

- <span id="page-12-15"></span><span id="page-12-2"></span>[1] N. Alon and R. Yuster, H-factors in dense graphs. J. Combin. Theory Ser. B, 66 (1996), no. 2, 269–282.
- <span id="page-12-14"></span>[2] A. Bush and Y. Zhao, Minimum degree thresholds for bipartite graph tiling, J. Graph Theory, to appear.
- [3] A. Czygrinow, L. DeBiasio. A note on bipartite graph tiling, SIAM J. Discrete Math.
- <span id="page-12-6"></span>[4] G. A. Dirac. Some theorem on abstract graphs. Proc. London Math. Soc.  $(3)$ , 2,  $(1952)$  69–81.
- <span id="page-12-0"></span>[5] E. Fischer, Variants of the Hajnal-Szemerédi theorem. *J. Graph Theory* 31 (1999), no. 4, 275–282.
- [6] A. Hajnal and E. Szemerédi, Proof of a conjecture of P. Erdös. Combinatorial theory and its applications, II  $(Proc. Collog., Balatonfüred, 1969), pp. 601–623. North-Holland, Amsterdam, 1970.$
- <span id="page-12-16"></span>[7] H. Hàn, Y. Person and M. Schacht, On perfect matchings in uniform hypergraphs with large minimum vertex degree, SIAM J. Discrete Math. 23 (2009), 732–748.
- <span id="page-12-13"></span><span id="page-12-7"></span>[8] J. Hladký and M. Schacht. Note on bipartite graph tilings, SIAM J. Discrete Math, 24, (2010), no. 2, 357362.
- <span id="page-12-9"></span>[9] P. Keevash and R. Mycroft, A geometric theory for hypergraph matching, submitted.
- <span id="page-12-1"></span>[10] P. Keevash and R. Mycroft, A multipartite Hajnal-Szemeredi theorem, submitted.
- [11] H. Kierstead and A. Kostochka. A short proof of the Hajnal-Szemerédi theorem on equitable colouring. Combinatorics, Probability and Computing, 17 (2008), no. 2, 265–270.
- <span id="page-12-11"></span>[12] J. Komlós, Tiling Turán theorems. Combinatorica  $20$  (2000), no. 2, 203–218.
- [13] J. Komlós, G. N. Sárközy, E. Szemerédi. On the square of a Hamiltonian cycle in dense graphs. Random Struct. Algorithms 9 (1996) 193–211.
- <span id="page-12-12"></span><span id="page-12-3"></span>[14] J. Komlós, G. N. Sárközy and E. Szemerédi, Blow-up Lemma. Combinatorica  $17$  (1997), no. 1, 109–123.
- [15] J. Komlós, G. N. Sárközy and E. Szemerédi, Proof of the Alon-Yuster conjecture. Combinatorics (Prague, 1998). Discrete Math. 235 (2001), no. 1-3, 255–269.
- <span id="page-12-4"></span>[16] D. Kühn and D. Osthus, The minimum degree threshold for perfect graph packings, *Combinatorica* 29 (2009), 65–107.
- <span id="page-12-5"></span>[17] D. Kühn and D. Osthus, Embedding large subgraphs into dense graphs, Surveys in Combinatorics, Cambridge University Press, 2009, 137–167.
- <span id="page-12-10"></span>[18] I. Levitt, G. N. Sárközy, E. Szemerédi. How to avoid using the regularity lemma: Pósa's conjecture revisited. Discrete Math. 310 (2010), no. 3, 630–641.
- <span id="page-12-8"></span>[19] A. Lo and K. Markström, A multipartite version of the Hajnal-Szemerédi theorem for graphs and hypergraphs, preprint.

## 14 JIE HAN AND YI ZHAO

- <span id="page-13-1"></span>[20] Cs. Magyar, R. Martin, Tripartite version of the Corrádi-Hajnal theorem. Discrete Math. 254 (2002), no. 1-3, 289–308.
- [21] R. Martin, E. Szemerédi, Quadripartite version of the Hajnal-Szemerédi theorem. Discrete Math. 308 (2008), no. 19, 4337–4360.
- <span id="page-13-6"></span>[22] R. Martin, Y. Zhao, Tiling tripartite graphs with 3-colorable graphs. Electronic Journal of Combinatorics 16, (2009).
- <span id="page-13-4"></span>[23] V. Rödl, A. Ruciński, E. Szemerédi, A Dirac-type theorem for 3-uniform hypergraphs, Combin. Probab. Comput. 15 (2006), no. 1- 2, 229–251.
- <span id="page-13-3"></span>[24] A. Shokoufandeh, Y. Zhao, Proof of a tiling conjecture of Komlós, Random Structures Algorithms 23 (2003), no. 2, 180–205.
- <span id="page-13-2"></span>[25] E. Szemerédi, Regular partitions of graphs. Problèmes combinatoires et théorie des graphes (Colloq. Internat. CNRS, Univ. Orsay, Orsay, 1976), pp. 399–401, Colloq. Internat. CNRS, 260, CNRS, Paris, 1978.
- <span id="page-13-5"></span>[26] Y. Zhao, Tiling bipartite graphs, SIAM J. Discrete Math. 23 (2009), no 2, 888–900.

(Jie Han and Yi Zhao) DEPARTMENT OF MATHEMATICS AND STATISTICS, Georgia State University, Atlanta, GA 30303 E-mail address, Jie Han: jhan22@gsu.edu

E-mail address, Yi Zhao: yzhao6@gsu.edu

<span id="page-13-0"></span>