# The Automorphism Group of an Extremal [72 , 36 , 16] Code does not contain elements of order 6

Martino Borello

Dipartimento di Matematica e Applicazioni, Università Milano Bicocca 20125 Milano, Italy m.borello1@campus.unimib.it

Version of March 15 2012, printed August 29, 2018

#### Abstract

The existence of an extremal code of length  $72$  is a long-standing open problem. Let  $C$  be a putative extremal code of length 72 and suppose that  $\mathcal C$  has an automorphism  $g$  of order 6. We show that  $\mathcal{C}$ , as an  $\mathbb{F}_2\langle g \rangle$ -module, is the direct sum of two modules, one easily determinable and the other one which has a very restrictive structure. We use this fact to do an exhaustive search and we do not find any code. This prove that the automorphism group of an extremal code of length 72 does not contain elements of order 6 .

KEY WORDS: automorphism group, extremal code of length 72

### 1 Introduction

The existence of an extremal code of length 72 is a long-standing open problem [\[18\]](#page-14-0). A series of papers investigates the structure of its automorphism group excluding most of the subgroup of  $S_{72}$ . The most recent result, established in [\[9\]](#page-14-1) and [\[14\]](#page-14-2), is the following:

The automorphism group of a binary self-dual doubly-even  $[72, 36, 16]$  code C has order 5 or a divisor of 24. Furthermore, if 8 divides the order of Aut(C) then its Sylow 2-subgroup is either  $D_8$  or  $C_2 \times C_2 \times C_2$ .

In this paper we will prove that the automorphism group of a binary self-dual doubly-even [72, 36, 16] code does not contain elements of order 6 , obtaining the following.

**Theorem 1.1.** The automorphism group of a binary self-dual doubly-even [72,36,16] code is either trivial or one of the following:  $C_2, C_3, C_4, C_2 \times C_2, C_5, S_3, D_8, C_2 \times C_2 \times C_2, A_4$  or  $S_4$ .

Notations:  $C_n$  is the cyclic group of order  $n$ ;  $D_n$  is the dihedral group of order  $n$ ;  $\mathcal{A}_n$  and  $\mathcal{S}_n$  are, respectively, the alternating group and the symmetric group of degree n.

With methods similar to those introduced by O'Brien and Willems in [\[15\]](#page-14-3), we strongly use the fact that a binary code with automorphism group G is an  $\mathbb{F}_2G$ -module. Consider g, automorphism of order 6 : we use a variety of results, some new, of modular representation theory, to study the structure of  $\mathbb{F}_2\langle g \rangle$ -modules (for basic concepts of representation theory see the introduction of [\[7\]](#page-14-4)). In particular we show that our putative code is the direct sum of two modules: one is the subcode of words fixed by  $g^2$ , easily determinable, and the other one has a socle which belongs, up to equivalence, to a very small set. From the knowledge of the socle it is quite easy to do an exhaustive search.

All computations were carried out using MAGMA [\[3\]](#page-13-0).

### 2 Preliminaries

Let C be a binary  $[n, k]$  linear code and let  $( , ) : \mathbb{F}_2^n \times \mathbb{F}_2^n \to \mathbb{F}_2$  be an inner product on  $\mathbb{F}_2^n$ , where  $\mathbb{F}_2$  is the field with 2 elements. Then

$$
\mathcal{C}^{\perp} = \{ v \in \mathbb{F}_2^n \mid (v, c) = 0 \ \forall c \in \mathcal{C} \}
$$

is the dual of C, a binary  $[n, n - k]$  linear code. C terms self-orthogonal if  $C \subseteq C^{\perp}$  and self-dual if  $C = C^{\perp}$ . In this paper we are interested in binary linear codes in  $\mathbb{F}_2^n$  with the Euclidean inner product; i.e., if  $u = (u_1, u_2, \dots, u_n), v = (v_1, v_2, \dots, v_n) \in \mathbb{F}_2^n$ , then  $(u, v) = \sum_{i=0}^n u_i v_i$ .

A theorem of Gleason, Pierce and Turyn [\[2\]](#page-13-1) guarantees that if  $s > 1$  divides the weight of every codeword in a nontrivial binary self-dual code, then  $s = 2$  or 4. Binary self-dual codes automatically satisfy this condition with  $s = 2$ . A binary doubly-even code is a binary linear code whose words have weight divisible by 4. Self-dual doubly-even codes exist only when n is a multiple of  $8 \text{ [10]}$  $8 \text{ [10]}$  $8 \text{ [10]}$ . A theorem of Mallows and Sloane [\[12\]](#page-14-6) shows that for a self-dual doubly-even code

$$
d\leq 4\left[\frac{n}{24}\right]+4
$$

where [x] is the integer part of x. If  $d = 4\left[\frac{n}{24}\right] + 4$ , C is called extremal. Extremal codes whose length is a multiple of 24 are very interesting, since, for example, all their codewords of a given weight support five-designs [\[1\]](#page-13-2), [\[17\]](#page-14-7). The only known extremal codes of length a multiple of 24 are the extended binary Golay code  $\mathcal{G}_{24}$  and the extended quadratic residue code  $QR_{48}$ , the unique (up to equivalence) extremal codes of length 24 and 48 respectively.

There is a natural (right) action of  $S_n$  on  $\mathbb{F}_2^n$  (action on the coordinates): if  $v = (v_1, v_2, \ldots, v_n)$  and  $g \in \mathcal{S}_n$ , define

$$
v^g = (v_{g^{-1}(1)}, v_{g^{-1}(2)}, \dots, v_{g^{-1}(n)}).
$$

If C is a binary code and  $c^g \in \mathcal{C}$  for all  $c \in \mathcal{C}$ , then g is an automorphism of C. We denote with Aut $(\mathcal{C})$  the group  $(\leq \mathcal{S}_n)$  of the automorphisms of  $\mathcal{C}$ .

When  $\mathcal C$  is a binary self-dual doubly-even [72, 36, 16] code, the possible structure of the elements of Aut(C) is well-known: we say that an automorphism h is of type  $p - (c, f)$  if h has c cycles of order p and f fixed points. Then we have the following result [\[6\]](#page-13-3), [\[5\]](#page-13-4), [\[8\]](#page-14-8).

<span id="page-1-0"></span>Proposition 2.1. Let h be an automorphism of prime order of a binary self-dual doubly-even [72, 36, 16] code. Then h can be only of the following types:  $2 - (36, 0)$ ,  $3 - (24, 0)$  or  $5 - (14, 2)$ .

Let us fix some notations that we will use throughout this paper.

If  $v = (v_1, \ldots, v_n) \in \mathbb{F}_2^n$  and  $\Omega = \{j_1, \ldots, j_m\} \subseteq \{1, \ldots, n\}$ , we define  $v_{|_{\Omega}} = (v_{j_1}, \ldots, v_{j_m})$ .

If W is a subspace of  $\mathbb{F}_2^n$  and  $h \in \mathcal{S}_n$  define

$$
\mathcal{W}(h) = \{ w \in \mathcal{W} \mid w^h = w \}.
$$

If  $\Omega_1, \ldots, \Omega_{n_h}$  are the orbits of the action of h on the coordinates, we say that a  $v \in V$  is constant on the orbits of h if  $v_{\vert \Omega_i}$  is either null or the all ones vector. Obviously,  $w \in \mathcal{W}$  belongs to  $\mathcal{W}(h)$  if and only if is constant on the orbits of  $h$ .

If  $W$  is a subspace of  $\mathbb{F}_2^n$ , we define

$$
\mathcal{W} \otimes \langle (\underbrace{1,\ldots,1}_{m \text{ times}}) \rangle
$$

to be the subspace of  $\mathbb{F}_2^{n \cdot m}$  obtained extending every vector of W substituting every one with  $m$ ones and every zero with  $m$  zeros.

### 3 Fixed codes

An usual starting point in the search of a code with certain automorphisms is the study of the subcode of words fixed by these automorphisms. This is, in general, an easy problem. In our case it will provide a fundamental tool to do the exhaustive search.

Let C be a self-dual doubly-even [72, 36, 16] binary code and suppose  $g \in Aut(\mathcal{C})$  such that  $o(g) = 6$ . By Proposition [2.1](#page-1-0) we have that the elements of order 2 and 3 in Aut( $\mathcal{C}$ ) have no fixed points, so g has no fixed point. Thus we can suppose, up to a relabelling of the coordinates,

$$
g = (1, 2, 3, 4, 5, 6) \dots (67, 68, 69, 70, 71, 72)
$$

We have the following known results about  $C(g^2)$  [\[9\]](#page-14-1) and  $C(g^3)$  [\[14\]](#page-14-2).

<span id="page-2-0"></span>**Proposition 3.1.** The code  $C(g^2)$  is equivalent to  $\mathcal{G}_{24} \otimes \langle (1,1,1) \rangle$ , with  $\mathcal{G}_{24}$  extended binary Golay code (remember that all the self-dual doubly-even [24, 12, 8] codes are equivalent and they are all called extended binary Golay codes).

<span id="page-2-1"></span>**Proposition 3.2.** The code  $C(g^3)$  is equivalent to  $K \otimes \langle (1,1) \rangle$ , with K one of the 41 self-dual [36, 18, 8] codes classified by Mechor and Gaborit [\[13\]](#page-14-9).

With argument similar to the ones used in [\[4\]](#page-13-5) and [\[14\]](#page-14-2) we prove the following result about  $\mathcal{C}(g)$ .

<span id="page-2-2"></span>**Proposition 3.3.** The code  $\mathcal{C}(g)$  is  $\mathcal{H} \otimes \langle (1, 1, 1, 1, 1, 1), \rangle$ , where H is equivalent to F, binary selfdual [12, 6, 4] code with generator matrix

$$
M = \left[\begin{smallmatrix} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \end{smallmatrix}\right].
$$

*Proof.* Let  $\pi_{12} : \mathbb{F}_2^{72} \to \mathbb{F}_2^{12}$  the projection such that

$$
(v_1, v_2, v_3, v_4, v_5, v_6, \ldots, v_{67}, v_{68}, v_{69}, v_{70}, v_{71}, v_{72}) \mapsto (v_1, \ldots, v_{67})
$$

and  $\phi: \mathbb{F}_2^{72} \to \mathbb{F}_2^{12}$  the map

$$
(v_1, v_2, v_3, v_4, v_5, v_6, \ldots, v_{67}, v_{68}, v_{69}, v_{70}, v_{71}, v_{72}) \mapsto \left(\sum_{i=1}^{6} v_i, \ldots, \sum_{i=67}^{72} v_i\right).
$$

The code  $\phi(\mathcal{C})$  is self-orthogonal: let  $c, c' \in \mathcal{C}$ , then

$$
(\phi(c), \phi(c')) = \sum_{j=1}^{12} \left( \sum_{i=6j-5}^{6j} c_i \right) \cdot \left( \sum_{i=6j-5}^{6j} c'_i \right) = \sum_{k=0}^{5} (c^{g^k}, c') = 0.
$$

Furthermore we have  $\pi_{12}(\mathcal{C}(g)) = \phi(\mathcal{C})^{\perp}$ : let  $f = (f_1, f_1, f_1, f_1, f_1, f_1, \ldots, f_{12}, f_{12}, f_{12}, f_{12}, f_{12}, f_{12}) \in \mathcal{C}(g)$  and  $c \in \mathcal{C}$ . Then

$$
(\pi_{12}(f), \phi(c)) = \sum_{j=1}^{12} f_j \cdot \left(\sum_{i=6j-5}^{6j} c_i\right) = (f, c) = 0,
$$

so  $\pi_{12}(\mathcal{C}(g)) \subseteq \phi(\mathcal{C})^{\perp}$ .

Viceversa, let  $v = (v_1, \ldots, v_{12}) \in \phi(\mathcal{C})^{\perp}$ . Let  $\overline{v} = (v_1, v_1, v_1, v_1, v_1, v_1, \ldots, v_{12}, v_{12}, v_{12}, v_{12}, v_{12}, v_{12})$ such that  $v = \pi_{12}(\overline{v})$ . We have

$$
0 = (v, \phi(c)) = (\overline{v}, c)
$$

for all  $c \in \mathcal{C} = \mathcal{C}^{\perp}$ , so  $\overline{v} \in \mathcal{C}$ .

Thus  $v \in \pi_{12}(\mathcal{C}(g))$  and so  $\pi_{12}(\mathcal{C}(g)) = \phi(\mathcal{C})^{\perp}$ .

 $\pi_{12}(\mathcal{C}(g))$  is the dual of a self-orthogonal code. It is, obviously, even and of minimum distance bigger or equal to 4, since every word in  $\mathcal{C}(g)$  has weight a multiple of 6. Thus it is a [12,  $\geq 6, \geq 4$ ] even code.

We have just proved that  $\phi(\mathcal{C}) \subseteq \pi_{12}(\mathcal{C}(g)) = \phi(\mathcal{C})^{\perp}$ . Let us suppose  $\phi(\mathcal{C}) \subsetneq \pi_{12}(\mathcal{C}(g)) = \phi(\mathcal{C})^{\perp}$ . So there exists  $v \in \phi(\mathcal{C})^{\perp} \setminus \phi(\mathcal{C})$ . Denote  $D = \langle \phi(\mathcal{C}), v \rangle$ . This is obviously a self-orthogonal code. If it is not self-dual we can repeat this algorithm. So we can find a self-dual code  $D'$  such that  $\phi(\mathcal{C}) \subset D' = D'^{\perp} \subset \phi(\mathcal{C})^{\perp}$ . We have that  $D'$  has minimum distance at least 4. There is, up to equivalence, only one self-dual [12, 6] code with minimum distance bigger or equal to 4 [\[16\]](#page-14-10), and has generator matrix

$$
M = \left[\begin{smallmatrix} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \end{smallmatrix}\right].
$$

This code has no overcode of minimum distance bigger or equal to 4. So  $\phi(\mathcal{C}) = \pi_{12}(\mathcal{C}(g)) = \phi(\mathcal{C})^{\perp}$ and it is equivalent to a self-dual code with generator matrix equivalent to M. Setting  $\mathcal{H} = \pi_{12}(\mathcal{C}(g))$ we have the thesis. П

Let us underline the following obvious fact:

<span id="page-3-0"></span>
$$
\mathcal{C}(g) = \mathcal{C}(g^2) \cap \mathcal{C}(g^3). \tag{1}
$$

This simply observation will be fundamental in section [6.](#page-8-0)

## <span id="page-3-1"></span>4 Decomposition of C as  $\mathbb{F}_2\langle q \rangle$ -module

From now on set  $V = \mathbb{F}_2^{72}$ , C our putative binary self-dual doubly-even [72, 36, 16] code and g an automorphism of  $C$  of order 6.

Following [\[15\]](#page-14-3), we observe that  $V$  is an  $\mathbb{F}_2\langle g \rangle$ -module defining the product in the natural way:

$$
v \cdot \left(\sum_{i=0}^{5} a_i g^i\right) = \sum_{i=0}^{5} a_i v^{g^i} \quad \text{for } v \in \mathcal{V} \text{ and } a_i \in \mathbb{F}_2.
$$

The code C is an  $\mathbb{F}_2\langle g \rangle$ -submodule of V, since g is an automorphism of the code. Obviously any  $\mathbb{F}_2\langle g \rangle$ -module is also an  $\mathbb{F}_2\langle g^2 \rangle$ -module.

In the previous section we considered the fixed codes  $\mathcal{C}(g)$ ,  $\mathcal{C}(g^2)$  and  $\mathcal{C}(g^3)$ . In particular  $\mathcal{C}(g^2)$ plays an important role in our method, since  $g^2$  has order 3 and we have a classical result of Huffman about the decomposition of binary codes with automorphisms of odd order.

**Lemma 4.1.** (cfr. Lemma 2 in [\[11\]](#page-14-11)) Let C be a binary linear code with an automorphism h of odd order. Then

$$
\mathcal{C} = \mathcal{C}(h) \oplus \mathcal{E}(h)
$$

where  $\mathcal{E}(h)$  is the subcode of C of words of even weight on the orbits of h, i.e.

$$
\mathcal{E}(h) = \{c \in \mathcal{C} \mid \text{wt}(c_{|\Omega_i}) \equiv 0 \ (mod \ 2), \ \forall i\}
$$

where  $\Omega_i$  are the orbits of h on the coordinates.

So, in our case,

$$
\mathcal{C} = \mathcal{C}(g^2) \oplus \mathcal{E}(g^2).
$$

Let us observe that  $\dim \mathcal{E}(g^2) = \dim \mathcal{C} - \dim \mathcal{C}(g^2) = 36 - 12 = 24$ .

In order to have more information about  $\mathcal{E}(g^2)$ , we need to reinterpret this decomposition in terms of representation theory. A basic result in this direction is given by Lemma [4.2](#page-4-0) below. We state it following the notations in [\[15\]](#page-14-3).

<span id="page-4-0"></span>**Lemma 4.2.** Let C be a binary linear code,  $G \le \text{Aut}(\mathcal{C})$  and

$$
1 = f_1 + \ldots + f_t
$$

be a decomposition of  $1 \in \mathbb{F}_2G$  into central orthogonal idempotents  $f_i \in \mathbb{F}_2G$ . Set  $\mathcal{V}_i = \mathcal{V}f_i$  and  $\mathcal{C}_i = \mathcal{C}f_i \subseteq \mathcal{V}_i$  for  $i \in \{1, ..., t\}$ . Then

$$
\mathcal{V} = \mathcal{V}_1 \oplus \ldots \oplus \mathcal{V}_t \qquad and \qquad \mathcal{C} = \mathcal{C}_1 \oplus \ldots \oplus \mathcal{C}_t
$$

as  $\mathbb{F}_2G$  -modules.

Let us take  $G = \langle g \rangle$ . Then

$$
f_1 = 1 + g^2 + g^4
$$
 and  $f_2 = g^2 + g^4$ 

is a decomposition of 1 into (central) orthogonal idempotents of  $\mathbb{F}_2\langle g \rangle$ .

<span id="page-4-2"></span>Remark 4.1. Observing the proof of Lemma 2 in [\[11\]](#page-14-11), it is easy to prove the following facts:

- 1.  $C_1 = C f_1 = C(g^2)$  and  $C_2 = C f_2 = \mathcal{E}(g^2)$ ;
- 2.  $\mathcal{V}_1 = \mathcal{V}f_1 = \mathcal{V}(g^2)$ , the subspace of all the vectors fixed by  $g^2$ ;
- 3.  $\mathcal{V}_2 = \mathcal{V}f_2$  is the set of vectors of even weight on the orbits of  $g^2$ .

They are all  $\mathbb{F}_2\langle g \rangle$  -modules.

<span id="page-4-1"></span>**Remark 4.2.** By direct calculations we deduce the following properties for the principal ideal  $\mathcal{I} =$  $(f_2) \subset \mathbb{F}_2\langle g \rangle$ :

- 1.  $\mathcal I$  is 4-dimensional, viewed as an  $\mathbb F_2$ -vector space.
- 2.  $\mathcal I$  has only one proper subideal, say  $\mathcal J$ . This is 2-dimensional as  $\mathbb F_2$ -vector space. In particular  $\mathcal{J} = \mathcal{I}(1+g^3)$  and  $\mathcal{J}$  is the set of elements of  $\mathcal{I}$  fixed by  $g^3$ , so that  $\mathcal{J}(1+g^3) = 0$ . If we consider  $\mathcal I$  as an  $\mathbb{F}_2\langle g \rangle$ -module, then obviously it is indecomposable and  $\mathcal J = \text{soc}(\mathcal I)$ .
- 3. As an  $\mathbb{F}_2\langle g^2 \rangle$ -module, *I* has exactly 5 irreducible  $\mathbb{F}_2\langle g^2 \rangle$ -submodules (all 2-dimensional), including  $\mathcal J$ . Calling  $\mathcal L_1, \mathcal L_2, \mathcal L_3, \mathcal L_4$  the others, it holds that  $\mathcal L_i(1+g^3)=\mathcal J$ , for all  $i \in \{1,\ldots,4\}$ . Obviously they have pairwise trivial intersection (as they are irreducibles), so that  $\mathcal{I} = \mathcal{J} \oplus \mathcal{L}_i$ , for all  $i \in \{1, ..., 4\}$ .

<span id="page-4-3"></span>**Remark 4.3.** Let  $v \in V_2 = Vf_2$  and consider  $\mathfrak{m} = v\mathbb{F}_2\langle g \rangle$ , the cyclic  $\mathbb{F}_2\langle g \rangle$ -module generated by v. An easy consequence of Remark [4.2](#page-4-1) is that, if  $v \neq 0$ , then only two possibilities occur:

- I.  $\mathfrak{m} \cong \mathcal{J}$ , so that  $\mathfrak{m}$  is an irreducible  $\mathbb{F}_2\langle g \rangle$ -module of dimension 2.
- II.  $\mathfrak{m} \cong \mathcal{I}$ , so that  $\mathfrak{m}$  is an indecomposable  $\mathbb{F}_2\langle g \rangle$ -module of dimension 4.

In the first case m will be called cyclic  $\mathbb{F}_2\langle g \rangle$  -module of type I, while in the second case m will be called cyclic  $\mathbb{F}_2\langle g \rangle$ -module of type II. They inherit by  $\mathcal J$  and  $\mathcal I$  all the properties stated in Remark [4.2.](#page-4-1) Let us underline that  $\mathfrak{m} = v \mathbb{F}_2 \langle g \rangle$  is a cyclic  $\mathbb{F}_2 \langle g \rangle$ -module of type I if and only if v is fixed by  $g^3$ . Since every  $v \in V_2$  belongs to a cyclic  $\mathbb{F}_2\langle g \rangle$ -submodule of  $V_2$  (the one generated by itself), all the irreducible  $\mathbb{F}_2\langle g \rangle$ -submodules of  $\mathcal{V}_2$  are cyclic of type I. For the same reason all the irreducible  $\mathbb{F}_2\langle g^2 \rangle$ -submodules of  $\mathcal{V}_2$  are of dimension 2.

Every  $\mathbb{F}_2\langle g \rangle$ -submodule of  $\mathcal{V}_2$  has even dimension: indeed it is also a  $\mathbb{F}_2\langle g^2 \rangle$ -submodule and so, by Maschke's Theorem, it is the direct sum of irreducible  $\mathbb{F}_2\langle g^2 \rangle$  modules, that have dimension 2.

Now, we give a lemma that relates the socle of  $\mathbb{F}_2\langle g \rangle$ -submodules of  $\mathcal{V}_2$  to elements fixed by  $g^3$ .

<span id="page-5-2"></span>**Lemma 4.3.** Let M an  $\mathbb{F}_2\langle g \rangle$ -submodule of  $\mathcal{V}_2$ . Then

$$
soc(\mathcal{M}) = \mathcal{M}(g^3)
$$

where  $\mathcal{M}(g^3)$  is the set of vectors in M fixed by  $g^3$ .

*Proof.* Since all the irreducible  $\mathbb{F}_2\langle g \rangle$ -submodules of  $\mathcal{V}_2$  are cyclic of type I, then all the elements of an irreducible  $\mathbb{F}_2\langle g \rangle$ -submodule of M are fixed by  $g^3$ . Viceversa, every element of  $\mathcal{M}(g^3)$  belongs necessarily to an irreducible  $\mathbb{F}_2\langle g \rangle$ -module, which is therefore contained in M. Finally, sum of elements fixed by  $g^3$  is again fixed by  $g^3$ . П

Thus  $\text{soc}(\mathcal{E}(g^2)) = (\mathcal{E}(g^2))(g^3)$ . For our purposes it is important the following lemma, which gives a characterization of the socle in terms of the subcodes fixed by  $g^2$  and  $g^3$ .

<span id="page-5-1"></span>Lemma 4.4. With the notations introduced before, it holds

$$
\operatorname{soc}(\mathcal{E}(g^2)) = (\mathcal{E}(g^2))(g^3) = (\mathcal{C}(g^2) + \mathcal{C}(g^3)) \cap \mathcal{V}_2.
$$

Proof. We have just proved the first equality. Now we prove the second one. Obviously  $(\mathcal{E}(g^2))(g^3) = (\mathcal{E}(g^2))(g^3) \cap \mathcal{V}_2 \subseteq \mathcal{C}(g^3) \cap \mathcal{V}_2 \subseteq (\mathcal{C}(g^2) + \mathcal{C}(g^3)) \cap \mathcal{V}_2$ . Viceversa, as we observed in Remark [4.1,](#page-4-2)  $C(g^2) = Cf_1$  is an  $\mathbb{F}_2\langle g \rangle$ -module. Also  $C(g^3)$  is an  $\mathbb{F}_2\langle g \rangle$ module, since g sends words constant on the orbits of  $g^3$  in words constant on the orbits of  $g^3$ . So  $\mathcal{C}(g^2) + \mathcal{C}(g^3)$  is a  $\mathbb{F}_2\langle g \rangle$ -module. Thus, by Lemma [4.2,](#page-4-0)

$$
\mathcal{C}(g^2) + \mathcal{C}(g^3) = (\mathcal{C}(g^2) + \mathcal{C}(g^3))f_1 \oplus (\mathcal{C}(g^2) + \mathcal{C}(g^3))f_2.
$$

Since

$$
\mathcal{C}(g^2) = \mathcal{C}(g^2)f_1 \subseteq (\mathcal{C}(g^2) + \mathcal{C}(g^3))f_1 \subseteq \mathcal{C}f_1 = \mathcal{C}(g^2),
$$

we have that

<span id="page-5-0"></span>
$$
\mathcal{C}(g^2) + \mathcal{C}(g^3) = \mathcal{C}(g^2) \oplus (\mathcal{C}(g^2) + \mathcal{C}(g^3))f_2 \tag{2}
$$

and  $(C(g^2) + C(g^3))f_2 = (C(g^2) + C(g^3)) \cap \mathcal{V}_2 \subseteq (C \cap \mathcal{V}_2) = \mathcal{E}(g^2)$ . If  $v \in (\mathcal{C}(g^2) + \mathcal{C}(g^3))f_2$  then there exist  $v_1 \in \mathcal{C}(g^2)$ ,  $v_2 \in \mathcal{C}(g^3)$  such that  $v = (v_1 + v_2)f_2$ . Obviously  $v_1 f_2 = 0$ , so  $v = v_2 f_2$  is fixed by  $g^3$ , like  $v_2$ .  $\Box$ 

Let us conclude with a observation that will be crucial in the next section.

**Remark 4.4.** We have, by [\(1\)](#page-3-0),  $\dim(\mathcal{C}(g^2) + \mathcal{C}(g^3)) = 12 + 18 - 6 = 24$ . So, by [\(2\)](#page-5-0),  $\dim((\mathcal{C}(g^2) + \mathcal{C}(g^3)) \cap \mathcal{V}_2) = \dim(\mathcal{C}(g^2) + \mathcal{C}(g^3)) - \dim(\mathcal{C}(g^2)) = 24 - 12$ . Thus, by Lemma [4.4,](#page-5-1)

<span id="page-5-3"></span>
$$
\dim(\operatorname{soc}(\mathcal{E}(g^2))) = 12 = \frac{\dim(\mathcal{E}(g^2))}{2}.
$$
\n(3)

# 5 On the structure of the  $\mathbb{F}_2\langle q \rangle$ -submodules of  $\mathcal{V}_2$

In this section we will prove a theorem that is a refinement of the Krull-Schmidt Theorem in a very particular case of  $\mathbb{F}_2\langle g \rangle$ -modules. This result gives us a tool to do an exhaustive search for our code. In this section we strongly use Remark [4.2](#page-4-1) and Remark [4.3.](#page-4-3) Let us state the theorem.

<span id="page-6-2"></span>**Theorem 5.1.** Let M be a  $\mathbb{F}_2\langle g \rangle$  -submodule of  $\mathcal{V}_2$  such that

$$
\dim(\mathcal{M}) = 2 \dim(\text{soc}(\mathcal{M})) = 4m.
$$

Then for every decomposition

$$
\operatorname{soc}({\cal M})={\frak p}_1\oplus\ldots\oplus{\frak p}_m
$$

of the socle in irreducible  $\mathbb{F}_2\langle g \rangle$  -submodules, there exist  $\mathfrak{q}_1,\ldots,\mathfrak{q}_m$ , cyclic  $\mathbb{F}_2\langle g \rangle$  -submodules of type II of M with  $\operatorname{soc}(\mathfrak{q}_i) = \mathfrak{p}_i$  for all  $i \in \{1, \ldots, m\}$ , such that

$$
\mathcal{M}=\mathfrak{q}_1\oplus\ldots\oplus\mathfrak{q}_m.
$$

Before proving the theorem, we need some lemmas.

<span id="page-6-1"></span>**Lemma 5.1.** Let  $\mathfrak{p}$  an irreducible  $\mathbb{F}_2\langle g \rangle$ -submodule of type I of  $\mathcal{V}_2$  and let  $\mathfrak{q}_1, \ldots, \mathfrak{q}_n$  be distinct cyclic  $\mathbb{F}_2\langle g \rangle$  -submodules of type II of  $\mathcal{V}_2$ , all having p as socle, such that

 $\mathfrak{q}_1 + \ldots + \mathfrak{q}_n = \mathfrak{p} \oplus \mathfrak{h}_1 \oplus \ldots \oplus \mathfrak{h}_n$ 

(that is equivalent to ask that  $\dim(\mathfrak{q}_1 + \ldots + \mathfrak{q}_n) = 2 + 2n$ ), where  $\mathfrak{q}_i = \mathfrak{p} \oplus \mathfrak{h}_i$  for all  $i \in \{1, \ldots, n\}$ . Then

$$
\dim(\text{soc}(\mathfrak{q}_1 + \ldots + \mathfrak{q}_n)) = 2n.
$$

Furthermore, every cyclic  $\mathbb{F}_2\langle g \rangle$  -submodule of type II of  $\mathfrak{q}_1 + \ldots + \mathfrak{q}_n$  has socle  $\mathfrak{p}$ .

Proof. Let

$$
m_{1+g^3}: \quad \mathfrak{q}_1 + \ldots + \mathfrak{q}_n \quad \to \quad \mathfrak{q}_1 + \ldots + \mathfrak{q}_n
$$

$$
v \quad \mapsto \quad v(1+g^3)
$$

be the linear map of multiplication by  $1 + g^3$ . Since  $\mathfrak{h}_1(1+g^3) = \ldots = \mathfrak{h}_n(1+g^3) = \mathfrak{p}$  and  $\mathfrak{p}(1+g^3) = 0$ , we have

<span id="page-6-0"></span>
$$
\operatorname{im}(m_{1+g^3}) = \mathfrak{p}.\tag{4}
$$

This implies that  $\ker(m_{1+g^3})$  has dimension  $(2+2n)-2=2n$ . The element  $v \in \mathfrak{q}_1 + \ldots + \mathfrak{q}_n$  is fixed by  $g^3$  if and only if  $v \in \text{ker}(m_{1+g^3})$ , so that, by Lemma [4.3,](#page-5-2)

$$
soc(\mathfrak{q}_1 + \ldots + \mathfrak{q}_n) = \ker(m_{1+g^3}).
$$

Furthermore, as for every q, cyclic  $\mathbb{F}_2\langle g\rangle$  -submodule of type II of  $\mathcal{V}_2$ ,  $\text{soc}(\mathfrak{q}) = m_{1+g^3}(\mathfrak{q})$ , we get the equality  $\operatorname{soc}(\mathfrak{q}) = \mathfrak{p}$  by [\(4\)](#page-6-0).  $\Box$ 

<span id="page-6-3"></span>**Corollary 5.1.** Every  $\mathbb{F}_2\langle g \rangle$  -submodule M of  $\mathcal{V}_2$  with

$$
\dim(\text{soc}(\mathcal{M})) = 2m
$$

has at most  $2^{2m-2}$  cyclic  $\mathbb{F}_2\langle g \rangle$  -submodules of type II with the same socle.

*Proof.* Let us fix an irreducible  $\mathbb{F}_2\langle g^2 \rangle$ -submodule  $\mathfrak{p}$  of soc(M) and let  $\mathcal{K}_{\mathfrak{p}} = {\mathfrak{q}_1, \ldots, \mathfrak{q}_N}$  be the set of all cyclic  $\mathbb{F}_2\langle g \rangle$ -submodules of type II of M that have socle  $\mathfrak{p}$ . If  $N \neq 0$ , let  $\mathcal{N} = \mathfrak{q}_1 + \ldots + \mathfrak{q}_N$ . We have that  $\dim(\mathcal{N}) = 2 + 2n$  for a certain integer  $n \geq 1$ . Obviously there are  $\mathfrak{q}'_1, \ldots, \mathfrak{q}'_n \in \mathcal{K}_{\mathfrak{p}}$ such that

$$
\mathcal{N}=\mathfrak{q}'_1+\ldots+\mathfrak{q}'_n.
$$

By Lemma [5.1,](#page-6-1) dim(soc(N)) = 2n and every cyclic  $\mathbb{F}_2\langle g \rangle$ -submodule of type II of N has socle p (and so it is contained in  $\mathcal{K}_{\mathfrak{p}}$ ). Since every element of  $\mathcal{N} \setminus \text{soc}(\mathcal{N})$  is obviously contained in a cyclic  $\mathbb{F}_2\langle g \rangle$ -submodule of type II of N and  $\mathfrak{q}_i \cap \mathfrak{q}_j = \mathfrak{p}$  for  $i \neq j$ ,  $i, j \in \{1, ..., N\}$ , it is easy to observe that

$$
N = \frac{|\mathcal{N}| - |\text{soc}(\mathcal{N})|}{|\mathfrak{q}| - |\text{soc}(\mathfrak{q})|} = \frac{2^{2n+2} - 2^{2n}}{2^4 - 2^2} = 2^{2n-2}.
$$

Obviously  $\operatorname{soc}(\mathcal{N}) \subseteq \operatorname{soc}(\mathcal{M})$ . Then the maximum for N is reached when  $\dim(\mathcal{N}) = 2 + 2m$ .  $\Box$  <span id="page-7-0"></span>**Corollary 5.2.** Every  $\mathbb{F}_2\langle g \rangle$  -submodule M of  $\mathcal{V}_2$  such that

$$
\dim(\mathcal{M}) = 2 \dim(\text{soc}(\mathcal{M})) = 4m
$$

has exactly  $2^{2m-2}$  cyclic  $\mathbb{F}_2\langle g \rangle$ -submodules of type II with socle p, for every irreducible  $\mathbb{F}_2\langle g \rangle$ submodule  $\mathfrak{p} \subset \text{soc}(\mathcal{M})$ .

*Proof.* The number of all the irreducible  $\mathbb{F}_2\langle g \rangle$  -submodules of M is equal to

$$
N_{\text{irr}} = \frac{|\text{soc}(\mathcal{M})| - 1}{2^2 - 1} = \frac{2^{2m} - 1}{2^2 - 1},
$$

since obviously every element of the socle belongs to an irreducible  $\mathbb{F}_2\langle g \rangle$ -module and the irreducibles have pairwise trivial intersection. With similar arguments it is easy to prove that the number of all the cyclic  $\mathbb{F}_2\langle g \rangle$  -submodules of type II of  $\mathcal M$  is equal to

$$
N_{\rm ind} = \frac{|\mathcal{M}| - |\text{soc}(\mathcal{M})|}{2^4 - 2^2} = \frac{2^{4m} - 2^{2m}}{2^4 - 2^2}.
$$

Then the average of indecomposables for each irreducible is

$$
\frac{N_{\text{ind}}}{N_{\text{irr}}} = 2^{2m-2},
$$

 $\Box$ that is also the maximum. So the maximum is reached for every irreducible  $\mathbb{F}_2\langle g \rangle$ -submodule.

<span id="page-7-1"></span>**Lemma 5.2.** Let  $q_1, \ldots, q_n$  be cyclic  $\mathbb{F}_2\langle g \rangle$ -submodules of type II of  $\mathcal{V}_2$  such that  $\mathfrak{p}_1 + \ldots + \mathfrak{p}_n$  is a direct sum (i.e.  $\dim(\mathfrak{p}_1 \oplus \ldots \oplus \mathfrak{p}_n) = 2n$ ), where  $\mathfrak{p}_i = \text{soc}(\mathfrak{q}_i)$  for every  $i \in \{1, \ldots, n\}$ . Then

1.  $\mathfrak{q}_1 + \ldots + \mathfrak{q}_n$  is a direct sum (i.e.  $\dim(\mathfrak{q}_1 \oplus \ldots \oplus \mathfrak{q}_n) = 4n$ ),

$$
2. \ \operatorname{soc}(\mathfrak{q}_1 + \ldots + \mathfrak{q}_n) = \mathfrak{p}_1 \oplus \ldots \oplus \mathfrak{p}_n.
$$

*Proof.* For every  $i \in \{1, ..., n\}$ , let  $\mathfrak{h}_i$  irreducible  $\mathbb{F}_2\langle g^2 \rangle$ -submodule such that  $\mathfrak{q}_i = \mathfrak{p}_i \oplus \mathfrak{h}_i$ . We make induction on  $n \geq 2$ .

Let  $n = 2$ . Obviously  $\mathfrak{q}_1 \cap \mathfrak{q}_2 = \{0\}$ . So  $\mathfrak{q}_1 + \mathfrak{q}_2 = \mathfrak{p}_1 \oplus \mathfrak{h}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{h}_2$ . Arguing as in Lemma [5.1,](#page-6-1) we can consider the linear map  $m_{1+g^3}$  and we can prove  $\text{soc}(\mathfrak{p}_1 \oplus \mathfrak{h}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{h}_2) = \text{ker}(m_{1+g^3})$  has dimension 4. Since  $\mathfrak{p}_1 \oplus \mathfrak{p}_2 \subseteq \text{soc}(\mathfrak{p}_1 \oplus \mathfrak{h}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{h}_2)$ , the equality holds.

Let us suppose 1. and 2. true for  $n-1$ . Then  $\mathfrak{q}_n \cap (\mathfrak{q}_1 \oplus \ldots \oplus \mathfrak{q}_{n-1}) = \{0\}$ , since  $\operatorname{soc}(\mathfrak{q}_n) = \mathfrak{p}_n$  has trivial intersection with  $\operatorname{soc}(\mathfrak{q}_1 \oplus \ldots \oplus \mathfrak{q}_{n-1})$ . Thus 1. is true for n. 2. can be proved as in the basis of the induction.  $\Box$ 

We have now all the ingredients to prove the theorem.

*Proof of Theorem* [5.1.](#page-6-2) Corollary [5.2](#page-7-0) implies that the set  $\mathcal{K}_{\mathfrak{p}_i}$  of cyclic  $\mathbb{F}_2\langle g \rangle$ -submodules of type II of M with socle  $\mathfrak{p}_i$  is non-empty for all  $i \in \{1, ..., m\}$ . Choose  $\mathfrak{q}_i \in \mathcal{K}_{\mathfrak{p}_i}$  for every  $i \in \{1, ..., m\}$ . Then, for Lemma [5.2,](#page-7-1)  $q_1 + \ldots + q_m$  ( $\subseteq \mathcal{M}$ ) has dimension  $4n$ . So the equality holds.

**Remark 5.1.** For the following we point out that every choice of  $\mathfrak{q}_i \in \mathcal{K}_{\mathfrak{p}_i}$   $(i \in \{1, ..., m\})$  is allowed.

Coming back to our problem, we observe that  $\mathcal{E}(g^2)$  satisfies, by [\(3\)](#page-5-3), the hypothesis of the theorem:

$$
24 = \dim(\mathcal{E}(g^2)) = 2 \cdot \dim(\text{soc}(\mathcal{E}(g^2))) = 4 \cdot 6.
$$

In section [6](#page-8-0) we will determine the possible socles of  $\mathcal{E}(g^2)$  and in section [7,](#page-11-0) using Theorem [5.1](#page-6-2) and the results of section [6,](#page-8-0) we will describe the exhaustive search.

# <span id="page-8-0"></span>6 Restrictions on  $\operatorname{soc}(\mathcal{E}(g^2))$

By Lemma [4.4](#page-5-1) we know that  $\operatorname{soc}(\mathcal{E}(g^2)) = (\mathcal{C}(g^2) + \mathcal{C}(g^3)) \cap \mathcal{V}_2$ . This suggests us to determine which code can be  $\mathcal{C}(g^2) + \mathcal{C}(g^3)$  in order to get soc $(\mathcal{E}(g^2))$ .

We have seen that we can suppose  $g = (1, 2, 3, 4, 5, 6) \dots (67, 68, 69, 70, 71, 72)$ , so that

$$
g^2 = (1, 3, 5)(2, 4, 6) \dots (67, 69, 71)(68, 70, 72)
$$
 and  $g^3 = (1, 4)(2, 5)(3, 6) \dots (67, 70)(68, 71)(69, 72).$ 

Let us recall the results and the notations of Proposition [3.1,](#page-2-0) [3.2](#page-2-1) and [3.3:](#page-2-2)  $\mathcal{C}(g^2)$  is equivalent to  $\mathcal{G}_{24} \otimes \langle (1,1,1) \rangle$ , with  $\mathcal{G}_{24}$  extended binary Golay code;  $\mathcal{C}(g^3)$  equivalent to  $\mathcal{K} \otimes \langle (1,1) \rangle$ , with  $\mathcal{K}$  one of the 41 self-dual [36, 18, 8] codes classified by Mechor and Gaborit;  $\mathcal{C}(g) = \mathcal{H} \otimes \langle (1, 1, 1, 1, 1, 1), \rangle$ with H equivalent to F, self-dual  $[12, 6, 4]$  code with generator matrix M.

Define

$$
\pi_{24} : \mathcal{C}(g^2) \subset \mathbb{F}_2^{72} \rightarrow \mathbb{F}_2^{24}
$$
  

$$
(c_1, c_2, c_1, c_2, c_1, c_2, \dots, c_{23}, c_{24}, c_{23}, c_{24}, c_{23}, c_{24}) \rightarrow (c_1, c_2, \dots, c_{23}, c_{24})
$$

and

$$
\pi_{36} : \mathcal{C}(g^3) \subset \mathbb{F}_2^{72} \rightarrow \mathbb{F}_2^{36}
$$
  

$$
(c_1, c_2, c_3, c_1, c_2, c_3, \dots, c_{34}, c_{35}, c_{36}, c_{34}, c_{35}, c_{36}) \rightarrow (c_1, c_2, c_3 \dots, c_{34}, c_{35}, c_{36})
$$

Let us call

$$
\overline{g}_{24} = (1, 2) \dots (23, 24)
$$
 and  $\overline{g}_{36} = (1, 2, 3) \dots (34, 35, 36).$ 

It is easy to notice that

$$
(\pi_{24}(c))^{\overline{g}_{24}} = \pi_{24}(c^g)
$$
 and  $(\pi_{36}(c))^{\overline{g}_{36}} = \pi_{36}(c^g)$ 

for c in  $C(g^2)$  and c in  $C(g^3)$  respectively. So

$$
\overline{g}_{24} \in \text{Aut}(\pi_{24}(\mathcal{C}(g^2))) \quad \text{and} \quad \overline{g}_{36} \in \text{Aut}(\pi_{36}(\mathcal{C}(g^3)))
$$

This observation implies that  $\pi_{36}(\mathcal{C}(g^3))$  has an automorphism of order 3 and degree 36. Only 13 out of the 41 codes classified by Mechor and Gaborit have such an automorphism.

It is easy to see that

$$
\pi_{24}(\mathcal{C}(g)) = (\pi_{24}(\mathcal{C}(g^2)))(\overline{g}_{24})
$$
 and  $\pi_{36}(\mathcal{C}(g)) = (\pi_{36}(\mathcal{C}(g^3)))(\overline{g}_{36}).$ 

Another important observation is that

$$
\pi_{24}(\mathcal{C}(g)) = \mathcal{H} \otimes \langle (1,1) \rangle \quad \text{and} \quad \pi_{36}(\mathcal{C}(g)) = \mathcal{H} \otimes \langle (1,1,1) \rangle
$$

These remarks together with the lemmas below allow us to deduce the following result.

<span id="page-8-1"></span>**Theorem 6.1.**  $C(g^2) + C(g^3)$  belongs, up to equivalence, to a set  $\mathcal L$  of cardinality 38. Every element of  $\mathcal L$  is a binary self-orthogonal doubly-even [72, 24, 16] code.

We describe the algorithm which proves the theorem.

**Step 1.** Choose a particular extended binary Golay code, say  $\mathcal{G}$ , and find all its subcodes equivalent to  $\mathcal{F} \otimes \langle (1, 1) \rangle$ , say  $\mathcal{M}_1, \ldots, \mathcal{M}_m$ .

For each  $\mathcal{M}_i$  denote with  $h_i$  an element of  $\mathcal{S}_{24}$  such that  $\mathcal{M}_i{}^{h_i} = \mathcal{F} \otimes \langle (1,1) \rangle$  and set  $\mathcal{G}_i = \mathcal{G}^{h_i}$ . Denote  $AF = \text{Aut}(\mathcal{F} \otimes \langle (1,1) \rangle)$  and  $\mathcal{AG} = \mathcal{G}_1^{AF}$ , i.e. the orbit of  $\mathcal{G}_1$  under the action of  $AF$ . Fact: we have  $\mathcal{G}_i^{AF} = \mathcal{AG}$  for all  $i \in \{1, ..., m\}$  (since, by direct calculations,  $\mathcal{G}_i \in \mathcal{AG}$ ).

**Lemma 6.1.** AG is the set of all extended binary Golay codes that has  $\mathcal{F} \otimes \langle (1, 1) \rangle$  as subcode.

*Proof.* Take an extended binary Golay code  $\mathcal J$  with  $\mathcal F \otimes \langle (1, 1) \rangle$  as subcode. Then there exists  $h \in \mathcal S_{24}$ such that  $\mathcal{J}^h = \mathcal{G}$ . There exists  $j \in \{1, ..., m\}$  such that  $(\mathcal{F} \otimes \langle (1, 1) \rangle)^h = \mathcal{M}_j$ . Then  $((\mathcal{F} \otimes \langle (1,1) \rangle)^{h})^{h_j} = \mathcal{M}_j^{h_j} = \mathcal{F} \otimes \langle (1,1) \rangle$  and so  $hh_j \in AF$ . But  $\mathcal{J}^{hh_j} = \mathcal{G}_j$  and so  $\mathcal{J} \in \mathcal{G}^{AF}_j = \ \mathcal{AG}$  . П

Step 2. Take the 13 codes of the classification of Mechor and Gaborit which have automorphisms of order 3 and degree 36, say  $\mathcal{D}_1, \ldots, \mathcal{D}_{13}$ . Denote

$$
\{e_{1,1}, \ldots, e_{1,n_1}\} \subset \text{Aut}(\mathcal{D}_1),
$$
  

$$
\vdots
$$
  

$$
\{e_{13,1}, \ldots, e_{13,n_{13}}\} \subset \text{Aut}(\mathcal{D}_{13})
$$

the sets of automorphisms of order 3 and degree 36 of  $\mathcal{D}_1, \ldots, \mathcal{D}_{13}$  respectively.

Find  $h_{i,j} \in \mathcal{S}_{36}$  such that  $h_{i,j}^{-1}e_{i,j}h_{i,j} = \overline{g}_{36}$  and set  $\mathcal{D}_{i,j} = \mathcal{D}_i^{h_{i,j}}$  (so that  $\overline{g}_{36}$  is an automorphism of  $\mathcal{D}_i^{h_{i,j}}$  ).

Find all the subcodes of  $\mathcal{D}_{i,j}$  equivalent to  $\mathcal{F} \otimes \langle (1,1,1) \rangle$  and which are fixed word by word by  $\overline{g}_{36}$ , say  $\mathcal{N}_{(i,j)_1}, \ldots, \mathcal{N}_{(i,j)_{n_{i,j}}}$ .

Denote with  $l_{(i,j)_k}$  an element of  $S_{36}$  such that  $\mathcal{N}_{(i,j)_k}l_{(i,j)_k} = \mathcal{F} \otimes \langle (1,1,1) \rangle$  and that commutes with  $\overline{g}_{36}$ . Let us show its existence: every element  $l \in S_{36}$  for which  $\mathcal{N}_{(i,j)_k}^{\phantom{1}}^{\phantom{1}} = \mathcal{F} \otimes \langle (1,1,1) \rangle$  is such that  $l^{-1}\overline{g}_{36}l$  has order 3 and fixes every word of  $\mathcal{F}\otimes\langle (1,1,1)\rangle$ ; thus, by direct calculations, it has the same orbits (on the coordinates, as an element of  $S_{36}$ ) of  $\overline{g}_{36}$ . So  $l^{-1}\overline{g}_{36}l = \overline{g}_{36}$  or  $l^{-1}\overline{g}_{36}l = \overline{g}_{36}^{-1}$ . In the first case take  $l_{(i,j)_k} = l$ , in the second case take  $l_{(i,j)_k} = ls$  where s is an automorphism of  $\mathcal{F} \otimes \langle (1, 1, 1) \rangle$  of order 2 (a relabelling of the coordinates of each orbit of  $\overline{g}_{36}$ ) such that  $s^{-1}\overline{g}_{36}^{-1}s = \overline{g}_{36}$ .

Set  $\mathcal{D}_{(i,j)_k} = \mathcal{D}_{i,j}^{l_{(i,j)_k}}$  and denote with C36 the set of all the codes  $\mathcal{D}_{(i,j)_k}$ .

C36 is a proper subset of all the codes equivalent to  $\mathcal{D}_1, \ldots, \mathcal{D}_{13}$  which contain  $\mathcal{F} \otimes \langle (1, 1, 1) \rangle$ . The following lemma shows that C36 is big enough to allow us to determine all the possible  $C(g^2) + C(g^3)$ .

**Lemma 6.2.** There exist  $\mathcal{B}_3 \in \mathcal{AG}$  and  $\mathcal{B}_2 \in \mathcal{C}36$  such that  $\mathcal{C}(g^2) + \mathcal{C}(g^3)$  is equivalent to

$$
{\pi_{24}}^{-1}(\mathcal{B}_3) + {\pi_{36}}^{-1}(\mathcal{B}_2).
$$

Proof. Up to equivalence, we can suppose

$$
\mathcal{C}(g^2) \cap \mathcal{C}(g^3) = \mathcal{F} \otimes \langle (1,1,1,1,1,1) \rangle.
$$

There exists  $i \in \{1, \ldots, 13\}$  and  $h \in S_{36}$  such that  $\pi_{36}(\mathcal{C}(g^3))^h = \mathcal{D}_i$ . There exists  $j \in \{i, \ldots, n_i\}$  such that  $h^{-1}\overline{g}_{36}h = e_{i,j}$ . So

$$
\pi_{36}(\mathcal{C}(g^3))^{hh_{i,j}} = \mathcal{D}_{i,j} \quad \text{and} \quad \overline{g}_{36} = h_{i,j}^{-1}e_{i,j}h_{i,j} = h_{i,j}^{-1}h^{-1}\overline{g}_{36}hh_{i,j}.
$$

There exists  $k \in \{1, \ldots, n_{i,j}\}\$  such that  $(\mathcal{F} \otimes \langle (1,1,1) \rangle)^{hh_{i,j}} = \mathcal{N}_{(i,j)_k}$ . So

$$
\pi_{36}(\mathcal{C}(g^3))^{hh_{i,j}l_{(i,j)_k}} = \mathcal{D}_{(i,j)_k} \quad \text{and} \quad \overline{g}_{36} = l_{(i,j)_k}^{-1} \overline{g}_{36}l_{(i,j)_k}.
$$

Thus, if we set  $\bar{t} = hh_{i,j}l_{(i,j)_k}$ , we have

1.  $\pi_{36}(\mathcal{C}(g^3))^{\overline{t}} = \mathcal{D}_{(i,j)_k};$ 2.  $\overline{g}_{36} = \overline{t}^{-1} \overline{g}_{36} \overline{t}$ ; 3.  $\overline{t} \in \text{Aut}(\mathcal{F} \otimes \langle (1, 1, 1) \rangle)$ .

It is now possible to construct an element  $t \in S_{72}$  such that

1.  $\pi_{36}(c^t) = (\pi_{36}(c))^{\overline{t}}$  for all  $c \in C(g^3)$ ; 2.  $g = t^{-1}gt;$ 3.  $t \in \text{Aut}(\mathcal{F} \otimes \langle (1, 1, 1, 1, 1, 1)\rangle)$ .

The construction of t will be done in Remark [6.1.](#page-10-0) Condition 1. implies that  $\pi_{36}(\mathcal{C}(g^3)^t) \in \mathcal{C}36$ . Condition 2. implies that  $\mathcal{C}(g^2)^t$  is equal to  $\mathcal{C}^t(g^2)$ . Actually, if  $c \in \mathcal{C}(g^2)$ , then  $(c^t)^{g^2} = (c^t)^{t^{-1}g^2t} =$  $c^t$ , so every word of  $C(g^2)^t$  is fixed by  $g^2$ . So  $C(g^2)^t \subseteq C^t(g^2)$ .  $C(g^2)^t$  has obviously dimension 12.  $\mathcal{C}^t$  is a binary self-dual doubly-even [72, 36, 16] code with g as automorphism, so  $\mathcal{C}^t(g^2)$  has dimension 12 too, and thus it holds the equality. This implies that  $\pi_{24}(\mathcal{C}(g^2)^t)$  is an extended binary Golay code. Condition 3. implies that  $\pi_{24}(\mathcal{C}(g^2)^t)$  has  $\mathcal{F} \otimes \langle (1,1) \rangle$  as subcode. Indeed,

$$
\mathcal{F}\otimes \langle (1,1)\rangle=\pi_{24}(\mathcal{F}\otimes \langle (1,1,1,1,1,1))\rangle)=\pi_{24}((\mathcal{F}\otimes \langle (1,1,1,1,1,1))\rangle)^t)).
$$

Thus  $\pi_{24}(\mathcal{C}(g^2)^t) \in \mathcal{AG}$ .

<span id="page-10-0"></span>**Remark 6.1.** It is easy to convince themselves that  $t$  exists. For reader's convenience we give an explicit construction of t through wreath product.

Let  $\Delta = \{1, 2\}$  and  $\Gamma = \{1, 2, 3\}$ . We have  $S_{\Delta} = S_2$  and  $S_{\Gamma} = S_3$ . We want to explain how the wreath product  $S_{\Delta} \wr S_{\Gamma}$  acts on the coordinates of  $\mathbb{F}_2^6$ .

Firstly, we can see

$$
\Delta \times \Gamma = \Delta_1 \cup \Delta_2 \cup \Delta_3
$$

with  $\Delta_1 = \{1, 4\}$ ,  $\Delta_2 = \{2, 5\}$  and  $\Delta_3 = \{3, 6\}$ . This can be send in the ordered set  $\Omega =$  $\{1, 2, 3, 4, 5, 6\}$  in a natural way, that is sending the first element of  $\Delta_1$  in the first element of  $\Omega$ , the second element of  $\Delta_1$  in the fourth element of  $\Omega$ , the first element of  $\Delta_2$  in the second element of  $\Omega$  and so on, i.e. by sending i in i, in the ordered set  $\Omega$ . We denote this map

$$
\varphi : \Delta_1 \cup \Delta_2 \cup \Delta_3 \to \Omega.
$$

An element h of  $S_{\Delta} \wr S_{\Gamma}$  has the shape

$$
h=(\delta_1,\delta_2,\delta_3,\gamma)\in\mathcal{S}_{\Delta}\times\mathcal{S}_{\Delta}\times\mathcal{S}_{\Delta}\times\mathcal{S}_{\Gamma}
$$

and acts on  $\Delta_1 \cup \Delta_2 \cup \Delta_3$  in the following way:

$$
(\Delta_1 \cup \Delta_2 \cup \Delta_3)^h = (\Delta_{\gamma^{-1}(1)})^{\delta_{\gamma^{-1}(1)}} \cup (\Delta_{\gamma^{-1}(2)})^{\delta_{\gamma^{-1}(2)}} \cup (\Delta_{\gamma^{-1}(3)})^{\delta_{\gamma^{-1}(3)}}.
$$

With this notation it is possible to check that, for example,

$$
(1, 2, 3, 4, 5, 6) = \varphi^{-1}(\text{Id}, \text{Id}, (1, 2), (1, 2, 3))\varphi.
$$

In a similar way, we have that

$$
\mathcal{S}_2 \wr \mathcal{S}_{36} = \mathcal{S}_{\Delta} \wr \mathcal{S}_{\Gamma_{36}},
$$

where  $\Gamma_{36} = \{1, \ldots, 36\}$ , acts on the coordinates of  $\mathbb{F}_2^{72}$ , thanks to a suitable

$$
\varphi_{36} : \underbrace{\{1,4\} \cup \{2,5\} }_{\Delta_1} \cup \underbrace{\{3,6\} \cup \ldots \cup \{67,70\} }_{\Delta_3} \cup \underbrace{\{68,71\} }_{\Delta_{34}} \cup \underbrace{\{69,72\} }_{\Delta_{35}} \rightarrow \Omega_{72}
$$

where  $\Omega_{72} = \{1, \ldots, 72\}$ . With this notation we have that

$$
g = \varphi_{36}^{-1}(\text{Id}, \text{Id}, (1, 2), \dots, \text{Id}, \text{Id}, (1, 2), \overline{g}_{36})\varphi_{36}.
$$

Now, the  $t \in S_{72}$  that we were looking for is

$$
t = \varphi_{36}^{-1}(\text{Id}, \text{Id}, (1, 2), \dots, \text{Id}, \text{Id}, (1, 2), \overline{t})\varphi_{36}.
$$

This t has all the desired properties (it is checkable by hand).

 $\Box$ 

**Step 3.** Construct the set of all  $\pi_{24}^{-1}(\mathcal{B}_3) + \pi_{36}^{-1}(\mathcal{B}_2)$ , with  $\mathcal{B}_3 \in \mathcal{AG}$  and  $\mathcal{B}_2 \in \mathcal{C}36$  and take one representant for each equivalence class of this set. Collect them in the set  $\mathcal{L}$ , that satisfies, obviously, the requested properties.

<span id="page-11-2"></span>**Remark 6.2.** We have proved that, if C does exist, there are  $L \in \mathcal{L}$  and  $r \in \mathcal{S}_{72}$  such that

$$
(\mathcal{C}(g^2) + \mathcal{C}(g^3))^r = L.
$$

The problem is that it is not clear, a priori, whether  $r^{-1}gr = g$  or not. This is very important, since  $V_2 = V(g^2 + g^4)$  depends strongly on g.

The element  $r^{-1}gr$  is an automorphism of L of order 6 and it has the same cycle structure of g. Furthermore  $L = L(r^{-1}g^2r) + L(r^{-1}g^3r)$ . There are not many elements with these features in Aut(L),  $L \in \mathcal{L}$ . Using this fact we construct a new list  $\mathcal{L}'$  (of 40 elements) with the following property: there exist  $L \in \mathcal{L}'$  and  $s \in \mathcal{S}_{72}$  such that

$$
(\mathcal{C}(g^2) + \mathcal{C}(g^3))^s = L
$$
 and  $s^{-1}gs = g$ .

<span id="page-11-0"></span>This guarantees that  $\operatorname{soc}(\mathcal{E}(g^2)))^r = L \cap \mathcal{V}_2$ .

# 7 Construction of a putative  $\mathcal{E}(g^2)$  from its socle

Now let us conclude the construction. Following the track laid out in Theorem [5.1,](#page-6-2) we define, for every **p**, irreducible  $\mathbb{F}_2\langle g \rangle$  -submodule of  $\operatorname{soc}(\mathcal{E}(g^2))$ , the set

 $\mathcal{H}_{\mathfrak{p}} = \{\mathfrak{h} \mid \mathfrak{h} \text{ irreducible } \mathbb{F}_2\langle g^2 \rangle\text{-submodules of } \mathcal{V}_2 \text{ such that } \mathfrak{h}(1 + g^3) = \mathfrak{p}\}.$ 

We will now take a decomposition of  $\operatorname{soc}(\mathcal{E}(g^2))$  into irreducible  $\mathbb{F}_2\langle g \rangle$ -submodules

$$
\text{soc}(\mathcal{E}(g^2))=\mathfrak{p}_1\oplus\mathfrak{p}_2\oplus\mathfrak{p}_3\oplus\mathfrak{p}_4\oplus\mathfrak{p}_5\oplus\mathfrak{p}_6
$$

and add 6 irreducibles  $\mathfrak{h}_1 \in \mathcal{H}_{\mathfrak{p}_1}, \mathfrak{h}_2 \in \mathcal{H}_{\mathfrak{p}_2}, \mathfrak{h}_3 \in \mathcal{H}_{\mathfrak{p}_3}, \mathfrak{h}_4 \in \mathcal{H}_{\mathfrak{p}_4}, \mathfrak{h}_5 \in \mathcal{H}_{\mathfrak{p}_5}, \mathfrak{h}_6 \in \mathcal{H}_{\mathfrak{p}_6}$ , to this decomposition, in all the possible combinations. This is obviously equivalent to take the direct sum of  $\mathfrak{q}_1,\ldots,\mathfrak{q}_6$ , cyclic  $\mathbb{F}_2\langle g\rangle$ -modules of type II such that  $\operatorname{soc}(\mathfrak{q}_i) = \mathfrak{p}_i$  for all  $i \in \{1,\ldots,6\}$ , in all the possible combinations.

Now we do some considerations important for the computational part. Firstly we determine the cardinalities of  $\mathcal{H}_{\mathfrak{p}}$ .

<span id="page-11-1"></span>**Lemma 7.1.** For every irreducible  $\mathbb{F}_2\langle g \rangle$  -submodule  $\mathfrak{p}$  we have  $|\mathcal{H}_{\mathfrak{p}}| = 4^{12}$ .

*Proof.* We give a constructive proof. As we have already observed in section [4,](#page-3-1) every irreducible  $\mathbb{F}_2\langle g \rangle$  submodule  $\mathfrak p$  of  $\mathcal V_2$  is cyclic and fixed by  $g^3$ . Furthermore it is also a cyclic  $\mathbb F_2\langle g^2\rangle$ -submodule. Say

$$
\mathfrak{p} = \{0, v, v^g, v^{g^2}\} \quad \text{with } v \in \mathcal{V}_2(g^3).
$$

Consider the sets  $\Omega_i = \{6i-5, 6i-4, 6i-3, 6i-2, 6i-1, 6i\}$ , which we call blocks, for  $i \in \{1, ..., 12\}$ . It is easy to observe that  $v_{\vert_{\Omega_i}}$  can be only one of the following



since v is of even weight on the orbits of  $g^2$  and fixed by  $g^3$ . Every  $\mathfrak{h} \in \mathcal{H}_{\mathfrak{p}}$  is cyclic. Thus we can choose its generator z so that  $z(1 + g^3) = v$ . Since  $z_{|_{\Omega_i}}(1+g^3)=v_{|_{\Omega_i}}$  , the following possibilities for  $\ z_{|_{\Omega_i}}$  occur:

 $z_{\vert_{\Omega_i}} \in \{[0,0,0,0,0,0], [1,1,0,1,1,0], [0,1,1,0,1,1], [1,0,1,1,0,1]\}$  if  $v_{\vert_{\Omega_i}}$  is **A.**,

$$
\begin{aligned} & \boldsymbol{z}_{|\boldsymbol{\alpha}_i} \in \{[1,0,0,0,1,0], [0,1,0,1,0,0], [1,1,1,0,0,1], [0,0,1,1,1,1]\} \text{ if } v_{|\boldsymbol{\alpha}_i} \text{ is } \mathbf{B}, \\ & \boldsymbol{z}_{|\boldsymbol{\alpha}_i} \in \{[0,1,0,0,0,1], [0,0,1,0,1,0], [1,1,1,1,0,0], [1,0,0,1,1,1]\} \text{ if } v_{|\boldsymbol{\alpha}_i} \text{ is } \mathbf{C}, \\ & \boldsymbol{z}_{|\boldsymbol{\alpha}_i} \in \{[1,0,1,0,0,0], [0,0,0,1,0,1], [0,1,1,1,1,0], [1,1,0,0,1,1]\} \text{ if } v_{|\boldsymbol{\alpha}_i} \text{ is } \mathbf{D}. \end{aligned}
$$

We have 4 choices for every block and the blocks are 12. So  $|\mathcal{H}_{\mathfrak{p}}| = 4^{12}$ .

So, apparently, we have  $(4^{12})^6$  calculations to do, a number that would make unfeasible our search. The point is that two modules that "make  $\operatorname{soc}(\mathcal{E}(g^2))$  grow in the same way" are equal from our point of view. More precisely, we are interested only in the representatives of the equivalence classes of the equivalence relation over  $\mathcal{H}_{\mathfrak{p}}$  defined as following:

$$
\mathfrak{h}_1 \sim \mathfrak{h}_2
$$
 if and only if 
$$
\operatorname{soc}(\mathcal{E}(g^2)) + \mathfrak{h}_1 = \operatorname{soc}(\mathcal{E}(g^2)) + \mathfrak{h}_2
$$

Lemma 7.2. Each equivalence class is composed by 4096 elements.

*Proof.* Let us fix  $\mathfrak{h} \in \mathcal{H}_{\mathfrak{p}}$ , for  $\mathfrak{p} \subseteq \text{soc}(\mathcal{E}(g^2))$ . With arguments similar to the ones used in Lemma [5.1,](#page-6-1) Corollary [5.1](#page-6-3) and Corollary [5.2,](#page-7-0) we can prove that all the indecomposable  $\mathbb{F}_2\langle g \rangle$ -modules in soc( $\mathcal{E}(g^2)$ ) + h have socle p and they are  $2^{2\cdot 6-2} = 1024$ . Since every indecomposable  $\mathbb{F}_2\langle g \rangle$ -module, as we have observed in Section [4,](#page-3-1) contains 4 elements of  $\mathcal{H}_{p}$ , and the indecomposable  $\mathbb{F}_{2}(g)$ -modules have pairwise intersection  $\mathfrak p$ , then there are exactly  $4 \cdot 1024 = 4096$  elements of  $\mathcal{H}_{\mathfrak p}$  in  $\operatorname{soc}(\mathcal{E}(g^2)) + \mathfrak h$ . The thesis follows easily. П

Thus the number of classes is  $4^{12}/4096 = 4096$ , a more practical number to do calculations. Unfortunately  $4096^6$  is still too big. However, we are interested only in the representatives that give us a doubly-even module. It is easy to prove (thanks to the construction in Lemma [7.1\)](#page-11-1) that exactly the half of the elements of  $\mathcal{H}_{\mathfrak{p}}$  are doubly-even. Moreover, if  $\operatorname{soc}(\mathcal{E}(g^2)) + \mathfrak{h}$  is doubly-even then all the elements of the class of h are doubly-even. It happens that the number of doubly-even representatives is at most  $4^{11}/4096 = 2048$ .

We are to explain our algorithm (we will use the notation  $d(\mathcal{C})$  to indicate the minimum distance of a code  $\mathcal{C}$  ):

**Step 1.** Take the set  $\mathcal{L}' = \{L_1, \ldots, L_{40}\}\$ and set  $F_i = L_i \cap \mathcal{V}_2$ .

Do the following steps for every  $i \in \{1, \ldots, 40\}$ .

**Step 2.** Find 6 irreducible  $\mathbb{F}_2\langle g \rangle$ -modules, say

$$
P_{i,1},\ldots,P_{i,6}
$$

such that  $F_i = P_{i,1} \oplus \ldots \oplus P_{i,6}$ .

**Step 3.** For every  $P_{i,j}$ ,  $j \in \{1, ..., 6\}$ , find a set of representatives of the 2048 equivalence classes of  $\mathcal{H}_{P_{i,j}}$  consisting of doubly-even modules. Then find the subset of  $\mathcal{H}_{i,j}$  defined in the following way

$$
\mathcal{H}'_{i,j} = \{ H \in \mathcal{H}_{i,j} \mid d(H + L_i) \ge 16 \}
$$
  
=  $\{ H_{(i,j)_1}, \dots, H_{(i,j)_{n_{i,j}}} \}$ 

**Step 4.** Find the subset of  $\mathcal{H}'_{i,1} \times \mathcal{H}'_{i,2}$  defined in the following way

$$
\mathcal{P}_i = \{ (H_1, H_2) \mid d(L_i + H_1 + H_2) \ge 16 \}.
$$

Find the subset of  $\mathcal{P}_i \times \mathcal{H}'_{i,3}$  defined in the following way

$$
\mathcal{T}_i = \{ (P, H) \mid d(L_i + P_1 + P_2 + H) \ge 16 \}.
$$

 $\Box$ 

Find the subset of  $\mathcal{T}_i \times \mathcal{H}'_{i,4}$  defined in the following way

$$
Q_i = \{(T, H) \mid d(L_i + T_1 + T_2 + T_3 + H) \ge 16\}.
$$

Find the subset of  $\mathcal{Q}_i \times \mathcal{H}'_{i,5}$  defined in the following way

$$
\mathcal{F}_i = \{ (Q, H) \mid d(L_i + Q_1 + \ldots + Q_4 + H) \ge 16 \}.
$$

Find the subset of  $\mathcal{F}_i \times \mathcal{H}'_{i,6}$  defined in the following way

$$
S_i = \{ (F, H) \mid d(L_i + F_1 + \ldots + F_5 + H) \ge 16 \}.
$$

#### 8 Conclusions

Theorems [6.1](#page-8-1) tells us that, if a binary self-dual doubly-even [72, 36, 16] code with automorphism of order 6 exists, then it has a subcode equivalent to one of the 38 codes in  $\mathcal{L}$ . Remark [6.2](#page-11-2) and Theorem [5.1](#page-6-2) imply that the eventual code can be found in the sets

$$
\{L_i + S_1 + \ldots + S_6 \mid S \in S_i\}_{i \in \{1, \ldots, 40\}}
$$

where  $\mathcal{L}' = \{L_1, ..., L_{40}\}.$ 

MAGMA calculations find  $S_i$  empty, for all  $i \in \{1, \ldots, 40\}$ . So a binary self-dual doubly-even [72, 36, 16] code with automorphism of order 6 does not exist.

#### Acknowledgements

The author would like to thank his advisors, Prof. Francesca Dalla Volta and Prof. Massimiliano Sala, for the insightful suggestion of the problem and for the long and fruitful discussions. Laboratorio di Matematica Industriale e Crittografia (CryptoLab) of Università degli Studi di Trento deserves thanks for the great help in the computational part.

#### <span id="page-13-2"></span>References

- <span id="page-13-1"></span>[1] E. F. Assmuss, H.F. Mattson, New 5 -designs, J. Combin. Theory 6 (1969) 122–151.
- [2] E.F. Assmus, H.F. Mattson, R.J. Turyn, Research to develop the algebraic theory of codes, Air Force Cambridge Res. Labs., Belford, MA, Report AFCRL-67-0365 (1967)
- <span id="page-13-0"></span>[3] W. Bosma, J. Cannon, C. Playoust, The MAGMA algebra system I: The user language, J. Symbol. Comput. 24 (1997) 235–265.
- <span id="page-13-5"></span>[4] S. Bouyuklieva, A method for constructing self-dual codes with an automorphism of order 2, IEEE Trans. Inform. Theory 46 (2000) 496–504.
- <span id="page-13-4"></span>[5] S. Bouyuklieva, On the automorphism group of a doubly even (72, 36, 16) code, IEEE Trans. Inform. Theory 50 (2004) 544–547.
- <span id="page-13-3"></span>[6] S. Bouyuklieva, On the automorphisms of order 2 with fixed points for the extremal self-dual codes of length 24m , Des. Codes Cryptogr. 25 (2002) 5–13.
- <span id="page-14-8"></span><span id="page-14-4"></span>[7] C.W. Curtis, I. Reiner, Methods of representation theory with applications to finite groups and orders, Vol. I, Wiley Classics Library, (1990).
- <span id="page-14-1"></span>[8] R. Dontcheva, A.J.V. Zanten, S. Dodunekov, Binary self-dual codes with automorphism of composite order, IEEE Trans. Inform. Theory 50 (2004) 311–318.
- <span id="page-14-5"></span>[9] T. Feulner, G. Nebe, The automorphism group of an extremal [72, 36, 16] code does not contain  $\mathbb{Z}_7$  ,  $\mathbb{Z}_3 \times \mathbb{Z}_3$  ,  $\emph{or}~ D_{10}$  , IEEE Trans. Inform. Theory
- <span id="page-14-11"></span>[10] A.M. Gleason, Weight polynomials of self-dual codes and MacWilliams identies, 1970 Actes Congrés Internl. de Mathematique 3 (1971) 211–215.
- <span id="page-14-6"></span>[11] W.C. Huffman, Automorphisms of codes with application to extremal doubly even codes of length 48 , IEEE Trans. Inform. Theory IT-28 (1982) 511–521.
- [12] C.L. Mallows, N.J.A. Sloane, An upper bound for self-dual codes, Information and Control 22 (1973) 188–200.
- <span id="page-14-9"></span><span id="page-14-2"></span>[13] C.A. Mechor, P. Gaborit, On the classification of extremal [36, 18, 8] binary self-dual codes, IEEE Trans. Inform. Theory 54 (2008) 4743–4750.
- <span id="page-14-3"></span>[14] G. Nebe, An extremal [72, 36, 16] binary code has no automorphism group containing  $\mathbb{Z}_2 \times \mathbb{Z}_4$ ,  $Q_8$ , or  $\mathbb{Z}_{10}$ , Finite Fields and their applications 18 (2012) 563–566.
- [15] E.A. O'Brein, W. Willems, On the automorphism group of a binary self-dual doubly-even [72, 36, 16] *code*, IEEE Trans. Inform. Theory 57 (2011) 4445-4451.
- <span id="page-14-10"></span><span id="page-14-7"></span>[16] V. Pless, A classification of self-orthogonal codes over  $\mathbb{F}_2$ , Discrete Math. 3 (1972) 209–246.
- <span id="page-14-0"></span>[17] N.J.A. Sloane, Binary codes, lattices, and sphere-packings, Combinatorial Surveys, P.J. Cameron, Ed. (1977) 117–164.
- [18] N.J.A. Sloane, Is there a  $(72; 36)$   $d = 16$  self-dual code?, IEEE Trans. Inform. Theory 2 (1973) 251.