

The Area of a Polygon with an Inscribed Circle

Marshall W. Buck* Robert L. Siddon†

April 2005

Abstract

Heron's formula states that the area K of a triangle with sides a , b , and c is given by

$$K = \sqrt{s(s-a)(s-b)(s-c)}$$

where s is the semiperimeter $(a + b + c)/2$. Brahmagupta, Robbins, Roskies, and Maley generalized this formula for polygons of up to eight sides inscribed in a circle. In this paper we derive formulas giving the areas of any n -gon, with odd n , in terms of the ordered list of side lengths, if the n -gon is circumscribed about a circle (instead of being inscribed in a circle). Unlike the cyclic polygon problem, where the order of the sides does not matter, for the inscribed circle problem (our case) it does matter. The solution is much easier than for the cyclic polygon problem, but it does generalize easily to all odd n . We also provide necessary and sufficient conditions for there to be solutions in the case of even n .

*Center for Communications Research, 805 Bunn Dr., Princeton, NJ 08540, buck@idaccr.org

†2706 Post Oak Ct., Annapolis, MD 21401, rsiddon@aahs.org

1 Introduction

Robbins wrote in 1992 [6]:

Since a triangle is determined by the lengths, a, b, c of its three sides, the area K of the triangle is determined by these three lengths. The well-known formula

$$K = \sqrt{s(s-a)(s-b)(s-c)}, \quad (1.1)$$

where s is the semiperimeter $(a + b + c)/2$, makes this dependence explicit. (This formula is usually ascribed to Heron of Alexandria, c. 60 BC, although some attribute it to Archimedes.)

For polygons of more than three sides, the lengths of the sides do not determine the polygon or its area. However, if we impose the condition that the polygon be convex and *cyclic*, (i.e., inscribed in a circle) then the area of the polygon is uniquely determined. Moreover, it is a symmetric function of the sides. ... Given positive real numbers a_1, \dots, a_n , one can construct a convex n -gon with the a_j 's as the lengths of the sides provided that the largest a_j is smaller than the sum of the remaining ones. In this case it is also possible to construct a convex cyclic n -gon with the same sides (this is not quite so easy to establish as one might imagine) and this cyclic n -gon has the largest area of all n -gons with the given side lengths.

Heron's formula has been generalized for cyclic polygons of up to eight sides [6, 5].

In this paper, we are interested in finding areas of polygons with n given side lengths subject to a different constraint. Instead of asking that the *vertices* of the polygon lie on a circle, we require that the *sides* of the polygon lie on (are tangent to) a circle.

There are several differences with the cyclic polygon problem. First, in the cases where n is even, there can be infinitely many solutions. Second, in the case where n is odd, there are only a finite number of solutions, but different orderings of the side lengths generally produce solutions with different areas. In both cases, for there to be any solutions at all, certain equalities and inequalities in the side lengths a_j must be satisfied.

2 Heron's Formula

Let us review the simplest case, $n = 3$, producing a simple proof of Heron's formula that will generalize to $n > 3$.

We are given three side lengths a, b, c , so the perimeter is $a + b + c$ and *semi-perimeter* is $s = (a + b + c)/2$. For there to be a triangle with those side lengths, the following (triangle) inequalities must be satisfied:

$$a \leq b + c$$

$$b \leq c + a$$

$$c \leq a + b.$$

For there to be an inscribed circle, and for the area of the triangle to be nonzero, we need strict inequalities above.




Figure 1: Diagram for proof of Heron's formula

Refer to Figure 1. Denote by A the vertex opposite the side of length a , by B the vertex opposite the side of length b , and by C the vertex opposite the side of length c . Thus, the side BC has length a , (more succinctly, $|BC| = a$); furthermore, $|AC| = b$ and $|AB| = c$. Let P be the center of the inscribed circle, and let D, E, F be the feet of perpendiculars dropped from P to the sides BC, CA, AB respectively. (Thus, D, E, F are the points of tangency of the triangle with the inscribed circle.) These perpendiculars are radii of the

circle, and all have the same length r . The area K of the triangle $\triangle ABC$ is the sum of the areas of the three triangles $\triangle ABP$, $\triangle BCP$, and $\triangle CAP$, having respective areas $cr/2$, $ar/2$, and $br/2$, summing to $(a + b + c)r/2$. Thus, we have the well-known formula relating the radius of the inscribed circle, the area of the triangle, and the semi-perimeter s :

$$K = sr. \tag{1}$$

The two tangents from A to the circle, AE and AF , have the same length t_A . Similarly, the tangents from B have length t_B , and the tangents from C have length t_C . Meanwhile, each side of the triangle is composed of two of these tangent lines. Thus, we have

$$\begin{aligned} a &= |BC| = |BD| + |DC| = t_B + t_C \\ b &= |AC| = |AE| + |EC| = t_A + t_C \\ c &= |AB| = |AF| + |FB| = t_A + t_B \\ a + b + c &= 2(t_A + t_B + t_C) \\ s &= t_A + t_B + t_C \\ t_A &= s - a \\ t_B &= s - b \\ t_C &= s - c. \end{aligned}$$

Let $\theta_A, \theta_B, \theta_C$ denote the angles $\angle APE = \angle APF$, $\angle BPD = \angle BPF$, and $\angle CPD = \angle CPE$ respectively. We must have $\theta_A + \theta_B + \theta_C = \pi$. Meanwhile, we observe that $\theta_A, \theta_B, \theta_C$ are the arguments of the three complex numbers $r + it_A, r + it_B, r + it_C$, respectively. The argument of the complex number $(r + it_A)(r + it_B)(r + it_C)$ must then be $\theta_A + \theta_B + \theta_C = \pi$, which means that that the imaginary part of $(r + it_A)(r + it_B)(r + it_C)$ must be zero. But

$$\Im \{(r + it_A)(r + it_B)(r + it_C)\} = (t_A + t_B + t_C)r^2 - t_A t_B t_C,$$

so

$$r^2 = \frac{t_A t_B t_C}{t_A + t_B + t_C} = \frac{(s - a)(s - b)(s - c)}{s}. \tag{2}$$

Combining (1) and (2), we have Heron's formula

$$K^2 = s^2 r^2 = s^2 \frac{(s - a)(s - b)(s - c)}{s} = s(s - a)(s - b)(s - c).$$

3 Generalization to $n > 3$: Finding the Tangent Lengths

Our solution involves two steps: (1) determine the tangent lengths from the side lengths; (2) determine the radius from the tangent lengths. The first step depends upon the ordering of the side lengths, but the second step produces a polynomial equation of degree $\lfloor (n-1)/2 \rfloor$ satisfied by r^2 , where the coefficients of the polynomial are elementary symmetric functions of the tangent lengths.

Denote the side lengths in cyclic order by a_1, a_2, \dots, a_n , and the tangent lengths in order by t_1, t_2, \dots, t_n , designated so that we have

$$\begin{aligned} a_1 &= t_1 + t_2 \\ a_2 &= t_2 + t_3 \\ &\vdots \\ a_n &= t_n + t_1. \end{aligned} \tag{3}$$

Again, we define the semi-perimeter

$$\begin{aligned} s &= (a_1 + a_2 + \dots + a_n)/2 \\ &= t_1 + t_2 + \dots + t_n, \end{aligned}$$

and (1) still holds in the general case.

If n is odd, we can determine the tangents t_i from the side lengths by inverting the linear equations above, getting each t_i as an alternating sum:

$$\begin{aligned} t_1 &= (a_1 - a_2 + a_3 - \dots + a_n)/2 \\ t_2 &= (a_2 - a_3 + a_4 - \dots - a_n + a_1)/2 \\ &\vdots \\ t_n &= (a_n - a_1 + a_2 - \dots + a_{n-1})/2. \end{aligned}$$

In this case, the condition that there be a polygon circumscribing a circle is that each of the t_i is positive, which converts into the following n inequalities in the side lengths:

$$a_1 + a_3 + a_5 + \dots + a_n > a_2 + a_4 + \dots + a_{n-1} \tag{4}$$

and all other cyclic rotations of the indices.

If n is even, we wish to find necessary and sufficient conditions on the a_i values and ordering, for there to be a corresponding set of positive tangent lengths satisfying the equations (3). First, we can write down equations determining each t_i in terms of a single one, t_1 :

$$\begin{aligned}
t_2 &= a_1 - t_1 \\
t_3 &= a_2 - t_2 = a_2 - a_1 + t_1 \\
t_4 &= a_3 - t_3 = a_3 - a_2 + a_1 - t_1 \\
&\vdots \\
t_n &= a_{n-1} - t_{n-1} = a_{n-1} - a_{n-2} + \cdots + a_1 - t_1 \\
t_1 &= a_n - a_{n-1} + \cdots + a_2 - a_1 + t_1.
\end{aligned}$$

For there to be any solution at all, it is necessary that the sum of the a_i for odd i is equal to the sum of the a_i for even i . Beyond that, it is necessary that we be able to choose t_1 to make all of the t_i positive. We can do that if and only if we can choose t_1 less than all of the following:

$$\begin{aligned}
&a_1 \\
&a_1 - a_2 + a_3 \\
&a_1 - a_2 + a_3 - a_4 + a_5 \\
&\vdots \\
&a_1 - a_2 + \cdots - a_{n-2} + a_{n-1},
\end{aligned}$$

and, simultaneously, greater than all of the following:

$$\begin{aligned}
&0 \\
&a_1 - a_2 \\
&a_1 - a_2 + a_3 - a_4 \\
&\vdots \\
&a_1 - a_2 + \cdots + a_{n-3} - a_{n-2}.
\end{aligned}$$

In other words, we need

$$\begin{aligned}
&\min(a_1, a_1 - a_2 + a_3, a_1 - a_2 + a_3 - a_4 + a_5, \dots, a_1 - a_2 + \cdots - a_{n-2} + a_{n-1}) > \\
&\max(0, a_1 - a_2, a_1 - a_2 + a_3 - a_4, \dots, a_1 - a_2 + \cdots + a_{n-3} - a_{n-2}).
\end{aligned}$$

This happens if the alternating sum of the a_i of length n vanishes, and if every contiguous, odd length, alternating sum is positive. Of course, we are given that the length one sums, like a_2 , are positive. But, we also need (if $n \geq 6$) the additional alternating sum conditions like $a_1 - a_2 + a_3 > 0$. Not all of these are needed, even of those length $n/2$ or less. The vanishing of the alternating sum of length n allows us to derive the alternating sum condition for a chain of length k from the complementary chain of length $n - k$. If we count the conditions $a_j > 0$, there are a total of $n^2/4$ independent inequalities needed, and these can be taken to be alternating sums of any odd length, starting from an a_j for odd j . It is easy to show that all the inequalities are necessary. Special cases of this result for $n = 4, 6, 8$ are stated below.

Proposition 1 *There will be a circle-enclosing quadrilateral with given side lengths a_1, a_2, a_3, a_4 if and only if the equality*

$$a_1 + a_3 = a_2 + a_4,$$

and the following four inequalities are satisfied:

$$a_1 > 0, a_2 > 0, a_3 > 0, a_4 > 0.$$

Proposition 2 *There will be a circle-enclosing hexagon with given side lengths a_1, a_2, \dots, a_6 if and only if the equality*

$$a_1 + a_3 + a_5 = a_2 + a_4 + a_6,$$

and the following nine inequalities are satisfied:

$$a_1 > 0, a_2 > 0, \dots, a_6 > 0$$

$$a_1 - a_2 + a_3 > 0$$

$$a_3 - a_4 + a_5 > 0$$

$$a_5 - a_6 + a_1 > 0.$$

Proposition 3 *There will be a circle-enclosing octagon with given side lengths a_1, a_2, \dots, a_8 if and only if the equality*

$$a_1 + a_3 + a_5 + a_7 = a_2 + a_4 + a_6 + a_8,$$

and the following 16 inequalities are satisfied:

$$\begin{aligned} a_1 > 0, a_2 > 0, \dots, a_8 > 0 \\ a_1 - a_2 + a_3 > 0 \\ a_2 - a_3 + a_4 > 0 \\ \vdots \\ a_7 - a_8 + a_1 > 0 \\ a_8 - a_1 + a_2 > 0. \end{aligned}$$

Computationally, rather than check the large number of inequalities ($n^2/4$ such) it is better to proceed as follows. From left to right, incrementally calculate the sum $v = a_1 - a_2 + a_3 - \dots - a_n$. We calculate the allowed range for t_1 as the interval (l, r) , starting with the interval $(0, a_1)$ and then executing the following pseudo C-code:

```
double u[n+1]; // u[1],...,u[n] are side lengths
double r, ell, v=0.0; /* (ell,r) will be the choice interval for t_1
ell = 0.0; r = u[1]; /* initialize choice interval to (0,u_1))
for(int i = 1; i <= n; i++){
    if( i&1 ) { /* i is odd */
        v += u[i];
        if( v < r ) r = v;
    } else { /* i is even */
        v -= u[i];
        if( v > ell ) ell = v;
    }
}
if( v != 0.0 ) printf("The alternating sum is not zero as required\n");
if( ell >= r ) printf("The choice interval for t_1 is empty\n");
```

(In an actual program, we would not compare a floating point number to “0.0”.)

4 Generalization to $n > 3$: from Tangent Lengths to the Radius of the Inscribed Circle

Given the tangent lengths t_1, t_2, \dots, t_n (summing to the semiperimeter s , and satisfying the edge length constraints $a_1 = t_1 + t_2$, $a_2 = t_2 + t_3$, etc.) we may now construct the polygon about any circle of radius r , by using complex numbers $q_1 = r, q_2, \dots, q_n$ to represent the points of tangency and the complex numbers p_1, p_2, \dots, p_n to represent the vertices of the polygon. From each q_j we produce the next p_j point, in the counterclockwise direction, and having tangent length t_j , by multiplying q_j by $(r + it_j)/r$. Similarly, we calculate the next q_{j+1} by multiplying p_j by $r/(r - it_j)$. Thus, we have

$$\begin{aligned}
 q_1 &= r \\
 p_1 &= \frac{r + it_1}{r} q_1 \\
 q_2 &= \frac{r + it_2}{r - it_1} p_1 \\
 p_2 &= \frac{r + it_3}{r} q_2 \\
 q_3 &= \frac{r + it_3}{r - it_2} p_2 \\
 p_3 &= \frac{r + it_4}{r} q_3 \\
 &\vdots \\
 p_n &= \frac{r + it_n}{r} q_n \\
 q_1 &= \frac{r + it_1}{r - it_n} p_n.
 \end{aligned}$$

For the polygon to close, it is necessary and sufficient that

$$\prod_{j=1}^n \frac{r + it_j}{r - it_j} = 1.$$

This happens if and only if the product

$$\prod_{j=1}^n (r + it_j)$$

is real. Define the elementary symmetric functions $\sigma_1, \sigma_2, \dots, \sigma_n$ of the tangent lengths t_1, t_2, \dots, t_n as usual. In particular, $\sigma_1 = t_1 + t_2 + \dots + t_n = s$ and $\sigma_n = t_1 t_2 \dots t_n$. Also, define $\sigma_0 = 1$. These can be calculated from the identity

$$\prod_{j=1}^n (x + t_j) = \sum_{j=0}^n \sigma_j x^{n-j}.$$

Rewriting the identity, we obtain

$$\prod_{j=1}^n (r + it_j) = \sum_{j=0}^n i^j \sigma_j r^{n-j}.$$

Meanwhile, we want the imaginary part of the left-hand side to vanish, so we can rewrite our condition as

$$\sigma_1 r^{n-1} - \sigma_3 r^{n-3} + \dots = 0. \quad (5)$$

When $n = 2k + 1$ is odd, r^2 must be a root of the polynomial

$$\sigma_1 x^k - \sigma_3 x^{k-1} + \dots + (-1)^k \sigma_n. \quad (6)$$

When $n = 2k + 2$ is even, we can take out one factor of r from (5), and say that r^2 must be a root of the polynomial

$$\sigma_1 x^k - \sigma_3 x^{k-1} + \dots + (-1)^k \sigma_{n-1}. \quad (7)$$

In both cases there will be exactly k roots $r_1^2 > r_2^2 > \dots > r_k^2$. The largest radius, r_1 , will give rise to a convex polygon enclosing the circle of radius r_1 . The other roots will have the polygon going around the circle multiple times. For example, for $n = 5$ and $k = 2$, the radius r_2 will be that of a circle inscribed in a pentagonal star.

One way to see that there are k solutions is to consider the sum of the central angles that needs to be a multiple of π . Let

$$f(r) = \tan^{-1} \frac{t_1}{r} + \tan^{-1} \frac{t_2}{r} + \dots + \tan^{-1} \frac{t_n}{r}.$$

The continuous function $f(r)$ on the interval $r \in (0, \infty)$ converges to $\frac{n\pi}{2}$ as $r \rightarrow 0$, converges to 0 as $r \rightarrow \infty$, and monotonically decreases as r goes from 0 to ∞ . Thus, by the intermediate value theorem, f takes on the values $\pi, 2\pi, \dots, k\pi$ at $r = r_1, r = r_2, \dots, r = r_k$.

Since general polynomial equations of degree up to four are solvable in radicals, we can similarly solve our problem for $k \leq 4$, thus for $n \leq 10$.

5 Quadrilaterals

Given four sides a_1, a_2, a_3, a_4 , the largest area of a quadrilateral is for the cyclic quadrilateral, and that area is given by Brahmagupta's formula

$$K = \sqrt{(s - a_1)(s - a_2)(s - a_3)(s - a_4)}. \quad (8)$$

Meanwhile, if $a_1 + a_3 = a_2 + a_4$, we will be able to inscribe circles in various ways. We would like to find the quadrilateral of largest area, among those containing a circle tangent to all four sides.

In this case, because of the constraint on the a_j 's, Brahmagupta's formula simplifies. For example,

$$s - a_1 = (-a_1 + a_2 + a_3 + a_4)/2 = (-a_1 + a_2 + a_1 + a_2)/2 = a_2.$$

So,

$$K = \sqrt{a_1 a_2 a_3 a_4} \quad (9)$$

in this special case. (See also [1, Exercise 1, page 60].)

If we set the tangent lengths to be

$$\begin{aligned} t_1 &= \frac{a_1 a_4}{a_1 + a_3} \\ t_2 &= \frac{a_2 a_1}{a_1 + a_3} \\ t_3 &= \frac{a_3 a_2}{a_1 + a_3} \\ t_4 &= \frac{a_4 a_3}{a_1 + a_3} \end{aligned} \quad (10)$$

it is easy to check that the conditions $a_1 = t_1 + t_2$, etc. are satisfied (making use of $a_1 + a_3 = a_2 + a_4$). Furthermore, one can verify that $t_1 t_3 = t_2 t_4$, and $s = a_1 + a_3 = a_2 + a_4$. Now we can calculate the area using our earlier results. First, r^2 satisfies $\sigma_1 r^2 = \sigma_3$, $\sigma_1 = s$, and

$$\begin{aligned} \sigma_3 &= t_1 t_2 t_3 + t_1 t_2 t_4 + t_1 t_3 t_4 + t_2 t_3 t_4 \\ &= t_1 t_3 (t_2 + t_4) + t_2 t_4 (t_1 + t_3) \\ &= t_1 t_3 (t_1 + t_2 + t_3 + t_4) \\ &= s t_1 t_3 \\ &= s^{-1} a_1 a_2 a_3 a_4, \end{aligned}$$

so we have

$$\begin{aligned} K^2 &= r^2 s^2 = s^2(\sigma_3/\sigma_1) \\ &= s\sigma_3 = a_1 a_2 a_3 a_4. \end{aligned}$$

This construction of the quadrilateral achieves the maximum area, so it simultaneously is inscribed in a circle and has a circle inscribed in it.

Samuel Kutin [4] remarks that the solution (10) can be derived simply as follows. He notes that in the case of a quadrilateral inscribed in a circle, opposite internal angles are supplementary, so opposite half internal angles are complementary. If there is an inscribed circle of radius r , then there are similar right triangles telling us that $t_1/r = r/t_3$ and $t_2/r = r/t_4$, and therefore $t_1 t_3 = t_2 t_4$. But, from this condition alone, we can then derive (10). For example,

$$\begin{aligned} a_1 a_2 &= (t_1 + t_2)(t_2 + t_3) \\ &= t_1 t_2 + t_1 t_3 + t_2^2 + t_2 t_3 \\ &= t_1 t_2 + t_2 t_4 + t_2^2 + t_2 t_3 \\ &= t_2(t_1 + t_4 + t_2 + t_3) \\ &= t_2((t_1 + t_2) + (t_3 + t_4)) \\ &= t_2(a_1 + a_3), \end{aligned}$$

giving $t_2 = a_1 a_2 / (a_1 + a_3)$, the second line of (10).

Stephen DiPippo [2] gives a nice way to construct examples (see Figure 2). He starts with any right triangle, makes a mirror image copy of the triangle about the hypotenuse, joins the two triangles to form a quadrilateral with an inscribed circle. Meanwhile, the hypotenuse is a diameter of a circle containing all four vertices of the quadrilateral. This makes one example of such a quadrilateral, but it is “too symmetric”. Then he remarks that one can apply Poncelet’s Porism [3], and know that one can start at *any* point on the outer circle, drop a tangent to the inner circle, intersect that tangent with the outer circle, drop another tangent, and continue, forming a new quadrilateral having the same pair of circles on the inside and outside. By such means, he constructs a general example.




Figure 2: Construction of quadrilaterals via Poncelet's Porism

References

- [1] H. S. M. Coxeter and S. L. Greitzer. *Geometry Revisited*, The Mathematical Association of America, 1967.
- [2] Stephen A. DiPippo. Personal Communication.
- [3] Phillip Griffiths and Joseph Harris. On Cayley's explicit solution to Poncelet's Porism, *L'Enseignement Mathématique*, 24(1-2):31–40, January–June 1978.
- [4] Samuel A. Kutin. Personal Communication.
- [5] F. Miller Maley, David P. Robbins, and Julie Roskies. On the areas of cyclic and semicyclic polygons, *Advances in Applied Mathematics*, 34(4):669–689, 2005.
- [6] David P. Robbins. Areas of polygons inscribed in a circle, *Discrete and Computational Geometry*, 12:223–236, 1994.

Keywords

polygon, cyclic, circle, inscribed, circumscribed, area, radius, Heron, Brahmagupta, Robbins, Maley, Litman, Roskies, Hero