

# DE–SUSPENSION OF FREE $S^3$ –ACTIONS ON HOMOTOPY SPHERES

Issam. H. Kaddoura.\*†  
Lebanese International University  
Department of Mathematics

## Abstract

In the paper of Montgomery, *D.* and Yang, *C.T.* [5], they discuss the de–suspension of smooth free actions of  $S^1$  on  $(2n + 1)$ –dimensional homotopy spheres. In this paper we discuss the de-suspension of smooth free actions of  $S^3$  on  $(4n + 3)$ –dimensional homotopy spheres.

*Keywords:* Free action, quaterionic projective space, homotopy sphere, de–suspension, principal fibration, orbit space, framed surgery, Index, Lie groups, imbedding, tangent and normal bundles, exact homology and cohomology sequences, attaching a handle, tubular neighborhood, deformation retract, lefschetz duality, Poincare duality, mapping cylinder, bilinear form, Hurwicz homomorphism.

## 1 Introduction

Through this paper,  $R^n$  denotes the Euclidean  $n$ –space,  $S^n$  denotes the unit  $n$ –sphere in  $R^{n+1}$  and  $QP(n)$  the quaterionic projective space, all having the usual differentiable structures. By a homotopy  $n$ –sphere abbreviated by  $\Sigma^n$ , we mean a closed differentiable  $n$ –manifold having the homotopy type of  $S^n$ , and by a homotopy quaterionic projective  $n$ –space, abbreviated by  $HQP(n)$ , we mean a closed differentiable  $4n$ –manifold having the homotopy type of  $QP(n)$ .  $\pi_n(M)$  denotes the  $n^{\text{th}}$ –homotopy group of  $M$ ,  $H_i(M)$  and  $H^i(M)$  denote the homology and cohomology of a space  $M$ , and assumed to be satisfying the Eilenberg–Steenrod axioms, See [[13],p.6]. It is well known that  $S^1$  and  $S^3$  are the only compact connected Lie groups which have free differentiable actions on homotopy spheres, [8, 6]. It follows from Gleasons lemma [1] that such an action is always a principal fibration which is homotopically equivalent to the classical Hopf fibration. In fact, there are always infinitely many differentiable distinct free actions of  $S^3$  on  $\Sigma^{4n+3}$  for  $n \geq 2$ , [16], such that for any differentiable action of the group  $S^3$  on  $\Sigma^{4n+3}$ , the orbit space  $\Sigma^{4n+3}/S^3$  is a  $HQP(n)$ , and that for any  $HQP(n)$  there is, up to differentiable equivalence a unique free differentiable action of  $S^3$  on  $\Sigma^{4n+3}$ , such that  $\Sigma^{4n+3}/S^3$  is diffeomorphic to  $HQP(n)$ . Hence, there is one to one correspondence between differentiable equivalence classes of free

---

\*e.mail : issam.kaddoura@liu.edu.lb.

†Faculty of Science Baghdad University.

differentiable actions of  $S^3$  on  $\sum^{4n+3}$  and diffeomorphism classes of  $HQP(n)$ 's. If we let  $M = \sum^{4n+3}/S^3$ , there is a differentiable homotopy equivalence  $f : M \rightarrow QP(n)$  which is transverse regular on  $QP(n-1)$ , denoted  $f \pitchfork QP(n-1)$ . Therefore  $N = f^{-1}(QP(n-1))$  is a differentiable submanifold of  $M$  of codimension 4. By means of framed surgery as devised in Browder [15] and Novikov [9], it is always possible to kill the kernel of  $f_{*i} : \pi_i(N) \rightarrow \pi_i(QP(n-1))$  for  $i = 0, 1, \dots, 2n-3$  (below the middle dimension). That means, there is a differentiable homotopy equivalence  $f^1 : M \rightarrow QP(n)$  which is homotopic to  $f$  and  $f^1 \pitchfork QP(n-1)$  such that if we let  $N^1 = f^{-1}(QP(n-1))$ , then  $f_{*i}^1 : \pi_i(N^1) \rightarrow \pi_i(QP(n-1))$  has a trivial kernel for  $i = 0, 1, 2, \dots, 2n-3$ . But for  $i = 2n-2$  (at the middle dimension) there is an obstruction called the index. In fact a necessary and sufficient condition for killing the kernel of  $f_{*i}$  for  $i = 2n-2$ , is that this index vanishes. This index turns out to be dependent only on the action  $(S^3, \sum^{4n+3})$  see [7]. We note that if we succeed to kill the kernel of  $f_{*i} : \pi_i(N) \rightarrow \pi_i(QP(n-1))$  for  $i = 0, 1, 2, \dots, 2n-2$ , hence by Poincare duality on the kernels, we succeed to kill the kernel of  $f_{*i}$  for all  $i = 0, 1, 2, \dots, 4n-4$ , then there is a differentiably imbedded  $HQP(n-1)$  in  $HQP(n)$  and hence there is a differentiably imbedded  $S^3$ -invariant  $\sum^{4n-1}$  in  $\sum^{4n+3}$  called a characteristic homotopy  $(4n-1)$ -sphere this what is called a de-suspension of the action.

## 2 Basic Construction

In this section, we give a construction which will be needed in our work. It is essentially the same construction as seen in [15] and [9] with certain modifications are made as in Haefliger [2], Levine [9], Montgomery and Yang [5]. For any manifold  $M$ ,  $\tau(M)$  denotes the tangent bundle of  $M$  and if  $N$  is a submanifold of  $M$ ,  $\nu(N, M)$  denotes the normal bundle of  $N$  in  $M$ . By an  $r$ -frame on  $N$ , we mean an  $r$ -tuple of linearly independent vector fields in  $\tau(M)|_N$ .  $D^n$  denotes the closed unit disk in  $R^n$  and  $B^n = D^n - S^{n-1}$ . Let  $M$  and  $M^1$  be closed  $\ell$ -manifolds,  $N^1$  a closed  $m$ -submanifold of  $M^1$ ,  $f : M \rightarrow M^1$  a map which is transverse regular on  $N^1$ ,  $N = f^{-1}(N^1)$  and  $h : S^{k-1} \rightarrow N$  is an imbedding. We now state assumptions as by Haefliger in [2] and Levine in [9] and exactly as Montgomery and Yang in [5]. Later these assumptions will be verified for the construction made in this paper.

- (A)  $0 \leq k-1 \leq \frac{m}{2}$  and  $2 \leq m+2 \leq \ell$ .
- (B)  $f \circ h : S^{k-1} \rightarrow N^1$  is null homotopic and in fact  $f \circ h : (S^{k-1}) = y$  a point in  $N^1$ .
- (C)  $\pi_i(M - N) = 0$  for  $i \leq k-1$ .
- (D)  $\pi_i(M^1 - N^1) = 0$  for  $i \leq \ell$ .

Choose an  $(\ell-m)$  frame of  $\nu(M^1 - N^1)$  at  $y$  and let  $(v_1, v_2, \dots, v_{\ell-m})$  be its inuced  $(\ell-m)$  frame on  $h(S^{k-1})$  by  $f$ . Then  $h$  can be extended to an imbedding of  $D^k$  into  $M$ , also denoted by  $h$ , such that  $h(B^k) \subset M - N$  and  $v_1$  points radically into  $h(D^k)$ . we also assume :

- (E) The frame  $(v_1, v_2, \dots, v_{\ell-m})$  can be extended to an  $(\ell-m-1)$  frame on  $h(D^k)$  in  $\nu(h(D^k), M)$ . The obstruction to this extention is an element  $\lambda \in \pi_{k-1}(V_{\ell-m-1}(R^{\ell-k}))$

where  $V_{\ell-m-1}(R^{\ell-k})$  is the stiefel manifold of orthonormal  $(\ell-m-1)$  frames in  $R^{\ell-k}$  [[4],p.83]. Hence (E) means  $\lambda = 0$ . Notice that when  $h : S^{k-1} \rightarrow V_{\ell-m-1}(R^{\ell-k})$ , then  $h$  can be extended to an imbedding  $h^1 : D^k \rightarrow V_{\ell-m-1}(R^{\ell-k})$ , provided the obstruction which is an element  $\lambda \in H^k(D^k, S^{k-1}, \pi_{k-1}(V_{\ell-m-1}(R^{\ell-k})))$  vanishes. Using the exact cohomology sequence of the pair  $(D^k, S^{k-1})$ :

$$\rightarrow H^{k-1}(D^k, G) \rightarrow H^{k-1}(S^{k-1}, G) \rightarrow H^k(D^k, S^{k-1}, G) \rightarrow H^k(S^{k-1}, G) \rightarrow$$

where  $G = \pi_{k-1}(V_{\ell-m-1}(R^{\ell-k}))$ , we get :

$$0 \rightarrow \pi_{k-1}(V_{\ell-m-1}(R^{\ell-k})) \rightarrow H^k(D^k, S^{k-1}, \pi_{k-1}(V_{\ell-m-1}(R^{\ell-k}))) \rightarrow 0$$

$$\text{Hence, } H^k(D^k, S^{k-1}, \pi_{k-1}(V_{\ell-m-1}(R^{\ell-k}))) = \pi_{k-1}(V_{\ell-m-1}(R^{\ell-k})).$$

Moreover, if  $2(k-1) \leq m$ , then  $\pi_{k-1}(V_{\ell-m-1}(R^{\ell-k})) \cong 0$ , i.e, (E) holds for  $2(k-1) \leq m$ . Under these assumptions Haefliger in [2] has shown that the map  $f$  can be extended to a map  $F : M \times [0, 1] \rightarrow M^1$  such that  $F \pitchfork N^1$  and  $B = F^{-1}(N^1)$  is a compact  $(m+1)$ -manifold in  $M \times [0, 1]$  which is a framed cobordism from  $N$  to  $N^1 = B \cap (M \times \{1\})$  obtained by attaching a handle to  $N \times [0, 1]$  at  $h(S^{k-1})$ . Let  $f_1 = F|_{M \times \{1\}}$ , which is also regarded as the map of  $M$  into  $M^1$  given by  $x \rightarrow F(x, 1)$ . Then  $f_1$  is a homotopy equivalence such that  $f \sim f_1$ ,  $f_1 \pitchfork N^1$  and  $N_1 = f_1^{-1}(N^1)$  is obtained from  $N = f^{-1}(N^1)$  by performing a framed surgery at  $h(S^{k-1})$ .

### 3 Surgery below the middle dimension

Suppose that a free differentiable action of  $S^3$  on  $\sum^{4n+3}$  is given,  $n \geq 2$ . Let  $M = \sum^{4n+3}/S^3$ , then there is a homotopy equivalence  $f : M \rightarrow QP(n)$ , let  $\sigma : S^i \rightarrow M$  be an imbedding representing a generator of  $\pi_i(M)$ , by the relative transversality theorem, we may assume  $f$  is transverse regular on  $QP(n-1)$ , hence  $N = f^{-1}(QP(n-1))$  is a closed manifold of  $M$  of codimension 4 and containing  $\sigma(S^i)$  for  $i < 2n-1$ .

#### Lemma 3.1

- (a)  $M - N$  is connected, i.e.,  $\pi_0(M - N) = 0$ .
- (b)  $\pi_i(M - N) = 0$  for  $i = 1, 2$ .

#### Proof.

(a) let  $x, y \in M - N$  then  $x, y \in M$ , but  $M$  is connected then there exist a path  $\alpha : [0, 1] \rightarrow M$  such that  $\alpha(0) = x$  and  $\alpha(1) = y$ . By transversality homotopy theorem, see [[14],p.70], there exist a path  $\alpha^1 : [0, 1] \rightarrow M$  such that  $\alpha^1(0) = x$  and  $\alpha^1(1) = y$  and  $\alpha^1 \pitchfork N$ . But  $\dim \alpha^1([0, 1]) + \dim N < \dim M$ , thus  $\alpha^1 \cap N = \emptyset$  and  $\alpha^1([0, 1]) \subset M - N$ , hence  $M - N$  is pathwise connected and therefore is connected.

(b) Let  $\alpha : S^1 \rightarrow M - N$  be an imbedding representing a generator of  $\pi_1(M - N)$ ,  $\alpha$  can be regarded as an imbedding into  $M$ ,  $\alpha : S^1 \rightarrow M$ . But  $\pi_1(M) = 0$  then  $\alpha$  can be extended to  $\alpha^1 : D^2 \rightarrow M$  such that  $\alpha^1|_{\partial D^2} = \alpha$ ,  $\alpha$  is homotopic to  $\alpha^1$  and  $\alpha^1 \pitchfork N$ .

But  $\dim \alpha^1(D^2) + \dim N < \dim M$  thus  $\alpha^1(D^2) \cap N = \phi$ , which implies  $\alpha^1(D^2) \subset M - N$  therefore  $\pi_1(M - N) = 0$ , similarly  $\pi_2(M - N) = 0$ .

**Lemma 3.2 :**  $f_{*i} : H_i(N) \rightarrow H_i(QP(n-1))$  is onto  $\forall i$ .

In particular  $f_{*4n-4} : H_{4n-4}(N) \rightarrow H_{4n-4}(QP(n-1))$  is an isomorphism.

**proof.** Let  $A^1$  be the complement of an open tubular neighborhood of  $QP(n-1)$  in  $QP(n)$  such that  $A = f^{-1}(A^1)$  is the complement of open tubular neighborhood of  $N$  in  $M$ . It is clear that  $f^* : H^{4n}(QP(n)) \rightarrow H^{4n}(M)$  is an isomorphism. Consider the cohomology exact sequence of the pair  $(QP(n), A^1)$  :

$$\rightarrow H^{4n-1}(A^1) \rightarrow H^{4n}(QP(n), A^1) \rightarrow H^{4n}(QP(n)) \rightarrow H^{4n}(A^1) \rightarrow$$

But,  $QP(n) = QP(n-1) \cup D^{4n}$ ,  $A^1$  is a deformation retract of  $QP(n) - QP(n-1)$ , thus  $A^1$  has the homotopy type of  $D^{4n}$ , hence  $H^{4n-1}(A^1) = 0$  and  $H^{4n}(A^1) = 0$ . Therefore  $H^{4n}(QP(n), A^1) \rightarrow H^{4n}(QP(n))$  is an isomorphism. Moreover, the cohomology exact sequence of  $(M, A)$ :

$$\rightarrow H^{4n-1}(A) \rightarrow H^{4n}(M, A) \rightarrow H^{4n}(M) \rightarrow H^{4n}(A) \rightarrow$$

$A$  is a deformation retract of  $M - N$ , and by Hurewicz theorem, [[10],p.349]:

$H_1(A) \cong \pi_1(A) \cong \pi_1(M - N) = 0$ . We get  $H^{4n-1}(A) = 0$  and  $H^{4n}(A) = 0$ . Therefore  $H^{4n}(M, A) \rightarrow H^{4n}(M)$  is an isomorphism. By the isomorphism theorem of Thom, see [6],  $H^{4n-4}(QP(n-1)) \rightarrow H^{4n}(QP(n), A^1)$  and  $H^{4n-4}(N) \rightarrow H^{4n}(M, A)$  are isomorphisms.

It follows that  $f^* : H^{4n-4}(QP(n-1)) \rightarrow H^{4n-4}(N)$  is an isomorphism.

Claim:  $f^{*i} : H^i(QP(n-1)) \rightarrow H^i(N)$  is one to one  $\forall i$ .

**Proof.** Let  $\alpha \neq 0 \in H^i(QP(n-1))$ , then  $\exists \beta \in H^{4n-4-i}(QP(n-1))$  such that  $\alpha \cup \beta = d \neq 0 \in H^{4n-4}(QP(n-1))$ . This implies  $f_{*4n-4}^*(\alpha \cup \beta) = f^*(\alpha) \cup f^*(\beta) = f^*(d) \neq 0$ , i.e.,  $f^{*i} : H^i(QP(n-1)) \rightarrow H^i(N)$  is one to one  $\forall i$ . Then by duality between homology and cohomology  $f_{*i} : H_i(N) \rightarrow H_i(QP(n-1))$  is onto  $\forall i$

**Lemma 3.3** If  $k \leq 2n - 1$  such that  $f_{*i} : H_i(N) \rightarrow H_i(QP(n-1))$  is 1 - 1 for  $i < k - 1$ , then  $H_{k-1}(M - N) = 0$ .

**Proof.** By lemma 2.2 and by the hypothesis we get  $f_{*i} : H_i(N) \rightarrow H_i(QP(n-1))$  is an isomorphism for  $i < k - 1$ . Therefore, by Poincaré duality [[12],p.138],  $f^* : H^{4n-4-i}(QP(n-1)) \rightarrow H^{4n-4-i}(N)$  is an isomorphism for  $i < k - 1$ , so that  $H^{4n-4-i}(M) \rightarrow H^{4n-4-i}(N)$  induced by the inclusion map is an isomorphism for  $i < k - 1$ , [12]. Consider the exact cohomology sequence of  $(M, N)$ :

$$\rightarrow H^{4n-4-i}(M, N) \rightarrow H^{4n-4-i}(M) \rightarrow H^{4n-4-i}(N) \rightarrow H^{4n-3-i}(M, N) \rightarrow$$

for  $i = k - 4$  :

$$\rightarrow H^{4n-k}(M, N) \rightarrow H^{4n-k}(M) \rightarrow H^{4n-k}(N) \rightarrow H^{4n-k+1}(M, N) \rightarrow$$

$H^{4n-4}(M) \cong H^{4n-k}(N)$  and  $H^{4n-k+1}(M) \cong H^{4n-k+1}(N)$  hence,  $H^{4n-k+1}(M, N) \cong 0$  by Lefschetz duality [11], we get  $H_{k-1}(M - N) \cong H^{4n-k+1}(M, N) \cong 0$ .

**Lemma 3.4** If  $k \leq 2n - 1$  such that  $f_{*i} : \pi_i(N) \rightarrow \pi_i(QP(n - 1))$  is 1 - 1 for  $i < k - 1$ , then both  $f_{*i} : \pi_i(N) \rightarrow \pi_i(QP(n - 1))$ , and  $f_{*i} : H_i(N) \rightarrow H_i(QP(n - 1))$  are isomorphisms for  $i < k - 1$ . Moreover, if  $\rho : \pi_{k-1}(N) \rightarrow H_{k-1}(N)$  is the Hurewicz homomorphism,  $\ker(f_{*\pi_{k-1}})$  is the kernel of  $f_* : \pi_{k-1}(N) \rightarrow \pi_{k-1}(QP(n - 1))$  and  $\ker(f_{*H_{k-1}})$  is the kernel of  $f_* : H_{k-1}(N) \rightarrow H_{k-1}(QP(n - 1))$ , then for  $k \neq 2$   $\rho(\ker(f_{*\pi_{k-1}})) \cong \ker(f_{*H_{k-1}})$  and for  $k = 2$ ,  $\rho$  is onto and its kernel is the commutator subgroup of  $\ker(f_{*\pi_{k-1}})$ . Furthermore  $\pi_i(M - N) = 0$  for  $i \leq k - 1$ .

**Proof.** Let  $\sigma : S^i \rightarrow M$  be an imbedding representing a generator of  $\pi_i(M)$ ,  $\sigma(S^i)$  is contained in  $N$  as our assumption at the beginning of this section for  $i < 2n - 1$ ,  $f\rho : S^i \rightarrow QP(n - 1)$  represents a generator of  $\pi_i(QP(n - 1))$  for  $i \leq k - 1$ , so that  $f_{*i} : \pi_i(N) \rightarrow \pi_i(QP(n - 1))$  is an isomorphism for  $i < k - 1$ . and onto for  $i \leq k - 1$ . Let  $C_f$  be the mapping cylinder of  $f : N \rightarrow QP(n - 1)$ , i.e.,  $C_f = N \times I \cup_f QP(n - 1)$ , it is clear that  $QP(n - 1)$  is homotopically equivalent to  $C_f$ . See[[11],p.283]. We consider the two exact sequences which together with the Hurewicz homomorphism form the commutative diagram:

$$\begin{array}{ccccccc} \rightarrow \pi_i(N) & \xrightarrow{f_*} & \pi_i(QP(n - 1)) & \rightarrow & \pi_i(C_f, N) & \rightarrow & \pi_{i-1}(N) \rightarrow \\ & \rho \downarrow & \rho \downarrow & & \rho \downarrow & & \rho \downarrow \\ \rightarrow H_i(N) & \longrightarrow & H_i(QP(n - 1)) & \rightarrow & H_i(C_f, N) & \rightarrow & H_{i-1}(N) \rightarrow \end{array}$$

It is clear that  $\pi_i(C_f, N) = 0$  for  $i \leq k - 1$ , hence  $H_i(C_f, N) = 0$  for  $i \leq k - 1$ , and  $\pi_k(C_f, N) \cong H_k(C_f, N)$  ( By Hurewicz theorem ), [3]. Moreover,  $f_{*i} : H_i(N) \rightarrow H_i(QP(n - 1))$  is an isomorphism for  $i < k - 1$ , and onto for  $i \leq k - 1$ .

Now, the following diagram is commutative and its rows are exact :

$$\begin{array}{ccccccc} \rightarrow \pi_k(C_f, N) & \rightarrow & \ker(f_{*\pi_{k-1}}) & \rightarrow & 0 & & \\ & \rho \downarrow & \rho \downarrow & & & & \\ \rightarrow H_k(C_f, N) & \rightarrow & \ker(f_{*H_{k-1}}) & \rightarrow & 0 & & \end{array}$$

We get  $\rho(\ker(f_{*\pi_{k-1}})) \cong \ker(f_{*H_{k-1}})$ , see[10]. since  $f_{*i} : H_i(N) \rightarrow H_i(QP(n - 1))$  is an isomorphism for  $i < k - 1$ . and onto for  $i \leq k - 1$ , [13], so as  $H_i(N) \rightarrow H_i(M)$  induced by the inclusion map, similarly as in (Lemma 3.3) we get  $H_i(M - N) = 0$  for  $i \leq k - 1$ . But  $\pi_0(M - N) = 0$  and  $\pi_1(M - N) = 0$  (By Lemma 3.1), hence by Hurewicz theorem  $\pi_i(M - N) = 0$  for  $i \leq k - 1$ .

Now, we are in position to use the construction of the preceding section to kill the kernel of  $f_{*i} : \pi_i(N) \rightarrow \pi_i(QP(n - 1))$  for  $i < 2n - 2$ .

Consider  $f_* : \pi_0(N) \rightarrow \pi_0(QP(n - 1))$ , if the kernel is not trivial ,i.e.,  $N$  has more than one component, so there is an imbedding  $h : S^0 \rightarrow N$  such that  $h(S^0)$  is not contained in the same component of  $N$ . If we put  $M, QP(n), QP(n - 1), f$  and  $h$  in place of  $M, M^1, N^1, f$  and  $h$  respectively. We have to satisfy the required assumptions, then (A), (B), (C), (D) and (E) are easily verified. Therefore, we obtain a homology equivalence  $f_1 : M \rightarrow QP(n)$  such that  $f \sim f_1$  and  $f_1 \pitchfork QP(n - 1)$  such that

$f_1^{-1}(QP(n-1))$  has one less component than  $f^{-1}(QP(n-1))$ . Repeating this construction, if necessary, we can finally have a homotopy equivalence  $f^1 : M \rightarrow QP(n)$  such that  $f \sim f^1$ ,  $f^1 \pitchfork QP(n-1)$  and  $f^{1^{-1}}(QP(n-1))$  is connected and thus the kernel of  $f_*^1 : \pi_0(f^{1^{-1}}(QP(n-1))) \rightarrow \pi_0(QP(n-1))$  is trivial, that is we kill the kernel of  $f_* : \pi_0(N) \rightarrow \pi_0(QP(n-1))$  by using  $f^1$  to replace  $f$ . Consider  $f_* : \pi_1(N) \rightarrow \pi_1(QP(n-1))$  such that the kernel of  $f_* : \pi_0(N) \rightarrow \pi_0(QP(n-1))$  is already killed. Given an element  $\mu$  of the kernel of  $f_* : \pi_1(N) \rightarrow \pi_1(QP(n-1))$ , there is an imbedding  $h : S^1 \rightarrow N$  representing  $\mu$  with  $M$ ,  $QP(n)$ ,  $QP(n-1)$ ,  $f$  and  $h$  in place of  $M$ ,  $M^1$ ,  $N^1$ ,  $f$  and  $h$  respectively. we have to satisfy the required assumptions: (A), (B), (C), (D) and (E) for  $k = 2$ ,  $m = 4n - 4$  and  $\ell = 4n$ .

(A) It is obvious that  $0 \leq k - 1 \leq \frac{m}{2}$  and  $5 \leq m + 2 \leq \ell$ .

(B)  $\mu \in \ker f_*$  implies  $f_*(\mu) = 0$ , hence  $f \circ h : S^1 \rightarrow QP(n-1)$  is null homotopic.

(C)  $\pi_0(M - N) = 0$  and  $\pi_1(M - N) = 0$  (by lemma 3.1).

(D)  $\pi_i(QP(n) - QP(n-1)) = 0$ ,  $\forall i \leq 4n$ .

In fact,  $QP(n) - QP(n-1)$  has the homotopy type of  $D^{4n}$ . Moreover  $h : S^1 \rightarrow N$  can be extended to an imbedding  $h : D^2 \rightarrow M$  such that  $h(B^2) \subset M - N$  and  $v_1$  points radically into  $h(D^2)$ , where  $v_1$  as in section 2.

(E)  $\mu \in \pi_1(V_3(R^{4n-2})) = 0$ . See [[9],p.83].

Hence, there is a homotopy equivalence  $f_1 : M \rightarrow QP(n)$  such that  $f \sim f_1$ ,  $f_1 \pitchfork QP(n-1)$  and such that the kernel of  $f_{1*} : \pi_0(f_1^{-1}(QP(n-1))) \rightarrow \pi_0(QP(n-1))$  is trivial and the kernel of  $f_{1*} : \pi_1(f_1^{-1}(QP(n-1))) \rightarrow \pi_1(QP(n-1))$  is the quotient group of the kernel of  $f_* : \pi_1(f^{-1}(QP(n-1))) \rightarrow \pi_1(QP(n-1))$  by the normal subgroup generated by  $\mu$ . Repeating the same procedure if necessary, the kernel of  $f_* : \pi_1(f^{-1}(QP(n-1))) \rightarrow \pi_1(QP(n-1))$  can be killed also. Similarly, if  $k$  is an integer such that  $1 < k - 1 < 2n - 2$  and the kernel of  $f_* : \pi_i(N) \rightarrow \pi_i(QP(n-1))$  has been killed for  $i < k - 1$ , we can kill the kernel of  $f_* : \pi_{k-1}(N) \rightarrow \pi_{k-1}(QP(n-1))$  by using the same argument. Finally, we have a homotopy equivalence  $f^1 : M \rightarrow QP(n)$  such that  $f \sim f^1$ ,  $f^1 \pitchfork QP(n-1)$  and the kernel of  $f_{*i}^1 : \pi_i(f^{1^{-1}}(QP(n-1))) \rightarrow \pi_i(QP(n-1))$  is trivial for  $i < 2n - 2$ .

## 4 The obstruction at the middle dimension (The index)

Assume that a free differentiable action of the group  $S^3$  on  $\sum^{4n-3}$  is given,  $n \geq 2$  and let  $M = \sum^{4n-3}/S^3$  and  $f : M \rightarrow QP(n)$  is a homotopy equivalence which is transverse regular on  $QP(n-1)$  and let  $N = f^{-1}(QP(n-1))$ . As in the preceding section, we may assume that the kernel of  $f_{*i} : \pi_i(N) \rightarrow \pi_i(QP(n-1))$  has been killed for  $i < 2n - 2$ . Therefore by lemma 3.4,  $f_{*i} : \pi_i(N) \rightarrow \pi_i(QP(n-1))$  and  $f_{*i} : H_i(N) \rightarrow H_i(QP(n-1))$  are isomorphisms for  $i < 2n - 2$  and the Hurewicz homomorphism  $\sigma$  maps  $\ker(f_{*\pi_{2n-2}})$  isomorphically onto  $\ker(f_{*H_{2n-2}})$ . Consider the bilinear form  $F : \ker(f_{*H_{2n-2}}) \otimes \ker(f_{*H_{2n-2}}) \rightarrow Z$  which maps each  $\xi \otimes \mu \in \ker(f_{*H_{2n-2}}) \otimes \ker(f_{*H_{2n-2}})$  into the intersection number  $\xi \cdot \mu$ . It is symmetric and has a signature  $I$ . The integer  $I$  is independent of the choice of  $f$ . For more

details see [5].  $I$  is called the index of the action  $(S^3, \sum^{4n+3})$  and is written as  $I(S^3, \sum^{4n+3})$ .

**Lemma 4.1** Suppose  $n = 2s$  and the group  $\ker(f_{*\pi_s})$  is trivial for  $i < s$ . Then  $\ker(f_{*H_s})$  is free abelian, and the matrix of intersections of the base cycles of the group  $\ker(f_{*H_s})$  is unimodular. [[11],p.283].

**Theorem 4.1** For any integer  $n \geq 2$ , the index  $I$  vanishes iff there is a homotopy equivalence  $f : M \rightarrow QP(n)$  which is transverse regular on  $QP(n-1)$  and such that if  $N = f^{-1}(QP(n-1))$ , then  $f : N \rightarrow QP(n-1)$  is a homotopy equivalence.

**Proof.** If  $f : M \rightarrow QP(n)$  is a homotopy equivalence such that  $f : N \rightarrow QP(n-1)$  is a homotopy equivalence then  $\ker(f_{*H_i}) = 0 \quad \forall i$  and hence  $I$  vanishes. Conversely, assume that  $I$  vanishes and let  $f : M \rightarrow QP(n)$  is a homotopy equivalence described at the beginning of this section. Using lemma 4.1,  $\ker(f_{*H_{2n-2}})$  is a free abelian group which has the basis  $\{\xi_1, \xi_2, \dots, \xi_{2r-1}, \xi_{2r}\}$  such that

$$\xi_i \cdot \xi_j = \begin{cases} 1 & \text{if } (i, j) = (2t-1, 2t) \text{ for some } t \\ 0 & \text{otherwise} \end{cases}$$

Let  $h : S^{2n-2} \rightarrow N$  be an imbedding representing  $\xi_1$ . We can use the basic construction of section 2 to kill  $\xi_1$  and  $\xi_2$  at the same time. For  $\ell = 4n$ ,  $k = 2n-1$ ,  $m = 4n-4$ ,  $M = \sum^{4n+3}/S^3$ ,  $M^1 = QP(n)$ ,  $N^1 = QP(n-1)$  and  $N = f^{-1}(QP(n-1))$ .

- (A) It is obvious that  $0 \leq k-1 \leq \frac{m}{2}$  and  $5 \leq m+2 \leq \ell$ .
- (B)  $\xi_1 \in \ker(f_{*H_{2n-2}})$ , hence  $f \circ h : S^{2n-2} \rightarrow N^1$  is null homotopic.
- (C)  $\pi_i(M - N) = 0$  for  $i \leq 2n-2$ , by lemma 3.4
- (D)  $\pi_i(QP(n) - QP(n-1)) = 0$  for  $i \leq 4n$ .
- (E)  $\lambda \in \pi_{2n-2}(V_3(R^{2n+1})) = 0$ , [[4],p83].

This shows that  $\xi_1$  and  $\xi_2$  can be killed by the basic construction.

Similarly  $\xi_3, \xi_4, \dots, \xi_{2r-1}, \xi_{2r}$  can be killed two at the same time. Hence  $\ker(f_{*H_i})$  is trivial for  $i = 1, 2, \dots, 2n-2$ , and then  $\ker(f_{*\pi_i})$  is trivial for  $i = 1, 2, \dots, 2n-2$ . By Poincare duality on the kernels, we deduce that  $\ker(f_{*\pi_i})$  is trivial for  $i = 1, 2, \dots, 4n-4$ . Therefore,  $f : N \rightarrow QP(n-1)$  is a homotopy equivalence. Finally, if  $I(S^3, \sum^{4n+3}) = 0$ , then there is a differentiably imbedded  $HQP(n-1)$  in  $HQP(n)$  and hence there is a differentiably imbedded  $S^3$ -invariant  $\sum^{4n-1}$  in  $\sum^{4n+3}$  which is the de-suspension of the action.

**Acknowledgement** I am very grateful to professor Adil G. Naoum who guided me in different topics of this paper.

## References

- [1] A. Gleason , *Spaces with a compact Lie group of transformations*, proc. Amer. Math. Soc. I (1950), 35-43.
- [2] A. Haefliger, *Knotted  $(4k - 1)$  spheres in  $6k$  space*, Ann. of Math.75 (1962), 452-466.
- [3] C.R. Maunder, *Algebraic topology*. Van Nostrand Reinhold Comp.1970.
- [4] D. Husemoller, *Fiber bundles*, McGraw-Hill, Book Company, New York,(1966).
- [5] D.Montgomery, and C.T. Yang, *Free differentiable actions on homotopy spheres*. Proc. Con. on transformation groups, Springer, New York, (1968) 175-192.
- [6] H.T. Ku, *A note on Semi-free actions on homotopy spheres*, Proc. Amer. Math. Soc.22(1969),614-617.
- [7] H.T. Ku,and M.C.Ku,*Characteristic spheres of free differentiable actions of  $S^1$  and  $S^3$  on homotopy spheres*. Trans. Amer. Math. Soc.156 (1971), 493-504.
- [8] H.T. Ku,and M.C.Ku, *Free differentiable actions of  $S^1$  and  $S^3$  on homotopy spheres*. Proc. Amer. Math. Soc.25(1970),864-869.
- [9] J.Levine, *A classification of differentiable knots*, Ann. of Math. ,82 (1965), 15-50.
- [10] P.J. Hilton, and S.Wylie, *Homology Theory*, Cambridge Univ. Press, London and New York,(1960).
- [11] S.P. Novikov, *Homotopically equivalent smooth manifolds*, Translations Amer. Math. Soc.48 (1965), 271-396.
- [12] S.T.Hu, *Cohomology theory*, Markham Publishing Comp. (1968).
- [13] S.T. Hu, *Homology theory*, Holden-day, Inc.(1970).
- [14] V. Guillemin, and A. Pollack, *Differential topology*, Prentice-Hall, Inc, New Jersey, (1974).
- [15] W.Browder , *Homotopy type of differentiable manifolds*, Colloquium on algebraic topology, Aarhus,(1962),42-46.
- [16] W.C.Hsiang, *A note on free differentiable actions of  $S^1$  and  $S^3$  on homotopy spheres*, Ann. of Math.83 (1966), 266-272.