

A Phase Transition for Circle Maps and Cherry Flows

Liviana Palmisano
Université Paris-Sud
91405 Orsay, France

July 31, 2018

Abstract

We study C^2 weakly order preserving circle maps with a flat interval. The main result of the paper is about a sharp transition from degenerate geometry to bounded geometry depending on the degree of the singularities at the boundary of the flat interval. We prove that the non-wandering set has zero Hausdorff dimension in the case of degenerate geometry and it has Hausdorff dimension strictly greater than zero in the case of bounded geometry. Our results about circle maps allow to establish a sharp phase transition in the dynamics of Cherry flows.

1 Introduction

1.1 Almost Smooth Maps with a Flat Interval

We consider the class \mathcal{L} of continuous circle endomorphisms f of degree one for which an open arc U exists such that the following properties hold:

1. The image of U is one point.
2. The restriction of f to $\mathbb{S}^1 \setminus \overline{U}$ is a C^2 -diffeomorphism onto its image.

3. Let (a, b) be a preimage of U under the projection of the real line of \mathbb{S}^1 . On some right-sided neighborhood of b , f can be represented as

$$h_r \left((x - b)^{l_r} \right),$$

where h_r is a C^2 -diffeomorphism on an open neighborhood of b . Analogously, there exists a C^2 -diffeomorphism on an left-sided neighborhood of a such that f is of the form

$$h_l \left((a - x)^{l_l} \right).$$

The ordered pair (l_l, l_r) will be called the *critical exponent* of the map. If $l_l = l_r$ the map will be referred to as *symmetric*.

1.2 Rotation Number

Let $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ be a map in \mathcal{L} and $F : \mathbb{R} \rightarrow \mathbb{R}$ a lift of f . Then

$$\rho(F) := \lim_{n \rightarrow \infty} \frac{1}{n} (F^n(x) - x)$$

exists for all $x \in \mathbb{R}$.

$\rho(F)$ is independent of x and well defined up to an integer; that is, if \tilde{F} is another lift of f , then $\rho(F) - \rho(\tilde{F}) = F - \tilde{F} \in \mathbb{Z}$.

This remark justifies the following terminology:

Definition 1.1. $\rho(f) := [\rho(F)]$ is called the *rotation number* of f .

In the discussion that follows in this paper, it will often be convenient to identify f and F and subsets of \mathbb{S}^1 with the corresponding subsets of \mathbb{R} . Also we permanently assume that the rotation number is *irrational*.

1.3 Combinatorics

The irrational rotation number $\rho(f)$ can be written as an infinite continued fraction

$$\rho(f) = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots}}}.$$

where a_i are positive integers.

If we cut off the portion of the continued fraction beyond the n -th position, and write the resulting fraction in lowest terms as $\frac{p_n}{q_n}$ then the numbers q_n for $n \geq 1$ satisfy the recurrence relation

$$q_{n+1} = a_{n+1}q_n + q_{n-1}; \quad q_0 = 1; \quad q_1 = a_1.$$

The number q_n is the iterate of the rotation by $\rho(f)$ for which the orbit of any point makes the closest return so far to the point itself.

1.4 Discussion of the Results

Our class \mathcal{L} is designed to comprise the first return maps for smooth Cherry flows (see section 6). Following the terminology for these flows we will distinguish three cases and we will classify the maps in \mathcal{L} depending on their critical exponent (l, l) :

- the non-dissipative case if $l < 1$;
- the conservative case if $l = 1$;
- the dissipative case if $l > 1$.

In [8] which deals with the non-dissipative case it is proved that the non-wandering set Ω has zero Hausdorff dimension. In the conservative case, $\dim_{\text{H}}(\Omega) = 0$, [10]. The dissipative case is studied in [5] and [7]. In particular in [7], it is proved that Ω has zero Lebesgue measure. In the dissipative case, under the assumption that the rotation number is of bounded type (i.e. $\sup_i a_i < \infty$) $\dim_{\text{H}}(\Omega) < 1$, [3].

1.5 Statement of Results

We investigate symmetric almost smooth maps with a flat interval and with critical exponent (l, l) , $l > 1$ (dissipative case) and we get results about their non-wandering set, which, by Theorem C in [5], is exactly $\mathbb{S}^1 \setminus \bigcup_{i=0}^{\infty} f^{-i}(U)$. Also, we permanently assume that the rotation number is irrational and of bounded type.

Theorem 1.2. *If f is a map from the class \mathcal{L} with the critical exponent $l \in (1, 2]$, then the Hausdorff dimension of the non-wandering set is equal to 0.*

Theorem 1.3. *If f is a map from the class \mathcal{L} with the critical exponent $l > 2$, then the Hausdorff dimension of the non-wandering set is strictly greater than 0.*

These results show a transition of geometry in the dissipative case as the critical exponent of our map passes through 2.

As a corollary to Theorem 1.3 we find an important property of Cherry flows:

Theorem 1.4. *Let X be a Cherry vector field with $\lambda_1 > 0 > \lambda_2$ as eigenvalues of the saddle point and let f be its first return map.*

If $|\lambda_2| > 2\lambda_1$ and if f has rotation number of bounded type, then the quasi-minimal set has Hausdorff dimension strictly greater than 1.

2 Technical Tools

2.1 Basic Notations

We will introduce a simplified notation for backward and forward images of the flat interval U . Instead of $f^i(U)$ we will simply write \underline{i} ; for example, $\underline{0} = U$. Thus, for us, underlined positive integers represent points, and underlined non-positive integers represent intervals.

2.2 Distance between Points

We denote by $(a, b) = (b, a)$ the open shortest interval between a and b regardless of the order of these two points. The length of that interval in the natural metric on the circle will be denoted by $|a - b|$. Following [3], let us adopt now these conventions of notation:

- $|\underline{-i}|$ stands for the length of the interval $\underline{-i}$.
- Consider a point x and an interval $\underline{-i}$ not containing it. Then the distance from x to the closer endpoint of $\underline{-i}$ will be denoted by $|(x, \underline{-i})|$, and the distance to the more distant endpoint by $|(x, \underline{-i}]|$.
- We define the distance between the endpoints of two intervals $\underline{-i}$ and $\underline{-j}$ analogously. For example, $|(\underline{-i}, \underline{-j})|$ denotes the distance between the closest endpoints of these two intervals while $|[\underline{-i}, \underline{-j})|$ stands for $|\underline{-i}| + |(\underline{-i}, \underline{-j})|$.

3 Scaling near Critical Point

We define a sequence of scalings

$$\tau_n := \frac{|(\underline{0}, \underline{q}_n)|}{|(\underline{0}, \underline{q}_{n-2})|}.$$

These quantities measure ‘the geometry’ near to a critical point.

When $\tau_n \rightarrow 0$ we say that the geometry of the mapping is ‘degenerate’.

When τ_n is bounded away from zero we say that the geometry is ‘bounded’.

The following sequence will also be frequently used:

$$\alpha_n := \frac{|(-q_n, \underline{0})|}{|[-q_n, \underline{0}]|}.$$

Along with τ_n these ratios serve as scalings relating the geometries of successive dynamic partitions and moreover $\alpha_n > \tau_n$.

The proof of our results is based on the following theorem.

Theorem 3.1. *1. If the critical exponent $l \leq 2$, then the scalings τ_n tend to zero at least exponentially fast.*

2. For maps with rotation number of bounded type and with critical exponent $l > 2$ the sequence τ_n is bounded away from zero.

Proof. The second claim is completely proved in [3], while the first claim is proved under the additional assumption that the Schwarzian derivative is strictly smaller than zero. For the general case, the details are provided in the Appendix. \square

The next proposition will be used to prove the mains results of this paper.

Proposition 1. *Let f be a function in \mathcal{L} and let $J \subset T$ be two intervals of the circle.*

Suppose that, for some $n \in \mathbb{N}$

- *f^n is a diffeomorphism on T ;*
- *$\sum_{i=0}^{n-1} |f^i(J)|$ is bounded;*
- *$|f^n(J)| \leq K \text{dist}(f^n(J), \partial f^n(T))$ with K a positive constant.*

Then, there exists a constant C such that, for every two intervals A and B in J

$$\frac{|f^n(A)|}{|f^n(B)|} \geq C \frac{|A|}{|B|}.$$

Proof. It follows from the Koebe principle of [4]. □

3.1 Continued Fractions and Partitions

Since f is order-preserving and has no periodic points, there exists an order-preserving and continuous map $h : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ such that $h \circ f = R_\rho \circ h$, where ρ is the rotation number of f and R_ρ is the rigid rotation by ρ .

In particular, the order of points in an orbit of f is the same as the order of points in an orbit of R_ρ . Therefore, results about R_ρ can be translated into results about f , via the semiconjugacy h .

We can build the so called dynamical partitions \mathcal{P}_n of \mathbb{S}^1 to study the geometric properties of f , see [2].

\mathcal{P}_n is generated by the first $q_n + q_{n+1}$ preimages of U and consists of

$$\mathcal{I}_n := \{\underline{-i} : 0 \leq i \leq q_{n+1} + q_n - 1\},$$

together with the gaps between these intervals.

There are two kinds of gaps:

- The ‘long’ gaps are the interval I_0^n , which is the interval between $\underline{-q_n}$ and $\underline{0}$ for n even or the interval between $\underline{0}$ and $\underline{-q_n}$ for n odd, with its preimages,

$$I_i^n := f^{-i}(I_0^n), i = 0, 1, \dots, q_{n+1} - 1.$$

- The ‘short’ gaps are the interval I_0^{n+1} , which is the interval between $\underline{0}$ and $\underline{-q_{n+1}}$ for n even or the interval between $\underline{-q_{n+1}}$ and $\underline{0}$ for n odd, with its preimages,


$$I_i^{n+1} := f^{-i}(I_0^{n+1}), i = 0, 1, \dots, q_n - 1.$$

We will briefly explain the structure of the partitions.


Take two consecutive dynamical partitions of order n and $n + 1$. The latter is clearly a refinement of the former. All ‘short’ gaps of \mathcal{P}_n become ‘long’ gaps of \mathcal{P}_{n+1} while all ‘long’ gaps of \mathcal{P}_n split into a_{n+2} preimages of U and a_{n+2} ‘long’ gaps and one ‘short’ gap of the next partition \mathcal{P}_{n+1} :

$$I_i^n = \bigcup_{j=1}^{a_{n+2}} f^{-i-q_n-jq_{n+1}}(U) \cup \bigcup_{j=0}^{a_{n+2}-1} I_{i+q_n+jq_{n+1}}^{n+1} \cup I_i^{n+2}. \quad (3.2)$$

Several of the proofs in the following will depend strongly on the relative positions of the points and intervals of \mathcal{P}_n . In reading the proofs the reader is advised to keep the following pictures in mind, which show some of these objects near the flat interval $\underline{0}$.



In the next picture we have enlarged the right-hand part of this picture to show the location of the points $\underline{q_n}$, $\underline{2q_n}$ and $\underline{3q_n}$ for the case n even and $a_n = 1$.



Now we present a few results from [3] which are used throughout the paper:

Proposition 2. *Let $A \in \mathcal{I}_n$ be a preimage of U and B any of two gaps adjacent to A . Then there exists a constant C such that, for every $n \in \mathbb{N}$, $\frac{|A|}{|B|} \geq C$.*

Corollary 3.3. *The lengths of the gaps of the dynamical partition \mathcal{P}_n tend to zero at least exponentially fast with n .*

Lemma 3.4. *The ratio*

$$\frac{\left| \frac{-q_n + iq_{n-1}}{\left[\frac{-q_n + iq_{n-1}}{\mathbb{Q}} \right]} \right|}{\left| \frac{-q_n + iq_{n-1}}{\mathbb{Q}} \right|}$$

is bounded away from zero by a uniform constant for all $i = 0, \dots, a_n$.

For the proof of Proposition 2, Corollary 3.3 and Lemma 3.4 see [3].

Lemma 3.5. *For every $i \in \{0, \dots, a_n - 1\}$ the ratio*

$$\omega_i = \frac{\left| \left(\frac{-q_n + (i+1)q_{n-1}}{\mathbb{Q}}, \frac{-q_n + iq_{n-1}}{\mathbb{Q}} \right) \right|}{\left| \frac{-q_n + iq_{n-1}}{\mathbb{Q}} \right|}$$

is uniformly comparable to α_{n-1} .

Proof. For every $i \in \{1, \dots, a_n - 1\}$, we apply Proposition 1 to

- $T = \left[\frac{-q_n + (i+1)q_{n-1}}{\mathbb{Q}}, \frac{-q_n + (i-1)q_{n-1}}{\mathbb{Q}} \right]$,
- $J = \left(\frac{-q_n + (i+1)q_{n-1}}{\mathbb{Q}}, \frac{-q_n + (i-1)q_{n-1}}{\mathbb{Q}} \right)$
- $f^{q_n - (i+1)q_{n-1}}$.

Observe that the hypotheses are satisfied:

- T doesn't contain any preimage of the flat interval of the type f^{-j} with $j < q_n - (i+1)q_{n-1}$, so $f^{q_n - (i+1)q_{n-1}}$ is a diffeomorphism on T ;
- the set $\bigcup_{j=0}^{q_n - (i+1)q_{n-1} - 1} f^j(J)$ covers each point of \mathbb{S}^1 at most twice;
- if n is large enough, by Lemma 3.4 there exists a constant K such that $\left| f^{q_n - (i+1)q_{n-1}}(J) \right| = \left| \left(\mathbb{Q}, \frac{-2q_{n-1}}{\mathbb{Q}} \right) \right| \leq K \left| \frac{-2q_{n-1}}{\mathbb{Q}} \right|$.

So, by Proposition 1 and Proposition 2, ω_i is comparable to α_{n-1} .

For $i = 0$ we apply Proposition 1 to

- $T = \left[\frac{-q_n + q_{n-1} + 1}{\mathbb{Q}}, \frac{-q_n + 1}{\mathbb{Q}} \right]$,
- $J = \left(\frac{-q_n + q_{n-1} + 1}{\mathbb{Q}}, \frac{-q_n + 1}{\mathbb{Q}} \right)$

- $f^{q_n - q_{n-1} - 1}$.

As before, the hypotheses are satisfied (the intervals $f^j(J)$ for $j \in \{1, \dots, q_n - q_{n-1} - 2\}$ are disjoint).

So, for n large

$$\omega_0^l = \frac{\left| \left(\underline{-q_n + q_{n-1} + 1}, \underline{-q_n + 1} \right) \right|}{\left| \underline{-q_n + 1} \right|}$$

is uniformly comparable to α_{n-1} .

This concludes the proof. \square

4 Proof of Theorem 1.2

In the proof of Theorem 2 in [3] it is shown that, if the critical exponent of the map f is less than or equal to 2, then there exists $0 < \lambda < 1$ such that, for every n large enough

$$\alpha_n = \frac{\left| \left(\underline{-q_n}, \underline{0} \right) \right|}{\left| \underline{-q_n}, \underline{0} \right|} \leq \lambda^n. \quad (4.1)$$

Lemma 4.2. *There is a positive constant C such that each gap A^{n-2} of \mathcal{P}_{n-2} is split into at least two gaps A_i^n of \mathcal{P}_n whose lengths satisfy $|A_i^n| < C\lambda^{n-1}|A^{n-2}|$.*

Proof. Because of the symmetry, it is enough to suppose that n is even, that is $\underline{-q_n}$ is to the left of $\underline{0}$.

Let A^{n-2} be a long gap of \mathcal{P}_{n-2} ;

$$A^{n-2} = \left(\underline{-q_{n-2}}, \underline{0} \right).$$

As explained by the formula (3.2), A^{n-2} is subdivided into a_n long gaps and one short gap of the dynamical partition \mathcal{P}_{n-1} . In our case these gaps are

$$A_i^n = \left(\underline{-q_n + (i+1)q_{n-1}}, \underline{-q_n + iq_{n-1}} \right)$$

for $i \in \{0, \dots, a_n - 1\}$.

By Lemma 3.5 and the inequality (4.1), there exists a constant C such that, for every $i \in \{0, \dots, a_n - 1\}$

$$\frac{|A_i^n|}{|A^{n-2}|} < \frac{\left| \left(\underline{-q_n + (i+1)q_{n-1}}, \underline{-q_n + iq_{n-1}} \right) \right|}{\left| \underline{-q_n + iq_{n-1}} \right|} \leq C\alpha_{n-1} \leq C\lambda^{n-1}.$$

Since the shorts gaps of \mathcal{P}_{n-2} become the longs gaps of the partition \mathcal{P}_{n-1} , we can use the estimates for long gaps of the partition \mathcal{P}_{n-1} to conclude the proof. \square

Proof of Theorem 1.2 Observe that $\Omega_{n-2} = \bigcup_{A^{n-2} \in \mathcal{P}_{n-2}} A^{n-2}$ and $\Omega_n = \bigcup_{A^n \in \mathcal{P}_n} A^n$ are two covers of K . We analyze the relations between them. By Lemma 4.2 and the formula (3.2), we can deduce that:

1. Each interval A^{n-2} of Ω_{n-2} is split into at least two intervals A_i^n of Ω_n whose lengths satisfy $|A_i^n| < C\lambda^{n-1}|A^{n-2}|$.
2. Each interval A^{n-2} is split into at most $a_n(a_{n+1} + 1) + 1$ sub-intervals.

Consequently,

$$\sum_{A^n \in \mathcal{P}_n} |A^n|^s \leq C^s (a_n(a_{n+1} + 1) + 1) \lambda^{(n-1)s} \sum_{A^{n-2} \in \mathcal{P}_{n-2}} |A^{n-2}|^s.$$

Since the maximum diameter of the covering $\{A^n\}$ goes to zero as n increases, the Hausdorff s -measure of K must be zero for every $s > 0$. This implies the theorem.

5 Proof of Theorem 1.3

In this section we will assume that the critical exponent (l, l) of f is strictly greater than 2.

Lemma 5.1. *Two adjacent gaps of the dynamical partition \mathcal{P}_n which are contained in the same gap of \mathcal{P}_{n-1} are comparable.*

Proof. Because of the symmetry, without loss of generality, we may assume that n is even; so $\underline{-q_n}$ is to the left of the flat interval.

For every $i \in \{0, \dots, a_{n+1} - 1\}$ let

$$A_i = \underline{-q_{n+1} + iq_n}$$

and let

$$B_i = (\underline{-q_{n+1} + iq_n}, \underline{-q_{n+1} + (i+1)q_n}).$$

By Lemma 3.5, there exists a constant C such that, for every $i \in \{0, \dots, a_{n+1} - 1\}$

$$\frac{|B_i|}{|A_i|} \geq C\alpha_n > C\tau_n \tag{5.2}$$

is bounded away from zero by Theorem 3.1.

So, by the formula (5.2) and Proposition 2, there exists a constant $K > 0$ such that

$$\frac{|B_{i-1}|}{|B_i|} \leq K.$$

In order to compare B_{i-1} and A_i , we observe that, for every $i \in \{1, \dots, a_{n+1} - 1\}$

$$\frac{|A_i|}{|\underline{-q_{n+1}}|} \leq K_1 \frac{|A_i|}{\left| \left(\underline{0}, \underline{-q_{n+1}} \right) \right|} \leq K_1 \frac{|(\underline{0}, A_i)|}{\left| \left(\underline{0}, \underline{-q_{n+1}} \right) \right|} \leq K_1 \frac{\left| \left(\underline{0}, \underline{-q_{n+1}} \right) \right|}{\left| \left(\underline{0}, \underline{-q_{n+1}} \right) \right|} \leq K_2 \quad (5.3)$$

where the first inequality follows from the Proposition 2 and the last one from Theorem 3.1.

Therefore, using the inequality (5.2), Proposition 2 and the inequality (5.3), we have that

$$\frac{|B_{i-1}|}{|A_i|} \geq C \frac{|A_{i-1}|}{|A_i|} \geq C_1 \frac{|B_{i-2}|}{|A_i|} \geq \dots \geq C_i \frac{|\underline{-q_{n+1}}|}{|A_i|} \geq \frac{C_i}{K_2}. \quad (5.4)$$

Moreover, $\left| \left(\underline{0}, \underline{-q_n} \right) \right|$ is bigger than a uniform constant multiple of $|\underline{-q_n}|$ as the sequence τ_n is bounded away from zero for $l > 2$.

Finally, by the formula (5.4) and Proposition 2, there exists a constant K_1 such that

$$\frac{|B_{i-1}|}{|B_i|} \geq K_1.$$

So, B_{i-1} and B_i are comparable. □

Lemma 5.5. *There exists a constant C such that*

$$\left| \left(\underline{q_n}, \underline{-q_{n+1} + q_n} \right) \right| \geq C \left| \left(\underline{-q_{n+1}}, \underline{-q_{n+1} + q_n} \right) \right|$$

Proof. Apply Proposition 1 to

- $T = \left[\underline{-q_{n+1}}, \underline{-q_{n+1} + q_n} \right]$
- $J = \left(\underline{-q_{n+1}}, \underline{-q_{n+1} + q_n} \right)$
- $f^{q_{n+1} - q_n}$.

Also in this case the hypotheses are satisfied:

- $f^{q_{n+1}-q_n}$ is a diffeomorphism on T ;
- the intervals $f^i(J)$ for $i \in \{1, \dots, q_{n+1} - q_n - 1\}$ are disjoint;
- for n large enough, by Proposition 2 there exists a constant K such that

$$|f^{q_{n+1}-q_n}(J)| = |(\underline{-q_n}, \underline{0})| \leq K |\underline{-q_n}| = K \text{dist}(f^{q_{n+1}-q_n}(J), \partial f^{q_{n+1}-q_n}(T)).$$

So, there exists a constant C such that


$$\frac{\left| \left(\underline{q_n}, \underline{-q_{n+1} + q_n} \right) \right|}{\left| \left(\underline{-q_{n+1}}, \underline{-q_{n+1} + q_n} \right) \right|} \geq C \frac{\left| \left(\underline{q_{n+1}}, \underline{0} \right) \right|}{\left| \left(\underline{-q_n}, \underline{0} \right) \right|} > C \frac{\left| \left(\underline{-q_{n+2}}, \underline{0} \right) \right|}{\left| \left(\underline{-q_n}, \underline{0} \right) \right|}$$

which is bounded away from zero by Theorem 3.1. □

Lemma 5.6. *Two gaps of the same dynamical partition which are adjacent to the flat interval are comparable.*

Proof. Because of the symmetry, it is enough to suppose that n is even.

Let $A = (\underline{-q_n}, \underline{0})$ and let $B = (\underline{0}, \underline{-q_{n+1}})$; the initial situation is explained in the following figure.



The idea is to apply Proposition 1 to

- $T = [\underline{-q_n + 1}, \underline{-q_{n+1} + 1}]$
- $J = (\underline{-q_n + 1}, \underline{-q_{n+1} + 1})$
- f^{q_n-1} .

So, there exists a constant C such that

$$\frac{|f(B)|}{|f(A)|} \geq C \frac{\left| \left(\underline{q_n}, \underline{-q_{n+1} + q_n} \right) \right|}{\left| \left(\underline{0}, \underline{q_n} \right) \right|} \geq C \frac{\left| \left(\underline{q_n}, \underline{-q_{n+1} + q_n} \right) \right|}{\left| \left(\underline{0}, \underline{-q_{n-1}} \right) \right|}. \quad (5.7)$$

By Theorem 3.1 and by Lemma 5.5 there exist two constants K and K_1 such that

$$\frac{\left| \left(\underline{q_n}, \underline{-q_{n+1} + q_n} \right) \right|}{\left| \left(\underline{0}, \underline{-q_{n-1}} \right) \right|} \geq K \frac{\left| \left(\underline{q_n}, \underline{-q_{n+1} + q_n} \right) \right|}{\left| \left(\underline{0}, \underline{-q_{n+1}} \right) \right|} \geq K_1 K \frac{\left| \left(\underline{-q_{n+1}}, \underline{-q_{n+1} + q_n} \right) \right|}{\left| \left(\underline{0}, \underline{-q_{n+1}} \right) \right|}.$$

Observe that the last intervals are contained in the same gap of the partition n ; so by Lemma 5.1 they are comparable.

Similarly, there exists $C > 0$

$$\frac{|f(A)|}{|f(B)|} \geq C \frac{|\underline{(0, q_n)}|}{\left| \left(\underline{q_n}, \underline{-q_{n+1} + q_n} \right) \right|} \geq C \frac{|\underline{(0, q_n)}|}{\left| \left(\underline{0}, \underline{q_{n-2}} \right) \right|} \geq C\tau_n. \quad (5.8)$$

which is bounded away from zero by Theorem 2 in [3].


So, $f(A)$ and $f(B)$ are comparable.

Moreover, we recall that near to the flat interval $f(x) = x^l$ hence also A and B are comparable. □

Proposition 3. *Two adjacent gaps of the dynamical partition \mathcal{P}_n are comparable.*

Proof. By the symmetry, without loss of generality, we can suppose that n is even.

Case 1. For $0 \leq i < q_n$ we have



Let $A = (\underline{-i - q_n}, \underline{-i})$ and let $B = (\underline{-i}, \underline{-i - q_{n+1}})$.

For $i = 0$ we find the two gaps adjacent to the flat interval which are uniformly comparable, see Lemma 5.6.

For $i \neq 0$ apply Proposition 1 to

- $T = \left[\underline{-i - q_n}, \underline{-i - q_{n+1}} \right],$
- $J = \left(\underline{-i - q_n}, \underline{-i - q_{n+1}} \right),$

- f^i .

Hence, by Proposition 1 and the inequality (5.7) there exist two constants C and C_1 such that

$$\frac{|A|}{|B|} \leq C \frac{|(-q_n + 1, \underline{1})|}{|(\underline{1}, -q_{n+1} + 1)|} \leq CC_1.$$

Similarly, Proposition 1 and the inequality (5.8) imply that

$$\frac{|B|}{|A|} \leq C \frac{|(\underline{1}, -q_{n+1} + 1)|}{|(-q_n + 1, \underline{1})|} \leq CC_2.$$

where C and C_2 are uniform constants.

Case 2. For $q_n \leq i < q_{n+1}$ we have

$$\begin{array}{ccc} \underline{-i - q_n} & \underline{-i} & \underline{-i + q_n} \\ \downarrow & \downarrow & \downarrow \\ \hline \end{array}$$

Let $A = (\underline{-i - q_n}, \underline{-i})$ and let $B = (\underline{-i}, \underline{-i + q_n})$.

We apply Proposition 1 to

- $T = [\underline{-i - q_n}, \underline{-i + q_n}]$,
- $J = (\underline{-i - q_n}, \underline{-i + q_n})$,
- $f^{i - q_n}$.

There exists a constant C such that


$$\frac{|A|}{|B|} \leq C \frac{|(-2q_n, -q_n)|}{|(-q_n, \underline{0})|} \leq C \frac{|(-q_{n-2}, \underline{0})|}{|(-q_n, \underline{0})|} \leq CC_1.$$

C_1 is a constant that comes from Theorem 3.1.

Moreover by Lemma 5.1 and Lemma 3.4 there exist two constants K_1 and K_2 such that

$$\frac{|B|}{|A|} \leq C \frac{|(-q_n, \underline{0})|}{|(-2q_n, -q_n)|} \leq CK_1 \frac{|(-q_n, \underline{0})|}{| -2q_n |} \leq CK_1 \frac{|[-2q_n, \underline{0}]|}{| -2q_n |} \leq CK_1 K_2.$$

Case 3. For $q_{n+1} \leq i < q_n + q_{n+1}$ we have



Let $A = (\underline{-i + q_{n+1}}, \underline{-i})$ and let $B = (\underline{-i}, \underline{-i + q_n})$.

We apply Proposition 1 to

- $T = [\underline{-i + q_{n+1}}, \underline{-i + q_n}]$,
- $J = (\underline{-i + q_{n+1}}, \underline{-i + q_n})$,
- $f^{i - q_{n+1}}$.

There exists a constant C such that

$$\frac{|A|}{|B|} \leq C \frac{\left| (\underline{0}, \underline{-q_{n+1}}) \right|}{\left| (\underline{-q_{n+1}}, \underline{-q_{n+1} + q_n}) \right|}.$$

Since $(\underline{0}, \underline{-q_{n+1}})$ and $(\underline{-q_{n+1}}, \underline{-q_{n+1} + q_n})$ are contained in the same gap of the partition $n - 1$, we can use Lemma 5.1 to obtain a uniform upper bound.

Similarly, by Proposition 1 and Lemma 5.1, there exist two constants C and C_1 such that

$$\frac{|B|}{|A|} \leq C \frac{\left| (\underline{-q_{n+1}}, \underline{-q_{n+1} + q_n}) \right|}{\left| (\underline{0}, \underline{-q_{n+1}}) \right|} \leq CC_1.$$

All the possibilities have been analysed, so the proof is complete. \square

Proof. Proof of Theorem 1.3.

Let \mathcal{A}_n be the algebra generated by the set of all gaps belonging to the partition \mathcal{P}_n . We define a probability measure μ on \mathcal{A}_n .

For every gap $I \in \mathcal{P}_1$ set $\mu(I) = \frac{1}{\#\mathcal{P}_1}$. Suppose μ is already constructed on

\mathcal{A}_n and we fix μ on \mathcal{A}_{n+1} .

If I_i^n is a long gap of \mathcal{P}_n then

$$I_i^n = \bigcup_{j=1}^{a_{n+2}} f^{-i-q_n-jq_{n+1}}(U) \cup \bigcup_{j=0}^{a_{n+2}-1} I_{i+q_n+jq_{n+1}}^{n+1} \cup I_i^{n+2}$$

and we set, for every $j \in \{0, \dots, a_{n+2} - 1\}$,

$$\mu(I_{i+q_n+jq_{n+1}}^{n+1}) = \frac{\mu(I_i^n)}{2a_{n+2}}$$

and

$$\mu(I_i^{n+2}) = \frac{\mu(I_i^n)}{2}.$$

If I_i^{n+1} is a short gap of \mathcal{P}_n then $\mu(I_i^{n+1})$ is already defined.

By Carathéodory's extension theorem, there exists a measure (that we will call again μ) which extends μ on $\sigma(\mathcal{A}_1, \mathcal{A}_2, \dots)$.

By the definition, $\mu(I_i^n) \leq \frac{1}{2^{\frac{1}{2}}}$. Proposition 3 and Corollary 3.3 imply that there exist $\lambda_1, \lambda_2 \in (0, 1)$ such that all gaps satisfy $\lambda_1^n \leq |I_i^n| \leq \lambda_2^n$. Therefore,

$$\mu(I_i^n) \leq |I_i^n|^\alpha$$

with $\alpha = \log_{\lambda_1} \frac{1}{\sqrt{2}}$.

Let I be an arbitrary interval and let I_i^{n+1} be a gap contained in I with n as small as possible. Then I is covered by at most two gaps of the n^{th} partition I_j^n and $I_{j'}^n$ and by preimages of U (which are of μ -measure zero). By the definition of μ

$$\mu(I) \leq \mu(I_j^n) + \mu(I_{j'}^n) \leq C\mu(I_j^n) \leq C'\mu(I_i^{n+1}) \leq C'|I_i^{n+1}|^\alpha \leq C'|I|^\alpha \quad (5.9)$$

Finally, let \mathcal{K} be an ϵ -cover of the non-wandering set K and let $0 < \alpha < 1$.

By the inequality (5.9),

$$\sum_{I \in \mathcal{K}} |I|^\alpha \geq \frac{1}{C'} \sum_{I \in \mathcal{K}} \nu(I) \geq \frac{1}{C'} \nu(K) = \frac{1}{C'}.$$

This establishes the theorem. □

6 Applications: Cherry Flows

6.1 Basic Definitions

Let X be a C^∞ vector field on the torus T^2 . Denote the flow through a point x by $t \rightarrow X_t(x)$ and by $Sing(X)$ the set of singularities of X . The positive and negative *semi-trajectories* are the sets

$$l^+(x) = \{X_t(x) : t \geq 0\}$$

$$l^-(x) = \{X_t(x) : t \leq 0\}.$$

The set $l(x) = l^+(x) \cup l^-(x)$ is called the *trajectory* through the point x . If $l(x) = x$ then x is called a *fixed point* or a *singularity*. If $x \in T^2$ is not a fixed point then it is called a *regular point* and $l(x)$ will be called a *one-dimensional trajectory*. If $l(x)$ is homeomorphic to a circle \mathbb{S}^1 , then $l(x)$ is a *closed trajectory* or a *periodic trajectory*. The trajectory $l(x)$ is called a *non-closed trajectory* if $l(x)$ is neither a fixed point nor a periodic trajectory. The ω -*limit set* of the positive semi-trajectory $l^+(x)$ is the set

$$\omega[l^+(x)] = \{y; \exists t_n \rightarrow \infty \text{ with } X_{t_n}(x) \rightarrow y\},$$

and α -*limit set* of the negative semi-trajectory $l^-(x)$ is

$$\alpha[l^-(x)] = \{y; \exists t_n \rightarrow \infty \text{ with } X_{-t_n}(x) \rightarrow y\}.$$

The ω -limit set (α -limit set) of any positive (negative) semi-trajectory of the trajectory l is called ω -limit set $\omega(l)$ of l (α -limit set $\alpha(l)$ of l). The union of ω -limit sets of all trajectories of the flow X_t is called the ω -limit set $\omega(X_t)$ of X_t (analogously for the α -limit set $\alpha(X_t)$ of X_t). The union $\lim(X_t) = \omega(X_t) \cup \alpha(X_t)$ is said to be the *limit set* of X_t .

The trajectory is ω -recurrent (α -recurrent), if it is contained in its ω -limit set (α -limit set). The trajectory is *recurrent* if it is both ω -recurrent and α -recurrent. A recurrent trajectory is *non-trivial* if it is neither a fixed point nor a periodic trajectory.

Let S be a subset of T^2 . We denote by $W^s(S)$ the set of points in T^2 that have S as ω -limit (it is called the *stable manifold* of S) and by $W^u(S)$ the set of points that have S as α -limit (it is called the *unstable manifold* of S).

Definition 6.1. A *Cherry field* is a C^∞ vector field on the torus T^2 without closed trajectories which has exactly two singularities, a sink and a saddle, both hyperbolic.

The first example of such a field was given by Cherry in 1938.

Definition 6.2. *The Poincaré global section is a transversal, simple, closed C^∞ curve Σ which intersects every one-dimensional trajectory of the flow.*

Fact 6.3. *It is proved in [9] that every Cherry flow admits a Poincaré global section.*

Definition 6.4. *Let l be a non-trivial recurrent trajectory. Then the closure \bar{l} of l is called a quasi-minimal set.*

The following fact was proved in [9].

Fact 6.5. *Every Cherry field has a quasi-minimal set which is locally homeomorphic to the Cartesian product of a Cantor set and a segment.*

6.2 First Return Map

Let X be a Cherry field with a global Poincaré section Σ . Notice that $T^2 \setminus \Sigma$ is C^∞ -equivalent to an annulus $\mathbb{S}^1 \times (0, 1)$ and we can write $T^2 \cong \mathbb{S}^1 \times [0, 1] / \sim$, where $(s, 0) \sim (s, 1)$. Consider X as a flow on $T^2 \cong \mathbb{S}^1 \times [0, 1]$ where we identify $\mathbb{S}^1 \times \{0\}$ and $\mathbb{S}^1 \times \{1\}$.

After this change of coordinates, the resulting field is a Cherry field.

We denote by W the set of points $x \in \mathbb{S}^1 \times \{0\}$ with $X_t(x) \in \mathbb{S}^1 \times \{1\}$ for some $t > 0$. For $x \in W$, let $t(x)$ be the minimal $t > 0$ such that $X_{t(x)}(x) \in \mathbb{S}^1 \times \{1\}$ and define the map $f : W \rightarrow \mathbb{S}^1$ by $f(x) = X_{t(x)}(x)$. This map is called the *first return map* to \mathbb{S}^1 .

We define f on $U = (\mathbb{S}^1 \setminus W)$ as $f(U) = v$ where $v = W^u(S) \cap \mathbb{S}^1 \times \{1\}$.

For every point $x \in (\mathbb{S}^1 \setminus U) \times \{0\}$ near ∂U , there exists $t > 0$ such that $X_t(x)$ intersects $\mathbb{S}^1 \times \{1\}$ near v (for more details see the section 2 of the chapter 7 in [1]). In particular $\lim_{x \rightarrow \partial U, x \notin U} f(x)$ consists of a single point.

So, we have a function f of the circle, which is everywhere continuous and C^∞ outside the boundary points of $\mathbb{S}^1 \setminus U$.

6.3 Properties of the First Return Map

Let X be a Cherry vector field, with a saddle point that has eigenvalues $\lambda_1 > 0 > \lambda_2$. Let \mathbb{S}^1 be a global Poincaré section and let f be the first return map to \mathbb{S}^1 .

Then:

1. f is order preserving.
2. It is constant on an interval U .

3. The restriction of f to $\mathbb{S}^1 \setminus \overline{U}$ is a C^∞ -diffeomorphism onto its image.

4. Let $\pi(a, b) = U$. On some right-sided neighborhood of b , f can be represented as

$$h_r((x - b)^\alpha),$$

where $\alpha = \frac{|\lambda_2|}{\lambda_1}$ and h_r is a C^∞ -diffeomorphism on a neighborhood of b . Analogously, on a left-sided neighborhood of a , f is

$$h_l((a - x)^\alpha).$$

5. Since X has no periodic trajectories, the rotation number of f , $\rho(f)$, is irrational.

These are well-known properties of f , the reader can consult for example [1] or [9].

The first return map f belongs to the class \mathcal{L} from Section 1.1.

Lemma 6.6. *There exists a Cherry vector field with first return map which has rotation number of bounded type.*

Proof. Let X be a Cherry vector field with a global Poincaré section \mathbb{S}^1 . For every $\epsilon > 0$, we define Y_ϵ as a constant horizontal vector field with all vectors of the same length ϵ . Let $X_\epsilon = X + Y_\epsilon$ and denote by f_ϵ the first return map of X_ϵ to \mathbb{S}^1 .

The family (f_ϵ) is increasing and hence $\rho_\epsilon = \rho(f_\epsilon)$ is an increasing and continuous function. Since ρ_0 is irrational, ρ_ϵ is strictly increasing at zero and hence, there is a ϵ_0 so that ρ_{ϵ_0} is an irrational number of bounded type. \square

6.4 Proof of Theorem 1.4

Proof. We denote by $\dim_{\mathbb{H}}(M)$ the Hausdorff dimension of a set M and by Q the quasi-minimal set of X . By Fact 6.5, in a small neighborhood of the Poincaré global section, Q is equivalent, by a C^2 diffeomorphism, to $I \times \Omega$ with Ω the non-wandering set.

So, by Theorem 1.3 and the product formula, Theorem 8.10 in [6],

$$\dim_{\mathbb{H}}(Q) \geq \dim_{\mathbb{H}}(I \times \Omega) \geq \dim_{\mathbb{H}}(I) + \dim_{\mathbb{H}}(\Omega) > 1.$$

\square

A Appendix

In the following we explain how to remove the hypothesis on the Schwarzian derivative in the first claim of Theorem 2 in [3].

The negative Schwarzian is used only to prove the expansion property of the cross-ratio **Poin** which is defined on ordered quadruples $a < b < c < d$ by

$$\mathbf{Poin}(a, b, c, d) := \frac{|d - a| |b - c|}{|c - a| |d - b|}.$$

Diffeomorphisms with negative Schwarzian derivative increase the cross-ratio **Poin**:

$$\mathbf{Poin}(f(a), f(b), f(c), f(d)) > \mathbf{Poin}(a, b, c, d).$$

In general, without the assumption of negative Schwarzian, the following holds.

Theorem A.1. *Let f be a C^2 map with no flat critical points. There exists a bounded increasing function $\sigma : [0, \infty) \rightarrow \mathbb{R}_+$ with $\sigma(t) \rightarrow 0$ as $t \rightarrow 0$ with the following property. Let $[b, c] \subset [a, d]$ be intervals such that $f^n|_{[a, d]}$ is a diffeomorphism. Then*

$$\mathbf{Poin}(f^n(a), f^n(b), f^n(c), f^n(d)) \geq \exp\{-\sigma(\tau) \sum_{i=0}^{n-1} |f^i([a, b])|\} \mathbf{Poin}(a, b, c, d),$$

where $\tau = \max_{i=0, \dots, n-1} |f^i((c, d])|$.

We recall the notation used in [3], let

$$\alpha_n = \frac{|(-\underline{q}_n, \underline{0})|}{|[-\underline{q}_n, \underline{0})|},$$

$$\sigma_n = \frac{|(\underline{0}, \underline{q}_n)|}{|(\underline{0}, \underline{q}_{n-1})|}$$

and

$$s_n = \frac{|[-\underline{q}_{n-2}, \underline{0})|}{|\underline{0}|}.$$

Theorem A.2. For n sufficiently large the following inequality holds:

$$(\alpha_n)^l \leq K_{0,n} K_{1,n}, \dots, K_{a_n-1,n} C_n \tilde{M}_n(l) \alpha_{n-2}^2, \quad (\text{A.3})$$

where for every $i \in 0, \dots, a_n - 1$ if we denote by

$$\tau_{i,n} = \max_{j=0, \dots, q_{n-1}-2} \left| f^j \left(\left(\underline{iq_{n-1} + 1}, \underline{-q_{n-1} + 1} \right) \right) \right|$$

and by

$$\rho_n = \max_{j=0, \dots, q_{n-2}-2} \left| f^j \left(\left(\underline{a_n q_{n-1} + 1}, \underline{1} \right) \right) \right|$$

then

$$\begin{aligned} K_{i,n} &= e^{\sigma(\tau_{i,n}) \sum_{j=0}^{q_{n-1}-2} |f^j(\underline{-q_n + iq_{n-1} + 1})|}, \\ C_n &= e^{\sigma(\rho_n) \sum_{j=0}^{q_{n-2}-2} |f^j(\underline{-q_{n-2} + 1})|} \end{aligned} \quad (\text{A.4})$$

and

$$\tilde{M}_n(l) = s_{n-1}^2 \cdot \frac{2}{l} \cdot \left(\frac{1}{1 + \sqrt{1 - \frac{2(l-1)}{l} C_n s_{n-1} \alpha_{n-1}}} \right) \cdot \frac{1}{1 - \alpha_{n-2}} \cdot \frac{\sigma_n}{\sigma_{n-2}}.$$

Proof. For large n ,

$$\alpha_n^l = \frac{|\underline{(-q_n + 1, \underline{1})}|}{|\underline{[-q_n + 1, \underline{1}]}|},$$

which is certainly less than

$$\mathbf{Poin} \left(\underline{-q_n + 1}, \underline{(\underline{1}, -q_{n-1} + 1)} \right).$$

By Theorem A.1 there exists a constant

$$K_{0,n} = e^{\sigma(\max_{j=0, \dots, q_{n-1}-2} |f^j(\underline{(\underline{1}, -q_{n-1} + 1)})|) \sum_{j=0}^{q_{n-1}-2} |f^j(\underline{-q_n + 1})|}$$

such that

$$\alpha_n^l \leq K_{0,n} \delta_n(1) \cdot s_n(1), \quad (\text{A.5})$$

with

$$\delta_n(k) = \frac{\left| \underline{(-q_n + kq_{n-1}, kq_{n-1})} \right|}{\left| \underline{[-q_n + kq_{n-1}, kq_{n-1}]} \right|}$$

and

$$s_n(k) = \frac{\left| \underline{[-q_n + kq_{n-1}, \underline{0}]} \right|}{\left| \underline{(-q_n + kq_{n-1}, \underline{0})} \right|}.$$

By the Mean Value Theorem, f transforms the intervals defining $\delta_n(k)$ into a pair whose ratio is

$$\left(\frac{u_k}{v_k}\right) \delta_n(k)$$

where u_k is the derivative of x^l at a point in the interval U_k between $\underline{-q_n + kq_{n-1}}$ and $\underline{kq_{n-1}}$, while v_k is the derivative at a point in

$$V_k = U_k \cup \underline{-q_n + kq_{n-1}}.$$

Observe that for n sufficiently large,

$$u_1 < v_1 < u_2 < v_2 < \cdots < u_{a_n} < v_{a_n}.$$

Moreover

$$f(\delta_n(k)) \leq \mathbf{Poin} \left(\underline{-q_n + kq_{n-1} + 1}, \underline{(kq_{n-1} + 1, -q_{n-1} + 1]} \right)$$

which by Theorem A.1 is less than

$$K_{k,n} \mathbf{Poin} \left(\underline{-q_n + (k+1)q_{n-1}}, \underline{((k+1)q_{n-1}, \underline{0}]} \right).$$

So, the sequence of inequalities above can be rewritten in the form

$$\frac{u_k}{v_k} \delta_n(k) \leq K_{k,n} \cdot s_n(k+1) \cdot \delta_n(k+1). \quad (\text{A.6})$$

Now, starting from inequality (A.5) and applying $a_n - 1$ times (A.6) we obtain

$$(\alpha_n)^l \leq K_{0,n} \cdots K_{a_n-1,n} \delta_n(a_n) \cdot \frac{v_{a_n-1}}{u_1} \cdot s_n(1) \cdots s_n(a_n).$$

Note that $s_n(1) \cdots s_n(a_n) \leq s_n$ and that

$$\frac{v_{a_n-1}}{u_1} \leq \left(\frac{\left| \underline{(-q_{n-2}, \underline{0})} \right|}{\left| \underline{(q_{n-1}, \underline{0})} \right|} \right)^{l-1} \leq \frac{\left| \underline{(-q_{n-2}, \underline{0})} \right|}{\left| \underline{(q_{n-1}, \underline{0})} \right|}$$

hence

$$(\alpha_n)^l \leq K_{0,n} \cdots K_{a_n-1,n} \delta_n(a_n) \cdot \frac{\left| \underline{(-q_{n-2}, \underline{0})} \right|}{\left| \underline{(q_{n-1}, \underline{0})} \right|} \cdot s_n$$

which can be rewritten in the form

$$(\alpha_n)^l \leq K_{0,n} \cdots K_{a_n-1,n} s_n \nu_{n-2} \mu_{n-2} \alpha_{n-2}, \quad (\text{A.7})$$

where

$$\nu_{n-2} = \frac{\left| \left[\frac{-q_{n-2}}{\underline{0}} \right] \right|}{\left| \left(\frac{q_{n-1}}{\underline{0}} \right) \right|} \cdot \frac{\left| \left[\frac{-q_{n-2}}{\underline{0}} \right] \right|}{\left| \left[\frac{-q_{n-2}}{\underline{a_n q_{n-1}}} \right] \right|},$$

and

$$\mu_{n-2} = \frac{\left| \left(\frac{-q_{n-2}}{\underline{a_n q_{n-1}}} \right) \right|}{\left| \left(\frac{-q_{n-2}}{\underline{0}} \right) \right|}.$$

To conclude the proof we have to estimate ν_{n-2} and μ_{n-2} . For ν_{n-2} , observe that

$$\left| \left[\frac{-q_{n-2}}{\underline{0}} \right] \right| \leq \left| \left(\frac{q_{n-3}}{\underline{0}} \right) \right|,$$

so that

$$\nu_{n-2} \leq \frac{1}{\sigma_{n-1} \sigma_{n-2}} \cdot \frac{1}{1 - \alpha_{n-2}}. \quad (\text{A.8})$$

To estimate μ_{n-2} , we use an elementary lemma (Lemma 3.1 in [3]).

Lemma A.9. *For any pair of numbers $x > y$ the inequality*

$$\frac{x^l - y^l}{x^l} \geq \left(\frac{x - y}{x} \right) \left(l - \frac{l(l-1)}{2} \left(\frac{x - y}{x} \right) \right)$$

holds.

Now, apply f to the intervals defining the ratio μ_{n-2} . By the previous lemma, the resulting ratio is larger than

$$\mu_{n-2} \left(l - \frac{l(l-1)}{2} \mu_{n-2} \right).$$

The cross-ratio

$$\mathbf{Poin} \left(\frac{-q_{n-2} + 1}{\underline{a_n q_{n-1} + 1}}, \frac{1}{\underline{1}} \right)$$

is larger again; so, by Theorem A.1, there exists a constant C_n such that

$$\mu_{n-2} \left(l - \frac{l(l-1)}{2} \mu_{n-2} \right) \leq C_n \cdot s_{n-1} \cdot \sigma_n \sigma_{n-1}. \quad (\text{A.10})$$

Solving this quadratic inequality we have that

$$\mu_{n-2} \leq \frac{2}{l} \cdot \left(\frac{1}{1 + \sqrt{1 - \frac{2(l-1)}{l} C_n s_{n-1} \sigma_n \sigma_{n-1}}} \right) \cdot C_n \cdot s_{n-1} \cdot \frac{\sigma_n}{\sigma_{n-2}}. \quad (\text{A.11})$$

Combining (A.7), (A.8), (A.10) and (A.11) we conclude the proof. \square

A.1 Convergence of the Sequence α_n

The convergence of $\prod_{j=0}^n K_{i,j}$ and $\prod_{j=0}^n C_j$ is assured by the following lemma:

Lemma A.12. *There exists $0 < \lambda < 1$ such that for every $i \in \{0, \dots, a_n - 1\}$ and for every n large enough, $K_{i,n} \leq e^{\sigma(\lambda^{n-2})\lambda^{n-2}}$.*

Proof. Observe that, for every $j \in \{0, \dots, q_{n-1} - 2\}$

1. the intervals $f^j(\underline{-q_n + iq_{n-1} + 1})$ are disjoint and $\sum_{j=0}^{q_{n-1}-2} \left| f^j(\underline{-q_n + iq_{n-1} + 1}) \right|$ is contained in a gap of the partition \mathcal{P}_{n-1} ,
2. every interval $f^j(\underline{(iq_{n-1} + 1, -q_{n-1} + 1]})$ is contained in a gap of the partition \mathcal{P}_{n-2} .

So, by Corollary 3.3, for large n

$$\sum_{j=0}^{q_{n-1}-2} \left| f^j(\underline{-q_n + iq_{n-1} + 1}) \right| < \lambda^{n-2}$$

and

$$\sigma \left(\max_{j=0, \dots, q_{n-1}-2} \left| f^j \left(\underline{(iq_{n-1} + 1, -q_{n-1} + 1]} \right) \right| \right) < \sigma(\lambda^{n-2}).$$

□

A similar argument shows that also $C_n \leq e^{\sigma(\lambda^{n-2})\lambda^{n-2}}$ for large n .

To establish the convergence of the sequence α_n we have to prove that $\prod_{i=0}^n \tilde{M}_i$ tends to zero.

Observe that $\tilde{M}_n(l) = M_n(l) T_n(l)$ with

$$M_n(l) = s_{n-1}^2 \cdot \frac{2}{l} \cdot \left(\frac{1}{1 + \sqrt{1 - \frac{2(l-1)}{l} s_{n-1} \alpha_{n-1}}} \right) \cdot \frac{1}{1 - \alpha_{n-2}} \cdot \frac{\sigma_n}{\sigma_{n-2}}$$

and

$$T_n(l) = \frac{1 + \sqrt{1 - \frac{2(l-1)}{l} s_{n-1} \alpha_{n-1}}}{1 + \sqrt{1 - \frac{2(l-1)}{l} C_n s_{n-1} \alpha_{n-1}}}.$$

The proof that $\prod_{j=0}^n M_j$ tends to zero is exactly the same as in [3]. It remains to determine the size of $T_n(l)$.

Noting that

$$T_n(l) \leq \max \left(1, \frac{\sqrt{1 - \frac{2(l-1)}{l} s_{n-1} \alpha_{n-1}}}{\sqrt{1 - \frac{2(l-1)}{l} C_n s_{n-1} \alpha_{n-1}}} \right)$$

it all comes down to the study of the function

$$f(x) = \frac{1 - x}{1 - C_n x}$$

for $C_n > 1$ tending to one and $x = \frac{2(l-1)}{l} s_{n-1} \alpha_{n-1}$.

Observe that $f'(x) > 0$ on $(0, c_n^{-1})$ and that by the a priori estimates of [3], $x \leq 0.55$.

Therefore,

$$T_n(l) \leq 1 + 1.3(C_n - 1).$$

So, by Lemma A.12, $\prod_{i=0}^n T_i$ is bounded.

So, for $l \leq 2$ the recursive inequality A.3 implies that the sequence α_n goes to zero at least exponentially fast. Since $\alpha_n > \tau_n$ the same holds also for the scalings τ_n .

Acknowledgments

I would sincerely thank Prof. J. Graczyk for introducing me to the subject of this paper, his valuable advice and continuous encouragement. I am also very grateful to Prof. G. Świątek for many helpful discussions.

References

- [1] S. Kh. Aranson, G. R. Belitskii, E.V. Zhuzhoma: *Introduction to the Qualitative Theory of Dynamical Systems on Surfaces*. American Mathematical Society, (1996)
- [2] J. Graczyk: *Dynamics of Circle Maps with Flat Spots*. *Fundamenta Mathematicae* **209** (2010)
- [3] J. Graczyk, L. B. Jonker, G. Świątek, F. M. Tangerman and J. J. P. Veerman: *Differentiable Circle Maps with a Flat Interval*. *Comm. Math. Phys.* **173**, 599-622 (1995)

- [4] J. Graczyk, D. Sands and G. Świątek: *Metric Attractors for Smooth Unimodal Maps*. *Annals of Mathematics* **159**, 725-740 (2004)
- [5] M. Martens, S. Van Strien, W. De Melo and P. Mendes: *On Cherry flows*. *Ergodic Theory and Dynamical Systems* **10**, 531-554 (1990)
- [6] P. Mattila: *Geometry of Sets and Measures in Euclidean Spaces*. Cambridge University Press, (1995)
- [7] P. Mendes: *A metric property of Cherry vector fields on the torus*. *Journal of Differential Equations* **89**, 305-316 (1990)
- [8] P. C. Moreira and A. A. Gaspar: *Metric Properties of Cherry Flows*. *Journal of Differential Equations* **97**, 16-26 (1992)
- [9] I. Nikolaev, E. Zhuzhoma: *Flows on 2-dimensional Manifolds*. Springer-Verlag Berlin Heidelberg, (1999)
- [10] J. J. P. Vermann: *Irrational rotation number*. *Nonlinearity* **2**, 419-428 (1989)