

HYPERBOLIC LAMBERT QUADRILATERALS AND QUASICONFORMAL MAPPINGS

MATTI VUORINEN AND GENDI WANG

ABSTRACT. We prove sharp bounds for the product and the sum of two hyperbolic distances between the opposite sides of hyperbolic Lambert quadrilaterals in the unit disk. Furthermore, we study the images of Lambert quadrilaterals under quasiconformal mappings from the unit disk onto itself and obtain sharp results in this case, too.

KEYWORDS. Hyperbolic quadrilateral, quasiconformal mapping

2010 MATHEMATICS SUBJECT CLASSIFICATION. 51M09(51M15)

1. INTRODUCTION

Given a pair of points in the closure of the unit disk \mathbb{B}^2 , there exists a unique hyperbolic geodesic line joining these two points. Hyperbolic lines are simply sets of the form $C \cap \mathbb{B}^2$, where C is a circle perpendicular to the unit circle, or a Euclidean diameter of \mathbb{B}^2 . For a quadruple of four points $\{a, b, c, d\}$ in the closure of the unit disk, we can draw these hyperbolic lines joining each of the four pairs of points $\{a, b\}$, $\{b, c\}$, $\{c, d\}$, and $\{d, a\}$. If these hyperbolic lines bound a domain $D \subset \mathbb{B}^2$ such that the points $\{a, b, c, d\}$ are in the positive order on the boundary of the domain, then we say that the quadruple of points $\{a, b, c, d\}$ determines a hyperbolic quadrilateral $Q(a, b, c, d)$ and that the points a, b, c, d are its vertices. A hyperbolic quadrilateral with angles equal to $\pi/2, \pi/2, \pi/2, \phi$ ($0 \leq \phi < \pi/2$), is called a hyperbolic *Lambert* quadrilateral [Be, p. 156], see Figure 1. Observe that one of the vertices of a Lambert quadrilateral may be on the unit circle, in which case the angle at that vertex is $\phi = 0$.

In this paper, we study bounds for the product and the sum of two hyperbolic distances between the opposite sides of hyperbolic Lambert quadrilaterals in the unit disk. Also, we consider the same product expression for the images of these hyperbolic Lambert quadrilaterals under quasiconformal mappings from the unit disk onto itself. In particular, we obtain similar results for ideal hyperbolic quadrilaterals, i.e., in the case when all the vertices are on the unit circle and all the angles are zero. This follows, because an ideal hyperbolic quadrilateral can be subdivided into four Lambert quadrilaterals.

For the formulation of our main results we introduce some notation – further notation will be given below in Section 2. Let $J^*[a, b]$ be the hyperbolic geodesic line with end points $a, b \in \partial\mathbb{B}^2$, and let $J[a, b]$ be the hyperbolic geodesic segment joining a and b when $a, b \in \mathbb{B}^2$, or the hyperbolic geodesic ray when one of the two points a, b is on $\partial\mathbb{B}^2$.

Given two nonempty subsets A, B of \mathbb{B}^2 (or of the upper half plane \mathbb{H}^2), let $d_\rho(A, B)$ denote the hyperbolic distance between them, defined as

$$d_\rho(A, B) = \inf_{\substack{x \in A \\ y \in B}} \rho(x, y),$$

where $\rho(x, y)$ stands for the hyperbolic distance (2.4) (or (2.3) in the case $A, B \subset \mathbb{H}^2$).

We now formulate our main results.

Theorem 1.1. *Let $Q(v_a, v_b, v_c, v_d)$ be a hyperbolic Lambert quadrilateral in \mathbb{B}^2 and let the quadruple of interior angles $(\frac{\pi}{2}, \frac{\pi}{2}, \phi, \frac{\pi}{2})$, $\phi \in [0, \pi/2)$, correspond to the quadruple (v_a, v_b, v_c, v_d) of vertices. Let $d_1 = d_\rho(J[v_a, v_d], J[v_b, v_c])$, $d_2 = d_\rho(J[v_a, v_b], J[v_c, v_d])$ (see Figure 1), and let $L = \text{th}\rho(v_a, v_c) \in (0, 1]$. Then*

$$d_1 d_2 \leq \left(\text{arth} \left(\frac{\sqrt{2}}{2} L \right) \right)^2.$$

The equality holds if and only if v_c is on the bisector of the interior angle at v_a .




FIGURE 1. A hyperbolic Lambert quadrilateral in \mathbb{B}^2 .

Theorem 1.2. *Let $Q(v_a, v_b, v_c, v_d)$, d_1 , d_2 and L be as in Theorem 1.1. Let $m = \sqrt{(2 - L^2)(3L^2 - 2)}$, $r_0 = \sqrt{\frac{1 - m/L^2}{2}}$ and $r'_0 = \sqrt{1 - r_0^2}$.*

(1) *If $0 < L \leq \sqrt{\frac{2}{3}}$, then*

$$\text{arth } L < d_1 + d_2 \leq \text{arth} \left(\frac{2\sqrt{2}L}{2 + L^2} \right).$$

The equality holds in the right-hand side if and only if v_c is on the bisector of the interior angle at v_a .

(2) *If $\sqrt{\frac{2}{3}} < L < \sqrt{2(\sqrt{2} - 1)}$, then*

$$\text{arth } L < d_1 + d_2 \leq \text{arth} \left(\frac{L(r_0 + r'_0)}{1 + L^2 r_0 r'_0} \right).$$

The equality holds in the right-hand side if and only if the interior angle between $J[v_a, v_b]$ and $J[v_a, v_c]$ is $\arccos r_0$ or $\arccos r'_0$.

(3) *If $\sqrt{2(\sqrt{2} - 1)} \leq L < 1$, then*

$$\text{arth} \left(\frac{2\sqrt{2}L}{2 + L^2} \right) \leq d_1 + d_2 \leq \text{arth} \left(\frac{L(r_0 + r'_0)}{1 + L^2 r_0 r'_0} \right).$$

The equality holds in the left-hand side if and only if v_c is on the bisector of the interior angle at v_a . The equality holds in the right-hand side if and only if the interior angle between $J[v_a, v_b]$ and $J[v_a, v_c]$ is $\arccos r_0$ or $\arccos r'_0$.

(4) If $L = 1$, then

$$d_1 + d_2 \geq \operatorname{arth} \left(\frac{2\sqrt{2}}{3} \right).$$

The equality holds if and only if v_c is on the bisector of the interior angle at v_a .

In a Lambert quadrilateral, the angle ϕ is related to the lengths d_1, d_2 of the sides "opposite" to it as follows [Be, Theorem 7.17.1]:

$$\operatorname{sh} d_1 \operatorname{sh} d_2 = \cos \phi.$$

See also the recent paper of A. F. Beardon and D. Minda [BM, Lemma 5]. The proof of Theorem 1.1 yields the following corollary, which provides a connection between d_1, d_2 and $L = \operatorname{th} \rho(v_a, v_c)$.

Corollary 1.3. *Let L, d_1 and d_2 be as in Theorem 1.1. Then*

$$\operatorname{th}^2 d_1 + \operatorname{th}^2 d_2 = L^2.$$

By Theorem 1.1 and Theorem 1.2, we obtain the following corollary which deals with the ideal hyperbolic quadrilaterals.

Corollary 1.4. *Let $Q(a, b, c, d)$ be an ideal hyperbolic quadrilateral in \mathbb{B}^2 . Let $d_1 = d_\rho(J^*[a, d], J^*[b, c])$ and $d_2 = d_\rho(J^*[a, b], J^*[c, d])$ (see Figure 2). Then*

$$d_1 d_2 \leq \left(2 \log(\sqrt{2} + 1) \right)^2$$

and

$$d_1 + d_2 \geq 4 \log(\sqrt{2} + 1).$$

In both cases the equalities hold if and only if $|a, b, c, d| = 2$.




FIGURE 2. An ideal hyperbolic quadrilateral in \mathbb{B}^2 .

Remark 1.5. $|a, b, c, d| = 2$ means that there exists a Möbius transformation f such that $f(a) = 1, f(b) = i, f(c) = -1, f(d) = -i$, see (2.2).

Theorem 1.6. Let $f : \mathbb{B}^2 \rightarrow \mathbb{B}^2$ be a K -quasiconformal mapping with $f\mathbb{B}^2 = \mathbb{B}^2$ and let $Q(v_a, v_b, v_c, v_d)$, d_1, d_2 , L be as in Theorem 1.1. Let $A(K)$ be as in Lemma 4.2 and $f_L(r)$ be as in Lemma 2.9(1) by taking $c = L$. Denote $D_1 = d_\rho(f(J[v_a, v_d]), f(J[v_b, v_c]))$ and $D_2 = d_\rho(f(J[v_a, v_b]), f(J[v_c, v_d]))$.

(1) If $0 < L \leq \frac{e^2-1}{e^2+1} \approx 0.761594$, then

$$D_1 D_2 \leq A(K)^2 \left(\operatorname{arth} \left(\frac{\sqrt{2}}{2} L \right) \right)^{2/K}.$$

(2) If $\frac{e^2-1}{e^2+1} < L \leq 1$, then let $r_L = \frac{1}{L} \frac{e^2-1}{e^2+1} \approx \frac{0.761594}{L}$ and

$$M_L = \frac{f_L(\sqrt{1-r_L^2})}{f_L(r_L)} > 1.$$

Let $r_L(K)$ be the unique solution r to the equation $K f_L(r) = f_L(\sqrt{1-r^2})$ with $r_L < r < 1$. Further, define

$$T(x, L) = \operatorname{arth}(Lx) \left(\operatorname{arth} \left(L\sqrt{1-x^2} \right) \right)^{1/K}, \quad 0 < x < 1.$$

Then

$$D_1 D_2 \leq A(K)^2 \max \left\{ T(r_L(K), L), \left(\operatorname{arth} \left(\frac{\sqrt{2}}{2} L \right) \right)^{2/K} \right\}$$

if $K > M_L$, and

$$D_1 D_2 \leq A(K)^2 \max \left\{ T(r_L, L), \left(\operatorname{arth} \left(\frac{\sqrt{2}}{2} L \right) \right)^{2/K} \right\}$$

if $1 \leq K \leq M_L$.

Corollary 1.7. Let $f : \mathbb{B}^2 \rightarrow \mathbb{B}^2$ be a K -quasiconformal mapping with $f\mathbb{B}^2 = \mathbb{B}^2$ and let $Q(a, b, c, d)$, d_1, d_2 be as in Corollary 1.4. Let $A(K)$ be as in Lemma 4.2 and $f_1(r)$ be as in Lemma 2.9(1) by taking $c = 1$. Denote $D_1 = d_\rho(f(J^*[a, d]), f(J^*[b, c]))$ and $D_2 = d_\rho(f(J^*[a, b]), f(J^*[c, d]))$. Further denote $r_1 = \frac{2\sqrt{e}}{e+1} \approx 0.886819$ and

$$M_1 = \frac{(e-1)(\log(\sqrt{e}+1) - \log(\sqrt{e}-1))}{\sqrt{e}} \approx 1.46618$$

and define $r_1(K)$ to be the unique solution r to the equation $K f_1(r) = f_1(\sqrt{1-r^2})$ with $r_1 < r < 1$. With the notation

$$T(x) = \operatorname{arth}(x) \left(\operatorname{arth}(\sqrt{1-x^2}) \right)^{1/K}, \quad 0 < x < 1,$$

we have

$$D_1 D_2 \leq A(K)^2 \max \left\{ 2^{1+1/K} T(r_1(K)), \left(2 \log(\sqrt{2}+1) \right)^2 \right\}$$

if $K > M_1$, and

$$D_1 D_2 \leq A(K)^2 \max \left\{ 2^{1+1/K} T(r_1), \left(2 \log(\sqrt{2}+1) \right)^2 \right\}$$

if $1 \leq K \leq M_1$.

2. PRELIMINARIES

It is assumed that the reader is familiar with basic definitions of geometric function theory and quasiconformal mapping theory, see e.g. [Be, V]. We recall here some basic information on hyperbolic geometry [Be].

The chordal metric is defined by

$$(2.1) \quad \begin{cases} q(x, y) = \frac{|x-y|}{\sqrt{1+|x|^2}\sqrt{1+|y|^2}}, & x, y \neq \infty, \\ q(x, \infty) = \frac{1}{\sqrt{1+|x|^2}}, & x \neq \infty, \end{cases}$$

for $x, y \in \overline{\mathbb{R}^2}$.

For an ordered quadruple a, b, c, d of distinct points in $\overline{\mathbb{R}^2}$ we define the absolute ratio by

$$|a, b, c, d| = \frac{q(a, c)q(b, d)}{q(a, b)q(c, d)}.$$

It follows from (2.1) that for distinct points $a, b, c, d \in \mathbb{R}^2$

$$(2.2) \quad |a, b, c, d| = \frac{|a-c||b-d|}{|a-b||c-d|}.$$

The most important property of the absolute ratio is Möbius invariance, see [Be, Theorem 3.2.7], i.e., if f is a Möbius transformation, then

$$|f(a), f(b), f(c), f(d)| = |a, b, c, d|,$$

for all distinct $a, b, c, d \in \overline{\mathbb{R}^2}$.

For a domain $G \subsetneq \mathbb{R}^2$ and a continuous weight function $w : G \rightarrow (0, \infty)$, we define the weighted length of a rectifiable curve $\gamma \subset G$ to be

$$\ell_w(\gamma) = \int_{\gamma} w(z)|dz|$$

and the weighted distance between two points $x, y \in G$ by

$$d_w(x, y) = \inf_{\gamma} \ell_w(\gamma),$$

where the infimum is taken over all rectifiable curves in G joining x and y ($x = (x_1, x_2)$, $y = (y_1, y_2)$). It is easy to see that d_w defines a metric on G and (G, d_w) is a metric space. We say that a curve $\gamma : [0, 1] \rightarrow G$ is a geodesic joining $\gamma(0)$ and $\gamma(1)$ if for all $t \in (0, 1)$, we have

$$d_w(\gamma(0), \gamma(1)) = d_w(\gamma(0), \gamma(t)) + d_w(\gamma(t), \gamma(1)).$$

The hyperbolic distance in \mathbb{H}^2 and \mathbb{B}^2 is defined in terms of the weight functions $w_{\mathbb{H}^2}(x) = 1/x_2$ and $w_{\mathbb{B}^2}(x) = 2/(1 - |x|^2)$, resp. We also have the corresponding explicit formulas

$$(2.3) \quad \cosh \rho_{\mathbb{H}^2}(x, y) = 1 + \frac{|x - y|^2}{2x_2y_2}$$

for all $x, y \in \mathbb{H}^2$ [Be, p.35], and

$$(2.4) \quad \operatorname{th} \frac{\rho_{\mathbb{B}^2}(x, y)}{2} = \frac{|x - y|}{\sqrt{|x - y|^2 + (1 - |x|^2)(1 - |y|^2)}}$$

for all $x, y \in \mathbb{B}^2$ [Be, p.40]. In particular, for $t \in (0, 1)$,

$$(2.5) \quad \rho_{\mathbb{B}^2}(0, te_1) = \log \frac{1+t}{1-t} = 2\operatorname{arth}t.$$

There is a third equivalent way to express the hyperbolic distances. Let $G \in \{\mathbb{H}^2, \mathbb{B}^2\}$, $x, y \in G$ and let L be an arc of a circle perpendicular to ∂G with $x, y \in L$ and let $\{x_*, y_*\} = L \cap \partial G$, the points being labelled so that x_*, x, y, y_* occur in this order on L . Then by [Be, (7.2.6)]

$$(2.6) \quad \rho_G(x, y) = \sup\{\log |a, x, y, b| : a, b \in \partial G\} = \log |x_*, x, y, y_*|.$$

We will omit the subscript G if it is clear from the context. The hyperbolic distance is invariant under Möbius transformations of G onto G' for $G, G' \in \{\mathbb{H}^2, \mathbb{B}^2\}$.

Hyperbolic geodesics are arcs of circles which are orthogonal to the boundary of the domain. More precisely, for $a, b \in \mathbb{B}^2$ (or \mathbb{H}^2), the hyperbolic geodesic segment joining a to b is an arc of a circle orthogonal to S^1 (or $\partial\mathbb{H}^2$). In a limiting case the points a and b are located on a Euclidean line through 0 (or located on a normal of $\partial\mathbb{H}^2$), see [Be]. Therefore, the points x_* and y_* are the end points of the hyperbolic geodesic. For any two distinct points the hyperbolic geodesic segment is unique (see Figure 3 and 4). For basic facts about hyperbolic geometry we refer the interested reader to [A], [Be] and [KL].




FIGURE 3. Hyperbolic geodesic segments in \mathbb{H}^2 .




FIGURE 4. Hyperbolic geodesic segments in \mathbb{B}^2 .

By [K, Exercise 1.1.27] and [KV, Lemma 2.2], for $x, y \in \mathbb{R}^2 \setminus \{0\}$ such that $0, x, y$ are noncollinear, the circle $S^1(a, r_a)$ containing x, y is orthogonal to the unit circle, where

$$(2.7) \quad a = i \frac{y(1 + |x|^2) - x(1 + |y|^2)}{2(x_2y_1 - x_1y_2)} \quad \text{and} \quad r_a = \frac{|x - y||x|y|^2 - y|}{2|y||x_1y_2 - x_2y_1|}.$$

For $r, s \in (0, +\infty)$, the *Hölder mean of order p* is defined by

$$H_p(r, s) = \left(\frac{r^p + s^p}{2} \right)^{1/p} \quad \text{for } p \neq 0, \quad H_0(r, s) = \sqrt{rs}.$$

For $p = 1$, we get the arithmetic mean $A = H_1$; for $p = 0$, the geometric mean $G = H_0$; and for $p = -1$, the harmonic mean $H = H_{-1}$. It is well-known that $H_p(r, s)$ is continuous

and increasing with respect to p . Many interesting properties of Hölder means are given in [Bu] and [HLP].

A function $f : I \rightarrow J$ is called $H_{p,q}$ -convex (concave) if it satisfies

$$f(H_p(r, s)) \leq (\geq) H_q(f(r), f(s))$$

for all $r, s \in I$, and *strictly* $H_{p,q}$ -convex (concave) if the inequality is strict except for $r = s$. For $H_{p,q}$ -convexity of some special functions the reader is referred to [AVV2, Ba, WZJ1, WZJ2, ZWC].

Some other notation is also needed in the paper. Let $[a, b]$ be the Euclidean segment with end points a and b . Let X, Y be the real axis and imaginary axis, resp. Let $\text{Arc}(abc)$ be the circular arc with end points a, c and through b , and $\text{SArc}(ac)$ be semicircle with end point a, c .

The next lemma, so-called *monotone form of l'Hôpital's rule*, has found recently numerous applications in proving inequalities. See the extensive bibliography of [AVZ].

Lemma 2.8. [AVV1, Theorem 1.25] *For $-\infty < a < b < \infty$, let $f, g : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$, and be differentiable on (a, b) , and let $g'(x) \neq 0$ on (a, b) . If $f'(x)/g'(x)$ is increasing (decreasing) on (a, b) , then so are*

$$\frac{f(x) - f(a)}{g(x) - g(a)} \quad \text{and} \quad \frac{f(x) - f(b)}{g(x) - g(b)}.$$

If $f'(x)/g'(x)$ is strictly monotone, then the monotonicity in the conclusion is also strict.

From now on we let $r' = \sqrt{1 - r^2}$ for $0 < r < 1$.

Lemma 2.9. *Let $c \in (0, 1]$ and $r \in (0, 1)$.*

(1) *The function $f_c(r) \equiv \frac{1 - (cr')^2}{\text{arth}(cr)}$ is strictly decreasing and concave with range $(0, 1)$ if $c = 1$, and strictly decreasing with range $(0, \infty)$ if $0 < c < 1$.*

(2) *The function $F_c(r) \equiv \text{arth}(cr)\text{arth}(cr')$ is strictly increasing on $(0, \frac{\sqrt{2}}{2}]$ and strictly decreasing on $[\frac{\sqrt{2}}{2}, 1)$ with maximum value $(\text{arth}(\frac{\sqrt{2}}{2}c))^2$.*

Proof. (1) If $c = 1$, then $f_1(r) = \frac{r}{\text{arth}r}$. By differentiation,

$$f_1'(r) = \frac{h_{11}(r)}{h_{12}(r)},$$

where $h_{11}(r) = \text{arth}r - \frac{r}{r^2}$ and $h_{12}(r) = (\text{arth}r)^2$. It is easy to see that $h_{11}(0^+) = h_{12}(0^+) = 0$. Then

$$\frac{h'_{11}(r)}{h'_{12}(r)} = -\frac{h_{13}(r)}{h_{14}(r)},$$

where $h_{13}(r) = (\frac{r}{r'})^2$ and $h_{14}(r) = \text{arth}r$. Then $h_{13}(0^+) = h_{14}(0^+) = 0$. By differentiation, we have

$$\frac{h'_{13}(r)}{h'_{14}(r)} = \frac{2r}{r'^2},$$

which is strictly increasing. Hence by Lemma 2.8, $\frac{h'_{11}(r)}{h'_{12}(r)}$ is strictly decreasing and so is f_1' with $f_1'(r) < f_1'(0^+) = 0$. Therefore, f_1 is strictly decreasing and concave on $(0, 1)$. The limiting value $f_1(1^-) = 0$ is clear and $f_1(0^+) = 1$ by l'Hôpital's Rule.

If $0 < c < 1$, then

$$f_c(r) = f_1(cr)h(r),$$

where $h(r) = \frac{1-(cr')^2}{cr^2}$. By differentiation,

$$h'(r) = -\frac{2(1-c^2)}{cr^3} < 0.$$

Therefore, f_c is strictly decreasing. The limiting values are clear.

(2) By differentiation,

$$F'_c(r) = \frac{c}{r'f_c(r)} \left(\frac{f_c(r)}{f_c(r')} - 1 \right).$$

$\frac{f_c(r)}{f_c(r')}$ is strictly decreasing from $(0, 1)$ onto $(0, \infty)$ by (1) and $F'_c(\frac{\sqrt{2}}{2}) = 0$. Then F_c is strictly increasing on $(0, \frac{\sqrt{2}}{2}]$ and strictly decreasing on $[\frac{\sqrt{2}}{2}, 1)$ with maximum value $F_c(\frac{\sqrt{2}}{2})$. \square

Lemma 2.10. *Let $c \in (0, 1]$, $r \in (0, 1)$, $m = \sqrt{(2-c^2)(3c^2-2)}$ and $r_0 = \sqrt{\frac{1-m/c^2}{2}}$. Let*

$$G_c(r) \equiv \operatorname{arth}(cr) + \operatorname{arth}(cr').$$

(1) *If $0 < c \leq \sqrt{\frac{2}{3}}$, then the range of G_c is $(\operatorname{arth} c, \operatorname{arth}(\frac{2\sqrt{2}c}{2+c^2}))$.*

(2) *If $\sqrt{\frac{2}{3}} < c < \sqrt{2(\sqrt{2}-1)}$, then the range of G_c is $(\operatorname{arth} c, \operatorname{arth}(\frac{c(r_0+r'_0)}{1+c^2r_0r'_0}))$.*

(3) *If $\sqrt{2(\sqrt{2}-1)} \leq c < 1$, then the range of G_c is $[\operatorname{arth}(\frac{2\sqrt{2}c}{2+c^2}), \operatorname{arth}(\frac{c(r_0+r'_0)}{1+c^2r_0r'_0})]$.*

(4) *If $c = 1$, then the range of G_c is $[\operatorname{arth}(\frac{2\sqrt{2}}{3}), \infty)$.*

Proof. It is clear that the limiting values

$$G_c(0^+) = G_c(1^-) = \begin{cases} \operatorname{arth} c, & 0 < c < 1, \\ \infty, & c = 1. \end{cases}$$

Let

$$g_c(r) = \operatorname{th}G_c(r) = \frac{c(r+r')}{1+c^2rr'}.$$

By differentiation, we have

$$r'(1+c^2rr')^2 g'_c(r) = c(r'-r)(1-c^2-c^2rr').$$

Making substitution of $x = r^2$, we get

$$(2.11) \quad \begin{aligned} 1 - c^2 - c^2rr' &= 0 \\ \Leftrightarrow c^4x^2 - c^4x + (1 - c^2)^2 &= 0. \end{aligned}$$

Therefore, equation (2.11) has no root if $0 < c^2 < \frac{2}{3}$ or $c = 1$, only one root $\frac{\sqrt{2}}{2}$ if $c^2 = \frac{2}{3}$, and two different roots r_0, r'_0 if $\frac{2}{3} < c^2 < 1$. It is obvious that $r_0 \in (0, \frac{\sqrt{2}}{2})$.

It is easy to see that $c = g_c(0) < g_c(\frac{\sqrt{2}}{2}) = \frac{2\sqrt{2}}{2+c^2}c$ if and only if $c^2 < 2(\sqrt{2}-1) \approx 0.828427$.

If $\sqrt{\frac{2}{3}} < c < 1$, g_c is increasing on $(0, r_0)$, $(\frac{\sqrt{2}}{2}, r'_0)$ and decreasing on $(r_0, \frac{\sqrt{2}}{2})$, $(r'_0, 1)$.

Now we get the following conclusions.

(1) If $0 < c \leq \sqrt{\frac{2}{3}}$, $g'_c(r) = 0$ is equivalent to $r = \frac{\sqrt{2}}{2}$. Then

$$\operatorname{arth} c = \operatorname{arth}(g_c(0)) < G_c(r) \leq \operatorname{arth}(g_c(\sqrt{2}/2)) = \operatorname{arth} \left(\frac{2\sqrt{2}c}{2+c^2} \right).$$

(2) If $\sqrt{\frac{2}{3}} < c < \sqrt{2(\sqrt{2}-1)}$, then

$$\operatorname{arth} c = \operatorname{arth}(g_c(0)) < G_c(r) \leq \operatorname{arth}(g_c(r_0)) = \operatorname{arth} \left(\frac{c(r_0 + r'_0)}{1 + c^2 r_0 r'_0} \right).$$

(3) If $\sqrt{2(\sqrt{2}-1)} \leq c < 1$, then

$$\operatorname{arth} \left(\frac{2\sqrt{2}c}{2 + c^2} \right) = \operatorname{arth}(g_c(\sqrt{2}/2)) \leq G_c(r) \leq \operatorname{arth}(g_c(r_0)) = \operatorname{arth} \left(\frac{c(r_0 + r'_0)}{1 + c^2 r_0 r'_0} \right).$$

(4) If $c = 1$, then

$$G_c(r) \geq \operatorname{arth}(g_1(\sqrt{2}/2)) = \operatorname{arth}(2\sqrt{2}/3).$$

This completes the proof. \square

Lemma 2.12. *Let $r \in (0, 1)$.*

(1) *The function $h_1(r) \equiv \frac{r'}{\operatorname{arth} r'}$ is strictly increasing and concave with range $(0, 1)$.*

(2) *The function $h(r) \equiv \frac{r}{\operatorname{arth} r} + \frac{r'}{\operatorname{arth} r'}$ is strictly increasing on $(0, \frac{\sqrt{2}}{2}]$, strictly decreasing on $[\frac{\sqrt{2}}{2}, 1)$, and concave on $(0, 1)$ with range $(1, \frac{\sqrt{2}}{\log(\sqrt{2}+1)})$.*

Proof. (1) The monotonicity and the limiting values of h_1 can be easily obtained by Lemma 2.9(1).

Now we prove the concavity of h_1 . By differentiation,

$$h'_1(r) = \frac{r' - r^2 \operatorname{arth} r'}{r r' (\operatorname{arth} r')^2}$$

and

$$(2.13) \quad h''_1(r) r^2 r'^3 (\operatorname{arth} r')^3 = 2r'^2 - r' \operatorname{arth} r' - r^2 (\operatorname{arth} r')^2 \equiv \psi_1(r').$$

Then $\psi_1(r) = 2r^2 - r \operatorname{arth} r - r'^2 (\operatorname{arth} r)^2$ and by differentiation

$$\psi'_1(r) = r(4 - \psi_2(r)),$$

where $\psi_2(r) = \frac{1}{r'^2} + 3 \frac{\operatorname{arth} r}{r} - 2(\operatorname{arth} r)^2$. Since

$$\operatorname{arth} r = \frac{1}{2} \log \frac{1+r}{1-r} = \sum_{n=0}^{\infty} \frac{r^{2n+1}}{2n+1},$$

we have

$$\begin{aligned} r^2 r'^4 \psi'_2(r) &= (r^4 + 2r^2 - 3) \operatorname{arth} r - r^3 + 3r \\ &= \sum_{n=0}^{\infty} \frac{r^{2n+5}}{2n+1} + 2 \sum_{n=1}^{\infty} \frac{r^{2n+3}}{2n+1} - 3 \sum_{n=2}^{\infty} \frac{r^{2n+1}}{2n+1} \\ &= \sum_{n=2}^{\infty} \frac{16(n-1)}{(2n-3)(2n-1)(2n+1)} r^{2n+1} > 0. \end{aligned}$$

Hence ψ_2 is increasing with $\psi_2(0^+) = 4$ and ψ_1 is decreasing with $\psi_1(0^+) = 0$. Therefore by (2.13), h''_1 is negative and h'_1 is decreasing. Then h_1 is concave on $(0, 1)$.

(2) By differentiation,

$$h'(r) = f'_1(r) + h'_1(r) = f'_1(r) - \frac{r}{r'} f'_1(r'),$$

where $f_1(r) = \frac{r}{\operatorname{arth} r} = h_1(r')$. It is easy to see that $h'(\frac{\sqrt{2}}{2}) = 0$. Therefore, h is strictly increasing on $(0, \frac{\sqrt{2}}{2}]$ and strictly decreasing on $[\frac{\sqrt{2}}{2}, 1)$. By (1) and Lemma 2.9(1), h' is strictly decreasing and $h(0^+) = h(1^-) = 1$. Hence h is concave and

$$1 < h(r) \leq h(\sqrt{2}/2) = \frac{\sqrt{2}}{\log(\sqrt{2} + 1)}.$$

This completes the proof. \square

Lemma 2.14. *Let $p \in \mathbb{R}$, $r \in (0, 1)$, and $C = 1 - \frac{\log(\sqrt{2}+1)}{\sqrt{2}} \approx 0.376775$. Let*

$$g(r) = \frac{r}{r'} \left(\frac{\operatorname{arth} r}{\operatorname{arth} r'} \right)^{p-1}.$$

(1) g is strictly decreasing if $p \leq 0$ and strictly increasing if $p \geq C$.

(2) If $p \in (0, C)$, then there exists exactly one point $r_0 \in (0, \frac{\sqrt{2}}{2})$ such that g is increasing on $(0, r_0)$, $(r'_0, 1)$ and decreasing on (r_0, r'_0) .

(3) If $p \in (0, C)$, then $g(0^+) = 0$ and $g(1^-) = \infty$.

Proof. (1) By logarithmic differentiation,

$$\begin{aligned} \frac{rr'^2 \operatorname{arth} r \operatorname{arth} r'}{\operatorname{arth} r' + r' \operatorname{arth} r} \cdot \frac{g'(r)}{g(r)} &= p - 1 + \frac{\operatorname{arth} r \operatorname{arth} r'}{r \operatorname{arth} r' + r' \operatorname{arth} r} \\ &= p - \left(1 - \frac{1}{h(r)} \right), \end{aligned}$$

where $h(r)$ is as in Lemma 2.12(2). Since

$$0 < 1 - \frac{1}{h(r)} \leq C,$$

we see that g is strictly increasing if $p \geq C$ and decreasing if $p \leq 0$.

(2) If $p \in (0, C)$, then there exists exactly one point $r_0 \in (0, \frac{\sqrt{2}}{2})$ such that $g'(r_0) = g'(r'_0) = 0$ because $1 - \frac{1}{h(r)}$ is increasing on $(0, \frac{\sqrt{2}}{2})$ and decreasing on $(\frac{\sqrt{2}}{2}, 1)$ by Lemma 2.12(2). Therefore, $g' > 0$ if $r \in (0, r_0) \cup (r'_0, 1)$ and $g' < 0$ if $r \in (r_0, r'_0)$. Hence g is increasing on $(0, r_0)$, $(r'_0, 1)$ and decreasing on (r_0, r'_0) .

(3) Since $0 < p < C < 1$ and

$$\lim_{r \rightarrow 0^+} \frac{(\operatorname{arth} r')^{1-p}}{r^{-p}} = \lim_{r \rightarrow 0^+} \frac{1-p}{p} \cdot \frac{r^p}{r'(\operatorname{arth} r')^p} = 0,$$

we have

$$\lim_{r \rightarrow 0^+} g(r) = \lim_{r \rightarrow 0^+} \left(\frac{r}{\operatorname{arth} r} \right)^{1-p} \cdot \left(\frac{1}{r'} \right) \cdot \frac{(\operatorname{arth} r')^{1-p}}{r^{-p}} = 0$$

and

$$\lim_{r \rightarrow 1^-} g(r) = \lim_{r \rightarrow 1^-} \left(\frac{\operatorname{arth} r'}{r'} \right)^{1-p} \cdot r \cdot \frac{r'^{-p}}{(\operatorname{arth} r)^{1-p}} = \infty.$$

This completes the proof. \square

Theorem 2.15. *Let C be as in Lemma 2.14. Then for all $r \in (0, 1)$,*

$$(2.16) \quad H_p(\operatorname{arth} r, \operatorname{arth} r') \leq \operatorname{arth} \left(\frac{\sqrt{2}}{2} \right)$$

holds if and only if $p \leq 0$, and

$$(2.17) \quad H_p(\operatorname{arth} r, \operatorname{arth} r') \geq \operatorname{arth} \left(\frac{\sqrt{2}}{2} \right)$$

holds if and only if $p \geq C$. The equalities hold if and only if $r = r' = \frac{\sqrt{2}}{2}$ and all inequalities are sharp in both cases.

Proof. We immediately obtain the inequality (2.16) if $p = 0$ by Lemma 2.9(2). Therefore, it suffices to discuss the case $p \neq 0$.

Let

$$f(r) = \frac{1}{p} \log \frac{(\operatorname{arth} r)^p + (\operatorname{arth} r')^p}{2}, \quad p \neq 0.$$

By differentiation,

$$f'(r) = \frac{(\operatorname{arth} r')^{p-1}}{rr'((\operatorname{arth} r)^p + (\operatorname{arth} r')^p)} (g(r) - 1),$$

where $g(r)$ is as in Lemma 2.14 and it is easy to see that $g(\frac{\sqrt{2}}{2}) = 1$.

Case 1. $p < 0$. f is strictly increasing on $(0, \frac{\sqrt{2}}{2})$ and strictly decreasing on $(\frac{\sqrt{2}}{2}, 1)$ by Lemma 2.14(1). Therefore, $f(r) \leq f(\frac{\sqrt{2}}{2})$ for all $p < 0$.

Case 2. $p \geq C$. By Lemma 2.14(1), f is strictly decreasing on $(0, \frac{\sqrt{2}}{2})$ and strictly increasing on $(\frac{\sqrt{2}}{2}, 1)$, and hence $f(r) \geq f(\frac{\sqrt{2}}{2})$.

Case 3. $p \in (0, C)$. By Lemma 2.14(2), there exists exactly one point $r_1 \in (0, r_0)$ such that $g(r_1) = g(r'_1) = 1$, where $r_0 \in (0, \frac{\sqrt{2}}{2})$ is as in Lemma 2.14(2). Then f is strictly decreasing on $(0, r_1)$, $(\frac{\sqrt{2}}{2}, r'_1)$ and strictly increasing on $(r_1, \frac{\sqrt{2}}{2})$, $(r'_1, 1)$. Thus,

$$f(r_1) = f(r'_1) < f(\sqrt{2}/2).$$

Since $f(0^+) = f(1^-) = \infty$, there exists $r_2 \in (0, r_1) \cup (r'_1, 1)$ such that

$$f(\sqrt{2}/2) < f(r_2).$$

Therefore, neither (2.16) nor (2.17) holds for all $r \in (0, 1)$.

This completes the proof of Theorem 2.15. □

Lemma 2.18. *Let $r \in (0, 1)$.*

(1) *The function $f(r) \equiv \frac{r'^4 \operatorname{arth} r - r(1+r^2)}{r'^2((1+r^2)\operatorname{arth} r - r)}$ is strictly decreasing with range $(-\infty, -2)$.*

(2) *For $p \in \mathbb{R}$ define*

$$h_p(r) \equiv 1 + pr'^2 \frac{\operatorname{arth} r}{r} - (1 + r^2) \frac{\operatorname{arth} r}{r}.$$

(i) *If $p \geq -2$, then the range of h_p is $(-\infty, p)$.*

(ii) *If $p < -2$, then the range of h_p is $(-\infty, C(p)]$, where $C(p) \equiv \sup_{0 < r < 1} h_p(r) \in (p, -1)$*

with $\lim_{p \rightarrow -2} C(p) = -2$ and $\lim_{p \rightarrow -\infty} C(p) = -\infty$.

Proof. (1) Let $f_1(r) = r'^2 \operatorname{arth} r - \frac{r(1+r^2)}{r'^2}$ and $f_2(r) = (1+r^2) \operatorname{arth} r - r$, then $f_1(0^+) = f_2(0^+) = 0$. By differentiation, we have

$$\frac{f_1'(r)}{f_2'(r)} = -1 - \frac{2r}{r'^2(r + r'^2 \operatorname{arth} r)} = -1 - \frac{2}{r'^2 \left(1 + \frac{f_3(r)}{f_4(r)}\right)},$$

where $f_3(r) = \operatorname{arth} r$ and $f_4(r) = \frac{r}{r'^2}$. It is easy to see that $f_3(0^+) = f_4(0^+) = 0$, then

$$\frac{f_3'(r)}{f_4'(r)} = \frac{r'^2}{1+r^2},$$

which is strictly decreasing. Hence by Lemma 2.8, $\frac{f_1'(r)}{f_2'(r)}$ is strictly decreasing and so is f with $f(0^+) = -2$. Since $(1+r^2) \operatorname{arth} r - r > 0$ and by l'Hôpital's Rule,

$$(2.19) \quad \lim_{r \rightarrow 1^-} \frac{\operatorname{arth} r}{r'^{-2}} = \lim_{r \rightarrow 1^-} \frac{r'^2}{2r} = 0,$$

we get $f(1^-) = -\infty$.

(2) By Lemma 2.12(1) and (2.19), it is easy to see that $h_p(0^+) = p$ and $h_p(1^-) = -\infty$. Next by differentiation, we have

$$h_p'(r) = \frac{1}{r} \left((1+r^2) \frac{\operatorname{arth} r}{r} - 1 \right) (f(r) - p),$$

where $f(r)$ is as in (1).

If $p \geq -2$, and by (1), we see that $p > f(r)$, which implies that h_p is strictly decreasing and hence $h_p(r) < p$.

If $p < -2$, since the range of f is $(-\infty, -2)$, we see that there exists exactly one point $r_0 \in (0, 1)$ such that $p = f(r_0)$. Then h_p is increasing on $(0, r_0)$ and decreasing on $(r_0, 1)$. Since

$$h_p(r) = -1 + 2 \left(1 - \frac{\operatorname{arth} r}{r} \right) + (p+1)r'^2 \frac{\operatorname{arth} r}{r} < -1,$$

by the continuity of h_p , there is a continuous function

$$C(p) \equiv \sup_{0 < r < 1} h_p(r)$$

with $p < C(p) < -1$, $\lim_{p \rightarrow -2} C(p) = -2$ and $\lim_{p \rightarrow -\infty} C(p) = -\infty$. \square

Lemma 2.20. *Let $p, q \in \mathbb{R}$, $r \in (0, 1)$, and let $C(p)$ be as in Lemma 2.18(2). Let*

$$g_{p,q}(r) \equiv \frac{\operatorname{arth}^{q-1} r}{r^{p-1} r'^2}.$$

(1) *If $p \geq -2$, then $g_{p,q}$ is strictly increasing for each $q \geq p$, and $g_{p,q}$ is not monotone for any $q < p$.*

(2) *If $p < -2$, then $g_{p,q}$ is strictly increasing for each $q \geq C(p)$, and $g_{p,q}$ is not monotone for any $q < C(p)$.*

Proof. By logarithmic differentiation in r ,

$$\frac{g_{p,q}'(r)}{g_{p,q}(r)} = \frac{1}{r'^2 \operatorname{arth} r} (q - h_p(r)),$$

where $h_p(r)$ is as in Lemma 2.18(2). Hence the results immediately follow from Lemma 2.18(2). \square

The following theorem studies the $H_{p,q}$ -convexity of arth .

Theorem 2.21. *The inverse hyperbolic tangent function arth is strictly $H_{p,q}$ -convex on $(0, 1)$ if and only if $(p, q) \in D_1 \cup D_2$, where*

$$D_1 = \{(p, q) \mid -2 \leq p < +\infty, p \leq q < +\infty\},$$

$$D_2 = \{(p, q) \mid -\infty < p < -2, C(p) \leq q < +\infty\},$$

and $C(p)$ is a continuous function as in Lemma 2.18(2). There are no values of p and q for which arth is $H_{p,q}$ -concave on the whole interval $(0, 1)$.

Proof. The proof is divided into the following four cases.

Case 1. $p \neq 0$ and $q \neq 0$. We may suppose that $0 < x \leq y < 1$. Define

$$F(x, y) = \text{arth}^q(H_p(x, y)) - \frac{\text{arth}^q x + \text{arth}^q y}{2}.$$

Let $t = H_p(x, y)$, then $\frac{\partial t}{\partial x} = \frac{1}{2}(\frac{x}{t})^{p-1}$. If $x < y$, we see that $t > x$. By differentiation, we have

$$\frac{\partial F}{\partial x} = \frac{q}{2}x^{p-1} \left(\frac{\text{arth}^{q-1} t}{t^{p-1}t'^2} - \frac{\text{arth}^{q-1} x}{x^{p-1}x'^2} \right).$$

Case 1.1. $p \geq -2$, $q \geq p$, and $pq \neq 0$. By Lemma 2.20(1), $\frac{\partial F}{\partial x} < 0$ if $q < 0$ and $\frac{\partial F}{\partial x} > 0$ if $q > 0$. Then $F(x, y)$ is strictly decreasing and $F(x, y) \geq F(y, y) = 0$ if $q < 0$, and $F(x, y)$ is strictly increasing and $F(x, y) \leq F(y, y) = 0$ if $q > 0$. Hence we have

$$\text{arth}(H_p(x, y)) \leq H_q(\text{arth } x, \text{arth } y)$$

with equality if and only if $x = y$.

In conclusion, arth is strictly $H_{p,q}$ -convex on $(0, 1)$ for $(p, q) \in \{(p, q) \mid -2 \leq p < 0, p \leq q < 0\} \cup \{(p, q) \mid -2 \leq p < 0, q > 0\} \cup \{(p, q) \mid 0 < p < +\infty, q \geq p\}$.

Case 1.2. $p \geq -2$, $q < p$, and $pq \neq 0$. By Lemma 2.20(1), with an argument similar to Case 1.1, it is easy to see that arth is neither $H_{p,q}$ -concave nor $H_{p,q}$ -convex on the whole interval $(0, 1)$.

Case 1.3. $p < -2$, $q \geq C(p)$, and $pq \neq 0$. By Lemma 2.20(2), $\frac{\partial F}{\partial x} < 0$ if $q < 0$ and $\frac{\partial F}{\partial x} > 0$ if $q > 0$. Then $F(x, y)$ is strictly decreasing and $F(x, y) \geq F(y, y) = 0$ if $q < 0$, and $F(x, y)$ is strictly increasing and $F(x, y) \leq F(y, y) = 0$ if $q > 0$. Hence we have

$$\text{arth}(H_p(x, y)) \leq H_q(\text{arth } x, \text{arth } y)$$

with equality if and only if $x = y$.

In conclusion, arth is strictly $H_{p,q}$ -convex on $(0, 1)$ for $(p, q) \in \{(p, q) \mid p < -2, C(p) \leq q < 0\} \cup \{(p, q) \mid p < -2, q > 0\}$.

Case 1.4. $p < -2$, $q < C(p)$, and $pq \neq 0$. By Lemma 2.20(2), with an argument similar to Case 1.3, it is easy to see that arth is neither $H_{p,q}$ -concave nor $H_{p,q}$ -convex on the whole interval $(0, 1)$.

Case 2. $p \neq 0$ and $q = 0$. For $0 < x \leq y < 1$, let

$$F(x, y) = \frac{\text{arth}^2(H_p(x, y))}{\text{arth } x \text{arth } y},$$

and $t = H_p(x, y)$. If $x < y$, we see that $t > x$. By logarithmic differentiation, we obtain

$$\frac{1}{F} \frac{\partial F}{\partial x} = x^{p-1} \left(\frac{(\text{arth } t)^{-1}}{t^{p-1}t'^2} - \frac{(\text{arth } x)^{-1}}{x^{p-1}x'^2} \right).$$

Case 2.1. $-2 \leq p < 0$ and $q = 0 > p$. By Lemma 2.20(1), we have $\frac{\partial F}{\partial x} > 0$ and $F(x, y) \leq F(y, y) = 1$. Hence we have

$$\operatorname{arth}(H_p(x, y)) \leq \sqrt{\operatorname{arth} x \operatorname{arth} y}$$

with equality if and only if $x = y$.

In conclusion, arth is strictly $H_{p,q}$ -convex on $(0, 1)$ for $(p, q) \in \{(p, q) \mid -2 \leq p < 0, q = 0\}$.

Case 2.2. $p > 0$ and $q = 0 < p$. By Lemma 2.20(1), with an argument similar to Case 2.1, it is easy to see that arth is neither $H_{p,q}$ -concave nor $H_{p,q}$ -convex on the whole interval $(0, 1)$.

Case 2.3. $p < -2$. We have $q = 0 \geq C(p)$, and by Lemma 2.20(2), with an argument similar to Case 2.1, it is easy to see that arth is $H_{p,q}$ -convex on $(0, 1)$ for $(p, q) \in \{(p, q) \mid p < -2, q = 0\}$.

Case 3. $p = 0$ and $q \neq 0$. For $0 < x \leq y < 1$, let

$$F(x, y) = \operatorname{arth}^q(\sqrt{xy}) - \frac{\operatorname{arth}^q x + \operatorname{arth}^q y}{2},$$

and $t = \sqrt{xy}$. If $x < y$, we have that $t > x$. By differentiation, we obtain

$$\frac{\partial F}{\partial x} = \frac{q}{2x} \left(\frac{\operatorname{arth}^{q-1} t}{t^{-1} t'^2} - \frac{\operatorname{arth}^{q-1} x}{x^{-1} x'^2} \right).$$

Case 3.1. $q > p = 0$. By Lemma 2.20(1), we have $\frac{\partial F}{\partial x} > 0$ and $F(x, y) \leq F(y, y) = 0$. Hence we have

$$\operatorname{arth}(\sqrt{xy}) \leq H_q(\operatorname{arth} x, \operatorname{arth} y)$$

with equality if and only if $x = y$.

In conclusion, arth is strictly $H_{p,q}$ -convex on $(0, 1)$ for $(p, q) \in \{(p, q) \mid p = 0, q > 0\}$.

Case 3.2. $q < p = 0$. By Lemma 2.20(1), with an argument similar to Case 3.1, it is easy to see that arth is neither $H_{p,q}$ -concave nor $H_{p,q}$ -convex on the whole interval $(0, 1)$.

Case 4. $p = q = 0$. By Case 1.1, for all $x, y \in (0, 1)$, we have

$$\operatorname{arth}(H_p(x, y)) \leq H_p(\operatorname{arth} x, \operatorname{arth} y), \quad \text{for } p \geq -2 \quad \text{and} \quad p \neq 0.$$

By the continuity of H_p in p and arth in x , we have

$$\operatorname{arth}(H_0(x, y)) \leq H_0(\operatorname{arth} x, \operatorname{arth} y).$$

In conclusion, arth is strictly $H_{0,0}$ -convex on $(0, 1)$.

This completes the proof of Theorem 2.21. \square

Setting $p = 1 = q$ in Theorem 2.21, we easily obtain the convexity of arth .

Corollary 2.22. *The inverse hyperbolic tangent function arth is strictly convex on $(0, 1)$.*

By (2.5), Theorem 2.21 has a simple application to the hyperbolic metric.

Corollary 2.23. *Let $z \in S^1(H_p(|x|, |y|))$. Then for all $x, y \in \mathbb{B}^2 \setminus \{0\}$ and $p \geq -2$*

$$\rho(0, z) \leq H_p(\rho(0, x), \rho(0, y))$$

with equality if and only if $|x| = |y|$.

3. SOME PROPOSITIONS FOR HYPERBOLIC METRIC

Next we give some geometric propositions for the hyperbolic metric.

Proposition 3.1. *Let c, d be arbitrary two points on the unit circle such that $0, c, d$ are noncollinear. Let $b \in [c, d] \cap \mathbb{B}^2$, $\{a\} = [0, b] \cap J^*[c, d]$, and let s be the midpoint of the Euclidean segment $[c, d]$. Then*

- (1) *a is the hyperbolic midpoint of the hyperbolic segment $J[0, b]$;*
- (2) *s is on the hyperbolic circle $S_\rho(a, \rho(0, a)) = \{z | \rho(z, a) = \rho(0, a)\}$.*




FIGURE 5. The point a is the hyperbolic midpoint of $J[0, b]$. The circular arc (cad) is orthogonal to the unit circle. Here s is the Euclidean midpoint of the chord $[c, d]$.

Proof. Without loss of generality, we may assume $c = e^{i\alpha}$ and $d = e^{-i\alpha}$, where $0 < \alpha < \pi/2$.

- (1) The point a is the hyperbolic midpoint of the hyperbolic segment $J[0, b]$

$$(3.2) \quad \Leftrightarrow \rho(0, b) = 2\rho(0, a) \Leftrightarrow \log \frac{1 + |b|}{1 - |b|} = 2 \log \frac{1 + |a|}{1 - |a|} \Leftrightarrow |b| = \frac{2|a|}{1 + |a|^2}.$$

Let $b = |b|e^{i\beta}$ ($-\pi/2 < \beta < \pi/2$). Then by the orthogonality of $J^*[c, d]$ and the unit circle

$$s = \cos \alpha = |b| \cos \beta = |b| \frac{|a|^2 + |w|^2 - r^2}{2|a||w|},$$

where $w = 1/\cos \alpha$ and $r = \sqrt{|w|^2 - 1}$ are the center and the radius, resp., of the circle containing c, d and orthogonal to $\partial\mathbb{B}^2$. Therefore, the last equality in (3.2) holds and a is the hyperbolic midpoint of $J[0, b]$.

(2) It is easy to see that $[0, b]$ is also the Euclidean diameter of $S_\rho(a, \rho(0, a))$ by geometric observation and (1). Therefore, s is on the hyperbolic circle $S_\rho(a, \rho(0, a))$, see Figure 5. \square

Proposition 3.3. *Let $J_1 = J^*[e^{i\alpha}, -e^{-i\alpha}]$, $J_2 = J^*[-e^{i\alpha}, e^{-i\alpha}]$ be two hyperbolic geodesics in \mathbb{B}^2 , $0 < \alpha < \pi/2$. Let $\{te_2\} = J_1 \cap Y$, see Figure 6. Then*

$$(3.4) \quad d_\rho(J_1, J_2) = \rho(-te_2, te_2) = 2 \log \frac{1 + t}{1 - t}.$$

Proof. Let $g : \mathbb{B}^2 \rightarrow \mathbb{H}^2$ be a Möbius transformation which satisfies $g(i) = \infty$ and $g(-i) = 0$. Then $g([-te_2, te_2]) = [g(-te_2), g(te_2)]$ which is on the ray emanating from 0 and perpendicular to $\partial\mathbb{H}^2$. By the orthogonality of $J_i (i = 1, 2)$ and Y , we get

$$J_1^* = g(J_1) = \text{Arc}(g(-e^{-i\alpha})g(te_2)g(e^{i\alpha})) \quad , \quad J_2^* = g(J_2) = \text{Arc}(g(-e^{i\alpha})g(-te_2)g(e^{-i\alpha})).$$

Here $g(-e^{-i\alpha}) = -g(e^{i\alpha})$ and $g(-e^{i\alpha}) = -g(e^{-i\alpha})$, see Figure 7. For every rectifiable arc γ_{xy} , $x \in J_1^*$ and $y \in J_2^*$, by geometric observation we get

$$\int_{\gamma_{xy}} \frac{1}{d(z, \partial\mathbb{H}^2)} |dz| \geq \int_{[g(-te_2), g(te_2)]} \frac{1}{d(z, \partial\mathbb{H}^2)} |dz|,$$

and hence

$$(3.5) \quad d_\rho(J_1^*, J_2^*) = \rho(g(-te_2), g(te_2)).$$

Since the hyperbolic distance is invariant under Möbius transformations, by (3.5) we get

$$d_\rho(J_1, J_2) = \rho(-te_2, te_2) = 2 \log \frac{1+t}{1-t}.$$

□




FIGURE 6




FIGURE 7

The Möbius transformation g maps the unit disk in Figure 6 onto the upper half plane in Figure 7.

Let $J_3 = J^*[e^{i\alpha}, e^{-i\alpha}]$ and $J_4 = J^*[-e^{-i\alpha}, -e^{i\alpha}]$. Let $\{se_1\} = J_3 \cap X$. Since the hyperbolic distance is invariant under rotations, by Proposition 3.3 we also obtain

$$(3.6) \quad d_\rho(J_3, J_4) = \rho(-se_1, se_1).$$

By the proof of Proposition 3.3, $J_3^* = g(J_3) = \text{SArc}(g(e^{i\alpha})g(e^{-i\alpha}))$ and $J_4^* = g(J_4) = \text{SArc}(g(-e^{-i\alpha})g(-e^{i\alpha}))$. Then g maps $\text{Arc}((-i)(se_1)i)$ to the ray emanating from 0 and tangent to J_3^* at the point $g(se_1)$. Similarly, g maps $\text{Arc}((-i)(-se_1)i)$ to the ray emanating from 0 and tangent to J_4^* at the point $g(-se_1)$. Then we have

$$(3.7) \quad d_\rho(J_3^*, J_4^*) = \rho(g(-se_1), g(se_1)).$$

Remark 3.8. By (3.4) and (3.6), we find the two distances between the opposite sides for the hyperbolic quadrilaterals with four vertices $e^{i\alpha}$, $-e^{-i\alpha}$, $-e^{i\alpha}$, $e^{-i\alpha}$ on the unit circle and the counterpart for the upper half plane by (3.5) and (3.7).

4. PROOF OF MAIN RESULTS

Proof of Theorem 1.1. Since the hyperbolic distance is Möbius invariant, we may assume that $v_a = 0$, v_b is on X , v_d is on Y and $v_c = te^{i\theta}$, $0 < t \leq 1$ and $0 < \theta < \frac{\pi}{2}$ (see Figure 1). Then by (2.7) the circle $S^1(b, r_b)$ through v_c and \bar{v}_c is orthogonal to $\partial\mathbb{B}^2$, where

$$b = \frac{1+t^2}{2t \cos \theta} \quad \text{and} \quad r_b = \frac{\sqrt{(1+t^2)^2 - 4t^2 \cos^2 \theta}}{2t \cos \theta}.$$

By Proposition 3.3, we get

$$d_1 = \rho(0, v_b) = \log \frac{1 + (b - r_b)}{1 - (b - r_b)} = \operatorname{arth} \left(\frac{2t}{1+t^2} \cos \theta \right).$$

Similarly, we get

$$d_2 = \rho(0, v_d) = \operatorname{arth} \left(\frac{2t}{1+t^2} \sin \theta \right).$$

Then

$$d_1 d_2 = \operatorname{arth}(Lr) \operatorname{arth}(Lr'),$$

where $L = \frac{2t}{1+t^2} = \operatorname{th} \rho(0, v_c) \in (0, 1]$ and $r = \cos \theta \in (0, 1)$. By Lemma 2.9(2), we have

$$d_1 d_2 \leq \left(\operatorname{arth} \left(\frac{\sqrt{2}}{2} L \right) \right)^2.$$

This completes the proof of Theorem 1.1. \square

Proof of Theorem 1.2. By the proof of Theorem 1.1, we have

$$d_1 + d_2 = \operatorname{arth}(Lr) + \operatorname{arth}(Lr'),$$

where $L \in (0, 1]$ and $r \in (0, 1)$. Then by Lemma 2.10, the desired conclusion follows. \square

Proof of Corollary 1.4. There exists a Möbius transformation g which maps a, b, c, d to $e^{i\alpha}, -e^{-i\alpha}, -e^{i\alpha}, e^{-i\alpha}$, resp., where $\alpha = \arccos \sqrt{1/|a, b, c, d|} \in (0, \pi/2)$, see Figure 6. By Proposition 3.3 and the proof of Theorem 1.1, we have

$$(4.1) \quad d_1 = 2 \operatorname{arth} r \quad \text{and} \quad d_2 = 2 \operatorname{arth} r'$$

where $r = \cos \alpha$.

Therefore, by Theorem 1.1 and Theorem 1.2, we have

$$d_1 d_2 = 4(\operatorname{arth} r)(\operatorname{arth} r') \leq \left(2 \operatorname{arth}(\sqrt{2}/2) \right)^2$$

and

$$d_1 + d_2 = 2(\operatorname{arth} r + \operatorname{arth} r') \geq 4 \operatorname{arth}(\sqrt{2}/2).$$

The equalities hold if and only if $\alpha = \frac{\pi}{4}$, namely, $|a, b, c, d| = \frac{1}{\cos^2 \alpha} = 2$.

This completes the proof of Corollary 1.4. \square

Let G, G' be domains in $\overline{\mathbb{R}^n}$ and let $f : G \rightarrow G'$ be a homeomorphism. Then f is K -quasiconformal if

$$M(\Gamma)/K \leq M(f\Gamma) \leq KM(\Gamma)$$

for every curve family Γ in G , where $M(\Gamma)$ is the modulus of Γ , see [V, 10.9].

For $r \in (0, 1)$ and $K \geq 1$, we define the distortion function

$$\varphi_K(r) = \mu^{-1}(\mu(r)/K),$$

where $\mu(r)$ is the modulus of the planar Grötzsch ring, see [V, Exercise 5.61].

Lemma 4.2. [BV, Theorem 1.10] *Let $f : \mathbb{B}^2 \rightarrow \mathbb{B}^2$ be a K -quasiconformal mapping with $f\mathbb{B}^2 = \mathbb{B}^2$, and let ρ be the hyperbolic metric of \mathbb{B}^2 . Then*

$$\rho(f(x), f(y)) \leq A(K) \max\{\rho(x, y), \rho(x, y)^{1/K}\}$$

for all $x, y \in \mathbb{B}^2$, where $A(K) = 2\operatorname{arth}(\varphi_K(\operatorname{th}\frac{1}{2}))$ and

$$K \leq u(K-1) + 1 \leq \log(\operatorname{ch}(K\operatorname{arch}(e))) \leq A(K) \leq v(K-1) + K$$

with $u = \operatorname{arch}(e)\operatorname{th}(\operatorname{arch}(e)) > 1.5412$ and $v = \log(2(1 + \sqrt{1 - 1/e^2})) < 1.3507$. In particular, $A(1) = 1$.

Proof of Theorem 1.6. In the same way as in the proof of Theorem 1.1, we still assume that $v_a = 0$, v_b is on X , v_d is on Y and $v_c = te^{i\theta}$, $0 < t \leq 1$ and $0 < \theta < \frac{\pi}{2}$ (see Figure 1). Then

$$d_1 = \rho(0, v_b) = \operatorname{arth}(Lr) \quad \text{and} \quad d_2 = \rho(0, v_d) = \operatorname{arth}(Lr'),$$

where $0 < L \leq 1$ and $0 < r < 1$.

By Lemma 4.2, we have

$$\begin{aligned} D_1 D_2 &\leq \rho(f(0), f(v_b))\rho(f(0), f(v_d)) \\ &\leq A(K)^2 \max\{\rho(0, v_b), \rho(0, v_b)^{1/K}\} \cdot \max\{\rho(0, v_d), \rho(0, v_d)^{1/K}\} \\ &= A(K)^2 \max\{d_1, d_1^{1/K}\} \cdot \max\{d_2, d_2^{1/K}\}. \end{aligned}$$

(1) $0 < L \leq \frac{e^2-1}{e^2+1} \approx 0.761594$. This implies that $d_1 < 1$ and $d_2 < 1$. Then by Theorem 1.1,

$$D_1 D_2 \leq A(K)^2 (d_1 d_2)^{1/K} \leq A(K)^2 \left(\operatorname{arth} \left(\frac{\sqrt{2}}{2} L \right) \right)^{2/K}.$$

(2) $\frac{e^2-1}{e^2+1} < L \leq 1$.

Case 1. $\frac{0.761594}{L} \approx \frac{1}{L} \frac{e^2-1}{e^2+1} = r_L < r < 1$. This implies that $d_1 > 1$ and $d_2 < 1$. Then

$$d_1 d_2^{1/K} = \operatorname{arth}(Lr)(\operatorname{arth}(Lr'))^{1/K} \equiv F_{L,K}(r).$$

By logarithmic differentiation, we have

$$(4.3) \quad \frac{F'_{L,K}(r)}{F_{L,K}(r)} = \frac{Lr}{r'(1 - (Lr')^2)\operatorname{arth}(Lr')} \left(\frac{f_L(r)}{f_L(r')} - \frac{1}{K} \right),$$

where $f_L(r) = \frac{1-(Lr')^2}{r\operatorname{arth}(Lr)}$. By Lemma 2.9(1), $\frac{f_L(r)}{f_L(r')}$ is strictly decreasing from $(r_L, 1)$ onto $(0, \frac{1}{M_L})$. Here $M_L = \frac{f_L(r'_L)}{f_L(r_L)} > 1$ since $r_L \approx \frac{0.761594}{L} \geq 0.761594 > 0.707107 \approx \frac{\sqrt{2}}{2}$.

Case 1.1 $1 \leq K \leq M_L$. By (4.3), we have $F'_{L,K}(r) \leq 0$ and hence $F_{L,K}(r)$ is strictly decreasing on $(r_L, 1)$. Therefore,

$$(4.4) \quad d_1 d_2^{1/K} \leq F_{L,K}(r_L) = \operatorname{arth}(Lr_L)(\operatorname{arth}(Lr'_L))^{1/K}.$$

Case 1.2 $K > M_L$. There exists exactly one point $r_L(K) \in (r_L, 1)$ such that $\frac{f_L(r_L(K))}{f_L(\sqrt{1-r_L(K)^2})} = \frac{1}{K}$. Then $F_{L,K}$ is strictly increasing on $(r_L, r_L(K))$ and strictly decreasing on $(r_L(K), 1)$. Therefore,

$$(4.5) \quad d_1 d_2^{1/K} \leq F_{L,K}(r_L(K)) = \operatorname{arth}(Lr_L(K)) \left(\operatorname{arth} \left(L\sqrt{1-r_L(K)^2} \right) \right)^{1/K}.$$

Case 2. $\sqrt{1-r_L^2} < r < r_L$. This implies that $d_1 < 1$ and $d_2 < 1$. Since $\frac{\sqrt{2}}{2} \approx 0.707107 \in (\sqrt{1-r_L^2}, r_L)$, by Theorem 1.1 we have

$$(4.6) \quad (d_1 d_2)^{1/K} \leq \left(\operatorname{arth} \left(\frac{\sqrt{2}}{2} L \right) \right)^{2/K}.$$

Case 3. $0 < r < \sqrt{1-r_L^2}$. This implies that $d_1 < 1$ and $d_2 > 1$. Putting $p = r'$, we have $r_L < p < 1$ and

$$d_1^{1/K} d_2 = \operatorname{arth}(Lp)(\operatorname{arth}(Lp'))^{1/K}.$$

Hence Case 3 is the same as Case 1.

Therefore, by (4.4) and (4.6), we have if $1 \leq K \leq M_L$,

$$D_1 D_2 \leq A(K)^2 \max \left\{ \operatorname{arth}(Lr_L) \left(\operatorname{arth} \left(L\sqrt{1-r_L^2} \right) \right)^{1/K}, \left(\operatorname{arth} \left(\frac{\sqrt{2}}{2} L \right) \right)^{2/K} \right\}.$$

And by (4.5) and (4.6), we have if $K > M_L$,

$$D_1 D_2 \leq A(K)^2 \max \left\{ \operatorname{arth}(Lr_L(K)) \left(\operatorname{arth} \left(L\sqrt{1-r_L(K)^2} \right) \right)^{1/K}, \left(\operatorname{arth} \left(\frac{\sqrt{2}}{2} L \right) \right)^{2/K} \right\}.$$

This completes the proof of Theorem 1.6. \square

Proof of Corollary 1.7. First, let g be the same as in the proof of Corollary 1.4. Let $\{se_1\} = g(J^*[a, d]) \cap X$, $\{te_2\} = g(J^*[a, b]) \cap Y$ and denote

$$z_1 = g^{-1}(se_1), \quad z_2 = g^{-1}(te_2), \quad z_3 = g^{-1}(-se_1), \quad z_4 = g^{-1}(-te_2).$$

Since the hyperbolic distance is Möbius invariant, by the proof of Corollary 1.4, we get

$$d_1 = \rho(z_1, z_3) = 2\operatorname{arth} r \quad \text{and} \quad d_2 = \rho(z_2, z_4) = 2\operatorname{arth} r', \quad 0 < r < 1.$$

Then by Lemma 4.2, we have

$$\begin{aligned} D_1 D_2 &\leq \rho(f(z_1), f(z_3)) \rho(f(z_2), f(z_4)) \\ &\leq A(K)^2 \max\{\rho(z_1, z_3), \rho(z_1, z_3)^{1/K}\} \cdot \max\{\rho(z_2, z_4), \rho(z_2, z_4)^{1/K}\} \\ &= A(K)^2 \max\{d_1, d_1^{1/K}\} \cdot \max\{d_2, d_2^{1/K}\}. \end{aligned}$$

Case 1. $0.886819 \approx \frac{2\sqrt{e}}{e+1} = r_1 < r < 1$. This implies that $d_1 > 1$ and $d_2 < 1$. Then

$$d_1 d_2^{1/K} = 2^{1+1/K} \operatorname{arth} r (\operatorname{arth} r')^{1/K} \equiv 2^{1+1/K} F_{1,K}(r).$$

Let $M_1 = \frac{f_1(r_1)}{f_1(r)}$, where $f_1(r) = \frac{r}{\operatorname{arth} r}$. By the proof of Case 1 in Theorem 1.6, we have the following conclusions.

Case 1.1 $1 \leq K \leq M_1$.

$$(4.7) \quad d_1 d_2^{1/K} \leq 2^{1+1/K} F_{1,K}(r_1) = 2^{1+1/K} \operatorname{arth} r_1 (\operatorname{arth} r'_1)^{1/K}.$$

Case 1.2 $K > M_1$. There exists exactly one point $r_1(K) \in (r_1, 1)$ such that $\frac{f_1(r_1(K))}{f_1(\sqrt{1-r_1(K)^2})} = \frac{1}{K}$. Therefore,

$$(4.8) \quad d_1 d_2^{1/K} \leq 2^{1+1/K} F_{1,K}(r_1(K)) = 2^{1+1/K} \operatorname{arth} (r_1(K)) \left(\operatorname{arth} \left(\sqrt{1-r_1(K)^2} \right) \right)^{1/K}.$$

Case 2. $0.462117 \approx \frac{e-1}{e+1} = \sqrt{1-r_1^2} < r < r_1 \approx 0.886819$. This implies that $d_1 > 1$ and $d_2 > 1$. Since $\frac{\sqrt{2}}{2} \approx 0.707107 \in (\sqrt{1-r_1^2}, r_1)$, by Corollary 1.4 we have

$$(4.9) \quad d_1 d_2 \leq \left(2 \log(\sqrt{2} + 1)\right)^2.$$

Case 3. $0 < r < \sqrt{1-r_1^2} \approx 0.462117$. This implies that $d_1 < 1$ and $d_2 > 1$. Putting $p = r'$, we have $r_1 < p < 1$ and

$$d_1^{1/K} d_2 = 2^{1+1/K} \operatorname{arth} p (\operatorname{arth} p')^{1/K}.$$

Hence Case 3 is the same as Case 2.

Therefore, by (4.7) and (4.9), we have if $1 \leq K \leq M_1$,

$$D_1 D_2 \leq A(K)^2 \max \left\{ 2^{1+1/K} \operatorname{arth} r_1 (\operatorname{arth} r_1')^{1/K}, \left(2 \log(\sqrt{2} + 1)\right)^2 \right\}.$$

And by (4.8) and (4.9), we have if $K > M_1$,

$$D_1 D_2 \leq A(K)^2 \max \left\{ 2^{1+1/K} \operatorname{arth} (r_1(K)) \left(\operatorname{arth} \left(\sqrt{1-r_1(K)^2} \right) \right)^{1/K}, \left(2 \log(\sqrt{2} + 1)\right)^2 \right\}.$$

This completes the proof of Corollary 1.7. \square

Acknowledgments. The research of Matti Vuorinen was supported by the Academy of Finland, Project 2600066611. The research of Gendi Wang was supported by CIMO of Finland, Grant TM-10-7364.

REFERENCES

- [AVV1] G. D. ANDERSON, M. K. VAMANAMURTHY, AND M. VUORINEN, *Conformal Invariants, Inequalities and Quasiconformal Maps*, Canadian Mathematical Society Series of Monographs and Advanced Texts, a Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1997.
- [AVV2] G. D. ANDERSON, M. K. VAMANAMURTHY, AND M. VUORINEN, *Generalized convexity and inequalities*, J. Math. Anal. Appl. **335** (2007), 1294–1308.
- [AVZ] G. D. ANDERSON, M. VUORINEN, AND X.-H. ZHANG, *Topics in special functions III*, Analytic Number Theory, Approximation Theory and Special Functions, ed. by K. Alladi, G. Milovanovic, and M. Th. Rassias, Springer-Verlag, arXiv math 1209.1696 [math.CA].
- [A] J. W. ANDERSON, *Hyperbolic Geometry*, 2nd ed. Springer Undergraduate Mathematics Series, Springer-Verlag London, Ltd., London, 2005.
- [Ba] Á. BARICZ, *Convexity of the zero-balanced Gaussian hypergeometric functions with respect to Hölder means*, JIPAM. J. Inequal. Pure Appl. Math. **8** (2007), Article 40, 9 pp.
- [Be] A. F. BEARDON, *The Geometry of Discrete Groups*, Graduate Texts in Math., vol. **91**, Springer-Verlag, New York, 1983.
- [BM] A. F. BEARDON AND D. MINDA, *Conformal images and the angle of parallelism*, Comput. Methods Funct. Theory **12** (2012), 87–100.
- [BV] B. A. BHAYO AND M. VUORINEN, *On Mori's theorem for quasiconformal maps in the n -space*, Trans. Amer. Math. Soc. **363** (2011), 5703–5719.
- [Bu] P. S. BULLEN, *Handbook of Means and Their Inequalities*, Mathematics and its Applications, vol. **560**, Kluwer Academic Publishers Group, Dordrecht, 2003.
- [HLP] G. H. HARDY, J. E. LITTLEWOOD, AND G. PÓLYA, *Inequalities*, 2nd ed. Cambridge, at the University Press, 1952.
- [KL] L. KEEN AND N. LAKIC, *Hyperbolic Geometry from a Local Viewpoint*, London Mathematical Society Student Texts **68**, Cambridge University Press, Cambridge, 2007.
- [KV] R. KLÉN AND M. VUORINEN, *Apollonian circles and hyperbolic geometry*, J. Analysis **19** (2011), 41–60.

- [K] J. G. KRZYŻ, *Problems in Complex Variable Theory*, Translation of the 1962 Polish original, Modern Analytic and Computational Methods in Science and Mathematics, No. **36**, American Elsevier Publishing Co., Inc., New York; PWN-Polish Scientific Publishers, Warsaw, 1971.
- [V] M. VUORINEN, *Conformal Geometry and Quasiregular Mappings*, Lecture Notes in Mathematics **1319**, Springer-Verlag, Berlin, 1988.
- [WZJ1] G.-D. WANG, X.-H. ZHANG, AND Y.-P. JIANG, *Concavity with respect to Hölder mean involving generalized Grötzsch function*, J. Math. Anal. Appl. **379** (2011), 200–204.
- [WZJ2] G.-D. WANG, X.-H. ZHANG, AND Y.-P. JIANG, *Hölder concavity and inequalities of the Jacobian elliptic function*, Integral Transforms Spec. Funct. **23** (2012), 337–345.
- [ZWC] X.-H. ZHANG, G.-D. WANG, AND Y.-M. CHU, *Convexity with respect to Hölder mean involving zero-balanced hypergeometric functions*, J. Math. Anal. Appl. **353** (2009), 256–259.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TURKU, TURKU 20014, FINLAND
E-mail address: vuorinen@utu.fi

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TURKU, TURKU 20014, FINLAND
E-mail address: genwan@utu.fi