

# ON VOLUMES OF HYPERBOLIC COXETER POLYTOPES AND QUADRATIC FORMS

JOHN G. RATCLIFFE AND STEVEN T. TSCHANTZ

ABSTRACT. In this paper, we compute the covolume of the group of units of the quadratic form  $f_d^n(x) = x_1^2 + x_2^2 + \cdots + x_n^2 - dx_{n+1}^2$  with  $d$  an odd, positive, square-free integer. Mcleod has determined the hyperbolic Coxeter fundamental domain of the reflection subgroup of the group of units of the quadratic form  $f_3^n$ . We apply our covolume formula to compute the volumes of these hyperbolic Coxeter polytopes.

## 1. INTRODUCTION

In 1996, we computed the covolume of the group of units of the quadratic form

$$f_d^n(x) = x_1^2 + x_2^2 + \cdots + x_n^2 - dx_{n+1}^2$$

with  $d$  an odd, positive, square-free integer, in order to determine the volume of some hyperbolic Coxeter simplices, see [14] and §5 of [8]. We never got around to writing up our computation except for the case  $d = 1$  in [14]. Recently, several papers have been written concerning the group of units of the quadratic form  $f_3^n$ , and so we felt it was time to write up our more general computation. Our main result is Theorem 4 which gives an explicit formula for the covolume of the group of units of  $f_d^n$ .

J. Mcleod [10] has determined the reflection subgroup of the group of units of  $f_3^n$  and has shown that it has finite index if and only if  $n \leq 13$ . As an application of our computation, we determine the volume of the hyperbolic Coxeter fundamental domain of the reflection subgroup of the group of units of the quadratic form  $f_3^n$ , for  $n \leq 13$ , described in [10]. See Table 1 for the volumes of Mcleod's polytopes.

M. Belolipetsky and V. Emery [2] proved that for each odd dimension  $n \geq 5$ , there is a unique, orientable, noncompact, arithmetic, hyperbolic  $n$ -orbifold of smallest volume and determined its volume. V. Emery mentioned in [6] that for  $n \equiv 3 \pmod{4}$ , with  $n \geq 7$ , the corresponding arithmetic group is commensurable to the group of units of  $f_3^n$ . We determine the ratio between these two covolumes.

## 2. SIEGEL'S COVOLUME FORMULA FOR UNIT GROUPS

Let  $f$  be a quadratic form in  $n + 1$  real variables, with  $n \geq 2$ , that is equivalent over  $\mathbb{R}$  to the *Lorentzian quadratic form*

$$x_1^2 + \cdots + x_n^2 - x_{n+1}^2.$$

Let  $S$  be the matrix of the quadratic form  $f$ . We assume that all the entries of  $S$  are integers and  $d = |\det S|$  is an odd, square-free, positive integer.

The *group of units* of the form  $f$  is the group of all  $(n + 1) \times (n + 1)$  matrices  $A$ , with integral entries, such that  $A^t S A = S$ . A unit of  $f$  is said to be *positive* or *negative* according as  $A$  leaves invariant each of the two connected components

of  $\{x \in \mathbb{R}^{n+1} : x^t S x < 0\}$  or interchanges them. The group of positive units of  $f$  corresponds to a discrete group  $\Gamma$  of isometries of hyperbolic  $n$ -space

$$H^n = \{x \in E^{n+1} : x_1^2 + \cdots + x_n^2 - x_{n+1}^2 = -1 \text{ and } x_{n+1} > 0\}$$

under the equivalence of  $f$  with the Lorentzian quadratic form. For a discussion, see §1 of [9].

Let  $q$  be a positive integer and let  $\mathbb{Z}_q = \mathbb{Z}/q\mathbb{Z}$ . Let  $E_q(S)$  be the number of  $(n+1) \times (n+1)$  matrices  $A$  over  $\mathbb{Z}_q$  such that  $A^t S A = S$ . By Formula 82 in Siegel's paper [16], we have that

$$(1) \quad \text{vol}(H^n/\Gamma) = 4d^{\frac{n+2}{2}} \prod_{k=1}^n \pi^{-\frac{k}{2}} \Gamma\left(\frac{k}{2}\right) \cdot \lim_{q \rightarrow \infty} 2^{\omega(q)} \frac{q^{\frac{n(n+1)}{2}}}{E_q(S)}$$

where  $\Gamma(\frac{k}{2})$  is the gamma function evaluated at  $k/2$ , and  $\omega(q)$  is the number of distinct prime divisors of  $q$ , and  $q$  goes to infinity via the sequence  $2!, 3!, 4! \dots$ . In this paper a centered dot will delimit a product from the next factor.

We now evaluate the limit in Siegel's volume formula. Let  $q = p_1^{a_1} p_2^{a_2} \cdots p_r^{a_r}$  be the prime factorization of  $q$ . Now  $E_q(S)$  is a multiplicative function of  $q$  by Lemma 18 of Siegel's paper [15], and so we have

$$(2) \quad \frac{2^{\omega(q)} q^{\frac{n(n+1)}{2}}}{E_q(S)} = \prod_{i=1}^r \frac{2p_i^{\frac{a_i n(n+1)}{2}}}{E_{p_i^{a_i}}(S)}.$$

Henceforth  $p$  is a prime number. By Lemma 18 of [15], we have

$$(3) \quad \frac{2p^{\frac{an(n+1)}{2}}}{E_{p^a}(S)} = \begin{cases} \frac{2p^{\frac{3n(n+1)}{2}}}{E_{p^3}(S)} & \text{if } p \mid 2d \text{ and } a \geq 3, \\ \frac{2p^{\frac{n(n+1)}{2}}}{E_p(S)} & \text{if } p \nmid 2d \text{ and } a \geq 1. \end{cases}$$

Therefore, we have

$$(4) \quad \lim_{q \rightarrow \infty} 2^{\omega(q)} \frac{q^{\frac{n(n+1)}{2}}}{E_q(S)} = \prod_{p \mid 2d} \frac{2p^{\frac{3n(n+1)}{2}}}{E_{p^3}(S)} \prod_{p \nmid 2d} \frac{2p^{\frac{n(n+1)}{2}}}{E_p(S)}.$$

First assume that  $n$  is even and  $p \nmid 2d$ . Then by Lemma 18 of [15], we have that

$$(5) \quad \frac{2p^{\frac{n(n+1)}{2}}}{E_p(S)} = \prod_{k=1}^{\frac{n}{2}} (1 - p^{-2k})^{-1}.$$

Hence we have

$$(6) \quad \prod_{p \nmid 2d} \frac{2p^{\frac{n(n+1)}{2}}}{E_p(S)} = \prod_{k=1}^{\frac{n}{2}} \prod_{p \nmid 2d} (1 - p^{-2k})^{-1}.$$

Using the product formula for the Riemann zeta function  $\zeta(2k)$ , we have

$$(7) \quad \prod_{p \nmid 2d} \frac{2p^{\frac{n(n+1)}{2}}}{E_p(S)} = \prod_{k=1}^{\frac{n}{2}} \left( \prod_{p \nmid 2d} (1 - p^{-2k}) \right) \zeta(2k).$$

Now assume that  $n$  is odd and  $p \nmid 2d$ . Then by Lemma 18 of [15], we have that

$$(8) \quad \frac{2p^{\frac{n(n+1)}{2}}}{E_p(S)} = \left( 1 - \left( \frac{(-1)^{\frac{n-1}{2}} d}{p} \right) p^{-\left(\frac{n+1}{2}\right)} \right)^{-1} \prod_{k=1}^{\frac{n-1}{2}} (1 - p^{-2k})^{-1}$$

where  $(\pm d/p)$  is a Legendre symbol. Define a fundamental discriminant  $D$  by the formula

$$(9) \quad D = \begin{cases} (-1)^{\frac{n-1}{2}} d & \text{if } (-1)^{\frac{n-1}{2}} d \equiv 1 \pmod{4}, \\ 4(-1)^{\frac{n-1}{2}} d & \text{if } (-1)^{\frac{n-1}{2}} d \equiv 3 \pmod{4}. \end{cases}$$

For all odd primes  $p$ , we have

$$(10) \quad \left( \frac{(-1)^{\frac{n-1}{2}} d}{p} \right) = \left( \frac{D}{p} \right).$$

Hence we have

$$(11) \quad \prod_{p \nmid 2d} \frac{2p^{\frac{n(n+1)}{2}}}{E_p(S)} = \prod_{p \nmid 2d} \left( 1 - \left( \frac{D}{p} \right) p^{-\left(\frac{n+1}{2}\right)} \right)^{-1} \prod_{k=1}^{\frac{n-1}{2}} (1 - p^{-2k})^{-1}.$$

Consider the Dirichlet  $L$ -series

$$(12) \quad L(s, D) = \sum_{k=1}^{\infty} \left( \frac{D}{k} \right) k^{-s}$$

where  $(D/k)$  is a Kronecker symbol. This series converges absolutely for  $s > 1$ . The Kronecker symbol  $(D/k)$  is a completely multiplicative function of  $k$  by Theorem 1.4.9 in Cohen [4]. By Theorem 11.7 in Apostol [1], this  $L$ -function has the product formula

$$(13) \quad L(s, D) = \prod_p \left( 1 - \left( \frac{D}{p} \right) p^{-s} \right)^{-1}.$$

Using the product formulas for  $\zeta(2k)$  and  $L(\frac{n+1}{2}, D)$ , we obtain the formula

$$(14) \quad \prod_{p \nmid 2d} \frac{2p^{\frac{n(n+1)}{2}}}{E_p(S)} = \left( 1 - \left( \frac{D}{2} \right) 2^{-\left(\frac{n+1}{2}\right)} \right) L\left(\frac{n+1}{2}, D\right) \prod_{k=1}^{\frac{n-1}{2}} \left( \prod_{p \mid 2d} (1 - p^{-2k}) \right) \zeta(2k).$$

It remains only to compute  $E_{p^3}(S)$  for each prime number  $p$  dividing  $2d$ .

### 3. THE COMPUTATION OF $E_8(S)$

From now on, we assume that  $f = f_d^n$ . Then  $S$  is the  $(n+1) \times (n+1)$  diagonal matrix  $\text{diag}(1, \dots, 1, -d)$ . Let  $k$  be a positive integer, and let  $\mathbb{Z}_k = \mathbb{Z}/k\mathbb{Z}$ . Let  $O(n+1, \mathbb{Z}_k)$  be the group of  $(n+1) \times (n+1)$  matrices  $A$  over the ring  $\mathbb{Z}_k$  such that  $A^t A = I$ . Let  $J$  be the  $(n+1) \times (n+1)$  diagonal matrix  $\text{diag}(1, \dots, 1, -1)$ . Let  $O(n, 1; \mathbb{Z}_k)$  be the group of  $(n+1) \times (n+1)$  matrices  $A$  over the ring  $\mathbb{Z}_k$  such that  $A^t J A = J$ .

Let  $O(S, \mathbb{Z}_k)$  be the group of invertible  $(n+1) \times (n+1)$  matrices  $A$  over the ring  $\mathbb{Z}_k$  such that  $A^t S A = S$ . Then  $E_8(S)$  is the order of  $O(S, \mathbb{Z}_8)$ , since  $d$  is odd.

Let  $O^*(n+1, \mathbb{Z}_2)$  be the image of  $O(n+1, \mathbb{Z}_4)$  in  $O(n+1, \mathbb{Z}_2)$  under the homomorphism induced by the ring homomorphism from  $\mathbb{Z}_4$  to  $\mathbb{Z}_2$ , and let  $O^*(n, 1; \mathbb{Z}_2)$

be the image of  $O(n, 1; \mathbb{Z}_4)$  in  $O(n+1, \mathbb{Z}_2)$  under the homomorphism induced by the ring homomorphism from  $\mathbb{Z}_4$  to  $\mathbb{Z}_2$ .

We know that reduction modulo 2 maps  $O(S, \mathbb{Z}_8)$  into  $O^*(n, 1; \mathbb{Z}_2)$  if  $d \equiv 1 \pmod{4}$  and into  $O^*(n+1, \mathbb{Z}_2)$  if  $d \equiv -1 \pmod{4}$ , since the image of  $O(S, \mathbb{Z}_8)$  factors through  $O(S, \mathbb{Z}_4)$ . We next show that  $O(S, \mathbb{Z}_8)$  maps onto the appropriate  $O^*$  subgroup of  $O(n+1, \mathbb{Z}_2)$  under reduction modulo 2.

**Lemma 1.** *If  $d \equiv -1 \pmod{4}$ , then  $O(S, \mathbb{Z}_8)$  maps onto  $O^*(n+1, \mathbb{Z}_2)$  under reduction modulo 2.*

*Proof.* By the discussion on page 60 of [14], the group  $O^*(n+1, \mathbb{Z}_2)$  is generated by the permutation matrices and, if  $n+1 \geq 6$ , a matrix  $C$  which is the identity except for a lower corner  $6 \times 6$  block with all 0 diagonal and all 1 off-diagonal entries.

First assume that  $d \equiv -1 \pmod{8}$ . Then  $O(S, \mathbb{Z}_8) = O(n+1, \mathbb{Z}_8)$ . The permutation matrices in  $O^*(n+1, \mathbb{Z}_2)$  obviously lift to  $O(n+1, \mathbb{Z}_8)$ . If  $n+1 \geq 6$ , the matrix  $C$  lifts to  $O(n+1, \mathbb{Z}_8)$ , since the identity part of  $C$  lifts as is, and the  $6 \times 6$  block lifts to a block with all 2 diagonal and all 1 off-diagonal entries.

Now assume that  $d \equiv 3 \pmod{8}$ . First we argue that all the permutation matrices lift to  $O(S, \mathbb{Z}_8)$ . Clearly the permutations of the first  $n$  coordinates lift. Thus it suffices to show that the transposition of the last two coordinates also lifts, and the lift is the  $2 \times 2$  block with diagonal 2, 6 and all 1 off-diagonal entries, that is,

$$\begin{pmatrix} 2 & 1 \\ 1 & 6 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -3 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ 1 & 6 \end{pmatrix} \equiv \begin{pmatrix} 1 & 0 \\ 0 & -3 \end{pmatrix} \pmod{8}.$$

We also need our lower  $6 \times 6$  corner block to lift. The lift has diagonal entries 0, 0, 0, 0, 0, 4 and all 1 off-diagonal entries.  $\square$

**Lemma 2.** *If  $d \equiv 1 \pmod{4}$ , then  $O(S, \mathbb{Z}_8)$  maps onto  $O^*(n, 1; \mathbb{Z}_2)$  under reduction modulo 2.*

*Proof.* The proof for the case  $d \equiv 1 \pmod{8}$  follows from Lemma 4 in [14]. Now assume  $d \equiv -3 \pmod{8}$ . Identify  $O^*(n, \mathbb{Z}_2)$  with the subgroup of  $O^*(n, 1; \mathbb{Z}_2)$  fixing the last standard basis vector  $e_{n+1}$ . By the proof of Lemma 4 in [14], the group  $O^*(n, 1; \mathbb{Z}_2)$  is generated by the subgroup  $O^*(n, \mathbb{Z}_2)$  and a matrix  $C$  which is the identity except for a lower corner  $4 \times 4$  block with all 0 diagonal and all 1 off-diagonal entries. The subgroup  $O^*(n, \mathbb{Z}_2)$  lifts to  $O(S, \mathbb{Z}_8)$  as in the case  $d \equiv -1 \pmod{8}$  in Lemma 1. The  $4 \times 4$  lower corner block of  $C$  lifts to the  $4 \times 4$  block with diagonal 2, 2, 2, 4 and all 1 off-diagonal entries.  $\square$

**Lemma 3.** *Let  $\eta : O(S, \mathbb{Z}_8) \rightarrow O(S, \mathbb{Z}_2)$  be the homomorphism induced by reduction modulo 2. The kernel of  $\eta$  has order  $2^{n^2+2n+2}$ .*

*Proof.* Each matrix  $M$  in  $O(S, \mathbb{Z}_8)$  can be written uniquely in the form  $N+2U+4V$  where  $N, U$ , and  $V$  are zero-one matrices and we are interested in counting the  $M$  with  $N = I$ . Now observe that

$$M^t S M = S + 2(U^t S + S U) + 4(U^t S U + V^t S + S V).$$

Using the fact that  $4(-d) \equiv 4 \pmod{8}$ , the last equation simplifies to

$$M^t S M = S + 2(U^t S + S U) + 4(U^t U + V^t + V).$$

First assume that  $d \equiv 1 \pmod{4}$ . Then  $2(-d) \equiv 2(-1) \pmod{8}$  and the last equation simplifies to

$$M^t S M = S + 2(U^t J + J U) + 4(U^t U + V^t + V).$$

The proof now proceeds as in the proof of Theorem 5 of [14].

Now assume that  $d \equiv -1 \pmod{4}$ . Then  $2(-d) \equiv 2 \pmod{8}$  and the next to last equation simplifies to

$$M^t S M = S + 2(U^t + U) + 4(U^t U + V^t + V).$$

The proof now proceeds as in the proof of Theorem 5 of [14] with the simplification that  $W = 0$ .  $\square$

Let  $\epsilon_2(k) = 1$  if  $k$  is even and 0 if  $k$  is odd and define

$$(15) \quad \alpha(n) = \prod_{k=1}^n (2^k - \epsilon_2(k)).$$

Define

$$(16) \quad \beta(n) = 2^{n-2} + 2^{(n-2)/2} \cos(n\pi/4),$$

and

$$(17) \quad \gamma(n) = 2^n + 2^{n/2} \cos(n\pi/4).$$

Note that the function  $f(n) = \cos(n\pi/4)$  is periodic with period 8. The values of  $f(n)$  for  $n = 0, 1, \dots, 7$  are  $1, 2^{-1/2}, 0, -2^{-1/2}, -1, -2^{-1/2}, 0, 2^{-1/2}$ , respectively.

**Theorem 1.** *The value of  $E_8(S)$  is given by*

$$E_8(S) = \begin{cases} 2^{n^2+2n+2} \alpha(n) / \beta(n+1) & \text{if } d \equiv -1 \pmod{4}, \\ 2^{n^2+2n+2} \alpha(n) / \gamma(n-1) & \text{if } d \equiv 1 \pmod{4}. \end{cases}$$

*Proof.* By Lemmas 2 of [14], the order of  $O^*(n+1, \mathbb{Z}_2)$  is  $\alpha(n)/\beta(n+1)$ , and by Lemma 3 of [14], the order  $O^*(n, 1; \mathbb{Z}_2)$  is  $\alpha(n)/\gamma(n-1)$ . The Theorem now follows from Lemmas 1, 2, 3.  $\square$

**Corollary 1.** *We have that*

$$\frac{2 \cdot 2^{\frac{3n(n+1)}{2}}}{E_8(S)} = \frac{2^{\frac{n-1}{2}} + \cos((n + (-1)^{\frac{d+1}{2}})\pi/4)}{2^{\frac{n+3}{2}} \prod_{k=1}^{\lfloor \frac{n}{2} \rfloor} (1 - 2^{-2k})}.$$

*Proof.* By Theorem 1, we have that

$$\frac{2 \cdot 2^{\frac{3n(n+1)}{2}}}{E_8(S)} = \frac{2 \cdot 2^{\frac{3n(n+1)}{2}} (2^{n-1} + 2^{\frac{n-1}{2}} \cos((n + (-1)^{\frac{d+1}{2}})\pi/4))}{2^{n^2+2n+2} \prod_{k=1}^n (2^k - \epsilon_2(k))}.$$

If  $n$  is odd, we have that

$$\begin{aligned} \prod_{k=1}^n (2^k - \epsilon_2(k)) &= 2 \prod_{k=1}^{\frac{n-1}{2}} 2^{2k+1} (2^{2k} - 1) \\ &= 2 \prod_{k=1}^{\frac{n-1}{2}} 2^{4k+1} (1 - 2^{-2k}) = 2^{\frac{n(n+1)}{2}} \prod_{k=1}^{\frac{n-1}{2}} (1 - 2^{-2k}). \end{aligned}$$

If  $n$  is even, we have that

$$\begin{aligned} \prod_{k=1}^n (2^k - \epsilon_2(k)) &= \prod_{k=1}^{\frac{n}{2}} 2^{2k-1} (2^{2k} - 1) \\ &= \prod_{k=1}^{\frac{n}{2}} 2^{4k-1} (1 - 2^{-2k}) = 2^{\frac{n(n+1)}{2}} \prod_{k=1}^{\frac{n}{2}} (1 - 2^{-2k}). \end{aligned}$$

Therefore, we have that

$$\begin{aligned} \frac{2 \cdot 2^{\frac{3n(n+1)}{2}}}{E_8(S)} &= \frac{2 \cdot 2^{\frac{3n(n+1)}{2}} 2^{\frac{n-1}{2}} (2^{\frac{n-1}{2}} + \cos((n + (-1)^{\frac{d+1}{2}})\pi/4))}{2^{n^2+2n+2} 2^{\frac{n(n+1)}{2}} \prod_{k=1}^{\lfloor \frac{n}{2} \rfloor} (1 - 2^{-2k})} \\ &= \frac{2^{\frac{n-1}{2}} + \cos((n + (-1)^{\frac{d+1}{2}})\pi/4)}{2^{\frac{n+3}{2}} \prod_{k=1}^{\lfloor \frac{n}{2} \rfloor} (1 - 2^{-2k})}. \end{aligned}$$

□

#### 4. THE COMPUTATION OF $E_{p^3}(S)$

Let  $p$  be a prime number that divides the odd, positive, square-free integer  $d$ . In this section, we determine the value of  $E_{p^3}(S)$ .

Let  $a$  be a positive integer with  $a \geq 2$ , and let  $M \in \text{GL}(n+1, \mathbb{Z}_{p^a})$ . Define

$$(18) \quad \alpha(M) = \text{diag}(1, 1, \dots, 1, 0) \cdot M,$$

$$(19) \quad \beta(M) = \text{diag}(0, 0, \dots, 0, 1) \cdot M.$$

Then  $\alpha(M)$  is the matrix  $M$  with its last row set to zero, and  $\beta(M)$  is  $M$  with its first  $n$  rows set to zero. Note that  $M = \alpha(M) + \beta(M)$ .

Define  $\bar{\beta}(M) = \beta(M)$  reduced modulo  $p^{a-1}$  with entries in the range  $[0, p^{a-1})$ . Define  $\bar{M} = \alpha(M) + \bar{\beta}(M)$ . Then  $M - \bar{M} = p^{a-1}U$  for  $U = \beta(U)$  with entries in the range  $[0, p)$ . Define

$$(20) \quad \bar{O}(S, \mathbb{Z}_{p^a}) = \{\bar{M} : M \in O(S, \mathbb{Z}_{p^a})\}.$$

**Lemma 4.** *Let  $M \in \text{GL}(n+1, \mathbb{Z}_{p^a})$  with  $a \geq 2$ . Then  $M \in O(S, \mathbb{Z}_{p^a})$  if and only if  $\bar{M} \in O(S, \mathbb{Z}_{p^a})$ . Moreover*

$$E_{p^a}(S) = |O(S, \mathbb{Z}_{p^a})| = p^{n+1} |\bar{O}(S, \mathbb{Z}_{p^a})|.$$

*Proof.* Let  $M$  be an  $(n+1) \times (n+1)$  matrix over  $\mathbb{Z}$  such that  $M^t S M \equiv S \pmod{p^a}$ . Then  $(\det(M))^2(-d) \equiv -d \pmod{p^a}$ . Hence  $(\det(M))^2 \equiv 1 \pmod{p^{a-1}}$ , and so  $\det(M)$  is invertible modulo  $p^a$ . Hence the reduction of  $M$  modulo  $p^a$  is invertible. Therefore  $E_{p^a}(S) = |O(S, \mathbb{Z}_{p^a})|$ .

Now suppose  $M$  is an  $(n+1) \times (n+1)$  matrix over  $\mathbb{Z}_{p^a}$ . Observe that

$$\begin{aligned} M^t S M &= (\bar{M} + p^{a-1}U)^t S (\bar{M} + p^{a-1}U) \\ &= \bar{M}^t S \bar{M} + p^{a-1}U^t S \bar{M} + p^{a-1} \bar{M}^t S U + p^{2a-2}U^t S U \\ &= \bar{M}^t S \bar{M} + p^{a-1}(-d)U^t \bar{M} + p^{a-1}(-d)\bar{M}^t U \\ &= \bar{M}^t S \bar{M}. \end{aligned}$$

Thus  $M \in \mathrm{O}(S, \mathbb{Z}_{p^a})$  if and only if  $\overline{M} \in \mathrm{O}(S, \mathbb{Z}_{p^a})$ . In the equation  $M - \overline{M} = p^{a-1}U$ , there are  $p^{n+1}$  choices for  $U$ . Hence for each  $\overline{M}$  there are  $p^{n+1}$  choices for  $M$ . Therefore  $|\mathrm{O}(S, \mathbb{Z}_{p^a})| = p^{n+1}|\overline{\mathrm{O}}(S, \mathbb{Z}_{p^a})|$ .  $\square$

**Lemma 5.** *If  $a \geq 2$ , we have that  $|\overline{\mathrm{O}}(S, \mathbb{Z}_{p^{a+1}})| = p^{\frac{n(n+1)}{2}}|\overline{\mathrm{O}}(S, \mathbb{Z}_{p^a})|$ .*

*Proof.* If  $M \in \overline{\mathrm{O}}(S, \mathbb{Z}_{p^{a+1}})$ , then  $N = \overline{(M \bmod p^a)} \in \overline{\mathrm{O}}(S, \mathbb{Z}_{p^a})$ . We want to count

$$M = N + p^a \alpha(U) + p^{a-1} \beta(U)$$

for  $U \bmod p$  and  $N \in \overline{\mathrm{O}}(S, \mathbb{Z}_{p^a})$  such that  $M \in \overline{\mathrm{O}}(S, \mathbb{Z}_{p^{a+1}})$ . Observe that

$$\begin{aligned} M^t S M - S &= N^t S N - S \\ &+ p^a (\alpha(U)^t S N + N^t S \alpha(U)) \\ &+ p^{a-1} (\beta(U)^t S N + N^t S \beta(U)) \\ &+ p^{2a-1} (\alpha(U)^t S \beta(U) + \beta(U)^t S \alpha(U)) \\ &+ p^{2a} (\alpha(U)^t S \alpha(U)) + p^{2a-2} (\beta(U)^t S \beta(U)) \\ &= N^t S N - S \\ &+ p^a (\alpha(U)^t N + N^t \alpha(U)) \\ &+ p^a (-d/p) (\beta(U)^t N + N^t \beta(U)). \end{aligned}$$

Let  $V = \alpha(U) + (-d/p)\beta(U)$ . Then  $\alpha(U) = \alpha(V)$  and  $\beta(U) \equiv (-d/p)^{-1}\beta(V) \bmod p$ .

Now  $M^t S M = S$  if and only if

$$V^t N + N^t V \equiv \frac{S - N^t S N}{p^a} \bmod p$$

where we regard the entries of  $N$  to be integers in the range  $[0, p^a)$ . Let  $W = N^t V \bmod p$ . Then  $V = (N^t)^{-1}W \bmod p$ . Note that  $N^t$  is invertible mod  $p$ , since  $N$  is invertible mod  $p$  by Lemma 4. Now  $M^t S M = S$  if and only if

$$W^t + W \equiv \frac{S - N^t S N}{p^a} \bmod p.$$

This congruence determines the diagonal of  $W$  and below diagonal entries in terms of arbitrary above diagonal entries for  $p^{n(n+1)/2}$  choices for  $W$  with entries mod  $p$ . Thus we have  $p^{n(n+1)/2}$  choices for  $U$  and as many  $M \in \overline{\mathrm{O}}(S, \mathbb{Z}_{p^{a+1}})$  for each  $N \in \overline{\mathrm{O}}(S, \mathbb{Z}_{p^a})$ .  $\square$

**Lemma 6.** *We have that  $|\overline{\mathrm{O}}(S, \mathbb{Z}_{p^2})| = 2p^{\frac{n(n+1)}{2}}|\mathrm{O}(n, \mathbb{Z}_p)|$ .*

*Proof.* Let  $M \in \overline{\mathrm{O}}(S, \mathbb{Z}_{p^2})$  and decompose  $M$  into blocks

$$M = \left( \begin{array}{c|c} M_0 & v \\ \hline w^t & x \end{array} \right)$$

with  $v \in \mathbb{Z}_{p^2}^n$ ,  $w \in \mathbb{Z}_p^n$  and  $x \in \mathbb{Z}_p$ .

Observe that  $M^t S M = S$  if and only if

$$\begin{cases} M_0^t M_0 - d w w^t &\equiv I \bmod p^2, \\ M_0^t v - d x w &\equiv 0 \bmod p^2, \\ v^t v - d x^2 &\equiv -d \bmod p^2. \end{cases}$$

The above system of congruences implies that

$$\begin{cases} M_0^t M_0 \equiv I \pmod{p}, \\ v \equiv 0 \pmod{p}, \\ x^2 \equiv 1 \pmod{p}. \end{cases}$$

Let  $N_0 = (M_0 \pmod{p})$ . Then  $M_0 = N_0 + pU_0$  for  $U_0 \pmod{p}$  and  $N_0 \in \mathrm{O}(n, \mathbb{Z}_p)$ . For each such  $N_0$  choose  $x = \pm 1 \pmod{p}$  and  $w \in \mathbb{Z}_p^n$  arbitrarily. We need to count the  $U_0$  and find  $v$  so that  $M \in \overline{\mathrm{O}}(S, \mathbb{Z}_{p^2})$ . Note that  $v = py$  where  $M_0^t y \equiv (d/p)xw \pmod{p}$ , and so  $v$  is determined from  $x, w$  and  $N_0$ . Thus it suffices to count  $U_0$  such that

$$(N_0 + pU_0)^t (N_0 + p^b U_0) - dww^t \equiv I \pmod{p^2}.$$

The above congruence is equivalent to

$$U_0^t N_0 + N_0^t U_0 \equiv \frac{I - N_0^t N_0}{p} + (d/p)ww^t \pmod{p}.$$

Let  $V_0 = N_0^t U_0$ . Then  $U_0 = (N_0^t)^{-1} V_0$ . The congruence

$$V_0^t + V_0 \equiv \frac{I - N_0^t N_0}{p} + (d/p)ww^t \pmod{p}$$

determines the diagonal of  $V_0$  and below diagonal entries from the above diagonal entries. Hence there are  $p^{n(n-1)/2}$  choices for  $V_0$  and so for  $U_0$  given  $w$  and  $x$ . Thus there are  $2p^{n(n-1)/2+n}$  possibilities for  $M \in \overline{\mathrm{O}}(S, \mathbb{Z}_{p^2})$  for each  $N_0 \in \mathrm{O}(n, \mathbb{Z}_p)$ .  $\square$

**Theorem 2.** *The values of  $E_{p^2}(S)$  and  $E_{p^3}(S)$  are given by*

$$\begin{aligned} E_{p^2}(S) &= 2p^{\frac{(n+1)(n+2)}{2}} |\mathrm{O}(n, \mathbb{Z}_p)|, \\ E_{p^3}(S) &= p^{\frac{n(n+1)}{2}} E_{p^2}(S). \end{aligned}$$

*Proof.* By Lemmas 4 and 6, we have that

$$\begin{aligned} E_{p^2}(S) &= p^{n+1} |\overline{\mathrm{O}}(S, \mathbb{Z}_{p^2})| \\ &= p^{n+1} 2p^{\frac{n(n+1)}{2}} |\mathrm{O}(n, \mathbb{Z}_p)| = 2p^{\frac{(n+1)(n+2)}{2}} |\mathrm{O}(n, \mathbb{Z}_p)|. \end{aligned}$$

By Lemmas 4 and 5, we have that

$$\begin{aligned} E_{p^3}(S) &= p^{n+1} |\overline{\mathrm{O}}(S, \mathbb{Z}_{p^3})| \\ &= p^{n+1} p^{\frac{n(n+1)}{2}} |\mathrm{O}(S, \mathbb{Z}_{p^2})| = p^{\frac{n(n+1)}{2}} E_{p^2}(S). \quad \square \end{aligned}$$

**Corollary 2.** *We have that*

$$\frac{2p^{\frac{3n(n+1)}{2}}}{E_{p^3}(S)} = \frac{2p^{\frac{2n(n+1)}{2}}}{E_{p^2}(S)} = \frac{p^{\frac{(n+1)(n-2)}{2}}}{|\mathrm{O}(n, \mathbb{Z}_p)|}.$$

We need the following classical theorem, see Theorem 172 of Dickson [5].

**Theorem 3.** *If  $p$  be an odd prime number, then the order of  $\mathrm{O}(n, \mathbb{Z}_p)$  is given by*

$$|\mathrm{O}(n, \mathbb{Z}_p)| = \begin{cases} 2 \prod_{k=1}^{n-1} p^k - \epsilon_2(k) & n \text{ odd}, \\ 2p^{\frac{n}{2}-1} \left( p^{\frac{n}{2}} - \left( \frac{-1}{p} \right)^{\frac{n}{2}} \right) \prod_{k=1}^{n-2} p^k - \epsilon_2(k) & n \text{ even}, \end{cases}$$

where  $(-1/p)$  is a Legendre symbol.



**Corollary 3.** *We have that*

$$\frac{2p^{\frac{3n(n+1)}{2}}}{E_{p^3}(S)} = \frac{p^{\frac{n}{2}} + \epsilon_2(n) \left(\frac{-1}{p}\right)^{\frac{n}{2}}}{2p^{\frac{n+2}{2}} \prod_{k=1}^{\lfloor \frac{n}{2} \rfloor} (1 - p^{-2k})}.$$

*Proof.* First assume that  $n$  is odd. By Theorem 3 and Corollary 2, we have that

$$\begin{aligned} \frac{2p^{\frac{3n(n+1)}{2}}}{E_{p^3}(S)} &= \frac{p^{\frac{(n+1)(n-2)}{2}}}{\frac{n-1}{2 \prod_{k=1} p^k - \epsilon_2(k)}} \\ &= \frac{p^{\frac{(n+1)(n-2)}{2}}}{\frac{\frac{n-1}{2}}{2 \prod_{k=1} p^{2k-1} (p^{2k} - 1)}} \\ &= \frac{p^{\frac{(n+1)(n-2)}{2}}}{\frac{\frac{n-1}{2}}{2 \prod_{k=1} p^{4k-1} (1 - p^{-2k})}} \\ &= \frac{p^{\frac{(n+1)(n-2)}{2}}}{2p^{\frac{(n-1)n}{2}} \prod_{k=1} (1 - p^{-2k})} = \frac{1}{2p^{\frac{n-1}{2}} \prod_{k=1} (1 - p^{-2k})}. \end{aligned}$$

Now assume that  $n$  is even. By Theorem 3 and Corollary 2, we have that

$$\begin{aligned} \frac{2p^{\frac{3n(n+1)}{2}}}{E_{p^3}(S)} &= \frac{p^{\frac{(n+1)(n-2)}{2}}}{2p^{\frac{n}{2}-1} \left(p^{\frac{n}{2}} - \left(\frac{-1}{p}\right)^{\frac{n}{2}}\right) \prod_{k=1}^{n-2} p^k - \epsilon_2(k)} \\ &= \frac{p^{\frac{(n+1)(n-2)}{2}} \left(p^{\frac{n}{2}} + \left(\frac{-1}{p}\right)^{\frac{n}{2}}\right)}{2p^{\frac{n}{2}-1} (p^n - 1) \prod_{k=1}^{n-2} p^k - \epsilon_2(k)} \\ &= \frac{p^{\frac{(n+1)(n-2)}{2}} \left(p^{\frac{n}{2}} + \left(\frac{-1}{p}\right)^{\frac{n}{2}}\right)}{2p^{-\frac{n}{2}} \prod_{k=1}^n p^k - \epsilon_2(k)} \\ &= \frac{p^{\frac{(n+1)(n-2)}{2}} \left(p^{\frac{n}{2}} + \left(\frac{-1}{p}\right)^{\frac{n}{2}}\right)}{2p^{-\frac{n}{2}} p^{\frac{n(n+1)}{2}} \prod_{k=1}^{\frac{n}{2}} (1 - p^{-2k})} = \frac{p^{\frac{n}{2}} + \left(\frac{-1}{p}\right)^{\frac{n}{2}}}{2p^{\frac{n+2}{2}} \prod_{k=1}^{\frac{n}{2}} (1 - p^{-2k})}. \quad \square \end{aligned}$$

5. THE COVOLUME OF THE GROUP OF UNITS OF  $f_d^n$ 

Let  $\Gamma_d^n$  be the discrete group of isometries of hyperbolic  $n$ -space  $H^n$  that corresponds to the group of positive units of the quadratic form  $f_d^n$ . In this section, we give an explicit formula for  $\text{vol}(H^n/\Gamma_d^n)$ .

From Formula (6) of [14], we have

$$(21) \quad \prod_{k=1}^n \pi^{-\frac{k}{2}} \Gamma\left(\frac{k}{2}\right) = \begin{cases} \prod_{k=1}^{\frac{n-1}{2}} \frac{2(2k-1)!}{(2\pi)^{2k}} & n \text{ odd,} \\ \prod_{k=1}^{\frac{n}{2}} \frac{2(2k-1)!}{(2\pi)^{2k}} \cdot \frac{(2\pi)^{\frac{n}{2}}}{(n-1)!!} & n \text{ even.} \end{cases}$$

By Theorems 12.17 and 12.18 of [1], we have that

$$(22) \quad \zeta(2k) = \frac{(2\pi)^{2k} |B_{2k}|}{2(2k)!}$$

for every positive integer  $k$  where  $B_{2k}$  is the  $(2k)$ th Bernoulli number.

Define a function  $B$  of  $n$  by the formula

$$(23) \quad B = \prod_{k=1}^{\lfloor \frac{n}{2} \rfloor} \frac{|B_{2k}|}{2k}.$$

Then we have that

$$(24) \quad \prod_{k=1}^n \pi^{-\frac{k}{2}} \Gamma\left(\frac{k}{2}\right) \prod_{k=1}^{\lfloor \frac{n}{2} \rfloor} \zeta(2k) = \begin{cases} B & n \text{ odd,} \\ B \cdot \frac{(2\pi)^{\frac{n}{2}}}{(n-1)!!} & n \text{ even.} \end{cases}$$

Define a function  $C$  of  $n$  and  $d$  by the formula

$$(25) \quad C = \cos\left((n + (-1)^{\frac{d+1}{2}})\pi/4\right).$$

**Theorem 4.** *Let  $d$  be an odd, square-free, positive integer, and let  $\Gamma_d^n$  be the discrete group of isometries of hyperbolic  $n$ -space  $H^n$  corresponding to the group of positive units of the quadratic form  $f_d^n$ . The volume of  $H^n/\Gamma_d^n$  is given by*

$$\text{vol}(H^n/\Gamma_d^n) = \begin{cases} \frac{d^{\frac{n-1}{2}} B}{2^{n+\omega(d)}} (2^{\frac{n-1}{2}} + C) (2^{\frac{n+1}{2}} - \frac{D}{2}) \sqrt{d} \cdot L\left(\frac{n+1}{2}, D\right) & n \text{ odd,} \\ \frac{B}{2^{\frac{n}{2}+\omega(d)}} (2^{\frac{n}{2}} + 2^{\frac{1}{2}} C) \prod_{p|d} \left(p^{\frac{n}{2}} + \left(\frac{-1}{p}\right)^{\frac{n}{2}}\right) \cdot \frac{(2\pi)^{\frac{n}{2}}}{(n-1)!!} & n \text{ even.} \end{cases}$$

*Proof.* First assume that  $n$  is odd. From Formulas (1), (4), (14), (24) and Corollaries 1 and 3, we have that

$$\begin{aligned}
 \text{vol}(H^n/\Gamma_d^n) &= 4d^{\frac{n+2}{2}} \prod_{k=1}^n \pi^{-\frac{k}{2}} \Gamma\left(\frac{k}{2}\right) \cdot \lim_{q \rightarrow \infty} 2^{\omega(q)} \frac{q^{\frac{n(n+1)}{2}}}{E_q(S)} \\
 &= 4d^{\frac{n+2}{2}} \prod_{k=1}^n \pi^{-\frac{k}{2}} \Gamma\left(\frac{k}{2}\right) \cdot \prod_{p|2d} \frac{2p^{\frac{3n(n+1)}{2}}}{E_{p^3}(S)} \cdot \prod_{p \nmid 2d} \frac{2p^{\frac{n(n+1)}{2}}}{E_p(S)} \\
 &= 4d^{\frac{n+2}{2}} B \prod_{p|2d} \frac{2p^{\frac{3n(n+1)}{2}}}{E_{p^3}(S)} \left(1 - \left(\frac{D}{2}\right) 2^{-\left(\frac{n+1}{2}\right)}\right) L\left(\frac{n+1}{2}, D\right) \prod_{k=1}^{\frac{n-1}{2}} \prod_{p|2d} (1 - p^{-2k}) \\
 &= 4d^{\frac{n+2}{2}} B \frac{\left(2^{\frac{n-1}{2}} + C\right)}{2^{\frac{n+3}{2}} 2^{\omega(d)} d} \left(1 - \left(\frac{D}{2}\right) 2^{-\left(\frac{n+1}{2}\right)}\right) L\left(\frac{n+1}{2}, D\right) \\
 &= \frac{d^{\frac{n-1}{2}} B}{2^{n+\omega(d)}} \left(2^{\frac{n-1}{2}} + C\right) \left(2^{\frac{n+1}{2}} - \left(\frac{D}{2}\right)\right) \sqrt{d} \cdot L\left(\frac{n+1}{2}, D\right).
 \end{aligned}$$

Now assume that  $n$  is even. From Formulas (1), (4), (7), (24) and Corollaries 1 and 3, we have that

$$\begin{aligned}
 \text{vol}(H^n/\Gamma_d^n) &= 4d^{\frac{n+2}{2}} \prod_{k=1}^n \pi^{-\frac{k}{2}} \Gamma\left(\frac{k}{2}\right) \cdot \prod_{p|2d} \frac{2p^{\frac{3n(n+1)}{2}}}{E_{p^3}(S)} \cdot \prod_{p \nmid 2d} \frac{2p^{\frac{n(n+1)}{2}}}{E_p(S)} \\
 &= 4d^{\frac{n+2}{2}} B \cdot \frac{(2\pi)^{\frac{n}{2}}}{(n-1)!!} \prod_{p|2d} \frac{2p^{\frac{3n(n+1)}{2}}}{E_{p^3}(S)} \cdot \prod_{k=1}^{\frac{n}{2}} \prod_{p|2d} (1 - p^{-2k}) \\
 &= 4d^{\frac{n+2}{2}} B \cdot \frac{(2\pi)^{\frac{n}{2}}}{(n-1)!!} \frac{\left(2^{\frac{n-1}{2}} + C\right)}{2^{\frac{n+3}{2}}} \prod_{p|d} \frac{\left(p^{\frac{n}{2}} + \left(\frac{-1}{p}\right)^{\frac{n}{2}}\right)}{2p^{\frac{n+2}{2}}} \\
 &= \frac{B}{2^{\frac{n}{2}+\omega(d)}} \left(2^{\frac{n}{2}} + 2^{\frac{1}{2}} C\right) \prod_{p|d} \left(p^{\frac{n}{2}} + \left(\frac{-1}{p}\right)^{\frac{n}{2}}\right) \cdot \frac{(2\pi)^{\frac{n}{2}}}{(n-1)!!}.
 \end{aligned}$$

□

**Corollary 4.** *If  $n$  is odd, then the Euler characteristic of  $\Gamma_d^n$  is zero. If  $n$  is even, then the Euler characteristic of  $\Gamma_d^n$  is given by*

$$\chi(\Gamma_d^n) = \frac{(-1)^{\frac{n}{2}} B}{2^{\frac{n}{2}+\omega(d)}} \left(2^{\frac{n}{2}} + 2^{\frac{1}{2}} C\right) \prod_{p|d} \left(p^{\frac{n}{2}} + \left(\frac{-1}{p}\right)^{\frac{n}{2}}\right).$$

*Proof.* The group  $\Gamma_d^n$  is finitely generated, and so  $\Gamma_d^n$  has a torsion-free subgroup of finite index by Selberg's Lemma. If  $n$  is odd, then  $\chi(\Gamma_d^n) = 0$ , since odd dimensional hyperbolic manifolds of finite volume have zero Euler characteristic. If  $n$  is even, then the Gauss-Bonnet Theorem implies that

$$(26) \quad \text{vol}(H^n/\Gamma_d^n) = (-1)^{\frac{n}{2}} \chi(\Gamma_d^n) \cdot \frac{(2\pi)^{\frac{n}{2}}}{(n-1)!!}.$$

The result now follows immediately from Theorem 4. □

Using the formulas in Theorem 4, we can easily compute the volume of  $H^n/\Gamma_d^n$ . For example, we have

$$\begin{aligned}\text{vol}(H^3/\Gamma_3^3) &= \frac{5\sqrt{3}}{64}L(2, -3), \\ \text{vol}(H^3/\Gamma_7^3) &= \frac{7\sqrt{7}}{64}L(2, -7), \\ \text{vol}(H^5/\Gamma_5^5) &= \frac{15\sqrt{5}}{2048}L(3, 5), \\ \text{vol}(H^7/\Gamma_7^7) &= \frac{49\sqrt{7}}{98304}L(4, -7).\end{aligned}$$

These four volumes were reported on pages 344 and 345 of [8], with some sign differences due to different definitions of  $\Gamma_d^n$  and  $L(s, D)$ . In [8], the group  $\Gamma_d^n$  was denoted by  $\Gamma_{-d}^n$ , and  $L$ -functions were defined via Legendre symbols rather than Kronecker symbols.

## 6. THE VOLUMES OF MCLEOD'S POLYTOPES

J. Mcleod [10] has determined the reflection subgroup of  $\Gamma_3^n$  and has shown that it has finite index if and only if  $n \leq 13$ . Let  $P^n$  be the hyperbolic Coxeter fundamental domain of the reflection subgroup of  $\Gamma_3^n$  determined by Mcleod in [10]. Mcleod showed that the reflection subgroup of  $\Gamma_3^n$  has index equal to the order of the symmetry group  $\text{Sym}(P^n)$  of  $P^n$ ; moreover, Mcleod determined  $\text{Sym}(P)$  for  $n \leq 13$ . Hence, for  $n \leq 13$ , we have that

$$(27) \quad \text{vol}(P^n) = |\text{Sym}(P^n)|\text{vol}(H^n/\Gamma_3^n).$$

Table 1 lists the volumes of  $H^n/\Gamma_3^n$  and  $P^n$  for  $n \leq 13$  computed using our formulas.

The polygon  $P^2$  is  $30^\circ - 45^\circ$  right triangle, and so we can compute the area of  $P^2$  by the classical angle defect formula

$$\text{vol}(P^2) = \pi - \left(\frac{\pi}{2} - \frac{\pi}{4} - \frac{\pi}{6}\right) = \frac{\pi}{12}.$$

The polyhedron  $P^3$  is an orthotetrahedron with angles  $\frac{\pi}{6}, \frac{\pi}{3}, \frac{\pi}{4}$ . By Theorem 10.4.5 and the duplication formulas 10.4.9 and 10.4.10 in [13], we have that

$$\text{vol}(P^3) = \frac{5}{16}\mathcal{J}\left(\frac{\pi}{3}\right)$$

where  $\mathcal{J}(\theta)$  is the Lobachevsky function. By Formula 2 in Milnor [11], we have that

$$(28) \quad \mathcal{J}(\theta) = \frac{1}{2} \sum_{k=1}^{\infty} \sin(2k\theta)k^{-2}.$$

Observe that

$$(29) \quad \sin(2k\pi/3) = \frac{\sqrt{3}}{2} \left(\frac{-3}{k}\right).$$

Hence we have that

$$\text{vol}(P^3) = \frac{5\sqrt{3}}{64}L(2, -3).$$

$n$	$\text{vol}(\Gamma_3^n)$	$ \text{Sym}(P^n) $	$\text{vol}(P^n)$	$ \text{vol}(P^n) $
2	$\frac{\pi}{12}$	1	$\frac{\pi}{12}$	$2.617993878 * 10^{-1}$
3	$\frac{5\sqrt{3}}{64}L(2, -3)$	1	$\frac{5\sqrt{3}}{64}L(2, -3)$	$1.057230840 * 10^{-1}$
4	$\frac{\pi^2}{288}$	1	$\frac{\pi^2}{288}$	$3.426945973 * 10^{-2}$
5	$\frac{\sqrt{3}}{320}L(3, 12)$	1	$\frac{\sqrt{3}}{320}L(3, 12)$	$5.358748797 * 10^{-3}$
6	$\frac{13\pi^3}{604800}$	1	$\frac{13\pi^3}{604800}$	$6.664708943 * 10^{-4}$
7	$\frac{51\sqrt{3}}{1146880}L(4, -3)$	1	$\frac{51\sqrt{3}}{1146880}L(4, -3)$	$7.240232999 * 10^{-5}$
8	$\frac{697\pi^4}{9144576000}$	1	$\frac{697\pi^4}{9144576000}$	$7.424525364 * 10^{-6}$
9	$\frac{L(5,12)}{716800\sqrt{3}}$	1	$\frac{L(5,12)}{716800\sqrt{3}}$	$8.051559421 * 10^{-7}$
10	$\frac{341\pi^5}{987614208000}$	2	$\frac{341\pi^4}{493807104000}$	$2.113228256 * 10^{-7}$
11	$\frac{403L(6,-3)}{12918456320\sqrt{3}}$	2	$\frac{403L(6,-3)}{6459228160\sqrt{3}}$	$3.546550442 * 10^{-8}$
12	$\frac{50443\pi^6}{12428137193472000}$	2	$\frac{50443\pi^6}{6214068596736000}$	$7.804122909 * 10^{-9}$
13	$\frac{691L(7,12)}{344408064000\sqrt{3}}$	4	$\frac{691L(7,12)}{86102016000\sqrt{3}}$	$4.633381297 * 10^{-9}$

TABLE 1. The volumes of Mcleod’s hyperbolic Coxeter polytopes

For even  $n$ , we can compute the volume of the polytope  $P^n$  using the Gauss-Bonnet Theorem which implies that

$$(30) \quad \text{vol}(P^n) = (-1)^{\frac{n}{2}} \chi(P^n) \cdot \frac{(2\pi)^{\frac{n}{2}}}{(n-1)!!}$$

where  $\chi(P^n)$  is the Euler characteristic of the Coxeter group  $P^n$  defined by  $P^n$ . By Proposition 3 of Chiswell [3], we have that

$$(31) \quad \chi(P^n) = \sum_{\Delta} \frac{(-1)^{|\Delta|}}{|C_{\Delta}|}$$

where the sum is over all subgraphs  $\Delta$  of the Coxeter graph of  $P^n$  such that the Coxeter group  $C_{\Delta}$  defined by  $\Delta$  is finite. Here  $|\Delta|$  is the number of vertices of  $\Delta$ . From the Coxeter graph of  $P^n$  given in Mcleod [10], we computed the volumes of  $P^n$  for all even  $n \leq 12$  using Chiswell’s Euler characteristic formula and the Gauss-Bonnet Theorem. The volumes agree with the volumes given in Table 1.

For odd  $n$ , we computed the volume of  $P^n$  by numerical integration. We found very close agreement with the numerical values in Table 1 in dimensions 3, 5, 7, 9. We found close agreement in dimensions 11 and 13, but the accuracy deteriorated to a 1% discrepancy in dimension 11 and a 3% discrepancy in dimension 13, which is not bad considering we numerically integrated an  $n$ -fold integral for  $n = 11, 13$ . After the agreement of all these alternate calculations of the volume of  $P^n$  for all  $n \leq 13$ , we are confident with the correctness of our volume formulas derived using Siegel’s analytic theory of quadratic forms.

## 7. A COMMENSURABILITY RATIO

M. Belolipetsky and V. Emery [2] proved that for each odd dimension  $n \geq 5$  there is a unique orientable, noncompact, arithmetic, hyperbolic  $n$ -orbifold  $H^n/\Delta_n$  of smallest volume. If  $n \equiv 3 \pmod{4}$ , they proved that the volume of  $\Delta_n$  is given by

$$(32) \quad \text{vol}(H^n/\Delta_n) = \frac{3^{\frac{n}{2}}}{2^{\frac{n-1}{2}}} L\left(\frac{n+1}{2}, -3\right) \prod_{k=1}^{\frac{n-1}{2}} \frac{(2k-1)!}{(2\pi)^{2k}} \zeta(2k).$$

In terms of Bernoulli numbers, the above formula can be rewritten as

$$(33) \quad \text{vol}(H^n/\Delta_n) = \frac{3^{\frac{n}{2}} B}{2^{n-1}} L\left(\frac{n+1}{2}, -3\right).$$

In [6], V. Emery mentioned that  $\Delta_n$  is commensurable to  $\Gamma_3^n$  when  $n \equiv 3 \pmod{4}$ . We now compute the ratio  $\text{vol}(H^n/\Gamma_3^n)/\text{vol}(H^n/\Delta_n)$  for  $n \equiv 3 \pmod{4}$ . By Theorem 4, we have that

$$(34) \quad \text{vol}(H^n/\Gamma_3^n) = \frac{3^{\frac{n}{2}} B}{2^{n+1}} (2^{\frac{n-1}{2}} + (-1)^{\frac{n+1}{4}}) (2^{\frac{n+1}{2}} + 1) L\left(\frac{n+1}{2}, -3\right).$$

Therefore, we have

$$(35) \quad \frac{\text{vol}(H^n/\Gamma_3^n)}{\text{vol}(H^n/\Delta_n)} = \frac{1}{4} (2^{\frac{n-1}{2}} + (-1)^{\frac{n+1}{4}}) (2^{\frac{n+1}{2}} + 1).$$

Although  $\Delta_n$  is considered in [2] only for  $n \geq 5$ , we can define  $\Delta_3$  to be  $\text{PGL}(2, \mathcal{O}_3)$  where  $\mathcal{O}_3$  is the ring of integers of  $\mathbb{Q}(\sqrt{-3})$ . Then  $\Delta_3$  is arithmetic and Meyerhoff [12] proved that  $H^3/\Delta_3$  has minimum volume among all orientable, noncompact, hyperbolic 3-orbifolds. The group  $\Delta_3$  is the orientation preserving subgroup of a hyperbolic Coxeter group of type  $[3, 3, 6]$ . The above formula for  $\text{Vol}(H^3/\Delta_3)$  gives the correct volume for  $H^3/\Delta_3$ . Hence the above formula for  $\text{vol}(H^3/\Gamma_3^3)/\text{vol}(H^3/\Delta_3)$  gives the correct value  $5/4$ .

The groups  $\Gamma_3^3$  and  $\Delta_3$  are commensurable. The relationship between  $\Gamma_3^3$  and  $\Delta_3$  is explained by the commensurability diagram on page 130 of Johnson et al. [9]. The group  $\Gamma_3^3$  is a hyperbolic Coxeter group of type  $[4, 3, 6]$ . The Coxeter group  $[4, 3, 6]$  has a Coxeter subgroup of type  $[6, 3^{1,1}]$  of index 2. Hence the orientation preserving subgroup of  $[6, 3^{1,1}]$  has index 4 in  $[4, 3, 6]$ . Now the Coxeter tetrahedron of type  $[6, 3^{1,1}]$  can be subdivided into 5 copies of the Coxeter tetrahedron of type  $[3, 3, 6]$ . See Figure 1. Hence  $[6, 3^{1,1}]$  is conjugate to a subgroup of  $[3, 3, 6]$  of index 5. Therefore, the orientation preserving subgroup of  $[6, 3^{1,1}]$  is conjugate to a subgroup of  $\Delta_3$  of index 5. Thus, the ratio  $5/4$  faithfully represents the commensurability relationship between  $\Gamma_3^3$  and  $\Delta_3$ , that is,  $\Gamma_3^3$  has a subgroup of index 4 that is conjugate to a subgroup of  $\Delta_3$  of index 5.

We now turn our attention to dimension 7. Our volume ratio formula gives that

$$\text{vol}(H^7/\Gamma_3^7)/\text{vol}(H^7/\Delta_7) = 153/4.$$

By the main theorem of Hild [7], the group  $\Delta_7$  is a subgroup of index two of a discrete group  $\hat{\Delta}_7$  of isometries of  $H^7$  such that  $\Delta_7$  is the orientation preserving subgroup of  $\hat{\Delta}_7$ . The group  $\hat{\Delta}_7$  is generated by a Coxeter subgroup  $\bar{\Delta}_7$  of index 2 of type  $[3^{3,2,2}]$  and an involution  $\sigma$  that acts as the mirror symmetry of the Coxeter system defining  $\bar{\Delta}_7$ . The involution  $\sigma$  is orientation preserving, since  $\sigma$  transposes two pairs of Coxeter generators of  $\bar{\Delta}_7$ . Hence  $\Delta_7$  is generated by the orientation preserving subgroup of  $\bar{\Delta}_7$  and  $\sigma$ .




FIGURE 1. The subdivision of the Coxeter tetrahedron of type  $[6, 3^{1,1}]$  into 5 copies of the Coxeter tetrahedron of type  $[3, 3, 6]$  shown in the projective ball model of hyperbolic 3-space with the coordinate axes shown.

We explicitly defined the group  $\bar{\Delta}_7$  to be the group generated by the reflections in the sides of the hyperbolic Coxeter 7-simplex  $\Delta$  whose sides have Lorentz normal unit vectors listed in the columns of the matrix

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & -\frac{1}{2} & 0 & 1 \\ 0 & -\frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & 0 & -\frac{1}{2} & 0 & \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & 0 & 0 & \frac{1}{2} & 0 & 0 & -1 & \frac{1}{2} \\ 0 & 0 & 0 & \frac{1}{2} & -1 & \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & -1 & \frac{1}{2} & 0 & 0 & 0 & 0 \\ -1 & \frac{1}{2} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{\sqrt{3}}{2} \end{pmatrix}.$$

The involution  $\sigma$  acts as a symmetry of  $\Delta$ . The Lorentzian matrix of  $\sigma$  is

$$\begin{pmatrix} -2 & 0 & 0 & 0 & 0 & 0 & 0 & \sqrt{3} \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ -\sqrt{3} & 0 & 0 & 0 & 0 & 0 & 0 & 2 \end{pmatrix}.$$

In 1996, we discovered that  $\Gamma_3^7$  is a Coxeter reflection group defined by the Coxeter graph in Figure 1 of [10], for  $n = 7$ , using Vinberg's algorithm [17]. We explicitly defined the group  $\Gamma_3^7$  to be the group generated by the reflections in the sides of the hyperbolic Coxeter 7-dimensional polytope  $P^7$  whose sides have Lorentz

normal unit vectors listed in the columns of the matrix

$$\begin{pmatrix} -\frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 & 0 & \sqrt{\frac{3}{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} \\ 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} \\ 0 & 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} \\ 0 & 0 & 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 & 0 & 0 & \frac{1}{\sqrt{2}} \\ 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} & \sqrt{\frac{3}{2}} \end{pmatrix}.$$

In 1996, we discovered by a coset enumeration program that  $\bar{\Delta}_7 \cap \Gamma_3^7$  has index 2295 in  $\bar{\Delta}_7$  and index 60 in  $\Gamma_3^7$ . The involution  $\sigma$  is an element of  $\Gamma_3^7$  and so  $\Delta_7 \cap \Gamma_3^7$  has index 2295 in  $\Delta_7$  and index 60 in  $\Gamma_3^7$ . The commensurability ratio is  $2295/60 = 153/4$ . We were disappointed that we could not find representations of  $\Delta_7$  and  $\Gamma_3^7$  such that  $\Delta_7 \cap \Gamma_3^7$  has index 153 in  $\Delta_7$  and index 4 in  $\Gamma_3^7$ , but we suspect that the representations that we found give the smallest possible indices.

In 1996, we found a discrete group  $\Gamma$  of isometries of  $H^7$  that corresponds to the group of positive units of the quadratic form defined by the diagonal matrix  $\text{diag}(1, 1, 1, 1, 1, 1, 3, -1)$  such that  $\bar{\Delta}_7 \cap \Gamma$  has index 119 in  $\bar{\Delta}_7$  and index 4 in  $\Gamma$ . Moreover  $\text{vol}(H^7/\Gamma)/\text{vol}(H^7/\bar{\Delta}_7) = 119/4$ , and so the ratio  $119/4$  faithfully represents the commensurability relationship between  $\bar{\Delta}_7$  and  $\Gamma$ . This computation was reported on page 345 of [8]. The story of the computation of the volume of  $H^7/\Gamma$  will have to wait for another day.

#### REFERENCES

- [1] T. M. Apostol, *Introduction to Analytic Number Theory*, Undergraduate Texts in Math., Springer-Verlag, New York, 1976.
- [2] M. Belolipetsky and V. Emery, On volumes of arithmetic quotients of  $\text{PO}(n, 1)^\circ$ ,  $n$  odd, preprint, 2010.
- [3] I. M. Chiswell, The Euler characteristic of graph products and of Coxeter groups, In: *Discrete Groups and Geometry*, London Math. Soc. Lec. Notes Ser. 173 (1992), 36-46.
- [4] H. Cohen, *A Course in Computational Algebraic Number Theory*, Graduate Texts in Math., vol. 138, Springer-Verlag, Berlin, Heidelberg, and New York, 1993.
- [5] L. E. Dickson, *Linear Groups with an Exposition of the Galois Field Theory*, B. G. Teubner, Leipzig, 1901.
- [6] V. Emery, Even unimodular Lorentzian lattices and hyperbolic volume, preprint 2011.
- [7] T. Hild, The cusped hyperbolic orbifolds of minimal volume in dimensions less than ten, *J. Algebra* 313 (2007), 208-222.
- [8] N. W. Johnson, R. Kellerhals, J. G. Ratcliffe, and S. T. Tschantz, The size of a hyperbolic Coxeter simplex, *Transform. Groups* 4 (1999), 329-353.
- [9] N. W. Johnson, R. Kellerhals, J. G. Ratcliffe, and S. T. Tschantz, Commensurability classes of hyperbolic Coxeter groups, *Linear Algebra Appl.* 345 (2002), 119-147.
- [10] J. McLeod, Hyperbolic reflection groups associated to the quadratic forms  $-3x_0^2 + x_1^2 + \dots + x_n^2$ , *Geom. Dedicata* 152 (2011), 1-16.
- [11] J. Milnor, Hyperbolic geometry: the first 150 years, *Bull. Amer. Math. Soc.* 6 (1982), 9 - 24.
- [12] R. Meyerhoff, The cusped hyperbolic 3-orbifold of minimum volume, *Bull. Amer. Math. Soc.* 13 (1985), 154-156.
- [13] J.G. Ratcliffe, *Foundations of Hyperbolic Manifolds, Second Edition*, Graduate Texts in Math., vol. 149, Springer-Verlag, Berlin, Heidelberg, and New York, 2006.



- [14] J. G. Ratcliffe and S. T. Tschantz, Volumes of integral congruence hyperbolic manifolds *J. Reine Angew. Math.* 488 (1997), 55-78.
- [15] C. L. Siegel, Über die analytische Theorie der quadratischen Formen, *Ann. Math.* 36 (1935), 527-606.
- [16] C. L. Siegel, Über die analytische Theorie der quadratischen Formen II, *Ann. Math.* 37 (1936), 230-263.
- [17] È. B. Vinberg, On groups of unit elements of certain quadratic forms, *Math. Sbornik* 16, (1972), 17-35.

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TN 37240

*E-mail address:* `j.g.ratcliffe@vanderbilt.edu`