

BRAUER'S GENERALIZED DECOMPOSITION NUMBERS AND UNIVERSAL DEFORMATION RINGS

FRAUKE M. BLEHER

ABSTRACT. The versal deformation ring $R(G, V)$ of a mod p representation V of a profinite group G encodes all isomorphism classes of lifts of V to representations of G over complete local commutative Noetherian rings. We introduce a new technique for determining $R(G, V)$ when G is finite which involves Brauer's generalized decomposition numbers.

1. INTRODUCTION

Let k be an algebraically closed field of positive characteristic p , let $W = W(k)$ be the ring of infinite Witt vectors over k , and let G be a finite group. An important question in the representation theory of G is whether a finitely generated kG -module V can be lifted to W . For example, Green's lifting theorem shows that this is possible if there are no non-trivial 2-extensions of V by itself. A natural generalization of this question is to consider the functor which sends each complete local commutative Noetherian ring R with residue field k to the set of isomorphism classes of lifts of V over R . If this functor is represented by a ring $R(G, V)$, we say that $R(G, V)$ is the universal deformation ring of V . More generally, one can always associate to V a versal deformation ring $R(G, V)$, whose precise definition is recalled in Section 2. It was shown in [5, Prop. 2.1] that if the stable endomorphism ring $\underline{\text{End}}_{kG}(V)$ is isomorphic to k , then the versal deformation ring $R(G, V)$ is always universal.

Apart from the fact that universal deformation rings of representations of finite groups give more insight into the representation theory of these finite groups, there is another important motivation for studying these universal deformation rings. Namely, universal deformation rings for finite groups provide a good test case for various conjectures concerning the ring theoretic properties of universal deformation rings for profinite Galois groups. For example, Flach asked whether there could be universal deformation rings which are not complete intersections (see [21]). In [6, 7] it was shown that the universal deformation ring of the non-trivial irreducible mod 2 representation of the symmetric group S_4 is not a complete intersection; in fact it is not even Cohen-Macaulay. This led to infinitely many examples of real quadratic fields L such that the universal deformation ring of the inflation of this representation to the Galois group over L of the maximal totally unramified extension of L is not a complete intersection. In [8], examples of finite groups G and mod p representations of G for every odd prime number p were constructed such that the universal deformation rings of these representations are not complete intersections. The main advantage of computing universal deformation rings for representations of finite groups is that one can use deep results from modular representation theory due to Brauer, Erdmann [27], Linckelmann [33, 34], Carlson-Thévenaz [19, 20], and others.

In this paper, we propose a new method of determining universal deformation rings, using Brauer's generalized decomposition numbers. Brauer generalized the usual decomposition numbers in [11] to be able to express the values of ordinary irreducible characters not only on p -regular elements but also on p -singular elements in terms of Brauer characters. We will show how to use

2000 *Mathematics Subject Classification.* Primary 20C20; Secondary 20C15, 16G10.

Key words and phrases. Universal deformation rings; Brauer's generalized decomposition numbers; tame blocks; dihedral defect groups; semidihedral defect groups; generalized quaternion defect groups.

The author was supported in part by NSA Grant H98230-11-1-0131.

these generalized decomposition numbers to determine the universal deformation rings $R(G, V)$ for certain mod p representations V of finite groups G whose stable endomorphisms are all given by scalars. The V we consider are those for which Brauer's generalized decomposition numbers carry the most information; these V are called maximally ordinary below. For maximally ordinary V , the generalized decomposition numbers enable us to find a family of Galois orbits of ordinary irreducible characters of G which can be used to construct lifts of V to large local rings in characteristic 0. This provides a powerful tool for computing universal deformation rings. The need for the use of Brauer's generalized decomposition numbers is related to how much fusion of conjugacy classes of p -power order elements occurs in G .

Suppose V is a kG -module whose stable endomorphism ring is isomorphic to k . Then V has a unique non-projective indecomposable summand (up to isomorphism). Since the universal deformation ring $R(G, V)$ only depends on this indecomposable direct summand by [5, Cor. 2.7], we may assume that V is indecomposable. Hence there exists a unique p -block B to which V belongs. The case when B has finite representation type has been fully studied in [5]. Therefore, we concentrate in this paper on the case when B has infinite tame representation type. Because the study of local blocks requires different arguments, as was demonstrated for example in [2], we further assume that B is not local. Let D be a defect group of B . We say V is maximally ordinary if the Brauer character of V is the restriction of an ordinary irreducible character χ such that for every $\sigma \in D$ of maximal p -power order, Brauer's generalized decomposition numbers corresponding to σ and χ do not all lie in $\{0, \pm 1\}$ (see Definition 3.2). This condition ensures that Brauer's generalized decomposition numbers carry enough information for our method to be applied to V .

The following theorem summarizes our main results; more precise statements can be found in Sect. 6, and in particular in Corollary 6.2 and Theorem 6.6.

Theorem 1.1. *Suppose G is a finite group, B is a non-local block of kG of infinite tame representation type, and D is a defect group of B of order p^n . There exists an indecomposable kG -module V belonging to B whose stable endomorphism ring is isomorphic to k and which is maximally ordinary if and only if $n \geq 4$. Moreover, the isomorphism class of every such V can be described explicitly. Suppose V is such a maximally ordinary module. There exists a monic polynomial $q_n(t) \in W[t]$ of degree $p^{n-2} - 1$ which depends on D but not on V and which can be given explicitly such that either*

- (i) $R(G, V)/pR(G, V) \cong k[[t]]/(t^{p^{n-2}-1})$, in which case $R(G, V) \cong W[[t]]/(q_n(t))$, or
- (ii) $R(G, V)/pR(G, V) \cong k[[t]]/(t^{p^{n-2}})$, in which case $R(G, V) \cong W[[t]]/(t q_n(t), p q_n(t))$.

In all cases, the ring $R(G, V)$ is isomorphic to a subquotient ring of WD , and it is a complete intersection if and only if we are in case (i).

In particular, Theorem 1.1 gives a positive answer to [5, Question 1.1] for all B and V considered in the theorem. A precise description of the maximally ordinary modules V belonging to B is given in Corollary 6.2. A formula for the polynomials $q_n(t)$ can be found in Definition 5.3 and Remark 5.4.

The use of Brauer's generalized decomposition numbers provides a correction to some arguments in [3, Sect. 3.4], [4, Sect. 5], and [9, Sect. 3.2]. Namely, in these papers a formula for the values of the ordinary irreducible characters on elements in D of maximal p -power order was assumed, which is true for principal infinite tame blocks but cannot be verified for arbitrary infinite tame blocks because there may be more fusion of D -conjugacy classes in G when the blocks are not principal (see Remark 5.2).

The paper is organized as follows. In Section 2, we recall the definitions of deformations and deformation rings. In Section 3, we give a brief introduction to Brauer's generalized decomposition numbers, as introduced in [11, 12]. Using [27], we describe in Section 4 the quivers and relations of the basic algebras of all non-local blocks B of infinite tame representation type, and provide their decomposition matrices. In Section 5, we describe results of [16, 37] about the ordinary irreducible characters of G belonging to B . In Section 6, we prove Theorem 1.1.

2. PRELIMINARIES

In this section, we give a brief introduction to versal and universal deformation rings and deformations. For more background material, we refer the reader to [36] and [23].

Let k be an algebraically closed field of characteristic $p > 0$, and let W be the ring of infinite Witt vectors over k . Let $\hat{\mathcal{C}}$ be the category of all complete local commutative Noetherian rings with residue field k . Note that all rings in $\hat{\mathcal{C}}$ have a natural W -algebra structure. The morphisms in $\hat{\mathcal{C}}$ are continuous W -algebra homomorphisms which induce the identity map on k .

Suppose G is a finite group and V is a finitely generated kG -module. A *lift* of V over an object R in $\hat{\mathcal{C}}$ is a pair (M, ϕ) where M is a finitely generated RG -module which is free over R , and $\phi : k \otimes_R M \rightarrow V$ is an isomorphism of kG -modules. Two lifts (M, ϕ) and (M', ϕ') of V over R are isomorphic if there is an isomorphism $f : M \rightarrow M'$ with $\phi = \phi' \circ (k \otimes f)$. The isomorphism class $[M, \phi]$ of a lift (M, ϕ) of V over R is called a *deformation* of V over R , and the set of all such deformations is denoted by $\text{Def}_G(V, R)$. The deformation functor

$$\hat{F}_V : \hat{\mathcal{C}} \rightarrow \text{Sets}$$

is a covariant functor which sends an object R in $\hat{\mathcal{C}}$ to $\text{Def}_G(V, R)$ and a morphism $\alpha : R \rightarrow R'$ in $\hat{\mathcal{C}}$ to the map $\text{Def}_G(V, R) \rightarrow \text{Def}_G(V, R')$ defined by $[M, \phi] \mapsto [R' \otimes_{R, \alpha} M, \phi_\alpha]$, where $\phi_\alpha = \phi$ after identifying $k \otimes_{R'} (R' \otimes_{R, \alpha} M)$ with $k \otimes_R M$.

Suppose there exists an object $R(G, V)$ in $\hat{\mathcal{C}}$ and a deformation $[U(G, V), \phi_U]$ of V over $R(G, V)$ with the following property: For each R in $\hat{\mathcal{C}}$ and for each lift (M, ϕ) of V over R there exists a morphism $\alpha : R(G, V) \rightarrow R$ in $\hat{\mathcal{C}}$ such that $\hat{F}_V(\alpha)([U(G, V), \phi_U]) = [M, \phi]$, and moreover α is unique if R is the ring of dual numbers $k[\epsilon]/(\epsilon^2)$. Then $R(G, V)$ is called the *versal deformation ring* of V and $[U(G, V), \phi_U]$ is called the *versal deformation* of V . If the morphism α is unique for all R and all lifts (M, ϕ) of V over R , then $R(G, V)$ is called the *universal deformation ring* of V and $[U(G, V), \phi_U]$ is called the *universal deformation* of V . In other words, $R(G, V)$ is universal if and only if $R(G, V)$ represents the functor \hat{F}_V in the sense that \hat{F}_V is naturally isomorphic to the Hom functor $\text{Hom}_{\hat{\mathcal{C}}}(R(G, V), -)$.

Note that the above definition of deformations can be weakened as follows. Given a lift (M, ϕ) of V over a ring R in $\hat{\mathcal{C}}$, define the corresponding weak deformation to be the isomorphism class of M as an RG -module, without taking into account the specific isomorphism $\phi : k \otimes_R M \rightarrow V$. In general, a weak deformation of V over R identifies more lifts than a deformation of V over R that respects the isomorphism ϕ of a representative (M, ϕ) . However, if the stable endomorphism ring $\underline{\text{End}}_{kG}(V)$ is isomorphic to k , these two definitions of deformations coincide (see [4, Remark 2.1]).

By [36], every finitely generated kG -module V has a versal deformation ring. Since G is a finite group, we have the following sufficient criterion for the universality of $R(G, V)$:

Proposition 2.1. ([5, Prop. 2.1]) *Suppose V is a finitely generated kG -module whose stable endomorphism ring $\underline{\text{End}}_{kG}(V)$ is isomorphic to k . Then V has a universal deformation ring $R(G, V)$.*

3. BRAUER'S GENERALIZED DECOMPOSITION NUMBERS

In this section, we give a brief introduction to Brauer's generalized decomposition numbers, emphasizing the results needed in this paper. Throughout this section, let p be a fixed prime number and let G be a finite group such that p divides $\#G$. Let P be a fixed Sylow p -subgroup of G .

Brauer introduced generalized decomposition numbers in [11] to be able to express the values of the ordinary irreducible characters of G on all conjugacy classes by means of the irreducible p -modular characters of certain subgroups of G . More precisely, let $g \in G$ and write g (uniquely) as $g = uv$ where u is a p -element of order p^α and v is a p -regular element in the centralizer $C_G(u)$ (where we allow u or v to be identity elements). Let $\text{IBr}(C_G(u))$ denote the set of distinct irreducible p -modular characters of $C_G(u)$. By [11, Sect. 1], if χ is an ordinary irreducible character of G then

we have a formula

$$(3.1) \quad \chi(uv) = \sum_{\varphi \in \text{IBr}(C_G(u))} d_{\chi, \varphi}^{(u)} \varphi(v)$$

where the numbers $d_{\chi, \varphi}^{(u)}$, $\varphi \in \text{IBr}(C_G(u))$, are algebraic integers in the field of p^α -th roots of unity which do not depend on v . These are called the *generalized decomposition numbers* of G corresponding to u and χ .

If $u = 1_G$, then the generalized decomposition numbers corresponding to u are the usual decomposition numbers, which are non-negative integers, and (3.1) is the usual formula for the restriction to p -regular elements of an ordinary irreducible character χ in terms of an integral combination of the irreducible p -modular characters of $G = C_G(1_G)$.

We can use these generalized decomposition numbers to express the ordinary character table of G as the product of two square matrices as follows. Recall that we fixed a Sylow p -subgroup P of G . Let u_0, u_1, \dots, u_h be a complete system of representatives of G -conjugacy classes of p -elements in G with $u_0 = 1_G$ and $u_i \in P$ for all $1 \leq i \leq h$. For each $0 \leq i \leq h$, let $v_{i,1}, \dots, v_{i,\ell_i}$ be a complete system of representatives of $C_G(u_i)$ -conjugacy classes of p -regular elements in $C_G(u_i)$ with $v_{i,1} = 1_G$. Then $\{u_i v_{i,j} \mid 0 \leq i \leq h, 1 \leq j \leq \ell_i\}$ is a complete set of representatives of the conjugacy classes of G . Moreover, for each $0 \leq i \leq h$, there are precisely ℓ_i irreducible p -modular characters of $C_G(u_i)$, which we denote by $\varphi_{i,1}, \dots, \varphi_{i,\ell_i}$. Let $\{\chi_1, \dots, \chi_c\}$ be a complete set of representatives of the ordinary irreducible characters of G . Then Brauer's above formula (3.1) can be written as

$$(3.2) \quad \chi_s(u_i v_{i,j}) = \sum_{t=1}^{\ell_i} d_{\chi_s, \varphi_{i,t}}^{(u_i)} \varphi_{i,t}(v_{i,j})$$

for all $1 \leq s \leq c$, $0 \leq i \leq h$, $1 \leq j \leq \ell_i$, where $d_{\chi_s, \varphi_{i,t}}^{(u_i)} = d_{\chi_s, \varphi_{i,t}}^{(u_i)}$ for all $1 \leq t \leq \ell_i$. Write the conjugacy class representatives in the order

$$u_0 v_{0,1}, \dots, u_0 v_{0,\ell_0}, u_1 v_{1,1}, \dots, u_1 v_{1,\ell_1}, \dots, u_h v_{h,1}, \dots, u_h v_{h,\ell_h}.$$

Using (3.2), we obtain that the ordinary character table \mathcal{X} can be written as a product $\mathcal{X} = \Delta \cdot \Phi$, where Δ contains the generalized decomposition numbers and Φ is a block diagonal matrix

$$(3.3) \quad \Phi = \begin{pmatrix} \Phi_0 & & 0 \\ & \ddots & \\ 0 & & \Phi_h \end{pmatrix}$$

with

$$\Phi_i = \begin{pmatrix} \varphi_{i,1}(v_{i,1}) & \cdots & \varphi_{i,1}(v_{i,\ell_i}) \\ \vdots & & \vdots \\ \varphi_{i,\ell_i}(v_{i,1}) & \cdots & \varphi_{i,\ell_i}(v_{i,\ell_i}) \end{pmatrix}$$

for all $0 \leq i \leq h$. By [12, Sect. 7, p. 45], the square of the determinant of Δ is $\pm p^a$ for some $a \in \mathbb{Z}^+$, and the square of the determinant of Φ is an integer which is relatively prime to p .

Fix now a p -modular block B of G and suppose that there are $k(B)$ ordinary irreducible characters belonging to B . Reorder χ_1, \dots, χ_c such that $\chi_1, \dots, \chi_{k(B)}$ belong to B . For each $0 \leq i \leq h$, reorder $\varphi_{i,1}, \dots, \varphi_{i,\ell_i}$ such that the first m_i characters, $\varphi_{i,1}, \dots, \varphi_{i,m_i}$, are precisely the irreducible p -modular characters belonging to blocks of $C_G(u_i)$ whose Brauer correspondents in G are equal to B . (Note that for each block b of $C_G(u_i)$, its Brauer correspondent b^G in G is well-defined since the centralizer in G of a defect group of b is contained in $C_G(u_i)$, see [1, Sect. 14].) It follows from [12, Sect. 6] that for $1 \leq s \leq k(B)$, (3.2) can be rewritten as

$$(3.4) \quad \chi_s(u_i v_{i,j}) = \sum_{t=1}^{m_i} d_{\chi_s, \varphi_{i,t}}^{(u_i)} \varphi_{i,t}(v_{i,j})$$

for all $0 \leq i \leq h$, $1 \leq j \leq \ell_i$. In particular, we obtain for all $0 \leq i \leq h$ that

$$(3.5) \quad \mathcal{X}_{B,i} = \begin{pmatrix} \chi_1(u_i v_{i,1}) & \cdots & \chi_1(u_i v_{i,\ell_i}) \\ \vdots & & \vdots \\ \chi_{k(B)}(u_i v_{i,1}) & \cdots & \chi_{k(B)}(u_i v_{i,\ell_i}) \end{pmatrix} = \Delta_{B,i} \cdot \Phi_i$$

where

$$\Delta_{B,i} = \begin{pmatrix} d_{1,1}^i & \cdots & d_{1,m_i}^i & 0 & \cdots & 0 \\ \vdots & & \vdots & \vdots & & \vdots \\ d_{k(B),1}^i & \cdots & d_{k(B),m_i}^i & 0 & \cdots & 0 \end{pmatrix}$$

and we assume, as above, that $\varphi_{i,1}, \dots, \varphi_{i,m_i}$ are precisely the irreducible p -modular characters belonging to blocks of $C_G(u_i)$ whose Brauer correspondents in G are equal to B .

Let now k be an algebraically closed field of characteristic p and view B as a block of kG . Let $W = W(k)$ be the ring of infinite Witt vectors over k and let F be the fraction field of W . In particular, W contains all roots of unity of order not divisible by p . Let \overline{F} be a fixed algebraic closure of F , and let ξ be a root of unity in \overline{F} whose order is a power of p such that $F(\xi)$ is a splitting field for G . Then $W[\xi]$ is the ring of integers of $F(\xi)$, and we can view the ordinary character table \mathcal{X} of G and the matrix Δ of generalized decomposition numbers as taking values in $W[\xi]$, and the matrix Φ in (3.3) as taking values in W . Since the square of the determinant of Φ is an integer that is relatively prime to p , it follows that the determinant of Φ , and hence the determinant of Φ_i for all $0 \leq i \leq h$, is a unit in W . Hence we can solve (3.5) for $\Delta_{B,i}$ to obtain

$$(3.6) \quad \Delta_{B,i} = \mathcal{X}_{B,i} \cdot \Psi_i$$

for all $0 \leq i \leq h$, where $\Psi_i = \Phi_i^{-1}$ is a matrix with values in W . In particular, (3.6) shows that if we reduce our discussion to a single block B then we can replace ξ by a p -power order root of unity ζ in \overline{F} such that all ordinary irreducible characters of G belonging to B are realizable over $F(\zeta)$, i.e. they correspond to absolutely irreducible $F(\zeta)G$ -modules.

Remark 3.1. Equations (3.1) and (3.4) can be rewritten to reflect the influence of fusion of P -conjugacy classes in G (see [14, Sect. 1]). As before, let u be a p -element of G and let v be a p -regular element in $C_G(u)$. Assuming the notation of the previous paragraph, let χ be an irreducible $F(\xi)$ -character which belongs to a block B of kG . Recall that a subsection (y, b_y) for B is a pair consisting of a p -element y of G and a block b_y of $C_G(y)$ with $b_y^G = B$. We obtain

$$(3.7) \quad \chi(uv) = \sum_{(y, b_y)} \sum_{\varphi \in \text{IBr}(b_y)} d_{\chi, \varphi}^{(y)} \varphi(z_y v z_y^{-1})$$

where (y, b_y) ranges over a system of representatives for the conjugacy classes of subsections for B such that y is conjugate to u in G , say $u = z_y^{-1} y z_y$. For each (y, b_y) , φ ranges over the irreducible p -modular characters associated with b_y .

Definition 3.2. Let k be an algebraically closed field of characteristic p , let W be the ring of infinite Witt vectors over k , and let \overline{F} be a fixed algebraic closure of the fraction field F of W . Suppose B is a block of kG and that D is a defect group of B . Let $\zeta \in \overline{F}$ be a root of unity of p -power order such that all ordinary irreducible characters of G belonging to B are realizable over $F(\zeta)$. In particular, the ordinary (resp. p -modular) characters belonging to B can be viewed as taking values in $W[\zeta]$ (resp. W). Suppose V is an indecomposable kG -module whose stable endomorphism ring is isomorphic to k and which belongs to B . We say V is *maximally ordinary* if the p -modular character of V is the restriction to the p -regular elements of an ordinary irreducible character χ such that for every $\sigma \in D$ of maximal p -power order there exists an irreducible p -modular character φ of $C_G(\sigma)$ such that $d_{\chi, \varphi}^{(\sigma)} \notin \{0, \pm 1\}$.

4. TAME BLOCKS

For the remainder of this paper, we make the following assumptions:

Hypothesis 4.1. Let k be an algebraically closed field of positive characteristic p , let $W = W(k)$ be the ring of infinite Witt vectors over k , let F be the fraction field of W , and let \overline{F} be a fixed algebraic closure of F . Suppose G is a finite group, B is a non-local block of kG of infinite tame representation type, and D is a defect group of B of order p^n . Let ζ be a primitive p^{n-1} -th root of unity in \overline{F} .

It follows from [10, 17, 29] that $p = 2$, $n \geq 2$, and D is dihedral, semidihedral or generalized quaternion. In particular, $n \geq 2$ if D is dihedral, $n \geq 3$ if D is generalized quaternion, and $n \geq 4$ if D is semidihedral. Moreover, Brauer and Olsson proved that all ordinary irreducible characters of G belonging to B take values in $F(\zeta)$ (see [15, Prop. (7D)], [16, Prop. (5A)] and [37, Prop. 4.1]). In fact, all ordinary irreducible characters of G belonging to B can be realized by simple $F(\zeta)G$ -modules (see Section 5).

Assume Hypothesis 4.1. By [16, 37], it follows that there are at most three isomorphism classes of simple B -modules. From Erdmann's classification of all blocks of tame representation type in [27], it follows that the quiver and relations of the basic algebra of B can be given explicitly and that, up to Morita equivalence, there are 24 families of blocks B .

Using [24, 26, 27, 30], we now give a description of these families as follows. By [27, pp. 294–306], there are 12 possible quivers Q which can occur for basic algebras of dihedral, semidihedral or quaternion type: $2\mathcal{A}$, $2\mathcal{B}$, $3\mathcal{A}$, $3\mathcal{B}$, $3\mathcal{C}$, $3\mathcal{D}$, $3\mathcal{F}$, $3\mathcal{H}$, $3\mathcal{K}$, $3\mathcal{L}$, $3\mathcal{Q}$, $3\mathcal{R}$. For each such quiver Q , we combine Erdmann's results in [26, 27] and [30, Prop. 4.2] with Eisele's results in [24] to provide the most accurate description of a full set of representatives of basic algebras $\Lambda = kQ/I$ for the Morita equivalence classes of blocks B as in Hypothesis 4.1 with Ext quiver Q .

We also provide the decomposition matrix for each block B . For better readability, all decomposition matrices appear at the end of the paper. As will be discussed in Section 5, B always contains exactly 4 ordinary irreducible characters of height 0 and, unless D is quaternion of order 8, exactly $2^{n-2} - 1$ ordinary irreducible characters of height 1. If D is quaternion of order 8, B contains exactly 3 ordinary irreducible characters of height 1. If B is generalized quaternion or semidihedral, there may be additional ordinary irreducible characters of height $n - 2$. In the decomposition matrices, we list first the 4 ordinary irreducible characters of height 0, then the family of $2^{n-2} - 1$ ordinary irreducible characters of height 1, and then the ordinary irreducible characters of height $n - 2$ if they exist. Note that the family of $2^{n-2} - 1$ characters of height 1 all define the same Brauer character on restricting to the 2-regular conjugacy classes of G . If B has three isomorphism classes of simple modules, we moreover order the ordinary irreducible characters according to the sign conventions described in [16, Thm. 5] and [37, Thms. 4.10, 4.11 and 4.15].

To distinguish between different defect groups, we use the notation $D(Q)$ (resp. $SD(Q)$, resp. $Q(Q)$) to mean that $\Lambda = kQ/I$ is Morita equivalent to a block B with dihedral (resp. semidihedral, resp. generalized quaternion) defect groups.

4.1. Blocks with quiver $2\mathcal{A}$.

$$2\mathcal{A} = \alpha \begin{array}{c} \circlearrowleft \\ \bullet \xrightarrow{\beta} \bullet \\ \bullet \xleftarrow{\gamma} \bullet \end{array}$$

By [24] and [27, p. 294], if B has dihedral defect groups and Ext quiver $2\mathcal{A}$, then B is Morita equivalent to $D(2\mathcal{A}) = k[2\mathcal{A}]/I_{D(2\mathcal{A})}$ where

$$I_{D(2\mathcal{A})} = \langle \beta\gamma, \alpha^2, (\gamma\beta\alpha)^{2^{n-2}} - (\alpha\gamma\beta)^{2^{n-2}} \rangle.$$

The corresponding decomposition matrix is given in Figure 1.

By [27, p. 298], if B has semidihedral defect groups and Ext quiver $2\mathcal{A}$, then there exists $c \in k$ such that B is Morita equivalent to either $SD(2\mathcal{A})_1(c) = k[2\mathcal{A}]/I_{SD(2\mathcal{A})_1,c}$ or $SD(2\mathcal{A})_2(c) =$

$k[2\mathcal{A}]/I_{\text{SD}(2\mathcal{A})_2,c}$ where

$$\begin{aligned} I_{\text{SD}(2\mathcal{A})_1,c} &= \langle \alpha^2 - c(\gamma\beta\alpha)^{2^{n-2}}, \beta\gamma\beta - \beta\alpha(\gamma\beta\alpha)^{2^{n-2}-1}, \gamma\beta\gamma - \alpha\gamma(\beta\alpha\gamma)^{2^{n-2}-1}, \\ &\quad \alpha(\gamma\beta\alpha)^{2^{n-2}} \rangle, \\ I_{\text{SD}(2\mathcal{A})_2,c} &= \langle \beta\gamma, \alpha^2 - \gamma\beta(\alpha\gamma\beta)^{2^{n-2}-1} - c(\gamma\beta\alpha)^{2^{n-2}}, (\gamma\beta\alpha)^{2^{n-2}} - (\alpha\gamma\beta)^{2^{n-2}} \rangle. \end{aligned}$$

The decomposition matrix for $\text{SD}(2\mathcal{A})_1(c)$ is given in Figure 2, and the decomposition matrix for $\text{SD}(2\mathcal{A})_2(c)$ is given in Figure 1.

By [27, p. 303], if B has generalized quaternion defect groups and Ext quiver $2\mathcal{A}$, then there exists $c \in k$ such that B is Morita equivalent to $\text{Q}(2\mathcal{A})(c) = k[2\mathcal{A}]/I_{\text{Q}(2\mathcal{A}),c}$ where

$$\begin{aligned} I_{\text{Q}(2\mathcal{A}),c} &= \langle \alpha^2 - \gamma\beta(\alpha\gamma\beta)^{2^{n-2}-1} - c(\alpha\gamma\beta)^{2^{n-2}}, \beta\gamma\beta - \beta\alpha(\gamma\beta\alpha)^{2^{n-2}-1}, \\ &\quad \gamma\beta\gamma - \alpha\gamma(\beta\alpha\gamma)^{2^{n-2}-1}, \beta\alpha^2 \rangle. \end{aligned}$$

The decomposition matrix for $\text{Q}(2\mathcal{A})(c)$ is given in Figure 2.

4.2. Blocks with quiver $2\mathcal{B}$.

$$2\mathcal{B} = \alpha \begin{array}{c} \circlearrowleft \\ \bullet \\ \circlearrowright \end{array} \begin{array}{c} 0 \\ \xrightarrow{\beta} \\ 1 \\ \xleftarrow{\gamma} \end{array} \begin{array}{c} \bullet \\ \circlearrowright \\ \bullet \end{array} \eta$$

By [24] and [27, p. 295], if B has dihedral defect groups and Ext quiver $2\mathcal{B}$, then B is Morita equivalent to $\text{D}(2\mathcal{B}) = k[2\mathcal{B}]/I_{\text{D}(2\mathcal{B})}$ where

$$I_{\text{D}(2\mathcal{B})} = \langle \eta\beta, \gamma\eta, \beta\gamma, \alpha^2, \gamma\beta\alpha - \alpha\gamma\beta, \eta^{2^{n-2}} - \beta\alpha\gamma \rangle.$$

The corresponding decomposition matrix is given in Figure 3.

By [26, Lemmas (8.11) and (8.15)], [27, p. 299] and [30, Prop. 4.2], if B has semidihedral defect groups and Ext quiver $2\mathcal{B}$, then there exists $c \in k$ such that B is Morita equivalent to either $\text{SD}(2\mathcal{B})_1(c) = k[2\mathcal{B}]/I_{\text{SD}(2\mathcal{B})_1,c}$ or $\text{SD}(2\mathcal{B})_2(c) = k[2\mathcal{B}]/I_{\text{SD}(2\mathcal{B})_2,c}$ or $\text{SD}(2\mathcal{B})_4(c) = k[2\mathcal{B}]/I_{\text{SD}(2\mathcal{B})_4,c}$ where

$$\begin{aligned} I_{\text{SD}(2\mathcal{B})_1,c} &= \langle \eta\beta, \gamma\eta, \beta\gamma, \alpha^2 - \gamma\beta - c\alpha\gamma\beta, \gamma\beta\alpha - \alpha\gamma\beta, \eta^{2^{n-2}} - \beta\alpha\gamma \rangle, \\ I_{\text{SD}(2\mathcal{B})_2,c} &= \langle \eta\beta - \beta\alpha(\gamma\beta\alpha), \gamma\eta - \alpha\gamma(\beta\alpha\gamma), \alpha^2 - c(\gamma\beta\alpha)^2, \beta\gamma - \eta^{2^{n-2}-1}, \\ &\quad \eta^2\beta, \gamma\eta^2 \rangle, \\ I_{\text{SD}(2\mathcal{B})_4,c} &= \langle \gamma\eta - \alpha\gamma, \beta\alpha - \eta\beta, \alpha^{2^{n-2}+1}, \eta^{2^{n-2}+1}, \beta\alpha^{2^{n-2}-1}, \alpha^{2^{n-2}-1}\gamma, \\ &\quad \gamma\eta^{2^{n-2}-1}, \eta^{2^{n-2}-1}\beta, \gamma\beta - \alpha^2, \beta\gamma - \eta^2(1 + c\eta^{2^{n-2}-2}) \rangle. \end{aligned}$$

The decomposition matrix for $\text{SD}(2\mathcal{B})_1(c)$ is given in Figure 3, the decomposition matrix for $\text{SD}(2\mathcal{B})_2(c)$ is given in Figure 4, and the decomposition matrix for $\text{SD}(2\mathcal{B})_4(c)$ is given in Figure 5.

By [27, IX.4.1 and pp. 303–304], if B has generalized quaternion defect groups and Ext quiver $2\mathcal{B}$, then there exists $c \in k$, respectively $p(t) \in k[t]$ with $p(0) = 1$ and $a, c \in k$ with $a \neq 0$, such that B is Morita equivalent to either $\text{Q}(2\mathcal{B})_1(c) = k[2\mathcal{B}]/I_{\text{Q}(2\mathcal{B})_1,c}$ or $\text{Q}(2\mathcal{B})_2(p, a, c) = k[2\mathcal{B}]/I_{\text{Q}(2\mathcal{B})_2,p,a,c}$ where

$$\begin{aligned} I_{\text{Q}(2\mathcal{B})_1,c} &= \langle \eta\beta - \beta\alpha(\gamma\beta\alpha), \gamma\eta - \alpha\gamma(\beta\alpha\gamma), \alpha^2 - \gamma\beta(\alpha\gamma\beta) - c(\alpha\gamma\beta)^2, \\ &\quad \beta\gamma - \eta^{2^{n-2}-1}, \beta\alpha^2 \rangle, \\ I_{\text{Q}(2\mathcal{B})_2,p,a,c} &= \langle \gamma\eta - \alpha\gamma, \beta\alpha - \eta\beta, \alpha^{2^{n-2}+1}, \eta^{2^{n-2}+1}, \beta\alpha^{2^{n-2}-1}, \alpha^{2^{n-2}-1}\gamma, \\ &\quad \gamma\beta - p(\alpha)\alpha^2, \beta\gamma - p(\eta)\eta^2 - a\eta^{2^{n-2}-1} - c\eta^{2^{n-2}} \rangle. \end{aligned}$$

The decomposition matrix for $\text{Q}(2\mathcal{B})_1(c)$ is given in Figure 4, and the decomposition matrix for $\text{Q}(2\mathcal{B})_2(p, a, c)$ is given in Figure 5. Note that by [37, Lemma 3.3], $\text{Q}(2\mathcal{B})_2(p, a, c)$ can actually not occur as a block.

4.3. Blocks with quiver $3\mathcal{A}$.

$$3\mathcal{A} = 1 \bullet \begin{array}{c} \xrightarrow{\beta} 0 \\ \xleftarrow{\gamma} \end{array} \bullet \begin{array}{c} \xrightarrow{\delta} \\ \xleftarrow{\eta} \end{array} \bullet 2$$

By [27, IX.5.4 and p. 295], if B has dihedral defect groups and Ext quiver $3\mathcal{A}$, then B is Morita equivalent to $D(3\mathcal{A})_1 = k[3\mathcal{A}]/I_{D(3\mathcal{A})_1}$ where

$$I_{D(3\mathcal{A})_1} = \langle \gamma\beta, \delta\eta, (\eta\delta\beta\gamma)^{2^{n-2}} - (\beta\gamma\eta\delta)^{2^{n-2}} \rangle.$$

The corresponding decomposition matrix is given in Figure 6.

By [27, IX.5.3 and pp. 299–300], if B has semidihedral defect groups and Ext quiver $3\mathcal{A}$, then B is Morita equivalent to $SD(3\mathcal{A})_1 = k[3\mathcal{A}]/I_{SD(3\mathcal{A})_1}$ where

$$I_{SD(3\mathcal{A})_1} = \langle \gamma\beta, \delta\eta\delta - \delta\beta\gamma(\eta\delta\beta\gamma)^{2^{n-2}-1}, \eta\delta\eta - \beta\gamma\eta(\delta\beta\gamma\eta)^{2^{n-2}-1} \rangle.$$

The decomposition matrix for $SD(3\mathcal{A})_1$ is given in Figure 7.

By [27, IX.5.2 and pp. 304–305], if B has generalized quaternion defect groups and Ext quiver $3\mathcal{A}$, then B is Morita equivalent to $Q(3\mathcal{A})_2 = k[3\mathcal{A}]/I_{Q(3\mathcal{A})_2}$ where

$$I_{Q(3\mathcal{A})_2} = \langle \beta\gamma\beta - \eta\delta\beta(\gamma\eta\delta\beta)^{2^{n-2}-1}, \gamma\beta\gamma - \gamma\eta\delta(\beta\gamma\eta\delta)^{2^{n-2}-1}, \delta\beta\gamma\beta, \\ \delta\eta\delta - \delta\beta\gamma(\eta\delta\beta\gamma)^{2^{n-2}-1}, \eta\delta\eta - \beta\gamma\eta(\delta\beta\gamma\eta)^{2^{n-2}-1}, \gamma\eta\delta\eta \rangle.$$

The decomposition matrix for $Q(3\mathcal{A})_2$ is given in Figure 8.

4.4. Blocks with quiver $3\mathcal{B}$.

$$3\mathcal{B} = \alpha \circlearrowleft \bullet \begin{array}{c} \xrightarrow{\beta} 0 \\ \xleftarrow{\gamma} \end{array} \bullet \begin{array}{c} \xrightarrow{\delta} \\ \xleftarrow{\eta} \end{array} \bullet 2$$

By [27, IX.5.4 and pp. 295–296], if B has dihedral defect groups and Ext quiver $3\mathcal{B}$, then B is Morita equivalent to $D(3\mathcal{B})_1 = k[3\mathcal{B}]/I_{D(3\mathcal{B})_1}$ where

$$I_{D(3\mathcal{B})_1} = \langle \beta\alpha, \alpha\gamma, \gamma\beta, \delta\eta, \eta\delta\beta\gamma - \beta\gamma\eta\delta, \alpha^{2^{n-2}} - \gamma\eta\delta\beta \rangle.$$

The corresponding decomposition matrix is given in Figure 9.

By [27, p. 300], if B has semidihedral defect groups and Ext quiver $3\mathcal{B}$, then B is Morita equivalent to either $SD(3\mathcal{B})_1 = k[3\mathcal{B}]/I_{SD(3\mathcal{B})_1}$ or $SD(3\mathcal{B})_2 = k[3\mathcal{B}]/I_{SD(3\mathcal{B})_2}$ where

$$I_{SD(3\mathcal{B})_1} = \langle \beta\alpha, \alpha\gamma, \gamma\beta, \delta\eta\delta - \delta\beta\gamma, \eta\delta\eta - \beta\gamma\eta, \alpha^{2^{n-2}} - \gamma\eta\delta\beta \rangle, \\ I_{SD(3\mathcal{B})_2} = \langle \delta\eta, \gamma\beta - \alpha^{2^{n-2}-1}, \alpha\gamma - \gamma\eta\delta(\beta\gamma\eta\delta), \beta\alpha - \eta\delta\beta(\gamma\eta\delta\beta) \rangle.$$

The decomposition matrix for $SD(3\mathcal{B})_1$ is given in Figure 10, and the decomposition matrix for $SD(3\mathcal{B})_2$ is given in Figure 11.

By [27, p. 305], if B has generalized quaternion defect groups and Ext quiver $3\mathcal{B}$, then B is Morita equivalent to $Q(3\mathcal{B}) = k[3\mathcal{B}]/I_{Q(3\mathcal{B})}$ where

$$I_{Q(3\mathcal{B})} = \langle \gamma\beta - \alpha^{2^{n-2}-1}, \alpha\gamma - \gamma\eta\delta(\beta\gamma\eta\delta), \beta\alpha - \eta\delta\beta(\gamma\eta\delta\beta), \delta\eta\delta - \delta\beta\gamma(\eta\delta\beta\gamma), \\ \eta\delta\eta - \beta\gamma\eta(\delta\beta\gamma\eta), \beta\alpha^2, \delta\eta\delta\beta \rangle.$$

The decomposition matrix for $Q(3\mathcal{B})$ is given in Figure 12.

4.5. Blocks with quiver $3\mathcal{C}$.

$$3\mathcal{C} = \begin{array}{c} \bullet \\ \xrightarrow{\beta} 0 \xrightarrow{\delta} \bullet \\ \xleftarrow{\gamma} \bullet \xleftarrow{\eta} \bullet \\ \rho \end{array}$$

By [27, VI.5] (resp. by [27, IX.5.3 and p. 305]), there are no blocks B with dihedral (resp. generalized quaternion) defect groups that have Ext quiver $3\mathcal{C}$.

By [27, IX.5.3 and pp. 300–301], if B has semidihedral defect groups and Ext quiver $3\mathcal{C}$, then B is Morita equivalent to either $\text{SD}(3\mathcal{C})_{2,1} = k[3\mathcal{C}]/I_{\text{SD}(3\mathcal{C})_{2,1}}$ or $\text{SD}(3\mathcal{C})_{2,2} = k[3\mathcal{C}]/I_{\text{SD}(3\mathcal{C})_{2,2}}$ where

$$\begin{aligned} I_{\text{SD}(3\mathcal{C})_{2,1}} &= \langle \rho\beta, \delta\rho, \rho\eta, \gamma\rho, \beta\gamma - \eta\delta, (\beta\gamma)^2 - \rho^{2^{n-2}}, \delta\beta\gamma\beta, \gamma\eta\delta\eta \rangle, \\ I_{\text{SD}(3\mathcal{C})_{2,2}} &= \langle \rho\beta, \delta\rho, \rho\eta, \gamma\rho, \beta\gamma - \eta\delta, (\beta\gamma)^{2^{n-2}} - \rho^2, \delta\beta(\gamma\beta)^{2^{n-2}-1}, \\ &\quad \gamma\eta(\delta\eta)^{2^{n-2}-1} \rangle. \end{aligned}$$

The decomposition matrix for $\text{SD}(3\mathcal{C})_{2,1}$ is given in Figure 13, and the decomposition matrix for $\text{SD}(3\mathcal{C})_{2,2}$ is given in Figure 14.

4.6. Blocks with quiver $3\mathcal{D}$.

$$3\mathcal{D} = \begin{array}{c} \bullet \\ \xrightarrow{\beta} 1 \xrightarrow{\delta} 0 \xrightarrow{\delta} 2 \\ \xleftarrow{\gamma} \bullet \xleftarrow{\eta} \bullet \\ \alpha \quad \xi \end{array}$$

By [27, IX.5.1, IX.5.4 and p. 296] (resp. by [27, IX.5.1 and p. 306]), there are no blocks B with dihedral (resp. generalized quaternion) defect groups that have Ext quiver $3\mathcal{D}$.

By [27, p. 301], if B has semidihedral defect groups and Ext quiver $3\mathcal{D}$, then B is Morita equivalent to $\text{SD}(3\mathcal{D}) = k[3\mathcal{D}]/I_{\text{SD}(3\mathcal{D})}$ where

$$I_{\text{SD}(3\mathcal{D})} = \langle \xi\delta, \eta\xi, \delta\eta, \gamma\beta - \alpha^{2^{n-2}-1}, \alpha\gamma - \gamma\eta\delta, \beta\alpha - \eta\delta\beta, \xi^2 - \delta\beta\gamma\eta \rangle.$$

The decomposition matrix for $\text{SD}(3\mathcal{D})$ is given in Figure 10.

4.7. Blocks with quiver $3\mathcal{F}$. By [27, VI.5] (resp. by [27, VII.4]), there are no blocks B with dihedral (resp. generalized quaternion) defect groups that have Ext quiver $3\mathcal{F}$. By [27, IX.5.2 and p. 301], there are also no blocks B with semidihedral defect groups that have Ext quiver $3\mathcal{F}$.

4.8. Blocks with quiver $3\mathcal{H}$.

$$3\mathcal{H} = \begin{array}{c} \bullet \\ \xrightarrow{\beta} 0 \xrightarrow{\gamma} \bullet \\ \xleftarrow{\lambda} \bullet \xleftarrow{\eta} \bullet \\ \delta \end{array}$$


By [27, VI.5] (resp. by [27, VII.4]), there are no blocks B with dihedral (resp. generalized quaternion) defect groups that have Ext quiver $3\mathcal{H}$.

By [27, p. 301], if B has semidihedral defect groups and Ext quiver $3\mathcal{H}$, then B is Morita equivalent to $\text{SD}(3\mathcal{H})_1 = k[3\mathcal{H}]/I_{\text{SD}(3\mathcal{H})_1}$ or $\text{SD}(3\mathcal{H})_2 = k[3\mathcal{H}]/I_{\text{SD}(3\mathcal{H})_2}$ where

$$\begin{aligned} I_{\text{SD}(3\mathcal{H})_1} &= \langle \lambda\delta - \gamma\beta\gamma, \beta\lambda - \eta(\delta\eta)^{2^{n-2}-1}, \eta\delta\beta, \delta\beta\gamma, \gamma\eta \rangle, \\ I_{\text{SD}(3\mathcal{H})_2} &= \langle \lambda\delta - \gamma(\beta\gamma)^{2^{n-2}-1}, \beta\lambda - \eta\delta\eta, \eta\delta\beta, \delta\beta\gamma, \gamma\eta \rangle. \end{aligned}$$

The decomposition matrix for $\text{SD}(3\mathcal{H})_1$ is given in Figure 15, and the decomposition matrix for $\text{SD}(3\mathcal{H})_2$ is given in Figure 16.

4.9. Blocks with quiver $3\mathcal{K}$.



By [27, IX.5.2 and p. 302], there are no blocks B with semidihedral defect groups that have Ext quiver $3\mathcal{K}$.

By [27, p. 296], if B has dihedral defect groups and Ext quiver $3\mathcal{K}$, then B is Morita equivalent to $D(3\mathcal{K}) = k[3\mathcal{K}]/I_{D(3\mathcal{K})}$ where

$$I_{D(3\mathcal{K})} = \langle \delta\beta, \lambda\delta, \beta\lambda, \kappa\gamma, \eta\kappa, \gamma\eta, \gamma\beta - \lambda\kappa, \kappa\lambda - (\delta\eta)^{2^{n-2}}, (\eta\delta)^{2^{n-2}} - \beta\gamma \rangle.$$

The corresponding decomposition matrix is given in Figure 17.

By [27, p. 306], if B has generalized quaternion defect groups and Ext quiver $3\mathcal{K}$, then B is Morita equivalent to $Q(3\mathcal{K}) = k[3\mathcal{K}]/I_{Q(3\mathcal{K})}$ where

$$I_{Q(3\mathcal{K})} = \langle \delta\beta - \kappa\lambda\kappa, \gamma\eta - \lambda\kappa\lambda, \lambda\delta - \gamma\beta\gamma, \eta\kappa - \beta\gamma\beta, \beta\lambda - \eta(\delta\eta)^{2^{n-2}-1}, \kappa\gamma - \delta(\eta\delta)^{2^{n-2}-1}, \delta\beta\gamma, \gamma\eta\delta, \eta\kappa\lambda \rangle.$$

The decomposition matrix for $Q(3\mathcal{K})$ is given in Figure 18.

4.10. Blocks with quivers $3\mathcal{L}$, $3\mathcal{Q}$ or $3\mathcal{R}$. By [27, VIII.2.0] (resp. by [27, VII.4]), there are no blocks B with semidihedral (resp. generalized quaternion) defect groups that have Ext quivers $3\mathcal{L}$, $3\mathcal{Q}$ or $3\mathcal{R}$. By [27, IX.5.4], there are also no blocks B with dihedral defect groups that have Ext quivers $3\mathcal{L}$, $3\mathcal{Q}$ or $3\mathcal{R}$.

5. ORDINARY CHARACTERS BELONGING TO TAME BLOCKS

Assume Hypothesis 4.1. In particular, $p = 2$, $n \geq 2$, and D is dihedral, semidihedral or generalized quaternion.

For $2 \leq \ell \leq n-1$, define $\zeta_\ell = \zeta^{2^{n-1-\ell}}$, so that ζ_ℓ is a primitive 2^ℓ -th root of unity. It follows from [15, Sect. VII], [16, 37] and [28] (see below) that every ordinary irreducible character of G which belongs to B is realizable over $F(\zeta)$, i.e. it corresponds to an absolutely irreducible $F(\zeta)G$ -module. In particular, if we are only interested in the ordinary irreducible characters of G belonging to B , we can replace ξ in the paragraph before Remark 3.1 by ζ .

In [16] (resp. [37]), the ordinary irreducible characters of G belonging to B were analyzed if $n \geq 3$ and D is dihedral (resp. semidihedral or generalized quaternion). In the notation of [16, Sect. 4] (resp. [37, Sect. 2]), this means that we are either in Case (aa) or in Case (ab) or (ba), see [16, Thm. 2] (resp. [37, Thms. 3.14-3.17]). Note that Case (ba) can only occur when D is semidihedral. In particular, in Case (aa) (resp. Case (ab) or (ba)) there are precisely three (resp. two) isomorphism classes of simple B -modules.

Remark 5.1. (a) If D is dihedral of order 4, i.e. $n = 2$, it follows from [25] that B is Morita equivalent to either $D(3\mathcal{A})_1$ or $D(3\mathcal{K})$. Moreover, there are precisely 4 ordinary irreducible characters belonging to B and they all have height 0. In the decomposition matrices in Figures 6 and 17, these characters are $\chi_1, \chi_2, \chi_3, \chi_4$. By [15, Prop. (7D)] and [28], it follows that all these characters are realizable over F , i.e. they correspond to absolutely irreducible FG -modules.

(b) If D is quaternion of order 8, i.e. $n = 3$, it follows from [37, p. 220 and Thm. 3.17] that we are in Case (aa) and that there are precisely 4 (resp. 3) ordinary irreducible characters of height 0 (resp. $1 = n - 2$) belonging to B . Moreover, B is Morita equivalent to one of

the algebras in $\{Q(3\mathcal{A})_2, Q(3\mathcal{B}), Q(3\mathcal{K})\}$ and in the decomposition matrices in Figures 8, 12 and 18, the ordinary irreducible characters of height 0 (resp. 1) are

$$\chi_1, \chi_2, \chi_3, \chi_4 \quad (\text{resp. } \chi_{5,1}, \chi_6, \chi_7).$$

By [37, Prop. 4.2] and [28], all these characters are realizable over F , i.e. they correspond to absolutely irreducible FG -modules.

- (c) If D is not quaternion of order 8, then there are precisely 4 (resp. $2^{n-2} - 1$) ordinary irreducible characters of height 0 (resp. 1) belonging to B . If D is dihedral, these are all ordinary irreducible characters belonging to B . If D is semidihedral, there is 0 (resp. 1) additional ordinary irreducible character of height $n - 2$ belonging to B if we are in Case (ab) (resp. Cases (ba) or (aa)). If D is generalized quaternion of order ≥ 16 , there are 1 (resp. 2) additional ordinary irreducible characters of height $n - 2$ belonging to B if we are in Case (ab) (resp. Case (aa)). In the decomposition matrices in Figures 1–18, the ordinary irreducible characters of height 0 (resp. 1) are

$$\chi_1, \chi_2, \chi_3, \chi_4 \quad (\text{resp. } \chi_{5,i} \text{ for } 1 \leq i \leq 2^{n-2} - 1),$$

whereas the ordinary irreducible characters of height $n - 2$ are χ_6 or χ_6, χ_7 , provided they exist.

Let $n \geq 3$, and let σ be an element of order 2^{n-1} in D . By [16, 37], there exists a block b_σ of $kC_G(\sigma)$ with $b_\sigma^G = B$ which contains a unique irreducible Brauer character $\varphi^{(\sigma)}$ such that the following is true. There is an ordering of $(1, 2, \dots, 2^{n-2} - 1)$ such that for $1 \leq i \leq 2^{n-2} - 1$, the generalized decomposition number of G corresponding to σ , $\chi_{5,i}$ and $\varphi^{(\sigma)}$ has the form

$$(5.8) \quad d_{\chi_{5,i}, \varphi^{(\sigma)}}^{(\sigma)} = \begin{cases} \zeta^i + \zeta^{-i} & \text{if } D \text{ is dihedral or generalized quaternion,} \\ & \text{or if } D \text{ is semidihedral and } i \text{ is even,} \\ \zeta^i - \zeta^{-i} & \text{if } D \text{ is semidihedral and } i \text{ is odd with } i \leq 2^{n-3} - 1, \\ \zeta^{-i} - \zeta^i & \text{if } D \text{ is semidihedral and } i \text{ is odd with } i \geq 2^{n-3} + 1. \end{cases}$$

Note that the formulas in (5.8) for D semidihedral and i odd follow since the D -conjugacy classes of elements of order 2^{n-1} in D are represented by

$$\sigma^1, \sigma^3, \dots, \sigma^{2^{n-3}-1}, \quad \sigma^{-(2^{n-3}+1)}, \sigma^{-(2^{n-3}+3)}, \dots, \sigma^{-(2^{n-2}-1)}.$$

If D is quaternion of order 8, we have that $d_{\chi, \varphi^{(\sigma)}}^{(\sigma)} = \zeta + \zeta^{-1} = 0$ for $\chi \in \{\chi_{5,1}, \chi_6, \chi_7\}$.

For $2 \leq \ell \leq n - 2$ define $\nu_\ell = \zeta_\ell + \zeta_\ell^{-1}$, and define

$$(5.9) \quad \nu_{n-1} = \begin{cases} \zeta + \zeta^{-1} & \text{if } D \text{ is dihedral or generalized quaternion,} \\ \zeta - \zeta^{-1} & \text{if } D \text{ is semidihedral.} \end{cases}$$

Note that W contains all roots of unity of order not divisible by 2. Hence by [16, 37] and by [28], the ordinary irreducible characters of height 0 or $n - 2$ belonging to B correspond to simple FG -modules. On the other hand, the characters $\chi_{5,i}$, $i = 1, \dots, 2^{n-2} - 1$, fall into $n - 2$ Galois orbits $\mathcal{O}_2, \dots, \mathcal{O}_{n-1}$ under the action of $\text{Gal}(F(\nu_{n-1})/F)$. Namely for $2 \leq \ell \leq n - 1$,

$$\mathcal{O}_\ell = \{\chi_{5,2^{n-1-\ell}(2u-1)} \mid 1 \leq u \leq 2^{\ell-2}\}.$$

The field generated by the character values of each $\xi_\ell \in \mathcal{O}_\ell$ over F is $F(\nu_\ell)$. Hence by [28], each ξ_ℓ corresponds to an absolutely irreducible $F(\nu_\ell)G$ -module X_ℓ . By [31, Satz V.14.9], this implies that for $2 \leq \ell \leq n - 1$, the Schur index of each $\xi_\ell \in \mathcal{O}_\ell$ over F is 1. Hence we obtain $n - 2$ non-isomorphic simple FG -modules V_2, \dots, V_{n-1} with characters $\rho_2, \dots, \rho_{n-1}$ satisfying

$$(5.10) \quad \rho_\ell = \sum_{\xi_\ell \in \mathcal{O}_\ell} \xi_\ell = \sum_{u=1}^{2^{\ell-2}} \chi_{5,2^{n-1-\ell}(2u-1)} \quad \text{for } 2 \leq \ell \leq n - 1.$$

By [31, Hilfssatz V.14.7], $\text{End}_{FG}(V_\ell)$ is a commutative F -algebra isomorphic to the field generated over F by the character values of any $\xi_\ell \in \mathcal{O}_\ell$. This means

$$(5.11) \quad \text{End}_{FG}(V_\ell) \cong F(\nu_\ell) \quad \text{for } 2 \leq \ell \leq n-1.$$

Suppose $v_1, \dots, v_{l(\sigma)}$ form a complete system of representatives of $C_G(\sigma)$ -conjugacy classes of 2-regular elements in $C_G(\sigma)$ with $v_1 = 1_G$. By (3.6), for all $1 \leq i \leq 2^{n-2} - 1$, the generalized decomposition number of G corresponding to $\chi_{5,i}$, σ and $\varphi^{(\sigma)}$ can be written as a W -linear combination of $\chi_{5,i}(\sigma v_1), \dots, \chi_{5,i}(\sigma v_{l(\sigma)})$, say

$$(5.12) \quad d_{\chi_{5,i}, \varphi^{(\sigma)}}^{(\sigma)} = \tilde{w}_1 \cdot \chi_{5,i}(\sigma v_1) + \dots + \tilde{w}_{l(\sigma)} \cdot \chi_{5,i}(\sigma v_{l(\sigma)})$$

for certain $\tilde{w}_1, \dots, \tilde{w}_{l(\sigma)} \in W$.

By [16, Thm. 5] and [37, Prop. 4.6], the characters $\chi_{5,i}$ have the same degree x and they are all of height 1 for $1 \leq i \leq 2^{n-2} - 1$. Hence $x = 2^{a-n+1}x^*$ where $\#G = 2^a \cdot g^*$ and x^* and g^* are odd. Since the centralizer $C_G(\sigma)$ contains $\langle \sigma \rangle$, we have $\#C_G(\sigma) = 2^{n-1} \cdot 2^b \cdot m^*$ where $b \geq 0$ and m^* is odd.

For $1 \leq j \leq l(\sigma)$, let C_j be the conjugacy class in G of σv_j , and let $t(C_j) \in WG$ be the class sum of C_j . Since for all j , the centralizer $C_G(\sigma v_j)$ contains $\langle \sigma \rangle$, we have $\#C_G(\sigma v_j) = 2^{n-1} \cdot 2^{b_j} \cdot m_j^*$ where $b_j \geq 0$ and m_j^* is odd. We want to determine the action of $t(C_j)$ on V_ℓ for $2 \leq \ell \leq n-1$. For this, we identify $\text{End}_{FG}(V_\ell) \cong F(\nu_\ell)$ with $\text{End}_{F(\nu_\ell)G}(X_\ell)$ for one particular absolutely irreducible $F(\nu_\ell)G$ -constituent X_ℓ of V_ℓ with character ξ_ℓ . Using (5.10), we choose $\xi_\ell = \chi_{5,2^{n-1-\ell}}$. Then for $2 \leq \ell \leq n-1$, the action of $t(C_j)$ on V_ℓ is given as multiplication by $\mu_j(\ell)$, where

$$(5.13) \quad \mu_j(\ell) = \frac{\#C_j}{\xi_\ell(1)} \cdot \xi_\ell(\sigma v_j) = 2^{-b_j} \frac{g^*}{m_j^* \cdot x^*} \cdot \xi_\ell(\sigma v_j).$$

For $1 \leq j \leq l(\sigma)$, define $w_j = 2^{b_j} \frac{m_j^* \cdot x^*}{g^*} \tilde{w}_j$. Then $w_j \in W$ since g^* is odd, and w_j does not depend on ℓ . By (5.8) and (5.9),

$$d_{\xi_\ell, \varphi^{(\sigma)}}^{(\sigma)} = d_{\chi_{5,2^{n-1-\ell}}, \varphi^{(\sigma)}}^{(\sigma)} = \nu_\ell.$$

Therefore, (5.12) and (5.13) imply that

$$(5.14) \quad \nu_\ell = w_1 \cdot \mu_1(\ell) + \dots + w_{l(\sigma)} \cdot \mu_{l(\sigma)}(\ell)$$

where $w_1, \dots, w_{l(\sigma)} \in W$ are independent of $\ell \in \{2, \dots, n-1\}$.

Remark 5.2. If B is a principal block, the above formulas simplify considerably due to the fact that there is very little fusion of D -conjugacy classes in G in this case. More precisely, following Brauer's arguments in [13, Sect. VII], suppose D is a Sylow 2-subgroup of G and let $S = \langle \sigma \rangle$. If σ^λ is not in the center of D , then S is a Sylow 2-subgroup of $C_G(\sigma^\lambda)$. Hence if σ^λ and σ^μ are conjugate in G , we can use Sylow's theorems to see that they are conjugate in $N_G(S)$. Since $N_G(S)/C_G(S)$ is a 2-group, it must be of order 2. Thus $N_G(S)$ is generated by D and $C_G(S)$, which implies that σ^λ and σ^μ are conjugate in D . Using (3.7) together with [16, Prop. (4A)] and [37, Prop. 2.10], we obtain that

$$(5.15) \quad \chi_{5,i}(\sigma) = d_{\chi_{5,i}, \varphi^{(\sigma)}}^{(\sigma)} \varphi^{(\sigma)}(1_G)$$

where $\varphi^{(\sigma)}$ is as in (5.8). Moreover, since $\varphi^{(\sigma)}$ is the unique Brauer character belonging to the block b_σ of $kC_G(\sigma)$ satisfying $b_\sigma^G = B$, it follows by Brauer's Third Main Theorem (see e.g. [1, Thm. 16.1]) that b_σ is the principal block of $kC_G(\sigma)$, which implies that $\varphi^{(\sigma)}$ is the trivial character of b_σ . Putting (5.15) into (5.13) for $j = 1$ therefore implies that if $\omega = 2^{b_1} \frac{m_1^* \cdot x^*}{g^*}$, which lies in W , then for all $2 \leq \ell \leq n-1$

$$\nu_\ell = d_{\chi_{5,2^{n-1-\ell}}, \varphi^{(\sigma)}}^{(\sigma)} = \omega \cdot \mu_1(\ell).$$

Note that in [3, Sect. 3.4], [4, Sect. 5] and [9, Sect. 3.2], it was assumed that the formula (5.15) was also true for non-principal blocks. However, this may not be true since there could be more fusion of D -conjugacy classes in G in this case. More precisely, let Y be a full set of representatives

of D -conjugacy classes of the elements of the set $\{\sigma^r \mid r \text{ odd}\}$, and let Y_σ be the set of all $y \in Y$ which are conjugate to σ in G . If B is not principal, then $|Y_\sigma|$ may be strictly greater than 1. By [16, Prop. (4A)] and [37, Prop. 2.10], the set $\{(y, b_\sigma) \mid y \in Y_\sigma\}$ is a system of representatives for the conjugacy classes of subsections for B such that y is conjugate to σ in G . Hence it follows from Remark 3.1 that for non-principal blocks B , (5.15) has to be replaced by the formula

$$\chi_{5,i}(\sigma) = \sum_{y \in Y_\sigma} d_{\chi_{5,i}, \varphi^{(\sigma)}}^{(y)} \varphi^{(\sigma)}(1_G).$$

Definition 5.3. Use the notation introduced above, and in particular (5.9). Assume $n \geq 3$.

(i) Define

$$q_n(t) = \prod_{\ell=2}^{n-1} \text{min.pol.}_F(\nu_\ell)$$

and let $R' = W[[t]]/(q_n(t))$.

(ii) Let $Z = \langle \sigma \rangle$, so that Z is a cyclic group of order $2^{n-1} > 2$. Let $\tau : Z \rightarrow Z$ be the group automorphism which sends σ to σ^{-1} if D is dihedral or generalized quaternion, and which sends σ to $\sigma^{-1+2^{n-2}}$ if D is semidihedral. Then τ can be extended to a W -algebra automorphism of the group ring WZ which will again be denoted by τ . Let $T(\sigma^2) = 1 + \sigma^2 + \sigma^4 + \dots + \sigma^{2^{n-1}-2}$, and define

$$S' = (WZ)^{\langle \tau \rangle} / (T(\sigma^2), \sigma T(\sigma^2)).$$

Remark 5.4. The minimal polynomial $\text{min.pol.}_F(\nu_\ell)$ for $2 \leq \ell \leq n-1$ is as follows:

$$\begin{aligned} \text{min.pol.}_F(\nu_2)(t) &= t, \\ \text{min.pol.}_F(\nu_\ell)(t) &= \text{min.pol.}_F(\nu_{\ell-1})(t^2 - 2) \quad \text{for } 3 \leq \ell \leq n-2, \\ \text{min.pol.}_F(\nu_{n-1})(t) &= \begin{cases} \text{min.pol.}_F(\nu_{n-2})(t^2 - 2) & \text{if } D \text{ is dihedral or} \\ & \text{generalized quaternion,} \\ \text{min.pol.}_F(\nu_{n-2})(t^2 + 2) & \text{if } D \text{ is semidihedral.} \end{cases} \end{aligned}$$

The W -algebra R' from Definition 5.3 is a complete local commutative Noetherian ring with residue field k . Moreover,

$$\begin{aligned} F \otimes_W R' &\cong \prod_{\ell=2}^{n-1} F(\nu_\ell) \quad \text{as } F\text{-algebras,} \\ k \otimes_W R' &\cong k[t]/(t^{2^n-2}-1) \quad \text{as } k\text{-algebras.} \end{aligned}$$

Additionally, R' is isomorphic to the W -subalgebra of $\prod_{\ell=2}^{n-1} W[\nu_\ell]$ generated by the element $(\nu_\ell)_{\ell=2}^{n-1}$.

Lemma 5.5. *Using the notation of Definition 5.3, there is a continuous W -algebra isomorphism $h : R' \rightarrow S'$ with $h(t) = \sigma + \tau(\sigma)$. In particular, R' is isomorphic to a subquotient algebra of the group algebra WD .*

Moreover, if D is dihedral or semidihedral, then the ring $W[[t]]/(t q_n(t), 2 q_n(t))$ is also isomorphic to a subquotient algebra of the group algebra WD .

Proof. If D is dihedral or generalized quaternion, this follows from [3, Lemma 2.3.6] and [4, Lemma 5.3].

For the remainder of the proof, assume that D is semidihedral (in particular, $n \geq 4$). Let $J = \sigma^{2^{n-2}}$ so that $\tau(\sigma) = J\sigma^{-1}$. Note that $(WZ)^{\langle \tau \rangle}$ is generated as a W -algebra by $(\sigma + J\sigma^{-1})$ and J . Moreover, $(WZ)^{\langle \tau \rangle}$ is a free W -module of rank $2^{n-2} + 1$ with W -basis given by

$$(5.16) \quad \begin{aligned} &(\sigma^{\pm 1} + J\sigma^{\mp 1}), (\sigma^{\pm 3} + J\sigma^{\mp 3}), \dots, (\sigma^{\pm(2^{n-3}-1)} + J\sigma^{\mp(2^{n-3}-1)}), \\ &1, J, (\sigma^2 + \sigma^{-2}), (\sigma^4 + \sigma^{-4}), \dots, (\sigma^{2^{n-2}-2} + \sigma^{-(2^{n-2}-2)}). \end{aligned}$$

Note that the W -subalgebra of $(WZ)^{\langle \tau \rangle}$ generated by $(\sigma + J\sigma^{-1})$ is a free W -module of the same rank $2^{n-2} + 1$ and with almost the same W -basis as in (5.16) except that J must be replaced by $2J$. It follows that $S' = (WZ)^{\langle \tau \rangle} / (T(\sigma^2), \sigma T(\sigma^2))$ is generated as a W -algebra by the image of $(\sigma + J\sigma^{-1})$ in S' . Hence we have a surjective W -algebra homomorphism

$$f : W[[t]] \rightarrow S' = (WZ)^{\langle \tau \rangle} / (T(\sigma^2), \sigma T(\sigma^2))$$

sending t to the image in S' of $\sigma + J\sigma^{-1} = \sigma + \tau(\sigma)$. Using the injective W -algebra homomorphism

$$\begin{aligned} \iota : WZ &\rightarrow W \times W \times \prod_{\ell=2}^{n-1} W[\zeta_\ell] \\ \sigma &\mapsto \left(1, -1, (\zeta_\ell)_{\ell=2}^{n-1} \right) \end{aligned}$$

it is straightforward to prove that there exists an odd integer c_n such that $q_n(\sigma + J\sigma^{-1}) = c_n \sigma T(\sigma^2)$. More precisely, $c_4 = 3$ and $c_n = 2c_{n-1}^2 - 1$ for $n \geq 5$. Thus $q_n(t)$ lies in the kernel of f . Since both $R' = W[[t]]/(q_n(t))$ and S' are free as W -modules of the same rank $2^{n-2} - 1$, it follows that $R' = W[[t]]/(q_n(t)) \cong S'$ as W -algebras.

To finish the proof of Lemma 5.5, it suffices to show that the ring $W[[t]]/((t-2)q_n(t))$ is isomorphic to a subquotient algebra of WD . Define

$$\Theta = (WZ)^{\langle \tau \rangle} / (T(\sigma^2) - \sigma T(\sigma^2)).$$

Then Θ is isomorphic to a subquotient algebra of WD and it is generated as a W -algebra by the image of $(\sigma + J\sigma^{-1})$ in Θ . Moreover, Θ is a free W -module of rank 2^{n-2} , since the ideal $(T(\sigma^2) - \sigma T(\sigma^2))$ is generated over W by $T(\sigma^2) - \sigma T(\sigma^2)$. Define a surjective W -algebra homomorphism $\theta : W[[t]] \rightarrow \Theta$ by sending t to the image in Θ of $\sigma + J\sigma^{-1} = \sigma + \tau(\sigma)$. Using the above calculations, we see that

$$\begin{aligned} \theta((t-2)q_n(t)) &= ((\sigma + J\sigma^{-1}) - 2)q_n(\sigma + J\sigma^{-1}) \\ &= ((\sigma + J\sigma^{-1}) - 2)c_n \sigma T(\sigma^2) \\ &= 2c_n [T(\sigma^2) - \sigma T(\sigma^2)] \end{aligned}$$

which is zero in Θ . Hence $(t-2)q_n(t)$ lies in the kernel of θ . Since both $W[[t]]/((t-2)q_n(t))$ and Θ are free over W of rank 2^{n-2} , it follows that $W[[t]]/((t-2)q_n(t))$ is isomorphic to Θ , which completes the proof of Lemma 5.5. \square

The following result gives a correction of [4, Lemma 5.4] and generalizes the corrected result to all tame blocks with at least two isomorphism classes of simple modules.

Lemma 5.6. *Assume Hypothesis 4.1 and use the notation introduced above, and in particular (5.9), (5.10) and Definition 5.3. Assume $n \geq 3$. If D is quaternion of order 8, let U' be a WG -module which is free over W and whose F -character is either $\chi_{5,1}$, or χ_6 , or χ_7 . If D is not quaternion of order 8, let U' be a WG -module which is free over W and whose F -character is equal to*

$$\sum_{\ell=2}^{n-1} \rho_\ell = \sum_{i=1}^{2^{n-2}-1} \chi_{5,i}.$$

- (i) *There exists a WG -module endomorphism α of U' such that the W -algebra $W[\alpha]$ generated by α is isomorphic to R' .*
- (ii) *Suppose either that $n = 3$, or that $n \geq 4$ and $\text{End}_{kG}(U'/2U') \cong R'/2R'$ and $U'/2U'$ is free as a module for $\text{End}_{kG}(U'/2U')$ of rank $\deg(\chi_{5,1})$. Then $\text{End}_{WG}(U') = W[\alpha] \cong R'$ and U' is free as a module for $\text{End}_{WG}(U')$.*

Proof. Suppose first that $n = 3$. Then $q_n(t) = t$ and $R' \cong W$. Hence we can choose the WG -module endomorphism α of U' to be the zero endomorphism. Since U' is free as a W -module and $F \otimes_W \text{End}_{WG}(U') \cong \text{End}_{FG}(F \otimes_W U') \cong F$, it follows that $\text{End}_{WG}(U') \cong W \cong R'$.

For the remainder of the proof, assume $n \geq 4$. We first construct a WG -module endomorphism α of U' as in part (i) of the lemma. As before, let $\sigma \in D$ be an element of order 2^{n-1} and let $\{v_1, \dots, v_{l(\sigma)}\}$ be a complete system of representatives of $C_G(\sigma)$ -conjugacy classes of 2-regular elements in $C_G(\sigma)$. For $1 \leq j \leq l(\sigma)$, let C_j be the conjugacy class in G of σv_j , and let $t(C_j) \in WG$ be the class sum of C_j .

Let $1 \leq j \leq l(\sigma)$. Since $t(C_j)$ lies in the center of WG , multiplication by $t(C_j)$ defines a WG -module endomorphism of U' . Since U' is free as a W -module, $\text{End}_{WG}(U')$ can naturally be identified with a W -subalgebra of

$$F \otimes_W \text{End}_{WG}(U') \cong \text{End}_{FG}(F \otimes_W U') \cong \prod_{\ell=2}^{n-1} \text{End}_{FG}(V_\ell) \cong \prod_{\ell=2}^{n-1} F(\nu_\ell).$$

Therefore, $t(C_j)$ acts on U' as multiplication by a scalar λ_j in the maximal W -order $\prod_{\ell=2}^{n-1} W[\nu_\ell]$ in $\prod_{\ell=2}^{n-1} F(\nu_\ell)$. Moreover, λ_j can be determined from the action of $t(C_j)$ on $F \otimes_W U' \cong \bigoplus_{\ell=2}^{n-1} V_\ell$, which implies by (5.13) that $\lambda_j = (\mu_j(\ell))_{\ell=2}^{n-1}$. By (5.14), there exist elements $w_1, \dots, w_{l(\sigma)} \in W$ such that

$$\nu_\ell = w_1 \cdot \mu_1(\ell) + \dots + w_{l(\sigma)} \cdot \mu_{l(\sigma)}(\ell)$$

for $2 \leq \ell \leq n-1$. Define α to be the WG -module endomorphism of U' given by multiplication by the element $\sum_{j=1}^{l(\sigma)} w_j t(C_j)$ in the center of WG . Then α acts on U' as multiplication by the scalar $(\nu_\ell)_{\ell=2}^{n-1} \in \prod_{\ell=2}^{n-1} W[\nu_\ell]$, which implies $W[\alpha] \cong R'$ (see Remark 5.4).

Let now $\overline{U'} = U'/2U'$ and suppose that $\text{End}_{kG}(\overline{U'}) \cong R'/2R' \cong k[t]/(t^{2^{n-2}-1})$ and that $\overline{U'}$ is free as a module for $\text{End}_{kG}(U'/2U')$ of rank $x = \deg(\chi_{5,1})$. Note that $x = \deg(\chi_{5,i})$ for all $1 \leq i \leq 2^{n-2} - 1$ and that $\dim_{F(\nu_\ell)} V_\ell = x$ for all $2 \leq \ell \leq n-1$, which implies that $F \otimes_W U'$ is a free module of rank x for $\text{End}_{FG}(F \otimes_W U')$. We have a short exact sequence of W -modules

$$0 \longrightarrow \text{End}_W(U') \xrightarrow{\cdot 2} \text{End}_W(U') \xrightarrow{\text{mod } 2} \text{End}_k(\overline{U'}) \longrightarrow 0.$$

Considering the G -action on these modules, we obtain an exact sequence of W -modules

$$(5.17) \quad 0 \longrightarrow \text{End}_{WG}(U') \xrightarrow{\cdot 2} \text{End}_{WG}(U') \xrightarrow{\text{mod } 2} \text{End}_{kG}(\overline{U'}) \longrightarrow H^1(G, \text{End}_W(U')).$$

Because $\text{End}_{WG}(U')$ is a W -submodule of the free W -module $\text{End}_W(U')$, it follows that $\text{End}_{WG}(U')$ is a free W -module of rank

$$\dim_F(F \otimes_W \text{End}_{WG}(U')) = \dim_F \text{End}_{FG}(F \otimes_W U') = 2^{n-2} - 1.$$

Since by assumption $\dim_k \text{End}_{kG}(\overline{U'}) = \dim_k(R'/2R') = 2^{n-2} - 1$, it follows that (5.17) induces a short exact sequence of W -modules

$$(5.18) \quad 0 \longrightarrow \text{End}_{WG}(U') \xrightarrow{\cdot 2} \text{End}_{WG}(U') \xrightarrow{\text{mod } 2} \text{End}_{kG}(\overline{U'}) \longrightarrow 0.$$

Let $\overline{\beta}$ be a generator of $\text{End}_{kG}(\overline{U'})$ as a k -algebra. By (5.18), there exists $\beta \in \text{End}_{WG}(U')$ whose induced kG -module endomorphism of $\overline{U'}$ is equal to $\overline{\beta}$. Using Nakayama's lemma, we see that $\text{End}_{WG}(U') = W[\beta]$. Since the maximal ideal $\mathfrak{m}_{W[\beta]}$ of $W[\beta]$ is generated by 2 and β and the maximal ideal $\mathfrak{m}_{k[\overline{\beta}]}$ of $k[\overline{\beta}]$ is generated by $\overline{\beta}$, it follows that

$$U'/\mathfrak{m}_{W[\beta]}U' \cong \overline{U'}/\overline{\beta}(\overline{U'}) \cong \overline{U'}/\mathfrak{m}_{k[\overline{\beta}]} \overline{U'}$$

where the latter has k -dimension $x = \deg(\chi_{5,1})$ by assumption. By Nakayama's lemma, we can lift a k -basis $\{\overline{s}_1, \dots, \overline{s}_x\}$ of $U'/\mathfrak{m}_{W[\beta]}U'$ to a set $\{s_1, \dots, s_x\}$ of generators of U' over $W[\beta] = \text{End}_{WG}(U')$. Because $F \otimes_W U'$ is a free module of rank x for $\text{End}_{FG}(F \otimes_W U') \cong F \otimes_W \text{End}_{WG}(U')$, it follows that s_1, \dots, s_x are linearly independent over $\text{End}_{WG}(U')$. Hence U' is free as a module for $\text{End}_{WG}(U')$ of rank $x = \deg(\chi_{5,1})$.

It remains to show that $\text{End}_{WG}(U') = W[\alpha]$, i.e. we need to show $W[\alpha] = W[\beta]$. We identify both $W[\alpha]$ and $W[\beta]$ with W -subalgebras of the maximal W -order $\prod_{\ell=2}^{n-1} W[\nu_\ell]$ in $\prod_{\ell=2}^{n-1} F(\nu_\ell) \cong$

$F \otimes_W \text{End}_{WG}(U')$. Since $W[\alpha] \subseteq W[\beta]$, there exists a polynomial $q(X) \in W[X]$ such that $\alpha = q(\beta)$. Because α is not a unit, the constant coefficient of $q(X)$ must be divisible by 2. Write

$$q(X) = 2\tilde{a}_0 + a_1X + a_2X^2 + \cdots + a_dX^d$$

for certain $\tilde{a}_0, a_1, \dots, a_d \in W$ and $d \geq 1$. Suppose $a_1 = 2\tilde{a}_1$ for some $\tilde{a}_1 \in W$. Consider the natural projection

$$\pi_{n-1} : \prod_{\ell=2}^{n-1} W[\nu_\ell] \rightarrow W[\nu_{n-1}],$$

and let \mathfrak{m} be the maximal ideal of $W[\nu_{n-1}]$. Since $n \geq 4$, either $\sqrt{2}$ or $\sqrt{-2}$ lies in \mathfrak{m} . Hence $2 \in \mathfrak{m}^2$. Since $2\pi_{n-1}(\beta), \pi_{n-1}(\beta)^2, \dots, \pi_{n-1}(\beta)^d$ all lie in \mathfrak{m}^2 , it follows that $\pi_{n-1}(\alpha) \in \mathfrak{m}^2$. However, we have seen that $\pi_{n-1}(\alpha) = \nu_{n-1} \notin \mathfrak{m}^2$. Hence a_1 is not divisible by 2. But then the kG -module endomorphism $\bar{\alpha}$ of $\overline{U'}$ which is induced by α has the form

$$\bar{\alpha} = \bar{a}_1\bar{\beta} + \bar{a}_2\bar{\beta}^2 + \cdots + \bar{a}_d\bar{\beta}^d$$

where $\bar{a}_1 \in k^*$ and $\bar{a}_2, \dots, \bar{a}_d \in k$. This means that $k[\bar{\alpha}] = k[\bar{\beta}] = \text{End}_{kG}(\overline{U'})$, which implies by Nakayama's lemma that $\text{End}_{WG}(U') = W[\alpha]$. \square

6. UNIVERSAL DEFORMATION RINGS

Assume Hypothesis 4.1 and the notation introduced in Section 4 and Section 5. In particular, $p = 2$, $n \geq 2$, and D is dihedral, semidihedral or generalized quaternion. In this section, we want to determine $R(G, V)$ for any maximally ordinary kG -module V in the sense of Definition 3.2.

We first determine all indecomposable kG -modules V belonging to B whose stable endomorphism rings are isomorphic to k and whose Brauer characters are restrictions of ordinary irreducible characters of height 1. By Remark 5.1, the ordinary irreducible characters of height 1 belonging to B are (using the notation from Section 5):

- none if $n = 2$,
- $\chi_{5,1}, \chi_6, \chi_7$ if D is quaternion of order 8, and
- $\chi_{5,i}$ for all $1 \leq i \leq 2^{n-2} - 1$ if $n \geq 3$ and D is not quaternion of order 8.

Lemma 6.1. *Assume Hypothesis 4.1, and assume $n \geq 3$. Let $\Lambda = kQ/I$ be a basic algebra such that B is Morita equivalent to Λ . For each vertex j in Q , let T_j denote the simple B -module corresponding to the simple Λ -module belonging to j . Let V be an indecomposable kG -module belonging to B such that $\underline{\text{End}}_{kG}(V) \cong k$ and such that the Brauer character of V is equal to the restriction to the 2-regular conjugacy classes of an ordinary irreducible character of G of height 1.*

- (a) *Suppose D is quaternion of order 8. If $Q = 3A$ or $Q = 3B$ then V is either isomorphic to T_1 or T_2 , or V is a uniserial module of length 4 whose radical quotient or socle is isomorphic to T_0 . If $Q = 3K$ then V is an arbitrary uniserial module of length 2.*
- (b) *Suppose D is not quaternion of order 8.*
 - (i) *If $Q = 2A$ then V is a uniserial module with descending composition factors T_0, T_0, T_1 , or T_1, T_0, T_0 .*
 - (ii) *If $Q = 2B$ and B is Morita equivalent to neither $\text{SD}(2B)_4(c)$ nor $\text{Q}(2B)_2(p, a, c)$ then V is isomorphic to T_1 . If B is Morita equivalent to $\text{SD}(2B)_4(c)$ or $\text{Q}(2B)_2(p, a, c)$ then V is a uniserial module with descending composition factors T_0, T_1 , or T_1, T_0 .*
 - (iii) *If $Q = 3A$ then V is a uniserial module with descending composition factors T_0, T_1, T_0, T_2 , or T_2, T_0, T_1, T_0 , or T_0, T_2, T_0, T_1 , or T_1, T_0, T_2, T_0 .*
 - (iv) *If $Q \in \{3B, 3D\}$ then V is isomorphic to T_1 .*
 - (v) *If B is Morita equivalent to $\text{SD}(3C)_{2,1}$ then V is isomorphic to T_0 . If B Morita equivalent to $\text{SD}(3C)_{2,2}$ then V is indecomposable with descending radical factors $T_0, T_1 \oplus T_2$, or $T_1 \oplus T_2, T_0$, or V is uniserial with descending composition factors T_1, T_0, T_2 , or T_2, T_0, T_1 .*

- (vi) If $Q \in \{3\mathcal{H}, 3\mathcal{K}\}$ and B is not Morita equivalent to $\text{SD}(3\mathcal{H})_2$ then V is a uniserial module with descending composition factors T_1, T_2 , or T_2, T_1 . If B is Morita equivalent to $\text{SD}(3\mathcal{H})_2$ then V is a uniserial module with descending composition factors T_0, T_1 , or T_1, T_0 .

Conversely, if V is as in (a) or (b), then $\text{End}_{kG}(V) \cong k$ and the Brauer character of V is equal to the restriction to the 2-regular conjugacy classes of an ordinary irreducible character of height 1. Moreover, $\text{Ext}_{kG}^1(V, V) = 0$ if D is quaternion of order 8, and $\text{Ext}_{kG}^1(V, V) \cong k$ in all other cases.

Proof. Lemma 6.1 is proved using the description of the basic algebra $\Lambda = kQ/I$ of the block B together with its decomposition matrix as provided in Section 4. To give an idea of the arguments, we discuss the case when $\Lambda = \text{SD}(3\mathcal{A})_1$ in part (b)(iii). It follows from the decomposition matrix in Figure 7 that V is an indecomposable B -module with composition factors T_0, T_0, T_1, T_2 . Let t_V (resp. s_V) be the composition series length of the radical quotient (resp. socle) of V . If $t_V = 3$ then V has radical series length 2 and $s_V = 1$. But then V is a submodule of a projective indecomposable B -module which is impossible for $t_V = 3$. If $t_V = 2$ then V has radical series length at most 3 and $s_V \leq 2$. Using that $\text{Ext}_B^1(T_i, T_j)$ is one-dimensional over k if $(i, j) \in \{(0, 1), (0, 2), (1, 0), (2, 0)\}$ and zero otherwise, it follows that there are no indecomposable B -modules V with $t_V = 2$ and radical series length 2. Analyzing $\text{Ext}_B^1(T_{i_1} \oplus T_{i_2}, V')$ for appropriate indecomposable B -modules V' of length 2, we see that the only indecomposable B -modules V with $t_V = 2$ and radical series length 3 are submodules of the projective cover P_{T_0} of the form

$$\begin{array}{ccc} T_0 & & T_0 \\ T_1 & T_2 & \text{or} & T_1 & T_2 \\ T_0 & & & T_0 & \end{array} .$$

However, the stable endomorphism ring of each of these modules has k -dimension 2. If $t_V = 1$ then V has radical series length at most 4 and $s_V \leq 3$. Since V is a quotient module of a projective indecomposable B -module in this case, there are no indecomposable B -modules V with $t_V = 1$ and radical series length 2. If $t_V = 1$ and the radical series length of V is 3, we use similar Ext^1 arguments as above to see that the only indecomposable B -modules V with these properties are quotient modules of the projective cover P_{T_0} with descending radical factors $T_0, T_1 \oplus T_2, T_0$. But all such modules have an endomorphism which factors through T_0 and which does not factor through a projective module, meaning that the stable endomorphism ring of each of these modules has k -dimension 2. Finally, if $t_V = 1$ and the radical series length of V is 4, then V is one of the uniserial modules described in part (b)(iii) of Lemma 6.1. This description shows directly that $\text{End}_B(V) \cong k$. Moreover, using the projective indecomposable B -modules, we see that $\text{Ext}_B^1(V, V) \cong k$. \square

Corollary 6.2. *Assume Hypothesis 4.1. Let V be an indecomposable kG -module belonging to B such that $\underline{\text{End}}_{kG}(V) \cong k$. The following statements are equivalent:*

- (i) V is maximally ordinary in the sense of Definition 3.2;
- (ii) $n \geq 4$ and the Brauer character of V is equal to the restriction to the 2-regular conjugacy classes of an ordinary irreducible character of G of height 1;
- (iii) $n \geq 4$ and V is as in part (b) of Lemma 6.1.

Proof. If $n = 2$ (resp. $n = 3$), it follows from [15, Sect. VII] (resp. [16, 37]) that all generalized decomposition numbers corresponding to maximal 2-power order elements in D lie in $\{0, \pm 1\}$. Hence there are no maximally ordinary kG -modules belonging to B if $n \leq 3$.

Now suppose $n \geq 4$. Let $\sigma \in D$ be of maximal 2-power order, and let χ be an ordinary irreducible character belonging to B . It follows from [16, 37] that if χ has height 0, then the non-zero generalized decomposition numbers corresponding to σ and χ are ± 1 . Also, if χ has height $n - 2$, then all generalized decomposition numbers corresponding to σ and χ are zero. On the other hand, we see from (5.8), since $n \geq 4$, that if χ has height 1, then $d_{\chi, \varphi(\sigma)}^{(\sigma)} \notin \{0, \pm 1\}$ if and only if $\chi = \chi_{5,i}$ for $i \in \{1, 2, \dots, 2^{n-2} - 1\} - \{2^{n-3}\}$. Note that the restriction of $\chi_{5,i}$ to the 2-regular conjugacy classes is the same for all $1 \leq i \leq 2^{n-2} - 1$. Therefore it follows that if $n \geq 4$ then V

is maximally ordinary if and only if its Brauer character is equal to the restriction to the 2-regular conjugacy classes of an ordinary irreducible character of height 1. Hence Corollary 6.2 follows from Lemma 6.1. \square

The lifts of some of the kG -modules V from Lemma 6.1 are connected to 3-tubes in the stable Auslander-Reiten quiver of B as follows.

Definition 6.3. Assume Hypothesis 4.1, assume $n \geq 3$, and let V be as in Lemma 6.1. We say V corresponds to a 3-tube if there exists an indecomposable quotient module \overline{U} of the projective kG -module cover P_V of V such that

- (a) \overline{U} defines a lift of V over $k[t]/(t^{2^n-2})$, and
- (b) \overline{U} belongs to a 3-tube of the stable Auslander-Reiten quiver of B .

The following lemma determines which modules V from Lemma 6.1 correspond to 3-tubes.

Lemma 6.4. Assume Hypothesis 4.1, assume $n \geq 3$, and let V be as in Lemma 6.1. If D is dihedral then V corresponds to a 3-tube, and if D is generalized quaternion then V does not correspond to a 3-tube. If D is semidihedral then V corresponds to a 3-tube if and only if either

- B is Morita equivalent to one of the algebras in

$$\{\text{SD}(2\mathcal{A})_2(c), \text{SD}(2\mathcal{B})_1(c), \text{SD}(3\mathcal{B})_1, \text{SD}(3\mathcal{C})_{2,1}\}$$

and V is arbitrary; or

- B is Morita equivalent to $\text{SD}(3\mathcal{A})_1$ and V is such that its radical quotient or its socle is isomorphic to T_1 ; or
- B is Morita equivalent to $\text{SD}(3\mathcal{C})_{2,2}$ and V is such that its radical quotient or its socle is isomorphic to T_0 ; or
- B is Morita equivalent to $\text{SD}(3\mathcal{H})_1$ and V is such that its radical quotient is isomorphic to T_1 ;
- B is Morita equivalent to $\text{SD}(3\mathcal{H})_2$ and V is such that its radical quotient is isomorphic to T_0 .

In all cases, if V corresponds to a 3-tube, then \overline{U} from Definition 6.3 belongs to the boundary of its 3-tube.

Proof. If D is generalized quaternion, Lemma 6.4 follows from [27, V.4.3].

Suppose next that D is dihedral, and let $\Lambda = kQ/I$ be a basic algebra such that B is Morita equivalent to Λ . By [27, VI.10.1], it follows that $\Lambda/\text{soc}(\Lambda)$ is special biserial. Therefore, we can use the techniques described in [18, Sect. 3] to determine the 3-tubes of the stable Auslander-Reiten quiver of B . Note that the modules at the boundaries of the 3-tubes are either maximal uniserial or simple. Using the list of the possible V in Lemma 6.1, we see by direct inspection that V always corresponds to a 3-tube and that the module \overline{U} from Definition 6.3 always belongs to the boundary of its 3-tube.

Finally, suppose that D is semidihedral. By [27, V.4.2], the stable Auslander-Reiten quiver of B has at most one 3-tube. To give an idea of the arguments, we discuss the case when B is Morita equivalent to $\Lambda = \text{SD}(3\mathcal{A})_1$. By Lemma 6.1(b)(iii), V is a uniserial module with descending composition factors T_0, T_1, T_0, T_2 , or T_2, T_0, T_1, T_0 , or T_0, T_2, T_0, T_1 , or T_1, T_0, T_2, T_0 . Since V is uniserial with $\text{End}_{kG}(V) \cong k \cong \text{Ext}_{kG}^1(V, V)$, we can use [4, Lemma 2.5] to show that $R(G, V)/2R(G, V)$ is isomorphic to $k[t]/(t^{2^n-2})$ (resp. $k[t]/(t^{2^n-2}-1)$) if the radical quotient or the socle of V is isomorphic to T_1 (resp. T_2). Hence by Definition 6.3, V can only correspond to a 3-tube if the radical quotient or the socle of V is isomorphic to T_1 . Since the projective cover P_{T_1} is uniserial, we obtain from [4, Lemma 2.5] that $\overline{U} = \Omega^{-1}(T_1)$ (resp. $\overline{U} = \Omega(T_1)$) defines a lift of V over $k[t]/(t^{2^n-2})$ if the radical quotient (resp. the socle) of V is isomorphic to T_1 . Since $\Omega^3(\overline{U}) \cong \overline{U}$, it follows that \overline{U} belongs to a 3-tube of the stable Auslander-Reiten quiver of B . Moreover, using the criterion given in [18, Sect. 1] for almost split sequences to have an indecomposable middle term, we see that \overline{U} belongs to the boundary of this 3-tube. \square

The following result shows that if V corresponds to a 3-tube, then the module \overline{U} from Definition 6.3 has a universal deformation ring $R(G, \overline{U}) \cong k$.

Proposition 6.5. *Assume Hypothesis 4.1, assume $n \geq 3$, and let V be as in Lemma 6.1. Moreover suppose that V corresponds to a 3-tube. Let \overline{U} be the kG -module from Definition 6.3 which belongs to a 3-tube of the stable Auslander-Reiten quiver of B . Then $\underline{\text{End}}_{kG}(\overline{U}) \cong k$ and $R(G, \overline{U}) \cong k$.*

Proof. If D is dihedral, this follows from [3, Sect. 5.2] and [9, Prop. 6.3]. For the remainder of the proof, assume that D is semidihedral.

Let \mathfrak{T} be the 3-tube of the stable Auslander-Reiten quiver of B to which \overline{U} belongs. By Lemma 6.4, \overline{U} belongs to the boundary of \mathfrak{T} . Going through the cases described in Lemma 6.4, it is straightforward to show that $\underline{\text{End}}_{kG}(\overline{U}) \cong k$ and $\text{Ext}_{kG}^1(\overline{U}, \overline{U}) = 0$.

Let K be a vertex of \overline{U} . Because of the work in [27, Chapter V], and in particular [27, V.4.2.1 and proof of V.4.2], we have the following facts:

- (i) The group K is a Klein four group and the quotient group $N_G(K)/C_G(K)$ is isomorphic to a symmetric group S_3 .
- (ii) There is a block b of $kN_G(K)$ with $b^G = B$ such that the Green correspondent $f\overline{U}$ of \overline{U} belongs to the boundary of a 3-tube in the stable Auslander-Reiten quiver of b . Moreover, b is Morita equivalent to kS_4 modulo the socle.

Using these facts in lieu of [3, Facts 5.2.1], we can use similar arguments as in the proofs of [3, Prop. 5.2.4 and Cor. 5.2.5] to show that $R(G, \overline{U}) \cong k$. \square

By Corollary 6.2, Theorem 1.1 is a consequence of the following result.

Theorem 6.6. *Assume Hypothesis 4.1, and assume $n \geq 3$. Let V be an indecomposable kG -module belonging to B such that $\underline{\text{End}}_{kG}(V) \cong k$ and such that the Brauer character of V is equal to the restriction to the 2-regular conjugacy classes of an ordinary irreducible character of G of height 1. Then*

$$R(G, V) \cong \begin{cases} W[[t]]/(t q_n(t), 2 q_n(t)) & \text{if } V \text{ corresponds to a 3-tube,} \\ W[[t]]/(q_n(t)) & \text{if } V \text{ does not correspond to a 3-tube,} \end{cases}$$

where $q_n(t)$ is as in Definition 5.3. In all cases, the ring $R(G, V)$ is a subquotient ring of WD , and it is a complete intersection ring if and only if V does not correspond to a 3-tube.

Proof. By Lemma 6.1, V is one of the modules in Lemma 6.1, listed in part (a) or part (b).

Suppose first that D is quaternion of order 8. By Lemma 6.1(a), $\text{Ext}_{kG}^1(V, V) = 0$, which implies that $R(G, V)$ is isomorphic to a quotient algebra of W . Using the decomposition matrices in Figures 8, 12 and 18 together with [22, Prop. (23.7)], we see that in all cases V can be lifted over W , which implies that $R(G, V) \cong W \cong W[[t]]/(q_3(t))$.

For the remainder of the proof, assume that D is not quaternion of order 8. In particular, the Brauer character of V is the restriction of $\chi_{5,1}$ to the 2-regular conjugacy classes of G . By Lemma 6.1(b), $\text{Ext}_{kG}^1(V, V) \cong k$, which implies that $R(G, V)$ is isomorphic to a quotient algebra of $W[[t]]$ but not to a quotient algebra of W . Let P_V be the projective kG -module cover of V .

Claim 0. There exists an indecomposable quotient module \overline{U}' of P_V such that \overline{U}' defines a lift of V over $k[t]/(t^{2^{n-2}-1})$ and $\overline{U}'/t^2\overline{U}'$ is an indecomposable kG -module. Moreover, if V corresponds to a 3-tube then the indecomposable kG -module \overline{U} of Definition 6.3, which is also an indecomposable quotient module of P_V , defines a lift of V over $k[t]/(t^{2^{n-2}})$ and satisfies $\overline{U}/t^{2^{n-2}-1}\overline{U} \cong \overline{U}'$.

Proof of Claim 0. Suppose first that B is Morita equivalent neither to $\text{SD}(2\mathcal{B})_4(c)$, nor to $\text{Q}(2\mathcal{B})_2(p, a, c)$, nor to $\text{SD}(3\mathcal{C})_{2,2}$. Then V is uniserial of length $\ell \leq 4$. Using the quiver and relations of the basic algebra of B , as provided in Section 4, it follows that in all cases P_V has an indecomposable quotient module \overline{U}' (resp. \overline{U} , provided it exists) which is uniserial of length $\ell(2^{n-2}-1)$ (resp. $\ell 2^{n-2}$). Moreover, \overline{U}' (resp. \overline{U}) can be pictured as having $2^{n-2}-1$ (resp. 2^{n-2}) copies of V stacked on top of each other. This means that the action of t on \overline{U}' (resp. \overline{U}) is given

by an automorphism of \overline{U}' (resp. \overline{U}) which is unique up to multiplication by a non-zero scalar and which factors through $\text{rad}^\ell(\overline{U}')$ (resp. $\text{rad}^\ell(\overline{U})$). This implies Claim 0 in this case.

If B is Morita equivalent to $\text{SD}(2\mathcal{B})_4(c)$ or $\text{Q}(2\mathcal{B})_2(p, a, c)$, it follows from Lemma 6.1(b)(ii) that V is uniserial of length 2 with descending composition factors T_u, T_v , where $\{u, v\} = \{0, 1\}$. Using the relations in Section 4.2, we see that the projective cover $P_V = P_{T_u}$ has a unique submodule K_u which is uniserial with descending composition factors T_u, T_u . It follows that $\overline{U}' = P_{T_u}/K_u$, which can be visualized as in (6.19), where T_u (resp. T_v) occurs $2^{n-2} - 1$ times. This implies Claim 0 in this case, since V does not correspond to a 3-tube, so \overline{U} does not exist.

$$(6.19) \quad \overline{U}' = \begin{array}{ccc} & T_u & \\ & & \\ & T_u & T_v \\ & \times & \times \\ & T_u & T_v \\ & \times & \times \\ & \vdots & \vdots \\ & & \\ & T_u & T_v \\ & \times & \times \\ & T_u & T_v \\ & & \\ & T_v & \end{array}$$

Finally, suppose B is Morita equivalent to $\text{SD}(3\mathcal{C})_{2,2}$. By Lemma 6.1(b)(v), V is indecomposable with descending radical factors $T_0, T_1 \oplus T_2$, or $T_1 \oplus T_2, T_0$, or V is uniserial with descending composition factors T_1, T_0, T_2 , or T_2, T_0, T_1 . Using the relations in Section 4.5, we see that there exists a unique uniserial B -module T_{00} , up to isomorphism, with descending composition factors T_0, T_0 . Also, for $u \in \{1, 2\}$, there exists a unique uniserial B -module T_{u0u} , up to isomorphism, with descending composition factors T_u, T_0, T_u . Moreover, the projective cover P_{T_0} has a unique submodule isomorphic to T_{00} , and for $u \in \{1, 2\}$, the projective cover P_{T_u} has a unique submodule isomorphic to T_{u0u} . If the radical quotient (resp. socle) of V is T_0 , it follows that \overline{U} is isomorphic to $\Omega^{-1}(T_{00})$ (resp. $\Omega(T_{00})$), and \overline{U}' is isomorphic to $\text{rad}^2(\overline{U})$. In particular, the action of t on \overline{U}' (resp. \overline{U}) is given by an automorphism of \overline{U}' (resp. \overline{U}) which is unique up to multiplication by a non-zero scalar and which factors through $\text{rad}^2(\overline{U}')$ (resp. $\text{rad}^2(\overline{U})$). If the radical quotient of V is T_u for $u \in \{1, 2\}$, it follows that \overline{U} does not exist and \overline{U}' is isomorphic to $\Omega^{-1}(T_{u0u})$. This completes the proof of Claim 0.

Claim 1. The universal mod 2 deformation ring $R(G, V)/2R(G, V)$ is isomorphic to $k[t]/(t^{2^{n-2}})$ (resp. $k[t]/(t^{2^{n-2}-1})$) and the universal mod 2 deformation of V is isomorphic to \overline{U} (resp. \overline{U}') if V corresponds to a 3-tube (resp. does not correspond to a 3-tube).

Proof of Claim 1. Suppose first that B is Morita equivalent neither to $\text{SD}(2\mathcal{B})_4(c)$, nor to $\text{Q}(2\mathcal{B})_2(p, a, c)$, nor to $\text{SD}(3\mathcal{C})_{2,2}$. As seen in the proof of Claim 0, V is uniserial in this case. Moreover, it is straightforward to check that the projective cover of V satisfies the hypotheses of [4, Lemma 2.5]. Hence we can use [4, Lemma 2.5] to prove Claim 1 in this case.

If B is Morita equivalent to $\text{SD}(2\mathcal{B})_4(c)$ or $\text{Q}(2\mathcal{B})_2(p, a, c)$, it follows from Lemma 6.1(b)(ii) that V is uniserial with descending composition factors T_u, T_v where $\{u, v\} = \{0, 1\}$. As seen in the proof of Claim 0, the projective cover $P_V = P_{T_u}$ has a unique submodule K_u which is uniserial with descending composition factors T_u, T_u , and $\overline{U}' = P_{T_u}/K_u$ can be visualized as in (6.19). Using the relations in Section 4.2, we see that there is a unique B -submodule V' of \overline{U}' which is isomorphic to V . Moreover, if $T = \overline{U}'/V'$ then there is a unique B -submodule T' of \overline{U}' such that there are kG -module isomorphisms $\varphi : \overline{U}'/T' \rightarrow V$ and $\psi : \overline{U}'/V' = T \rightarrow T'$. Since $\text{Ext}_{kG}^1(\overline{U}', V) = 0$ and since every surjective kG -module homomorphism $\overline{U}' \rightarrow V$ must have kernel equal to T , we can argue as in the proof of [4, Lemma 2.5] to show that $R(G, V)/2R(G, V)$ is isomorphic to $k[t]/(t^{2^{n-2}-1})$ and that the universal mod 2 deformation of V is isomorphic to \overline{U}' . This implies Claim 1 in this case.

If B is Morita equivalent to $\text{SD}(3\mathcal{C})_{2,2}$, we use again similar arguments as in the proof of [4, Lemma 2.5] to prove Claim 1. The main point is that, as seen in the proof of Claim 0, $\overline{U'}$ and \overline{U} are suitable submodules (resp. suitable quotient modules) of the projective cover P_{T_0} if the radical quotient (resp. socle) of V is T_0 , and $\overline{U'}$ is a suitable quotient module of P_{T_u} if the radical quotient of V is T_u for $u \in \{1, 2\}$. This completes the proof of Claim 1.

Claim 2. In all cases, $\text{End}_{kG}(\overline{U'}) \cong k[t]/(t^{2^{n-2}-1})$.

Proof of Claim 2. Suppose first that B is Morita equivalent neither to $\text{SD}(2\mathcal{B})_4(c)$, nor to $\text{Q}(2\mathcal{B})_2(p, a, c)$, nor to $\text{SD}(3\mathcal{C})_{2,2}$. As seen in the proof of Claim 0, $\overline{U'}$ is a uniserial module which can be pictured as having $2^{n-2} - 1$ copies of V stacked on top of each other. Since $\overline{U'}$ is isomorphic to a tree module in the sense of [32], we can use the main result of [32] to prove Claim 2.

If B is Morita equivalent to $\Lambda \in \{\text{SD}(2\mathcal{B})_4(c), \text{Q}(2\mathcal{B})_2(p, a, c), \text{SD}(3\mathcal{C})_{2,2}\}$, we use the relations in Section 4.2 and Section 4.5 to analyze the possible kG -module endomorphisms of $\overline{U'}$, where we use the description of $\overline{U'}$ as given in the proof of Claim 0. Using explicit k -bases for the Λ -module $\overline{U'_\Lambda}$ corresponding to $\overline{U'}$ under the Morita equivalence, a straightforward linear algebra calculation shows that $\text{End}_\Lambda(\overline{U'_\Lambda}) \cong k[t]/(t^{2^{n-2}-1})$. This proves Claim 2.

Claim 3. In all cases, $\overline{U'}$ has a lift U' over W such that the F -character of U' is equal to

$$\sum_{\ell=2}^{n-1} \rho_\ell = \sum_{i=1}^{2^{n-2}-1} \chi_{5,i}$$

where ρ_ℓ is as in (5.10).

Proof of Claim 3. In all cases, we use the description of $\overline{U'}$ as given in the proof of Claim 0. Suppose first that either B is not Morita equivalent to any of the algebras in

$$\{\text{SD}(2\mathcal{A})_1(c), \text{Q}(2\mathcal{A})(c), \text{SD}(3\mathcal{A})_1, \text{Q}(3\mathcal{A})_2\}$$

and V is arbitrary or B is Morita equivalent to $\text{SD}(3\mathcal{A})_1$ and V is such that its radical quotient (resp. socle) is isomorphic to T_1 . Then Claim 3 follows by using the decomposition matrix of B together with [22, Prop. (23.7)].

Suppose next that B is Morita equivalent to one of the algebras in $\{\text{SD}(2\mathcal{A})_1(c), \text{Q}(2\mathcal{A})(c)\}$. If the radical quotient (resp. socle) of V is isomorphic to T_1 , then $\Omega(\overline{U'}) \cong Z_{001}$ (resp. $\Omega^{-1}(\overline{U'}) \cong Z_{100}$) where

$$Z_{001} = \begin{array}{cc} & T_1 \\ T_1 & T_0 \\ & T_1 \end{array}, \quad Z_{100} = \begin{array}{cc} & T_1 \\ T_0 & T_0 \\ T_1 & T_1 \end{array}$$

and Z_{001} is a submodule of P_{T_1} and Z_{100} is a quotient module of P_{T_1} . Using [3, Lemma 2.3.2], it follows that Z_{001} and Z_{100} each have a lift over W such that the F -character of this lift is $\chi_3 + \chi_4 + \chi_6$ (using the notation in Figure 2), which implies Claim 3 in this case.

Finally, suppose that B is Morita equivalent to $\text{SD}(3\mathcal{A})_1$ and $u = 2$ and $v = 1$, or B is Morita equivalent to $\text{Q}(3\mathcal{A})_2$ and $\{u, v\} = \{1, 2\}$. If V is such that its radical quotient (resp. socle) is isomorphic to T_u then $\Omega(\overline{U'}) = Z_{0v0u}$ (resp. $\Omega^{-1}(\overline{U'}) = Z_{u0v0}$) where

$$Z_{0v0u} = \begin{array}{ccc} & & T_u \\ & & T_0 \\ T_u & T_v & \\ & T_0 & \\ & & T_u \end{array}, \quad Z_{u0v0} = \begin{array}{ccc} & & T_u \\ & & T_0 \\ T_u & T_v & \\ & T_0 & \\ & & T_u \end{array}$$

and Z_{0v0u} is a submodule of P_{T_u} and Z_{u0v0} is a quotient module of P_{T_u} . Using [3, Lemma 2.3.2], it follows that Z_{0v0u} and Z_{u0v0} each have a lift over W such that the F -character of this lift is

$\chi_2 + \chi_3 + \chi_6$ if B is Morita equivalent to $\text{SD}(3\mathcal{A})_1$ (using the notation in Figure 7) and the F -character of this lift is $\chi_2 + \chi_4 + \chi_6$ (resp. $\chi_2 + \chi_3 + \chi_7$) if B is Morita equivalent to $\text{Q}(3\mathcal{A})_2$ and $u = 1$ (resp. $u = 2$) (using the notation in Figure 8). This completes the proof of Claim 3.

Claim 4. The universal deformation ring $R(G, V)$ is as stated in Theorem 6.6.

Proof of Claim 4. In all cases, it follows by Lemma 5.6 that U' from Claim 3 is an $R'G$ -module. More precisely, there exists a WG -module endomorphism α of U' such that $W[\alpha] \cong R'$. By Claim 2, we have $\text{End}_{kG}(\overline{U}') \cong k[t]/(t^{2^{n-2}-1}) \cong R'/2R'$. Moreover, since \overline{U}' is a lift of V over $k[t]/(t^{2^{n-2}-1})$, \overline{U}' is free as a module for $\text{End}_{kG}(\overline{U}')$ of rank $\dim_k V = \deg(\chi_{5,1})$. Hence it follows by Lemma 5.6 that $\text{End}_{WG}(U') = W[\alpha] \cong R'$ and U' is free as a module for $\text{End}_{WG}(U')$.

In other words, U' defines a lift of V over R' . Let $\tau : R(G, V) \rightarrow R'$ be the unique continuous W -algebra homomorphism relative to the lift defined by U' . Since $R'/(\mathfrak{m}_{R'}^2 + 2R') \cong k[t]/(t^2)$, τ is surjective if and only if $R'/(\mathfrak{m}_{R'}^2 + 2R') \otimes_{R'} U'$ does not define the trivial lift of V over $k[t]/(t^2)$. However,

$$R'/(\mathfrak{m}_{R'}^2 + 2R') \otimes_{R'} U' \cong U'/(\alpha^2(U') + 2U') \cong \overline{U}'/\overline{\alpha}^2(\overline{U}') \cong \overline{U}'/t^2\overline{U}'$$

is an indecomposable kG -module, which implies that this does not define the trivial lift of V over $k[t]/(t^2)$. Hence τ is surjective and induces a surjective k -algebra homomorphism

$$\overline{\tau} : R(G, V)/2R(G, V) \rightarrow R'/2R'.$$

Suppose first that V does not correspond to a 3-tube. Then $R(G, V)/2R(G, V)$ and $R'/2R'$ are isomorphic and finite dimensional over k , which implies that $\overline{\tau}$ is an isomorphism. Because R' is a free W -module of finite rank, it follows that τ is an isomorphism. By Lemma 5.5, R' is isomorphic to a subquotient ring of WD . This proves Claim 4 and completes the proof of Theorem 6.6 if V does not correspond to a 3-tube.

Suppose next that V corresponds to a 3-tube. Then the universal mod 2 deformation ring $R(G, V)/2R(G, V)$ is isomorphic to $k[t]/(2^{n-2})$ and the universal mod 2 deformation is isomorphic to \overline{U} . By [3, Lemma 2.3.3], it follows that $R(G, V) \cong W[[t]]/(q_n(t)(t - 2\mu), a2^m q_n(t))$ for certain $\mu \in W$, $a \in \{0, 1\}$ and $0 < m \in \mathbb{Z}$. If $a = 0$ then $R(G, V) \cong W[[t]]/(q_n(t)(t - 2\mu))$ is free over W . If $a = 1$ then $R(G, V)/2^m R(G, V) \cong (W/2^m W)[[t]]/(q_n(t)(t - 2\mu))$ is free over $W/2^m W$. Therefore it follows that if $a = 0$ (resp. $a = 1$), then there is a lift of \overline{U} , when regarded as a kG -module, over W (resp. over $W/2^m W$). But by Proposition 6.5 we have $R(G, \overline{U}) \cong k$, which means we must have $a = 1$ and $m = 1$. Since V corresponds to a 3-tube, D is dihedral or semidihedral. Hence by Lemma 5.5, the ring $W[[t]]/(t q_n(t), 2 q_n(t))$ is isomorphic to a subquotient ring of WD . This proves Claim 4 and completes the proof of Theorem 6.6 if V corresponds to a 3-tube. \square

REFERENCES

- [1] J. L. Alperin, Local representation theory. Modular representations as an introduction to the local representation theory of finite groups. Cambridge Studies in Advanced Mathematics, vol. 11, Cambridge University Press, Cambridge, 1986.
- [2] F. M. Bleher, Universal deformation rings for dihedral 2-groups. J. London Math Soc. 79, (2009), 225–237.
- [3] F. M. Bleher, Universal deformation rings and dihedral defect groups. Trans. Amer. Math. Soc. 361 (2009), 3661–3705.
- [4] F. M. Bleher, Universal deformation rings and generalized quaternion defect groups. Adv. Math. 225 (2010), 1499–1522.
- [5] F. M. Bleher and T. Chinburg, Universal deformation rings and cyclic blocks. Math. Ann. 318 (2000), 805–836.
- [6] F. M. Bleher and T. Chinburg, Universal deformation rings need not be complete intersections. C. R. Math. Acad. Sci. Paris 342 (2006), 229–232.
- [7] F. M. Bleher and T. Chinburg, Universal deformation rings need not be complete intersections. Math. Ann. 337 (2007), 739–767.
- [8] F. M. Bleher, T. Chinburg and B. de Smit, Inverse Problems for deformation rings. Accepted by Trans. Amer. Math. Soc., 2012, [arXiv:1012.1290](https://arxiv.org/abs/1012.1290)
- [9] F. M. Bleher, G. Lloset and J. B. Schaefer, Universal deformation rings and dihedral blocks with two simple modules. J. Algebra 345 (2011), 49–71.
- [10] V. M. Bondarenko and J. A. Drozd, The representation type of finite groups. Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 71 (1977), 24–41. English translation: J. Soviet Math. 20 (1982), 2515–2528.

- [11] R. Brauer, On the Connection Between the Ordinary and The Modular Characters of Groups of Finite Order. *Ann. Math.* (2) (1942), 926–935.
- [12] R. Brauer, Zur Darstellungstheorie der Gruppen endlicher Ordnung. II. *Math. Zeitschr.* 72 (1959), 25–46.
- [13] R. Brauer, Some applications of the theory of blocks of characters of finite groups. III. *J. Algebra* 3 (1966), 225–255.
- [14] R. Brauer, On blocks and sections in finite groups. II. *Amer. J. Math.* 90 (1968), 895–925.
- [15] R. Brauer, Some applications of the theory of blocks of characters of finite groups. IV. *J. Algebra* 17 (1971), 489–521.
- [16] R. Brauer, On 2-blocks with dihedral defect groups. *Symposia Mathematica*, vol. XIII (Convegno di Gruppi e loro Rappresentazioni, INDAM, Rome, 1972), pp. 367–393, Academic Press, London, 1974.
- [17] S. Brenner, Modular representations of p groups. *J. Algebra* 15 (1970) 89–102.
- [18] M. C. R. Butler and C. M. Ringel, Auslander-Reiten sequences with few middle terms and applications to string algebras. *Comm. Algebra* 15 (1987), 145–179.
- [19] J. F. Carlson and J. Thévenaz, The classification of endo-trivial modules. *Invent. Math.* 158 (2004), 389–411.
- [20] J. F. Carlson and J. Thévenaz, The classification of torsion endo-trivial modules. *Ann. of Math.* (2) 162 (2005), 823–883.
- [21] T. Chinburg, Can deformation rings of group representations not be local complete intersections? In: *Problems from the Workshop on Automorphisms of Curves*. Edited by Gunther Cornelissen and Frans Oort, with contributions by I. Bouw, T. Chinburg, Cornelissen, C. Gasbarri, D. Glass, C. Lehr, M. Matignon, Oort, R. Pries and S. Wewers. *Rend. Sem. Mat. Univ. Padova* 113 (2005), 129–177.
- [22] C. W. Curtis and I. Reiner, *Methods of representation theory*. Vols. I and II. With applications to finite groups and orders. John Wiley & Sons, Inc., New York, 1981 and 1987.
- [23] B. de Smit and H. W. Lenstra, Explicit construction of universal deformation rings. In: *Modular Forms and Fermat's Last Theorem* (Boston, 1995), Springer-Verlag, Berlin-Heidelberg-New York, 1997, pp. 313–326.
- [24] F. Eisele, p -adic lifting problems and derived equivalences. *J. Algebra* 356 (2012), 90–114.
- [25] K. Erdmann, Blocks whose defect groups are Klein four groups: a correction. *J. Algebra* 76 (1982), 505–518.
- [26] K. Erdmann, Algebras and semidihedral defect groups I. *Proc. London Math. Soc.* (3) 57 (1988), 109–150.
- [27] K. Erdmann, *Blocks of Tame Representation Type and Related Algebras*. *Lecture Notes in Mathematics*, vol. 1428, Springer-Verlag, Berlin-Heidelberg-New York, 1990.
- [28] P. Fong, A note on splitting fields of representations of finite groups. *Illinois J. Math.* 7 (1963), 515–520.
- [29] D. Higman, Indecomposable representations at characteristic p . *Duke Math. J.* 21 (1954), 377–381.
- [30] T. Holm, Derived equivalence classification of algebras of dihedral, semidihedral, and quaternion type. *J. Algebra* 211 (1999), 159–205.
- [31] B. Huppert, *Endliche Gruppen. I. Die Grundlehren der Mathematischen Wissenschaften, Band 134*, Springer-Verlag, Berlin-New York, 1967.
- [32] H. Krause, Maps between tree and band modules. *J. Algebra* 137 (1991), 186–194.
- [33] M. Linckelmann, A derived equivalence for blocks with dihedral defect groups. *J. Algebra* 164 (1994), 244–255.
- [34] M. Linckelmann, The source algebras of blocks with a Klein four defect group, *J. Algebra* 167 (1994), 821–854.
- [35] M. Linckelmann, Introduction to fusion systems. In: *Group representation theory*, EPFL Press, Lausanne, 2007, pp. 79–113.
- [36] B. Mazur, Deforming Galois representations. In: *Galois groups over \mathbb{Q}* (Berkeley, 1987), Springer-Verlag, Berlin-Heidelberg-New York, 1989, pp. 385–437.
- [37] J. B. Olsson, On 2-blocks with quaternion and quasidihedral defect groups. *J. Algebra* 36 (1975), 212–241.

Frauke M. Bleher
 Department of Mathematics
 University of Iowa
 Iowa City, IA 52242-1419
 U.S.A.
 frauke-bleher@uiowa.edu

APPENDIX: DECOMPOSITION MATRICES FOR THE ALGEBRAS IN SECTION 4

FIGURE 1. The decomposition matrix for blocks of type $D(2\mathcal{A})$ or $SD(2\mathcal{A})_2(c)$.

$$\begin{array}{c}
\varphi_0 \varphi_1 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i}
\end{array}
\begin{array}{c}
\left[\begin{array}{cc}
1 & 0 \\
1 & 0 \\
1 & 1 \\
1 & 1 \\
2 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 2. The decomposition matrix for blocks of type $SD(2\mathcal{A})_1(c)$ or $Q(2\mathcal{A})(c)$.

$$\begin{array}{c}
\varphi_0 \varphi_1 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\left[\begin{array}{cc}
1 & 0 \\
1 & 0 \\
1 & 1 \\
1 & 1 \\
2 & 1 \\
0 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 3. The decomposition matrix for blocks of type $D(2\mathcal{B})$ or $SD(2\mathcal{B})_1(c)$.

$$\begin{array}{c}
\varphi_0 \varphi_1 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i}
\end{array}
\begin{array}{c}
\left[\begin{array}{cc}
1 & 0 \\
1 & 0 \\
1 & 1 \\
1 & 1 \\
0 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 4. The decomposition matrix for blocks of type $SD(2\mathcal{B})_2(c)$ or $Q(2\mathcal{B})_1(c)$.

$$\begin{array}{c}
\varphi_0 \varphi_1 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\left[\begin{array}{cc}
1 & 0 \\
1 & 0 \\
1 & 1 \\
1 & 1 \\
0 & 1 \\
2 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 5. The decomposition matrix for blocks of type $\text{SD}(2\mathcal{B})_4(c)$ or $\text{Q}(2\mathcal{B})_2(p, a, c)$.

$$\begin{array}{c}
\varphi_0 \ \varphi_1 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i}
\end{array}
\begin{array}{c}
\left[\begin{array}{cc}
1 & 0 \\
1 & 0 \\
0 & 1 \\
0 & 1 \\
1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 6. The decomposition matrix for blocks of type $\text{D}(3\mathcal{A})_1$.

$$\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i}
\end{array}
\begin{array}{c}
\left[\begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 1 \\
1 & 0 & 1 \\
1 & 1 & 0 \\
2 & 1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 7. The decomposition matrix for blocks of type $\text{SD}(3\mathcal{A})_1$.

$$\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\left[\begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 1 \\
1 & 0 & 1 \\
1 & 1 & 0 \\
2 & 1 & 1 \\
0 & 0 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 8. The decomposition matrix for blocks of type $\text{Q}(3\mathcal{A})_2$.

$$\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6 \\
\chi_7
\end{array}
\begin{array}{c}
\left[\begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 1 \\
1 & 0 & 1 \\
1 & 1 & 0 \\
2 & 1 & 1 \\
0 & 1 & 0 \\
0 & 0 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 9. The decomposition matrix for blocks of type $D(3\mathcal{B})_1$.

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i}
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[\begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 0 \\
1 & 0 & 1 \\
1 & 1 & 1 \\
0 & 1 & 0
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 10. The decomposition matrix for blocks of type $SD(3\mathcal{B})_1$ or $SD(3\mathcal{D})$.

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[\begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 0 \\
1 & 0 & 1 \\
1 & 1 & 1 \\
0 & 1 & 0 \\
0 & 0 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 11. The decomposition matrix for blocks of type $SD(3\mathcal{B})_2$.

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[\begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 0 \\
1 & 1 & 1 \\
1 & 0 & 1 \\
0 & 1 & 0 \\
2 & 1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 12. The decomposition matrix for blocks of type $Q(3\mathcal{B})$.

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6 \\
\chi_7
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[\begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 0 \\
1 & 1 & 1 \\
1 & 0 & 1 \\
0 & 1 & 0 \\
0 & 0 & 1 \\
2 & 1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 13. The decomposition matrix for blocks of type $\text{SD}(3\mathcal{C})_{2,1}$.

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[\begin{array}{ccc}
0 & 1 & 0 \\
1 & 1 & 0 \\
1 & 0 & 1 \\
0 & 0 & 1 \\
1 & 0 & 0 \\
1 & 1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 14. The decomposition matrix for blocks of type $\text{SD}(3\mathcal{C})_{2,2}$.

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[\begin{array}{ccc}
0 & 1 & 0 \\
1 & 0 & 1 \\
1 & 1 & 0 \\
0 & 0 & 1 \\
1 & 1 & 1 \\
1 & 0 & 0
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 15. The decomposition matrix for blocks of type $\text{SD}(3\mathcal{H})_1$.

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[\begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 1 \\
0 & 1 & 0 \\
0 & 0 & 1 \\
0 & 1 & 1 \\
1 & 1 & 0
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 16. The decomposition matrix for blocks of type $\text{SD}(3\mathcal{H})_2$.

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[\begin{array}{ccc}
1 & 0 & 0 \\
0 & 1 & 0 \\
1 & 1 & 1 \\
0 & 0 & 1 \\
1 & 1 & 0 \\
0 & 1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 17. The decomposition matrix for blocks of type $D(3\mathcal{K})$.

$$\begin{array}{c}
 \chi_1 \\
 \chi_2 \\
 \chi_3 \\
 \chi_4 \\
 \chi_{5,i}
 \end{array}
 \begin{array}{c}
 \varphi_0 \ \varphi_1 \ \varphi_2 \\
 \left[\begin{array}{ccc}
 1 & 0 & 0 \\
 1 & 1 & 1 \\
 0 & 1 & 0 \\
 0 & 0 & 1 \\
 0 & 1 & 1
 \end{array} \right]
 \end{array}
 \quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 18. The decomposition matrix for blocks of type $Q(3\mathcal{K})$.

$$\begin{array}{c}
 \chi_1 \\
 \chi_2 \\
 \chi_3 \\
 \chi_4 \\
 \chi_{5,i} \\
 \chi_6 \\
 \chi_7
 \end{array}
 \begin{array}{c}
 \varphi_0 \ \varphi_1 \ \varphi_2 \\
 \left[\begin{array}{ccc}
 1 & 0 & 0 \\
 1 & 1 & 1 \\
 0 & 1 & 0 \\
 0 & 0 & 1 \\
 0 & 1 & 1 \\
 1 & 0 & 1 \\
 1 & 1 & 0
 \end{array} \right]
 \end{array}
 \quad 1 \leq i \leq 2^{n-2} - 1$$