

# BRAUER'S GENERALIZED DECOMPOSITION NUMBERS AND UNIVERSAL DEFORMATION RINGS

FRAUKE M. BLEHER

**ABSTRACT.** We study the problem of lifting to local rings certain mod 2 representations  $V$  of finite groups  $G$  which belong to 2-modular tame blocks  $B$  of  $G$  having at least two isomorphism classes of simple modules. Green's lifting theorem determines when such a  $V$  may be lifted to the ring of infinite Witt vectors. We generalize this result by determining the full universal deformation ring of such  $V$  using Brauer's generalized decomposition numbers.

## 1. INTRODUCTION

Let  $k$  be an algebraically closed field of positive characteristic  $p$ , let  $W = W(k)$  be the ring of infinite Witt vectors over  $k$ , and let  $G$  be a finite group. An important question in the representation theory of  $G$  is whether a finitely generated  $kG$ -module  $V$  can be lifted to  $W$ . For example, Green's lifting theorem shows that this is possible if there are no non-trivial 2-extensions of  $V$  by itself. The most natural generalization of such results is to determine the full versal deformation ring  $R(G, V)$  of  $V$ . The topological ring  $R(G, V)$  is characterized by the property that every lift of  $V$  over a complete local commutative Noetherian ring  $R$  with residue field  $k$  arises from a local ring homomorphism  $\alpha : R(G, V) \rightarrow R$  and that  $\alpha$  is unique if  $R$  is the ring of dual numbers  $k[\epsilon]/(\epsilon^2)$ . In case  $\alpha$  is unique for all  $R$ ,  $R(G, V)$  is called the universal deformation ring of  $V$ . For more precise definitions, see §2. Note that all these rings  $R$ , including  $R(G, V)$ , have a natural structure as  $W$ -algebras. It was shown in [5, Prop. 2.1] that if the stable endomorphism ring  $\underline{\text{End}}_{kG}(V)$  is isomorphic to  $k$ , then the versal deformation ring  $R(G, V)$  is always universal.

In this paper, we determine  $R(G, V)$  for certain mod 2 representations  $V$  of finite groups  $G$ . The  $V$  we consider are those for which Brauer's generalized decomposition numbers carry the most information. Namely, their Brauer characters are restrictions of the ordinary irreducible characters for which Brauer's generalized decomposition numbers are generically not zero on maximal 2-power order elements. We will show that this non-triviality provides a powerful tool for computing universal deformation rings. Brauer generalized the usual decomposition numbers to be able to express the values of ordinary irreducible characters on  $p$ -singular elements in terms of Brauer characters. These generalized decomposition numbers enable us to find a family of Galois orbits of ordinary irreducible characters of  $G$  which can be used to construct lifts of  $V$  to large local rings in characteristic 0. The need for the use of Brauer's generalized decomposition numbers is related to how much fusion of conjugacy classes of 2-power order elements occurs in  $G$ .

Apart from the fact that universal deformation rings of representations of finite groups give more insight into the representation theory of these finite groups, there is another important motivation for studying these universal deformation rings. Namely, universal deformation rings for finite groups provide a good test case for various conjectures concerning the ring theoretic properties of universal deformation rings for profinite Galois groups. For example, Flach asked whether there could be universal deformation rings which are not complete intersections (see [17]). In [6, 7] it was shown that the universal deformation ring of the non-trivial irreducible mod 2 representation of the symmetric group  $S_4$  is not a complete intersection; in fact it is not even Cohen-Macaulay. This led to infinitely

---

2000 *Mathematics Subject Classification.* Primary 20C20; Secondary 20C15, 16G10.

*Key words and phrases.* Universal deformation rings; Brauer's generalized decomposition numbers; tame blocks; dihedral defect groups; semidihedral defect groups; generalized quaternion defect groups.

The author was supported in part by NSA Grant H98230-11-1-0131.

many examples of real quadratic fields  $L$  such that the universal deformation ring of the inflation of this representation to the Galois group over  $L$  of the maximal totally unramified extension of  $L$  is not a complete intersection. In [8], examples of finite groups  $G$  and mod  $p$  representations of  $G$  for every odd prime number  $p$  were constructed such that the universal deformation rings of these representations are not complete intersections. The main advantage of computing universal deformation rings for representations of finite groups is that one can use deep results from modular representation theory due to Brauer, Erdmann [22], Linckelmann [27, 28], Carlson-Thévenaz [15, 16], and others.

In this paper, we consider the case when  $k$  has characteristic 2 and  $B$  is an arbitrary tame block of  $kG$  with at least two isomorphism classes of simple modules. Note that tame blocks with exactly one isomorphism class of simple modules seem to require slightly different arguments, as was demonstrated for example in [2]. In [14, 31], it was proved that a tame block contains at most three simple modules up to isomorphism. Hence we consider the case when there are either precisely two or precisely three isomorphism classes of simple  $B$ -modules.

The following theorem summarizes our main results; more precise statements can be found in §6, and in particular in Theorem 6.5. Let  $F$  be the fraction field of  $W$  and let  $\overline{F}$  be a fixed algebraic closure of  $F$ .

**Theorem 1.1.** *Suppose  $k$  has characteristic 2,  $G$  is a finite group, and  $B$  is a tame block of  $kG$  with a defect group  $D$  of order  $2^n \geq 8$  such that there are at least two isomorphism classes of simple  $B$ -modules. Let  $V$  be an indecomposable  $kG$ -module belonging to  $B$  whose stable endomorphism ring is isomorphic to  $k$  and whose Brauer character is the restriction to the 2-regular conjugacy classes of an irreducible  $\overline{F}$ -character of  $G$  of height 1. Then there exists a monic polynomial  $p_n(t) \in W[t]$  of degree  $2^{n-2} - 1$ , depending on  $D$ , whose non-leading coefficients are all divisible by 2 such that either*

- (i)  $R(G, V)/2R(G, V) \cong k[[t]]/(t^{2^{n-2}-1})$ , in which case  $R(G, V) \cong W[[t]]/(p_n(t))$ , or
- (ii)  $R(G, V)/2R(G, V) \cong k[[t]]/(t^{2^{n-2}})$ , in which case  $R(G, V) \cong W[[t]]/(t p_n(t), 2 p_n(t))$ .

*In all cases, the ring  $R(G, V)$  is isomorphic to a subquotient ring of  $WD$ , and it is a complete intersection if and only if we are in case (i).*

In particular, Theorem 1.1 gives a positive answer to [5, Question 1.1] for all  $B$  and  $V$  considered in the theorem.

In [14, 31], Brauer, respectively Olsson, proved that, except for the case when  $D$  is quaternion of order 8, there are precisely  $2^{n-2} - 1$  irreducible  $\overline{F}$ -characters of  $G$  of height 1 belonging to  $B$ . Moreover, these characters fall into  $n - 2$  Galois orbits under the action of  $\text{Gal}(\overline{F}/F)$ . Brauer's generalized decomposition numbers are used to give a formula for the values of these characters on the 2-singular elements of  $G$  in terms of the irreducible Brauer characters belonging to  $B$ . Depending on the fusion of  $D$ -conjugacy classes in  $G$ , this formula is more or less complicated (see Remarks 3.1 and 5.2). This intricate relationship between ordinary and 2-modular characters enables us to construct a lift of  $V$  over  $W[[t]]/(p_n(t))$ . More precisely, we first construct an indecomposable  $B$ -module  $\overline{U}'$  which defines a lift of  $V$  over  $k[t]/(t^{2^{n-2}-1})$  and whose Brauer character is the sum of the restrictions of  $2^{n-2} - 1$  irreducible  $\overline{F}$ -characters of height 1 to the 2-regular conjugacy classes of  $G$ . Then we construct an indecomposable  $WG$ -module  $U'$  which defines a lift of  $\overline{U}'$  over  $W$  and whose  $F$ -character is the sum of  $2^{n-2} - 1$  irreducible  $\overline{F}$ -characters of height 1. We use Brauer's generalized decomposition numbers corresponding to an element  $\sigma \in D$  of order  $2^{n-1}$  to analyze the  $WG$ -module endomorphisms of  $U'$  and to prove that  $U'$  defines a lift of  $V$  over  $W[[t]]/(p_n(t))$ . If the stable Auslander-Reiten quiver of  $B$  contains no 3-tubes then  $W[[t]]/(p_n(t))$  is the universal deformation ring of  $V$ . Otherwise the universal deformation ring of  $V$  is isomorphic to  $W[[t]]/(t p_n(t), 2 p_n(t))$ . Note that if  $D$  is dihedral (resp. generalized quaternion) then the stable Auslander-Reiten quiver of  $B$  has at least one (resp. no) 3-tube. On the other hand, if  $D$  is semidihedral then the stable Auslander-Reiten quiver of  $B$  may or may not contain 3-tubes (see Lemma 6.3).

The use of Brauer's generalized decomposition numbers provides a correction to some arguments in [3, §3.4], [4, §5], and [9, §3.2]. Namely, in these papers a simplistic formula for the values of the irreducible  $\overline{F}$ -characters on elements in  $D$  of maximal 2-power order was assumed, which is true for principal tame blocks but cannot be verified for arbitrary tame blocks because there may be more fusion of  $D$ -conjugacy classes in  $G$  when the blocks are not principal (see Remark 5.2).

The paper is organized as follows. In §2, we recall the definitions of deformations and deformation rings. In §3, we give a brief introduction to Brauer's generalized decomposition numbers, as introduced in [10, 11]. In §4, we describe the quivers and relations of the basic algebras of all tame blocks  $B$  considered in this paper, and provide their decomposition matrices. In §5, we describe results of [14, 31] about the irreducible  $\overline{F}$ -characters of  $G$  belonging to  $B$ . Moreover, we show how Brauer's generalized decomposition numbers are used to analyze the endomorphism ring of a  $WG$ -module  $U'$  which is free over  $W$  and whose  $F$ -character is the sum of  $2^{n-2} - 1$  irreducible  $\overline{F}$ -characters of height 1 belonging to  $B$  (see Lemma 5.6). In §6, we prove Theorem 1.1.

## 2. PRELIMINARIES

In this section, we give a brief introduction to versal and universal deformation rings and deformations. For more background material, we refer the reader to [30] and [19].

Let  $k$  be an algebraically closed field of characteristic  $p > 0$ , and let  $W$  be the ring of infinite Witt vectors over  $k$ . Let  $\hat{\mathcal{C}}$  be the category of all complete local commutative Noetherian rings with residue field  $k$ . The morphisms in  $\hat{\mathcal{C}}$  are continuous  $W$ -algebra homomorphisms which induce the identity map on  $k$ .

Suppose  $G$  is a finite group and  $V$  is a finitely generated  $kG$ -module. A lift of  $V$  over an object  $R$  in  $\hat{\mathcal{C}}$  is a pair  $(M, \phi)$  where  $M$  is a finitely generated  $RG$ -module which is free over  $R$ , and  $\phi : k \otimes_R M \rightarrow V$  is an isomorphism of  $kG$ -modules. Two lifts  $(M, \phi)$  and  $(M', \phi')$  of  $V$  over  $R$  are isomorphic if there is an isomorphism  $f : M \rightarrow M'$  with  $\phi = \phi' \circ (k \otimes f)$ . The isomorphism class  $[M, \phi]$  of a lift  $(M, \phi)$  of  $V$  over  $R$  is called a deformation of  $V$  over  $R$ , and the set of all such deformations is denoted by  $\text{Def}_G(V, R)$ . The deformation functor

$$\hat{F}_V : \hat{\mathcal{C}} \rightarrow \text{Sets}$$

is a covariant functor which sends an object  $R$  in  $\hat{\mathcal{C}}$  to  $\text{Def}_G(V, R)$  and a morphism  $\alpha : R \rightarrow R'$  in  $\hat{\mathcal{C}}$  to the map  $\text{Def}_G(V, R) \rightarrow \text{Def}_G(V, R')$  defined by  $[M, \phi] \mapsto [R' \otimes_{R, \alpha} M, \phi_\alpha]$ , where  $\phi_\alpha = \phi$  after identifying  $k \otimes_{R'} (R' \otimes_{R, \alpha} M)$  with  $k \otimes_R M$ .

Suppose there exists an object  $R(G, V)$  in  $\hat{\mathcal{C}}$  and a deformation  $[U(G, V), \phi_U]$  of  $V$  over  $R(G, V)$  with the following property: For each  $R$  in  $\hat{\mathcal{C}}$  and for each lift  $(M, \phi)$  of  $V$  over  $R$  there exists a morphism  $\alpha : R(G, V) \rightarrow R$  in  $\hat{\mathcal{C}}$  such that  $\hat{F}_V(\alpha)([U(G, V), \phi_U]) = [M, \phi]$ , and moreover  $\alpha$  is unique if  $R$  is the ring of dual numbers  $k[\epsilon]/(\epsilon^2)$ . Then  $R(G, V)$  is called the versal deformation ring of  $V$  and  $[U(G, V), \phi_U]$  is called the versal deformation of  $V$ . If the morphism  $\alpha$  is unique for all  $R$  and all lifts  $(M, \phi)$  of  $V$  over  $R$ , then  $R(G, V)$  is called the universal deformation ring of  $V$  and  $[U(G, V), \phi_U]$  is called the universal deformation of  $V$ . In other words,  $R(G, V)$  is universal if and only if  $R(G, V)$  represents the functor  $\hat{F}_V$  in the sense that  $\hat{F}_V$  is naturally isomorphic to the Hom functor  $\text{Hom}_{\hat{\mathcal{C}}}(R(G, V), -)$ .

Note that the above definition of deformations can be weakened as follows. Given a lift  $(M, \phi)$  of  $V$  over a ring  $R$  in  $\hat{\mathcal{C}}$ , define the corresponding weak deformation to be the isomorphism class of  $M$  as an  $RG$ -module, without taking into account the specific isomorphism  $\phi : k \otimes_R M \rightarrow V$ . In general, a weak deformation of  $V$  over  $R$  identifies more lifts than a deformation of  $V$  over  $R$  that respects the isomorphism  $\phi$  of a representative  $(M, \phi)$ . However, if the stable endomorphism ring  $\underline{\text{End}}_{kG}(V)$  is isomorphic to  $k$ , these two definitions of deformations coincide (see [4, Remark 2.1]).

By [30], every finitely generated  $kG$ -module  $V$  has a versal deformation ring. Since  $G$  is a finite group, we have the following sufficient criterion for the universality of  $R(G, V)$ :

**Proposition 2.1.** ([5, Prop. 2.1]) *Suppose  $V$  is a finitely generated  $kG$ -module whose stable endomorphism ring  $\underline{\text{End}}_{kG}(V)$  is isomorphic to  $k$ . Then  $V$  has a universal deformation ring  $R(G, V)$ .*

## 3. BRAUER'S GENERALIZED DECOMPOSITION NUMBERS

In this section, we give a brief introduction to Brauer's generalized decomposition numbers, emphasizing the results needed in this paper. Throughout this section, let  $p$  be a fixed prime number and let  $G$  be a finite group such that  $p$  divides  $\#G$ . Let  $P$  be a fixed Sylow  $p$ -subgroup of  $G$ .

Brauer introduced generalized decomposition numbers in [10] to be able to express the values of the ordinary irreducible characters of  $G$  on all conjugacy classes by means of the irreducible  $p$ -modular characters of certain subgroups of  $G$ . More precisely, let  $g \in G$  and write  $g$  (uniquely) as  $g = uv$  where  $u$  is a  $p$ -element of order  $p^\alpha$  and  $v$  is a  $p$ -regular element of  $G$  (where we allow  $u$  or  $v$  to be identity elements). Let  $\varphi_1^{(u)}, \dots, \varphi_{\ell(u)}^{(u)}$  be the irreducible  $p$ -modular characters of  $C_G(u)$ . By [10, Sect. 1], if  $\chi$  is an ordinary irreducible character of  $G$  then we have a formula

$$(3.1) \quad \chi(uv) = \sum_{t=1}^{\ell(u)} \left( d_{u, \varphi_t^{(u)}} \right)_\chi \varphi_t^{(u)}(v)$$

where  $\left( d_{u, \varphi_1^{(u)}} \right)_\chi, \dots, \left( d_{u, \varphi_{\ell(u)}^{(u)}} \right)_\chi$  are algebraic integers in the field of  $p^\alpha$ -th roots of unity which do not depend on  $v$ . These are called the generalized decomposition numbers of  $G$  corresponding to  $u$ .

We can use these generalized decomposition numbers to express the ordinary character table of  $G$  as the product of two square matrices as follows. Recall that we fixed a Sylow  $p$ -subgroup  $P$  of  $G$ . Let  $u_1, \dots, u_r$  be a complete system of representatives of  $G$ -conjugacy classes of  $p$ -elements in  $G$  with  $u_i \in P$  for all  $1 \leq i \leq r$  and  $u_1 = 1_G$ . For each  $1 \leq i \leq r$ , let  $v_{i,1}, \dots, v_{i,\ell_i}$  be a complete system of representatives of  $C_G(u_i)$ -conjugacy classes of  $p$ -regular elements in  $C_G(u_i)$  with  $v_{i,1} = 1_{C_G(u_i)}$ . Then  $\{u_i v_{i,j} \mid 1 \leq i \leq r, 1 \leq j \leq \ell_i\}$  is a complete set of representatives of the conjugacy classes of  $G$ . Moreover, for each  $1 \leq i \leq r$ , there are precisely  $\ell_i$  irreducible  $p$ -modular characters of  $C_G(u_i)$ , which we denote by  $\varphi_{i,1}, \dots, \varphi_{i,\ell_i}$ . Let  $\{\chi_1, \dots, \chi_h\}$  be a complete set of representatives of the ordinary irreducible characters of  $G$ . Then Brauer's above formula (3.1) can be written as

$$(3.2) \quad \chi_s(u_i v_{i,j}) = \sum_{t=1}^{\ell_i} (d_{i,t})_{\chi_s} \varphi_{i,t}(v_{i,j})$$

for all  $1 \leq s \leq h$ ,  $1 \leq i \leq r$ ,  $1 \leq j \leq \ell_i$ , where  $\varphi_{i,t} = \varphi_t^{(u_i)}$  and  $d_{i,t} = d_{u_i, \varphi_{i,t}}$  for all  $1 \leq t \leq \ell_i$ . Write the conjugacy class representatives in the order

$$u_1 v_{1,1}, \dots, u_1 v_{1,\ell_1}, u_2 v_{2,1}, \dots, u_2 v_{2,\ell_2}, \dots, u_r v_{r,1}, \dots, u_r v_{r,\ell_r}.$$

Using (3.2), we obtain that the ordinary character table  $\mathcal{X}$  can be written as a product  $\mathcal{X} = D \cdot \Phi$ , where  $D$  contains the generalized decomposition numbers and  $\Phi$  is a block diagonal matrix

$$(3.3) \quad \Phi = \begin{pmatrix} \Phi_1 & & 0 \\ & \ddots & \\ 0 & & \Phi_r \end{pmatrix}$$

with

$$\Phi_i = \begin{pmatrix} \varphi_{i,1}(v_{i,1}) & \cdots & \varphi_{i,1}(v_{i,\ell_i}) \\ \vdots & & \vdots \\ \varphi_{i,\ell_i}(v_{i,1}) & \cdots & \varphi_{i,\ell_i}(v_{i,\ell_i}) \end{pmatrix}$$

for all  $1 \leq i \leq r$ . By [11, Sect. 7, p. 45], the square of the determinant of  $D$  is  $\pm p^a$  for some  $a \in \mathbb{Z}^+$ , and the square of the determinant of  $\Phi$  is an integer which is relatively prime to  $p$ .

Fix now a  $p$ -modular block  $B$  of  $G$  and suppose that there are  $k(B)$  ordinary irreducible characters belonging to  $B$ . Reorder  $\chi_1, \dots, \chi_h$  such that  $\chi_1, \dots, \chi_{k(B)}$  belong to  $B$ . For each  $1 \leq i \leq r$ , reorder  $\varphi_{i,1}, \dots, \varphi_{i,\ell_i}$  such that each of the first  $m_i$  characters,  $\varphi_{i,1}, \dots, \varphi_{i,m_i}$ , belongs to a block of  $C_G(u_i)$

whose Brauer correspondent in  $G$  is equal to  $B$ . (Note that for each block  $b$  of  $C_G(u_i)$ , its Brauer correspondent  $b^G$  in  $G$  is well-defined since the centralizer in  $G$  of a defect group of  $b$  is contained in  $C_G(u_i)$ , see [1, Sect. 14].) It follows from [11, Sect. 6] that for  $1 \leq s \leq k(B)$ , (3.2) can be rewritten as

$$(3.4) \quad \chi_s(u_i v_{i,j}) = \sum_{t=1}^{m_i} (d_{i,t})_{\chi_s} \varphi_{i,t}(v_{i,j})$$

for all  $1 \leq i \leq r$ ,  $1 \leq j \leq \ell_i$ . In particular, we obtain for all  $1 \leq i \leq r$  that

$$(3.5) \quad \mathcal{X}_{B,i} = \begin{pmatrix} \chi_1(u_i v_{i,1}) & \cdots & \chi_1(u_i v_{i,\ell_i}) \\ \vdots & & \vdots \\ \chi_{k(B)}(u_i v_{i,1}) & \cdots & \chi_{k(B)}(u_i v_{i,\ell_i}) \end{pmatrix} = D_{B,i} \cdot \Phi_i$$

where

$$D_{B,i} = \begin{pmatrix} (d_{i,1})_{\chi_1} & \cdots & (d_{i,m_i})_{\chi_1} & 0 & \cdots & 0 \\ \vdots & & \vdots & \vdots & & \vdots \\ (d_{i,1})_{\chi_{k(B)}} & \cdots & (d_{i,m_i})_{\chi_{k(B)}} & 0 & \cdots & 0 \end{pmatrix}$$

and we assume, as above, that  $\varphi_{i,1}, \dots, \varphi_{i,m_i}$  belong to blocks of  $C_G(u_i)$  whose Brauer correspondents in  $G$  are each equal to  $B$ .

Let now  $k$  be an algebraically closed field of characteristic  $p$  and view  $B$  as a block of  $kG$ . Let  $W = W(k)$  be the ring of infinite Witt vectors over  $k$  and let  $F$  be the fraction field of  $W$ . In particular,  $W$  contains all roots of unity of order not divisible by  $p$ . Let  $\overline{F}$  be a fixed algebraic closure of  $F$ , and let  $\xi$  be a primitive  $|P|$ -th root of unity, where  $P$  is our fixed Sylow  $p$ -subgroup of  $G$ . Then  $F(\xi)$  is a splitting field for  $G$  with ring of integers  $W[\xi]$ . We can thus view the ordinary character table  $\mathcal{X}$  of  $G$  and the matrix  $D$  of generalized decomposition numbers as taking values in  $W[\xi]$ , and the matrix  $\Phi$  in (3.3) as taking values in  $W$ . Since the square of the determinant of  $\Phi$  is an integer that is relatively prime to  $p$ , it follows that the determinant of  $\Phi$ , and hence the determinant of  $\Phi_i$  for all  $1 \leq i \leq r$ , is a unit in  $W$ . Hence we can solve (3.5) for  $D_{B,i}$  to obtain

$$(3.6) \quad D_{B,i} = \mathcal{X}_{B,i} \cdot \Psi_i$$

for all  $1 \leq i \leq r$ , where  $\Psi_i = \Phi_i^{-1}$  is a matrix with values in  $W$ .

*Remark 3.1.* Equations (3.1) and (3.4) can be rewritten to reflect the influence of fusion of  $P$ -conjugacy classes in  $G$  (see [13, §1]). As before, let  $u$  be a  $p$ -element of  $G$  and let  $v$  be a  $p$ -regular element in  $C_G(u)$ . Assuming the notation of the previous paragraph, let  $\chi$  be an irreducible  $F(\xi)$ -character which belongs to a block  $B$  of  $kG$ . Recall that a subsection  $(y, b_y)$  for  $B$  is a pair consisting of a  $p$ -element  $y$  of  $G$  and a block  $b_y$  of  $C_G(y)$  with  $b_y^G = B$ . We obtain

$$(3.7) \quad \chi(uv) = \sum_{(y, b_y)} \sum_{\varphi} (d_{y, \varphi})_{\chi} \varphi(z_y v z_y^{-1}).$$

Here  $(y, b_y)$  ranges over a system of representatives for the conjugacy classes of subsections for  $B$  such that  $y$  is conjugate to  $u$  in  $G$ , say  $u = z_y^{-1} y z_y$ . For each  $(y, b_y)$ ,  $\varphi$  ranges over the irreducible Brauer characters associated with  $b_y$ .

#### 4. TAME BLOCKS

For the remainder of this paper, we make the following assumptions.

**Hypothesis 4.1.** Let  $k$  be an algebraically closed field of characteristic 2, and let  $n \geq 3$  be an integer. Let  $W = W(k)$  be the ring of infinite Witt vectors over  $k$ , let  $F$  be the fraction field of  $W$ , and let  $\overline{F}$  be a fixed algebraic closure of  $F$ . Let  $\zeta$  be a primitive  $2^{n-1}$ -th root of unity in  $\overline{F}$ . Suppose  $G$  is a finite group and  $B$  is a block of  $kG$  of tame representation type with a defect group  $D$  of order  $2^n$ . Assume that  $n \geq 3$  if  $D$  is dihedral or generalized quaternion, and that  $n \geq 4$  if  $D$  is semidihedral. Suppose further that  $B$  contains at least two isomorphism classes of simple modules.

Note that Brauer [14] and Olsson [31] proved that a tame block  $B$  as in Hypothesis 4.1 contains at most three simple modules up to isomorphism. Hence  $B$  contains either exactly two or exactly three isomorphism classes of simple modules. By [14, Prop. (5A)] and [31, Prop. 4.1], all ordinary irreducible characters of  $G$  belonging to  $B$  take values in  $F(\zeta)$ . In fact, all ordinary irreducible characters of  $G$  belonging to  $B$  can be realized by simple  $F(\zeta)G$ -modules (see §5).

Assume Hypothesis 4.1. From Erdmann's classification of all blocks of tame representation type in [22], it follows that the quiver and relations of the basic algebra of  $B$  can be given explicitly and that, up to Morita equivalence, there are 24 families of blocks  $B$ .

Using [20, 21, 22, 24], we now give a description of these families as follows. By [22, pp. 294–306], there are 12 possible quivers  $Q$  which can occur for basic algebras of dihedral, semidihedral or quaternion type:  $2\mathcal{A}$ ,  $2\mathcal{B}$ ,  $3\mathcal{A}$ ,  $3\mathcal{B}$ ,  $3\mathcal{C}$ ,  $3\mathcal{D}$ ,  $3\mathcal{F}$ ,  $3\mathcal{H}$ ,  $3\mathcal{K}$ ,  $3\mathcal{L}$ ,  $3\mathcal{Q}$ ,  $3\mathcal{R}$ . For each such quiver  $Q$ , we combine Erdmann's results in [21, 22] and [24, Prop. 4.2] with Eisele's results in [20] to provide the most accurate description of a full set of representatives of basic algebras  $\Lambda = kQ/I$  for the Morita equivalence classes of blocks  $B$  as in Hypothesis 4.1 with Ext quiver  $Q$ .

We also provide the decomposition matrix for each block  $B$ . For better readability, all decomposition matrices appear at the end of the paper. As will be discussed in §5,  $B$  always contains exactly 4 ordinary irreducible characters of height 0 and, unless  $D$  is quaternion of order 8, exactly  $2^{n-2} - 1$  ordinary irreducible characters of height 1. If  $D$  is quaternion of order 8,  $B$  contains exactly 3 ordinary irreducible characters of height 1. If  $B$  is generalized quaternion or semidihedral, there may be additional ordinary irreducible characters of height  $n - 2$ . In the decomposition matrices, we list first the 4 characters of height 0, then the family of  $2^{n-2} - 1$  characters of height 1 which all reduce to the same Brauer character on restricting to the 2-regular conjugacy classes of  $G$ , and then the ordinary irreducible characters of height  $n - 2$  if they exist. Moreover, if  $B$  has three isomorphism classes of simple modules, we order the ordinary irreducible characters according to the sign conventions described in [14, Thm. 5] and [31, Thms. 4.10, 4.11 and 4.15].

To distinguish between different defect groups, we use the notation  $D(Q)$  (resp.  $SD(Q)$ , resp.  $Q(Q)$ ) to mean that  $\Lambda = kQ/I$  is Morita equivalent to a block  $B$  with dihedral (resp. semidihedral, resp. generalized quaternion) defect groups.

#### 4.1. Blocks with quiver $2\mathcal{A}$ .

$$2\mathcal{A} = \alpha \begin{array}{c} \circlearrowleft \\ \bullet \xrightarrow{\beta} \bullet \\ \bullet \xleftarrow{\gamma} \bullet \end{array}$$

By [20] and [22, p. 294], if  $B$  has dihedral defect groups and Ext quiver  $2\mathcal{A}$ , then  $B$  is Morita equivalent to  $D(2\mathcal{A}) = k[2\mathcal{A}]/I_{D(2\mathcal{A})}$  where

$$I_{D(2\mathcal{A})} = \langle \beta\gamma, \alpha^2, (\gamma\beta\alpha)^{2^{n-2}} - (\alpha\gamma\beta)^{2^{n-2}} \rangle.$$

The corresponding decomposition matrix is given in Figure 1.

By [22, p. 298], if  $B$  has semidihedral defect groups and Ext quiver  $2\mathcal{A}$ , then there exists  $c \in k$  such that  $B$  is Morita equivalent to either  $SD(2\mathcal{A})_1(c) = k[2\mathcal{A}]/I_{SD(2\mathcal{A})_1,c}$  or  $SD(2\mathcal{A})_2(c) = k[2\mathcal{A}]/I_{SD(2\mathcal{A})_2,c}$  where

$$I_{SD(2\mathcal{A})_1,c} = \langle \alpha^2 - c(\gamma\beta\alpha)^{2^{n-2}}, \beta\gamma\beta - \beta\alpha(\gamma\beta\alpha)^{2^{n-2}-1}, \gamma\beta\gamma - \alpha\gamma(\beta\alpha\gamma)^{2^{n-2}-1}, \alpha(\gamma\beta\alpha)^{2^{n-2}} \rangle,$$

$$I_{SD(2\mathcal{A})_2,c} = \langle \beta\gamma, \alpha^2 - \gamma\beta(\alpha\gamma\beta)^{2^{n-2}-1} - c(\gamma\beta\alpha)^{2^{n-2}}, (\gamma\beta\alpha)^{2^{n-2}} - (\alpha\gamma\beta)^{2^{n-2}} \rangle.$$

The decomposition matrix for  $SD(2\mathcal{A})_1(c)$  is given in Figure 2, and the decomposition matrix for  $SD(2\mathcal{A})_2(c)$  is given in Figure 1.

By [22, p. 303], if  $B$  has generalized quaternion defect groups and Ext quiver  $2\mathcal{A}$ , then there exists  $c \in k$  such that  $B$  is Morita equivalent to  $Q(2\mathcal{A})(c) = k[2\mathcal{A}]/I_{Q(2\mathcal{A}),c}$  where

$$I_{Q(2\mathcal{A}),c} = \langle \alpha^2 - \gamma\beta(\alpha\gamma\beta)^{2^{n-2}-1} - c(\alpha\gamma\beta)^{2^{n-2}}, \beta\gamma\beta - \beta\alpha(\gamma\beta\alpha)^{2^{n-2}-1}, \gamma\beta\gamma - \alpha\gamma(\beta\alpha\gamma)^{2^{n-2}-1}, \beta\alpha^2 \rangle.$$

The decomposition matrix for  $Q(2\mathcal{A})(c)$  is given in Figure 2.

#### 4.2. Blocks with quiver $2\mathcal{B}$ .

$$2\mathcal{B} = \alpha \begin{array}{c} \circlearrowleft \\ \bullet \\ \circlearrowright \end{array} \begin{array}{c} 0 \\ \xrightarrow{\beta} \\ 1 \\ \xleftarrow{\gamma} \end{array} \begin{array}{c} \bullet \\ \circlearrowright \\ \bullet \end{array} \eta$$

By [20] and [22, p. 295], if  $B$  has dihedral defect groups and Ext quiver  $2\mathcal{B}$ , then  $B$  is Morita equivalent to  $D(2\mathcal{B}) = k[2\mathcal{B}]/I_{D(2\mathcal{B})}$  where

$$I_{D(2\mathcal{B})} = \langle \eta\beta, \gamma\eta, \beta\gamma, \alpha^2, \gamma\beta\alpha - \alpha\gamma\beta, \eta^{2^{n-2}} - \beta\alpha\gamma \rangle.$$

The corresponding decomposition matrix is given in Figure 3.

By [21, Lemmas (8.11) and (8.15)], [22, p. 299] and [24, Prop. 4.2], if  $B$  has semidihedral defect groups and Ext quiver  $2\mathcal{B}$ , then there exists  $c \in k$  such that  $B$  is Morita equivalent to either  $SD(2\mathcal{B})_1(c) = k[2\mathcal{B}]/I_{SD(2\mathcal{B})_1,c}$  or  $SD(2\mathcal{B})_2(c) = k[2\mathcal{B}]/I_{SD(2\mathcal{B})_2,c}$  or  $SD(2\mathcal{B})_4(c) = k[2\mathcal{B}]/I_{SD(2\mathcal{B})_4,c}$  where

$$\begin{aligned} I_{SD(2\mathcal{B})_1,c} &= \langle \eta\beta, \gamma\eta, \beta\gamma, \alpha^2 - \gamma\beta - c\alpha\gamma\beta, \gamma\beta\alpha - \alpha\gamma\beta, \eta^{2^{n-2}} - \beta\alpha\gamma \rangle, \\ I_{SD(2\mathcal{B})_2,c} &= \langle \eta\beta - \beta\alpha(\gamma\beta\alpha), \gamma\eta - \alpha\gamma(\beta\alpha\gamma), \alpha^2 - c(\gamma\beta\alpha)^2, \beta\gamma - \eta^{2^{n-2}-1}, \\ &\quad \eta^2\beta, \gamma\eta^2 \rangle, \\ I_{SD(2\mathcal{B})_4,c} &= \langle \gamma\eta - \alpha\gamma, \beta\alpha - \eta\beta, \alpha^{2^{n-2}+1}, \eta^{2^{n-2}+1}, \beta\alpha^{2^{n-2}-1}, \alpha^{2^{n-2}-1}\gamma, \\ &\quad \gamma\eta^{2^{n-2}-1}, \eta^{2^{n-2}-1}\beta, \gamma\beta - \alpha^2, \beta\gamma - \eta^2(1 + c\eta^{2^{n-2}-2}) \rangle. \end{aligned}$$

The decomposition matrix for  $SD(2\mathcal{B})_1(c)$  is given in Figure 3, the decomposition matrix for  $SD(2\mathcal{B})_2(c)$  is given in Figure 4, and the decomposition matrix for  $SD(2\mathcal{B})_4(c)$  is given in Figure 5.

By [22, IX.4.1 and pp. 303–304], if  $B$  has generalized quaternion defect groups and Ext quiver  $2\mathcal{B}$ , then there exists  $c \in k$ , respectively  $p(t) \in k[t]$  with  $p(0) = 1$  and  $a, c \in k$  with  $a \neq 0$ , such that  $B$  is Morita equivalent to either  $Q(2\mathcal{B})_1(c) = k[2\mathcal{B}]/I_{Q(2\mathcal{B})_1,c}$  or  $Q(2\mathcal{B})_2(p, a, c) = k[2\mathcal{B}]/I_{Q(2\mathcal{B})_2,p,a,c}$  where

$$\begin{aligned} I_{Q(2\mathcal{B})_1,c} &= \langle \eta\beta - \beta\alpha(\gamma\beta\alpha), \gamma\eta - \alpha\gamma(\beta\alpha\gamma), \alpha^2 - \gamma\beta(\alpha\gamma\beta) - c(\alpha\gamma\beta)^2, \\ &\quad \beta\gamma - \eta^{2^{n-2}-1}, \beta\alpha^2 \rangle, \\ I_{Q(2\mathcal{B})_2,p,a,c} &= \langle \gamma\eta - \alpha\gamma, \beta\alpha - \eta\beta, \alpha^{2^{n-2}+1}, \eta^{2^{n-2}+1}, \beta\alpha^{2^{n-2}-1}, \alpha^{2^{n-2}-1}\gamma, \\ &\quad \gamma\beta - p(\alpha)\alpha^2, \beta\gamma - p(\eta)\eta^2 - a\eta^{2^{n-2}-1} - c\eta^{2^{n-2}} \rangle. \end{aligned}$$

The decomposition matrix for  $Q(2\mathcal{B})_1(c)$  is given in Figure 4, and the decomposition matrix for  $Q(2\mathcal{B})_2(p, a, c)$  is given in Figure 5. Note that by [31, Lemma 3.3],  $Q(2\mathcal{B})_2(p, a, c)$  can actually not occur as a block.

#### 4.3. Blocks with quiver $3\mathcal{A}$ .

$$3\mathcal{A} = 1 \bullet \begin{array}{c} \xrightarrow{\beta} \\ \bullet \\ \xleftarrow{\gamma} \end{array} \begin{array}{c} 0 \\ \xrightarrow{\delta} \\ \bullet \\ \xleftarrow{\eta} \end{array} \bullet 2$$

By [22, IX.5.4 and p. 295], if  $B$  has dihedral defect groups and Ext quiver  $3\mathcal{A}$ , then  $B$  is Morita equivalent to  $D(3\mathcal{A})_1 = k[3\mathcal{A}]/I_{D(3\mathcal{A})_1}$  where

$$I_{D(3\mathcal{A})_1} = \langle \gamma\beta, \delta\eta, (\eta\delta\beta\gamma)^{2^{n-2}} - (\beta\gamma\eta\delta)^{2^{n-2}} \rangle.$$

The corresponding decomposition matrix is given in Figure 6.

By [22, IX.5.3 and pp. 299–300], if  $B$  has semidihedral defect groups and Ext quiver  $3\mathcal{A}$ , then  $B$  is Morita equivalent to  $SD(3\mathcal{A})_1 = k[3\mathcal{A}]/I_{SD(3\mathcal{A})_1}$  where

$$I_{SD(3\mathcal{A})_1} = \langle \gamma\beta, \delta\eta\delta - \delta\beta\gamma(\eta\delta\beta\gamma)^{2^{n-2}-1}, \eta\delta\eta - \beta\gamma\eta(\delta\beta\gamma\eta)^{2^{n-2}-1} \rangle.$$

The decomposition matrix for  $SD(3\mathcal{A})_1$  is given in Figure 7.

By [22, IX.5.2 and pp. 304–305], if  $B$  has generalized quaternion defect groups and Ext quiver  $3\mathcal{A}$ , then  $B$  is Morita equivalent to  $Q(3\mathcal{A})_2 = k[3\mathcal{A}]/I_{Q(3\mathcal{A})_2}$  where

$$I_{Q(3\mathcal{A})_2} = \langle \beta\gamma\beta - \eta\delta\beta(\gamma\eta\delta\beta)^{2^{n-2}-1}, \gamma\beta\gamma - \gamma\eta\delta(\beta\gamma\eta\delta)^{2^{n-2}-1}, \delta\beta\gamma\beta, \\ \delta\eta\delta - \delta\beta\gamma(\eta\delta\beta\gamma)^{2^{n-2}-1}, \eta\delta\eta - \beta\gamma\eta(\delta\beta\gamma\eta)^{2^{n-2}-1}, \gamma\eta\delta\eta \rangle.$$

The decomposition matrix for  $Q(3\mathcal{A})_2$  is given in Figure 8.

#### 4.4. Blocks with quiver $3\mathcal{B}$ .

$$3\mathcal{B} = \alpha \begin{array}{c} \curvearrowright \\ \bullet \\ \leftarrow \gamma \end{array} \begin{array}{c} 1 \\ \xrightarrow{\beta} \\ \bullet \\ \xleftarrow{\eta} \end{array} \begin{array}{c} 0 \\ \xrightarrow{\delta} \\ \bullet \\ \xleftarrow{\eta} \end{array} \bullet 2$$

By [22, IX.5.4 and pp. 295–296], if  $B$  has dihedral defect groups and Ext quiver  $3\mathcal{B}$ , then  $B$  is Morita equivalent to  $D(3\mathcal{B})_1 = k[3\mathcal{B}]/I_{D(3\mathcal{B})_1}$  where

$$I_{D(3\mathcal{B})_1} = \langle \beta\alpha, \alpha\gamma, \gamma\beta, \delta\eta, \eta\delta\beta\gamma - \beta\gamma\eta\delta, \alpha^{2^{n-2}} - \gamma\eta\delta\beta \rangle.$$

The corresponding decomposition matrix is given in Figure 9.

By [22, p. 300], if  $B$  has semidihedral defect groups and Ext quiver  $3\mathcal{B}$ , then  $B$  is Morita equivalent to either  $SD(3\mathcal{B})_1 = k[3\mathcal{B}]/I_{SD(3\mathcal{B})_1}$  or  $SD(3\mathcal{B})_2 = k[3\mathcal{B}]/I_{SD(3\mathcal{B})_2}$  where

$$I_{SD(3\mathcal{B})_1} = \langle \beta\alpha, \alpha\gamma, \gamma\beta, \delta\eta\delta - \delta\beta\gamma, \eta\delta\eta - \beta\gamma\eta, \alpha^{2^{n-2}} - \gamma\eta\delta\beta \rangle, \\ I_{SD(3\mathcal{B})_2} = \langle \delta\eta, \gamma\beta - \alpha^{2^{n-2}-1}, \alpha\gamma - \gamma\eta\delta(\beta\gamma\eta\delta), \beta\alpha - \eta\delta\beta(\gamma\eta\delta\beta) \rangle.$$

The decomposition matrix for  $SD(3\mathcal{B})_1$  is given in Figure 10, and the decomposition matrix for  $SD(3\mathcal{B})_2$  is given in Figure 11.

By [22, p. 305], if  $B$  has generalized quaternion defect groups and Ext quiver  $3\mathcal{B}$ , then  $B$  is Morita equivalent to  $Q(3\mathcal{B}) = k[3\mathcal{B}]/I_{Q(3\mathcal{B})}$  where

$$I_{Q(3\mathcal{B})} = \langle \gamma\beta - \alpha^{2^{n-2}-1}, \alpha\gamma - \gamma\eta\delta(\beta\gamma\eta\delta), \beta\alpha - \eta\delta\beta(\gamma\eta\delta\beta), \delta\eta\delta - \delta\beta\gamma(\eta\delta\beta\gamma), \\ \eta\delta\eta - \beta\gamma\eta(\delta\beta\gamma\eta), \beta\alpha^2, \delta\eta\delta\beta \rangle.$$

The decomposition matrix for  $Q(3\mathcal{B})$  is given in Figure 12.

#### 4.5. Blocks with quiver $3\mathcal{C}$ .

$$3\mathcal{C} = 1 \bullet \begin{array}{c} \xrightarrow{\beta} \\ \bullet \\ \xleftarrow{\gamma} \end{array} \begin{array}{c} 0 \\ \xrightarrow{\delta} \\ \bullet \\ \xleftarrow{\eta} \end{array} \bullet 2$$

$\rho$

By [22, VI.5] (resp. by [22, IX.5.3 and p. 305]), there are no blocks  $B$  with dihedral (resp. generalized quaternion) defect groups that have Ext quiver  $3\mathcal{C}$ .

By [22, IX.5.3 and pp. 300–301], if  $B$  has semidihedral defect groups and Ext quiver  $3\mathcal{C}$ , then  $B$  is Morita equivalent to either  $SD(3\mathcal{C})_{2,1} = k[3\mathcal{C}]/I_{SD(3\mathcal{C})_{2,1}}$  or  $SD(3\mathcal{C})_{2,2} = k[3\mathcal{C}]/I_{SD(3\mathcal{C})_{2,2}}$  where

$$I_{SD(3\mathcal{C})_{2,1}} = \langle \rho\beta, \delta\rho, \rho\eta, \gamma\rho, \beta\gamma - \eta\delta, (\beta\gamma)^2 - \rho^{2^{n-2}}, \delta\beta\gamma\beta, \gamma\eta\delta\eta \rangle, \\ I_{SD(3\mathcal{C})_{2,2}} = \langle \rho\beta, \delta\rho, \rho\eta, \gamma\rho, \beta\gamma - \eta\delta, (\beta\gamma)^{2^{n-2}} - \rho^2, \delta\beta(\gamma\beta)^{2^{n-2}-1}, \\ \gamma\eta(\delta\eta)^{2^{n-2}-1} \rangle.$$

The decomposition matrix for  $SD(3\mathcal{C})_{2,1}$  is given in Figure 13, and the decomposition matrix for  $SD(3\mathcal{C})_{2,2}$  is given in Figure 14.



#### 4.6. Blocks with quiver $3\mathcal{D}$ .

$$3\mathcal{D} = \begin{array}{c} \alpha \curvearrowright \bullet \xrightarrow{\beta} \bullet \xrightarrow{\delta} \bullet \curvearrowright \xi \\ \xleftarrow{\gamma} \bullet \xleftarrow{\eta} \bullet \end{array}$$

By [22, IX.5.1, IX.5.4 and p. 296] (resp. by [22, IX.5.1 and p. 306]), there are no blocks  $B$  with dihedral (resp. generalized quaternion) defect groups that have Ext quiver  $3\mathcal{D}$ .

By [22, p. 301], if  $B$  has semidihedral defect groups and Ext quiver  $3\mathcal{D}$ , then  $B$  is Morita equivalent to  $\text{SD}(3\mathcal{D}) = k[3\mathcal{D}]/I_{\text{SD}(3\mathcal{D})}$  where

$$I_{\text{SD}(3\mathcal{D})} = \langle \xi\delta, \eta\xi, \delta\eta, \gamma\beta - \alpha^{2^{n-2}-1}, \alpha\gamma - \gamma\eta\delta, \beta\alpha - \eta\delta\beta, \xi^2 - \delta\beta\gamma\eta \rangle.$$

The decomposition matrix for  $\text{SD}(3\mathcal{D})$  is given in Figure 10.

**4.7. Blocks with quiver  $3\mathcal{F}$ .** By [22, VI.5] (resp. by [22, VII.4]), there are no blocks  $B$  with dihedral (resp. generalized quaternion) defect groups that have Ext quiver  $3\mathcal{F}$ . By [22, IX.5.2 and p. 301], there are also no blocks  $B$  with semidihedral defect groups that have Ext quiver  $3\mathcal{F}$ .

#### 4.8. Blocks with quiver $3\mathcal{H}$ .

$$3\mathcal{H} = \begin{array}{ccc} & \xrightarrow{\beta} & \\ 0 \bullet & & \bullet 1 \\ & \xleftarrow{\gamma} & \\ & & \nearrow \eta \\ \lambda & & \searrow \delta \\ & & 2 \bullet \end{array}$$

By [22, VI.5] (resp. by [22, VII.4]), there are no blocks  $B$  with dihedral (resp. generalized quaternion) defect groups that have Ext quiver  $3\mathcal{H}$ .

By [22, p. 301], if  $B$  has semidihedral defect groups and Ext quiver  $3\mathcal{H}$ , then  $B$  is Morita equivalent to  $\text{SD}(3\mathcal{H})_1 = k[3\mathcal{H}]/I_{\text{SD}(3\mathcal{H})_1}$  or  $\text{SD}(3\mathcal{H})_2 = k[3\mathcal{H}]/I_{\text{SD}(3\mathcal{H})_2}$  where

$$I_{\text{SD}(3\mathcal{H})_1} = \langle \lambda\delta - \gamma\beta\gamma, \beta\lambda - \eta(\delta\eta)^{2^{n-2}-1}, \eta\delta\beta, \delta\beta\gamma, \gamma\eta \rangle,$$

$$I_{\text{SD}(3\mathcal{H})_2} = \langle \lambda\delta - \gamma(\beta\gamma)^{2^{n-2}-1}, \beta\lambda - \eta\delta\eta, \eta\delta\beta, \delta\beta\gamma, \gamma\eta \rangle.$$

The decomposition matrix for  $\text{SD}(3\mathcal{H})_1$  is given in Figure 15, and the decomposition matrix for  $\text{SD}(3\mathcal{H})_2$  is given in Figure 16.

#### 4.9. Blocks with quiver $3\mathcal{K}$ .

$$3\mathcal{K} = \begin{array}{ccc} & \xrightarrow{\beta} & \\ 0 \bullet & & \bullet 1 \\ & \xleftarrow{\gamma} & \\ & \searrow \kappa & \nearrow \eta \\ \lambda & & \searrow \delta \\ & & 2 \bullet \end{array}$$

By [22, IX.5.2 and p. 302], there are no blocks  $B$  with semidihedral defect groups that have Ext quiver  $3\mathcal{K}$ .

By [22, p. 296], if  $B$  has dihedral defect groups and Ext quiver  $3\mathcal{K}$ , then  $B$  is Morita equivalent to  $\text{D}(3\mathcal{K}) = k[3\mathcal{K}]/I_{\text{D}(3\mathcal{K})}$  where

$$I_{\text{D}(3\mathcal{K})} = \langle \delta\beta, \lambda\delta, \beta\lambda, \kappa\gamma, \eta\kappa, \gamma\eta, \gamma\beta - \lambda\kappa, \kappa\lambda - (\delta\eta)^{2^{n-2}}, (\eta\delta)^{2^{n-2}} - \beta\gamma \rangle.$$

The corresponding decomposition matrix is given in Figure 17.

By [22, p. 306], if  $B$  has generalized quaternion defect groups and Ext quiver  $3\mathcal{K}$ , then  $B$  is Morita equivalent to  $Q(3\mathcal{K}) = k[3\mathcal{K}]/I_{Q(3\mathcal{K})}$  where

$$I_{Q(3\mathcal{K})} = \langle \delta\beta - \kappa\lambda\kappa, \gamma\eta - \lambda\kappa\lambda, \lambda\delta - \gamma\beta\gamma, \eta\kappa - \beta\gamma\beta, \beta\lambda - \eta(\delta\eta)^{2^{n-2}-1}, \\ \kappa\gamma - \delta(\eta\delta)^{2^{n-2}-1}, \delta\beta\gamma, \gamma\eta\delta, \eta\kappa\lambda \rangle.$$

The decomposition matrix for  $Q(3\mathcal{K})$  is given in Figure 18.

**4.10. Blocks with quivers  $3\mathcal{L}$ ,  $3\mathcal{Q}$  or  $3\mathcal{R}$ .** By [22, VIII.2.0] (resp. by [22, VII.4]), there are no blocks  $B$  with semidihedral (resp. generalized quaternion) defect groups that have Ext quivers  $3\mathcal{L}$ ,  $3\mathcal{Q}$  or  $3\mathcal{R}$ . By [22, IX.5.4], there are also no blocks  $B$  with dihedral defect groups that have Ext quivers  $3\mathcal{L}$ ,  $3\mathcal{Q}$  or  $3\mathcal{R}$ .

## 5. ORDINARY CHARACTERS BELONGING TO TAME BLOCKS

Assume Hypothesis 4.1. For  $2 \leq \ell \leq n-1$ , define  $\zeta_\ell = \zeta^{2^{n-1-\ell}}$ , so that  $\zeta_\ell$  is a primitive  $2^\ell$ -th root of unity. It follows from [14, 31] and [23] (see below) that every ordinary irreducible character of  $G$  which belongs to  $B$  can be realized by a simple  $F(\zeta)G$ -module. In particular, if we are only interested in the ordinary irreducible characters of  $G$  belonging to  $B$ , we can assume that  $\xi$  in §3 is equal to  $\zeta$ .

In [14] (resp. [31]), the ordinary irreducible characters of  $G$  belonging to  $B$  were analyzed if  $D$  is dihedral (resp. semidihedral or generalized quaternion). In the notation of [14, Sect. 4] (resp. [31, Sect. 2]), this means that we are either in Case  $(aa)$  or in Case  $(ab)$  or  $(ba)$ , see [14, Thm. 2] (resp. [31, Thms. 3.14-3.17]). Note that Case  $(ba)$  can only occur when  $D$  is semidihedral. In particular, in Case  $(aa)$  (resp. Case  $(ab)$  or  $(ba)$ ) there are precisely three (resp. two) isomorphism classes of simple  $B$ -modules.

*Remark 5.1.* (a) If  $D$  is quaternion of order 8, i.e.  $n = 3$ , it follows from [31, p. 220 and Thm. 3.17] that we are in Case  $(aa)$  and that there are precisely 4 (resp. 3) ordinary irreducible characters of height 0 (resp.  $1 = n - 2$ ) belonging to  $B$ . Moreover,  $B$  is Morita equivalent to one of the algebras in  $\{Q(3\mathcal{A})_2, Q(3\mathcal{B}), Q(3\mathcal{K})\}$  and in the decomposition matrices in Figures 8, 12 and 18, the ordinary irreducible characters of height 0 (resp. 1) are

$$\chi_1, \chi_2, \chi_3, \chi_4 \quad (\text{resp. } \chi_{5,1}, \chi_6, \chi_7).$$

By [31, Prop. 4.2], all ordinary irreducible characters are realizable over  $F$ , i.e. they correspond to absolutely irreducible  $FG$ -modules.

- (b) If  $D$  is not quaternion of order 8, then there are precisely 4 (resp.  $2^{n-2} - 1$ ) ordinary irreducible characters of height 0 (resp. 1) belonging to  $B$ . If  $D$  is dihedral, these are all ordinary irreducible characters belonging to  $B$ . If  $D$  is semidihedral, there is 0 (resp. 1) additional ordinary irreducible character of height  $n - 2$  belonging to  $B$  if we are in Case  $(ab)$  (resp. Cases  $(ba)$  or  $(aa)$ ). If  $D$  is generalized quaternion of order  $\geq 16$ , there are 1 (resp. 2) additional ordinary irreducible characters of height  $n - 2$  belonging to  $B$  if we are in Case  $(ab)$  (resp. Case  $(aa)$ ). In the decomposition matrices in Figures 1–18, the ordinary irreducible characters of height 0 (resp. 1) are

$$\chi_1, \chi_2, \chi_3, \chi_4 \quad (\text{resp. } \chi_{5,i} \text{ for } 1 \leq i \leq 2^{n-2} - 1),$$

whereas the ordinary irreducible characters of height  $n - 2$  are  $\chi_6$  or  $\chi_6, \chi_7$ , provided they exist.

Let  $\sigma$  be an element of order  $2^{n-1}$  in  $D$ . By [14, 31], there exists a block  $b_\sigma$  of  $kC_G(\sigma)$  with  $b_\sigma^G = B$  which contains a unique Brauer character  $\varphi^{(\sigma)}$  such that the following is true. There is an ordering of  $(1, 2, \dots, 2^{n-2} - 1)$  such that for  $1 \leq i \leq 2^{n-2} - 1$ , the generalized decomposition

number of  $G$  corresponding to  $\chi_{5,i}$ ,  $\sigma$  and  $\varphi^{(\sigma)}$  has the form

$$(5.8) \quad (d_{\sigma, \varphi^{(\sigma)}})_{\chi_{5,i}} = \begin{cases} \zeta^i + \zeta^{-i} & \text{if } D \text{ is dihedral or generalized quaternion,} \\ & \text{or if } D \text{ is semidihedral and } i \text{ is even,} \\ \zeta^i - \zeta^{-i} & \text{if } D \text{ is semidihedral and } i \text{ is odd with } i \leq 2^{n-3} - 1, \\ \zeta^{-i} - \zeta^i & \text{if } D \text{ is semidihedral and } i \text{ is odd with } i \geq 2^{n-3} + 1. \end{cases}$$

Note that the formulas in (5.8) for  $D$  semidihedral and  $i$  odd follow since the  $D$ -conjugacy classes of elements of order  $2^{n-1}$  in  $D$  are represented by

$$\sigma^1, \sigma^3, \dots, \sigma^{2^{n-3}-1}, \quad \sigma^{-(2^{n-3}+1)}, \sigma^{-(2^{n-3}+3)}, \dots, \sigma^{-(2^{n-2}-1)}.$$

If  $D$  is quaternion of order 8, we have that  $(d_{\sigma, \varphi^{(\sigma)}})_{\chi} = \zeta + \zeta^{-1} = 0$  for  $\chi \in \{\chi_{5,1}, \chi_6, \chi_7\}$ .

For  $2 \leq \ell \leq n-2$  define  $\nu_\ell = \zeta_\ell + \zeta_\ell^{-1}$ , and define

$$(5.9) \quad \nu_{n-1} = \begin{cases} \zeta + \zeta^{-1} & \text{if } D \text{ is dihedral or generalized quaternion,} \\ \zeta - \zeta^{-1} & \text{if } D \text{ is semidihedral.} \end{cases}$$

Note that  $W$  contains all roots of unity of order not divisible by 2. Hence by [14, 31] and by [23], the ordinary irreducible characters of height 0 or  $n-2$  belonging to  $B$  correspond to simple  $FG$ -modules. On the other hand, the characters  $\chi_{5,i}$ ,  $i = 1, \dots, 2^{n-2}-1$ , fall into  $n-2$  Galois orbits  $\mathcal{O}_2, \dots, \mathcal{O}_{n-1}$  under the action of  $\text{Gal}(F(\nu_{n-1})/F)$ . Namely for  $2 \leq \ell \leq n-1$ ,  $\mathcal{O}_\ell = \{\chi_{5,2^{n-1-\ell}(2u-1)} \mid 1 \leq u \leq 2^{\ell-2}\}$ . The field generated by the character values of each  $\xi_\ell \in \mathcal{O}_\ell$  over  $F$  is  $F(\nu_\ell)$ . Hence by [23], each  $\xi_\ell$  corresponds to an absolutely irreducible  $F(\nu_\ell)G$ -module  $X_\ell$ . By [25, Satz V.14.9], this implies that for  $2 \leq \ell \leq n-1$ , the Schur index of each  $\xi_\ell \in \mathcal{O}_\ell$  over  $F$  is 1. Hence we obtain  $n-2$  non-isomorphic simple  $FG$ -modules  $V_2, \dots, V_{n-1}$  with characters  $\rho_2, \dots, \rho_{n-1}$  satisfying

$$(5.10) \quad \rho_\ell = \sum_{\xi_\ell \in \mathcal{O}_\ell} \xi_\ell = \sum_{u=1}^{2^{\ell-2}} \chi_{5,2^{n-1-\ell}(2u-1)} \quad \text{for } 2 \leq \ell \leq n-1.$$

By [25, Hilfssatz V.14.7],  $\text{End}_{FG}(V_\ell)$  is a commutative  $F$ -algebra isomorphic to the field generated over  $F$  by the character values of any  $\xi_\ell \in \mathcal{O}_\ell$ . This means

$$(5.11) \quad \text{End}_{FG}(V_\ell) \cong F(\nu_\ell) \quad \text{for } 2 \leq \ell \leq n-1.$$

Suppose  $v_1, \dots, v_{l(\sigma)}$  form a complete system of representatives of  $C_G(\sigma)$ -conjugacy classes of 2-regular elements in  $C_G(\sigma)$  with  $v_1 = 1_G$ . By (3.6), for all  $1 \leq i \leq 2^{n-2}-1$ , the generalized decomposition number of  $G$  corresponding to  $\chi_{5,i}$ ,  $\sigma$  and  $\varphi^{(\sigma)}$  can be written as a  $W$ -linear combination of  $\chi_{5,i}(\sigma v_1), \dots, \chi_{5,i}(\sigma v_{l(\sigma)})$ , say

$$(5.12) \quad (d_{\sigma, \varphi^{(\sigma)}})_{\chi_{5,i}} = \tilde{w}_1 \cdot \chi_{5,i}(\sigma v_1) + \dots + \tilde{w}_{l(\sigma)} \cdot \chi_{5,i}(\sigma v_{l(\sigma)})$$

for certain  $\tilde{w}_1, \dots, \tilde{w}_{l(\sigma)} \in W$ .

By [14, Thm. 5] and [31, Prop. 4.6], the characters  $\chi_{5,i}$  have the same degree  $x$  and they are all of height 1 for  $1 \leq i \leq 2^{n-2}-1$ . Hence  $x = 2^{a-n+1}x^*$  where  $\#G = 2^a \cdot g^*$  and  $x^*$  and  $g^*$  are odd. Since the centralizer  $C_G(\sigma)$  contains  $\langle \sigma \rangle$ , we have  $\#C_G(\sigma) = 2^{n-1} \cdot 2^b \cdot m^*$  where  $b \geq 0$  and  $m^*$  is odd.

For  $1 \leq j \leq l(\sigma)$ , let  $C_j$  be the conjugacy class in  $G$  of  $\sigma v_j$ , and let  $t(C_j) \in WG$  be the class sum of  $C_j$ . Since for all  $j$ , the centralizer  $C_G(\sigma v_j)$  contains  $\langle \sigma \rangle$ , we have  $\#C_G(\sigma v_j) = 2^{n-1} \cdot 2^{b_j} \cdot m_j^*$  where  $b_j \geq 0$  and  $m_j^*$  is odd. We want to determine the action of  $t(C_j)$  on  $V_\ell$  for  $2 \leq \ell \leq n-1$ . For this, we identify  $\text{End}_{FG}(V_\ell) \cong F(\nu_\ell)$  with  $\text{End}_{F(\nu_\ell)G}(X_\ell)$  for one particular absolutely irreducible  $F(\nu_\ell)G$ -constituent  $X_\ell$  of  $V_\ell$  with character  $\xi_\ell$ . Using (5.10), we choose  $\xi_\ell = \chi_{5,2^{n-1-\ell}}$ . Then for  $2 \leq \ell \leq n-1$ , the action of  $t(C_j)$  on  $V_\ell$  is given as multiplication by  $\mu_j(\ell)$ , where

$$(5.13) \quad \mu_j(\ell) = \frac{\#C_j}{\xi_\ell(1)} \cdot \xi_\ell(\sigma v_j) = 2^{-b_j} \frac{g^*}{m_j^* \cdot x^*} \cdot \xi_\ell(\sigma v_j).$$

For  $1 \leq j \leq l(\sigma)$ , define  $w_j = 2^{b_j} \frac{m_j^+ \cdot x^*}{g^*} \tilde{w}_j$ . Then  $w_j \in W$  since  $g^*$  is odd, and  $w_j$  does not depend on  $\ell$ . By (5.8) and (5.9),

$$(d_{\sigma, \varphi(\sigma)})_{\xi_\ell} = (d_{\sigma, \varphi(\sigma)})_{\chi_{5, 2^{n-1-\ell}}} = \nu_\ell.$$

Therefore, (5.12) and (5.13) imply that

$$(5.14) \quad \nu_\ell = w_1 \cdot \mu_1(\ell) + \cdots + w_{l(\sigma)} \cdot \mu_{l(\sigma)}(\ell)$$

where  $w_1, \dots, w_{l(\sigma)} \in W$  are independent of  $\ell \in \{2, \dots, n-1\}$ .

*Remark 5.2.* If  $B$  is a principal block, the above formulas simplify considerably due to the fact that there is very little fusion of  $D$ -conjugacy classes in  $G$  in this case. More precisely, following Brauer's arguments in [12, §VII], suppose  $D$  is a Sylow 2-subgroup of  $G$  and let  $S = \langle \sigma \rangle$ . If  $\sigma^\lambda$  is not in the center of  $D$ , then  $S$  is a Sylow 2-subgroup of  $C_G(\sigma^\lambda)$ . Hence if  $\sigma^\lambda$  and  $\sigma^\mu$  are conjugate in  $G$ , we can use Sylow's theorems to see that they are conjugate in  $N_G(S)$ . Since  $N_G(S)/C_G(S)$  is a 2-group, it must be of order 2. Thus  $N_G(S)$  is generated by  $D$  and  $C_G(S)$ , which implies that  $\sigma^\lambda$  and  $\sigma^\mu$  are conjugate in  $D$ . Using (3.7) together with [14, Prop. (4A)] and [31, Prop. 2.10], we obtain that

$$(5.15) \quad \chi_{5,i}(\sigma) = (d_{\sigma, \varphi(\sigma)})_{\chi_{5,i}} \varphi^{(\sigma)}(1_G)$$

where  $\varphi^{(\sigma)}$  is as in (5.8). Moreover, since  $\varphi^{(\sigma)}$  is the unique Brauer character belonging to the block  $b_\sigma$  of  $kC_G(\sigma)$  satisfying  $b_\sigma^G = B$ , it follows by Brauer's Third Main Theorem (see e.g. [1, Thm. 16.1]) that  $b_\sigma$  is the principal block of  $kC_G(\sigma)$ , which implies that  $\varphi^{(\sigma)}$  is the trivial character of  $b_\sigma$ . Putting (5.15) into (5.13) for  $j = 1$  therefore implies that if  $\omega = 2^{b_1} \frac{m_1^+ \cdot x^*}{g^*}$ , which lies in  $W$ , then for all  $2 \leq \ell \leq n-1$

$$\nu_\ell = (d_{\sigma, \varphi(\sigma)})_{\chi_{5, 2^{n-1-\ell}}} = \omega \cdot \mu_1(\ell).$$

Note that in [3, §3.4], [4, §5] and [9, §3.2], it was assumed that the formula (5.15) was also true for non-principal blocks. However, this may not be true since there could be more fusion of  $D$ -conjugacy classes in  $G$  in this case. More precisely, let  $Y$  be a full set of representatives of  $D$ -conjugacy classes of the elements of the set  $\{\sigma^r \mid r \text{ odd}\}$ , and let  $Y_\sigma$  be the set of all  $y \in Y$  which are conjugate to  $\sigma$  in  $G$ . If  $B$  is not principal, then  $|Y_\sigma|$  may be strictly greater than 1. By [14, Prop. (4A)] and [31, Prop. 2.10], the set  $\{(y, b_\sigma) \mid y \in Y_\sigma\}$  is a system of representatives for the conjugacy classes of subsections for  $B$  such that  $y$  is conjugate to  $\sigma$  in  $G$ . Hence it follows from Remark 3.1 that for non-principal blocks  $B$ , (5.15) has to be replaced by the formula

$$\chi_{5,i}(\sigma) = \sum_{y \in Y_\sigma} (d_{y, \varphi(\sigma)})_{\chi_{5,i}} \varphi^{(\sigma)}(1_G).$$

**Definition 5.3.** Use the notation introduced above, and in particular (5.9).

(i) Define

$$p_n(t) = \prod_{\ell=2}^{n-1} \text{min.pol.}_F(\nu_\ell)$$

and let  $R' = W[[t]]/(p_n(t))$ .

(ii) Let  $Z = \langle \sigma \rangle$  be a cyclic group of order  $2^{n-1}$ . Let  $\tau : Z \rightarrow Z$  be the group automorphism which sends  $\sigma$  to  $\sigma^{-1}$  if  $D$  is dihedral or generalized quaternion, and which sends  $\sigma$  to  $\sigma^{-1+2^{n-2}}$  if  $D$  is semidihedral. Then  $\tau$  can be extended to a  $W$ -algebra automorphism of the group ring  $WZ$  which will again be denoted by  $\tau$ . Let  $T(\sigma^2) = 1 + \sigma^2 + \sigma^4 + \cdots + \sigma^{2^{n-1}-2}$ , and define

$$S' = (WZ)^{(\tau)} / (T(\sigma^2), \sigma T(\sigma^2)).$$

*Remark 5.4.* The minimal polynomial  $\text{min.pol.}_F(\nu_\ell)$  for  $2 \leq \ell \leq n-1$  is as follows:

$$\begin{aligned} \text{min.pol.}_F(\nu_2)(t) &= t, \\ \text{min.pol.}_F(\nu_\ell)(t) &= \text{min.pol.}_F(\nu_{\ell-1})(t^2 - 2) \quad \text{for } 3 \leq \ell \leq n-2, \\ \text{min.pol.}_F(\nu_{n-1})(t) &= \begin{cases} \text{min.pol.}_F(\nu_{n-2})(t^2 - 2) & \text{if } D \text{ is dihedral or} \\ & \text{generalized quaternion,} \\ \text{min.pol.}_F(\nu_{n-2})(t^2 + 2) & \text{if } D \text{ is semidihedral.} \end{cases} \end{aligned}$$

The  $W$ -algebra  $R'$  from Definition 5.3 is a complete local commutative Noetherian ring with residue field  $k$ . Moreover,

$$\begin{aligned} F \otimes_W R' &\cong \prod_{\ell=2}^{n-1} F(\nu_\ell) \quad \text{as } F\text{-algebras,} \\ k \otimes_W R' &\cong k[t]/(t^{2^{n-2}-1}) \quad \text{as } k\text{-algebras.} \end{aligned}$$

Additionally,  $R'$  is isomorphic to the  $W$ -subalgebra of  $\prod_{\ell=2}^{n-1} W[\nu_\ell]$  generated by the element  $(\nu_\ell)_{\ell=2}^{n-1}$ .

**Lemma 5.5.** *Using the notation of Definition 5.3, there is a continuous  $W$ -algebra isomorphism  $h : R' \rightarrow S'$  with  $h(t) = \sigma + \tau(\sigma)$ . In particular,  $R'$  is isomorphic to a subquotient algebra of the group algebra  $WD$ .*

*Moreover, if  $D$  is dihedral or semidihedral, then the ring  $W[[t]]/(t p_n(t), 2 p_n(t))$  is also isomorphic to a subquotient algebra of the group algebra  $WD$ .*

*Proof.* If  $D$  is dihedral or generalized quaternion, this follows from [3, Lemma 2.3.6] and [4, Lemma 5.3].

For the remainder of the proof, assume that  $D$  is semidihedral. Let  $J = \sigma^{2^{n-2}}$  so that  $\tau(\sigma) = J\sigma^{-1}$ . Note that  $(WZ)^{\langle \tau \rangle}$  is generated as a  $W$ -algebra by  $(\sigma + J\sigma^{-1})$  and  $J$ . Moreover,  $(WZ)^{\langle \tau \rangle}$  is a free  $W$ -module of rank  $2^{n-2} + 1$  with  $W$ -basis given by

$$(5.16) \quad \begin{aligned} &(\sigma^{\pm 1} + J\sigma^{\mp 1}), (\sigma^{\pm 3} + J\sigma^{\mp 3}), \dots, (\sigma^{\pm(2^{n-3}-1)} + J\sigma^{\mp(2^{n-3}-1)}), \\ &1, J, (\sigma^2 + \sigma^{-2}), (\sigma^4 + \sigma^{-4}), \dots, (\sigma^{2^{n-2}-2} + \sigma^{-(2^{n-2}-2)}). \end{aligned}$$

Note that the  $W$ -subalgebra of  $(WZ)^{\langle \tau \rangle}$  generated by  $(\sigma + J\sigma^{-1})$  is a free  $W$ -module of the same rank  $2^{n-2} + 1$  and with almost the same  $W$ -basis as in (5.16) except that  $J$  must be replaced by  $2J$ . It follows that  $S' = (WZ)^{\langle \tau \rangle} / (T(\sigma^2), \sigma T(\sigma^2))$  is generated as a  $W$ -algebra by the image of  $(\sigma + J\sigma^{-1})$  in  $S'$ . Hence we have a surjective  $W$ -algebra homomorphism

$$f : W[[t]] \rightarrow S' = (WZ)^{\langle \tau \rangle} / (T(\sigma^2), \sigma T(\sigma^2))$$

sending  $t$  to the image in  $S'$  of  $\sigma + J\sigma^{-1} = \sigma + \tau(\sigma)$ . Using the injective  $W$ -algebra homomorphism

$$\begin{aligned} \iota : WZ &\rightarrow W \times W \times \prod_{\ell=2}^{n-1} W[\zeta_\ell] \\ \sigma &\mapsto \left( 1, -1, (\zeta_\ell)_{\ell=2}^{n-1} \right) \end{aligned}$$

it is straightforward to prove that there exists an odd integer  $c_n$  such that  $p_n(\sigma + J\sigma^{-1}) = c_n \sigma T(\sigma^2)$ . More precisely,  $c_4 = 3$  and  $c_n = 2c_{n-1}^2 - 1$  for  $n \geq 5$ . Thus  $p_n(t)$  lies in the kernel of  $f$ . Since both  $R' = W[[t]]/(p_n(t))$  and  $S'$  are free as  $W$ -modules of the same rank  $2^{n-2} - 1$ , it follows that  $R' = W[[t]]/(p_n(t)) \cong S'$  as  $W$ -algebras.

To finish the proof of Lemma 5.5, it suffices to show that the ring  $W[[t]]/((t-2)p_n(t))$  is isomorphic to a subquotient algebra of  $WD$ . Define

$$\Theta = (WZ)^{\langle \tau \rangle} / (T(\sigma^2) - \sigma T(\sigma^2)).$$

Then  $\Theta$  is isomorphic to a subquotient algebra of  $WD$  and it is generated as a  $W$ -algebra by the image of  $(\sigma + J\sigma^{-1})$  in  $\Theta$ . Moreover,  $\Theta$  is a free  $W$ -module of rank  $2^{n-2}$ , since the ideal

$(T(\sigma^2) - \sigma T(\sigma^2))$  is generated over  $W$  by  $T(\sigma^2) - \sigma T(\sigma^2)$ . Define a surjective  $W$ -algebra homomorphism  $\theta : W[[t]] \rightarrow \Theta$  by sending  $t$  to the image in  $\Theta$  of  $\sigma + J\sigma^{-1} = \sigma + \tau(\sigma)$ . Using the above calculations, we see that

$$\begin{aligned} \theta((t-2)p_n(t)) &= ((\sigma + J\sigma^{-1}) - 2)p_n(\sigma + J\sigma^{-1}) \\ &= ((\sigma + J\sigma^{-1}) - 2)c_n \sigma T(\sigma^2) \\ &= 2c_n [T(\sigma^2) - \sigma T(\sigma^2)] \end{aligned}$$

which is zero in  $\Theta$ . Hence  $(t-2)p_n(t)$  lies in the kernel of  $\theta$ . Since both  $W[[t]]/((t-2)p_n(t))$  and  $\Theta$  are free over  $W$  of rank  $2^{n-2}$ , it follows that  $W[[t]]/((t-2)p_n(t))$  is isomorphic to  $\Theta$ , which completes the proof of Lemma 5.5.  $\square$

The following result gives a correction of [4, Lemma 5.4] and generalizes the corrected result to all tame blocks with at least two isomorphism classes of simple modules.

**Lemma 5.6.** *Assume Hypothesis 4.1 and use the notation introduced above, and in particular (5.9), (5.10) and Definition 5.3. If  $D$  is quaternion of order 8, let  $U'$  be a  $WG$ -module which is free over  $W$  and whose  $F$ -character is either  $\chi_{5,1}$ , or  $\chi_6$ , or  $\chi_7$ . If  $D$  is not quaternion of order 8, let  $U'$  be a  $WG$ -module which is free over  $W$  and whose  $F$ -character is equal to*

$$\sum_{\ell=2}^{n-1} \rho_\ell = \sum_{i=1}^{2^{n-2}-1} \chi_{5,i}.$$

- (i) *There exists a  $WG$ -module endomorphism  $\alpha$  of  $U'$  such that the  $W$ -algebra  $W[\alpha]$  generated by  $\alpha$  is isomorphic to  $R'$ .*
- (ii) *Suppose either that  $n = 3$ , or that  $n \geq 4$  and  $\text{End}_{kG}(U'/2U') \cong R'/2R'$  and  $U'/2U'$  is free as a module for  $\text{End}_{kG}(U'/2U')$  of rank  $\deg(\chi_{5,1})$ . Then  $\text{End}_{WG}(U') = W[\alpha] \cong R'$  and  $U'$  is free as a module for  $\text{End}_{WG}(U')$ .*

*Proof.* Suppose first that  $n = 3$ . Then  $p_n(t) = t$  and  $R' \cong W$ . Hence we can choose the  $WG$ -module endomorphism  $\alpha$  of  $U'$  to be the zero endomorphism. Since  $U'$  is free as a  $W$ -module and  $F \otimes_W \text{End}_{WG}(U') \cong \text{End}_{FG}(F \otimes_W U') \cong F$ , it follows that  $\text{End}_{WG}(U') \cong W \cong R'$ .

For the remainder of the proof, assume  $n \geq 4$ . We first construct a  $WG$ -module endomorphism  $\alpha$  of  $U'$  as in part (i) of the lemma. As before, let  $\sigma \in D$  be an element of order  $2^{n-1}$  and let  $\{v_1, \dots, v_{l(\sigma)}\}$  be a complete system of representatives of  $C_G(\sigma)$ -conjugacy classes of 2-regular elements in  $C_G(\sigma)$ . For  $1 \leq j \leq l(\sigma)$ , let  $C_j$  be the conjugacy class in  $G$  of  $\sigma v_j$ , and let  $t(C_j) \in WG$  be the class sum of  $C_j$ .

Let  $1 \leq j \leq l(\sigma)$ . Since  $t(C_j)$  lies in the center of  $WG$ , multiplication by  $t(C_j)$  defines a  $WG$ -module endomorphism of  $U'$ . Since  $U'$  is free as a  $W$ -module,  $\text{End}_{WG}(U')$  can naturally be identified with a  $W$ -subalgebra of

$$F \otimes_W \text{End}_{WG}(U') \cong \text{End}_{FG}(F \otimes_W U') \cong \prod_{\ell=2}^{n-1} \text{End}_{FG}(V_\ell) \cong \prod_{\ell=2}^{n-1} F(\nu_\ell).$$

Therefore,  $t(C_j)$  acts on  $U'$  as multiplication by a scalar  $\lambda_j$  in the maximal  $W$ -order  $\prod_{\ell=2}^{n-1} W[\nu_\ell]$  in  $\prod_{\ell=2}^{n-1} F(\nu_\ell)$ . Moreover,  $\lambda_j$  can be determined from the action of  $t(C_j)$  on  $F \otimes_W U' \cong \bigoplus_{\ell=2}^{n-1} V_\ell$ , which implies by (5.13) that  $\lambda_j = (\mu_j(\ell))_{\ell=2}^{n-1}$ . By (5.14), there exist elements  $w_1, \dots, w_{l(\sigma)} \in W$  such that

$$\nu_\ell = w_1 \cdot \mu_1(\ell) + \dots + w_{l(\sigma)} \cdot \mu_{l(\sigma)}(\ell)$$

for  $2 \leq \ell \leq n-1$ . Define  $\alpha$  to be the  $WG$ -module endomorphism of  $U'$  given by multiplication by the element  $\sum_{j=1}^{l(\sigma)} w_j t(C_j)$  in the center of  $WG$ . Then  $\alpha$  acts on  $U'$  as multiplication by the scalar  $(\nu_\ell)_{\ell=2}^{n-1} \in \prod_{\ell=2}^{n-1} W[\nu_\ell]$ , which implies  $W[\alpha] \cong R'$  (see Remark 5.4).

Let now  $\overline{U'} = U'/2U'$  and suppose that  $\text{End}_{kG}(\overline{U'}) \cong R'/2R' \cong k[t]/(t^{2^{n-2}-1})$  and that  $\overline{U'}$  is free as a module for  $\text{End}_{kG}(U'/2U')$  of rank  $x = \deg(\chi_{5,1})$ . Note that  $x = \deg(\chi_{5,i})$  for all

$1 \leq i \leq 2^{n-2} - 1$  and that  $\dim_{F(\nu_\ell)} V_\ell = x$  for all  $2 \leq \ell \leq n - 1$ , which implies that  $F \otimes_W U'$  is a free module of rank  $x$  for  $\text{End}_{FG}(F \otimes_W U')$ . We have a short exact sequence of  $W$ -modules

$$0 \longrightarrow \text{End}_W(U') \xrightarrow{\cdot 2} \text{End}_W(U') \xrightarrow{\text{mod } 2} \text{End}_k(\overline{U'}) \longrightarrow 0.$$

Considering the  $G$ -action on these modules, we obtain an exact sequence of  $W$ -modules

$$(5.17) \quad 0 \longrightarrow \text{End}_{WG}(U') \xrightarrow{\cdot 2} \text{End}_{WG}(U') \xrightarrow{\text{mod } 2} \text{End}_{kG}(\overline{U'}) \longrightarrow H^1(G, \text{End}_W(U')).$$

Because  $\text{End}_{WG}(U')$  is a  $W$ -submodule of the free  $W$ -module  $\text{End}_W(U')$ , it follows that  $\text{End}_{WG}(U')$  is a free  $W$ -module of rank

$$\dim_F(F \otimes_W \text{End}_{WG}(U')) = \dim_F \text{End}_{FG}(F \otimes_W U') = 2^{n-2} - 1.$$

Since by assumption  $\dim_k \text{End}_{kG}(\overline{U'}) = \dim_k(R'/2R') = 2^{n-2} - 1$ , it follows that (5.17) induces a short exact sequence of  $W$ -modules

$$(5.18) \quad 0 \longrightarrow \text{End}_{WG}(U') \xrightarrow{\cdot 2} \text{End}_{WG}(U') \xrightarrow{\text{mod } 2} \text{End}_{kG}(\overline{U'}) \longrightarrow 0.$$

Let  $\overline{\beta}$  be a generator of  $\text{End}_{kG}(\overline{U'})$  as a  $k$ -algebra. By (5.18), there exists  $\beta \in \text{End}_{WG}(U')$  whose induced  $kG$ -module endomorphism of  $\overline{U'}$  is equal to  $\overline{\beta}$ . Using Nakayama's lemma, we see that  $\text{End}_{WG}(U') = W[\beta]$ . Since the maximal ideal  $\mathfrak{m}_{W[\beta]}$  of  $W[\beta]$  is generated by 2 and  $\beta$  and the maximal ideal  $\mathfrak{m}_{k[\overline{\beta}]}$  of  $k[\overline{\beta}]$  is generated by  $\overline{\beta}$ , it follows that

$$U'/\mathfrak{m}_{W[\beta]}U' \cong \overline{U'}/\overline{\beta}(\overline{U'}) \cong \overline{U'}/\mathfrak{m}_{k[\overline{\beta}]} \overline{U'}$$

where the latter has  $k$ -dimension  $x = \deg(\chi_{5,1})$  by assumption. By Nakayama's lemma, we can lift a  $k$ -basis  $\{\overline{b}_1, \dots, \overline{b}_x\}$  of  $U'/\mathfrak{m}_{W[\beta]}U'$  to a set  $\{b_1, \dots, b_x\}$  of generators of  $U'$  over  $W[\beta] = \text{End}_{WG}(U')$ . Because  $F \otimes_W U'$  is a free module of rank  $x$  for  $\text{End}_{FG}(F \otimes_W U') \cong F \otimes_W \text{End}_{WG}(U')$ , it follows that  $b_1, \dots, b_x$  are linearly independent over  $\text{End}_{WG}(U')$ . Hence  $U'$  is free as a module for  $\text{End}_{WG}(U')$  of rank  $x = \deg(\chi_{5,1})$ .

It remains to show that  $\text{End}_{WG}(U') = W[\alpha]$ , i.e. we need to show  $W[\alpha] = W[\beta]$ . We identify both  $W[\alpha]$  and  $W[\beta]$  with  $W$ -subalgebras of the maximal  $W$ -order  $\prod_{\ell=2}^{n-1} W[\nu_\ell]$  in  $\prod_{\ell=2}^{n-1} F(\nu_\ell) \cong F \otimes_W \text{End}_{WG}(U')$ . Since  $W[\alpha] \subseteq W[\beta]$ , there exists a polynomial  $q(x) \in W[x]$  such that  $\alpha = q(\beta)$ . Because  $\alpha$  is not a unit, the constant coefficient of  $q(x)$  must be divisible by 2. Write

$$q(x) = 2a_0 + b_1x + b_2x^2 + \dots + b_dx^d$$

for certain  $a_0, b_1, \dots, b_d \in W$  and  $d \geq 1$ . Suppose  $b_1 = 2a_1$  for some  $a_1 \in W$ . Consider the natural projection

$$\pi_{n-1} : \prod_{\ell=2}^{n-1} W[\nu_\ell] \rightarrow W[\nu_{n-1}],$$

and let  $\mathfrak{m}$  be the maximal ideal of  $W[\nu_{n-1}]$ . Since  $n \geq 4$ , either  $\sqrt{2}$  or  $\sqrt{-2}$  lies in  $\mathfrak{m}$ . Hence  $2 \in \mathfrak{m}^2$ . Since  $2\pi_{n-1}(\beta), \pi_{n-1}(\beta)^2, \dots, \pi_{n-1}(\beta)^d$  all lie in  $\mathfrak{m}^2$ , it follows that  $\pi_{n-1}(\alpha) \in \mathfrak{m}^2$ . However, we have seen that  $\pi_{n-1}(\alpha) = \nu_{n-1} \notin \mathfrak{m}^2$ . Hence  $b_1$  is not divisible by 2. But then the  $kG$ -module endomorphism  $\overline{\alpha}$  of  $\overline{U'}$  which is induced by  $\alpha$  has the form

$$\overline{\alpha} = u\overline{\beta} + c_2\overline{\beta}^2 + \dots + c_d\overline{\beta}^d$$

where  $u \in k^*$  and  $c_2, \dots, c_d \in k$ . This means that  $k[\overline{\alpha}] = k[\overline{\beta}] = \text{End}_{kG}(\overline{U'})$ , which implies by Nakayama's lemma that  $\text{End}_{WG}(U') = W[\alpha]$ .  $\square$

## 6. UNIVERSAL DEFORMATION RINGS

Assume Hypothesis 4.1 and the notation introduced in §4 and §5. In this section, we want to determine  $R(G, V)$  for any indecomposable  $kG$ -module  $V$  belonging to  $B$  whose stable endomorphism ring is isomorphic to  $k$  and whose Brauer character equals the restriction to the 2-regular conjugacy classes of an irreducible  $\overline{F}$ -character of  $G$  of height 1. In other words, if  $D$  is not quaternion of order 8 then the Brauer character of  $V$  is the restriction to the 2-regular conjugacy classes of  $\chi_{5,i}$  for all  $1 \leq i \leq 2^{n-2} - 1$ , and if  $D$  is quaternion of order 8 then the Brauer character of  $V$  is the restriction to the 2-regular conjugacy classes of either  $\chi_{5,1}$ , or  $\chi_6$ , or  $\chi_7$ .

The following lemma gives a list of all such  $V$  together with  $\text{End}_{kG}(V)$  and  $\text{Ext}_{kG}^1(V, V)$ . This result is proved using the description of the quiver and relations of the basic algebra of  $B$  together with its decomposition matrix as provided in §4.

**Lemma 6.1.** *Assume Hypothesis 4.1. Let  $\Lambda = kQ/I$  be a basic algebra such that  $B$  is Morita equivalent to  $\Lambda$ . For each vertex  $j$  in  $Q$ , let  $T_j$  denote the simple  $B$ -module corresponding to the simple  $\Lambda$ -module associated to  $j$ . Let  $V$  be an indecomposable  $kG$ -module belonging to  $B$  such that  $\underline{\text{End}}_{kG}(V) \cong k$  and such that the Brauer character of  $V$  is equal to the restriction to the 2-regular conjugacy classes of an irreducible  $\overline{F}$ -character of  $G$  of height 1.*

- (a) *Suppose  $D$  is quaternion of order 8. If  $Q = 3A$  or  $Q = 3B$  then  $V$  is either isomorphic to  $T_1$  or  $T_2$ , or  $V$  is a uniserial module of length 4 whose radical quotient or socle is isomorphic to  $T_0$ . If  $Q = 3K$  then  $V$  is an arbitrary uniserial module of length 2.*
- (b) *Suppose  $D$  is not quaternion of order 8.*
  - (i) *If  $Q = 2A$  then  $V$  is a uniserial module with descending composition factors  $T_0, T_0, T_1$ , or  $T_1, T_0, T_0$ .*
  - (ii) *If  $Q = 2B$  and  $B$  is Morita equivalent to neither  $\text{SD}(2B)_4(c)$  nor  $\text{Q}(2B)_2(p, a, c)$  then  $V$  is isomorphic to  $T_1$ . If  $B$  is Morita equivalent to  $\text{SD}(2B)_4(c)$  or  $\text{Q}(2B)_2(p, a, c)$  then  $V$  is a uniserial module with descending composition factors  $T_0, T_1$ , or  $T_1, T_0$ .*
  - (iii) *If  $Q = 3A$  then  $V$  is a uniserial module with descending composition factors  $T_0, T_1, T_0, T_2$ , or  $T_2, T_0, T_1, T_0$ , or  $T_0, T_2, T_0, T_1$ , or  $T_1, T_0, T_2, T_0$ .*
  - (iv) *If  $Q \in \{3B, 3D\}$  then  $V$  is isomorphic to  $T_1$ .*
  - (v) *If  $B$  is Morita equivalent to  $\text{SD}(3C)_{2,1}$  then  $V$  is isomorphic to  $T_0$ . If  $B$  Morita equivalent to  $\text{SD}(3C)_{2,2}$  then  $V$  is indecomposable with descending radical factors  $T_0, T_1 \oplus T_2$ , or  $T_1 \oplus T_2, T_0$ , or  $V$  is uniserial with descending composition factors  $T_1, T_0, T_2$ , or  $T_2, T_0, T_1$ .*
  - (vi) *If  $Q \in \{3H, 3K\}$  and  $B$  is not Morita equivalent to  $\text{SD}(3H)_2$  then  $V$  is a uniserial module with descending composition factors  $T_1, T_2$ , or  $T_2, T_1$ . If  $B$  is Morita equivalent to  $\text{SD}(3H)_2$  then  $V$  is a uniserial module with descending composition factors  $T_0, T_1$ , or  $T_1, T_0$ .*

In all these cases,  $\text{End}_{kG}(V) \cong k$ . Moreover,  $\text{Ext}_{kG}^1(V, V) = 0$  if  $D$  is quaternion of order 8, and  $\text{Ext}_{kG}^1(V, V) \cong k$  in all other cases.

**Definition 6.2.** Assume Hypothesis 4.1, and let  $V$  be as in Lemma 6.1. We say  $V$  corresponds to a 3-tube if there exists an indecomposable quotient module  $\overline{U}$  of the projective  $kG$ -module cover  $P_V$  of  $V$  such that

- (a)  $\overline{U}$  defines a lift of  $V$  over  $k[t]/(t^{2^{n-2}})$ , and
- (b)  $\overline{U}$  belongs to a 3-tube of the stable Auslander-Reiten quiver of  $B$ .

The following result is again proved using the description of the quiver and relations of the basic algebra of  $B$  together with its decomposition matrix as provided in §4. For blocks with generalized quaternion defect groups, we also use [22, VII.1 and VII.1.1].

**Lemma 6.3.** *Assume Hypothesis 4.1, and let  $V$  be as in Lemma 6.1. If  $D$  is dihedral then  $V$  corresponds to a 3-tube, and if  $D$  is generalized quaternion then  $V$  does not correspond to a 3-tube. If  $D$  is semidihedral then  $V$  corresponds to a 3-tube if and only if either*



- $B$  is Morita equivalent to one of the algebras in

$$\{\text{SD}(2\mathcal{A})_2(c), \text{SD}(2\mathcal{B})_1(c), \text{SD}(3\mathcal{B})_1, \text{SD}(3\mathcal{C})_{2,1}\}$$

and  $V$  is arbitrary; or

- $B$  is Morita equivalent to  $\text{SD}(3\mathcal{A})_1$  and  $V$  is such that its radical quotient or its socle is isomorphic to  $T_1$ ; or
- $B$  is Morita equivalent to  $\text{SD}(3\mathcal{C})_{2,2}$  and  $V$  is such that its radical quotient or its socle is isomorphic to  $T_0$ ; or
- $B$  is Morita equivalent to  $\text{SD}(3\mathcal{H})_1$  and  $V$  is such that its radical quotient is isomorphic to  $T_1$ ;
- $B$  is Morita equivalent to  $\text{SD}(3\mathcal{H})_2$  and  $V$  is such that its radical quotient is isomorphic to  $T_0$ .

The following result shows that if  $V$  corresponds to a 3-tube, then the module  $\overline{U}$  from Definition 6.2 has a universal deformation ring  $R(G, \overline{U}) \cong k$ .

**Proposition 6.4.** *Assume Hypothesis 4.1, and let  $V$  be as in Lemma 6.1. Moreover suppose that  $V$  corresponds to a 3-tube. Let  $\overline{U}$  be the  $kG$ -module from Definition 6.2 which belongs to a 3-tube of the stable Auslander-Reiten quiver of  $B$ . Then  $\underline{\text{End}}_{kG}(\overline{U}) \cong k$  and  $R(G, \overline{U}) \cong k$ .*

*Proof.* If  $D$  is dihedral, this follows from [3, Sect. 5.2] and [9, Prop. 6.3]. For the remainder of the proof, assume that  $D$  is semidihedral.

Let  $\mathfrak{T}$  be the 3-tube of the stable Auslander-Reiten quiver of  $B$  to which  $\overline{U}$  belongs. Then  $\overline{U}$  belongs to the boundary of  $\mathfrak{T}$ . Going through the cases described in Lemma 6.3, it is straightforward to show that  $\underline{\text{End}}_{kG}(\overline{U}) \cong k$  and  $\text{Ext}_{kG}^1(\overline{U}, \overline{U}) = 0$ .

Let  $K$  be a vertex of  $\overline{U}$ . Because of the work in [22, Chapter V], and in particular [22, V.4.2.1 and proof of V.4.2], we have the following facts:

- The group  $K$  is a Klein four group and the quotient group  $N_G(K)/C_G(K)$  is isomorphic to a symmetric group  $S_3$ .
- There is a block  $b$  of  $kN_G(K)$  with  $b^G = B$  such that the Green correspondent  $f\overline{U}$  of  $\overline{U}$  belongs to the boundary of a 3-tube in the stable Auslander-Reiten quiver of  $b$ . Moreover,  $b$  is Morita equivalent to  $kS_4$  modulo the socle.

Using these facts in lieu of [3, Facts 5.2.1], we can use similar arguments as in the proofs of [3, Prop. 5.2.4 and Cor. 5.2.5] to show that  $R(G, \overline{U}) \cong k$ .  $\square$

**Theorem 6.5.** *Assume Hypothesis 4.1, and let  $V$  be as in Lemma 6.1. Then*

$$R(G, V) \cong \begin{cases} W[[t]]/(tp_n(t), 2p_n(t)) & \text{if } V \text{ corresponds to a 3-tube,} \\ W[[t]]/(p_n(t)) & \text{if } V \text{ does not correspond to a 3-tube,} \end{cases}$$

where  $p_n(t)$  is as in Definition 5.3. In all cases, the ring  $R(G, V)$  is a subquotient ring of  $WD$ , and it is a complete intersection ring if and only if  $V$  does not correspond to a 3-tube.

*Proof.* Suppose first that  $D$  is quaternion of order 8. By Lemma 6.1,  $\text{Ext}_{kG}^1(V, V) = 0$ , which implies that  $R(G, V)$  is isomorphic to a quotient algebra of  $W$ . Using the decomposition matrices in Figures 8, 12 and 18 together with [18, Prop. (23.7)], we see that in all cases  $V$  can be lifted over  $W$ , which implies that  $R(G, V) \cong W \cong W[[t]]/(p_3(t))$ .

For the remainder of the proof, assume that  $D$  is not quaternion of order 8. In particular, the Brauer character of  $V$  is the restriction of  $\chi_{5,1}$  to the 2-regular conjugacy classes of  $G$ . By Lemma 6.1,  $\text{Ext}_{kG}^1(V, V) \cong k$ , which implies that  $R(G, V)$  is isomorphic to a quotient algebra of  $W[[t]]$  but not to a quotient algebra of  $W$ .

Considering the projective  $kG$ -module cover  $P_V$  of  $V$ , we see that there exists an indecomposable quotient module  $\overline{U}'$  of  $P_V$  such that  $\overline{U}'$  defines a lift of  $V$  over  $k[t]/(t^{2^n-2}-1)$  and  $\overline{U}'/t^2\overline{U}'$  is an indecomposable  $kG$ -module. Moreover, if  $V$  corresponds to a 3-tube then the indecomposable  $kG$ -module  $\overline{U}$  of Definition 6.2 defines a lift of  $V$  over  $k[t]/(t^{2^n-2})$  and  $\overline{U}/t^{2^n-2}\overline{U} \cong \overline{U}'$ .

*Claim 1.* The universal mod 2 deformation ring  $R(G, V)/2R(G, V)$  is isomorphic to  $k[t]/(t^{2^{n-2}})$  (resp.  $k[t]/(t^{2^{n-2}-1})$ ) and the universal mod 2 deformation of  $V$  is isomorphic to  $\overline{U}$  (resp.  $\overline{U}'$ ) if  $V$  corresponds to a 3-tube (resp. does not correspond to a 3-tube).

*Proof of Claim 1.* If  $B$  is Morita equivalent neither to  $\text{SD}(2\mathcal{B})_4(c)$ , nor to  $\text{Q}(2\mathcal{B})_2(p, a, c)$ , nor to  $\text{SD}(3\mathcal{C})_{2,2}$ , then Claim 1 follows from [4, Lemma 2.5].

If  $B$  is Morita equivalent to  $\text{SD}(2\mathcal{B})_4(c)$  or  $\text{Q}(2\mathcal{B})_2(p, a, c)$ , it follows from Lemma 6.1(b)(ii) that  $V$  is uniserial with descending composition factors  $T_u, T_v$  where  $\{u, v\} = \{0, 1\}$ . The projective cover  $P_V = P_{T_u}$  has a unique submodule  $K_u$  which is uniserial with descending composition factors  $T_u, T_u$ , and  $\overline{U}' = P_{T_u}/K_u$  can be visualized as follows, where  $T_u$  (resp.  $T_v$ ) occurs  $2^{n-2} - 1$  times.

$$\overline{U}' = \begin{array}{ccc} & T_u & \\ & \diagdown & \diagup \\ T_u & & T_v \\ & \diagup & \diagdown \\ T_u & & T_v \\ & \diagdown & \diagup \\ \vdots & & \vdots \\ & \diagup & \diagdown \\ T_u & & T_v \\ & \diagdown & \diagup \\ T_u & & T_v \\ & T_v & \end{array}$$

Using the relations in §4.2, we see that there is a unique  $B$ -submodule  $V'$  of  $\overline{U}'$  which is isomorphic to  $V$ . Moreover, if  $T = \overline{U}'/V'$  then there is a unique  $B$ -submodule  $T'$  of  $\overline{U}'$  such that there are  $kG$ -module isomorphisms  $\varphi : \overline{U}'/T' \rightarrow V$  and  $\psi : \overline{U}'/V' = T \rightarrow T'$ . Since  $\text{Ext}_{kG}^1(\overline{U}', V) = 0$  and since every surjective  $kG$ -module homomorphism  $\overline{U}' \rightarrow V$  must have kernel equal to  $T$ , we can argue as in the proof of [4, Lemma 2.5] to show that  $R(G, V)/2R(G, V)$  is isomorphic to  $k[t]/(t^{2^{n-2}-1})$  and that the universal mod 2 deformation of  $V$  is isomorphic to  $\overline{U}'$ . This implies Claim 1 in case  $B$  is Morita equivalent to  $\text{SD}(2\mathcal{B})_4(c)$  or  $\text{Q}(2\mathcal{B})_2(p, a, c)$ .

If  $B$  is Morita equivalent to  $\text{SD}(3\mathcal{C})_{2,2}$ , we use again similar arguments as in the proof of [4, Lemma 2.5] to prove Claim 1, where we use as  $\overline{U}'$  and  $\overline{U}$  suitable submodules (resp. suitable quotient modules) of the projective cover  $P_{T_0}$  if the radical quotient (resp. socle) of  $V$  is  $T_0$ , and we use as  $\overline{U}' = \overline{U}$  a suitable quotient module of  $P_{T_1}$  (resp.  $P_{T_2}$ ) if the radical quotient of  $V$  is  $T_1$  (resp.  $T_2$ ). This proves Claim 1.

*Claim 2.* In all cases,  $\text{End}_{kG}(\overline{U}') \cong k[t]/(t^{2^{n-2}-1})$ .

*Proof of Claim 2.* If  $B$  is Morita equivalent neither to  $\text{SD}(2\mathcal{B})_4(c)$ , nor to  $\text{Q}(2\mathcal{B})_2(p, a, c)$ , nor to  $\text{SD}(3\mathcal{C})_{2,2}$ , then  $\overline{U}'$  is isomorphic to a tree module in the sense of [26]. Therefore, we can use the main result of [26] to prove Claim 2.

If  $B$  is Morita equivalent to  $\Lambda \in \{\text{SD}(2\mathcal{B})_4(c), \text{Q}(2\mathcal{B})_2(p, a, c), \text{SD}(3\mathcal{C})_{2,2}\}$ , we use the relations in §4.2 and §4.5 to analyze the possible  $kG$ -module endomorphisms of  $\overline{U}'$ . Using explicit  $k$ -bases for the  $\Lambda$ -module  $\overline{U}'_\Lambda$  corresponding to  $\overline{U}'$  under the Morita equivalence, a straightforward linear algebra calculation shows that  $\text{End}_\Lambda(\overline{U}'_\Lambda) \cong k[t]/(t^{2^{n-2}-1})$ . This proves Claim 2.

*Claim 3.* In all cases,  $\overline{U}'$  has a lift  $U'$  over  $W$  such that the  $F$ -character of  $U'$  is equal to

$$\sum_{\ell=2}^{n-1} \rho_\ell = \sum_{i=1}^{2^{n-2}-1} \chi_{5,i}$$

where  $\rho_\ell$  is as in (5.10).

*Proof of Claim 3.* If  $B$  is not Morita equivalent to any of the algebras in

$$\{\text{SD}(2\mathcal{A})_1(c), \text{Q}(2\mathcal{A})(c), \text{SD}(3\mathcal{A})_1, \text{Q}(3\mathcal{A})_2\}$$

and  $V$  is arbitrary or if  $B$  is Morita equivalent to  $\text{SD}(3\mathcal{A})_1$  and  $V$  is such that its radical quotient or its socle is isomorphic to  $T_1$ , then Claim 3 follows by using the decomposition matrix of  $B$  together with [18, Prop. (23.7)].

Suppose next that  $B$  is Morita equivalent to one of the algebras in  $\{\text{SD}(2\mathcal{A})_1(c), \text{Q}(2\mathcal{A})(c)\}$ . If the radical quotient (resp. socle) of  $V$  is isomorphic to  $T_1$ , then  $\Omega(\overline{U'}) \cong Z_{001}$  (resp.  $\Omega^{-1}(\overline{U'}) \cong Z_{100}$ ) where

$$Z_{001} = \begin{array}{cc} & T_1 \\ T_1 & T_0 \\ & T_0 \\ & T_1 \end{array}, \quad Z_{100} = \begin{array}{cc} & T_1 \\ & T_0 \\ T_1 & T_0 \\ & T_1 \end{array}$$

and  $Z_{001}$  is a submodule of  $P_{T_1}$  and  $Z_{100}$  is a quotient module of  $P_{T_1}$ . Using [3, Lemma 2.3.2], it follows that  $Z_{001}$  and  $Z_{100}$  each have a lift over  $W$  such that the  $F$ -character of this lift is  $\chi_3 + \chi_4 + \chi_6$  (using the notation in Figure 2), which implies Claim 3 in this case.

Finally, suppose that  $B$  is Morita equivalent to  $\text{SD}(3\mathcal{A})_1$  and  $u = 2$  and  $v = 1$ , or  $B$  is Morita equivalent to  $\text{Q}(3\mathcal{A})_2$  and  $\{u, v\} = \{1, 2\}$ . If  $V$  is such that its radical quotient (resp. socle) is isomorphic to  $T_u$  then  $\Omega(\overline{U'}) = Z_{0v0u}$  (resp.  $\Omega^{-1}(\overline{U'}) = Z_{u0v0}$ ) where

$$Z_{0v0u} = \begin{array}{cc} & T_u \\ & T_0 \\ T_u & T_v \\ & T_0 \\ & T_u \end{array}, \quad Z_{u0v0} = \begin{array}{cc} & T_u \\ & T_0 \\ T_u & T_v \\ & T_0 \\ & T_u \end{array}$$

and  $Z_{0v0u}$  is a submodule of  $P_{T_u}$  and  $Z_{u0v0}$  is a quotient module of  $P_{T_u}$ . Using [3, Lemma 2.3.2], it follows that  $Z_{0v0u}$  and  $Z_{u0v0}$  each have a lift over  $W$  such that the  $F$ -character of this lift is  $\chi_2 + \chi_3 + \chi_6$  if  $B$  is Morita equivalent to  $\text{SD}(3\mathcal{A})_1$  (using the notation in Figure 7) and the  $F$ -character of this lift is  $\chi_2 + \chi_4 + \chi_6$  (resp.  $\chi_2 + \chi_3 + \chi_7$ ) if  $B$  is Morita equivalent to  $\text{Q}(3\mathcal{A})_2$  and  $u = 1$  (resp.  $u = 2$ ) (using the notation in Figure 8). This finishes the proof of Claim 3.

*Claim 4.* The universal deformation ring  $R(G, V)$  is as stated in Theorem 6.5.

*Proof of Claim 4.* In all cases, it follows by Lemma 5.6 that  $U'$  from Claim 3 is an  $R'G$ -module. More precisely, there exists a  $WG$ -module endomorphism  $\alpha$  of  $U'$  such that  $W[\alpha] \cong R'$ . By Claim 2, we have  $\text{End}_{kG}(\overline{U'}) \cong k[t]/(t^{2^{n-2}-1}) \cong R'/2R'$ . Moreover, since  $\overline{U'}$  is a lift of  $V$  over  $k[t]/(t^{2^{n-2}-1})$ ,  $\overline{U'}$  is free as a module for  $\text{End}_{kG}(\overline{U'})$  of rank  $\dim_k V = \deg(\chi_{5,1})$ . Hence it follows by Lemma 5.6 that  $\text{End}_{WG}(U') = W[\alpha] \cong R'$  and  $U'$  is free as a module for  $\text{End}_{WG}(U')$ .

In other words,  $U'$  defines a lift of  $V$  over  $R'$ . Let  $\tau : R(G, V) \rightarrow R'$  be the unique continuous  $W$ -algebra homomorphism relative to the lift defined by  $U'$ . Since  $R'/(\mathfrak{m}_{R'}^2 + 2R') \cong k[t]/(t^2)$ ,  $\tau$  is surjective if and only if  $R'/(\mathfrak{m}_{R'}^2 + 2R') \otimes_{R'} U'$  does not define the trivial lift of  $V$  over  $k[t]/(t^2)$ . However,

$$R'/(\mathfrak{m}_{R'}^2 + 2R') \otimes_{R'} U' \cong U'/(\alpha^2(U') + 2U') \cong \overline{U'}/\overline{\alpha}^2(\overline{U'}) \cong \overline{U'}/t^2\overline{U'}$$

is an indecomposable  $kG$ -module, which implies that this does not define trivial lift of  $V$  over  $k[t]/(t^2)$ . Hence  $\tau$  is surjective and induces a surjective  $k$ -algebra homomorphism

$$\overline{\tau} : R(G, V)/2R(G, V) \rightarrow R'/2R'.$$

Suppose first that  $V$  does not correspond to a 3-tube. Then  $R(G, V)/2R(G, V)$  and  $R'/2R'$  are isomorphic and finite dimensional over  $k$ , which implies that  $\overline{\tau}$  is an isomorphism. Because  $R'$  is a free  $W$ -module of finite rank, it follows that  $\tau$  is an isomorphism. By Lemma 5.5,  $R'$  is isomorphic to a subquotient ring of  $WD$ . This proves Claim 4 and completes the proof of Theorem 6.5 if  $V$  does not correspond to a 3-tube.

Suppose next that  $V$  corresponds to a 3-tube. Then the universal mod 2 deformation ring  $R(G, V)/2R(G, V)$  is isomorphic to  $k[t]/(2^{n-2})$  and the universal mod 2 deformation is isomorphic to  $\overline{U}$ . By [3, Lemma 2.3.3], it follows that  $R(G, V) \cong W[[t]]/(p_n(t)(t - 2\mu), a2^m p_n(t))$  for certain  $\mu \in W$ ,  $a \in \{0, 1\}$  and  $0 < m \in \mathbb{Z}$ . If  $a = 0$  then  $R(G, V) \cong W[[t]]/(p_n(t)(t - 2\mu))$  is free over  $W$ . If

$a = 1$  then  $R(G, V)/2^m R(G, V) \cong (W/2^m W)[[t]]/(p_n(t)(t - 2\mu))$  is free over  $W/2^m W$ . Therefore it follows that if  $a = 0$  (resp.  $a = 1$ ), then there is a lift of  $\overline{U}$ , when regarded as a  $kG$ -module, over  $W$  (resp. over  $W/2^m W$ ). But by Proposition 6.4 we have  $R(G, \overline{U}) \cong k$ , which means we must have  $a = 1$  and  $m = 1$ . Since  $V$  corresponds to a 3-tube,  $D$  is dihedral or semidihedral. Hence by Lemma 5.5, the ring  $W[[t]]/(tp_n(t), 2p_n(t))$  is isomorphic to a subquotient ring of  $WD$ . This proves Claim 4 and completes the proof of Theorem 6.5 if  $V$  corresponds to a 3-tube.  $\square$

## REFERENCES

- [1] J. L. Alperin, Local representation theory. Modular representations as an introduction to the local representation theory of finite groups. Cambridge Studies in Advanced Mathematics, vol. 11, Cambridge University Press, Cambridge, 1986.
- [2] F. M. Bleher, Universal deformation rings for dihedral 2-groups. *J. London Math Soc.* 79, (2009), 225–237.
- [3] F. M. Bleher, Universal deformation rings and dihedral defect groups. *Trans. Amer. Math. Soc.* 361 (2009), 3661–3705.
- [4] F. M. Bleher, Universal deformation rings and generalized quaternion defect groups. *Adv. Math.* 225 (2010), 1499–1522.
- [5] F. M. Bleher and T. Chinburg, Universal deformation rings and cyclic blocks. *Math. Ann.* 318 (2000), 805–836.
- [6] F. M. Bleher and T. Chinburg, Universal deformation rings need not be complete intersections. *C. R. Math. Acad. Sci. Paris* 342 (2006), 229–232.
- [7] F. M. Bleher and T. Chinburg, Universal deformation rings need not be complete intersections. *Math. Ann.* 337 (2007), 739–767.
- [8] F. M. Bleher, T. Chinburg and B. de Smit, Inverse Problems for deformation rings. [arXiv:1012.1290](https://arxiv.org/abs/1012.1290)
- [9] F. M. Bleher, G. Lloset and J. B. Schaefer, Universal deformation rings and dihedral blocks with two simple modules. *J. Algebra* 345 (2011), 49–71.
- [10] R. Brauer, On the Connection Between the Ordinary and The Modular Characters of Groups of Finite Order. *Ann. Math.* (2) (1942), 926–935.
- [11] R. Brauer, Zur Darstellungstheorie der Gruppen endlicher Ordnung. II. *Math. Zeitschr.* 72 (1959), 25–46.
- [12] R. Brauer, Some applications of the theory of blocks of characters of finite groups. III. *J. Algebra* 3 (1966) 225–255.
- [13] R. Brauer, On blocks and sections in finite groups. II. *Amer. J. Math.* 90 (1968), 895–925.
- [14] R. Brauer, On 2-blocks with dihedral defect groups. *Symposia Mathematica*, vol. XIII (Convegno di Gruppi e loro Rappresentazioni, INDAM, Rome, 1972), pp. 367–393, Academic Press, London, 1974.
- [15] J. F. Carlson and J. Thévenaz, The classification of endo-trivial modules. *Invent. Math.* 158 (2004), 389–411.
- [16] J. F. Carlson and J. Thévenaz, The classification of torsion endo-trivial modules. *Ann. of Math.* (2) 162 (2005), 823–883.
- [17] T. Chinburg, Can deformation rings of group representations not be local complete intersections? In: *Problems from the Workshop on Automorphisms of Curves*. Edited by Gunther Cornelissen and Frans Oort, with contributions by I. Bouw, T. Chinburg, Cornelissen, C. Gasbarri, D. Glass, C. Lehr, M. Marnett, Oort, R. Pries and S. Wewers. *Rend. Sem. Mat. Univ. Padova* 113 (2005), 129–177.
- [18] C. W. Curtis and I. Reiner, *Methods of representation theory*. Vols. I and II. With applications to finite groups and orders. John Wiley & Sons, Inc., New York, 1981 and 1987.
- [19] B. de Smit and H. W. Lenstra, Explicit construction of universal deformation rings. In: *Modular Forms and Fermat’s Last Theorem* (Boston, 1995), Springer-Verlag, Berlin-Heidelberg-New York, 1997, pp. 313–326.
- [20] F. Eisele,  $p$ -adic lifting problems and derived equivalences. [arXiv:1102.1674v2](https://arxiv.org/abs/1102.1674v2)
- [21] K. Erdmann, Algebras and semidihedral defect groups I. *Proc. London Math. Soc.* (3) 57 (1988), 109–150.
- [22] K. Erdmann, *Blocks of Tame Representation Type and Related Algebras*. Lecture Notes in Mathematics, vol. 1428, Springer-Verlag, Berlin-Heidelberg-New York, 1990.
- [23] P. Fong, A note on splitting fields of representations of finite groups. *Illinois J. Math.* 7 (1963), 515–520.
- [24] T. Holm, Derived equivalence classification of algebras of dihedral, semidihedral, and quaternion type. *J. Algebra* 211 (1999), 159–205.
- [25] B. Huppert, *Endliche Gruppen*. I. Die Grundlehren der Mathematischen Wissenschaften, Band 134, Springer-Verlag, Berlin-New York, 1967.
- [26] H. Krause, Maps between tree and band modules. *J. Algebra* 137 (1991), 186–194.
- [27] M. Linckelmann, A derived equivalence for blocks with dihedral defect groups. *J. Algebra* 164 (1994), 244–255.
- [28] M. Linckelmann, The source algebras of blocks with a Klein four defect group, *J. Algebra* 167 (1994), 821–854.
- [29] M. Linckelmann, Introduction to fusion systems. In: *Group representation theory*, EPFL Press, Lausanne, 2007, pp. 79–113.
- [30] B. Mazur, Deforming Galois representations. In: *Galois groups over  $\mathbb{Q}$*  (Berkeley, 1987), Springer-Verlag, Berlin-Heidelberg-New York, 1989, pp. 385–437.
- [31] J. B. Olsson, On 2-blocks with quaternion and quasidihedral defect groups. *J. Algebra* 36 (1975), 212–241.

Frauke M. Bleher  
 Department of Mathematics  
 University of Iowa  
 Iowa City, IA 52242-1419  
 U.S.A.  
 frauke-bleher@uiowa.edu

FIGURE 1. The decomposition matrix for blocks of type  $D(2\mathcal{A})$  or  $SD(2\mathcal{A})_2(c)$ .

$$\begin{array}{c}
 \chi_1 \\
 \chi_2 \\
 \chi_3 \\
 \chi_4 \\
 \chi_{5,i}
 \end{array}
 \begin{array}{cc}
 \varphi_0 & \varphi_1 \\
 \left[ \begin{array}{cc}
 1 & 0 \\
 1 & 0 \\
 1 & 1 \\
 1 & 1 \\
 2 & 1
 \end{array} \right]
 \end{array}
 \quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 2. The decomposition matrix for blocks of type  $SD(2\mathcal{A})_1(c)$  or  $Q(2\mathcal{A})(c)$ .

$$\begin{array}{c}
 \chi_1 \\
 \chi_2 \\
 \chi_3 \\
 \chi_4 \\
 \chi_{5,i} \\
 \chi_6
 \end{array}
 \begin{array}{cc}
 \varphi_0 & \varphi_1 \\
 \left[ \begin{array}{cc}
 1 & 0 \\
 1 & 0 \\
 1 & 1 \\
 1 & 1 \\
 2 & 1 \\
 0 & 1
 \end{array} \right]
 \end{array}
 \quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 3. The decomposition matrix for blocks of type  $D(2\mathcal{B})$  or  $SD(2\mathcal{B})_1(c)$ .

$$\begin{array}{c}
 \chi_1 \\
 \chi_2 \\
 \chi_3 \\
 \chi_4 \\
 \chi_{5,i}
 \end{array}
 \begin{array}{cc}
 \varphi_0 & \varphi_1 \\
 \left[ \begin{array}{cc}
 1 & 0 \\
 1 & 0 \\
 1 & 1 \\
 1 & 1 \\
 0 & 1
 \end{array} \right]
 \end{array}
 \quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 4. The decomposition matrix for blocks of type  $\text{SD}(2\mathcal{B})_2(c)$  or  $\text{Q}(2\mathcal{B})_1(c)$ .

$$\begin{array}{c}
\varphi_0 \ \varphi_1 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\left[ \begin{array}{cc}
1 & 0 \\
1 & 0 \\
1 & 1 \\
1 & 1 \\
0 & 1 \\
2 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 5. The decomposition matrix for blocks of type  $\text{SD}(2\mathcal{B})_4(c)$  or  $\text{Q}(2\mathcal{B})_2(p, a, c)$ .

$$\begin{array}{c}
\varphi_0 \ \varphi_1 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i}
\end{array}
\begin{array}{c}
\left[ \begin{array}{cc}
1 & 0 \\
1 & 0 \\
0 & 1 \\
0 & 1 \\
1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 6. The decomposition matrix for blocks of type  $\text{D}(3\mathcal{A})_1$ .

$$\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i}
\end{array}
\begin{array}{c}
\left[ \begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 1 \\
1 & 0 & 1 \\
1 & 1 & 0 \\
2 & 1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 7. The decomposition matrix for blocks of type  $\text{SD}(3\mathcal{A})_1$ .

$$\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\left[ \begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 1 \\
1 & 0 & 1 \\
1 & 1 & 0 \\
2 & 1 & 1 \\
0 & 0 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 8. The decomposition matrix for blocks of type  $Q(3\mathcal{A})_2$ .

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6 \\
\chi_7
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[ \begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 1 \\
1 & 0 & 1 \\
1 & 1 & 0 \\
2 & 1 & 1 \\
0 & 1 & 0 \\
0 & 0 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 9. The decomposition matrix for blocks of type  $D(3\mathcal{B})_1$ .

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i}
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[ \begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 0 \\
1 & 0 & 1 \\
1 & 1 & 1 \\
0 & 1 & 0
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 10. The decomposition matrix for blocks of type  $SD(3\mathcal{B})_1$  or  $SD(3\mathcal{D})$ .

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[ \begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 0 \\
1 & 0 & 1 \\
1 & 1 & 1 \\
0 & 1 & 0 \\
0 & 0 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 11. The decomposition matrix for blocks of type  $SD(3\mathcal{B})_2$ .

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[ \begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 0 \\
1 & 1 & 1 \\
1 & 0 & 1 \\
0 & 1 & 0 \\
2 & 1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 12. The decomposition matrix for blocks of type  $Q(3\mathcal{B})$ .

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6 \\
\chi_7
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[ \begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 0 \\
1 & 1 & 1 \\
1 & 0 & 1 \\
0 & 1 & 0 \\
0 & 0 & 1 \\
2 & 1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 13. The decomposition matrix for blocks of type  $SD(3\mathcal{C})_{2,1}$ .

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[ \begin{array}{ccc}
0 & 1 & 0 \\
1 & 1 & 0 \\
1 & 0 & 1 \\
0 & 0 & 1 \\
1 & 0 & 0 \\
1 & 1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 14. The decomposition matrix for blocks of type  $SD(3\mathcal{C})_{2,2}$ .

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[ \begin{array}{ccc}
0 & 1 & 0 \\
1 & 0 & 1 \\
1 & 1 & 0 \\
0 & 0 & 1 \\
1 & 1 & 1 \\
1 & 0 & 0
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 15. The decomposition matrix for blocks of type  $SD(3\mathcal{H})_1$ .

$$\begin{array}{c}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[ \begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 1 \\
0 & 1 & 0 \\
0 & 0 & 1 \\
0 & 1 & 1 \\
1 & 1 & 0
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$



FIGURE 16. The decomposition matrix for blocks of type  $\text{SD}(3\mathcal{H})_2$ .

$$\begin{array}{l}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[ \begin{array}{ccc}
1 & 0 & 0 \\
0 & 1 & 0 \\
1 & 1 & 1 \\
0 & 0 & 1 \\
1 & 1 & 0 \\
0 & 1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 17. The decomposition matrix for blocks of type  $\text{D}(3\mathcal{K})$ .

$$\begin{array}{l}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i}
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[ \begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 1 \\
0 & 1 & 0 \\
0 & 0 & 1 \\
0 & 1 & 1
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$

FIGURE 18. The decomposition matrix for blocks of type  $\text{Q}(3\mathcal{K})$ .

$$\begin{array}{l}
\chi_1 \\
\chi_2 \\
\chi_3 \\
\chi_4 \\
\chi_{5,i} \\
\chi_6 \\
\chi_7
\end{array}
\begin{array}{c}
\varphi_0 \ \varphi_1 \ \varphi_2 \\
\left[ \begin{array}{ccc}
1 & 0 & 0 \\
1 & 1 & 1 \\
0 & 1 & 0 \\
0 & 0 & 1 \\
0 & 1 & 1 \\
1 & 0 & 1 \\
1 & 1 & 0
\end{array} \right]
\end{array}
\quad 1 \leq i \leq 2^{n-2} - 1$$