

A Weiszfeld-like algorithm for a Weber location problem constrained to a closed and convex set[☆]

Germán A. Torres^{a,*}

^a*Facultad de Matemática, Astronomía y Física, Universidad Nacional de Córdoba, CIEM (CONICET), Medina Allende s/n, Ciudad Universitaria (5000) Córdoba, Argentina*

Abstract

The Weber problem consists of finding a point in \mathbb{R}^n that minimizes the weighted sum of distances from m points in \mathbb{R}^n that are not collinear. An application that motivated this problem is the optimal location of facilities in the 2-dimensional case. A classical method to solve the Weber problem, proposed by Weiszfeld in 1937, is based on a fixed point iteration.

In this work a Weber problem constrained to a closed and convex set is considered. A Weiszfeld-like algorithm, well defined even when an iterate is a vertex, is presented. The iteration function Q that defines the proposed algorithm, is based mainly on an orthogonal projection over the feasible set, combined with the iteration function of a modified Weiszfeld algorithm presented by Vardi and Zhang in 2001.

It can be seen that the proposed algorithm generates a sequence of feasible iterates that have descent properties. Under certain hypotheses, the limit of this sequence satisfies the KKT optimality conditions, is a fixed point of the iteration function that defines the algorithm, and is the solution of the constrained minimization problem. Numerical experiments confirmed the theoretical results.

Keywords: location, Weber problem, Weiszfeld algorithm, fixed point iteration
2000 MSC: 90B85, 90C25, 90C30

1. Introduction

Let a^1, \dots, a^m be m distinct points in the space \mathbb{R}^n , called vertices, and positive numbers w_1, \dots, w_m , called weights. The function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ defined by

$$f(x) = \sum_{j=1}^m w_j \|x - a^j\|, \quad (1)$$

4 is called the Weber function, where $\|\cdot\|$ denotes the Euclidean norm. It is well-known that this function
5 is not differentiable at the vertices, and strictly convex if the vertices are not collinear (we will assume
6 this hypothesis from now on).

7 The Weber problem (also known as the Fermat-Weber problem) is to find a point in \mathbb{R}^n that min-
8 imizes the weighted sum of Euclidean distances from the m given points, that is, we have to find the
9 solution of the following unconstrained optimization problem:

$$\begin{aligned} & \underset{x}{\operatorname{argmin}} && f(x) \\ & \text{subject to} && x \in \mathbb{R}^n. \end{aligned} \quad (2)$$

[☆]Research partially supported by the CONICET, the SECYT-UNC and the ANPCyT.

*Corresponding author

Email address: torres@famaf.unc.edu.ar (Germán A. Torres)

10 This problem has a unique solution x^u in \mathbb{R}^n .

11 The problem was also stated as a pure mathematical problem by Fermat [44, 27], Cavalieri [37],
 12 Steiner [14], Fasbender [20] and many others. Several solutions, based on geometrical arguments, were
 13 proposed by Torricelli and Simpson. In [30] historical details and geometric aspects were presented by
 14 Kupitz and Martini. In [41] Weber formulated the problem (2) from an economical point of view. The
 15 vertices represent customers or demands, the solution to the problem denotes the location of a new
 16 facility, and the weights are costs associated with the interactions between the new facility and the
 17 customers.

18 Among several schemes to solve the Weber location problem (see [12, 19, 28, 34]), one of the most
 19 popular methods was presented by Weiszfeld in [42, 43]. The Weiszfeld algorithm is an iterative method
 20 based on the first-order necessary conditions for a stationary point of the objective function.

21 If we define $T_0 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by:

$$T_0(x) = \begin{cases} \frac{\sum_{j=1}^m \frac{w_j a^j}{\|x - a^j\|}}{\sum_{j=1}^m \frac{w_j}{\|x - a^j\|}}, & \text{if } x \neq a^1, \dots, a^m, \\ a^k, & \text{if } x = a^k, k = 1, \dots, m, \end{cases} \quad (3)$$

22 the Weiszfeld algorithm is:

$$x^{(l)} = T_0(x^{(l-1)}), \quad l \in \mathbb{N}, \quad (4)$$

23 where $x^{(0)} \in \mathbb{R}^n$ is a starting point.

24 The Weiszfeld algorithm (4), despite of its simplicity, has a serious problem if some $x^{(l)}$ lands ac-
 25 cidentally in a vertex a^k , because the algorithm gets stuck at a^k , even when a^k is not the solution of
 26 (2). Many authors studied the set of initial points for which the sequence generated by the Weiszfeld
 27 algorithm yields in a vertex (see [29, 11, 6, 9, 7, 3]). Vardi and Zhang [40] derived a simple but nontrivial
 28 modification of the Weiszfeld algorithm in which they solved the problem of landing in a vertex.

29 Generalizations and new techniques for the Fermat-Weber location problem have been developed in
 30 recent years. In [18] Eckhardt applied the Weiszfeld algorithm to generalized Weber problems in Banach
 31 spaces. An exact algorithm for a Weber problem with attraction and repulsion was presented by Chen et
 32 al. in [13]. Kaplan and Yang [24] proved a duality theorem which includes as special cases a great variety
 33 of choices of norms in the terms of the Fermat-Weber sum. In [10] Carrizosa et al. studied the so called
 34 Regional Weber Problem, which allows the demand not to be concentrated onto a finite set of points,
 35 but follows an arbitrary probability measure. In [17] Drezner and Wesolowsky studied the case where
 36 different l_p norms are used for each demand point. In [23] the so called Complementary Problem (the
 37 Weber problem with one negative weight) was studied by Jalal and Krarup, and geometrical solutions
 38 were given. In [15] Drezner presented a Weiszfeld-like iterative procedure and convergence is proved if
 39 appropriate conditions hold.

40 In some practical problems it is necessary to consider barriers (forbidden regions). Barriers were first
 41 introduced to location modeling by Katz and Cooper [25]. There exist several heuristic and iterative
 42 algorithms for single-facility location problems for distance computations in the presence of barriers (see
 43 [2, 8, 5, 4]). In [35] Pfeiffer and Klamroth presented a unified formulation for problems with barriers
 44 and network location problems. A complete reference to barriers in location problems can be found in
 45 [26]. Barriers can be applied to model real life problems where regions like lakes and mountains are
 46 forbidden.

47 On the other hand, there are location problems whose solution needs to lie within a closed set.
 48 For example, see [39] for a discussion of the case when the solution is constrained to be within a

49 maximum distance of each demand point. Drezner and Wesolowsky [16] studied the problem of locating
50 an obnoxious facility with rectangular distances (l_1 norm), where the facility must lie within some
51 prespecified region (linear constraints). A primal-dual algorithm to deal with the constrained Fermat-
52 Weber problem using mixed norms was developed in [33] by Idrissi et al.. In [21] Hansen et al. presented
53 an algorithm for solving the Weber problem when the set of feasible locations is the union of a finite
54 number of convex polygons. In [36] Pilotta and Torres considered a Weber location problem with box
55 constraints.

56 Constrained Weber problems arise when we require that the solution is in an area (feasible region)
57 determined by, for example, environmental and/or political reasons. It could be the case for a facil-
58 ity producing dangerous materials that must be installed in a restricted (constrained) area. Another
59 example could be the location of a plant in an industrial zone or of a hospital in a non-polluted area.

60 In this paper a constrained location problem is considered. An algorithm is proposed to solve the
61 following problem:

$$\begin{aligned} & \underset{x}{\operatorname{argmin}} && f(x) \\ & \text{subject to} && x \in \Omega, \end{aligned} \tag{5}$$

62 where Ω is a closed and convex set, generalizing the problem formulated in [36]. Problem (5) could be
63 seen as a nonlinear programming problem and solved by standard solvers, but they may fail since the
64 Weber function is not differentiable at the vertices.

65 It can be proved that problem (5) has a unique solution x^* , since the function f is strictly convex
66 and Ω is a closed and convex set. On the other hand, it is well-known that the convex hull of the given
67 vertices a^1, \dots, a^m contains the solution x^u of the unconstrained Weber problem (see for instance [29,
68 pp. 100]). If Ω contains the convex hull, both solutions x^* and x^u agree. In other cases, the solution
69 x^* is not necessarily a projection of x^u over Ω (see [36]). The algorithm is based basically on a slight
70 variation of an orthogonal projection of the Weiszfeld algorithm presented in [40], that is well defined
71 even when an iterate coincides with a vertex. Properties of the sequence generated by the proposed
72 algorithm related with the minimization problem 5 will be proved in the following sections.

73 The paper is structured as follows: Section 2 describes the results in [40] in which a modified
74 Weiszfeld algorithm is presented and some notation is introduced. In Section 3 the proposed algorithm
75 is defined. Section 4 is dedicated to definitions and technical lemmas. In Section 5 the main results
76 about convergence to optimality are presented. Numerical experiments are considered in Section 6.
77 Finally, conclusions are given in Section 7.

78 Some words about notation. As it was mentioned, we will call x^u the solution of problem (2) and
79 x^* the solution of problem (5). The symbols $\|\cdot\|$ and $\langle \cdot, \cdot \rangle$ will refer to the standard Euclidean norm
80 and standard inner product in \mathbb{R}^n , respectively. For a function $f : \mathbb{R} \rightarrow \mathbb{R}$ we will denote by $f'(a-)$ the
81 left-hand side derivative at a , and by $f'(a+)$ the right-hand side derivative at a .

82 2. The modified Weiszfeld algorithm

83 This section reviews the main results presented in [40] in which the authors generalize the Weiszfeld
84 algorithm for the case that an iterate lands on a vertex. From now on, this algorithm will be referred
85 to as the modified Weiszfeld algorithm.

86 In order to make notation easier, we define the function $A : \mathbb{R}^n \rightarrow \mathbb{R}$ by:

$$A(x) = \begin{cases} \sum_{j=1}^m \frac{w_j}{2\|x - a^j\|}, & \text{if } x \neq a^1, \dots, a^m, \\ \sum_{\substack{j=1 \\ j \neq k}}^m \frac{w_j}{2\|a^k - a^j\|}, & \text{if } x = a^k, k = 1, \dots, m. \end{cases} \tag{6}$$

87 Notice that $A(x) > 0$ for all $x \in \mathbb{R}^n$. In [40, pp. 563], the number $A(a^k)$ was called A_k .

88 A generalization for the iteration function T_0 , defined in (3), is given by $\tilde{T} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined as
 89 follows:

$$\tilde{T}(x) = \begin{cases} \frac{\sum_{j=1}^m \frac{w_j a^j}{\|x - a^j\|}}{2A(x)}, & \text{if } x \neq a^1, \dots, a^m, \\ \frac{\sum_{\substack{j=1 \\ j \neq k}}^m \frac{w_j a^j}{\|a^k - a^j\|}}{2A(a^k)}, & \text{if } x = a^k, k = 1, \dots, m. \end{cases} \quad (7)$$

90 Notice that \tilde{T} coincides with T_0 in $\mathbb{R}^n - \{a^1, \dots, a^m\}$.

91 Let $\tilde{R} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $r : \mathbb{R}^n \rightarrow \mathbb{R}$ be:

$$\tilde{R}(x) = \begin{cases} \sum_{j=1}^m \frac{w_j (a^j - x)}{\|x - a^j\|}, & \text{if } x \neq a^1, \dots, a^m, \\ \sum_{\substack{j=1 \\ j \neq k}}^m \frac{w_j (a^j - a^k)}{\|a^k - a^j\|}, & \text{if } x = a^k, k = 1, \dots, m, \end{cases} \quad (8)$$

$$r(x) = \|\tilde{R}(x)\|, \quad \forall x \in \mathbb{R}^n.$$

92 The function \tilde{R} generalizes the negative gradient of the Weber function since, for all $x \neq a^1, \dots, a^m$,

$$\nabla f(x) = -\tilde{R}(x). \quad (9)$$

93 The following lemma is very easy to prove (see [40, equation (14)]), and it relates the functionals \tilde{T}
 94 and \tilde{R} .

95 **Lemma 1.** *For all $x \in \mathbb{R}^n$ we have $\tilde{R}(x) = 2A(x) [\tilde{T}(x) - x]$.*

If we define $\gamma : \mathbb{R}^n \rightarrow \mathbb{R}$ by:

$$\gamma(x) = \begin{cases} 0, & \text{if } x \neq a^1, \dots, a^m, \\ 0, & \text{if } x = a^k \text{ and } r(a^k) = 0 \text{ for some } k = 1, \dots, m, \\ w_k/r(a^k), & \text{if } x = a^k \text{ and } r(a^k) \neq 0 \text{ for some } k = 1, \dots, m, \end{cases}$$

96 we can see that $\gamma(x) \geq 0$ for all $x \in \mathbb{R}^n$.

The modified Weiszfeld algorithm presented in [40] is defined by:

$$x^{(l)} = T(x^{(l-1)}), \quad l \in \mathbb{N},$$

97 where $x^{(0)} \in \mathbb{R}^n$ and $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is given by:

$$T(x) = (1 - \beta(x))\tilde{T}(x) + \beta(x)x, \quad (10)$$

98 where $\beta : \mathbb{R}^n \rightarrow \mathbb{R}$ is defined by $\beta(x) = \min\{1, \gamma(x)\}$.

99 **Remark 2.** (a) *If $x \neq a^1, \dots, a^m$, then $\beta(x) = 0$ because $\gamma(x) = 0$. So, we can deduce that $T(x) =$
 100 $\tilde{T}(x)$. Notice that this fact implies that the functional T is continuous in $\mathbb{R}^n - \{a^1, \dots, a^m\}$.*

- 101 (b) It can be seen that if $a^k \neq x^u$, then $0 < \beta(a^k) < 1$ (see [40, pp. 563]).
 102 (c) From equation (10) we obtain that $T(x) - x = (1 - \beta(x))(\tilde{T}(x) - x)$ for $x \in \mathbb{R}^n$.

103 The main result in [40, pp. 562] is:

104 **Theorem 3.** *The following propositions are equivalent:*

- 105 (a) $x = x^u$.
 106 (b) $T(x) = x$.
 107 (c) $r(x) \leq \eta(x)$.

where

$$\eta(x) = \begin{cases} 0, & \text{if } x \neq a^1, \dots, a^m, \\ w_k, & \text{if } x = a^k, k = 1, \dots, m. \end{cases}$$

108 3. The proposed algorithm

109 This section is dedicated to describe the proposed algorithm, introducing some definitions and re-
 110 marks.

111 First of all, we can notice that problem (5) has a unique solution, due to the fact that f is a
 112 non-negative, strictly convex, and continuous function, $\lim_{\|x\| \rightarrow \infty} f(x) = \infty$ and Ω is closed and convex.

In order to define the proposed algorithm at the vertices, we will need to determine which points of the segment that joins a^k and $T(a^k)$ are in the feasible set Ω . If $k = 1, \dots, m$, let the set \mathcal{S}_k be defined by:

$$\mathcal{S}_k = \{ \lambda \in [0, 1] : (1 - \lambda)T(a^k) + \lambda a^k \in \Omega \}.$$

Notice that \mathcal{S}_k could be equal to the empty set in case that a^k and $T(a^k)$ do not belong to Ω . On the other hand, if $a^k \in \Omega$, then $1 \in \mathcal{S}_k$, which means that $\mathcal{S}_k \neq \emptyset$. Thus, we can define:

$$\lambda(a^k) = \inf \mathcal{S}_k, \quad a^k \in \Omega.$$

113 In case a vertex a^k is not in Ω , there is no need to define the number $\lambda(a^k)$.

114 In the following lemma, a set of basic properties of $\lambda(a^k)$ are listed:

115 **Lemma 4.** *If $k = 1, \dots, m$ and $a^k \in \Omega$ then:*

- 116 (a) $\lambda(a^k) \in [0, 1]$.
 117 (b) If $T(a^k) \in \Omega$ then $\lambda(a^k) = 0$.
 118 (c) If $T(a^k) \notin \Omega$ then $\lambda(a^k) \in (0, 1]$.

119 **PROOF.** The proof of (a) follows from the definition of \mathcal{S}_k . If $T(a^k) \in \Omega$, then $0 \in \mathcal{S}_k$, so $\lambda(a^k) = 0$,
 120 and this proves (b). Finally, for item (c), let us consider that $T(a^k) \notin \Omega$. Since Ω is a closed set, there
 121 is an entire ball centered at $T(a^k)$ that does not intersect Ω , which implies that there exists ϵ such that
 122 $(1 - \lambda)T(a^k) + \lambda a^k \notin \Omega$ for all $\lambda \in [0, \epsilon]$. Thus, $\lambda(a^k) \in (0, 1]$ and this concludes the proof. \square

123 Let us call $P_\Omega : \mathbb{R}^n \rightarrow \Omega$ the orthogonal projection over Ω . Since Ω is a nonempty, closed and convex
 124 set, the operator P_Ω is a continuous function [1, pp. 99].

125 We define the iteration function $Q : \Omega \rightarrow \Omega$ by:

$$Q(x) = \begin{cases} P_\Omega \circ T(x), & \text{if } x \neq a^1, \dots, a^m, \\ (1 - \lambda(a^k))T(a^k) + \lambda(a^k)a^k, & \text{if } x = a^k \in \Omega, k = 1, \dots, m. \end{cases} \quad (11)$$

126 There will be no need to define Q outside Ω since the proposed algorithm generates a sequence of feasible
 127 points. The iteration function Q at $x \in \Omega$ coincides with the orthogonal projection of $T(x)$ over the
 128 feasible set when x is different from the vertices. Only when x is a vertex a^k belonging to Ω , $Q(x)$ is
 129 defined as the farthest possible feasible point of the segment that joins x with $T(x)$.

130 The following remark states some basic properties of the iteration function of the proposed algorithm.
 131

132 **Remark 5.**

133 (a) If $a^k \in \Omega$ and $T(a^k) \in \Omega$, then $Q(a^k) = T(a^k) = P_\Omega \circ T(a^k)$.

134 (b) If $a^k \in \Omega$, it can be seen that:

$$\begin{aligned} Q(a^k) - a^k &= (1 - \lambda(a^k)) (T(a^k) - a^k), \\ Q(a^k) - T(a^k) &= -\lambda(a^k) (T(a^k) - a^k). \end{aligned}$$

135 (c) The functional Q is continuous in $\mathbb{R}^n - \{a^1, \dots, a^m\}$.

136 **PROOF.** The proofs of (a) and (b) are straightforward. For (c), since P_Ω is continuous in \mathbb{R}^n (see [1,
 137 pp. 99]) and T is continuous in $\mathbb{R}^n - \{a^1, \dots, a^m\}$ (see Remark 2), we have that Q is continuous in
 138 $\mathbb{R}^n - \{a^1, \dots, a^m\}$. \square

139 The proposed algorithm is described below.

140 **Algorithm 6.** Let $\Omega \subset \mathbb{R}^n$ be a closed and convex set. Assume that $x^{(0)} \in \Omega$ is an initial approximation
 141 such that $f(x^{(0)}) \leq f(a^j)$ for all $j \in \{1, \dots, m\}$ and $a^j \in \Omega$. Given $\varepsilon > 0$ a tolerance and $x^{(l-1)} \in \Omega$, do
 142 the following steps to compute $x^{(l)}$:

143 **Step 1:** Compute:

$$x^{(l)} = Q(x^{(l-1)}). \quad (12)$$

Step 2: Stop the execution if

$$\|x^{(l)} - x^{(l-1)}\| < \varepsilon,$$

145 and declare $x^{(l)}$ as solution to the problem (5). Otherwise return to Step 1.

146 From the definition of Q it follows that Algorithm 6 generates a sequence of feasible iterates. Also
 147 notice that if there are vertices in the feasible set, $x^{(0)}$ can be one of them, for example, a vertex a_s such
 148 that $f(a_s) \leq f(a^j)$ for all $a^j \in \Omega$. On the other hand, if there are no vertices in the feasible set, $x^{(0)}$ can
 149 be chosen as the projection over Ω of the null vector.

150 4. Some definitions and technical results

151 The purpose of this section is to define some entities and prove technical lemmas that will be
 152 important in the proof of the main results.

First of all, we will define some useful operators for making notation easier. If $\mathcal{A} \subset \{1, \dots, n\}$, then
 we define $\|\cdot\|_{\mathcal{A}} : \mathbb{R}^n \rightarrow \mathbb{R}$ and $\langle \cdot, \cdot \rangle_{\mathcal{A}} : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ by:

$$\|x\|_{\mathcal{A}} = \sqrt{\sum_{j \in \mathcal{A}} x_j^2}, \quad \langle x, y \rangle_{\mathcal{A}} = \sum_{j \in \mathcal{A}} x_j y_j.$$

153 Notice that, when $\mathcal{A} \subsetneq \{1, \dots, n\}$, $\|\cdot\|_{\mathcal{A}}$ is not necessarily a norm and $\langle \cdot, \cdot \rangle_{\mathcal{A}}$ is not necessarily an inner
 154 product.

155 According to this definition, if \mathcal{A} and \mathcal{B} are sets such that $\mathcal{A} \cap \mathcal{B} = \emptyset$ and $\mathcal{A} \cup \mathcal{B} = \{1, \dots, n\}$, it can
 156 be seen that:

$$\|x\|^2 = \|x\|_{\mathcal{A}}^2 + \|x\|_{\mathcal{B}}^2, \quad (13)$$

$$\langle x, y \rangle = \langle x, y \rangle_{\mathcal{A}} + \langle x, y \rangle_{\mathcal{B}}, \quad (14)$$

$$c\langle x, y \rangle_{\mathcal{A}} = \langle cx, y \rangle_{\mathcal{A}} = \langle x, cy \rangle_{\mathcal{A}}. \quad (15)$$

157 For $x \in \Omega$, let us define the following sets of indices:

$$\mathcal{N}(x) = \{k \in \mathbb{N} : 1 \leq k \leq n, (T(x))_k \neq (Q(x))_k\},$$

$$\mathcal{E}(x) = \{k \in \mathbb{N} : 1 \leq k \leq n, (T(x))_k = (Q(x))_k\},$$

158 Notice that for all $x \in \mathbb{R}^n$ we have that $\mathcal{N}(x) \cap \mathcal{E}(x) = \emptyset$ and $\mathcal{N}(x) \cup \mathcal{E}(x) = \{1, \dots, n\}$.

159 Let $\alpha : \Omega \rightarrow \mathbb{R}^n$ be the following function:

160 • If $x \neq a^1, \dots, a^m$:

$$\alpha(x) = \sum_{j=1}^m \frac{w_j}{\|x - a^j\|} [Q(x) - a^j]. \quad (16)$$

161 • If $x = a^k \in \Omega$ for some $k = 1, \dots, m$:

$$\alpha(x) = \sum_{\substack{j=1 \\ j \neq k}}^m \frac{w_j}{\|a^k - a^j\|} [Q(a^k) - (1 - \beta(a^k))a^j - \beta(a^k)a^k]. \quad (17)$$

162 It can be seen that the function α is related to the iteration function Q of the proposed algorithm,
 163 and the iteration function T of the modified algorithm.

164 **Lemma 7.** *If $x \in \Omega$, then $\alpha(x) = 2A(x)[Q(x) - T(x)]$.*

165 **PROOF.** If $x \neq a^1, \dots, a^m$, then:

$$\begin{aligned} \alpha(x) &= \sum_{j=1}^m \frac{w_j}{\|x - a^j\|} [Q(x) - a^j] = \sum_{j=1}^m \frac{w_j Q(x)}{\|x - a^j\|} - \sum_{j=1}^m \frac{w_j a^j}{\|x - a^j\|} \\ &= \left(\sum_{j=1}^m \frac{w_j}{\|x - a^j\|} \right) \left[Q(x) - \frac{\sum_{j=1}^m \frac{w_j a^j}{\|x - a^j\|}}{\sum_{j=1}^m \frac{w_j}{\|x - a^j\|}} \right] = 2A(x) [Q(x) - \tilde{T}(x)] \\ &= 2A(x) [Q(x) - T(x)]. \end{aligned}$$

166 where in the last equalities we have used the definition of \tilde{T} as in (7), and the fact that $\tilde{T}(x) = T(x)$
 167 due to Remark 2.

168 If $x = a^k$ for some $k = 1, \dots, m$, we follow a similar procedure than in the previous case. \square

169 Now, we will define auxiliary functions that take into account the projection P_{Ω} in order to prove a
 170 descent property of f (see next sections). If $x \in \Omega$, we define:

171 (a) $E_x : \mathbb{R}^n \rightarrow \mathbb{R}^n$, where:

$$(E_x(y))_k = \begin{cases} (Q(x))_k, & \text{if } k \in \mathcal{N}(x), \\ y_k, & \text{if } k \in \mathcal{E}(x). \end{cases} \quad (18)$$

(b) If $\mathcal{E}(x) = \{i_1, \dots, i_r\} \neq \emptyset$ define $P_x : \mathbb{R}^n \rightarrow \mathbb{R}^r$ where:

$$(P_x(y))_k = y_{i_k}, \quad k = 1, \dots, r.$$

172 A useful property of E_x , that follows from the definition, is pointed out in the following remark.

173 **Remark 8.** *If $x \in \Omega$ then $E_x \circ Q(x) = Q(x)$.*

174 The iteration function Q inherits an important property from the orthogonal projection P_Ω .

175 **Lemma 9.** *If $x \in \Omega$ we have that $\langle Q(x) - x, Q(x) - T(x) \rangle \leq 0$.*

176 **PROOF.** If $x \neq a^1, \dots, a^m$, then $Q(x) = P_\Omega \circ T(x)$. By a property of the orthogonal projection [1, pp.
177 93] we have that $\langle Q(x) - x, Q(x) - T(x) \rangle \leq 0$.

If $x = a^k$ for some $k = 1, \dots, m$, Remark 5 and Lemma 4 imply:

$$\langle Q(a^k) - a^k, Q(a^k) - T(a^k) \rangle = -\lambda(a^k) (1 - \lambda(a^k)) \|T(a^k) - a^k\|^2 \leq 0,$$

178 and this concludes the proof. □

179 The next technical lemma will help us to save computations in other lemmas.

Lemma 10. *If $x \in \Omega$, \mathcal{A} is a subset of $\{1, \dots, n\}$ and $j \in \{1, \dots, m\}$, then:*

$$\|Q(x) - a^j\|_{\mathcal{A}}^2 = \|x - a^j\|_{\mathcal{A}}^2 - \|Q(x) - x\|_{\mathcal{A}}^2 + 2\langle Q(x) - x, Q(x) - a^j \rangle_{\mathcal{A}}.$$

180 **PROOF.** If $x \in \Omega$, we have:

$$\begin{aligned} \|Q(x) - a^j\|_{\mathcal{A}}^2 &= \langle Q(x) - a^j, Q(x) - a^j \rangle_{\mathcal{A}} \\ &= \langle Q(x) - x + x - a^j, Q(x) - x + x - a^j \rangle_{\mathcal{A}} \\ &= \|Q(x) - x\|_{\mathcal{A}}^2 + \|x - a^j\|_{\mathcal{A}}^2 + 2\langle Q(x) - x, x - a^j \rangle_{\mathcal{A}} \\ &= \|Q(x) - x\|_{\mathcal{A}}^2 + \|x - a^j\|_{\mathcal{A}}^2 + 2\langle Q(x) - x, x - Q(x) \rangle_{\mathcal{A}} \\ &\quad + 2\langle Q(x) - x, Q(x) - a^j \rangle_{\mathcal{A}} \\ &= \|x - a^j\|_{\mathcal{A}}^2 - \|Q(x) - x\|_{\mathcal{A}}^2 + 2\langle Q(x) - x, Q(x) - a^j \rangle_{\mathcal{A}}. \end{aligned}$$

181 □

182 If $x \in \Omega$, let us define $g_x : \mathbb{R}^n \rightarrow \mathbb{R}$ by:

$$g_x(y) = \begin{cases} \sum_{j=1}^m \frac{w_j}{2\|x - a^j\|} \|E_x(y) - a^j\|^2, & \text{if } x \neq a^1, \dots, a^m, \\ \sum_{\substack{j=1 \\ j \neq k}}^m \frac{w_j}{2\|a^k - a^j\|} \|y - a^j\|^2 + w_k \|y - a^k\|, & \text{if } x = a^k, \\ & k = 1, \dots, m. \end{cases} \quad (19)$$

183 The values that g_x assumes at x and $Q(x)$ will play an important role in the proof of a property of the
184 objective function f .

185 **Lemma 11.** *Let $x \in \Omega$ be.*

186 (a) If $x \neq a^1, \dots, a^m$ then:

$$\begin{aligned} g_x(x) &= \frac{1}{2}f(x) + 2A(x) \langle Q(x) - x, Q(x) - T(x) \rangle \\ &\quad - A(x) \|Q(x) - x\|_{\mathcal{N}(x)}^2. \end{aligned}$$

187 (b) If $x = a^k$ for some $k = 1, \dots, m$, then $g_{a^k}(a^k) = \frac{1}{2}f(a^k)$.

PROOF. Let us suppose that $x \neq a^1, \dots, a^m$. By property (13) and (18), we have for $j = 1, \dots, m$:

$$\|E_x(x) - a^j\|^2 = \|x - a^j\|_{\mathcal{E}(x)}^2 + \|Q(x) - a^j\|_{\mathcal{N}(x)}^2.$$

188 Using Lemma 10, we can see that:

$$\begin{aligned} g_x(x) &= \sum_{j=1}^m \frac{w_j}{2\|x - a^j\|} \left[\|x - a^j\|_{\mathcal{E}(x)}^2 + \|x - a^j\|_{\mathcal{N}(x)}^2 \right. \\ &\quad \left. - \|Q(x) - x\|_{\mathcal{N}(x)}^2 + 2 \langle Q(x) - x, Q(x) - a^j \rangle_{\mathcal{N}(x)} \right]. \end{aligned}$$

Due to (13), the definition of the Weber function f , the definition of A as in (6), the property (15) and the definition of α as in (16), we obtain:

$$g_x(x) = \frac{1}{2}f(x) - A(x)\|Q(x) - x\|_{\mathcal{N}(x)}^2 + \langle Q(x) - x, \alpha(x) \rangle_{\mathcal{N}(x)}.$$

189 By Lemma 7, the fact that $(Q(x))_i = (T(x))_i$ for all $i \in \mathcal{E}(x)$ and (14), we get:

$$\begin{aligned} g_x(x) &= \frac{1}{2}f(x) - A(x)\|Q(x) - x\|_{\mathcal{N}(x)}^2 \\ &\quad + 2A(x)\langle Q(x) - x, Q(x) - T(x) \rangle, \end{aligned}$$

190 which concludes the proof of (a).

Now, let us assume that $x = a^k$ for some $k = 1, \dots, m$. Then:

$$g_{a^k}(a^k) = \sum_{\substack{j=1 \\ j \neq k}}^m \frac{w_j}{2\|a^k - a^j\|} \|a^k - a^j\|^2 = \frac{1}{2} \sum_{\substack{j=1 \\ j \neq k}}^m w_j \|a^k - a^j\| = \frac{1}{2}f(a^k).$$

191 This concludes the proof of (b). □

192 The number $g_x(Q(x))$ can be computed in the next lemma.

193 **Lemma 12.** *Let $x \in \Omega$ be.*

194 (a) If $x \neq a^1, \dots, a^m$ then:

$$\begin{aligned} g_x(Q(x)) &= \frac{1}{2}f(x) + 2A(x) \langle Q(x) - x, Q(x) - T(x) \rangle \\ &\quad - A(x) \|Q(x) - x\|^2. \end{aligned}$$

195 (b) If $x = a^k$ for some $k = 1, \dots, m$, then

$$\begin{aligned} g_{a^k}(Q(a^k)) &= \frac{1}{2}f(a^k) - A(a^k) \|Q(a^k) - a^k\|^2 \\ &\quad + 2A(a^k) \langle Q(a^k) - a^k, Q(a^k) - T(a^k) \rangle \\ &\quad - 2\beta(a^k)A(a^k) \langle Q(a^k) - a^k, \tilde{T}(a^k) - a^k \rangle \\ &\quad + w_k \|Q(a^k) - a^k\|. \end{aligned}$$

PROOF. First, let us consider $x \neq a^1, \dots, a^m$. Due to Remark 8 we have:

$$g_x(Q(x)) = \sum_{j=1}^m \frac{w_j}{2\|x - a^j\|} \|Q(x) - a^j\|^2.$$

196 By Lemma 10 we obtain:

$$\begin{aligned} g_x(Q(x)) &= \sum_{j=1}^m \frac{w_j}{2\|x - a^j\|} [\|x - a^j\|^2 - \|Q(x) - x\|^2 \\ &\quad + 2\langle Q(x) - x, Q(x) - a^j \rangle]. \end{aligned}$$

Due to the definition of the Weber function f , the definition of A as in (6) and the definition of α as in (16), we deduce that:

$$g_x(Q(x)) = \frac{1}{2}f(x) - A(x)\|Q(x) - x\|^2 + \langle Q(x) - x, \alpha(x) \rangle.$$

By Lemma 7 we get:

$$g_{a^k}(Q(a^k)) = \frac{1}{2}f(a^k) - A(a^k)\|Q(a^k) - a^k\|^2 + 2A(a^k)\langle Q(a^k) - a^k, Q(a^k) - T(a^k) \rangle,$$

197 concluding the proof of (a).

Now, consider $x = a^k$ for some $k = 1, \dots, m$. Due to (19) we have:

$$g_{a^k}(Q(a^k)) = \sum_{\substack{j=1 \\ j \neq k}}^m \frac{w_j}{2\|a^k - a^j\|} \|Q(a^k) - a^j\|^2 + w_k \|Q(a^k) - a^k\|.$$

198 By Lemma 10, the definition of the Weber function f and the definition of A as in (6) we obtain:

$$\begin{aligned} g_{a^k}(Q(a^k)) &= \frac{1}{2}f(a^k) - A(a^k)\|Q(a^k) - a^k\|^2 \\ &\quad + \left\langle Q(a^k) - a^k, \sum_{\substack{j=1 \\ j \neq k}}^m \frac{w_j}{\|a^k - a^j\|} [Q(a^k) - a^j] \right\rangle \\ &\quad + w_k \|Q(a^k) - a^k\|. \end{aligned}$$

Manipulating algebraically,

$$Q(a^k) - a^j = Q(a^k) - (1 - \beta(a^k))a^j - \beta(a^k)a^k + \beta(a^k)(a^k - a^j).$$

199 Due to the definition of α (see (17)) and the definition of \tilde{R} (see (8)) we get:

$$\begin{aligned} g_{a^k}(Q(a^k)) &= \frac{1}{2}f(a^k) - A(a^k)\|Q(a^k) - a^k\|^2 \\ &\quad + \langle Q(a^k) - a^k, \alpha(a^k) \rangle - \beta(a^k) \langle Q(a^k) - a^k, \tilde{R}(a^k) \rangle \\ &\quad + w_k \|Q(a^k) - a^k\|. \end{aligned}$$

200 By Lemma 1 and Lemma 7 we have:

$$\begin{aligned} g_{a^k}(Q(a^k)) &= \frac{1}{2}f(a^k) - A(a^k)\|Q(a^k) - a^k\|^2 \\ &\quad + 2A(a^k) \langle Q(a^k) - a^k, Q(a^k) - T(a^k) \rangle \\ &\quad - 2A(a^k)\beta(a^k) \langle Q(a^k) - a^k, \tilde{T}(a^k) - a^k \rangle + w_k \|Q(a^k) - a^k\|. \end{aligned}$$

201 which concludes the proof. □

202 The next lemma deals with the last two terms of $g_{a^k}(Q(a^k))$.

Lemma 13. *If $a^k \in \Omega$ for some $k = 1, \dots, m$, the number*

$$z = w_k \left\| Q(a^k) - a^k \right\| - 2A(a^k)\beta(a^k) \left\langle Q(a^k) - a^k, \tilde{T}(a^k) - a^k \right\rangle,$$

203 *is equal to zero.*

204 **PROOF.** If $Q(a^k) = a^k$ the result is true. So, from now on, let us consider that $Q(a^k) \neq a^k$. First, let us
 205 check that $a^k \neq x^u$. In case that $a^k = x^u$, then $T(a^k) = a^k$ by Theorem 3. Since $a^k \in \Omega$, then $T(a^k) \in \Omega$.
 206 By Remark 5 we have that $Q(a^k) = T(a^k) = a^k$ which is a contradiction.

By Remark 2, we have that $\beta(a^k) \in (0, 1)$ (since $a^k \neq x^u$) and:

$$z = w_k \left\| Q(a^k) - a^k \right\| - \frac{2A(a^k)\beta(a^k)}{1 - \beta(a^k)} \left\langle Q(a^k) - a^k, T(a^k) - a^k \right\rangle.$$

207 Extracting common factors, using Remarks 2 and 5, the fact that $T(a^k) \neq a^k$ (if $T(a^k) = a^k$ then
 208 $a^k = x^u$ by Theorem 3), and the fact that $\tilde{T}(a^k) \neq a^k$ (if $\tilde{T}(a^k) = a^k$ then $T(a^k) = a^k$ by definition (10))
 209 we get that:

$$\begin{aligned} z &= 2A(a^k) \left\| Q(a^k) - a^k \right\| \left\| \tilde{T}(a^k) - a^k \right\| \left[\frac{w_k}{2A(a^k) \left\| \tilde{T}(a^k) - a^k \right\|} \right. \\ &\quad \left. - \beta(a^k) \left\langle \frac{(1 - \lambda(a^k))(T(a^k) - a^k)}{\left\| (1 - \lambda(a^k))(T(a^k) - a^k) \right\|}, \frac{T(a^k) - a^k}{\left\| T(a^k) - a^k \right\|} \right\rangle \right]. \end{aligned}$$

Simplifying and using the definition of $\beta(a^k)$ we have that:

$$z = 2A(a^k) \left\| Q(a^k) - a^k \right\| \left\| \tilde{T}(a^k) - a^k \right\| [\beta(a^k) - \beta(a^k)] = 0,$$

210 which concludes the proof. □

211 The purpose of the next two lemmas is to determine a strict inequality between the functions g_x and
 212 f at suitable points. First of all, we have to prove the following result.

213 **Lemma 14.** *Let $x \in \Omega$ be such that $x \neq Q(x)$.*

214 (a) *If $x \neq a^1, \dots, a^m$, then $g_x(Q(x)) \leq g_x(x)$. Besides that, if $\mathcal{E}(x) \neq \emptyset$ and $P_x \circ Q(x) \neq P_x(x)$, then*
 215 *$g_x(Q(x)) < g_x(x)$.*

216 (b) *If $x = a^k$ for some $k = 1, \dots, m$, then $g_{a^k}(Q(a^k)) < g_{a^k}(a^k)$.*

217 **PROOF.** If $x \neq a^1, \dots, a^m$, then $g_x(Q(x)) - g_x(x) = -A(x)\|Q(x) - x\|_{\mathcal{E}(x)}^2 \leq 0$, by Lemma 11 and
 218 Lemma 12. Besides that, if $\mathcal{E}(x) \neq \emptyset$ and $P_x \circ Q(x) \neq P_x(x)$ we deduce that $\|Q(x) - x\|_{\mathcal{E}(x)} \neq 0$. Thus,
 219 $g_x(Q(x)) < g_x(x)$.

220 If $x = a^k$ for some $k = 1, \dots, m$, by Lemmas 11, 12 and 13 we have:

$$\begin{aligned} g_{a^k}(Q(a^k)) - g_{a^k}(a^k) &= -A(a^k) \left\| Q(a^k) - a^k \right\|^2 \\ &\quad + 2A(a^k) \left\langle Q(a^k) - a^k, Q(a^k) - T(a^k) \right\rangle. \end{aligned}$$

Due to Lemma 9 and the fact that $A > 0$ we obtain:

$$g_{a^k}(Q(a^k)) - g_{a^k}(a^k) \leq -A(a^k) \left\| Q(a^k) - a^k \right\|^2 < 0,$$

221 and the proof is finished. □

222 **Lemma 15.** *Let $x \in \Omega$ be such that $x \neq Q(x)$. Then $g_x(Q(x)) < \frac{1}{2}f(x)$.*

223 **PROOF.** Let us consider the case when $x \neq a^1, \dots, a^m$. By Lemmas 9, 11 and 14 we have that:

$$\begin{aligned} g_x(Q(x)) &\leq g_x(x) = \frac{1}{2}f(x) + 2A(x) \langle Q(x) - x, Q(x) - T(x) \rangle \\ &\quad - A(x) \|Q(x) - x\|_{\mathcal{N}(x)}^2 \\ &\leq \frac{1}{2}f(x) - A(x) \|Q(x) - x\|_{\mathcal{N}(x)}^2. \end{aligned}$$

If $\mathcal{E}(x) = \emptyset$, then $\|\cdot\|_{\mathcal{N}(x)} = \|\cdot\|$. Therefore:

$$g_x(Q(x)) \leq \frac{1}{2}f(x) - A(x) \|Q(x) - x\|^2 < \frac{1}{2}f(x).$$

If $\mathcal{E}(x) \neq \emptyset$ and $P_x \circ Q(x) = P_x(x)$, then there exists an index $i \in \mathcal{N}(x)$ such that $x_i \neq (Q(x))_i$ since $x \neq Q(x)$. Thus, $\|Q(x) - x\|_{\mathcal{N}(x)} \neq 0$, which implies:

$$g_x(Q(x)) \leq \frac{1}{2}f(x) - A(x) \|Q(x) - x\|_{\mathcal{N}(x)}^2 < \frac{1}{2}f(x).$$

If $\mathcal{E}(x) \neq \emptyset$ and $P_x \circ Q(x) \neq P_x(x)$, due to Lemmas 9, 11 and 14, we have that:

$$g_x(Q(x)) < g_x(x) \leq \frac{1}{2}f(x) - A(x) \|Q(x) - x\|_{\mathcal{N}(x)}^2 \leq \frac{1}{2}f(x).$$

224 Now, when $x = a^k$ for some $k = 1, \dots, m$, $g_{a^k}(Q(a^k)) < g_{a^k}(a^k) = \frac{1}{2}f(a^k)$ due to Lemma 11 and
225 Lemma 14. □

226 The next lemma states an equality that relates the Weber function and g_x at appropriate points
227 when $x \neq a^1, \dots, a^m$. Besides that, this result will be crucial in the next section.

Lemma 16. *Let $x \neq a^1, \dots, a^m$ be such that $x \in \Omega$ and $x \neq Q(x)$. Then:*

$$g_x \circ Q(x) = \frac{1}{2}f(x) + (f(Q(x)) - f(x)) + \delta, \quad \delta \geq 0.$$

PROOF. Due to the definition of g_x as in (19) and Remark 8 we get that:

$$g_x \circ Q(x) = \sum_{j=1}^m \frac{w_j}{2\|x - a^j\|} \|Q(x) - a^j\|^2.$$

228 Adding and subtracting $\|x - a^j\|$ we have:

$$\begin{aligned} g_x \circ Q(x) &= \sum_{j=1}^m \frac{w_j}{2\|x - a^j\|} [\|x - a^j\| + (\|Q(x) - a^j\| - \|x - a^j\|)]^2 \\ &= \frac{1}{2} \sum_{j=1}^m w_j \|x - a^j\| + \sum_{j=1}^m w_j (\|Q(x) - a^j\| - \|x - a^j\|) \\ &\quad + \sum_{j=1}^m \frac{w_j}{2\|x - a^j\|} (\|Q(x) - a^j\| - \|x - a^j\|)^2. \end{aligned}$$

Notice that the first term of the last equality is the Weber function (divided by two), and the last term is a non-negative number, so we will define it as δ . So, using the definition of the Weber function in the middle term we obtain:

$$g_x \circ Q(x) = \frac{1}{2}f(x) + (f(Q(x)) - f(x)) + \delta.$$

229 □

230 **5. Convergence to optimality results**

231 This section states the main results about convergence of the sequence $\{x^{(l)}\}$ generated by Algorithm
 232 6. The next theorem establishes that if a point $x \in \Omega$ is not a fixed point of the iteration function, then
 233 the function f strictly decreases at the next iterate.

234 **Theorem 17.** *Let $x \in \Omega$ be such that $x \neq Q(x)$. Then $f(Q(x)) < f(x)$.*

PROOF. Let us consider that $x \neq a^1, \dots, a^m$. By Lemma 15, we have that:

$$g_x \circ Q(x) < \frac{1}{2}f(x).$$

By Lemma 16 we get that:

$$\frac{1}{2}f(x) + f(Q(x)) - f(x) + \delta < \frac{1}{2}f(x).$$

Simplifying the last expression we obtain:

$$f(Q(x)) - f(x) + \delta < 0.$$

Finally,

$$f(Q(x)) - f(x) \leq f(Q(x)) - f(x) + \delta < 0.$$

235 Therefore, $f(Q(x)) < f(x)$.

Now, consider that $x = a^k$ for some $k = 1, \dots, m$. Following a reasoning similar than in [40, pp. 564], using Lemma 14 we have that:

$$g_{a^k} \circ Q(a^k) - g_{a^k}(a^k) < 0.$$

236 By definition of g_{a^k} we know that:

$$\begin{aligned} g_{a^k} \circ Q(a^k) - g_{a^k}(a^k) &= w_k \|Q(a^k) - a^k\| \\ &+ \sum_{\substack{j=1 \\ j \neq k}}^m \frac{w_j}{2 \|a^k - a^j\|} \left(\|Q(a^k) - a^j\|^2 - \|a^k - a^j\|^2 \right). \end{aligned}$$

237 Using the fact that $(a^2 - b^2)/(2b) \geq a - b$ for $a = \|Q(a^k) - a^j\|^2 \geq 0$ and $b = \|a^k - a^j\|^2 > 0$ we obtain
 238 that:

$$\begin{aligned} g_{a^k} \circ Q(a^k) - g_{a^k}(a^k) &\geq w_k \|Q(a^k) - a^k\| \\ &- \sum_{\substack{j=1 \\ j \neq k}}^m w_j \|a^k - a^j\| + \sum_{\substack{j=1 \\ j \neq k}}^m w_j \|Q(a^k) - a^j\|. \end{aligned}$$

239 Rearranging terms we deduce that:

$$\begin{aligned} 0 &> g_{a^k} \circ Q(a^k) - g_{a^k}(a^k) \\ &= \sum_{j=1}^m w_j \|Q(a^k) - a^j\| - \sum_{j=1}^m w_j \|a^k - a^j\| = f(Q(a^k)) - f(a^k), \end{aligned}$$

240 and the proof is complete. □

241 **Corollary 18.** Let $\{x^{(l)}\}$ be the sequence generated by Algorithm 6. Then the sequence $\{f(x^{(l)})\}$ is
 242 not increasing. Even more, each time $x^{(l)} \neq Q(x^{(l)})$ the sequence strictly decreases at the next iterate.

243 If the sequence $\{x^{(l)}\}$ generated by Algorithm 6 were not bounded, then we could choose a subse-
 244 quence $\{y^{(l)}\}$ such that $y^{(l)} \rightarrow \infty$. But this implies that $f(y^{(l)}) \rightarrow \infty$, which is a contradiction since the
 245 sequence $\{f(x^{(l)})\}$ is not increasing.

246 **Remark 19.** The sequence $\{x^{(l)}\}$ generated by Algorithm 6 is bounded. So, there exists a subsequence
 247 convergent to a point $x^* \in \Omega$. Hence, x^* is a feasible point.

Due to the nondifferentiability of f at the vertices a^1, \dots, a^m , we can not use the KKT optimality conditions at a^k . Therefore, if a^k and z are in Ω , let us define $G_{a^k}^z : [0, 1] \rightarrow \mathbb{R}$ by:

$$G_{a^k}^z(t) = f(a^k + t(z - a^k)).$$

248 If $a^k \in \Omega$, $z \in \Omega$, $t \in [0, 1]$ and Ω convex, we have that $a^k + t(z - a^k) \in \Omega$. Notice that the right-hand
 249 side derivative $G_{a^k}^z(0+)$ (or the directional derivative of f in the direction of z) exists (see [22, pp. 33]).
 250 Besides that,

$$G_{a^k}^z(0+) = w_k \|z - a^k\| - \left\langle \tilde{R}(a^k), z - a^k \right\rangle. \quad (20)$$

251 The next lemma shows that if we are in a vertex a^k , the directional derivative of f at a^k in the
 252 direction of $Q(a^k)$ is a descent direction.

253 **Lemma 20.** Let $a^k \in \Omega$ be such that $T(a^k) \notin \Omega$. Then:

$$G_{a^k}^z(0+) \geq G_{a^k}^{Q(a^k)}(0+), \quad \forall z \in [a^k, T(a^k)], \quad (21)$$

254 where:

$$G_{a^k}^{Q(a^k)}(0+) = -2 [1 - \beta(a^k)] A(a^k) \left\| \tilde{T}(a^k) - a^k \right\| \|Q(a^k) - a^k\|. \quad (22)$$

255 **PROOF.** If $T(a^k) = a^k$ then $T(a^k) \in \Omega$, which is a contradiction. Besides that, if $\tilde{T}(a^k) = a^k$, we would
 256 have that $T(a^k) = a^k$ because of (10), and again it would be a contradiction. So, we will consider
 257 $\tilde{T}(a^k) \neq a^k$ and $T(a^k) \neq a^k$ for the rest of the proof. Since $T(a^k) \neq a^k$, then $\beta(a^k) \in (0, 1)$ (see Remark
 258 2 and Theorem 3).

Let us prove equation (22) first. Now, by (20) we can see that:

$$G_{a^k}^{Q(a^k)}(0+) = w_k \|Q(a^k) - a^k\| - \left\langle \tilde{R}(a^k), Q(a^k) - a^k \right\rangle.$$

Notice that if $Q(a^k) = a^k$, equation (22) holds. So, let us consider from now on that $Q(a^k) \neq a^k$. By using Lemma 1 we replace $\tilde{R}(a^k)$ and get:

$$G_{a^k}^{Q(a^k)}(0+) = w_k \|Q(a^k) - a^k\| - 2A(a^k) \left\langle \tilde{T}(a^k) - a^k, Q(a^k) - a^k \right\rangle.$$

259 Extracting common factors and using the definition of β when it belongs to $(0, 1)$ we obtain:

$$\begin{aligned} G_{a^k}^{Q(a^k)}(0+) &= 2A(a^k) \|Q(a^k) - a^k\| \left\| \tilde{T}(a^k) - a^k \right\| \left[\beta(a^k) \right. \\ &\quad \left. - \left\langle \frac{\tilde{T}(a^k) - a^k}{\left\| \tilde{T}(a^k) - a^k \right\|}, \frac{Q(a^k) - a^k}{\|Q(a^k) - a^k\|} \right\rangle \right]. \end{aligned}$$

By Remarks 2 and 5 the vectors $Q(a^k) - a^k$ and $\tilde{T}(a^k) - a^k$ are parallel, so:

$$G_{a^k}^{Q(a^k)'}(0+) = 2A(a^k) \|Q(a^k) - a^k\| \left\| \frac{\tilde{T}(a^k) - a^k}{\|Q(a^k) - a^k\|} \right\| [\beta(a^k) - 1].$$

260 which is equivalent to (22).

Now, let us prove (21). If $z = a^k$ then $G_{a^k}^z(t) = f(a^k)$ for all $t \in [0, 1]$, thus $G_{a^k}^{z'}(0) = 0$, and therefore the inequality (21) holds. So, let us assume that $z \neq a^k$ for the rest of the proof. Using (20) and due to Lemma 1 to replace $\tilde{R}(a^k)$:

$$G_{a^k}^z(0+) = w_k \|z - a^k\| - 2A(a^k) \left\langle \frac{\tilde{T}(a^k) - a^k}{\|\tilde{T}(a^k) - a^k\|}, z - a^k \right\rangle.$$

261 Extracting common factors:

$$\begin{aligned} G_{a^k}^z(0+) &= 2A(a^k) \|z - a^k\| \left\| \frac{\tilde{T}(a^k) - a^k}{\|\tilde{T}(a^k) - a^k\|} \right\| \left[\frac{w_k}{2A(a^k) \|\tilde{T}(a^k) - a^k\|} \right. \\ &\quad \left. - \left\langle \frac{\tilde{T}(a^k) - a^k}{\|\tilde{T}(a^k) - a^k\|}, \frac{z - a^k}{\|z - a^k\|} \right\rangle \right]. \end{aligned}$$

262 Using the expression for $\beta(a^k) \in (0, 1)$ we obtain:

$$\begin{aligned} G_{a^k}^z(0+) &= 2A(a^k) \|z - a^k\| \left\| \frac{\tilde{T}(a^k) - a^k}{\|\tilde{T}(a^k) - a^k\|} \right\| \left[\beta(a^k) \right. \\ &\quad \left. - \left\langle \frac{\tilde{T}(a^k) - a^k}{\|\tilde{T}(a^k) - a^k\|}, \frac{z - a^k}{\|z - a^k\|} \right\rangle \right]. \end{aligned}$$

If z belongs to the segment that joins a^k and $T(a^k)$ we have that $z - a^k$ and $\tilde{T}(a^k) - a^k$ are parallel vectors, then:

$$G_{a^k}^z(0+) \geq -2 [1 - \beta(a^k)] A(a^k) \|z - a^k\| \left\| \frac{\tilde{T}(a^k) - a^k}{\|\tilde{T}(a^k) - a^k\|} \right\|.$$

We can write $z = (1 - \lambda)T(a^k) + \lambda a^k$ where $\lambda \in [0, 1]$. Therefore:

$$G_{a^k}^z(0+) \geq -2 [1 - \beta(a^k)] A(a^k) (1 - \lambda) \|T(a^k) - a^k\| \left\| \frac{\tilde{T}(a^k) - a^k}{\|\tilde{T}(a^k) - a^k\|} \right\|.$$

for all $\lambda \in [0, 1]$. The minimum value of the right-hand side of the last expression happens when $\lambda = \lambda(a^k)$, so:

$$G_{a^k}^z(0+) \geq -2 [1 - \beta(a^k)] A(a^k) (1 - \lambda(a^k)) \|T(a^k) - a^k\| \left\| \frac{\tilde{T}(a^k) - a^k}{\|\tilde{T}(a^k) - a^k\|} \right\|.$$

Using Remark 5 we conclude that:

$$G_{a^k}^z(0+) \geq -2 [1 - \beta(a^k)] A(a^k) \|Q(a^k) - a^k\| \left\| \frac{\tilde{T}(a^k) - a^k}{\|\tilde{T}(a^k) - a^k\|} \right\|.$$

264 Now we will prove an equivalence that characterizes the solution of (5) in terms of the iteration
 265 function Q . Moreover, if x^* is a regular point that is not a vertex, then x^* is a KKT point.

266 From now on, let us consider that

$$\Omega = \{y \in \mathbb{R}^n : g(y) \leq 0, h(y) = 0\}, \quad (23)$$

267 where $g : \mathbb{R}^n \rightarrow \mathbb{R}^s$ is a convex function and $h : \mathbb{R}^n \rightarrow \mathbb{R}^p$ is an affine function.

268 **Theorem 21.** *Let Ω be defined as in (23) and $x \in \Omega$. Consider the following propositions:*

269 (a) x is a KKT point.

270 (b) x is the minimizer of the problem (5).

271 (c) $Q(x) = x$.

272 If $x \neq a^1, \dots, a^m$, g and h are continuously differentiable, and x is a regular point, then (a), (b) and
 273 (c) are equivalent.

274 If $x = a^k$ for some $k = 1, \dots, m$, then (b) implies (c).

275 **PROOF.** Let $x \neq a^1, \dots, a^m$ be. Since f is strictly convex and Ω is convex, the KKT optimality
 276 conditions are necessary and sufficient. Therefore, it holds that (a) is equivalent to (b).

277 Now we will prove that (b) implies (c). Let us suppose that x is the minimizer of the problem (5).
 278 If x were not a fixed point of the iteration function Q , we would have that $x \neq Q(x)$, which means that
 279 $f(Q(x)) < f(x)$ by Theorem 17. This contradicts the hypothesis.

To demonstrate that (c) implies (a), we will assume that x is a fixed point of Q , that is, $x = Q(x)$.
 Since $Q(x) = P_\Omega \circ T(x)$, x is the solution of:

$$\begin{aligned} \underset{z}{\operatorname{argmin}} \quad & F(z) = \frac{1}{2} \|z - T(x)\|^2 \\ \text{subject to} \quad & g(z) \leq 0, \\ & h(z) = 0. \end{aligned}$$

280 Since F and g are convex, h is affine, and x is a regular point, the KKT optimality conditions hold at
 281 x . That is, there exist multipliers $\{\mu_j\}_{j=1}^s$ and $\{\lambda_j\}_{j=1}^p$ such that (see [31, 38]):

$$\begin{aligned} x - T(x) + \sum_{j=1}^s \mu_j \nabla g_j(x) + \sum_{j=1}^p \lambda_j \nabla h_j(x) &= 0, \\ \mu_j g_j(x) &= 0, \quad j = 1, \dots, s, \\ \mu_j &\geq 0, \quad j = 1, \dots, s, \\ g(x) &\leq 0, \\ h(x) &= 0. \end{aligned}$$

282 Multiplying these equations by $2A(x)$, using equation (9), Lemma 1 and Remark 2, we obtain:

$$\begin{aligned} \nabla f(x) + \sum_{j=1}^s (2A(x)\mu_j) \nabla g_j(x) + \sum_{j=1}^p (2A(x)\lambda_j) \nabla h_j(x) &= 0, \\ (2A(x)\mu_j) g_j(x) &= 0, \quad j = 1, \dots, s, \\ (2A(x)\mu_j) &\geq 0, \quad j = 1, \dots, s, \\ g(x) &\leq 0, \\ h(x) &= 0. \end{aligned}$$

283 where $\{2A(x)\mu_j\}_{j=1}^s$ and $\{2A(x)\lambda_j\}_{j=1}^p$ are multipliers. Therefore, x is a KKT point of the problem (5)
 284 (see [31, 38]).

285 Now, let us suppose that $x = a^k$ for some $k = 1, \dots, m$. As before, if x is a minimizer of the problem
 286 (5), then $Q(a^k) = a^k$, otherwise $f(Q(a^k)) < f(a^k)$, which would be a contradiction. \square

287 **6. Numerical experiments.**

288 The purpose of this section is to discuss the efficiency and robustness of the proposed algorithm
 289 versus a solver for nonlinear programming problems.

290 A prototype code of Algorithm 6 was programmed in MATLAB (version R2011a) and executed in a
 291 PC running Linux OS, Intel(R) Core(TM) i7 CPU Q720, 1.60GHz.

We have considered a closed and convex set $\Omega \subset \mathbb{R}^2$ defined by the set $\Omega = \{y \in \mathbb{R}^n : g(y) \leq 0\}$, where g is given by:

$$g(x) = \begin{bmatrix} -4 - \frac{1}{8}x + \frac{7}{72}x^2 + \frac{1}{216}x^2(x-3) + y \\ \frac{4}{5}x + y - \frac{59}{10} \\ x - \frac{11}{2} \\ \frac{3}{2}x - y - \frac{35}{4} \\ x - y - \frac{13}{2} \\ -4 + \frac{1}{8}(x-1) + \frac{1}{16}(x-1)^2 + \frac{1}{32}(x-1)^2(x-3) - y \\ -\frac{1}{3}x - y - \frac{11}{3} \\ -\frac{2}{3}x - y - \frac{13}{3} \\ -4x + y - 19 \end{bmatrix}.$$

292 The feasible set is defined by linear and nonlinear constraints, as it can be seen in Figure 1.

293

294 We have built 1000 different experiments where for each one:

- 295 • The number of vertices was $m = 50$.
- 296 • The vertices were normally distributed random vectors, with mean equal to 0 and standard deviation equal to 10.
- 297
- 298 • The weights were uniformly distributed random positive numbers between 0 and 10.
- 299 • Tolerance was set to $\varepsilon = 0.00001$.

300 On one hand, each experiment was solved using Algorithm 6 and, on the other hand, it was considered
 301 as a nonlinear programming problem and solved using function *fmincon* (see [32] and references therein).
 302 Since the Weber function (1) is not differentiable at the vertices, nonlinear programming solvers may
 303 fail.

304 Let $x_m(i)$ be the solution of (5) obtained by *fmincon* in experiment i , and $f_m(i) = f(x_m(i))$.
 305 Analogously, let $x_p(i)$ be the solution of (5) obtained by Algorithm 6 in experiment i , and $f_p(i) =$
 306 $f(x_p(i))$. Figure 2 shows the difference between the arrays f_m and f_p . Both methods finished succesfully
 307 in all cases, however, Algorithm 6 found equal or better results for all experiments. For example,
 308 the difference $f_m - f_p$ was greater than 0.01 in 35 experiments (the maximum difference ocurred in
 309 experiment 506).



Figure 1: Feasible set Ω




Figure 2: Difference between minimum values found by Algorithm 6 and $fmincon$.




Figure 3: Feasibility of the solution $x_p(i)$ obtained by Algorithm 6.

310

311 Feasibility of the solutions $x_p(i)$ can be checked computing $\max(g(x_m(i)))$. Results can be seen in
 312 Figure 3

313

314 7. Conclusions

315 This paper proposes a Weiszfeld-like algorithm for solving the Weber problem constrained to a closed
 316 and convex set, and it is well defined even when an iterate is a vertex. The algorithm consists of two
 317 stages: first, iterate using the fixed point modified Weiszfeld iteration (10), and second, either project
 318 onto the set Ω when the iterate is different from the vertices, or, if the iterate is a vertex a^k , take the
 319 point belonging to the line that joins $T(a^k)$ with a^k as defined in (11).

320 It is proved that the constrained problem (5) has a unique solution. Besides that, the definition
 321 of the iteration function Q allows us to demonstrate that the proposed algorithm produces a sequence
 322 $\{x^{(l)}\}$ of feasible iterates. Moreover, the sequence $\{f(x^{(l)})\}$ is not increasing, and when $x^{(l)} \neq Q(x^{(l)})$,
 323 the sequence decreases at the next iterate. It can be seen that if a point x^* is the solution of the problem
 324 (5) then x^* is a fixed point of the iteration function Q . Even more, if x^* is different from the vertices,
 325 the fact of being x^* a fixed point of Q is equivalent to the fact that x^* satisfies the KKT optimality
 326 conditions, and equivalent to the fact that x^* is the solution of the problem (5). These properties allows
 327 us to connect the proposed algorithm with the minimization problem.

328 Numerical experiments showed that the proposed algorithm found equal or better solutions than a
 329 well-known standard solver, in a practical example with 1000 random choices of vertices and weights.
 330 That is due to the fact that the proposed algorithm does not use of the existence of derivatives at the
 331 vertices, because the Weber function is not differentiable at the vertices.

332 **References**

- 333 [1] N. Andréasson, A. Evgrafov, M. Patriksson, An introduction to continuous optimization, Stu-
334 dentlitteratur, 2007.
- 335 [2] Y.P. Aneja, M. Parlar, Technical note-Algorithms for Weber facility location in the presence of
336 forbidden regions and/or barriers to travel, *Transportation Science* 28 (1994) 70–76.
- 337 [3] R.J. Cañavate Bernal, Algunas cuestiones teóricas sobre la validez del algoritmo de Weiszfeld para
338 el problema de Weber, *Rect@ Actas_13* (2005) 24.
- 339 [4] M. Bischoff, T. Fleischmann, K. Klamroth, The multi-facility location-allocation problem with
340 polyhedral barriers, *Comput. Oper. Res.* 36 (2009) 1376–1392.
- 341 [5] M. Bischoff, K. Klamroth, An efficient solution method for Weber problems with barriers based on
342 genetic algorithms, *European J. Oper. Res.* 177 (2007) 22–41.
- 343 [6] J. Brimberg, The Fermat-Weber location problem revisited, *Mathematical Programming* 71 (1995)
344 71–76.
- 345 [7] J. Brimberg, Further notes on convergence of the Weiszfeld algorithm, *Yugosl. J. Oper. Res.* 13
346 (2003) 199–206.
- 347 [8] S.E. Butt, T.M. Cavalier, An efficient algorithm for facility location in the presence of forbidden
348 regions, *European J. Oper. Res.* 90 (1996) 56–70.
- 349 [9] L. Cánovas, R. Cañavate, A. Marín, On the convergence of the Weiszfeld algorithm, *Mathematical*
350 *Programming* 93 (2002) 327–330.
- 351 [10] E. Carrizosa, M. Muñoz-Márquez, J. Puerto, The Weber problem with regional demand, *European*
352 *J. Oper. Res.* 104 (1998) 358–365.
- 353 [11] R. Chandrasekaran, A. Tamir, Open questions concerning Weiszfeld’s algorithm for the Fermat-
354 Weber location problem, *Mathematical Programming* 44 (1989) 293–295.
- 355 [12] J.A. Chatelon, D.W. Hearn, T.J. Lowe, A subgradient algorithm for certain minimax and minisum
356 problems, *Mathematical Programming* 15 (1978) 130–145.
- 357 [13] P.C. Chen, P. Hansen, B. Jaumard, H. Tuy, Weber’s problem with attraction and repulsion, *Journal*
358 *of Regional Science* 32 (1992) 467–486.
- 359 [14] R. Courant, D. Hilbert, *Methoden der mathematischen Physik. I*, Springer-Verlag, Berlin, 1968.
360 Dritte Auflage, Heidelberger Taschenbucher, Band 30.
- 361 [15] Z. Drezner, On the convergence of the generalized Weiszfeld algorithm, *Ann. Oper. Res.* 167 (2009)
362 327–336.
- 363 [16] Z. Drezner, G.O. Wesolowsky, The location of an obnoxious facility with rectangular distances,
364 *Journal of Regional Science* 23 (1983) 241–248.
- 365 [17] Z. Drezner, G.O. Wesolowsky, Sensitivity analysis to the value of p of the l_p distance Weber problem,
366 *Ann. Oper. Res.* 111 (2002) 135–150.
- 367 [18] U. Eckhardt, Weber’s problem and Weiszfeld’s algorithm in general spaces, *Mathematical Program-*
368 *ming* 18 (1980) 186–196.

- 369 [19] J.W. Eyster, J.A. White, W.W. Wierwille, On solving multifacility location problems using a
370 hyperboloid approximation procedure, *A I I E Transactions* 5 (1973) 01–06.
- 371 [20] E. Fasbender, Über die gleichseitigen Dreiecke, welche um ein gegebenes Dreieck gelegt werden
372 können, *Journal für die reine und angewandte Mathematik* 30 (1846) 230–231.
- 373 [21] P. Hansen, D. Peeters, J. Thisse, An algorithm for a constrained Weber problem, *Management*
374 *Science* (1982) 1285–1295.
- 375 [22] J. Jahn, *Introduction to the theory of nonlinear optimization*, Springer Verlag, 2007.
- 376 [23] G. Jalal, J. Krarup, Geometrical solution to the Fermat problem with arbitrary weights, *Ann. Oper.*
377 *Res.* 123 (2003) 67–104.
- 378 [24] W. Kaplan, W. Yang, Duality theorem for a generalized Fermat-Weber problem, *Mathematical*
379 *Programming* 76 (1997) 285–297.
- 380 [25] I.N. Katz, L. Cooper, Facility location in the presence of forbidden regions. I. Formulation and the
381 case of Euclidean distance with one forbidden circle, *European J. Oper. Res.* 6 (1981) 166–173.
- 382 [26] K. Klamroth, *Single-facility location problems with barriers*, Springer Series in Operations Research,
383 Springer-Verlag, New York, 2002.
- 384 [27] J. Krarup, S. Vajda, On Torricelli’s geometrical solution to a problem of Fermat, *IMA J. Math.*
385 *Appl. Bus. Indust.* 8 (1997) 215–224. Duality in practice.
- 386 [28] H.W. Kuhn, On a pair of dual nonlinear programs, in: *Nonlinear Programming (NATO Summer*
387 *School, Menton, 1964)*, North-Holland, Amsterdam, 1967, pp. 37–54.
- 388 [29] H.W. Kuhn, A note on Fermat’s problem, *Mathematical Programming* 4 (1973) 98–107.
- 389 [30] Y.S. Kupitz, H. Martini, Geometric aspects of the generalized Fermat-Torricelli problem, in: *In-*
390 *tuitive geometry (Budapest, 1995)*, volume 6 of *Bolyai Soc. Math. Stud.*, János Bolyai Math. Soc.,
391 Budapest, 1997, pp. 55–127.
- 392 [31] D.G. Luenberger, Y. Ye, *Linear and nonlinear programming*, International Series in Operations
393 Research & Management Science, 116, Springer, New York, third edition, 2008.
- 394 [32] MATLAB, <http://www.mathworks.com/help/toolbox/optim/ug/fmincon.html>, 2012.
- 395 [33] C. Michelot, O. Lefebvre, A primal-dual algorithm for the Fermat-Weber problem involving mixed
396 gauges, *Mathematical Programming* 39 (1987) 319–335.
- 397 [34] M.L. Overton, A quadratically convergent method for minimizing a sum of Euclidean norms, *Math-*
398 *ematical Programming* 27 (1983) 34–63.
- 399 [35] B. Pfeiffer, K. Klamroth, A unified model for Weber problems with continuous and network dis-
400 tances, *Comput. Oper. Res.* 35 (2008) 312–326.
- 401 [36] E.A. Pilotta, G.A. Torres, A projected Weiszfeld algorithm for the box-constrained Weber location
402 problem, *Applied Mathematics and Computation* 218 (2011) 2932–2943.
- 403 [37] G. Polya, *Mathematik und plausible Schließen, Band 1, Induktion Und Analogie in Der Mathe-*
404 *matik*, Birkhauser, Basel, 1962.

- 405 [38] A. Quarteroni, R. Sacco, F. Saleri, Numerical mathematics, volume 37, Springer, 2006.
- 406 [39] M.K. Schaefer, A.P.J. Hurter, An algorithm for the solution of a location problem with metric
407 constraints, Naval Res. Logist. Quart. 21 (1974) 625–636.
- 408 [40] Y. Vardi, C.H. Zhang, A modified Weiszfeld algorithm for the Fermat-Weber location problem,
409 Mathematical Programming 90 (2001) 559–566.
- 410 [41] A. Weber, Über den Standort der Industrien, University of Chicago Press, Chicago, 1929. Translated
411 by C. J. Friederich: “Theory of the location of industries”.
- 412 [42] E. Weiszfeld, Sur le point par lequel la somme des distances de n points donnés est minimum,
413 Tohoku Mathematics Journal 43 (1937) 355–386.
- 414 [43] E. Weiszfeld, On the point for which the sum of the distances to n given points is minimum, Ann.
415 Oper. Res. 167 (2009) 7–41.
- 416 [44] G.O. Wesolowsky, The Weber problem: history and perspectives, Location Sci. 1 (1993) 5–23.