SEMI-CLASSICAL ORTHOGONAL POLYNOMIAL SYSTEMS ON NON-UNIFORM LATTICES, DEFORMATIONS OF THE ASKEY TABLE AND ANALOGS OF ISOMONODROMY

N.S. WITTE

ABSTRACT. A D-semi-classical weight is one which satisfies a particular linear, first order homogeneous equation in a divided-difference operator D. It is known that the system of polynomials, orthogonal with respect to this weight, and the associated functions satisfy a linear, first order homogeneous matrix equation in the divided-difference operator termed the spectral equation. Attached to the spectral equation is a structure which constitutes a number of relations such as those arising from compatibility with the three-term recurrence relation. Here this structure is elucidated in the general case of quadratic lattices. The simplest examples of the D-semi-classical orthogonal polynomial systems are precisely those in the Askey table of hypergeometric and basic hypergeometric orthogonal polynomials. However within the D-semi-classical class it is entirely natural to define a generalisation of the Askey table weights which involve a deformation with respect to new deformation variables. We completely construct the analogous structures arising from such deformations and their relations with the other elements of the theory. As an example we treat the first non-trivial deformation of the Askey-Wilson orthogonal polynomial system defined by the *q*-quadratic divided-difference operator, the Askey-Wilson operator, and derive the coupled first order divided-difference equations characterising its evolution in the deformation variable. We show that this system is a member of a sequence of classical solutions to the $E_7^{(1)} q$ -Painlevé system.

CONTENTS

1. Background and Motivation	1
2. Divided-Difference Calculus of the Quadratic Lattices	4
3. Orthogonal Polynomial Systems on the Non-uniform Lattice	7
3.1. General orthogonal polynomial systems	7
3.2. q-Quadratic lattice	11
4. Spectral Differences	13
5. $M = 2, L = 0$ case and the Askey-Wilson polynomials	22
6. Deformation Differences	26
7. $M = 3, L = 1$ case and deformation of the Askey-Wilson OPS	35
7.1. Moments and Integrals	35
7.2. Spectral Structure	38
7.3. Deformation Structure	41
7.4. Seed Solution	53
8. Acknowledgments	55
References	55

1. BACKGROUND AND MOTIVATION

We propose a method for constructing systems of linear divided-difference equations which are analogs of isomonodromic linear differential equations and therefore "isomonodromic interpretations" or Lax pairs of the known q-Painlevé and difference Painlevé equations, such as those in the Sakai scheme [86]. What is

²⁰⁰⁰ Mathematics Subject Classification. 39A05,42C05,34M55,34M56,33C45,37K35.

Key words and phrases. non-uniform lattices, divided-difference operators, orthogonal polynomials, semi-classical weights, isomonodromic deformations, Askey table, Askey-Wilson polynomials.

meant by an analog of a monodromy matrix turns out to be a connection matrix, appropriate to the class of lattices defining the divided-difference operators under consideration. In essence our method constructs a particular "isomonodromic analog" system from an orthogonal polynomial system, orthogonal with respect to a generalisation or deformation of a weight with discrete or countable support on a class of non-uniform quadratic lattices. We then deduce a number of linear divided-difference equations that this system satisfies and show that their pair-wise compatibility holds provided the coefficients of the linear system obey evolution equations of the difference or q-Painlevé type. This is a very natural extension of the Fokas-Its-Kitaev construction [49, 23, 24] at the heart of Riemann-Hilbert techniques.

Our method is independent of and distinct from other approaches which we briefly recount here. The first studies to construct Lax pairs for the q-Painlevé equations were those of Jimbo and Sakai [50] and Sakai [87, 88] using the Birkhoff theory of linear q-difference equations Y(qx) = A(x)Y(x) and imposing the condition that the connection matrix was independent of the zeros of det A(x). However this approach has not been extended beyond the $D_5^{(1)}$ or $E_6^{(1)}$ cases (the latter case only found from a degeneration of the two-variable extension of the former). Another distinct approach which is founded upon the notion of the τ -function of a rational d-connection is the Arinkin-Borodin theory [17, 2, 3], which has been applied to the difference Painlevé equations. Our approach is similar in spirit to that of Rains [85], who has treated the master elliptic Painlevé equation, in that an explicit construction is made of the solution to the linear problem which contains a multiple integral representation of the orthogonal rational function. However we will not make direct contact with this theory because it requires us to consider orthogonal rational functions on elliptic lattices, which is a generalisation beyond the class of lattices considered here. In addition we believe that the discrete Riemann-Hilbert approach, as formulated by Borodin in [16] and applied to two examples and extended to further cases in [18], shares many features with the present study and one should be able to construct a Riemann-Hilbert formulation of our own theory. Recently Yamada has constructed Lax pairs for the *q*-Painlevé equations for the systems with symmetries $E_8^{(1)}, E_7^{(1)}, E_6^{(1)}$ by a reduction from the elliptic form of the $E_8^{(1)}$ Painlevé equation [98, 99] however no theoretical construction from first principles was proposed in the individual cases.

The approach we propose here has been successfully employed for the isomonodromic systems and Painlevé equations, see for example [66, 61, 62, 13, 49, 19, 21, 26, 93, 27, 94], and while most of the findings are reproductions of known ones it has lead to novel results hitherto not found using other treatments, such as the discrete Garnier systems [95]. An important feature of our approach is that it is strongly motivated by a probabilistic setting, namely that of the theory of random matrices and more generally determinantal point processes where the classical weights in the Askey table appear in the one-body factors of the eigenvalue probability density functions [25]. Some preliminary exploration of the program we propose here has already been initiated but not carried through to its logical conclusion, and we will delay citing this work until the body of our paper where it is directly relevant. However we should point out that we are most indebted to the pioneering work of Magnus [60, 63].

The essential elements of our approach are the following.

(A) The classification of special non-uniform lattices SNUL of quadratic type [60], their associated divideddifference operators \mathbb{D}_x and \mathbb{M}_x and their rules of calculus, which applies to orthogonal or bi-orthogonal polynomial systems. Of significance is the fact that in general these lattices possess two fixed points which we denote x_L, x_R .

(B) The notion of a D-semi-classical weight w(x) which is characterised by an analog of the Pearson equation (see Definition 4.1) [60, 63]

(1.1)
$$\mathbb{D}_x w(x) = \frac{2V(x)}{W(x)} \mathbb{M}_x w(x),$$

where W(x), V(x) are polynomials in $\mathbb{C}[x]$.

(C) The orthogonal polynomial system (OPS) defined by such a weight on the SNUL $Y_n(x) \in \mathbb{C}^{2\times 2}$ satisfies a three-term recurrence relation, which in our context is a particular Schlesinger transformation $n \mapsto n+1$ (see 3.29)

(1.2)
$$Y_{n+1}(x) = K_n(x)Y_n(x).$$

(D) The semi-classical character implies, under fairly weak conditions, a spectral structure on the "isomonodromic" system $Y_n(x)$ [60, 63] (see Proposition 4.4), namely that it satisfies the linear divided-difference equation

(1.3)
$$\mathbb{D}_x Y_n(x) = A_n(x) \mathbb{M}_x Y_n(x).$$

Here $A_n(x)$ is rational in x and the degrees of its numerator and denominator with respect to x are independent of n.

(E) Parallel to the spectral structure is a deformation structure, whereby the weight and the system acquires a u dependence, constrained by another Pearson relation (see Definition 6.1)

(1.4)
$$\mathbb{D}_u w(x;u) = \frac{2S(x;u)}{R(x;u)} \mathbb{M}_u w(x;u),$$

where R(x; u), S(x; u) are polynomials in $\mathbb{C}[x]$.

(F) This also has a direct consequence for the "isomonodromic" system $Y_n(x; u)$, a deformation structure (see Proposition 6.6) and a second associated linear divided-difference equation

(1.5)
$$\mathbb{D}_u Y_n(x;u) = B_n(x;u) \mathbb{M}_u Y_n(x;u),$$

where $B_n(x; u)$ is also rational in x.

(G) The compatibility relations implied by this over-determined "isomonodromic" system $Y_n(x; u)$ then lead to a number of conclusions - the deformation matrix $B_n(x; u)$ is expressible in terms of the spectral matrix $A_n(x; u)$ at neighbouring lattice nodes which we refer to as *closure* (see Proposition 7.8), and furthermore relations exist between components of the spectral matrix at two consecutive nodes on the *u*-lattice which, given a suitable parameterisation of this matrix by appropriate co-ordinates, is an recurrence relation on the deformation lattice. Our key results for such recurrence relations are given in Propositions 7.9, 7.10, 7.11 and 7.12.

(H) In our approach we derive a preservation property for the connection matrix defined as

(1.6)
$$P(x;u) := (Y_R(x;u))^{-1} Y_L(x;u)$$

where $Y_{L,R}(x; u)$ are local fundamental solutions of the spectral equation about $x_{L,R}$. Specifically this implies

(1.7)
$$\mathbb{D}_x P(x; u) = 0, \qquad \mathbb{D}_u P(x; u) = 0,$$

as a consequence of our assumptions.

In our approach we can sidestep a number of issues, which are important to be sure, but don't affect the outcome. So we postpone deeper considerations of an analytic or algebraic theory of linear systems of divided-difference operators on these lattices for subsequent studies, however our present work will provide concrete illustrative examples for such an investigation. We are referring to, for example, issues of a Galois theory for D-difference equations, a Birkhoff theory for the local character of the solutions to systems of linear first order D-difference equations and analogs of monodromy for D-difference equations.

Our approach poses the question concerning a correspondence between the system of hypergeometric and basic hypergeometric orthogonal polynomial systems generalising the classical systems, known as the Askey table, as represented in reference work [56], and the most complete system of elliptic, q-difference and difference analogs of the Painlevé equations, the Sakai scheme [86]. A correspondence between the Sakai scheme and the Askey table would explain the occurrence of many features that have been discovered recently such as the appearance of basic hypergeometric functions in their classical solutions. We have already referred to the correspondence between the classical orthogonal polynomial systems (OPS) Hermite, Laguerre and Jacobi and the classical solutions to Painlevé IV, V and VI respectively. Amongst the difference and q-difference OPS the current evidence for such a correspondence can be summarised in the Table 1.

The layout of this work loosely follows the plan given above.

In Section 2 we describe the classification of the quadratic lattices and their divided-difference calculus, with special emphasis on the example of the master class, the q-quadratic lattice. Section 3 is devoted to the reformulation of orthogonal polynomial system theory necessary for systems with a weight having support on a general quadratic lattice, and some analytical results for the q-quadratic lattice. The spectral structures are laid out in Section4 for a general quadratic lattice, starting with the definition of a \mathbb{D} -semi-classical weight and developing the consequences of this for the orthogonal polynomial system. We also describe

Base OPS	Integrable system	Reference
little q -Jacobi	$D_5^{(1)}$, full $q - \mathbf{P}_{\mathrm{VI}}$	[79]
Pastro [80]	$D_5^{(1)}$, full $q - \mathbf{P}_{\mathrm{VI}}$	[14]
little q -Jacobi	$D_5^{(1)}$, special $q - P_{VI}$	[18]
q-Krawtchouk	$D_5^{(1)}$, special $q - P_{VI}$	[18]
q-Charlier	$D_5^{(1)}$, degenerate $q - P_{VI}$	[18]
q-Freud	$A_4^{(1)}$, degenerate $q - P_V$	[15]
Meixner, Krawtchouk	$d - P_V$	[18]
Charlier	$d - P_{IV}$	[18]

TABLE 1. The Askey table-Sakai scheme correspondence ranked according to the degeneration pathway from the master case. The affine Weyl group refers to the symmetry group of Bäcklund transformations for the integrable system. Only the discrete lattices are included here.

the compatibility relations of this structure with the three-term recurrence relations, and this leads to a generalisation of the Laguerre-Freud equations. An explicit example of the forgoing theory is the lowest case, and from our point of view the trivial case, of the q-quadratic lattice which is denoted by the label M = 2. This is dealt with in Section 5. We recover the Askey-Wilson OPS and demonstrate that every aspect of this system can be derived in an efficient manner using the theoretical tools developed in the previous section.

In Section 6 we introduce an (or many) auxiliary variable and demand that the weight satisfies an analogous \mathbb{D} -semi-classical relation to the spectral relation, on a general lattice not necessarily the same as that for the spectral variable. This constitutes a *deformation* of the weight which leaves it having the same structure as the original. From this we develop a parallel theoretical analysis of the orthogonal polynomial system with respect to this deformation. We now have two compatibility relations with this deformation structure - one with the three-term recurrence and other with the spectral structure. For the present purposes this completes our task for the general theory.

In Section 7 we treat a natural deformation of the Askey-Wilson weight, but not by any means the only one, to the q-quadratic lattice. Here we construct parameterisations of the spectral and deformation matrices, the closure relations between the two structures, determine the co-ordinate transformations to a set of variables and derive the evolution equations for this system (see Propositions 7.9, 7.10,7.11 and 7.12). Finally we offer evidence that our system is a classical solution of the $E_7^{(1)}$ q-Painlevé system.

2. DIVIDED-DIFFERENCE CALCULUS OF THE QUADRATIC LATTICES

Divided-difference operators associated with the special non-uniform lattices have appeared in many studies of orthogonal polynomials of a discrete variable. For example see the early studies by Hahn [36, 35, 37, 38], the foundational work by Askey and Wilson [4] and the monograph of Nikiforov, Suslov and Uvarov [78]. The employment of divided-difference operators such as the Askey-Wilson and Wilson operators has been common place in studies of the Askey table of hypergeometric orthogonal polynomial systems too numerous to cite here (see the monographs by Ismail [43] and Lesky [59]).

Let $\Pi_n[x]$ denote the linear space of polynomials in x over \mathbb{C} with degree at most $n \in \mathbb{Z}_{\geq 0}$. In pioneering investigations Magnus [60, 63] provided a geometrical understanding of these lattices and their divided-difference operators which we relate here briefly. If we define the *divided-difference operator* (DDO) \mathbb{D}_x by

(2.1)
$$\mathbb{D}_x f(x) = \frac{f(\iota_+(x)) - f(\iota_-(x))}{\iota_+(x) - \iota_-(x)},$$

then a simple consequence of the condition that $\mathbb{D}_x : \Pi_n[x] \to \Pi_{n-1}[x]$ for all $n \in \mathbb{N}$ is that $\iota_{\pm}(x)$ are the two *y*-roots of the quadratic equation

(2.2)
$$\mathcal{A}y^2 + 2\mathcal{B}xy + \mathcal{C}x^2 + 2\mathcal{D}y + 2\mathcal{E}x + \mathcal{F} = 0.$$

The functions $\iota_{\pm}(x)$ satisfy

(2.3)
$$\iota_{+}(x) + \iota_{-}(x) = -2\frac{\mathcal{B}x + \mathcal{D}}{\mathcal{A}},$$

(2.4)
$$\iota_+(x)\iota_-(x) = \frac{\mathcal{C}x^2 + 2\mathcal{E}x + \mathcal{F}}{\mathcal{A}},$$

and their inverse functions ι_{\pm}^{-1} are defined by $\iota_{\pm}^{-1}(\iota_{\pm}(x)) = x$. For a given y-value the quadratic (2.2) defines two x-roots if $\mathcal{C} \neq 0$, which are consecutive points on the x-lattice, $x_s := x(s), x_{s+1} := x(s+1)$ parameterised by the variable s and therefore defines a map $x_s \mapsto x_{s+1}$. Conversely for a given x-value the quadratic defines two y-roots if $\mathcal{A} \neq 0$ which are consecutive points on a dual lattice, the y-lattice, $y_s := y(s) = \iota_{-}(x(s)), y_{s+1} := y(s+1) = \iota_{+}(x(s))$ which generally is distinct from the x-lattice. We will employ an operator notation for the mappings from points on the direct lattice to the dual lattice $E_x^{\pm}f(x) := f(\iota_{\pm}(x))$ so that

(2.5)
$$\mathbb{D}_x f(x) := \frac{f(\iota_+(x)) - f(\iota_-(x))}{\iota_+(x) - \iota_-(x)} = \frac{E_x^+ f - E_x^- f}{E_x^+ x - E_x^- x},$$

for arbitrary functions f(x). The inverse functions $\iota_{\pm}^{-1}(x)$ define operators $(E^{\pm})^{-1}$ which map points on the dual lattice to the direct lattice and also an adjoint to the divided-difference operator \mathbb{D}_x

(2.6)
$$\mathbb{D}_x^* f(x) := \frac{f(\iota_+^{-1}(x)) - f(\iota_-^{-1}(x))}{\iota_+^{-1}(x) - \iota_-^{-1}(x)} = \frac{(E_x^+)^{-1} f - (E_x^-)^{-1} f}{(E_x^+)^{-1} x - (E_x^-)^{-1} x}$$

The composite operators $E_x := (E_x^-)^{-1}E_x^+$ and $E_x^{-1} = (E_x^+)^{-1}E_x^-$ map between consecutive points on the direct lattice.

However in the situation of a symmetric quadratic $\mathcal{A} = \mathcal{C}$ and $\mathcal{D} = \mathcal{E}$, which entails no loss of generality, then we have $(E_x^+)^{-1} = E_x^-$ and $(E_x^-)^{-1} = E_x^+$. Consequently there is no distinction between the divided-difference operator and its adjoint, and hereafter we adopt this simplification. For example one useful consequence of this choice is that the ratio

(2.7)
$$\frac{(x-E^+u)(x-E^-u)}{(E^-x-u)(E^+x-u)} = \frac{\mathcal{A}x^2 + 2\mathcal{B}xu + \mathcal{C}u^2 + 2\mathcal{D}x + 2\mathcal{E}u + \mathcal{F}}{\mathcal{A}u^2 + 2\mathcal{B}ux + \mathcal{C}x^2 + 2\mathcal{D}u + 2\mathcal{E}x + \mathcal{F}} = 1$$

for all x, u. A companion operator to the divided-difference operator \mathbb{D}_x is the mean or average operator \mathbb{M}_x defined by

(2.8)
$$\mathbb{M}_x f(x) = \frac{1}{2} \left[f(\iota_+(x)) + f(\iota_-(x)) \right],$$

so that the property $\mathbb{M}_x : \Pi_n[x] \to \Pi_n[x]$ is ensured by the condition we imposed upon \mathbb{D}_x .

Definition 2.1. Henceforth we use the shorthand for the difference in consecutive y points $\Delta y(x) := \iota_+(x) - \iota_-(x)$. We introduce the notion of "fixed points", whereby $\Delta y^2(x_F) = 0$, or $\iota_+(x_F) = \iota_-(x_F)$ and are given as the roots of $(\mathcal{B}^2 - \mathcal{AC})x_F^2 + 2(\mathcal{BD} - \mathcal{AE})x_F + \mathcal{D}^2 - \mathcal{AE} = 0$.

Assuming $\mathcal{AC} \neq 0$ one can classify these non-uniform quadratic lattices (or *SNUL*, special non-uniform lattices) according to two parameters - the discriminant $\mathcal{B}^2 - \mathcal{AC}$ and

(2.9)
$$\Theta = \det \begin{pmatrix} \mathcal{A} & \mathcal{B} & \mathcal{D} \\ \mathcal{B} & \mathcal{C} & \mathcal{E} \\ \mathcal{D} & \mathcal{E} & \mathcal{F} \end{pmatrix},$$

or $\mathcal{A}\Theta = (\mathcal{B}^2 - \mathcal{A}\mathcal{C})(\mathcal{D}^2 - \mathcal{A}\mathcal{F}) - (\mathcal{B}\mathcal{D} - \mathcal{A}\mathcal{E})^2$. There are four primary classes - the linear lattice, the linear q-lattice, the quadratic lattice and the q-quadratic lattice, which are given in Table 2. The q-quadratic lattice, in its general non-symmetrical form, is the most general case and the other lattices can be found from this by limiting processes.

This classification of lattices for polynomial systems can be extended to rational function systems [89, 90, 64] and in this case one has a lattice characterised by a bi-quadratic relation and parameterised by elliptic

$\mathcal{B}^2 - \mathcal{AC}$	Θ	conic	lattice	canonical DDO	Notes
0	0	parallel lines	linear	forward difference	
> 0	0	intersecting lines	q-linear	q-difference	
0	< 0	parabola	quadratic	Wilson	
> 0	< 0	hyperbola	q-quadratic	Askey-Wilson	q real
< 0	< 0	ellipse	q-quadratic	Askey-Wilson	q = 1

TABLE 2. The non-uniform lattices of quadratic type.

functions [39] (see also Section 15.10 of [12]). However for the purposes of studying the Askey table we will not pursue this direction.

For the quadratic class of lattices the parameterisation on s can be made explicit through the trigonometric/hyperbolic functions or their degenerations so we can employ a parameterisation such that $\iota_{-}(x(s)) = y(s) = x(s - 1/2)$ and $\iota_{+}(x(s)) = y(s + 1) = x(s + 1/2)$. We denote the totality of lattice points by $G[x] := \{x(s) : s \in \mathbb{Z}\}$ with the point x(0) = x as the *basal point*, and of the dual lattice by $\tilde{G}[x] := \{x(s) : s \in \mathbb{Z} + \frac{1}{2}\}.$

Having established the basic properties of the divided-difference operators we can deduce key elements of their calculus. A consequence of the general definition of the divided-difference operators are the following identities

(i) the product or Leibniz formulae

(2.10)
$$\mathbb{D}_x fg = \mathbb{D}_x f\mathbb{M}_x g + \mathbb{M}_x f\mathbb{D}_x g,$$

(2.11)
$$\mathbb{M}_x fg = \mathbb{M}_x f\mathbb{M}_x g + \frac{1}{4}\Delta y^2 \mathbb{D}_x f\mathbb{D}_x g$$

(ii) the inverse formulae

(2.12)
$$\mathbb{M}_x \frac{1}{f} = \frac{\mathbb{M}_x f}{E_x^+ f E_x^- f}, \qquad \mathbb{D}_x \frac{1}{f} = -\frac{\mathbb{D}_x f}{E_x^+ f E_x^- f}$$

(iii) and the commutativity formulae

(2.13)
$$\mathbb{M}_u \mathbb{M}_x = \mathbb{M}_x \mathbb{M}_u, \qquad \mathbb{M}_u \mathbb{D}_x = \mathbb{D}_x \mathbb{M}_u, \qquad \mathbb{D}_u \mathbb{D}_x = \mathbb{D}_x \mathbb{D}_u.$$

The other side of our divided-difference calculus concerns the definition and properties of analogs to integrals. We define the \mathbb{D} -Integral of a function defined on the x-lattice $f : G[x] \to \mathbb{C}$ with basal point x_0 by the Riemann sum over the lattice points

(2.14)
$$I[f](x_0) = \int_G \mathbb{D}x \, f(x) := \sum_{s \in \mathbb{Z}} (\iota_+(s) - \iota_-(s)) f(x(s)) = \sum_{s \in \mathbb{Z}} \Delta y(x_s) f(x_s),$$

where the sum is either a finite subset of \mathbb{Z} , namely $\{0, \ldots, \mathfrak{N}\}$, or $\mathbb{Z}_{\geq 0}$, \mathbb{Z} . This definition reduces to the usual definition of the difference integral and the Thomae-Jackson *q*-integrals in the canonical forms of the linear and *q*-linear lattices respectively. A number of properties flow from this definition -

(i) an analog of the fundamental theorem of calculus

(2.15)
$$\int_{x_0 \le x_s \le x_{\mathfrak{N}}} \mathbb{D}x \, \mathbb{D}_x f(x) = f(E_x^+ x_{\mathfrak{N}}) - f(E_x^- x_0),$$

(ii) an analog of integration by parts for two functions f(x), g(x)(2.16) $\int_{x_0 \le x_s \le x_{\mathfrak{N}}} \mathbb{D}x f(x) \mathbb{D}_x g(x) = -\int_{x_0 \le x_s \le x_{\mathfrak{N}}} \mathbb{D}(E_x^+ x) \mathbb{D}_x f(E_x^+ x) g(E_x^+ x) + f(E_x^{+2} x_{\mathfrak{N}}) g(E_x^+ x_{\mathfrak{N}}) - f(x_0) g(E_x^- x_0),$

(iii) and the parameterisation "invariance" property

(2.17)
$$\int_{x_0 \le x_s \le x_{\mathfrak{N}}} \mathbb{D}x \, f(x) = \int_{x_0 \le x_s \le x_{\mathfrak{N}}} \mathbb{D}(E_x^{+2}x) \, f(E_x^{+2}x) + \Delta y(x_0) f(x_0) - \Delta y(E_x^{+2}x_{\mathfrak{N}}) f(E_x^{+2}x_{\mathfrak{N}}),$$

(2.18)
$$= \int_{x_0 \le x_s \le x_{\mathfrak{N}}} \mathbb{D}(E_x^{-2}x) \, f(E_x^{-2}x) - \Delta y(E_x^{-2}x_0) f(E_x^{-2}x_0) + \Delta y(x_{\mathfrak{N}}) f(x_{\mathfrak{N}}).$$

We will apply our theory to the case of the q-quadratic lattice and the Askey-Wilson divided-difference calculus, and in order to simplify the description and to conform to convention we will employ the canonical, that is to say the centred and symmetrised forms of the lattice and the divided-difference operators. Let us define the base $q = \exp(2i\eta)$ although we will not restrict ourselves to q-domains such as $0 < \Re(q) < 1$ except to avoid special degenerate cases and to ensure convergence. Consider the projection map from the unit circle $z = e^{i\theta}$, $\theta \in [-\pi, \pi)$ onto [-1, 1] by $x = \frac{1}{2}(z + z^{-1}) = \cos \theta \in [-1, 1]$. We denote the unit circle by \mathbb{T} and the unit open disc by \mathbb{D} . The inverse of the projection map defines a two-sheeted Riemann surface, one of which corresponds to the interior of the unit circle, and the other to the exterior. Thus we take x-plane to be cut along [-1, 1] and will usually give results for the second sheet i.e. when $|x| \to \infty$ as $z \to \infty$. In the symmetrised and canonical form of the lattice we have $\mathcal{A} = \mathcal{C}$, arbitrary and non-zero, $\mathcal{B} = -\cos \eta \mathcal{A}$, $\mathcal{D} = \mathcal{E} = 0$, $\mathcal{F} = -\sin^2 \eta \mathcal{A}$, and $\theta = 2s\eta$. Define the shift operators E_x^{\pm} by $E_x^{\pm} f(x) = f(\frac{1}{2}[q^{1/2}z + q^{-1/2}z^{-1}])$ and set $y_{\pm} = E_x^{\pm} x$. This implies that

(2.19)
$$y_{+} + y_{-} = (q^{1/2} + q^{-1/2})x = 2\cos\eta x,$$

(2.20)
$$\Delta y := y_+ - y_- = \frac{1}{2}(q^{1/2} - q^{-1/2})(z - z^{-1}) = -2\sin\eta\sin\theta,$$

(2.21)
$$\Delta y^2 = (q^{1/2} - q^{-1/2})^2 (x^2 - 1),$$

(2.22)
$$y_+ y_- = x^2 + \frac{1}{4} (q^{1/2} - q^{-1/2})^2 = x^2 - \sin^2 \eta$$

The Askey-Wilson divided-difference operators are defined as

(2.23)
$$\mathbb{D}_{x}f(x) = \frac{f(\frac{1}{2}[q^{1/2}z + q^{-1/2}z^{-1}]) - f(\frac{1}{2}[q^{-1/2}z + q^{1/2}z^{-1}])}{\frac{1}{2}(q^{1/2} - q^{-1/2})(z - z^{-1})}$$
$$\mathbb{M}_{x}f(x) = \frac{1}{2}\left[f(\frac{1}{2}[q^{1/2}z + q^{-1/2}z^{-1}]) + f(\frac{1}{2}[q^{-1/2}z + q^{1/2}z^{-1}])\right]$$

There is an explicit parameterisation of the q-quadratic lattice

(2.24)
$$x(s) = \frac{1}{2}(q^s + q^{-s}) = \cos(2\eta s),$$

(2.25)
$$y_{\pm}(s) = \frac{1}{2}(q^{s\pm 1/2} + q^{-s\mp 1/2}) = \cos(2\eta[s\pm\frac{1}{2}]).$$

Here the direct lattice is $G[x = \frac{1}{2}(a + a^{-1})] = \{\frac{1}{2}(q^{r/2}a + q^{-r/2}a^{-1}) : r \in 2\mathbb{Z}\}$. For |q| = 1 and η not a rational multiple of π then the lattice densely fills the interval [-1, 1]. If η is a rational multiple of π then one has the root of unity case $q^N = 1$ and a finite lattice. In the generic case we will assume that we are dealing with functions f(x) in the class where

(2.26)
$$\int \mathbb{D}x f(x) = \frac{\sin \eta}{\eta} \int_{-1}^{1} dx f(x),$$

is applicable. The reader should note that we will not distinguish a function of x, f(x), from the function of z, $\check{f}(z) = f(\frac{1}{2}[z+z^{-1}])$ as done by some authors, and it should be clear from the context which is meant.

3. ORTHOGONAL POLYNOMIAL SYSTEMS ON THE NON-UNIFORM LATTICE

3.1. General orthogonal polynomial systems. Our study requires the revision of a number of standard results in orthogonal polynomial theory [92, 30, 43] so we recount our formulation. Let $\{l_n(x;a)\}_{n=0}^{\infty}$ be a polynomial basis of $L^2(w(x)\mathbb{D}x, G)$, where l_n is of exact degree n and the support is $G = \{E_x^{+k}x : k \in 2\mathbb{Z}\}$ or if finite $G = \{x_0, \ldots, x_{\mathfrak{N}}\}$ and a denotes the set of parameters characterising the lattice. The appropriate canonical basis is dependent on the lattice type through the general requirements that

(i) l_n is of precise degree n so that $l_n(x; a) = g_n(a)x^n + O(x^{n-1})$ with $g_n \neq 0$,

(ii) \mathbb{D}_x is an exact lowering operator in this basis

$$\mathbb{D}_x l_n(x;a) = c_n(a) l_{n-1}(x;a'),$$

where c_n is constant with respect to x and the transformed parameter set a' is related to the original a depending on the lattice type.

We will also require the linearisation formula $xl_n(x; a) = d_n(a)l_{n+1}(x; a) + e_n(a)l_n(x; a)$. A general solution to the two requirements above is the following product expression

(3.2)
$$l_n(x;a) = g_n(a) \prod_{k=0}^{n-1} [x - (E_x^+)^{2k} x(a)],$$

where the basal point x(a) is parameterised by a. We note that for some lattices the limit $\lim_{n\to\infty} l_n(x;a)$ exists and in this case we denote it by $l_{\infty}(x;a)$. For the classes of quadratic non-uniform lattices the basis choices are tabulated in Table 3.

DDO	Lattice Type	Basis l_r	Notes
$\frac{d}{dx}$	continuous	x^r	$c_r = r$
Δ_x	linear	$x^{(r)} = \prod_{k=0}^{r-1} (x-k) = \frac{\Gamma(x+1)}{\Gamma(x-r+1)}$	$c_r = r$
D_q	q-linear	$(ax;q)_r = \prod_{k=0}^{r-1} (1 - aq^k x) = \frac{(ax;q)_{\infty}}{(aq^r x;q)_{\infty}}$	$c_r = -\frac{1 - aq^r}{q - 1}$ $a' = qa$
W	quadratic	$\prod_{k=0}^{r-1} \left[x + (k+a)^2 \right] = \frac{\Gamma(r+a-i\sqrt{x})\Gamma(r+a+i\sqrt{x})}{\Gamma(a-i\sqrt{x})\Gamma(a+i\sqrt{x})}$	$c_r = r$ $a' = a + \frac{1}{2}$
\mathbb{D}_x	q-quadratic	$(az, az^{-1}; q)_r = \frac{(az, az^{-1}; q)_\infty}{(aq^r z, aq^r z^{-1}; q)_\infty}$	$c_{r} = -2a \frac{q^{r} - 1}{q - 1}$ $a' = q^{1/2}a$ $g_{r} = (-2a)^{r} q^{\frac{1}{2}r(r-1)}$ $d_{r} = -\frac{1}{2aq^{r}}$ $e_{r} = \frac{1}{2}(aq^{r} + a^{-1}q^{-r})$

TABLE 3. Canonical Bases for the non-uniform lattices of quadratic type.

Consider the general orthogonal polynomial system $\{p_n(x)\}_{n=0}^{\infty}$ defined by the orthogonality relations

(3.3)
$$\int_{G} \mathbb{D}x \ w(x)p_{n}(x) \ l_{m}(x;b) = \begin{cases} 0 & 0 \le m < n \\ h_{n}(b) & m = n \end{cases}, \quad n \ge 0,$$

with G denoting the support of the weight w(x). Our system of orthogonal polynomials and their associated functions (to be defined in (3.21)) have a distinguished singular point at $x = \infty$ and possess expansions about this point which can characterise solutions uniquely. This is related to the fact that orthogonal polynomials

are the denominators of single point Padé approximants and that point is conventionally set at $x = \infty$. We give special notation for the coefficients of x^n and x^{n-1} in $p_n(x)$,

(3.4)
$$p_n(x) = \gamma_n x^n + \gamma_{n,1} x^{n-1} + \dots, \quad n \ge 0.$$

The corresponding *monic* polynomials are then $\pi_n(x) = \gamma_n^{-1} p_n(x)$ given that $n \ge 0$. A consequence of the orthogonality relation is the three term recurrence relation

(3.5)
$$a_{n+1}p_{n+1}(x) = (x - b_n)p_n(x) - a_n p_{n-1}(x), \quad n \ge 0.$$

and we consider the set of orthogonal polynomials with initial values $p_{-1} = 0$ and $p_0 = \gamma_0$. The three term recurrence coefficients are related to the leading and sub-leading polynomial coefficients by [92, 30]

(3.6)
$$a_n = \frac{\gamma_{n-1}}{\gamma_n}, \quad b_n = \frac{\gamma_{n,1}}{\gamma_n} - \frac{\gamma_{n+1,1}}{\gamma_{n+1}}, \quad n \ge 1.$$

The initial values of the recurrence coefficients are

(3.7)
$$b_0 = -\frac{\gamma_{1,1}}{\gamma_1}, \quad \gamma_{0,1} = 0.$$

and where a_0 is not fixed by the initial polynomials but rather by the initial associated functions (see after (3.22).

The orthogonality relation (3.3) is derived from the linear functional on the space of polynomials $\mathcal{L} : p \in \Pi \mapsto \mathbb{C}$ and we employ our basis polynomials as an expansion basis although not necessarily with the same parameter as above

(3.8)
$$p_n(x) = \sum_{k=0}^n c_{n,k}(a) l_k(x;a) \quad n \ge 0.$$

Consequently we define the moments $\{m_{j,k}\}_{j,k=0,1,...,\infty}$ of the weight as the action of this functional on products of the basis polynomials, defined as

(3.9)
$$m_{j,k}(b,a) := \int_{G} \mathbb{D}x \ w(x) \ l_{j}(x;b) l_{k}(x;a), \quad j,k \ge 0.$$

Central objects in our theory are the Moment determinants

(3.10)
$$\Delta_n := \det[m_{j,k}]_{j,k=0,\dots,n-1}, \quad n \ge 1, \quad \Delta_0 := 1,$$

and

(3.11)
$$\Sigma_{n,j} := \det \begin{pmatrix} m_{0,0} & \cdots & m_{0,j-1} & [] & m_{0,j+1} & \cdots & m_{0,n} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ m_{n-1,0} & \cdots & m_{n-1,j-1} & [] & m_{n-1,j+1} & \cdots & m_{n-1,n} \end{pmatrix}, \quad n \ge 1, j = 0, \dots n - 1.$$

defined in terms of the moments above. Obviously $\Delta_n = \Sigma_{n,n}$ and we set $\Sigma_{0,0} := 0$. The expansion coefficients are given in terms of these determinants

(3.12)
$$c_{n,j}(a) = (-)^{n+j} h_n(b) \frac{\Sigma_{n,j}}{\Delta_{n+1}}, \quad c_{n,n}(a) = h_n(b) \frac{\Delta_n}{\Delta_{n+1}}.$$

It follows from (3.3) that

$$\int_G \mathbb{D}x \ w(x)[p_n(x)]^2 = c_{n,n}(b)h_n(b), \quad n \ge 0,$$

and thus for $p_n(x)$ to be normalised as well as orthogonal we set $c_{n,n}(a)h_n(a) = 1$. We have moment determinant representations of the polynomials

(3.13)
$$p_n(x) = \frac{c_{n,n}(a)}{\Delta_n} \det \begin{pmatrix} m_{0,0} & \cdots & m_{0,j} & \cdots & m_{0,n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ m_{n-1,0} & \cdots & m_{n-1,j} & \cdots & m_{n-1,n} \\ l_0 & \cdots & l_j & \cdots & l_n \end{pmatrix}, \quad n \ge 0.$$

The three-term recurrence coefficients are related to these determinants

(3.14)
$$a_n^2 = d_{n-1}(a)d_{n-1}(b)\frac{\Delta_{n+1}\Delta_{n-1}}{\Delta_n^2}, \quad n \ge 1,$$

(3.15)
$$b_n = e_n(a) + d_n(a) \frac{\Sigma_{n+1,n}}{\Delta_{n+1}} - d_{n-1}(a) \frac{\Sigma_{n,n-1}}{\Delta_n}, \quad n \ge 0,$$

(3.16)
$$\gamma_n^2 = g_n(a)g_n(b)\frac{\Delta_n}{\Delta_{n+1}}, \quad n \ge 0,$$

where each coefficient is independent of the choices of a, b as can be easily verified from their determinantal definitions given previously.

Another set of polynomial solutions to the three term recurrence relation are the associated polynomials ${p_n^{(1)}(x)}_{n=0}^{\infty}$, defined by

(3.17)
$$p_{n-1}^{(1)}(x) := \int_{G} \mathbb{D}y \ w(y) \frac{p_n(y) - p_n(x)}{y - x}, \quad n \ge 0.$$

In particular these polynomials satisfy

(3.18)
$$a_{n+1}p_n^{(1)}(x) = (x - b_n)p_{n-1}^{(1)}(x) - a_n p_{n-2}^{(1)}(x),$$

with the initial conditions $p_{-1}^{(1)}(x) = 0$, $p_0^{(1)}(x) = m_{0,0}\gamma_1$. Note the shift by one decrement in comparison to the three-term recurrence (3.5) for the polynomials $\{p_n(x)\}_{n=0}^{\infty}$. We also need the definition of the *Stieltjes* function

(3.19)
$$f(x) \equiv \int_{G} \mathbb{D}y \ \frac{w(y)}{x-y}, \quad x \notin G$$

which is a moment generating function in the following sense

(3.20)
$$f(x) = \frac{f_{\infty}(x;a)}{l_{\infty}(x;a)} + \sum_{n=0}^{\infty} \frac{m_{0,n}(a)}{d_n(a)l_{n+1}(x;a)}, \quad x \notin G, \quad x \to \infty,$$

and splits into two parts - one part being a series with inverse basis polynomials and a remainder, which may be absent for some lattices. We define non-polynomial associated functions or functions of the second kind $\{q_n(x)\}_{n=0}^{\infty}$ by

(3.21)
$$q_n(x) := f(x)p_n(x) - p_{n-1}^{(1)}(x), \quad n \ge 0,$$

which also satisfy the three term recurrence relation (3.5), namely

(3.22)
$$a_{n+1}q_{n+1}(x) = (x - b_n)q_n(x) - a_nq_{n-1}(x), \quad n \ge 0,$$

subject to the initial values $q_{-1}(x) = 1/a_0\gamma_0$, $q_0(x) = \gamma_0 f(x)$. The initial value of a_0 is irrelevant and therefore arbitrary in so far as the polynomials are concerned, however many relations for the whole system extend from $n \ge 1$ to include n = 0 if we allow this to be finite, non-zero and satisfying the above initial condition. The associated functions also have a determinantal representation

(3.23)
$$q_n(x) = \frac{c_{n,n}(a)}{\Delta_n} \det \begin{pmatrix} m_{0,0} & \cdots & m_{0,j} & \cdots & m_{0,n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ m_{n-1,0} & \cdots & m_{n-1,j} & \cdots & m_{n-1,n} \\ f_0 & \cdots & f_j & \cdots & f_n \end{pmatrix}, \quad n \ge 0,$$

where

(3.24)
$$f_j(x;a) := \int_G \mathbb{D}y \ w(y) \frac{l_j(y;a)}{x-y}, \quad j \ge 1, \quad f_0 = g_0 f_0$$

Likewise this function has an expansion analogous to (3.20)

(3.25)
$$f_j(x;b) = \frac{f_{\infty,j}(x;b)}{l_{\infty}(x;a)} + \sum_{n=0}^{\infty} \frac{m_{j,n}(b,a)}{d_n(a)l_{n+1}(x;a)}, \quad x \notin G, \quad x \to \infty.$$

The polynomials and their associated functions satisfy the Casoratian relation

(3.26)
$$p_n(x)q_{n-1}(x) - p_{n-1}(x)q_n(x) = \frac{1}{a_n}, \quad n \ge 0.$$

Central to our analysis is a composite of polynomial and non-polynomial solutions of (3.5), the 2×2 matrix variable

(3.27)
$$Y_n(x) = \begin{pmatrix} p_n(x) & \frac{q_n(x)}{w(x)} \\ p_{n-1}(x) & \frac{q_{n-1}(x)}{w(x)} \end{pmatrix}, \quad n \ge 0.$$

We will refer to this as the orthogonal polynomial system (OPS). From (3.26) we note that

(3.28)
$$\det Y_n(x) = \frac{1}{a_n w(x)}, \quad n \ge 0.$$

The three term recurrence is then recast as the matrix equation

(3.29)
$$Y_{n+1}(x) = K_n Y_n(x), \quad n \ge 0,$$

with the *recurrence matrix* given by

(3.30)
$$K_n(x) = \frac{1}{a_{n+1}} \begin{pmatrix} x - b_n & -a_n \\ a_{n+1} & 0 \end{pmatrix}, \quad \det K_n = \frac{a_n}{a_{n+1}}, \quad n \ge 0.$$

A well known consequence of (3.5) are the Christoffel-Darboux summation formulae

(3.31)
$$\sum_{j=0}^{n-1} p_j(x) p_j(y) = a_n \frac{[p_n(x)p_{n-1}(y) - p_{n-1}(x)p_n(y)]}{x - y}, \quad n \ge 0,$$

(3.32)
$$\sum_{j=0}^{n-1} q_j(x) p_j(y) = a_n \frac{[q_n(x)p_{n-1}(y) - q_{n-1}(x)p_n(y)]}{x - y} + \frac{1}{x - y}, \quad n \ge 0,$$

(3.33)
$$\sum_{j=0}^{n-1} q_j(x)q_j(y) = a_n \frac{[q_n(x)q_{n-1}(y) - q_{n-1}(x)q_n(y)]}{x - y} - \frac{f(x) - f(y)}{x - y}, \quad n \ge 0.$$

If one is only interested in the leading orders of the large x expansion rather than a systematic expansion then it is convenient to employ an expansion in monomials rather than in basis functions. Extending (3.4) we have expansions about the fixed singularity at $x = \infty$

(3.34)
$$p_n(x) = \gamma_n \left[x^n - \left(\sum_{i=0}^{n-1} b_i \right) x^{n-1} + \left(\sum_{0 \le i < j < n} b_i b_j - \sum_{i=1}^{n-1} a_i^2 \right) x^{n-2} + \mathcal{O}(x^{n-3}) \right],$$

valid for $n \ge 1$, while for the associated functions

(3.35)
$$q_n(x) = \gamma_n^{-1} \left[x^{-n-1} + \left(\sum_{i=0}^n b_i \right) x^{-n-2} + \left(\sum_{0 \le i \le j \le n} b_i b_j + \sum_{i=1}^{n+1} a_i^2 \right) x^{-n-3} + \mathcal{O}(x^{-n-4}) \right],$$

valid for $n \ge 0$.

3.2. *q*-Quadratic lattice. In Sections 5 and 7 we intend to apply our theory to the *q*-quadratic lattice and will draw upon numerous properties of the corresponding basis, which we discuss here. Firstly we recall the analytic continuation of $\phi_n(x; a)$ which is

(3.36)
$$\phi_r(x;a) = \frac{(az, az^{-1}; q)_{\infty}}{(aq^r z, aq^r z^{-1}; q)_{\infty}},$$

for all $r \in \mathbb{C}$ but subject to |q| < 1. We will employ the shorthand notation $(az^{\pm 1}; q)_{\infty} = (az, az^{-1}; q)_{\infty}$. Implicit in the above formula is the elliptic-like function $\phi_{\infty}(x; a)$ which has meaning for all $a, x \in \mathbb{C}$ for |q| < 1. The action of the divided-difference operators are

(3.37)
$$\mathbb{D}_x \phi_r(x;a) = 2a \frac{1-q^r}{q-1} \phi_{r-1}(x;q^{1/2}a)$$

(3.38)
$$\mathbb{M}_x \phi_r(x;a) = \frac{1}{2} (1+q^{-r}) \phi_r(x;q^{1/2}a) + \frac{1}{2} (1-q^{-r}) (1-a^2 q^{2r-1}) \phi_{r-1}(x;q^{1/2}a),$$

(3.39)
$$\mathbb{D}_x \phi_{\infty}(x;a) = \frac{2a}{q-1} \phi_{\infty}(x;q^{1/2}a)$$

(3.40)
$$\mathbb{M}_x \phi_\infty(x; a) = \phi_\infty(x; q^{-1/2}a).$$

The expansion theorem of Ismail [40, 41] states that

Proposition 3.1. Let p be a polynomial of degree n, then

(3.41)
$$p(x) = \sum_{k=0}^{n} p_k \phi_k(x;a)$$

for any $a \in \mathbb{C}$ where

(3.42)
$$p_k = \frac{(q-1)^k}{(2a)^k (q;q)_k} q^{-k(k-1)/4} (\mathbb{D}_x^k p)(x_k)$$

with $x_k = \frac{1}{2}(q^{k/2}a + q^{-k/2}a^{-1}).$

In these works, see (2.2) and the proof of Theorem 1.1 in [40], we also have the change of base formula

(3.43)
$$\phi_n(x;b) = \sum_{k=0}^n {n \brack k}_q (abq^k, b/a; q)_{n-k} \left(\frac{b}{a}\right)^k \phi_k(x;a),$$

where we use the standard definition of the q-binomial coefficient. This result allows us to derive the following linearisation formula.

Lemma 3.1. The product of two basis polynomials with the same base a has the following expansion in terms of the same basis

$$(3.44) \quad \phi_k(x;a)\phi_l(x;a) = (-)^{k+l}q^{-kl} \sum_{m=\max(k,l)}^{k+l} (-)^m (a^2q^m;q)_{k+l-m} \frac{(q;q)_l(q;q)_k}{(q;q)_{k+l-m}(q;q)_{m-k}(q;q)_{m-l}} \phi_m(x;a),$$

for all $k, l \in \mathbb{N}$.

Lastly we have the Cauchy expansion formula of Ismail and Stanton [47].

Proposition 3.2. The Cauchy kernel has the expansion

(3.45)
$$\frac{1}{y-x} = \frac{1}{y-x} \frac{\phi_{\infty}(x;a)}{\phi_{\infty}(y;a)} - 2a \sum_{n=0}^{\infty} \frac{\phi_n(x;a)}{\phi_{n+1}(y;a)} q^n,$$

for all y such that $y \neq x$ and $\phi_{\infty}(y;a) \neq 0$. The expansion also holds for $y = y_0$ with $\phi_{\infty}(y_0;a) = 0$ but $y_0 \neq x$ in the sense that the left-hand side $(y_0 - x)^{-1}$ equals the limit of the right-hand side as $y \to y_0$.

This implies the following expansion of the Stieltjes function.

Corollary 3.1. The Stieltjes function has the following expansion as $x \to \infty$ with $x \neq x_k(a)$, with $k \in 2\mathbb{Z}$ (*i.e.* $\phi_{\infty}(x; a) \neq 0$)

(3.46)
$$f(x) = \frac{f_{\infty}(x)}{\phi_{\infty}(x;a)} - 2a \sum_{n=0}^{\infty} \frac{q^n}{\phi_{n+1}(x;a)} m_{0,n}(a),$$

where

(3.47)
$$f_{\infty}(x) = \int_{G} \mathbb{D}y \ w(y) \frac{\phi_{\infty}(y;a)}{x-y}$$

Conforming with standard notation [31] we define the basic hypergeometric function $r_{+1}\varphi_r$ by the series

(3.48)
$$r_{+1}\varphi_r \begin{bmatrix} a_1, a_2, \dots, a_{r+1} \\ b_1, b_2, \dots, b_r \end{bmatrix} = \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_{r+1}; q)_n}{(q, b_1, b_2, \dots, b_r; q)_n} z^n,$$

which is convergent for |z| < 1. The very-well-poised basic hypergeometric function $_{r+1}W_r$ is a specialisation of the above

$$(3.49) \quad {}_{r+1}W_r(a_1; a_4, a_5, \dots, a_{r+1}; q, z) = {}_{r+1}\varphi_r \left[\begin{array}{cccc} a_1, & q\sqrt{a_1}, & -q\sqrt{a_1}, & a_4, & \dots, & a_{r+1} \\ \sqrt{a_1}, & -\sqrt{a_1}, & qa_1/a_4, & \dots, & qa_1/a_{r+1} \end{array} ; q, z \right]$$

so that $qa_1 = a_2b_1 = a_3b_2 = \cdots = a_{r+1}b_r$ and $a_2 = q\sqrt{a_1}$, $a_3 = -q\sqrt{a_1}$. The $_{r+1}\varphi_r$ or $_{r+1}W_r$ functions may also be *balanced*, whereby z = q and $\prod_{j=1}^r b_j = q \prod_{j=1}^{r+1} a_j$.

4. Spectral Differences

In this section we lay out the structures of the spectral divided-difference operator in the context of the orthogonal polynomial system for a general lattice type. Our starting point is the notion of the \mathbb{D} -semiclassical weight, as given by the following definition of Magnus [60].

Definition 4.1 ([60]). Let the \mathbb{D} -semi-classical weight satisfy

(4.1)
$$W\mathbb{D}_x w = 2V\mathbb{M}_x w$$

or equivalently

(4.2)
$$\frac{w(y_+)}{w(y_-)} = \frac{W + \Delta yV}{W - \Delta yV}(x),$$

for W(x), V(x) irreducible polynomials, which we will call spectral data polynomials. Furthermore we assume $W \pm \Delta y V \neq 0$ for all $x \in G$. For minimal degrees of W, V this is the analog of the Pearson equation.

Remark 4.1. On the finite lattice $x \in \{x_0, \ldots, x_{\mathfrak{N}}\}$ we naturally require $w(x) \neq 0$, however we will impose upper and lower terminating conditions

(4.3)
$$w(E_x^{+2}x_{\mathfrak{N}}) = w(E_x^{-2}x_0) = 0,$$

respectively. These are the conditions analogous to Eqs. (2.3.2) and (3.3.3) in Nikiforov, Suslov and Uvarov [78]. To be consistent with (4.2) this implies

(4.4)
$$(W + \Delta yV)(E_x^+ x_{\mathfrak{N}}) = (W - \Delta yV)(E_x^- x_0) = 0.$$

Remark 4.2. In a series of works Suslov and collaborators [5, 91, 6, 84, 83] have studied the Pearson equation for all of the lattices admissible in the classification, and sought solutions for the weight functions given suitable polynomials for W, V. However they limited their choices to examples of minimal degree for W, Vwhich made contact with those weights found in the Askey table. In terms of our own variables those found in [91] are given by

(4.5)
$$\sigma = E_x^-(W - \Delta yV), \qquad \tau = \frac{E_x^+(W + \Delta yV)(x) - E_x^-(W - \Delta yV)(x)}{\Delta y}.$$

From (3.20) we recognise f(x) as a moment generating function and a key element in our theory are the systems of linear divided-difference equations satisfied by the moments. Before we state such systems we need to note the following result.

Lemma 4.1. Let $l_k(x; a)$ be a canonical basis polynomial. For any $k \in \mathbb{Z}_{\geq 0}$ and $a \in \mathbb{C}$ the weight satisfies the integral equation

(4.6)
$$\int_{x_0 \le x \le x_{\mathfrak{N}}} \mathbb{D}x \, \frac{w(x)}{\Delta y(x)} \left\{ \left(l_k [W + \Delta yV] \right) \left(E_x^+ x \right) - \left(l_k [W - \Delta yV] \right) \left(E_x^- x \right) \right\} = 0.$$

Proof. We will prove this statement accounting for the boundary terms explicitly. We establish two preliminary identities first, namely

(4.7)
$$\int_{x_0 \le x \le x_{\mathfrak{N}}} \mathbb{D}x \, w(x) \left\{ \frac{1}{E_x^+ \Delta y} E_x^+ \left(l_k [W + \Delta y V] \right) - \frac{E_x^{-2} \Delta y}{\Delta y (E_x^- \Delta y)} E_x^- \left(l_k [W - \Delta y V] \right) \right\} = 0.$$

(4.8)
$$\int_{x_0 \le x \le x_{\mathfrak{N}}} \mathbb{D}x \, w(x) \left\{ \frac{E_x \ \Delta y}{\Delta y(E_x \ \Delta y)} E_x^+ \left(l_k [W + \Delta yV] \right) - \frac{1}{E_x \ \Delta y} E_x^- \left(l_k [W - \Delta yV] \right) \right\} = 0,$$

with $l_k := l_k(x; a)$. Taking the first of these we compute

$$\begin{split} 0 &= \int_{x_0 \le x \le x_{\Re}} \mathbb{D}x \, l_k(E_x^+ x) (W \mathbb{D}_x w - 2V M_x w) (E_x^+ x), \\ &= \sum_{0 \le s \le \Re} \Delta y(x_s) l_k(E_x^+ x_s) W(E_x^+ x_s) \frac{w(E_x^{+2} x_s) - w(x_s)}{\Delta y(E_x^+ x_s)} \\ &\quad - \sum_{0 \le s \le \Re} \Delta y(x_s) l_k(E_x^+ x_s) V(E_x^+ x_s) \left[w(E_x^{+2} x_s) + w(x_s) \right], \\ &= \sum_{0 \le s \le \Re} \frac{\Delta y(E_x^{-2} x_s)}{\Delta y(E_x^- x_s)} l_k(E_x^- x_s) W(E_x^- x_s) w(x_s) - \sum_{0 \le s \le \Re} \frac{\Delta y(x_s)}{\Delta y(E_x^+ x_s)} l_k(E_x^+ x_s) W(E_x^+ x_s) w(x_s) \\ &\quad + \frac{\Delta y(x_{\Re})}{\Delta y(E_x^- x_{\Re})} l_k(E_x^+ x_{\Re}) W(E_x^+ x_{\Re}) w(E_x^{+2} x_{\Re}) - \frac{\Delta y(E_x^{-2} x_0)}{\Delta y(E_x^- x_0)} l_k(E_x^- x_0) W(E_x^- x_0) w(x_0) \\ &\quad - \sum_{0 \le s \le \Re} \Delta y(E_x^{-2} x_s) l_k(E_x^- x_s) V(E_x^- x_s) w(x_s) - \sum_{0 \le s \le \Re} \Delta y(x_s) l_k(E_x^+ x_{\Re}) V(E_x^+ x_{\Re}) w(E_x^{+2} x_{\Re}) + \Delta y(E_x^{-2} x_0) l_k(E_x^- x_0) V(E_x^- x_0) w(x_0), \\ &= \sum_{0 \le s \le \Re} \Delta y(x_s) w(x_s) \left\{ - \left(l_k \left[\frac{W}{\Delta y} + V \right] \right) (E_x^+ x_s) + \frac{\Delta y(E_x^{-2} x_0)}{\Delta y(x_s)} \left(l_k \left[\frac{W}{\Delta y} - V \right] \right) (E_x^- x_s) \right\} \\ &\quad + \frac{\Delta y(x_{\Re})}{\Delta y(E_x^+ x_{\Re})} l_k(E_x^+ x_{\Re}) w(E_x^{+2} x_{\Re}) [W - \Delta yV](E_x^+ x_{\Re}) - \frac{\Delta y(E_x^{-2} x_0)}{\Delta y(E_x^- x_0)} l_k(E_x^- x_0) w(x_0) [W - \Delta yV](E_x^- x_0) d_x(E_x^- x_0) w(x_0)] W(x_x + 2 x_{\Re}) d_x(E_x^- x_0) d_x(x_x + 2 x_{\Re}) [W - \Delta yV](E_x^+ x_{\Re}) - \frac{\Delta y(E_x^{-2} x_0)}{\Delta y(E_x^- x_0)} l_x(E_x^- x_0) w(x_0)[W - \Delta yV](E_x^- x_0) d_x(x_x + 2 x_{\Re}) d_x(x_$$

where the boundary terms arise from using the integration by parts and the change of variables $x_s \mapsto E_x^{\pm 2} x_s$ formulae. Both boundary terms are in fact zero due to the upper and lower terminating conditions. The second formula (4.8) is just a variation of the first with the integrand evaluated at $E_x^- x_s$ instead. Now for a general quadratic lattice

$$\frac{\Delta y + E_x^{\pm 2} \Delta y}{E_x^{\pm} \Delta y} = -2\frac{\mathcal{B}}{\mathcal{A}},$$

so we simply add (4.7) and (4.8) and note that a common, constant factor appears in the integrand of precisely this form. Then (4.6) immediately follows. \Box

Remark 4.3. An equivalent system for general lattices has been derived by Suslov [91] in the special case of k = 0 of (4.6).

We also need to express $W \pm \Delta y V$ in terms of canonical basis polynomials and formulate the following definitions for the coefficients $\kappa_{N,l}(a), \delta_{N,l}(a)$

(4.9)
$$E_x^+(W + \Delta yV)(x) + E_x^-(W - \Delta yV)(x) =: 2\sum_{k=0}^N \kappa_{N,k}(a)l_k(x;a),$$

(4.10)
$$E_x^+(W + \Delta yV)(x) - E_x^-(W - \Delta yV)(x) =: \Delta y \sum_{k=0}^{N-1} \delta_{N,k}(a) l_k(x;a)$$

Here N is the cutoff determined by the degrees of W, V and the lattice type.

Corollary 4.1. Let us assume that $b' \neq a$, where b, b' are related as a, a' are by (3.1). For all $k \in \mathbb{Z}_{>0}$ the moments $m_{k,l}(b', a)$ are characterised by the linear, homogeneous recurrence relations

(4.11)
$$\left[d_k(b)c_{k+1}(b) + \frac{\mathcal{B}}{\mathcal{A}} d_{k-1}(b')c_k(b) \right] \sum_{l=0}^{N-1} \delta_{N,l}(a)m_{k,l}(b',a) + c_k(b) \sum_{l=0}^{N} \left[\kappa_{N,l}(a) + \left(e_k(b) + \frac{\mathcal{B}}{\mathcal{A}} e_{k-1}(b') + \frac{\mathcal{D}}{\mathcal{A}} \right) \delta_{N,l}(a) \right] m_{k-1,l}(b',a) = 0.$$

For k = 0 the last term is absent and therefore b, b' do not appear.

Proof. We start with the integral formula (4.6) and employ the resolutions of $E_x^+(W + \Delta yV)$ and $E_x^-(W - \Delta yV)$ ΔyV) as given by (4.9) and (4.10). In addition we require expressions for $\mathbb{D}_x l_k(x;b)$ and $\mathbb{M}_x l_k(x;b)$, and use (3.1) for the former and

$$\mathbb{M}_{x}l_{k}(x;b) = \left(d_{k}(b)c_{k+1}(b) + \frac{\mathcal{B}}{\mathcal{A}}d_{k-1}(b')c_{k}(b)\right)l_{k}(x;b') + c_{k}(b)\left(e_{k}(b) + \frac{\mathcal{B}}{\mathcal{A}}e_{k-1}(b') + \frac{\mathcal{D}}{\mathcal{A}}\right)l_{k-1}(x;b'),$$

the latter.

for the latter.

If a specialisation of the parameters b', a is made to effect some cancellation with corresponding factors in the weight then the moments are given by an integral whose integrand has the same structure and (4.11)is now a linear divided-difference equation with respect to the internal parameters of the weight.

Now we continue to develop the consequences of the D-semi-classical weight on the orthogonal polynomial system, and in particular for the Stieltjes function. The following result will be crucial for our approach.

Proposition 4.1. Given a \mathbb{D} -semi-classical class of an orthogonal polynomial system fulfilling (4.1) and the conditions therein then the Stieltjes function satisfies

$$W\mathbb{D}_x f = 2V\mathbb{M}_x f + U,$$

where U(x) in (4.13) is a polynomial in x (as are V(x) and W(x)).

Proof. Starting with the definition (3.19) we compute

$$\mathbb{D}_x f = \int_{u_0 \le u \le u_{\mathfrak{N}}} \mathbb{D}u \, w(u) \mathbb{D}_x \frac{1}{x - u}$$

and make the evaluation of the factor

$$\mathbb{D}_x \frac{1}{x-u} = -\frac{1}{(E_x^+ x - u)(E_x^- x - u)}.$$

However we note that

$$\mathbb{D}_u \frac{1}{x - u} = \frac{1}{(x - E_u^+ u)(x - E_u^- u)},$$

so that

$$\mathbb{D}_x \frac{1}{x-u} = -\frac{(x - E_u^+ u)(x - E_u^- u)}{(E_x^+ x - u)(E_x^- x - u)} \mathbb{D}_u \frac{1}{x-u} = -\mathbb{D}_u \frac{1}{x-u},$$

where we have assumed a symmetric quadratic lattice for simplicity, but point out that our argument could be extended to the general case. We deduce then that

$$\mathbb{D}_{x}f = -\int_{u_{0} \leq u \leq u_{\mathfrak{N}}} \mathbb{D}u \, w(u) \mathbb{D}_{u} \frac{1}{x-u},$$

= $\int_{u_{0} \leq u \leq u_{\mathfrak{N}}} \mathbb{D}(E_{u}^{+}u) \, \frac{1}{x-E_{u}^{+}u} \mathbb{D}_{u}w(E_{u}^{+}u) - \frac{w(E_{u}^{+}^{2}u_{\mathfrak{N}})}{x-E_{u}^{+}u_{\mathfrak{N}}} + \frac{w(u_{0})}{x-E_{u}^{-}u_{0}},$

after employing the formula for integration by parts. We next use the definition of the \mathbb{D} -semi-classical weight (4.1) and perform a sequence of subtractions in the numerator of the integrand

$$\begin{split} W(x)\mathbb{D}_{x}f(x) &= \int_{u_{0} \leq u \leq u_{\mathfrak{N}}} \mathbb{D}(E_{u}^{+}u) \frac{1}{x - E_{u}^{+}u} \mathbb{M}_{u} w(E_{u}^{+}u) \frac{2V(E_{u}^{+}u)}{W(E_{u}^{+}u)} W(x) - \frac{w(E_{u}^{+}^{+}u_{\mathfrak{N}})}{x - E_{u}^{+}u_{\mathfrak{N}}} W(x) + \frac{w(u_{0})}{x - E_{u}^{-}u_{0}} W(x) + \frac{w(u_{0})}{x - E_{u}^{-}u_{0}} W(x) + \frac{w(u_{0})}{x - E_{u}^{+}u_{\mathfrak{N}}} W(x$$

Since the third term on the right-hand side of the previous equation is clearly a polynomial in x we shall focus next on the second term. The last factor of this latter term can be written as

$$\begin{split} \mathbb{M}_{x}f(x) - \int_{u_{0} \leq u \leq u_{\mathfrak{N}}} \mathbb{D}(E_{u}^{+}u) \, \frac{\mathbb{M}_{u}w(E_{u}^{+}u)}{x - E_{u}^{+}u} &= \int_{u_{0} \leq u \leq u_{\mathfrak{N}}} \mathbb{D}u \, w(u) \frac{1}{2} \left\{ \frac{1}{E_{x}^{+}x - u} + \frac{1}{E_{x}^{-}x - u} \right\} \\ &- \frac{1}{2} \int_{u_{0} \leq u \leq u_{\mathfrak{N}}} \mathbb{D}(E_{u}^{+}u) \, \frac{w(E_{u}^{+}^{2}u)}{x - E_{u}^{+}u} - \frac{1}{2} \int_{u_{0} \leq u \leq u_{\mathfrak{N}}} \mathbb{D}(E_{u}^{+}u) \, \frac{w(u)}{x - E_{u}^{+}u}, \end{split}$$

The last two terms of the above equation are

_

$$\begin{split} \int_{u_0 \le u \le u_{\mathfrak{N}}} \mathbb{D}(E_u^+ u) \, \frac{w(E_u^{+2}u)}{x - E_u^+ u} + \int_{u_0 \le u \le u_{\mathfrak{N}}} \mathbb{D}(E_u^+ u) \, \frac{w(u)}{x - E_u^+ u} \\ &= \sum_{0 \le s \le \mathfrak{N}} [u_s - E_u^{-2}u_s] \frac{w(u_s)}{x - E_u^- u_s} + \sum_{0 \le s \le \mathfrak{N}} [E_u^{+2}u_s - u_s] \frac{w(u_s)}{x - E_u^+ u_s} + \Delta y(E_u^+ u_{\mathfrak{N}}) \frac{w(E_u^{+2}u_{\mathfrak{N}})}{x - E_u^+ u_{\mathfrak{N}}} - \Delta y(E_u^- u_0) \frac{w(u_0)}{x - E_u^- u_0}, \\ &= \sum_{0 \le s \le \mathfrak{N}} w(u_s) \left\{ \frac{E_u^{+2}u_s - u_s}{x - E_u^- u_s} + \frac{u_s - E_u^{-2}u_s}{x - E_u^- u_s} \right\} + ", \end{split}$$

where we have used the identity for integrals under the re-parameterisation of the lattice $u \mapsto E_u^{-2}u$ and introducing the additional boundary terms. Combining these results we have

$$\begin{split} \mathbb{M}_{x}f(x) &- \int_{u_{0} \leq u \leq u_{\mathfrak{N}}} \mathbb{D}(E_{u}^{+}u) \, \frac{\mathbb{M}_{u}w(E_{u}^{+}u)}{x - E_{u}^{+}u} \\ &= \frac{1}{2} \sum_{0 \leq s \leq \mathfrak{N}} w(u_{s}) \left\{ \left(E_{u}^{+}u_{s} - E_{u}^{-}u_{s}\right) \left(\frac{1}{E_{x}^{+}x - u_{s}} + \frac{1}{E_{x}^{-}x - u_{s}}\right) - \frac{E_{u}^{+2}u_{s} - u_{s}}{x - E_{u}^{+}u_{s}} - \frac{u_{s} - E_{u}^{-2}u_{s}}{x - E_{u}^{-}u_{s}} \right\} \\ &- \frac{1}{2} \Delta y(E_{u}^{+}u_{\mathfrak{N}}) \frac{w(E_{u}^{+2}u_{\mathfrak{N}})}{x - E_{u}^{+}u_{\mathfrak{N}}} + \frac{1}{2} \Delta y(E_{u}^{-}u_{0}) \frac{w(u_{0})}{x - E_{u}^{-}u_{0}}, \\ &= \frac{1}{2} \sum_{0 \leq s \leq \mathfrak{N}} w(u_{s}) \left\{ (E_{u}^{+}u_{s} - E_{u}^{-}u_{s}) \frac{(E_{x}^{+} + E_{x}^{-})x - 2u_{s}}{(x - E_{u}^{+}u_{s})(x - E_{u}^{-}u_{s})} - \frac{E_{u}^{+2}u_{s} - u_{s}}{x - E_{u}^{+}u_{s}} - \frac{u_{s} - E_{u}^{-2}u_{s}}{x - E_{u}^{-}u_{s}} \right\} + ", \end{split}$$

where we have used the ratio identity again. We observe that the summand appears to have two sets of simple poles at $x = E^{\pm}u_s$, however when we compute their residues they vanish identically, implying that the summand is in reality a polynomial in x. This leaves us with the two sets of boundary terms to account for. We find that their combination is

$$-\frac{w(E_u^{+2}u_{\mathfrak{N}})}{x-E_u^{+}u_{\mathfrak{N}}}[W(x) - \Delta y(E_u^{+}u_{\mathfrak{N}})V(x)] + \frac{w(u_0)}{x-E_u^{-}u_0}[W(x) - \Delta y(E_u^{-}u_0)V(x)] + \frac{w(u_0)}{x-E_u^{-}u_0$$

The first term vanishes under the upper terminating condition whilst the second is a polynomial in x due to the lower terminating condition. In summary this leads us to conclude that $W(x)\mathbb{D}_x f(x) - 2V(x)\mathbb{M}_x f(x)$ is a polynomial in x.

We offer the following analog to the concept of a regular semi-classical weight.

Definition 4.2. A generic or regular \mathbb{D} -semi-classical weight has two properties -

- (i) strict inequalities in the degrees of the spectral data polynomials, i.e. $\deg_x W = M$, $\deg_x V = M 1$ and $\deg_x U = M - 2$, and
- (ii) the lattice generated by any zero of $(W^2 \Delta y^2 V^2)(x)$ does not coincide with the lattice generated by another zero, i.e. for any \tilde{x} such that $(W^2 - \Delta y^2 V^2)(\tilde{x}) = 0$ then $(W^2 - \Delta y^2 V^2)(E_x^{2\mathbb{Z}} \tilde{x}) \neq 0$.

Conversely any weight with data not satisfying these conditions will be termed an *irregular* \mathbb{D} -semi-classical weight. Note that the zeros of W are not relevant and no interpretation as a singularity can be placed upon a zero of $(W^2 - \Delta y^2 V^2)(x)$.

Proposition 4.2 ([63]). The spectral coefficients $W_n(x)$, $\Theta_n(x)$ and $\Omega_n(x)$ are defined in terms of bilinear formulae of the polynomials and associated functions

$$(4.14) \quad \frac{2}{a_n} (2W_n(x) - W(x)) = (W + \Delta yV) \left[p_n(y_+)q_{n-1}(y_-) - p_{n-1}(y_+)q_n(y_-) \right] + (W - \Delta yV) \left[p_n(y_-)q_{n-1}(y_+) - p_{n-1}(y_-)q_n(y_+) \right], \quad n \ge 0,$$

(4.15)
$$\Delta y \Theta_n(x) = (W + \Delta y V) p_n(y_+) q_n(y_-) - (W - \Delta y V) p_n(y_-) q_n(y_+), \quad n \ge 0,$$

$$(4.16) \quad \frac{2\Delta y}{a_n} \left(\Omega_n(x) + V(x)\right) = (W + \Delta y V) \left[p_n(y_+)q_{n-1}(y_-) + p_{n-1}(y_+)q_n(y_-)\right] \\ + \left(W - \Delta y V\right) \left[-p_n(y_-)q_{n-1}(y_+) - p_{n-1}(y_-)q_n(y_+)\right], \quad n \ge 0.$$

For the \mathbb{D} -semi-classical class of weights these coefficients are polynomials in x.

Proof. Our proof will be an extension of Laguerre's reasoning for the differential case. However we first need to appreciate one important fact. Any polynomial in y_- and y_+ is an element of the Ring $\mathbb{C}[x] + \Delta y \mathbb{C}[x]$ because $y_{\pm} = \mathbb{M}_x x \pm \frac{1}{2} \Delta y$ and $\mathbb{M}_x x$ is a linear polynomial in x and Δy^2 is a quadratic polynomial in x. The starting point will be the definition for q_n , but inverted to solve for the Stieltjes function

(4.17)
$$f = \frac{q_n + p_{n-1}^{(1)}}{p_n},$$

for arbitrary $n \ge 0$. Now consider the formulation of (4.13) using this expression

$$0 = (W - \Delta yV)f(y_{+}) - (W + \Delta yV)f(y_{-}) - \Delta yU,$$

=
$$\frac{(W - \Delta yV)q_{n}(y_{+})p_{n}(y_{-}) - (W + \Delta yV)q_{n}(y_{-})p_{n}(y_{+})}{p_{n}(y_{+})p_{n}(y_{-})} + \frac{(W - \Delta yV)p_{n-1}^{(1)}(y_{+})p_{n}(y_{-}) - (W + \Delta yV)p_{n-1}^{(1)}(y_{-})p_{n}(y_{+}) - \Delta yUp_{n}(y_{+})p_{n}(y_{-})}{p_{n}(y_{+})p_{n}(y_{-})}.$$

Now $(W - \Delta yV)p_{n-1}^{(1)}(y_+)p_n(y_-) - (W + \Delta yV)p_{n-1}^{(1)}(y_-)p_n(y_+) - \Delta yUp_n(y_+)p_n(y_-) \in \mathbb{C}[x] + \Delta y\mathbb{C}[x]$ and furthermore because it is odd under $y_+ \leftrightarrow y_-$ this quantity is actually an element of $\Delta y\mathbb{C}[x]$ only. Consequently $(W - \Delta yV)q_n(y_+)p_n(y_-) - (W + \Delta yV)q_n(y_-)p_n(y_+) \in \Delta y\mathbb{C}[x]$ and we define the polynomial $\Theta_n(x)$ by

(4.18)
$$\Delta y \Theta_n(x) = (W + \Delta y V) q_n(y_-) p_n(y_+) - (W - \Delta y V) q_n(y_+) p_n(y_-),$$

which is (4.15). To find the other relations we employ the alternative form of $\Theta_n(x)$

(4.19)
$$\Delta y \Theta_n = (W - \Delta y V) p_{n-1}^{(1)}(y_+) p_n(y_-) - (W + \Delta y V) p_{n-1}^{(1)}(y_-) p_n(y_+) - \Delta y U p_n(y_+) p_n(y_-),$$

and multiply the left-hand sides by the Casoratians $1 = a_n [p_{n-1}(y_{\pm})p_{n-1}^{(1)}(y_{\pm}) - p_n(y_{\pm})p_{n-2}^{(1)}(y_{\pm})]$ so that upon re-organising we have

$$(4.20) \quad p_n(y_+) \left[(W + \Delta yV) p_{n-1}^{(1)}(y_-) + \Delta yU p_n(y_-) - \Delta y a_n \Theta_n p_{n-2}^{(1)}(y_+) \right] \\ = p_{n-1}^{(1)}(y_+) \left[(W - \Delta yV) p_n(y_-) - \Delta y a_n \Theta_n p_{n-1}(y_+) \right],$$

and

$$(4.21) \quad p_n(y_-) \left[(W - \Delta y V) p_{n-1}^{(1)}(y_+) - \Delta y U p_n(y_+) + \Delta y a_n \Theta_n p_{n-2}^{(1)}(y_-) \right] \\ = p_{n-1}^{(1)}(y_-) \left[(W + \Delta y V) p_n(y_+) + \Delta y a_n \Theta_n p_{n-1}(y_-) \right].$$

Relation (4.20) implies there must exist polynomials $\pi_1, \pi_2 \in \mathbb{C}[x]$ such that

$$(4.22) \quad p_n(y_+)p_{n-1}^{(1)}(y_+)(\pi_1 + \Delta y\pi_2) = p_n(y_+) \left[(W + \Delta yV)p_{n-1}^{(1)}(y_-) + \Delta yUp_n(y_-) - \Delta ya_n\Theta_n p_{n-2}^{(1)}(y_+) \right],$$

$$(4.23) \quad = p_{n-1}^{(1)}(y_+) \left[(W - \Delta yV)p_n(y_-) - \Delta ya_n\Theta_n p_{n-1}(y_+) \right],$$

while (4.21) implies an identical relation with $y_+ \leftrightarrow y_-$ and $\Delta y \mapsto -\Delta y$. Therefore we have the pair of relations

(4.24)
$$p_{n-1}^{(1)}(y_{+})(\pi_{1} + \Delta y\pi_{2}) = (W + \Delta yV)p_{n-1}^{(1)}(y_{-}) + \Delta yUp_{n}(y_{-}) - \Delta ya_{n}\Theta_{n}p_{n-2}^{(1)}(y_{+}),$$

(4.25)
$$p_{n}(y_{+})(\pi_{1} + \Delta y\pi_{2}) = (W - \Delta yV)p_{n}(y_{-}) - \Delta ya_{n}\Theta_{n}p_{n-1}(y_{+}),$$

and an analogous pair with $y_+ \leftrightarrow y_-$. We form the combination of $p_{n-1}(y_+) \times (4.24) - p_{n-2}^{(1)}(y_+) \times (4.25)$ and by employing the Casoratian and eliminating the associated polynomials we arrive at the solution

(4.26)
$$\pi_1 + \Delta y \pi_2 = -a_n (W + \Delta y V) p_{n-1}(y_+) q_n(y_-) + a_n (W - \Delta y V) p_n(y_-) q_{n-1}(y_+).$$

A similar procedure on the other pair yields the partner relation

(4.27)
$$\pi_1 - \Delta y \pi_2 = a_n (W + \Delta y V) p_n(y_+) q_{n-1}(y_-) - a_n (W - \Delta y V) p_{n-1}(y_-) q_n(y_+).$$

The sum and difference of these are simply (4.14) and (4.16) with the identifications $\pi_1 = 2W_n - W$ and $\pi_2 = -\Omega_n - V$.

Knowledge of the large x expansions of the spectral coefficients will be important in the ensuing analysis.

Proposition 4.3 ([60]). The spectral coefficients have terminating expansions about $x = \infty$ with leading order terms

(4.28)
$$W_n(x) = \frac{1}{2}W + \frac{1}{4}[W + \Delta yV] \left(\frac{y_+}{y_-}\right)^n + \frac{1}{4}[W - \Delta yV] \left(\frac{y_-}{y_+}\right)^n + O(x^{M-1}), \quad n \ge 0$$

(4.29)
$$\Theta_n(x) = \frac{1}{y_- \Delta y} [W + \Delta y V] \left(\frac{y_+}{y_-}\right)^n - \frac{1}{y_+ \Delta y} [W - \Delta y V] \left(\frac{y_-}{y_+}\right)^n + \mathcal{O}(x^{M-3}), \quad n \ge 0,$$

(4.30)
$$\Omega_n(x) + V(x) = \frac{1}{2\Delta y} [W + \Delta yV] \left(\frac{y_+}{y_-}\right)^n - \frac{1}{2\Delta y} [W - \Delta yV] \left(\frac{y_-}{y_+}\right)^n + \mathcal{O}(x^{M-2}), \quad n \ge 0.$$

Proof. These expansions follow from the substitution of the expansions for the polynomials (3.34) and associated functions (3.35) into the formulae (4.14)-(4.16). One should note that $y_+/y_- = O(1)$ as $x \to \infty$.

Corollary 4.2 ([60]). In the generic or regular \mathbb{D} -semi-classical case W_n, Θ_n, Ω_n are polynomials in x with fixed degrees independent of n, specifically $\deg_x W_n = M$, $\deg_x \Omega_n = M - 1$ and $\deg_x \Theta_n = M - 2$.

Proof. This follows from the previous proposition after recalling that $y_{\pm} = O(x)$ as $x \to \infty$.

The expressions for the spectral coefficients (4.14-4.16) can be inverted yielding a system of coupled first order linear divided-difference equations for the polynomials and associated functions with respect to x.

Proposition 4.4 ([60, 63]). The polynomials with a \mathbb{D} -semi-classical weight satisfy

(4.31)
$$W_n \mathbb{D}_x p_n = \Omega_n \mathbb{M}_x p_n - a_n \Theta_n \mathbb{M}_x p_{n-1}, \quad n \ge 0,$$

for certain spectral coefficients, $W_n(x), \Theta_n(x), \Omega_n(x)$. The partner equation to (4.31) is

(4.32)
$$W_n \mathbb{D}_x p_{n-1} = -(\Omega_n + 2V) \mathbb{M}_x p_{n-1} + a_n \Theta_{n-1} \mathbb{M}_x p_n, \quad n \ge 0.$$

The associated functions satisfy divided-difference equations

(4.33)
$$W_n \mathbb{D}_x q_n = (\Omega_n + 2V) \mathbb{M}_x q_n - a_n \Theta_n \mathbb{M}_x q_{n-1}, \quad n \ge 0,$$

(4.34)
$$W_n \mathbb{D}_x q_{n-1} = -\Omega_n \mathbb{M}_x q_{n-1} + a_n \Theta_{n-1} \mathbb{M}_x q_n, \quad n \ge 0,$$

identical in structure to (4.31) and (4.32) respectively. The coupled system (4.31, 4.32, 4.33, 4.34) can be written in matrix form as the spectral divided-difference equation

(4.35)
$$\mathbb{D}_x Y_n(x) := A_n \mathbb{M}_x Y_n(x) = \frac{1}{W_n(x)} \begin{pmatrix} \Omega_n(x) & -a_n \Theta_n(x) \\ a_n \Theta_{n-1}(x) & -\Omega_n(x) - 2V(x) \end{pmatrix} \mathbb{M}_x Y_n(x), \quad n \ge 0,$$

with A_n termed the spectral matrix.

Proof. Our derivation of the spectral divided-difference equations will employ a simple method - that of inversion of the relations for the spectral coefficients (4.14-4.16). Consider the following equivalent forms of those relations

$$(4.36) \qquad 2W_n - W + \Delta y(\Omega_n + V) = a_n \left[(W + \Delta y V) p_n(y_+) q_{n-1}(y_-) - (W - \Delta y V) p_{n-1}(y_-) q_n(y_+) \right],$$

$$(4.37) \qquad 2W_n - W - \Delta y(\Omega_n + V) = a_n \left[-(W + \Delta yV)p_{n-1}(y_+)q_n(y_-) + (W - \Delta yV)p_n(y_-)q_{n-1}(y_+) \right],$$

(4.38)
$$\Delta y \Theta_n = (W + \Delta y V) p_n(y_+) q_n(y_-) - (W - \Delta y V) p_n(y_-) q_n(y_+).$$

For our first task, that of deriving (4.31), we construct the combination $p_n(y_+) \times (4.37) - p_n(y_-) \times (4.36) + (p_{n-1}(y_+)+p_{n-1}(y_-))a_n \times (4.38)$. After effecting some initial cancellation we can then employ the Casoratian relation (3.26) to make further simplification and our final result is precisely (4.31) in finite difference form. The approach taken for the other relations (4.32)-(4.34) is the same. For (4.32) we use the combination $p_{n-1}(y_+) \times (4.36) - p_{n-1}(y_-) \times (4.37) - (p_n(y_+) + p_n(y_-))a_n \times (4.38)|_{n \mapsto n-1}$, while for (4.33) we use $q_n(y_+) \times (4.37) - q_n(y_-) \times (4.36) + (q_{n-1}(y_+) + q_{n-1}(y_-))a_n \times (4.38)$ and for (4.34) we use $q_{n-1}(y_+) \times (4.36) - q_{n-1}(y_-) \times (4.38)|_{n \mapsto n-1}$.

The initial values of the spectral coefficients are specified by the weight data W, V, U and the initial norm γ_0

(4.39)
$$W_{-1} = 0, \quad W_0 = W, \quad \Theta_{-1} = 0, \quad \Theta_0 = -\gamma_0^2 U, \quad \Omega_0 = 0, \quad \Omega_1 = -2V - \gamma_0^2 (\mathbb{M}_x x - b_0) U.$$

Remark 4.4. An alternative line of reasoning to achieve the results above, and in fact to generalise them beyond the \mathbb{D} -semi-classical class, is the following. By positing an explicit quadrature formula for the nonlinear lattice, one upon which the orthogonality condition is founded, one can derive (4.31), (4.32) and (4.35) assuming the existence of certain moments of "log-derivative" of the weight $\mathbb{D}_x w/\mathbb{M}_x w$. A byproduct of this approach is that one also constructs an explicit quadrature formula for the spectral coefficients, albeit one which involves the polynomials themselves.

The spectral structure in Proposition 4.4 has been derived before for OPS with respect to general classes of weights on the linear lattice by [29, 67, 44, 68] and on the q-linear lattice by [55, 67, 42, 54, 20, 32, 46, 70, 79]. This has also been done for specific weights on the q-linear lattice in [72, 15] and for OPS on the unit circle subject to q-difference relations in [48, 14]. In addition to the studies by Magnus the works [9, 28] have investigated the spectral structures for general SNUL and arbitrary $\deg_x W$, $\deg_x V$. Characterisation theorems for the classical OPS of the Askey table have been made by [22, 1] and others using an equivalent formulation to the spectral structures on general SNUL, but of course restricted to the $\deg_x W = 2$, $\deg_x V = 1$ case.

Compatibility relation between the matrix recurrence relation and the spectral divided-difference equation imposes the following conditions. The Cayley transform $(1 - \frac{1}{2}\Delta y A)^{-1} (1 + \frac{1}{2}\Delta y A)$ will figure prominently is the ensuing analysis.

Proposition 4.5. The spectral matrix and the recurrence matrix satisfy

(4.40)
$$A_{n+1} \cdot \mathbb{M}_x K_n - \mathbb{M}_x K_n \cdot A_n = \mathbb{D}_x K_n - \frac{1}{4} \Delta y^2 A_{n+1} \cdot \mathbb{D}_x K_n \cdot A_n, \quad n \ge 0,$$

or equivalently

(4.41)
$$K_n(y_+) \left(1 - \frac{1}{2}\Delta y A_n\right)^{-1} \left(1 + \frac{1}{2}\Delta y A_n\right) = \left(1 - \frac{1}{2}\Delta y A_{n+1}\right)^{-1} \left(1 + \frac{1}{2}\Delta y A_{n+1}\right) K_n(y_-), \quad n \ge 0.$$

Proof. The finite difference form of (4.35) is

(4.42)
$$\left(1 - \frac{1}{2}\Delta y A_n\right) Y_n(y_+) = \left(1 + \frac{1}{2}\Delta y A_n\right) Y_n(y_-),$$

which demonstrates why the Cayley transform arises. The second relation (4.41) is found by computing $Y_{n+1}(y_+)$ in two different ways from $Y_n(y_-)$ corresponding to the different orders of the operations $n \mapsto n+1$ and $y_- \mapsto y_+$. The first relation (4.40) is derived by computing $\mathbb{D}_x Y_{n+1}$ in terms of $\mathbb{M}_x Y_n$ in two different orders.

Compatibility of the three term recurrence relation and the spectral divided-difference equation equation implies the following result.

Proposition 4.6 ([60, 63]). The spectral coefficients arising in Proposition 4.4 satisfy recurrence relations in n,

(4.43)
$$W_{n+1} = W_n + \frac{1}{4}\Delta y^2 \Theta_n, \quad n \ge 0,$$

(4.44)
$$\Omega_{n+1} + \Omega_n + 2V = (\mathbb{M}_x x - b_n)\Theta_n, \quad n \ge 0,$$

(4.45)

$$(W_n\Omega_{n+1} - W_{n+1}\Omega_n)(\mathbb{M}_x x - b_n) = -\frac{1}{4}\Delta y^2 \Omega_{n+1}\Omega_n + W_n W_{n+1} + a_{n+1}^2 W_n \Theta_{n+1} - a_n^2 W_{n+1}\Theta_{n-1}, \quad n \ge 0.$$

Proof. We offer a proof in the spirit of our previous proofs. For the first relation (4.43) we construct $2W_{n+1} - 2W_n - \frac{1}{2}\Delta y^2\Theta_n$ and substitute the expressions (4.14) and (4.15) for W_n and Θ_n respectively. After collecting terms proportional to $W \pm \Delta y V$ we observe that the coefficients of these terms separate into two parts, in which the three term recurrence relation can be employed. The result we find is that they identically vanish. The second relation (4.44) can be found in an identical manner starting with $\Delta y [\Omega_{n+1} + \Omega_n + 2V - (\mathbb{M}_x x - b_n)\Theta_n]$.

The relations of Proposition 4.6 constitute the analogs of the Laguerre-Freud equations and have been studied in the special case of linear lattices by [29, 65, 44, 68] and in the q-linear case by [55, 54, 70, 32]. For the situation of OPS on the unit circle with q-difference operators some of these relations can be found in [48]. Following on from the Magnus studies the works [9, 28] have investigated these compatibility conditions for general SNUL. Together with Proposition 4.4 this result allows for the construction of ladder, or raising and lowering operators and these have been found in the q-linear lattice by [20, 46].

A consequence of Proposition 4.6 is the following.

Proposition 4.7 ([60]). The spectral coefficients satisfy the bilinear recurrence relation

$$(4.46) \quad W_n(W_n - W) = \frac{1}{4}\Delta y^2 \left[\Omega_n(\Omega_n + 2V) - a_n^2 \Theta_{n-1}\Theta_n\right] = -\frac{1}{4}\Delta y^2 \det \begin{pmatrix} \Omega_n & -a_n\Theta_n \\ a_n\Theta_{n-1} & -\Omega_n - 2V \end{pmatrix}, \quad n \ge 0.$$

Proof. Two methods of proof are available here. The first method starts from the observation that the above bilinear relation can be expressed as

(4.47)
$$[2W_n - W + \Delta y(\Omega_n + V)] [2W_n - W - \Delta y(\Omega_n + V)] + a_n^2 \Delta y^2 \Theta_n \Theta_{n-1} = W^2 - \Delta y^2 V^2$$

Now we have evaluations of the factors appearing in the left-hand side of the preceeding equation from the proof of Proposition 4.2, namely (4.26, 4.27, 4.18), in terms of products of a polynomial and associated function. Substituting these evaluations into the left-hand side, effecting considerable cancellation of terms and employing the Casoratian relation (3.26) the final result simplifies to the right-hand side.

The second method uses the compatibility relations (4.43- 4.45) and constructs an integral for this coupled system. First one multiplies the left-hand side of (4.44) by the left-hand side of (4.45) and cancels out the common factor of $\mathbb{M}x - b_n$ in the resulting equation. By judicious use of (4.43) one can effect the cancellation of four terms in this equation and a recasting of the terms linear in Θ_n . Consequently we arrive at the following equation

$$(4.48) \quad \frac{(\Omega_{n+1}+V)^2}{W_{n+1}} - \frac{(\Omega_n+V)^2}{W_n} = \frac{4}{\Delta y^2} (W_{n+1}-W_n) + V^2 \left[\frac{1}{W_{n+1}} - \frac{1}{W_n}\right] + a_{n+1}^2 \frac{\Theta_{n+1}\Theta_n}{W_{n+1}} - a_n^2 \frac{\Theta_n\Theta_{n-1}}{W_n},$$

which is manifestly a perfect difference equation in n. By summing this and employing the initial value evaluations (4.39) we have (4.46).

The above Proposition must be augmented with the following matrix identities.

Corollary 4.3. The matrix factor appearing in Proposition 4.5 has the determinant evaluation

(4.49)
$$\det(1 \pm \frac{1}{2}\Delta y A_n) = \frac{W \mp \Delta y V}{W_n}, \quad n \ge 0,$$

and its inverse is

(4.50)
$$(1 \pm \frac{1}{2}\Delta y A_n)^{-1} = \frac{1}{W \mp \Delta y V} \begin{pmatrix} W_n \mp \frac{1}{2}\Delta y (\Omega_n + 2V) & \pm \frac{1}{2}\Delta y a_n \Theta_n \\ \mp \frac{1}{2}\Delta y a_n \Theta_{n-1} & W_n \pm \frac{1}{2}\Delta y \Omega_n \end{pmatrix}, \quad n \ge 0.$$

Proof. From the form for the spectral matrix given by (4.35) we have

$$\det(1 \pm \frac{1}{2}\Delta y A_n) = \frac{1}{W_n^2} \left[(W_n \pm \frac{1}{2}\Delta y \Omega_n) (W_n \mp \frac{1}{2}\Delta y (\Omega_n + 2V)) + \frac{1}{4}\Delta y^2 a_n^2 \Theta_n \Theta_{n-1} \right]$$
$$= \frac{1}{W_n^2} \left[W_n^2 \mp \Delta y V W_n - \frac{1}{4}\Delta y^2 \left[\Omega_n (\Omega_n + 2V) - a_n^2 \Theta_n \Theta_{n-1} \right] \right]$$
$$= \frac{W \mp \Delta y V}{W_n},$$

where we have employed the previous Corollary in the last step. Constructing the matrix inverse given on the left-hand side of (4.50) and using our determinant formula we find (4.50).

Consequent to the results of Corollary 4.3 we require the matrix product found in (4.41).

Corollary 4.4. The matrix product appearing in (4.41) has the evaluation

$$(4.51) \quad \left(1 - \frac{1}{2}\Delta y A_n\right)^{-1} \left(1 + \frac{1}{2}\Delta y A_n\right) \\ = \frac{1}{W + \Delta y V} \begin{pmatrix} 2W_n - W + \Delta y (\Omega_n + V) & -\Delta y a_n \Theta_n \\ \Delta y a_n \Theta_{n-1} & 2W_n - W - \Delta y (\Omega_n + V) \end{pmatrix}, \quad n \ge 0, \\ = a_n \begin{pmatrix} p_n(y_+) \\ p_{n-1}(y_+) \end{pmatrix} \otimes \left(q_{n-1}(y_-), \quad -q_n(y_-)\right) - a_n \frac{W - \Delta y V}{W + \Delta y V} \begin{pmatrix} q_n(y_+) \\ q_{n-1}(y_+) \end{pmatrix} \otimes \left(p_{n-1}(y_-), \quad -p_n(y_-)\right)$$

Proof. Using (4.50) we multiply out the matrix product and employ the bilinear identity (4.46) to simplify the diagonal elements, resulting in (4.51). \Box

This result motivates the following definitions

(4.52)
$$\mathfrak{W}_{\pm} := 2W_n - W \pm \Delta y \,(\Omega_n + V), \quad n \ge 1,$$

(4.53)
$$\mathfrak{T}_{+} := \Delta y \, a_{n} \Theta_{n}, \quad \mathfrak{T}_{-} := \Delta y \, a_{n} \Theta_{n-1}, \quad n \ge 1,$$

whilst for n = 0 we have $\mathfrak{W}_{\pm}(n = 0) := W \pm \Delta y V$, $\mathfrak{T}_{+}(n = 0) := -\Delta y a_0 \gamma_0^2 U$, $\mathfrak{T}_{-}(n = 0) := 0$ together with

(4.54)
$$A_n^* := \begin{pmatrix} \mathfrak{W}_+ & -\mathfrak{T}_+ \\ \mathfrak{T}_- & \mathfrak{W}_- \end{pmatrix}.$$

The variables \mathfrak{W}_{\pm} are essentially the variables U_n, X_n employed by Magnus in [63] respectively whilst the variables \mathfrak{T}_{\pm} are directly related to his Y_n, Z_n variables. In these new variables the bi-linear relation (4.46) becomes

(4.55)
$$\det A_n^* = \mathfrak{W}_+ \mathfrak{W}_- + \mathfrak{T}_+ \mathfrak{T}_- = W^2 - \Delta y^2 V^2,$$

which will be the more useful form.

It is of advantage to write out recurrence-spectral compatibility equations in terms of $\mathfrak{W}_{\pm,n}, \mathfrak{T}_{\pm,n}$, where we append a subscript to indicate the dependence on the index n.

Corollary 4.5. Solving (4.41) for A_{n+1}^* we deduce

$$(4.56) \quad a_n a_{n+1}\mathfrak{T}_{+,n+1} = -a_n(y_+ - b_n)\mathfrak{W}_{+,n} + a_n(y_- - b_n)\mathfrak{W}_{-,n} + (y_+ - b_n)(y_- - b_n)\mathfrak{T}_{+,n} + a_n^2\mathfrak{T}_{-,n}$$

$$(4.57) a_n \mathfrak{T}_{-,n+1} = a_{n+1} \mathfrak{T}_{+,n},$$

(4.58)
$$a_n \mathfrak{W}_{+,n+1} = a_n \mathfrak{W}_{-,n} + (y_+ - b_n) \mathfrak{T}_{+,n},$$

(4.59)
$$a_n \mathfrak{W}_{-,n+1} = a_n \mathfrak{W}_{+,n} - (y_- - b_n) \mathfrak{T}_{+,n}$$

A significant result follows from the observation that the right-hand side of (4.55) is a polynomial in x with fixed degree independent of n and in fact contains no dependence on n. If we denote one of the zeros of the spectral polynomial $W^2 - \Delta y^2 V^2$ as x_j then we can apply the equality for $n \mapsto n+1$ and use (4.57) to deduce

(4.60)
$$\mathfrak{W}_{+,n+1}(x_j)\mathfrak{W}_{-,n+1}(x_j) + \frac{a_{n+1}}{a_n}\mathfrak{T}_{+,n+1}(x_j)\mathfrak{T}_{+,n}(x_j) = 0.$$

This then allows us to draw the following conclusion.

Proposition 4.8. The variable $\mathfrak{T}_{+,n+1}(x_j)$, when written in terms of the variables at n, factorises in the following way (4.61)

$$(1.01) - a_n a_{n+1} \mathfrak{T}_{+,n+1}(x_j) \mathfrak{T}_{+,n}(x_j) = [a_n \mathfrak{W}_{-,n}(x_j) + (y_{+,j} - b_n) \mathfrak{T}_{+,n}(x_j)] [a_n \mathfrak{W}_{+,n}(x_j) - (y_{-,j} - b_n) \mathfrak{T}_{+,n}(x_j)].$$

The coupled pair of first order divided-difference equations (4.31) and (4.32) imply a second order equation for one of the components, say p_n .

Proposition 4.9 ([60, 8]). The \mathbb{D} -semi-classical orthogonal polynomials or associated functions satisfy a second order divided-difference equation in the following two equivalent forms - either

(4.62)

$$\begin{bmatrix} \mathbb{M}_x \left(\frac{W_n}{\Theta_n} \right) + \frac{1}{4} \Delta y^2 \mathbb{D}_x \left(\frac{\Omega_n + 2V}{\Theta_n} \right) \end{bmatrix} \mathbb{D}_x^2 p_n + \begin{bmatrix} \mathbb{D}_x \left(\frac{W_n}{\Theta_n} \right) + \mathbb{M}_x \left(\frac{2V}{\Theta_n} \right) + \Delta y^2 \mathbb{M}_x \left(\frac{W - W_n}{\Delta y^2 \Theta_n} \right) \end{bmatrix} \mathbb{M}_x \mathbb{D}_x p_n + \begin{bmatrix} \mathbb{M}_x \left(4 \frac{W - W_n}{\Delta y^2 \Theta_n} \right) - \mathbb{D}_x \left(\frac{\Omega_n}{\Theta_n} \right) \end{bmatrix} \mathbb{M}_x^2 p_n = 0,$$

or alternatively

$$(4.63) \quad E_x^+ \left(\frac{W + \Delta yV}{\Delta y\Theta_n}\right) (E_x^+)^2 p_n + E_x^- \left(\frac{W - \Delta yV}{\Delta y\Theta_n}\right) (E_x^-)^2 p_n \\ + \left\{-E_x^+ \left(\frac{2W_n - W + \Delta y(\Omega_n + V)}{\Delta y\Theta_n}\right) - E_x^- \left(\frac{2W_n - W - \Delta y(\Omega_n + V)}{\Delta y\Theta_n}\right)\right\} p_n = 0.$$

Proof. Starting with (4.31) and (4.32), we can employ (4.46) to write an equation for each of $\mathbb{M}_x p_{n-1}$ and $\mathbb{D}_x p_{n-1}$ in terms of linear combinations of $\mathbb{M}_x p_n$ and $\mathbb{D}_x p_n$. Then utilising the fact that $\mathbb{M}_x \mathbb{D}_x p_{n-1} = \mathbb{D}_x \mathbb{M}_x p_{n-1}$ we arrive at (4.62). The equation (4.63) is the nodal equivalent of the former.

This concludes our discussion of the spectral structures for a general quadratic lattice and we now turn to the simplest explicit example on the q-quadratic lattice.

5. M = 2, L = 0 case and the Askey-Wilson polynomials

Here we will employ the theory of Section 4 and explicitly compute the spectral coefficients for the Askey-Wilson system itself because it serves both as an instructive example for the theory of the previous section and clarifies some confusion present in the literature. Virtually all of our results presented here have been found by earlier studies - firstly by Askey and Wilson [4], then most notably by the Soviet school of Nikiforov, Suslov and Uvarov [76, 73, 77, 74, 75] who were primarily concerned with hypergeometric type OPS on non-uniform lattices, and in the 1988 work of Magnus [60] whose work and intent is most similar to the spirit of our own.

We recall the Askey-Wilson weight [4] itself has degrees 2N = 4 or M = 2 and L = 0 with

(5.1)
$$w(x) = w(x; \{a_1, \dots, a_4\}) = \frac{(z^{\pm 2}; q)_{\infty}}{\sin(\theta) \prod_{j=1}^4 (a_j z^{\pm 1}; q)_{\infty}}, \quad G = (-1, 1).$$

Let the *j*-th elementary symmetric polynomial of a_1, \ldots, a_4 be denoted by σ_j for $j = 0, \ldots, 4$.

Proposition 5.1. Assume that $q \neq 1$ and $|q^{-1/2}a_j| \neq 1$ for j = 1, ..., 4. The spectral coefficients for the Askey-Wilson OP system are

(5.2)
$$W_n(x) = (1+q^{-n})(1+\sigma_4 q^{n-2})(x^2-1) - \left[q^{-1/2}\sigma_1 + q^{-3/2}\sigma_3\right]x + 1 + q^{-1}\sigma_2 + q^{-2}\sigma_4,$$

(5.3)

$$\Omega_{n}(x) + V(x) = 2\frac{q^{n-2}\sigma_{4} - q^{-n}}{q^{1/2} - q^{-1/2}}x + \frac{q^{-n-1/2}\sigma_{1} + (-2+q^{-n})q^{-3/2}\sigma_{3} + (-2+q^{n})q^{-5/2}\sigma_{1}\sigma_{4} + q^{n-7/2}\sigma_{3}\sigma_{4}}{(q^{1/2} - q^{-1/2})(q^{-n} - q^{n-2}\sigma_{4})},$$

$$(5.4) \qquad \qquad \Theta_{n}(x) = 4\frac{q^{n-3/2}\sigma_{4} - q^{-n-1/2}}{q^{1/2} - q^{-1/2}},$$

valid for $n \ge 0$. The three-term recurrence coefficients are found to be given by the standard expressions [56, 58]

(5.5)
$$a_n^2 = \frac{1}{4} \frac{(1-q^n)(1-\sigma_4 q^{n-2}) \prod_{k>j} (1-a_j a_k q^{n-1})}{(1-\sigma_4 q^{2n-3})(1-\sigma_4 q^{2n-2})^2 (1-\sigma_4 q^{2n-1})},$$

and

(5.6)
$$b_n = \left[\sigma_1(q + \sigma_4(q^{2n} - q^n - q^{n-1})) + \sigma_3(1 - q^n - q^{n+1} + \sigma_4 q^{2n-1})\right] \frac{q^{n-1}}{2(1 - \sigma_4 q^{2n})(1 - \sigma_4 q^{2n-2})},$$

for $n \ge 0$ where we assume $a_0 = 0$.

Proof. From the weight (5.1) we compute the spectral data

(5.7)
$$W \pm \Delta y V = z^{\mp 2} \prod_{j=1}^{4} (1 - a_j q^{-1/2} z^{\pm 1}),$$

from which we deduce

(5.8)
$$W(x) = 2(1 + \sigma_4 q^{-2})x^2 - \left[q^{-1/2}\sigma_1 + q^{-3/2}\sigma_3\right]x - 1 + q^{-1}\sigma_2 - q^{-2}\sigma_4,$$

(5.9)
$$V(x) = 2\frac{q^{-2}\sigma_4 - 1}{q^{1/2} - q^{-1/2}}x + \frac{q^{-1/2}\sigma_1 - q^{-3/2}\sigma_3}{q^{1/2} - q^{-1/2}}.$$

We will see that $W^2 - \Delta y^2 V^2$ will play a significant role and therefore define another set of elementary symmetric polynomials by

(5.10)
$$W^2 - \Delta y^2 V^2 = K^2 \left[x^4 - e_1 x^3 + e_2 x^2 - e_3 x + e_4 \right].$$

We note the evaluation $K = 4q^{\mu} = 4q^{-1}\sqrt{\sigma_4}$. We parameterise the spectral coefficients in the following way

(5.11)
$$2W_n - W = w_2 x^2 + w_1 x + w_0$$

(5.12)
$$\Theta_n = \overline{\omega}_+, \quad \Theta_{n-1} = \overline{\omega}_-,$$

$$(5.13)\qquad \qquad \Omega_n + V = v_1 x + v_0,$$

where we know from (4.28-4.30) that the leading coefficients are

(5.14)
$$w_2 = \frac{K}{2} \left(q^{n+\mu} + q^{-n-\mu} \right), \quad v_1 = \frac{K}{2} \frac{q^{n+\mu} - q^{-n-\mu}}{q^{1/2} - q^{-1/2}}, \quad \varpi_+ = K \frac{q^{n+\mu+1/2} - q^{-n-\mu-1/2}}{q^{1/2} - q^{-1/2}}.$$

From the fundamental bi-linear relation (4.46) we get a system of quadratic polynomial equalities

(5.15)
$$w_2^2 - \Delta v_1^2 = K^2,$$

(5.16)
$$2w_1w_2 - 2\Delta v_0v_1 = -K^2 e_1,$$

(5.17)
$$2w_0w_2 + w_1^2 - \Delta(v_0^2 - v_1^2) + a_n^2\Delta\varpi_+\varpi_- = K^2e_2,$$

(5.18)
$$2w_0w_1 + 2\Delta v_0v_1 = -K^2 e_3,$$

(5.19)
$$w_0^2 + \Delta v_0^2 - a_n^2 \Delta \varpi_+ \varpi_- = K^2 e_4,$$

where $\Delta := (q^{1/2} - q^{-1/2})^2$. Now (5.15,5.17,5.19) imply $w_1^2 + (w_0 + w_2)^2 = K^2(1 + e_2 + e_4)$ while (5.18,5.16) imply $2w_1(w_0 + w_2) = -K^2(e_1 + e_3)$. Forming the sum and difference of these two later relations we conclude that

(5.20)
$$w_1 = \frac{W(1) - W(-1)}{2}, \quad w_0 = -w_2 + \frac{W(1) + W(-1)}{2}$$

Using (5.16) for example, along with the above solutions, we find

(5.21)
$$v_0 = \frac{K^2 e_1 + w_2(W(1) - W(-1))}{2\Delta v_1}.$$

Therefore we have succeeded in relating the sub-leading coefficients in terms of explicitly known quantities and after simplification we arrive at (5.2-5.3). Finally, using (5.19), we can solve for a_n^2 and after observing that e_4 can be expressed as

(5.22)
$$e_4 = \left[\frac{q^{\mu} + q^{-\mu}}{2} - \frac{W(1) + W(-1)}{2K}\right]^2 + \frac{1}{(q^{\mu} - q^{-\mu})^2} \left[e_1 + \frac{q^{\mu} + q^{-\mu}}{2} \frac{W(1) - W(-1)}{K}\right]^2,$$

we can factorise the resulting four terms and arrive at (5.5). To find (5.6) we start with (4.44) and use our previous results for (5.4) and (5.3). We note the terms linear in x cancel identically, as they must, and after some factorisation we deduce (5.6).

It is the coefficients (5.2-5.3) along with first order divided-difference equation (4.31) that constitutes the structural relation for the Askey-Wilson polynomials, that is the analog of first order difference or differential relations for the classical orthogonal polynomials. The divided-difference relations reported in [58] are all of second order.

It is an easy task to evaluate the Askey-Wilson integral [4, 56] as a q-factorial and therefore compute the moments. The Askey-Wilson integral is defined by

(5.23)
$$I_2(a_1, a_2, a_3, a_4) := \int_{\mathbb{T}} \frac{dz}{2\pi i z} \frac{(z^{\pm 2}; q)_{\infty}}{\prod_{j=1}^4 (a_j z^{\pm 1}; q)_{\infty}}$$

with $|a_j| < 1$ for j = 1, ..., 4.

Our method is to apply the general system of moment recurrences to the case M = 2 and we find that it coincides with the recurrence of Kalnins and Miller [53] and Koelink and Koornwinder [57].

Proposition 5.2 ([97]). The Askey-Wilson integral satisfies the two-term linear recurrence

$$(5.24) \qquad (\sigma_4 - 1)I_2(qa_1, a_2, a_3, a_4) = (a_1a_2 - 1)(a_1a_3 - 1)(a_1a_4 - 1)I_2(a_1, a_2, a_3, a_4)$$

which is solved by

(5.25)
$$I_2(a_1, a_2, a_3, a_4) = 2 \frac{(\sigma_4; q)_{\infty}}{(q; q)_{\infty} \prod_{k>j} (a_j a_k; q)_{\infty}}.$$

Consequently the moments are given by

(5.26)
$$m_{0,n}(a_1) = \pi \frac{(a_1 a_2, a_1 a_3, a_1 a_4; q)_n}{(\sigma_4; q)_n} I_2(a_1, a_2, a_3, a_4)$$

(5.27)
$$= 2\pi \frac{(q^n \sigma_4; q)_{\infty}}{(q^n a_1 a_2, q^n a_1 a_3, q^n a_1 a_4, a_2 a_3, a_2 a_4, a_3 a_4, q; q)_{\infty}}$$

Proof. We only require the k = 0 case of (4.11) with $a = a_1$ and from (5.7) we compute the relevant coefficients as $\delta_{2,1}(a) = q^{-1}a^{-1}(1-\sigma_4)$ and $\delta_{2,0}(a) = q^{-1}(a+a^{-1})(\sigma_4-1) + q^{-1}(\sigma_1-\sigma_3)$. The solution of the recurrence (5.24) follows from the arguments given in [57].

The specialisation of the second-order divided-difference equation (4.63) with the M = 2 spectral coefficients of Proposition 5.1 is given by (5.28)

$$\frac{\prod_{j=1}^{4}(1-a_jz)}{qz^2-1}\left[(E_x^+)^2p_n-p_n\right] + \frac{\prod_{j=1}^{4}(z-a_j)}{z^2-q}\left[(E_x^-)^2p_n-p_n\right] + q^{-n-1}(1-q^n)(q^n\sigma_4-q)(z^2-1)p_n = 0.$$

This is soluble in terms of basic hypergeometric functions and their polynomial solutions have been given by [60] or using factorisation methods in [7, 11, 10] as Askey-Wilson polynomials. The Askey-Wilson polynomials have an explicit form as a balanced $_4\varphi_3$ function [56] with the manifest symmetry under $z \leftrightarrow z^{-1}$ and are given in the monic form by

(5.29)
$$\pi_n(x) = \frac{(a_1a_2, a_1a_3, a_1a_4; q)_n}{(2a_1)^n (q^{n-1}\sigma_4; q)_n} {}_4\varphi_3 \left(\begin{array}{c} q^{-n}, \sigma_4 q^{n-1}, a_1z, a_1z^{-1}\\ a_1a_2, a_1a_3, a_1a_4 \end{array}; q, q\right)$$

or the alternative form [34] which is manifestly symmetric under permutations of a_1, a_2, a_3, a_4

$$(5.30) \quad \pi_n(x) = (2z)^{-n} \frac{(a_1 z, a_2 z, a_3 z, a_4 z, \sigma_4 q^{-1}; q)_n}{(z^2; q)_n (\sigma_4 q^{-1}; q)_{2n}} {}_8W_7(q^{-n} z^{-2}; q^{-n}, a_1 z^{-1}, a_2 z^{-1}, a_3 z^{-1}, a_4 z^{-1}; \frac{q^{2-n}}{\sigma_4}).$$

From Eq. (2.6) of [82] or equivalently from Eq. (4.18) of [45] with $\alpha = 0$ and Eq. (III.23) of [31] we deduce the expression for the Stieltjes function as a very-well-poised ${}_{8}W_{7}$ (5.31)

$$f(x) = \frac{4\pi (q^{-1}\sigma_4; q)_{\infty}}{(q; q)_{\infty} \prod_{k>j} (a_j a_k; q)_{\infty}} \frac{(1 - qz^{-2})}{z \prod_{1 \le j \le 4} (1 - a_j z^{-1})} * W_7(qz^{-2}; \frac{q}{a_1} z^{-1}, \frac{q}{a_2} z^{-1}, \frac{q}{a_3} z^{-1}, \frac{q}{a_4} z^{-1}, q; q, q^{-1}\sigma_4),$$

which exhibits the parameter symmetry. This expression is the one valid on the second Riemann sheet of the cut x-plane or exterior to the unit circle in z, |z| > 1. There are evaluations of $q_n(x)$ in [82] and [45] that are simple generalisations of the above formulae but we do not need to discuss them here.

Having an explicit form for the Stieltjes function we have two tasks at hand - to verify that is satisfies the inhomogeneous divided-difference equation (4.13) and the large x expansion formula (3.46).

Proposition 5.3. The Stieltjes function given by Eq.(5.31) satisfies the inhomogeneous divided difference equation (4.13) with the constant

(5.32)
$$U = \frac{8\pi}{q-1} \frac{(q^{-1}\sigma_4; q)_{\infty}}{(q; q)_{\infty} \prod_{k>j} (a_j a_k; q)_{\infty}}$$

Proof. We require an alternative form

$$(5.33) \quad f(x) = 4\pi \frac{qa_1}{\sigma_4} \frac{1}{\left(\frac{q}{a_2a_3}, \frac{q}{a_2a_4}, \frac{q}{a_3a_4}; q\right)_{\infty}} \left\{ -\frac{\left(\frac{q}{a_2z}, \frac{q}{a_3z}, \frac{q}{a_4z}, \frac{\sigma_4}{qa_1z}, \frac{q^2a_1z}{\sigma_4}; q\right)_{\infty}}{(a_2a_3, a_2a_4, a_3a_4, a_1z^{-1}, a_2z^{-1}, a_3z^{-1}, a_4z^{-1}, a_1z; q)_{\infty}} + \frac{(q^{-1}\sigma_4, q\frac{a_1}{a_2}, q\frac{a_1}{a_3}, q\frac{a_1}{a_4}, \frac{q^2}{\sigma_4}; q)_{\infty}}{(q, qa_1^2; q)_{\infty} \prod_{k>j} (a_ja_k; q)_{\infty} (1 - a_1z)(1 - a_1z^{-1})} {}_8W_7(a_1^2; a_1z, a_1z^{-1}, a_1a_2, a_1a_3, a_1a_4; \frac{q^2}{\sigma_4}) \right\},$$

which makes the $z \leftrightarrow z^{-1}$ symmetry manifest in the second term only. This is derived from the previous expression by utilising Eq.(III.37) combined with Eq.(III.23) of [31], and Eq.(III.36) for the term which specialises. We are now in a position to verify that (5.33) satisfies (4.13). This relies on the fact that (4.13)

with the data (5.7) is precisely the contiguous relation, Eq. (2.2) of [45], which in our context states

$$(5.34) \quad z^{2} \frac{\prod_{j=2,3,4} (1-q^{-1/2}a_{j}z^{-1})}{1-q^{1/2}a_{1}z} {}_{8}W_{7}(a_{1}^{2};q^{1/2}a_{1}z,q^{-1/2}a_{1}z^{-1},a_{1}a_{2},a_{1}a_{3},a_{1}a_{4};\frac{q^{2}}{\sigma_{4}}) - z^{-2} \frac{\prod_{j=2,3,4} (1-q^{-1/2}a_{j}z)}{1-q^{1/2}a_{1}z^{-1}} {}_{8}W_{7}(a_{1}^{2};q^{-1/2}a_{1}z,q^{1/2}a_{1}z^{-1},a_{1}a_{2},a_{1}a_{3},a_{1}a_{4};\frac{q^{2}}{\sigma_{4}}) = q^{-3/2}a_{2}a_{3}a_{4}(1-\frac{q}{\sigma_{4}})(z-z^{-1})_{8}W_{7}(a_{1}^{2};q^{1/2}a_{1}z,q^{1/2}a_{1}z^{-1},a_{1}a_{2},a_{1}a_{3},a_{1}a_{4};\frac{q}{\sigma_{4}}).$$

We have also used the specialisation formula when aq = bc

(5.35)
$${}_{8}W_{7}(a;b,c,d,e,f;q,\frac{q^{2}a^{2}}{bcdef}) = \frac{(aq,aq/de,aq/df,aq/ef;q)_{\infty}}{(aq/d,aq/e,aq/f,aq/def;q)_{\infty}}$$

which applies to the right-hand side of the previous equation. A consequence one can draw from this calculation is the explicit evaluation of the constant spectral coefficient. \Box

Remark 5.1. The constant U can be found in another way. This is by noting the initial condition for Φ_0 is given by (4.39) and that Φ_n is given purely by the leading order term in (5.4), so that knowing the normalisation γ_0^2 allows us to make the evaluation.

Proposition 5.4. The Stieltjes function possesses the explicit large x moment generating function formula

$$(5.36) \quad f(x) = -\frac{4\pi a_1(\sigma_4; q)_{\infty}}{(q; q)_{\infty} \prod_{k>j} (a_j a_k; q)_{\infty} (1 - a_1 z)(1 - a_1 z^{-1})} {}_4 \varphi_3 \left(\begin{array}{c} q, a_1 a_2, a_1 a_3, a_1 a_4 \\ q a_1 z, q a_1 z^{-1}, \sigma_4 \end{array}; q, q \right) \\ - \frac{4\pi (a_2 a_3 a_4 z^{-1}, q z^{-2}; q)_{\infty}}{z(a_2 a_3, a_2 a_4, a_3 a_4, a_2 z^{-1}, a_3 z^{-1}, a_4 z^{-1}; q)_{\infty}} \frac{1}{\phi_{\infty}(x; a_1)} {}_3 \varphi_2 \left(\begin{array}{c} a_2 z^{-1}, a_3 z^{-1}, a_4 z^{-1} \\ a_2 a_3 a_4 z^{-1}, q z^{-2} \end{array}; q, q \right).$$

Proof. By applying the transformation formula (III.36) of [31] to the $_8W_7$ function in (5.33) we get a sum of two $_4\varphi_3$ functions one of which reduces to a $_3\varphi_2$ function. After some simplification we get (5.36). We observe that the $_4\varphi_3$ term is precisely the second term of (3.46) given our formula for the moments (5.27). This allows us to conclude that

$$(5.37) f_{\infty}(x) = -\frac{4\pi (a_2 a_3 a_4 z^{-1}, q z^{-2}; q)_{\infty}}{z(a_2 a_3, a_2 a_4, a_3 a_4, a_2 z^{-1}, a_3 z^{-1}, a_4 z^{-1}; q)_{\infty}} {}_{3}\varphi_2 \left(\begin{array}{c} a_2 z^{-1}, a_3 z^{-1}, a_4 z^{-1} \\ a_2 a_3 a_4 z^{-1}, q z^{-2} \end{array}; q, q\right).$$

6. Deformation Differences

In this section we discuss the deformation structures based upon a nonlinear lattice in the u variable (or any number of variables for that matter) and the corresponding divided-difference operators. We emphasise that many of the results for the spectral structure have a parallel result in the deformation structure although there will be crucial differences in the details. Denote the deformation variable u and its forward and backward shifts by $v_{\pm} = E_u^{\pm} u$ and their difference by $\Delta v := v_+ - v_-$. The reader should be aware that the deformation lattice does not have to be the same type as the spectral lattice. Our D-semi-classical weight w, defined by Definition 4.1, acquires an additional dependence on a deformation variable u and furthermore satisfies a deformation divided-difference equation with respect to u.

Definition 6.1. The deformed \mathbb{D} -semi-classical weight satisfies

(6.1)
$$R\mathbb{D}_u w = 2S\mathbb{M}_u w,$$

or equivalently

(6.2)
$$\frac{w(x;v_+)}{w(x;v_-)} = \frac{R + \Delta vS}{R - \Delta vS}(x;u).$$

for polynomials S(x; u), R(x; u) irreducible in x and $u \in J$. In addition we assume $R \pm \Delta vS \neq 0$ for all $x \in G, u \in J$ and that the deformed OPS exists, i.e. that $\gamma_n(v_{\pm}) \neq 0$ for all $n \in \mathbb{Z}_{\geq 0}$. We also require the condition $\gamma_0(v_{\pm})(R - \Delta vS) + \gamma_0(v_{\pm})(R + \Delta vS) \neq 0$.

There is a very simple motivation for this relation. In the process of extending the weights beyond those of the classical cases in the Askey table by increasing the degrees $\deg_x W \ge 3$, $\deg_x V \ge 2$ the weights acquire additional parameters which if suitably chosen sit in a completely reflexive or symmetric way to the spectral variable and therefore will satisfy an analogous linear, homogeneous and first order divided-difference equation in any of these new parameters.

There are conditions imposed on the weight given that it satisfies both (4.1) or (4.2) and (6.1) or (6.2).

Proposition 6.1. The spectral data polynomials W(x; u), V(x; u) and the deformation data polynomials R(x; u), S(x; u) satisfy the compatibility relations

$$(6.3) \left[\mathbb{D}_x \frac{2S}{R} - \mathbb{D}_u \frac{2V}{W} \right] \left[1 - \frac{1}{16} \Delta y^2 \Delta v^2 \mathbb{D}_x \frac{2S}{R} \mathbb{D}_u \frac{2V}{W} \right] = \frac{1}{4} \Delta y^2 \mathbb{D}_x \frac{2S}{R} \left(\mathbb{M}_u \frac{2V}{W} \right)^2 - \frac{1}{4} \Delta v^2 \mathbb{D}_u \frac{2V}{W} \left(\mathbb{M}_x \frac{2S}{R} \right)^2,$$

or alternatively

(6.4)
$$\frac{W + \Delta yV}{W - \Delta yV}(x; v_{+}) \frac{R + \Delta vS}{R - \Delta vS}(y_{-}; u) = \frac{W + \Delta yV}{W - \Delta yV}(x; v_{-}) \frac{R + \Delta vS}{R - \Delta vS}(y_{+}; u).$$

Proof. From the fact that the weight satisfies the over-determined system of first order divided-difference equations (4.1) and (6.1) we first compute that

(6.5)
$$\left[1 - \frac{1}{16}\Delta y^2 \Delta v^2 \mathbb{D}_x \frac{2S}{R} \mathbb{D}_u \frac{2V}{W}\right] \mathbb{M}_u \mathbb{D}_x w = \left[\mathbb{M}_u \frac{2V}{W} + \frac{1}{4}\Delta v^2 \mathbb{D}_u \frac{2V}{W} \mathbb{M}_x \frac{2S}{R}\right] \mathbb{M}_u \mathbb{M}_x w,$$

(6.6)
$$\left[1 - \frac{1}{16}\Delta y^2 \Delta v^2 \mathbb{D}_x \frac{2S}{R} \mathbb{D}_u \frac{2V}{W}\right] \mathbb{M}_x \mathbb{D}_u w = \left[\mathbb{M}_x \frac{2S}{R} + \frac{1}{4}\Delta y^2 \mathbb{D}_x \frac{2S}{R} \mathbb{M}_u \frac{2V}{W}\right] \mathbb{M}_x \mathbb{M}_u w,$$

using (2.11). Now we compute

(6.7)
$$\mathbb{D}_u \mathbb{D}_x w = \mathbb{D}_u \frac{2V}{W} \mathbb{M}_u \mathbb{M}_x w + \mathbb{M}_u \frac{2V}{W} \mathbb{D}_u \mathbb{M}_x w,$$

(6.8)
$$\mathbb{D}_x \mathbb{D}_u w = \mathbb{D}_x \frac{2S}{R} \mathbb{M}_x \mathbb{M}_u w + \mathbb{M}_x \frac{2S}{R} \mathbb{D}_x \mathbb{M}_u w,$$

using (2.10) and use the results of the previous set of equations to derive two independent relations linking $\mathbb{D}_u \mathbb{D}_x w$ and $\mathbb{M}_u \mathbb{M}_x w$. Comparison of these two latter relations leads to (6.3).

To establish (6.4) we note

(6.9)
$$w(y_+;v_+) = \frac{W + \Delta yV}{W - \Delta yV}(x;v_+)w(y_-;v_+) = \frac{W + \Delta yV}{W - \Delta yV}(x;v_+)\frac{R + \Delta vS}{R - \Delta vS}(y_-;u)w(y_-;v_-),$$

whereas

(6.10)
$$w(y_{+};v_{+}) = \frac{R + \Delta vS}{R - \Delta vS}(y_{+};u)w(y_{+};v_{-}) = \frac{R + \Delta vS}{R - \Delta vS}(y_{+};u)\frac{W + \Delta yV}{W - \Delta yV}(x;v_{-})w(y_{-};v_{-}).$$

Proposition 6.2. As a consequence of Definition 6.1 the Stieltjes transform satisfies the inhomogeneous equation

(6.11)
$$R\mathbb{D}_u f = 2S\mathbb{M}_u f + T.$$

The deformed \mathbb{D} -semi-classical class of orthogonal polynomial systems are characterised by the property that R(x; u), S(x; u) and T(x; u) in (6.11) are polynomials in x.

Proof. From the definition (3.19) and (6.1) we compute

(6.12)
$$\mathbb{D}_u f(x;u) = \int \mathbb{D}y \, \frac{\mathbb{M}_u w(y;u)}{x-y} \frac{2S(y;u)}{R(y;u)}$$

Now we observe that the rational function 2S(y; u)/[(x - y)R(y; u)] has the partial fraction expansion

(6.13)
$$\frac{2S(y;u)}{(x-y)R(y;u)} = \frac{2S(x;u)}{(x-y)R(x;u)} + \sum_{\substack{j,R(x_j)=0\\27}} \frac{2S(x_j;u)}{R'(x_j;u)(x-x_j)} \frac{1}{y-x_j}.$$

Consequently

(6.14)
$$\mathbb{D}_{u}f(x;u) = \frac{2S(x;u)}{R(x;u)} \int \mathbb{D}y \, \frac{\mathbb{M}_{u}w(y;u)}{x-y} + \sum_{j,R(x_{j})=0} \frac{2S(x_{j};u)}{R'(x_{j};u)(x-x_{j})} \int \mathbb{D}y \, \frac{\mathbb{M}_{u}w(y;u)}{y-x_{j}},$$

(6.15)
$$= \frac{2S(x;u)}{R(x;u)} \mathbb{M}_u f(x;u) - \sum_{j,R(x_j)=0} \frac{2S(x_j;u)}{R'(x_j;u)(x-x_j)} \mathbb{M}_u f(x_j;u).$$

We conclude that (6.11) follows with $\deg_x T \leq \deg_x R - 1$.

Proposition 6.3. Compatibility of (6.11) and (4.13) implies the following identity on U and T.

(6.16)
$$\Delta y \left[\frac{(W + \Delta y V)(x; v_{+})}{(W + \Delta y V)(x; v_{-})} (R + \Delta v S)(y_{-}; u) U(x; v_{-}) - (R - \Delta v S)(y_{-}; u) U(x; v_{+}) \right] \\ = \Delta v \left[(W + \Delta y V)(x; v_{+}) T(y_{-}; u) - (W - \Delta y V)(x; v_{+}) \frac{(R - \Delta v S)(y_{-}; u)}{(R - \Delta v S)(y_{+}; u)} T(y_{+}; u) \right].$$

Proof. We begin by defining the ratio

(6.17)
$$\chi \equiv \frac{(W + \Delta yV)(x; v_{+})}{(W + \Delta yV)(x; v_{-})} \frac{(R + \Delta vS)(y_{-}; u)}{(R + \Delta vS)(y_{+}; u)} = \frac{(W - \Delta yV)(x; v_{+})}{(W - \Delta yV)(x; v_{-})} \frac{(R - \Delta vS)(y_{-}; u)}{(R - \Delta vS)(y_{+}; u)}$$

by virtue of (6.4). Consider (4.13) in the form

(6.18)
$$(W - \Delta yV)(x;u)f(y_+;u) - (W + \Delta yV)(x;u)f(y_-;u) - \Delta yU(x;u) = 0,$$

and construct the combination $(W + \Delta yV)(x; v_+)/(W + \Delta yV)(x; v_-)(R + \Delta vS)(y_-; u) \times (6.18)(u \mapsto v_-) - (R - \Delta vS)(y_-; u) \times (6.18)(u \mapsto v_+)$. Using the above identity we find the two terms containing f possess the factors $(R + \Delta vS)(y_-; u)f(y_-; v_-) - (R - \Delta vS)(y_-; u)f(y_-; v_+)$ and $(R + \Delta vS)(y_+; u)f(y_+; v_-) - (R - \Delta vS)(y_+; u)f(y_+; v_+)$, into which we can apply (6.11) in the form

(6.19)
$$(R - \Delta vS)(x;u)f(x;v_{+}) - (R + \Delta vS)(x;u)f(x;v_{-}) - \Delta vT(x;u) = 0.$$

Then (6.16) immediately follows.

We can extend the notion of a generic or regular \mathbb{D} -semi-classical weight, given in Definition 4.2, to the deformed situation by the following definition

Definition 6.2. A regular, deformed \mathbb{D} -semi-classical weight is one that satisfies the requirements of the Definition 4.2 and $\deg_x R = \deg_x S = L$ and $\deg_x T = \deg_x R - 1$. Clearly M in Definition 4.2 and L are related, depending on the specific case on hand.

The analog of Proposition 4.2 is the following.

Proposition 6.4. Let the deformation coefficients $R_n(x; u)$, $\Gamma_n(x; u)$, $\Xi_n(x; u)$, $\Phi_n(x; u)$, $\Psi_n(x; u)$ be defined in terms of bilinear formulae involving products of the polynomials and associated functions by

$$(6.20) \quad \frac{2}{H_n} R_n = (R + \Delta vS) \left[-\frac{1}{a_n(v_-)} + p_{n-1}(;v_+)q_n(;v_-) - p_n(;v_+)q_{n-1}(;v_-) \right] \\ + (R - \Delta vS) \left[-\frac{1}{a_n(v_+)} + p_{n-1}(;v_-)q_n(;v_+) - p_n(;v_-)q_{n-1}(;v_+) \right],$$

(6.21)
$$\frac{\Delta v}{H_n} \Gamma_n = (R + \Delta vS) \left[\frac{1}{a_n(v_-)} - p_{n-1}(;v_+)q_n(;v_-) - p_n(;v_+)q_{n-1}(;v_-) \right] \\ + (R - \Delta vS) \left[-\frac{1}{a_n(v_+)} + p_{n-1}(;v_-)q_n(;v_+) + p_n(;v_-)q_{n-1}(;v_+) \right],$$

(6.22)
$$\frac{\Delta v}{H_n} \Xi_n = (R + \Delta vS) \left[\frac{1}{a_n(v_-)} + p_{n-1}(;v_+)q_n(;v_-) + p_n(;v_+)q_{n-1}(;v_-) \right] \\ + (R - \Delta vS) \left[-\frac{1}{a_n(v_+)} - p_{n-1}(;v_-)q_n(;v_+) - p_n(;v_-)q_{n-1}(;v_+) \right],$$

г		
L		
L		
L		

(6.23)
$$\frac{\Delta v}{2H_n} \Phi_n = (R + \Delta vS)p_n(;v_+)q_n(;v_-) - (R - \Delta vS)p_n(;v_-)q_n(;v_+),$$

and

(6.24)
$$\frac{\Delta v}{2H_n} \Psi_n = -(R + \Delta vS)p_{n-1}(;v_+)q_{n-1}(;v_-) + (R - \Delta vS)p_{n-1}(;v_-)q_{n-1}(;v_+),$$

for $n \ge 0$ and where the decoupling factor $H_n(u)$ has $\deg_x H_n = 0$. Then the deformation coefficients of the deformed \mathbb{D} -semi-classical class are polynomials in x.

Proof. We offer a proof in the spirit of that for Proposition 4.2, using Laguerre's method, and apply it to (6.23) first. From (6.11) and (3.21) we have

$$0 = (R - \Delta vS)f(;v_{+}) - (R + \Delta vS)f(;v_{-}) - \Delta vT,$$

$$= (R - \Delta vS)\frac{q_{n} + p_{n-1}^{(1)}}{p_{n}}(;v_{+}) - (R + \Delta vS)\frac{q_{n} + p_{n-1}^{(1)}}{p_{n}}(;v_{-}) - \Delta vT,$$

$$= \frac{(R - \Delta vS)p_{n}(;v_{-})p_{n-1}^{(1)}(;v_{+}) - (R + \Delta vS)p_{n}(;v_{+})p_{n-1}^{(1)}(;v_{-}) - \Delta vTp_{n}(;v_{+})p_{n}(;v_{-})}{p_{n}(;v_{+})p_{n}(;v_{-})}$$

$$(6.25) \qquad + \frac{(R - \Delta vS)p_{n}(;v_{-})q_{n}(;v_{+}) - (R + \Delta vS)p_{n}(;v_{+})q_{n}(;v_{-})}{p_{n}(;v_{+})p_{n}(;v_{-})}.$$

In this last expression the numerator of the first term is clearly a polynomial in x which implies the numerator of the second term is likewise. This latter numerator is an odd function under the exchange $v_+ \leftrightarrow v_-$ and we denote it by $\Delta v \Phi_n(x; u)/(2H_n)$, thus deriving (6.23). The relation (6.24) is simply the case of (6.23) under $n \mapsto n-1$.

The remaining relations may now be derived from the following argument. From the workings of the previous paragraph we know that

(6.26)
$$\frac{\Delta v \Phi_n}{2H_n} = (R - \Delta v S) p_n(; v_-) p_{n-1}^{(1)}(; v_+) - (R + \Delta v S) p_n(; v_+) p_{n-1}^{(1)}(; v_-) - \Delta v T p_n(; v_+) p_n(; v_-).$$

By multiplying the left-hand side of this relation by the Casoratian $a_n(v_+)[p_{n-1}p_{n-1}^{(1)} - p_np_{n-2}^{(1)}](;v_+) = 1$ we observe that the resulting equation separates into two terms with the factorisation

$$(6.27) \quad p_n(;v_+) \left[-a_n(v_+) \frac{\Delta v \Phi_n}{2H_n} p_{n-2}^{(1)}(;v_+) + (R + \Delta vS) p_{n-1}^{(1)}(;v_-) + \Delta vT p_n(;v_-) \right] \\ = p_{n-1}^{(1)}(;v_+) \left[-a_n(v_+) \frac{\Delta v \Phi_n}{2H_n} p_{n-1}(;v_+) + (R - \Delta vS) p_n(;v_-) \right],$$

which implies that this expression contains the polynomial factors $p_n(;v_+)p_{n-1}^{(1)}(;v_+)$ and therefore can be written as $p_n(;v_+)p_{n-1}^{(1)}(;v_+)\pi_1$, where π_1 is a polynomial in x. Given that the two factors $p_n(;v_+), p_{n-1}^{(1)}(;v_+)$ are non-zero this then leads to two evaluations for π_1 , which by constructing a suitable combination of these and employing the Casoratian once more we find, upon some simplification,

(6.28)
$$\pi_1 = a_n(v_+) \left[-(R + \Delta vS)p_{n-1}(;v_+)q_n(;v_-) + (R - \Delta vS)p_n(;v_-)q_{n-1}(;v_+) \right].$$

Alternatively we could have multiplied the left-hand side of (6.26) by the Casoratian $a_n(v_-)[p_{n-1}p_{n-1}^{(1)} - p_n p_{n-2}^{(1)}](;v_-) = 1$ and deduced the factorisation

(6.29)
$$p_{n}(;v_{-})\left[-a_{n}(v_{-})\frac{\Delta v\Phi_{n}}{2H_{n}}p_{n-2}^{(1)}(;v_{-}) - (R - \Delta vS)p_{n-1}^{(1)}(;v_{+}) + \Delta vTp_{n}(;v_{+})\right]$$
$$= p_{n-1}^{(1)}(;v_{-})\left[-a_{n}(v_{-})\frac{\Delta v\Phi_{n}}{2H_{n}}p_{n-1}(;v_{-}) - (R + \Delta vS)p_{n}(;v_{+})\right],$$

This fact then allows us to conclude that either of these expressions can be written as $p_n(;v_-)p_{n-1}^{(1)}(;v_-)\pi_2$, where π_2 is another polynomial in x. Using an identical procedure to that employed above we can infer that

(6.30)
$$\pi_2 = a_n(v_-) \left[-(R + \Delta vS)p_n(;v_+)q_{n-1}(;v_-) + (R - \Delta vS)p_{n-1}(;v_-)q_n(;v_+) \right].$$

Relations (6.20) and (6.21) then follow from the definitions

$$(6.31) \qquad 2R_n - \Delta v \Gamma_n + 2H_n \frac{R + \Delta v S}{a_n(v_-)} = -\frac{2H_n}{a_n(v_+)} \pi_1, \quad 2R_n + \Delta v \Gamma_n + 2H_n \frac{R - \Delta v S}{a_n(v_+)} = \frac{2H_n}{a_n(v_-)} \pi_2,$$

whereas (6.22) follows from $\Delta v (\Gamma_n + \Xi_n)/2H_n = (R + \Delta v S)/a_n(v_-) - (R - \Delta v S)/a_n(v_+).$

Analogous to the large x expansions of the spectral coefficients we have the following expansions for the deformation coefficients.

Proposition 6.5. Let $L = \max(\deg_x R, \deg_x S)$. As $x \to \infty$ we have the leading orders of the terminating expansions of the deformation coefficients

(6.32)
$$\frac{2}{H_n}R_n = -(\gamma_n(v_+) + \gamma_n(v_-)) \left[\frac{R - \Delta vS}{\gamma_{n-1}(v_+)} + \frac{R + \Delta vS}{\gamma_{n-1}(v_-)} \right] + \sum_{i=0}^{n-1} (b_i(v_+) - b_i(v_-)) \left[(R + \Delta vS) \frac{\gamma_n(v_+)}{\gamma_{n-1}(v_-)} - (R - \Delta vS) \frac{\gamma_n(v_-)}{\gamma_{n-1}(v_+)} \right] x^{-1} + \mathcal{O}(x^{L-2}) \quad n \ge 0,$$

$$(6.33) \quad \frac{\Delta v}{2H_n} \Phi_n = \left[(R + \Delta vS) \frac{\gamma_n(v_+)}{\gamma_n(v_-)} - (R - \Delta vS) \frac{\gamma_n(v_-)}{\gamma_n(v_+)} \right] x^{-1} \\ + \left((R + \Delta vS) \frac{\gamma_n(v_+)}{\gamma_n(v_-)} \left[\sum_{i=0}^n b_i(v_-) - \sum_{i=0}^{n-1} b_i(v_+) \right] + (R - \Delta vS) \frac{\gamma_n(v_-)}{\gamma_n(v_+)} \left[\sum_{i=0}^{n-1} b_i(v_-) - \sum_{i=0}^n b_i(v_+) \right] \right) x^{-2} \\ + O(x^{L-3}), \quad n \ge 0,$$

$$(6.34) \quad \frac{\Delta v}{2H_n} \Psi_n = -\left[(R + \Delta vS) \frac{\gamma_{n-1}(v_+)}{\gamma_{n-1}(v_-)} - (R - \Delta vS) \frac{\gamma_{n-1}(v_-)}{\gamma_{n-1}(v_+)} \right] x^{-1} \\ - \left((R + \Delta vS) \frac{\gamma_{n-1}(v_+)}{\gamma_{n-1}(v_-)} \left[\sum_{i=0}^{n-1} b_i(v_-) - \sum_{i=0}^{n-2} b_i(v_+) \right] + (R - \Delta vS) \frac{\gamma_{n-1}(v_-)}{\gamma_{n-1}(v_+)} \left[\sum_{i=0}^{n-2} b_i(v_-) - \sum_{i=0}^{n-1} b_i(v_+) \right] \right) x^{-2} \\ + O(x^{L-3}), \quad n \ge 0,$$

(6.35)
$$\frac{\Delta v}{H_n} \Gamma_n = (\gamma_n(v_-) - \gamma_n(v_+)) \left[\frac{R + \Delta vS}{\gamma_{n-1}(v_-)} + \frac{R - \Delta vS}{\gamma_{n-1}(v_+)} \right] \\ + \sum_{i=0}^{n-1} (b_i(v_+) - b_i(v_-)) \left[(R + \Delta vS) \frac{\gamma_n(v_+)}{\gamma_{n-1}(v_-)} + (R - \Delta vS) \frac{\gamma_n(v_-)}{\gamma_{n-1}(v_+)} \right] x^{-1} + \mathcal{O}(x^{L-2}), \quad n \ge 0,$$

and

(6.36)
$$\frac{\Delta v}{H_n} \Xi_n = (\gamma_n(v_-) + \gamma_n(v_+)) \left[\frac{R + \Delta vS}{\gamma_{n-1}(v_-)} - \frac{R - \Delta vS}{\gamma_{n-1}(v_+)} \right] \\ - \sum_{i=0}^{n-1} (b_i(v_+) - b_i(v_-)) \left[(R + \Delta vS) \frac{\gamma_n(v_+)}{\gamma_{n-1}(v_-)} + (R - \Delta vS) \frac{\gamma_n(v_-)}{\gamma_{n-1}(v_+)} \right] x^{-1} + \mathcal{O}(x^{L-2}), \quad n \ge 0.$$

Proof. These formulae follow from the substitution of the expansions (3.34) and (3.35) into the definitions (6.20-6.24).

Remark 6.1. Unlike the case of the spectral coefficients (see Proposition 4.3) the leading terms of the deformation coefficients are not determined by the weight data alone but depend non-trivially upon the three-term recurrence coefficients.

Corollary 6.1. In the regular, deformed \mathbb{D} -semi-classical case the deformation coefficients $R_n, \Gamma_n, \Phi_n, \Psi_n, \Xi_n$ are polynomials in the spectral variable x with degrees independent of n, $\deg_x R_n = \deg_x \Gamma_n = \deg_x \Xi_n = \max(\deg_x R, \deg_x S)$ and $\deg_x \Phi_n = \deg_x \Psi_n = \max(\deg_x R, \deg_x S) - 1$.

The expressions for the deformation coefficients (6.24-6.24) can be inverted yielding a system of linear divided-difference equations in the deformation variable for the polynomials and associated functions.

Proposition 6.6. The OPS corresponding to a deformed \mathbb{D} -semi-classical weight satisfies the deformation divided-difference equation

(6.37)
$$\mathbb{D}_{u}Y_{n} := B_{n}\mathbb{M}_{u}Y_{n} = \frac{1}{R_{n}}\begin{pmatrix}\Gamma_{n} & \Phi_{n}\\\Psi_{n} & \Xi_{n}\end{pmatrix}\mathbb{M}_{u}Y_{n}, \quad n \ge 0.$$

Proof. The essence of this result involves the inversion of Proposition 6.4 which is carried out in a manner analogous to the proof of Proposition 4.4.

Of the four coefficients $\Gamma_n, \Phi_n, \Psi_n, \Xi_n$ only two are independent because of the following relations.

Proposition 6.7. The deformation coefficients satisfy the linear identity

(6.38)
$$\Psi_n = -\frac{a_n}{a_{n-1}} \Phi_{n-1}, \quad n \ge 1,$$

and the trace identity

(6.39)
$$\Delta v \left(\Gamma_n + \Xi_n\right) = 2H_n \left[\frac{R + \Delta vS}{a_n(v_-)} - \frac{R - \Delta vS}{a_n(v_+)}\right], \quad n \ge 0.$$

Proof. The first relation follows by comparison of (6.24) with (6.23) and setting $H_n/H_{n-1} = a_n/a_{n-1}$. The second identity is easily seen from the expressions (6.21) and (6.22) and has already been observed in the conclusions of the workings for the proof of Proposition 6.4.

The initial values of the deformation coefficients are

(6.40)
$$R_0 = -\frac{1}{2}H_0[\gamma_0(v_-) + \gamma_0(v_+)] \left(\frac{R + \Delta vS}{a_0(v_-)\gamma_0(v_-)} + \frac{R - \Delta vS}{a_0(v_+)\gamma_0(v_+)}\right)$$

(6.41)
$$\Gamma_0 = \frac{1}{\Delta v} H_0[\gamma_0(v_-) - \gamma_0(v_+)] \left(\frac{R + \Delta vS}{a_0(v_-)\gamma_0(v_-)} + \frac{R - \Delta vS}{a_0(v_+)\gamma_0(v_+)} \right)$$

(6.42)
$$\Xi_0 = \frac{1}{\Delta v} H_0[\gamma_0(v_-) + \gamma_0(v_+)] \left(\frac{R + \Delta vS}{a_0(v_-)\gamma_0(v_-)} - \frac{R - \Delta vS}{a_0(v_+)\gamma_0(v_+)} \right)$$

(6.43)
$$\Phi_{-1} = 0, \qquad \Phi_0 = -2H_0\gamma_0(v_-)\gamma_0(v_+)T,$$

 $\Psi_0 = 0.$ (6.44)

Compatibility between the matrix recurrence relation and the deformation divided-difference equation implies the next result.

Proposition 6.8. The recurrence matrix and the deformation matrix satisfy

(6.45)
$$B_{n+1} \cdot \mathbb{M}_u K_n - \mathbb{M}_u K_n \cdot B_n = \mathbb{D}_u K_n - \frac{1}{4} \Delta v^2 B_{n+1} \cdot \mathbb{D}_u K_n \cdot B_n, \quad n \ge 0,$$

or equivalently

(6.46)
$$K_n(;v_+) \left(1 - \frac{1}{2}\Delta v B_n\right)^{-1} \left(1 + \frac{1}{2}\Delta v B_n\right) = \left(1 - \frac{1}{2}\Delta v B_{n+1}\right)^{-1} \left(1 + \frac{1}{2}\Delta v B_{n+1}\right) K_n(;v_-), \quad n \ge 0,$$

Proof. The first form of the compatibility relation is deduced by comparing

$$\mathbb{D}_{u}Y_{n+1} = B_{n+1}\mathbb{M}_{u}Y_{n+1}$$

= $B_{n+1}\mathbb{M}_{u}(K_{n}Y_{n})$
= $B_{n+1}\left(\mathbb{M}_{u}K_{n}\cdot\mathbb{M}_{u}Y_{n} + \frac{1}{4}\Delta v^{2}\mathbb{D}_{u}K_{n}\cdot\mathbb{D}_{u}Y_{n}\right)$
= $B_{n+1}\left(\mathbb{M}_{u}K_{n} + \frac{1}{4}\Delta v^{2}\mathbb{D}_{u}K_{n}\cdot B_{n}\right)\mathbb{M}_{u}Y_{n},$

and

$$(6.47) \qquad \qquad \mathbb{D}_u Y_{n+1} = \mathbb{D}_u K_n Y_n = (\mathbb{D}_u K_n \cdot \mathbb{M}_u Y_n + \mathbb{M}_u K_n \cdot \mathbb{D}_u Y_n) = (\mathbb{D}_u K_n + \mathbb{M}_u K_n \cdot B_n) \mathbb{M}_u Y_n.$$

The second form is found by computing $Y_{n+1}(x;v_+)$ in terms of $Y_n(x;v_-)$ in the two possible ways corresponding to the orders of the operations $n \mapsto n+1$ and $v_- \mapsto v_+$. Compatibility of the three term recurrence relation and the deformation divided-difference equation implies the following result which can be found by considering the representations of the deformation coefficients given in Proposition 6.4.

Proposition 6.9. The deformation coefficients R_n, Γ_n, Φ_n satisfy recurrence relations in n,

(6.48)
$$\frac{a_{n+1}(v_{-})}{H_{n+1}} (-2R_{n+1} + \Delta v \Gamma_{n+1}) + \frac{a_n(v_{-})}{H_n} (2R_n + \Delta v \Gamma_n) = -[x - b_n(v_{-})] \frac{\Delta v}{H_n} \Phi_n + 2a_n(v_{-}) \left(\frac{R + \Delta v S}{a_n(v_{-})} - \frac{R - \Delta v S}{a_n(v_{+})}\right), \quad n \ge 0,$$

(6.49)
$$\frac{a_{n+1}(v_{+})}{H_{n+1}}(2R_{n+1} + \Delta v\Gamma_{n+1}) + \frac{a_n(v_{+})}{H_n}(-2R_n + \Delta v\Gamma_n) = -[x - b_n(v_{+})]\frac{\Delta v}{H_n}\Phi_n + 2a_n(v_{+})\left(\frac{R + \Delta vS}{a_n(v_{-})} - \frac{R - \Delta vS}{a_n(v_{+})}\right), \quad n \ge 0,$$

Recurrences (6.48) and (6.49) are the analogs of combinations of (4.43) and (4.44).

Proof. The simplest proof of these relations is the one employing the definitions (6.20, 6.21, 6.23) of Proposition 6.4. Taking the first relation we deduce

$$\begin{split} & \frac{a_{n+1}(v_{-})}{H_{n+1}} (-2R_{n+1} + \Delta v \Gamma_{n+1}) + \frac{a_n(v_{-})}{H_n} (2R_n + \Delta v \Gamma_n) + [x - b_n(v_{-})] \frac{\Delta v}{H_n} \Phi_n \\ &= 2(R + \Delta vS) \left[1 - p_n(;v_{+})(a_{n+1}(v_{-})q_{n+1}(;v_{-}) + b_n(v_{-})q_n(v_{-}) + a_n(v_{-})q_{n-1}(v_{-}) - xq_n(v_{-}))\right] \\ &- 2(R - \Delta vS) \left[\frac{a_n(v_{-})}{a_n(v_{+})} - q_n(;v_{+})(a_{n+1}(v_{-})p_{n+1}(;v_{-}) + b_n(v_{-})p_n(v_{-}) + a_n(v_{-})p_{n-1}(v_{-}) - xp_n(v_{-}))\right] \\ &= 2a_n(v_{-}) \left(\frac{R + \Delta vS}{a_n(v_{-})} - \frac{R - \Delta vS}{a_n(v_{+})}\right), \end{split}$$

where essential use of the three-term recurrence relations (3.5, 3.22) has been made. The second relation can be found in an identical manner.

We find bilinear identities for the deformation matrix analogous to those of Proposition 4.7.

Proposition 6.10. The deformation coefficients satisfy the bilinear or determinantal identity

(6.50)
$$R_n^2 + \frac{1}{4}\Delta v^2 \left[\Gamma_n \Xi_n - \Phi_n \Psi_n\right] = -H_n R_n \left[\frac{R + \Delta vS}{a_n(v_-)} + \frac{R - \Delta vS}{a_n(v_+)}\right], \quad n \ge 0$$

Proof. A direct method of proof is possible substituting the expressions for the deformation coefficients $R_n, \Gamma_n, \Xi_n, \Phi_n, \Psi_n$ in terms of products of polynomials and associated functions as given by (6.20-6.24) into the left-hand side of the (6.50). After expansion and considerable cancellation we recognise the the form for the right-hand side.

From the above result we can deduce matrix identities analogous to Corollary 4.3.

Corollary 6.2. The matrix factor appearing in (6.46) has the determinant evaluation

(6.51)
$$\det(1 \pm \frac{1}{2}\Delta v B_n) = -\frac{2H_n(R \mp \Delta v S)}{R_n a_n(v_{\pm})}, \quad n \ge 0,$$

and its inverse is

(6.52)
$$(1 \pm \frac{1}{2}\Delta v B_n)^{-1} = -\frac{a_n(v_{\pm})}{2H_n(R \mp \Delta v S)} \begin{pmatrix} R_n \pm \frac{1}{2}\Delta v \Xi_n & \mp \frac{1}{2}\Delta v \Phi_n \\ \mp \frac{1}{2}\Delta v \Psi_n & R_n \pm \frac{1}{2}\Delta v \Gamma_n \end{pmatrix}, \quad n \ge 0.$$

Proof. We note from the matrix formula for B_n (6.37) that

$$\det(1 \pm \frac{1}{2}\Delta v B_n) = \frac{1}{R_n^2} \left\{ R_n^2 + \frac{1}{4}\Delta v^2 \left[\Gamma_n \Xi_n - \Phi_n \Psi_n\right] \pm \frac{1}{2}\Delta v R_n(\Gamma_n + \Xi_n) \right\}$$
$$= -\frac{2H_n}{R_n} \frac{R \mp \Delta v S}{a_n(v_{\pm})},$$

where we have used (6.50) and (6.39) in the last step. The inversion formula follows from this last result and the standard formula for an inverse.

Consequent to the results of Corollary 6.2 we have the following expression for the matrix product appearing in (6.46).

Corollary 6.3. The matrix product (6.46) has the evaluation

$$(6.53) \quad \left(1 - \frac{1}{2}\Delta v B_{n}\right)^{-1} \left(1 + \frac{1}{2}\Delta v B_{n}\right) \\ = \frac{a_{n}(v_{-})}{2H_{n}(R + \Delta v S)} \begin{pmatrix} 2R_{n} + 2H_{n} \frac{R - \Delta v S}{a_{n}(v_{+})} + \Delta v \Gamma_{n} & \Delta v \Phi_{n} \\ \Delta v \Psi_{n} & 2R_{n} + 2H_{n} \frac{R + \Delta v S}{a_{n}(v_{-})} - \Delta v \Gamma_{n}. \end{pmatrix}, \quad n \ge 0 \\ = -a_{n}(v_{-}) \begin{pmatrix} p_{n}(;v_{+}) \\ p_{n-1}(;v_{+}) \end{pmatrix} \otimes (q_{n-1}(;v_{-}), \quad -q_{n}(;v_{-})) + a_{n}(v_{-}) \frac{R - \Delta v S}{R + \Delta v S} \begin{pmatrix} q_{n}(;v_{+}) \\ q_{n-1}(;v_{+}) \end{pmatrix} \otimes (p_{n-1}(;v_{-}), \quad -p_{n}(;v_{-})) \end{pmatrix}$$

Proof. Using the inverse (6.52) we form the matrix product and employ (6.50) to simplify the diagonal elements, with the result given by (6.53).

This result also motivates the following definitions

(6.54)
$$\mathfrak{R}_{+} := 2R_n + 2H_n \frac{R - \Delta v S}{a_n(v_+)} + \Delta v \Gamma_n,$$

(6.55)
$$\mathfrak{R}_{-} := 2R_n + 2H_n \frac{R + \Delta v S}{a_n(v_-)} - \Delta v \Gamma_n$$

(6.56)
$$\mathfrak{P}_{+} := -\Delta v \, \Phi_{n}, \quad \mathfrak{P}_{-} := \Delta v \, \Psi_{n},$$

valid for $n \ge 1$, and for n = 0 we have

(6.57)
$$\mathfrak{R}_{+}(n=0) := -2\frac{H_{0}}{a_{0}(v_{-})}\frac{\gamma_{0}(v_{+})}{\gamma_{0}(v_{-})}(R+\Delta v\,S),$$

(6.58)
$$\mathfrak{R}_{-}(n=0) := -2\frac{H_{0}}{a_{0}(v_{+})}\frac{\gamma_{0}(v_{-})}{\gamma_{0}(v_{+})}(R - \Delta v S),$$

(6.59)
$$\mathfrak{P}_{+}(n=0) := 2\Delta v H_{0} \gamma_{0}(v_{+}) \gamma_{0}(v_{-}) T, \quad \mathfrak{P}_{-}(n=0) := 0,$$

together with

(6.60)
$$B_n^* := \begin{pmatrix} \mathfrak{R}_+ & -\mathfrak{P}_+\\ \mathfrak{P}_- & \mathfrak{R}_- \end{pmatrix}.$$

In these variables the deformation bi-linear relation (6.50) takes the form

(6.61)
$$\det B_n^* = \mathfrak{R}_+ \mathfrak{R}_- + \mathfrak{P}_+ \mathfrak{P}_- = \frac{4H_n^2}{a_n(v_+)a_n(v_-)} (R^2 - \Delta v^2 S^2).$$

It is of advantage to write out recurrence-deformation compatibility equations in terms of $\mathfrak{R}_{\pm,n}, \mathfrak{P}_{\pm,n}$, where again we append a subscript to indicate the dependence on the index n. We will henceforth denote variables evaluated at the advanced and retarded deformation variable by $\hat{a}_n = a_n(v_+), \check{a}_n = a_n(v_-)$. **Corollary 6.4.** Solving (6.46) for B_{n+1}^* we deduce

(6.62)

$$\frac{H_n}{H_{n+1}}\hat{a}_{n+1}\check{a}_{n+1}\mathfrak{P}_{+,n+1} = -\check{a}_n(x-\hat{b}_n)\mathfrak{R}_{+,n} + \hat{a}_n(x-\hat{b}_n)\mathfrak{R}_{-,n} + (x-\hat{b}_n)(x-\check{b}_n)\mathfrak{P}_{+,n} + \hat{a}_n\check{a}_n\mathfrak{P}_{-,n} \\
3) \frac{H_n}{H_{n+1}}\mathfrak{P}_{-,n+1} = \mathfrak{P}_{+,n},$$

(6.63)

(6.64)
$$\frac{H_n}{H_{n+1}}\mathfrak{R}_{+,n+1} = \frac{\hat{a}_n}{\hat{a}_{n+1}}\mathfrak{R}_{-,n} + \frac{x - \hat{b}_n}{\hat{a}_{n+1}}\mathfrak{P}_{+,n},$$

(6.65)
$$\frac{H_n}{H_{n+1}}\mathfrak{R}_{-,n+1} = \frac{\check{a}_n}{\check{a}_{n+1}}\mathfrak{R}_{+,n} - \frac{x-b_n}{\check{a}_{n+1}}\mathfrak{P}_{+,n}.$$

The compatibility relation between the spectral and deformation divided-difference equations takes the following form.

Proposition 6.11. The spectral matrix $A_n(x; u)$ and the deformation matrix $B_n(x; u)$ satisfy the \mathbb{D} -Schlesinger equation

$$(6.66) \quad \mathbb{D}_{u}A_{n} - \mathbb{D}_{x}B_{n} + \mathbb{M}_{u}A_{n} \cdot \mathbb{M}_{x}B_{n} - \mathbb{M}_{x}B_{n} \cdot \mathbb{M}_{u}A_{n}$$

$$= \frac{1}{4}\Delta v^{2}\mathbb{M}_{x}B_{n} \cdot \mathbb{D}_{u}A_{n} \cdot \left(1 - \frac{1}{16}\Delta y^{2}\Delta v^{2}\mathbb{D}_{x}B_{n} \cdot \mathbb{D}_{u}A_{n}\right)^{-1} \left(\mathbb{M}_{x}B_{n} + \frac{1}{4}\Delta y^{2}\mathbb{D}_{x}B_{n} \cdot \mathbb{M}_{u}A_{n}\right)$$

$$- \frac{1}{4}\Delta y^{2}\mathbb{M}_{u}A_{n} \cdot \mathbb{D}_{x}B_{n} \cdot \left(1 - \frac{1}{16}\Delta y^{2}\Delta v^{2}\mathbb{D}_{u}A_{n} \cdot \mathbb{D}_{x}B_{n}\right)^{-1} \left(\mathbb{M}_{u}A_{n} + \frac{1}{4}\Delta v^{2}\mathbb{D}_{u}A_{n} \cdot \mathbb{M}_{x}B_{n}\right),$$

or the equivalent form

$$(6.67) \quad \left(1 - \frac{1}{2}\Delta y A_n(;v_+)\right)^{-1} \left(1 + \frac{1}{2}\Delta y A_n(;v_+)\right) \left(1 - \frac{1}{2}\Delta v B_n(y_-;)\right)^{-1} \left(1 + \frac{1}{2}\Delta v B_n(y_-;)\right) \\ = \left(1 - \frac{1}{2}\Delta v B_n(y_+;)\right)^{-1} \left(1 + \frac{1}{2}\Delta v B_n(y_+;)\right) \left(1 - \frac{1}{2}\Delta y A_n(;v_-)\right)^{-1} \left(1 + \frac{1}{2}\Delta y A_n(;v_-)\right).$$

Proof. The first form follows from $\mathbb{D}_x \mathbb{D}_u Y_n = \mathbb{D}_u \mathbb{D}_x Y_n$ and the Leibniz formulae of (2.10) and (2.11), from which we deduce

$$\begin{bmatrix} 1 - \frac{1}{16}\Delta y^2 \Delta v^2 \mathbb{D}_u A_n \cdot \mathbb{D}_x B_n \end{bmatrix} \mathbb{D}_x \mathbb{M}_u Y = \begin{bmatrix} \mathbb{M}_u A_n + \frac{1}{4}\Delta v^2 \mathbb{D}_u A_n \cdot \mathbb{M}_x B_n \end{bmatrix} \mathbb{M}_u \mathbb{M}_x Y,$$
$$\begin{bmatrix} 1 - \frac{1}{16}\Delta y^2 \Delta v^2 \mathbb{D}_x B_n \cdot \mathbb{D}_u A_n \end{bmatrix} \mathbb{D}_u \mathbb{M}_x Y = \begin{bmatrix} \mathbb{M}_x B_n + \frac{1}{4}\Delta y^2 \mathbb{D}_x B_n \cdot \mathbb{M}_u A_n \end{bmatrix} \mathbb{M}_x \mathbb{M}_u Y.$$

Using equivalent forms of the spectral and deformation divided-difference equations (4.35) and (6.37) we see

$$Y(y_{+};v_{+}) = \left(1 - \frac{1}{2}\Delta y A_{n}(;v_{+})\right)^{-1} \left(1 + \frac{1}{2}\Delta y A_{n}(;v_{+})\right) Y(y_{-};v_{+})$$

= $\left(1 - \frac{1}{2}\Delta y A_{n}(;v_{+})\right)^{-1} \left(1 + \frac{1}{2}\Delta y A_{n}(;v_{+})\right) \left(1 - \frac{1}{2}\Delta v B_{n}(y_{-};)\right)^{-1} \left(1 + \frac{1}{2}\Delta v B_{n}(y_{-};)\right) Y(y_{-};v_{-}),$

whereas

$$Y(y_{+};v_{+}) = \left(1 - \frac{1}{2}\Delta v B_{n}(y_{+};)\right)^{-1} \left(1 + \frac{1}{2}\Delta v B_{n}(y_{+};)\right) Y(y_{+};v_{-})$$

= $\left(1 - \frac{1}{2}\Delta v B_{n}(y_{+};)\right)^{-1} \left(1 + \frac{1}{2}\Delta v B_{n}(y_{+};)\right) \left(1 - \frac{1}{2}\Delta y A_{n}(;v_{-})\right)^{-1} \left(1 + \frac{1}{2}\Delta y A_{n}(;v_{-})\right) Y(y_{-};v_{-}).$
his gives us the form (6.67).

This gives us the form (6.67).

Remark 6.2. Let us denote the two fixed points of the x-lattice by x_L and x_R . By analogy with the linear lattices let us conjecture the existence of fundamental solutions to (4.35) about $x = x_L, x_R$ which we denote by Y_L, Y_R respectively. Furthermore let us define the connection matrix

(6.68)
$$P(x;u) := (Y_R(x;u))^{-1} Y_L(x;u).$$

From (4.35) it is clear that P is a \mathbb{D} -constant function with respect to x, that is to say

$$P(y_+;u) = P(y_-;u).$$

In addition it is clear from (6.37) that this type of deformation is also a connection preserving deformation in the sense that

(6.69)
$$P(x; v_{+}) = P(x; v_{-}).$$

The compatibility relation (6.67) $\chi B_n^*(y_+; u) A_n^*(x; v_-) = A_n^*(x; v_+) B_n^*(y_-; u)$ may be written componentwise in the new variables in the more practical form as

$$(6.70) \qquad \chi \left[\mathfrak{W}_{+}(x;v_{-})\mathfrak{R}_{+}(y_{+};u) - \mathfrak{T}_{-}(x;v_{-})\mathfrak{P}_{+}(y_{+};u) \right] = \mathfrak{W}_{+}(x;v_{+})\mathfrak{R}_{+}(y_{-};u) - \mathfrak{T}_{+}(x;v_{+})\mathfrak{P}_{-}(y_{-};u) + \mathfrak{T}_{+}(x;v_{+})\mathfrak{R}_{+}(y_{+};u) \right] = \mathfrak{W}_{+}(x;v_{+})\mathfrak{R}_{+}(y_{+};u) - \mathfrak{T}_{+}(x;v_{+})\mathfrak{R}_{+}(y_{+};u) = \mathfrak{W}_{+}(x;v_{+})\mathfrak{R}_{+}(y_{+};u) + \mathfrak{T}_{+}(y_{+};u) = \mathfrak{W}_{+}(y_{+};u) + \mathfrak{W}_{+}(y_{+};u) = \mathfrak{W}_{+}(y$$

 $(6.71) \qquad \chi \left[\mathfrak{T}_{+}(x;v_{-})\mathfrak{R}_{+}(y_{+};u) + \mathfrak{W}_{-}(x;v_{-})\mathfrak{P}_{+}(y_{+};u) \right] = \mathfrak{T}_{+}(x;v_{+})\mathfrak{R}_{-}(y_{-};u) + \mathfrak{W}_{+}(x;v_{+})\mathfrak{P}_{+}(y_{-};u),$

$$(6.72) \qquad \chi \left[\mathfrak{T}_{-}(x;v_{-})\mathfrak{R}_{-}(y_{+};u) + \mathfrak{W}_{+}(x;v_{-})\mathfrak{P}_{-}(y_{+};u) \right] = \mathfrak{T}_{-}(x;v_{+})\mathfrak{R}_{+}(y_{-};u) + \mathfrak{W}_{-}(x;v_{+})\mathfrak{P}_{-}(y_{-};u),$$

$$(6.73) \qquad \chi \left[\mathfrak{W}_{-}(x;v_{-})\mathfrak{R}_{-}(y_{+};u) - \mathfrak{T}_{+}(x;v_{-})\mathfrak{P}_{-}(y_{+};u) \right] = \mathfrak{W}_{-}(x;v_{+})\mathfrak{R}_{-}(y_{-};u) - \mathfrak{T}_{-}(x;v_{+})\mathfrak{P}_{+}(y_{-};u),$$

where χ is defined by (6.17). The real content of (6.70-6.73) lie in the case $n \ge 1$ as the n = 0 evaluations of (6.72) is trivially satisfied, whereas that of (6.70) and (6.73) are identically satisfied due to the definition (6.17), and (6.71) is equivalent to the consistency identity (6.16).

Having assembled all of the ingredients of our theory we have to draw them together and to perform three tasks. The first is to parameterise the spectral coefficients in a minimal way consistent with the constraints of the spectral structures, namely (4.46). The second task is to close this system, that is to say relate the deformation coefficients to the spectral coefficients, preferably with the parameterisation found in the first step. The third task is to derive the dynamical equations for this parameterisation with respect to the deformation variable. There are two avenues of approach to the problem of extracting useful information from the above compatibility relations (6.70-6.73). One way is to clear the denominators on both sides and resolve the resulting expressions in terms of $\bigoplus_{k\geq 0} x^k + \Delta y \bigoplus_{k\geq 0} x^k$. This is useful for certain results, as we shall see in the application. The other method is to work with the evaluation of the spectral and deformation coefficients at certain key ordinates and construct a parameterisation based upon these variables. A fundamental and crucial role will be played by the zeros of the polynomials $W^2 - \Delta y^2 V^2$, Θ_n and χ as these ordinates, however these cannot be interpreted as singularities in the spectral plane.

Remark 6.3. The compatibility relations (6.70-6.73) are satisfied identically at the common zeros of $(W^2 - \Delta y^2 V^2)(\tilde{x}; v_{\pm})$ in the sense that both sides of the relations are identically zero, that is to say at those fixed zeros independent of u. In addition one can show that the left-hand side of the bilinear relation (6.61) vanishes at the movable zeros in either of the two above cases. This implies that when $(W^2 - \Delta y^2 V^2)(\tilde{x}; v_{\pm}) = 0$ then $(R^2 - \Delta v^2 S^2)(\tilde{y}_{\pm}; u) = 0$.

7. M = 3, L = 1 case and deformation of the Askey-Wilson OPS

The previous sections treated the D-semi-classical orthogonal polynomial system with general divideddifference operators D, M and arbitrary degrees M, L for the spectral and deformation coefficients respectively. Here we apply the forgoing theory to the symmetrised form of the q-quadratic lattice in the elliptic sub-case, which singles out the Askey-Wilson weight and the non-trivial examples of deformations of this weight. We will consider the situation of the deformation variable also on this sub-case of the q-quadratic lattice. The deformation variable is $u = (t + t^{-1})/2$ with $t = e^{i\phi}$ and the analogous relations for the nodes on the deformation lattice are

(7.1)
$$v_{+} + v_{-} = (q^{1/2} + q^{-1/2})u,$$

(7.2)
$$v_{+} - v_{-} = \frac{1}{2}(q^{1/2} - q^{-1/2})(t - t^{-1}),$$

(7.3)
$$\Delta v^2 = (q^{1/2} - q^{-1/2})^2 (u^2 - 1),$$

(7.4)
$$v_{+}v_{-} = u^{2} + \frac{1}{4}(q^{1/2} - q^{-1/2})^{2}.$$

7.1. Moments and Integrals. Our starting point is the simplest extension of the M = 2 case

(7.5)
$$W \pm \Delta y V = z^{\mp 3} \prod_{j=1}^{6} (1 - a_j q^{-1/2} z^{\pm 1}),$$

which is a minimal, natural extension of the Askey-Wilson case given by (5.7). The reader should appreciate that all of the conclusions we are going to draw will follow entirely from (7.5), so that any valid solution of the Pearson equation (4.2) for the weight with this data is as acceptable as any other. The most important consideration in selecting a solution is the support for the weight and existence of the moment data. As

we will see (i.e. in Propositions 7.2 and 7.7) the only place where a specific choice for the weight enters is in determining the initial values of recurrences or divided-difference equations. The four fixed parameters a_1, \ldots, a_4 appear in the same form as they do in the Askey-Wilson weight (M = 2) and so constitutes the base expression, and we seek to introduce an extra parameter and its associated deformation variable. We set $a_5 = \alpha t$, $a_6 = \alpha t^{-1}$ and designate α, t as the deformation parameter and variable respectively.

Remark 7.1. One might entertain the possibility of a weight with the spectral data

(7.6)
$$W \pm \Delta y V = z^{\mp 2} \prod_{j=1}^{4} (1 - q^{-1/2} a_j z^{\pm 1}) (1 - q^{-1/2} t z^{\pm 1}) (1 - q^{\alpha - 1/2} z^{\mp 1}),$$

which is a M = 3 case on a q-quadratic spectral lattice, however it is q-linear in either of the deformation variables t, q^{α} . This is in fact the spectral data for the weight in the integral representation of the very-wellpoised $_{8}W_{7}$ basic hypergeometric function, see Eq. (1.13) of [81]. This function is the classical seed solution for the $E_{7}^{(1)}$ q-Painlevé system as constructed by [52].

Our first undertaking is to derive linear recurrences for the moments which will be required in the verification of key identities from Sections 4 and 6, to make contact with the n = 0 seed solutions of the evolution equations derived later and for checking purposes. Through suitable choices of parameters k, b, a the moment recurrences can be recast as q-difference equations for an integral $I_3(a_1, \ldots, a_6)$ which is a generalisation of the Askey-Wilson integral (5.24).

Proposition 7.1 ([97]). Let σ_k denote the k-th elementary symmetric polynomial in a_1, \ldots, a_4 or a_1, \ldots, a_6 depending on the context. The integral $I_3(a_1, \ldots, a_6)$ satisfies a three-term recurrence in one variable, which we take without loss of generality with respect to a_1 ,

$$(7.7) \quad 0 = \prod_{j \neq 1} (a_1 a_j - 1) I_3(a_1, \ldots) + [1 + q^{-1} - a_1(\sum_{j \neq 1} a_j - q a_1) + \prod_{1}^6 a_j(a_1 \sum_{k \neq 1}^5 a_k^{-1} - q^{-1} - (q+1)a_1^2)] I_3(q a_1, \ldots) + q^{-1}(\prod_{1}^6 a_j - 1) I_3(q^2 a_1, \ldots).$$

In addition the integral $I_3(a_1, \ldots, a_6)$ satisfies a three-term recurrence in two variables, taken to be with respect to a_5, a_6 , which constitutes a pure recurrence in the deformation variable u

$$(7.8) \quad 0 = (a_5 - qa_6) \prod_{j=1}^{4} (1 - a_j a_6) I_3(\dots, q^2 a_5, a_6) - (a_5 - a_6) \left[(1 + q)(1 + qa_5 a_6 \sigma_2 + q^2 a_5^2 a_6^2 \sigma_4) - (qa_5 - a_6)(qa_6 - a_5)(q + \sigma_4) - q(a_5 + a_6)(\sigma_1 + qa_5 a_6 \sigma_3) \right] I_3(\dots, qa_5, qa_6) + (qa_5 - a_6) \prod_{i=1}^{4} (1 - a_j a_5) I_3(\dots, a_5, q^2 a_6).$$

Proof. The first recurrence can be read off from (4.11) specialised to k = 0, after evaluating the expansion coefficients

$$\begin{split} \delta_{3,2}(a) &= \frac{\sigma_6 - 1}{q^{5/2} a^2}, \\ \delta_{3,1}(a) &= \frac{1}{q^{5/2} a^2} [qa(\sigma_5 - \sigma_1) - (1 + q)(1 + qa^2)(\sigma_6 - 1)], \\ \delta_{3,0}(a) &= \frac{1}{q^{3/2} a^2} [(1 + a^2 + a^4)(\sigma_6 - 1) - (a + a^3)(\sigma_5 - \sigma_1) + a^2(\sigma_4 - \sigma_2)]. \end{split}$$

The second recurrence follows from combining the first with the general identity

$$a_k I(qa_j,\ldots,a_k) - a_j I(a_j,\ldots,qa_k) = (a_k - a_j)(1 - a_j a_k) I(a_j,\ldots,a_k),$$

which applies to any integral with products of $\phi_{\infty}(x; a_j)\phi_{\infty}(x; a_k)$ in the denominator of the integrand and any distinct pair $a_j \neq a_k$.

Remark 7.2. A particular solution of (7.8) which serves as a concrete example of a moment sequence is one taken from [97]

$$(7.9) \quad m_{0,0}(t) = t^{1/2} (q\alpha t, \alpha^{-1} t^{-1}, q^{1/2} \alpha t, q^{1/2} \alpha^{-1} t^{-1}; q)_{\infty} \frac{(a_1^{-1} \sigma_4 \alpha t, a_2^{-1} \sigma_4 \alpha t, a_3^{-1} \sigma_4 \alpha t, a_4^{-1} \sigma_4 \alpha t; q)_{\infty}}{(a_1 \alpha t, a_2 \alpha t, a_3 \alpha t, a_4 \alpha t, t^{-2}, \sigma_4 \alpha^2 t^2; q)_{\infty}} \times {}_8 W_7 (q^{-1} \sigma_4 \alpha^2 t^2; q^{-1} \sigma_4 \alpha^2, a_1 \alpha t, a_2 \alpha t, a_3 \alpha t, a_4 \alpha t; q \alpha^{-2}) + (t \mapsto t^{-1}),$$

although this is not a general solution of the q-difference equation, given subsequently as (7.78).

Henceforth we denote $\tilde{\sigma}_j$ as the *j*-th elementary symmetric function of the parameters $q^{-1/2}a_1, \ldots, q^{-1/2}a_6$. If σ_k denotes the *k*-th elementary symmetric function of a_1, a_2, a_3, a_4 then

- (7.10) $\tilde{\sigma}_1 = q^{-1/2} (\sigma_1 + 2\alpha u),$
- (7.11) $\tilde{\sigma}_2 = q^{-1}(\sigma_2 + \alpha^2 + 2\alpha\sigma_1 u),$
- (7.12) $\tilde{\sigma}_3 = q^{-3/2} (\sigma_3 + \alpha^2 \sigma_1 + 2\alpha \sigma_2 u),$

(7.13)
$$\tilde{\sigma}_4 = q^{-2}(\sigma_4 + \alpha^2 \sigma_2 + 2\alpha \sigma_3 u)$$

- (7.14) $\tilde{\sigma}_5 = q^{-5/2} (\alpha^2 \sigma_3 + 2\alpha \sigma_4 u),$
- (7.15) $\tilde{\sigma}_6 = q^{-3} \alpha^2 \sigma_4.$

We will find it advantageous to define two "analogs" of the integers or half-integers s by $[s] := q^s \tilde{\sigma}_6 - q^{-s}$ and $\{s\} := q^s \tilde{\sigma}_6 + q^{-s}$. For convenience we employ the notations w_{\pm} for the evaluations $w_{\pm} \equiv W(\pm 1) = \pm 1 - \tilde{\sigma}_1 \pm \tilde{\sigma}_2 - \tilde{\sigma}_3 \pm \tilde{\sigma}_4 - \tilde{\sigma}_5 \pm \tilde{\sigma}_6$. We also require the coordinates x_j and \tilde{x}_j defined in terms of the parameters by $x_j = \frac{1}{2}(a_j + a_j^{-1})$ and $\tilde{x}_j = \frac{1}{2}(q^{-1/2}a_j + q^{1/2}a_j^{-1})$. Finally let us define the polynomial $w(z) := \prod_{j=1}^4 (1 - q^{-1/2}a_j z)$ which is not to be confused with the weight w(x; u).

Definition 7.1. Note that we have six free parameters at our disposal, $a_1, \ldots, a_4, \alpha \in \mathbb{C}, n \in \mathbb{Z}_{\geq 0}$ and one variable t subject henceforth to the following generic conditions -

(i) $q \neq 1$, (ii) $|q^{-1/2}a_j| \neq 1$ so that $\tilde{x}_j \notin (-1, 1)$ for j = 1, ..., 4, (iii) $|q^{-1/2}\alpha| \neq 1$ so that $q^{-1/2}\alpha + q^{1/2}\alpha^{-1} \notin (-1, 1)$, (iv) $\alpha \neq 0, \pm q^{1/2}$, (v) $t \neq \pm 1$, (vi) $[n], [n + \frac{1}{2}] \neq 0$.

Situations where one or more of the above conditions are violated have to be treated separately, which we refrain from doing here.

As a consequence of (7.5) we have

(7.16)
$$W(x) = 4(1 + \tilde{\sigma}_6)x^3 - 2(\tilde{\sigma}_1 + \tilde{\sigma}_5)x^2 + (\tilde{\sigma}_2 + \tilde{\sigma}_4 - 3 - 3\tilde{\sigma}_6)x + \tilde{\sigma}_1 - \tilde{\sigma}_3 + \tilde{\sigma}_5,$$

(7.17)
$$V(x) = \frac{1}{a^{1/2} - a^{-1/2}} \left[-4(1 - \tilde{\sigma}_6)x^2 + 2(\tilde{\sigma}_1 - \tilde{\sigma}_5)x + 1 - \tilde{\sigma}_2 + \tilde{\sigma}_4 - \tilde{\sigma}_6 \right],$$

The theory detailed in Section 4 allow us to evaluate the third spectral data polynomial.

Proposition 7.2 ([97]). The polynomial U(x) is given by

(7.18)
$$U(x) = \frac{4}{(q^{1/2} - q^{-1/2})} \left[m_{0,0} \left(-2[\frac{1}{2}]x + q^{1/2}\tilde{\sigma}_5 - q^{-1/2}\tilde{\sigma}_1 \right) - [1](m_{0,+} + m_{0,-}) \right],$$

where $m_{0,\pm} = \int \mathbb{D}x w(x;u) z^{\pm}$.

Proof. We start with

(7.19)
$$\Delta y U(x) = (W - \Delta y V) f(y_+) - (W + \Delta y V) f(y_-),$$

and employ the proof in [97]. Having achieved the task of expanding the right-hand side of (7.19) in canonical basis polynomials, all we require are explicit expressions for the coefficients given below

(7.20)
$$\kappa_{3,2}(a) = \frac{1}{2q^{9/2}a^3} [(1+q+q^2)(1+q^2a^2)(1+\sigma_6) - q^2a(\sigma_1+\sigma_5)],$$

(7.21)
$$\kappa_{3,3}(a) = -\frac{1+\sigma_6}{2q^{9/2}a^3}$$

The result is (7.18).

7.2. Spectral Structure. Our next task is to construct a minimal parameterisation of the spectral matrix $A_n(x; u)$ and is the subject of the following proposition. We also recall the large x expansions for the spectral data and coefficients of a more explicit nature which will be of importance

(7.22)
$$W \pm \Delta y V \sim \left[w \pm (q^{1/2} - q^{-1/2}) v \right] x^M + \dots,$$

(7.23)
$$W_n \sim \frac{1}{4} \left[(1+q^n)(1+q^{-n})w + (q^{1/2}-q^{-1/2})(q^n-q^{-n})v \right] x^M + \dots$$

(7.24)
$$\Theta_n \sim \left[\frac{q^{n+1/2} - q^{-n-1/2}}{q^{1/2} - q^{-1/2}}w + (q^{n+1/2} + q^{-n-1/2})v\right]x^{M-2} + \dots,$$

(7.25)
$$\Omega_n + V \sim \frac{1}{2} \left[\frac{q^n - q^{-n}}{q^{1/2} - q^{-1/2}} w + (q^n + q^{-n}) v \right] x^{M-2} + \dots$$

for some constants w, v independent of n, x.

Proposition 7.3. Let one of our free variables, $\lambda_n(u)$, be the zero of $\Theta_n(x; u)$ with respect to x. Also define the other independent variables for $n \ge 0$ by

(7.26)
$$\nu_n(u) \equiv 2W_n(\lambda_n; u) - W(\lambda_n; u), \quad \mu_n(u) \equiv \Omega_n(\lambda_n; u) + V(\lambda_n; u),$$

which are related by the spectral conic equation

(7.27)
$$\nu_n^2 - W^2(\lambda_n) = \Delta(\lambda_n^2 - 1) \left[\mu_n^2 - V^2(\lambda_n) \right],$$

again valid for $n \ge 0$. Let us assume that $\lambda_n(u) \ne \pm 1$ for all n, u. The spectral coefficients for the M = 3 deformed Askey-Wilson OP system are parameterised by λ_n , and either ν_n or μ_n , through the expressions

(7.28)
$$2W_n(x;u) - W(x;u) = \frac{x^2 - 1}{\lambda_n^2 - 1}\nu_n + (x - \lambda_n) \left[4\{n\}(x^2 - 1) + \frac{1}{2}w_+ \frac{x + 1}{1 - \lambda_n} + \frac{1}{2}w_- \frac{x - 1}{1 + \lambda_n}\right],$$

$$(7.29) \quad \Omega_n(x;u) + V(x;u) = \mu_n + 4 \frac{[n]}{q^{1/2} - q^{-1/2}} x(x - \lambda_n) + \frac{16}{q^{1/2} - q^{-1/2}} \frac{\tilde{\sigma}_6}{[n]} \left[\frac{\tilde{\sigma}_1 \tilde{\sigma}_6 + \tilde{\sigma}_5}{4\tilde{\sigma}_6} - \lambda_n \right] (x - \lambda_n) \\ + \frac{1}{q^{1/2} - q^{-1/2}} \frac{\{n\}}{[n]} (x - \lambda_n) \left[\frac{\nu_n}{\lambda_n^2 - 1} + \frac{1}{2} w_+ \frac{1}{1 - \lambda_n} + \frac{1}{2} w_- \frac{1}{1 + \lambda_n} \right],$$

and

(7.30)
$$\Theta_n(x;u) = 8 \frac{[n+\frac{1}{2}]}{q^{1/2} - q^{-1/2}} (x - \lambda_n),$$

for $n \geq 0$.

Proof. Firstly we recall the spectral data polynomials given by (7.5, 7.16,7.17). The product $W^2 - \Delta y^2 V^2$ plays a significant role and therefore we define another set of elementary symmetric polynomials by

(7.31)
$$W^2 - \Delta y^2 V^2 = C_{\infty} \prod_{j=1}^{6} (x - \tilde{x}_j) = C_{\infty} \left[x^6 - e_1 x^5 + e_2 x^4 - e_3 x^3 + e_4 x^2 - e_5 x + e_6 \right],$$

٠,

where $C_{\infty} = 64\tilde{\sigma}_6$. We parameterise the spectral coefficients in the following way

(7.32)
$$2W_n - W = w_3 x^3 + w_2 x^2 + w_1 x + w_0,$$

(7.33)
$$\Theta_n = \varpi_+(x - \lambda_n), \quad \Theta_{n-1} = \varpi_-(x - \lambda_{n-1}),$$

(7.34)
$$\Omega_n + V = v_2 x^2 + v_1 x + v_0,$$

where the leading order coefficients in each are trivial and given by using (7.16) and (7.17) in the expansions (4.28), (4.29) and (4.30)

(7.35)
$$w_3 = 4\{n\}, \quad \varpi_+ = 8 \frac{\left[n + \frac{1}{2}\right]}{q^{1/2} - q^{-1/2}}, \quad v_2 = 4 \frac{\left[n\right]}{q^{1/2} - q^{-1/2}}.$$

From the fundamental bi-linear relation (4.46) we get a system of quadratic polynomial equalities

$$(7.36) w_3^2 - \Delta v_2^2 = C_\infty,$$

$$(7.37) 2w_2w_3 - 2\Delta v_1v_2 = -C_{\infty}e_1,$$

(7.38)
$$2w_1w_3 + w_2^2 - \Delta(v_1^2 - v_2^2 + 2v_0v_2) + a_n^2\Delta\varpi_+\varpi_- = C_\infty e_2,$$

(7.39)
$$2w_1w_2 + 2w_0w_3 - 2\Delta(v_0v_1 - v_1v_2) - a_n^2\Delta\varpi_+\varpi_-(\lambda_n + \lambda_{n-1}) = -C_{\infty}e_3,$$
(7.40)
$$w_1^2 + 2w_1w_2 - \Delta(w_1^2 - w_2^2 - 2w_1w_1) + w_2^2\Delta = -\omega_1(\lambda_n + \lambda_{n-1}) = -C_{\infty}e_3,$$

(7.40)
$$w_1^2 + 2w_0w_2 - \Delta(v_0^2 - v_1^2 - 2v_0v_2) + a_n^2\Delta\varpi_+\varpi_-(\lambda_n\lambda_{n-1} - 1) = C_\infty e_4,$$

(7.41)
$$2w_0w_1 + 2\Delta v_0v_1 + a_n^2 \Delta \varpi_+ \varpi_- (\lambda_n + \lambda_{n-1}) = -C_\infty e_5,$$

(7.42)
$$w_0^2 + \Delta v_0^2 - a_n^2 \Delta \varpi_+ \varpi_- \lambda_n \lambda_{n-1} = C_\infty e_6,$$

where Δ is, again, defined as $(q^{1/2} - q^{-1/2})^2$. Now (7.42,7.40,7.38,7.36) imply $(w_0 + w_2)^2 + (w_1 + w_3)^2 = C_{\infty}(1 + e_2 + e_4 + e_6)$ while (7.41,7.39,7.37) imply $2(w_0 + w_2)(w_1 + w_3) = -C_{\infty}(e_1 + e_3 + e_5)$. Forming the sum and difference of these two later relations we conclude that

(7.43)
$$w_1 + w_3 = \frac{\epsilon W(1) + \epsilon' W(-1)}{2}, \quad w_0 + w_2 = \frac{\epsilon W(1) - \epsilon' W(-1)}{2}$$

where $\epsilon, \epsilon' = \pm 1$ and are yet to be determined. Using (7.43) along with the above definition we can solve for w_0, w_2 , which only leaves the signs ϵ, ϵ' unresolved. These can be fixed by requiring that the n = 0evaluation of $2W_n - W$ precisely reproduces W. This is identically true for all free parameters a_1, \ldots, a_6 provided $\epsilon = +1, \epsilon' = -1$, and yields (7.46,7.44). The above coefficients are given by

(7.44)
$$w_2 = \frac{\nu_n}{\lambda_n^2 - 1} - w_3 \lambda_n - \frac{1}{2} W(1) \frac{1}{\lambda_n - 1} + \frac{1}{2} W(-1) \frac{1}{\lambda_n + 1}$$

(7.45)
$$w_1 = -w_3 + \frac{1}{2}W(1) - \frac{1}{2}W(-1),$$

(7.46)
$$w_0 = -\frac{\nu_n}{\lambda_n^2 - 1} + w_3 \lambda_n + \frac{1}{2} W(1) \frac{\lambda_n}{\lambda_n - 1} + \frac{1}{2} W(-1) \frac{\lambda_n}{\lambda_n + 1}.$$

We observe that this condition also gives $\nu_0 = W(\lambda_0)$, as it must. Through knowledge of w_2 and utilising (7.37) we can determine v_1 , which is given in (7.47).

(7.47)
$$\frac{\Delta v_2}{w_3}v_1 = \frac{\nu_n}{\lambda_n^2 - 1} - w_3\lambda_n + \frac{1}{2}\frac{C_\infty e_1}{w_3} - \frac{1}{2}W(1)\frac{1}{\lambda_n - 1} + \frac{1}{2}W(-1)\frac{1}{\lambda_n + 1},$$

However to find the remaining coefficient v_0 we require μ_n , and this result is given by (7.48).

(7.48)
$$\frac{\Delta v_2}{w_3}v_0 = \frac{\Delta v_2}{w_3}\mu_n - \frac{\lambda_n}{\lambda_n^2 - 1}\nu_n + \frac{C_\infty}{w_3}\lambda_n^2 - \frac{1}{2}\frac{C_\infty e_1}{w_3}\lambda_n + \frac{1}{2}W(1)\frac{\lambda_n}{\lambda_n - 1} - \frac{1}{2}W(-1)\frac{\lambda_n}{\lambda_n + 1}$$

When we examine $\Omega_n + V$ at n = 0 (recall that $\Omega_0 = 0$), we find in addition to previously found relations the equality $\mu_0 = V(\lambda_0)$, again confirming our definition. Therefore we have succeeded in relating the sub-leading coefficients explicitly in terms of two independent variables.

We now address the question of representations for the three-term recurrence coefficients. We shall find that another set of variables, although equivalent to ν_n and μ_n , will lead to the simplest forms for the relations we seek.

(7.49)
$$w_{2,n} := \frac{\nu_n}{\lambda_n^2 - 1} - 4\{n\}\lambda_n - \frac{1}{2}W(1)\frac{1}{\lambda_n - 1} + \frac{1}{2}W(-1)\frac{1}{\lambda_n + 1},$$

(7.50)
$$v_{0,n} := (q^{1/2} - q^{-1/2})\mu_n - 4[n]\lambda_n^2 - \frac{1}{[n]}\lambda_n \left(\{n\}w_{2,n} + 4(\tilde{\sigma}_1\tilde{\sigma}_6 + \tilde{\sigma}_5)\right)$$

To achieve this outcome we require the following recurrence relations.

Proposition 7.4. The dynamical variables $w_{2,n}(u)$ and $v_{0,n}(u)$ satisfy the following system of first order coupled recurrence relations in n

(7.51)
$$w_{2,n+1} - w_{2,n} = -4(q^{1/2} - q^{-1/2})[n + \frac{1}{2}]\lambda_n,$$

(7.52)
$$v_{0,n+1} + v_{0,n} = -\frac{[n+\frac{1}{2}]}{[n+1][n]} \left(2\{n+\frac{1}{2}\}\lambda_n w_{2,n} + 8[n+\frac{1}{2}][n]\lambda_n^2 + 4(q^{1/2}+q^{-1/2})(\tilde{\sigma}_1\tilde{\sigma}_6+\tilde{\sigma}_5)\lambda_n \right),$$

and valid for $n \ge 0$ and subject to the initial conditions $U(\lambda_0; u) = 0$, $w_{2,0} = -2(\tilde{\sigma}_1 + \tilde{\sigma}_5)$ and $v_{0,0} = 1 - \tilde{\sigma}_2 + \tilde{\sigma}_4 - \tilde{\sigma}_6$.

Proof. For the first recurrence we employ our expressions (7.28) and (7.30) in (4.43) and equate the coefficients. The coefficients of x^3 and x are identically satisfied whereas that of either x^2 or x^0 yields (7.51). The second recurrence follows from the examination of the x^0 terms in (4.44) and the employment of the first recurrence. We note the terms in x^2 cancel identically, as they must.

Explicit evaluations of the three term recurrence coefficients can be given in terms of this alternative set of variables.

Proposition 7.5. The three-term recurrence coefficients are found to be given by

$$(7.53) \quad 16[n+\frac{1}{2}][n]^{2}[n-\frac{1}{2}]a_{n}^{2} = \tilde{\sigma}_{6}w_{2,n}^{2} + 2(\tilde{\sigma}_{1}\tilde{\sigma}_{6}+\tilde{\sigma}_{5})\{n\}w_{2,n} + 2[n]^{3}v_{0,n} \\ + 2[n]^{2}\left(4\tilde{\sigma}_{6}+2\tilde{\sigma}_{4}+2\tilde{\sigma}_{2}\tilde{\sigma}_{6}+2\tilde{\sigma}_{1}\tilde{\sigma}_{5}-\{n\}(1+\tilde{\sigma}_{2}+\tilde{\sigma}_{4}+\tilde{\sigma}_{6})+2[n]^{2}\right) + 4(\tilde{\sigma}_{1}\tilde{\sigma}_{6}+\tilde{\sigma}_{5})^{2},$$

for $n \ge 0$ assuming $a_0^2 = 0$, and

(7.54)
$$[n+1][n]b_n = -\frac{1}{4}\{n+\frac{1}{2}\}w_{2,n} - [n+\frac{1}{2}][n]\lambda_n - \frac{1}{2}(q^{1/2}+q^{-1/2})(\tilde{\sigma}_1\tilde{\sigma}_6+\tilde{\sigma}_5)$$

again valid for $n \geq 0$.

Proof. Using (7.38) we can solve for a_n^2 . Observe that the right-hand side of this equation is independent of n whereas individual terms on the left-hand side are. If we assume $a_0^2 = 0$ then the right-hand side is equal to that of the left-hand side evaluated at n = 0, which we know because we can express it simply in terms of the parameters. This means a_n^2 is expressible as a sum of differences and after considerable simplification we arrive at the expression (7.53). To find (7.54) we start with (4.44) and use our previous results for (7.30) and (7.29). Examining the terms in x, we find an expression for b_n in terms of λ_n, ν_n and λ_{n+1}, ν_{n+1} , or equivalently in terms of $\lambda_n, w_{2,n}$ and $\lambda_{n+1}, w_{2,n+1}$. Using (7.51) we can eliminate the $w_{2,n+1}$ term and after some factorisation we arrive at (7.54).

We will not pursue the theory for the $n \mapsto n+1$ recurrences any further here but refer the reader to [96]. Consequently we will drop the n subscript from most variables until Subsection 7.4.

The forgoing parameterisation of our system given in Proposition 7.3 is not useful in the investigations of the u or t-evolution of our system and we require an alternative construction. In conformance with this we define the auxiliary variables l(t) and $\mathfrak{z}_{\pm}(t)$ by

(7.55)
$$\lambda := \frac{1}{2}(l+l^{-1}),$$

(7.56)
$$\mathfrak{z}_{\pm} := \nu \pm \frac{1}{2} (q^{1/2} - q^{-1/2}) [l - l^{-1}] \mu,$$

where, in the first case, the inversion is given by the branch whereby $l \to \infty$ when $\lambda \to \infty$ and an identical choice is made for the second case.

Proposition 7.6. The spectral matrix elements have an alternative parameterisation

(7.57)
$$\mathfrak{T}_{+}(z;t) = 2a_{n}[n+\frac{1}{2}](z-z^{-1})(z-l)(1-l^{-1}z^{-1})$$

and

$$(7.58) \quad [n]\mathfrak{W}_{+}(z;t) = \mathfrak{z}_{+} \frac{(z-z^{-1})(z-l^{-1})(q^{n}\tilde{\sigma}_{6}-q^{-n}lz^{-1})}{(l-l^{-1})^{2}} + \mathfrak{z}_{-} \frac{(z-z^{-1})(z-l)(q^{n}\tilde{\sigma}_{6}-q^{-n}l^{-1}z^{-1})}{(l-l^{-1})^{2}} \\ + [n](z-z^{-1})(z-l)(1-l^{-1}z^{-1})(q^{n}\tilde{\sigma}_{6}z-q^{-n}z^{-1}) \\ + [\tilde{\sigma}_{6}\tilde{\sigma}_{1}+\tilde{\sigma}_{5}-2\tilde{\sigma}_{6}(l+l^{-1})](z-z^{-1})(z-l)(1-l^{-1}z^{-1}) \\ + \frac{1}{2}w_{+}\frac{(z+1)(z-l)(z^{-1}-l)(q^{n}\tilde{\sigma}_{6}-q^{-n}z^{-1})}{(l-1)^{2}} - \frac{1}{2}w_{-}\frac{(z-1)(z-l)(z^{-1}-l)(q^{n}\tilde{\sigma}_{6}+q^{-n}z^{-1})}{(l+1)^{2}},$$

and

$$(7.59) \quad [n]\mathfrak{W}_{-}(z;t) = \mathfrak{z}_{-} \frac{(z-z^{-1})(z-l^{-1})(q^{n}\tilde{\sigma}_{6}lz^{-1}-q^{-n})}{(l-l^{-1})^{2}} + \mathfrak{z}_{+} \frac{(z-z^{-1})(z-l)(q^{n}\tilde{\sigma}_{6}l^{-1}z^{-1}-q^{-n})}{(l-l^{-1})^{2}} \\ - [n](z-z^{-1})(z-l)(1-l^{-1}z^{-1})(q^{n}\tilde{\sigma}_{6}z^{-1}-q^{-n}z) \\ - [\tilde{\sigma}_{6}\tilde{\sigma}_{1}+\tilde{\sigma}_{5}-2\tilde{\sigma}_{6}(l+l^{-1})](z-z^{-1})(z-l)(1-l^{-1}z^{-1}) \\ + \frac{1}{2}w_{+}\frac{(z+1)(z-l)(z^{-1}-l)(q^{n}\tilde{\sigma}_{6}z^{-1}-q^{-n})}{(l-1)^{2}} + \frac{1}{2}w_{-}\frac{(z-1)(z-l)(z^{-1}-l)(q^{n}\tilde{\sigma}_{6}z^{-1}+q^{-n})}{(l+1)^{2}}$$

Proof. These formulae can be viewed as Laurent interpolating polynomials satisfying the following evaluations at the given nodes

(7.60) $\mathfrak{T}_{\pm}(z^{-1};t) = -\mathfrak{T}_{\pm}(z;t),$

(7.61)
$$\mathfrak{T}_+(l^{\pm 1};t) = 0,$$

(7.62)
$$\mathfrak{T}_{+}(\pm 1; t) = 0,$$

(7.63)
$$\mathfrak{T}_+(z;t) \underset{z \to \infty}{\sim} 2a_n [n + \frac{1}{2}] z^2,$$

(7.64)
$$\mathfrak{T}_{+}(z;t) \underset{z \to 0}{\sim} -2a_{n}[n+\frac{1}{2}]z^{-2},$$

and

(7.65)
$$\mathfrak{W}_{\pm}(z^{-1};t) = \mathfrak{W}_{\mp}(z;t),$$

$$\mathfrak{W}_{\pm}(l;t) = \mathfrak{z}_{\pm}$$

$$\mathfrak{W}_{\pm}(\pm 1;t) = w_{\pm}$$

(7.68)
$$\mathfrak{W}_{+}(z;t) \underset{z \to \infty}{\sim} q^{n} \tilde{\sigma}_{6} z^{3}$$

(7.69)
$$\mathfrak{W}_+(z;t) \underset{z \to 0}{\sim} q^{-n} z^{-3}.$$

The above formulae are also a consequence of the parameterisation (7.28-7.29).

Remark 7.3. We note that the eigenvalues of $A_n^*(z;t)$ as $z \to 0, \infty$ are $q^n \tilde{\sigma}_6, q^{-n}$ whilst those at the fixed points of the lattice, of the matrix $A_n^*(\pm 1;t)$, are w_{\pm} . Under the mapping interchanging the interior and exterior of the unit circle in the spectral variable $z \mapsto 1/z$ we observe that $\mathfrak{T}_{\pm} \mapsto -\mathfrak{T}_{\pm}$ and $\mathfrak{W}_{\pm} \mapsto \mathfrak{W}_{\mp}$. As a consequence $A_n^*(z;)A_n^*(z^{-1};) = (W^2 - \Delta y^2 V^2)$ Id, and therefore the mapping corresponds to a reversal of direction on the spectral lattice. With respect to the mapping $l \mapsto l^{-1}$ we note that this must be taken with $\mathfrak{z}_+ \leftrightarrow \mathfrak{z}_-$, and conclude that $\mathfrak{T}_+, \mathfrak{W}_{\pm}$ are invariant under this type of transformation. In addition A_n^* is symmetrical under $t \mapsto t^{-1}$.

7.3. Deformation Structure. The deformation data polynomials R(x; u), S(x; u) are computed as

(7.70)
$$R \pm \Delta v S = (1 - q^{-1/2} \alpha t^{\pm 1} z) (1 - q^{-1/2} \alpha t^{\pm 1} z^{-1}) = \phi_1(x; q^{-1/2} \alpha t^{\pm 1})$$

which implies

(7.71)
$$R = 1 - \alpha^2 q^{-1} - 2\alpha q^{-1/2} x u + 2\alpha^2 q^{-1} u^2,$$

(7.72)
$$S = 2\frac{\alpha}{q-1}(\alpha q^{-1/2}u - x),$$

indicating that L = 1. We observe that the polynomial

(7.73)
$$R^2 - \Delta v^2 S^2 = 4q^{-1} \alpha^2 (x - \tilde{x}_5) (x - \tilde{x}_6),$$

divides

(7.74)
$$W^2 - \Delta y^2 V^2 = 64q^{-3}\sigma_4 \alpha^2 \prod_{j=1}^4 (x - \tilde{x}_j) \cdot (x - \tilde{x}_5)(x - \tilde{x}_6),$$

in conformity with Remark 6.3. The role of the common zeros \tilde{x}_5, \tilde{x}_6 will be crucial in the ensuing investigations. We also note that these points can be represented on a *u*-lattice thus $\tilde{x}_5 = E_u^- x_5$ and $\tilde{x}_6 = E_u^+ x_6$. We can compute the remaining deformation data polynomial T(x; u), which will be of degree zero.

Proposition 7.7 ([97]). The deformation data polynomial T(x; u) has the evaluation

(7.75)
$$T(x;u) = \frac{4\alpha}{q-1} \frac{t m_{0,0}(v_{-}) - t^{-1} m_{0,0}(v_{+})}{t-t^{-1}}$$

Proof. We start from the formula

(7.76)
$$\Delta vT(x;u) = (R - \Delta vS)f(x;v_+) - (R + \Delta vS)f(x;v_-),$$

and employ the expansion (3.46) for some parameter a. Setting a to one of a_1, a_2, a_3, a_4 and utilising the identity

$$\frac{\phi_1(x;b)}{\phi_{n+1}(x;a)} = \frac{(1-abq^n)(1-ba^{-1}q^{-n})}{\phi_{n+1}(x;a)} + \frac{ba^{-1}q^{-n}}{\phi_n(x;a)},$$

we find, by equating coefficients, the result (7.75). Note that $m_{0,0}$ doesn't depend on a. If one chooses a from a_5, a_6 then one requires the additional identity

$$\frac{1}{\phi_n(x;qa)} = \frac{(1-q^n)(a^2-q^{-n})}{\phi_{n+1}(x;a)} + \frac{q^{-n}}{\phi_n(x;a)},$$

in order to merge the two series in (7.76). Alternatively one can compute T from the initial value (6.43) using the asymptotic expression for Φ_0 as given by (6.33).

As a check we can verify that the spectral data polynomials W, V, U and the deformation data polynomials R, S, T satisfy all of the consistency relations formulated in Section 6.

Corollary 7.1. The spectral data polynomials W, V, U and the deformation data polynomials R, S, T satisfy the consistency relations (6.4) and (6.16).

Proof. The first task involving W, V and R, S is elementary, whereas the second requires a moment relation

$$(7.77) \quad [1](m_{0,+} + m_{0,-})(t) = \left[q^{-2}\alpha^2\sigma_3 - q^{-1}\sigma_1 + q^{-3}(q - \alpha^2)(q^2\alpha^{-1}t^{-1} + \sigma_4\alpha t)\right]m_{0,0}(t) \\ - \frac{w(q^{-1/2}\alpha t)}{\alpha t}\frac{q^{-1/2}t\,m_{0,0}(q^{-1}t) - q^{1/2}t^{-1}m_{0,0}(t)}{q^{-1/2}t - q^{1/2}t^{-1}},$$

and the moment q-difference equation in t

$$(7.78) \quad (q^{-1}t - qt^{-1})w(\alpha t^{-1})m_{0,0}(q^{1/2}t) - q^{-1/2}(q^{-1/2}t - q^{1/2}t^{-1}) \left[(1+q)(1+q^{-1}\alpha^2\sigma_2 + q^{-2}\alpha^4\sigma_4) + q^{-1}\alpha^2(q+\sigma_4)(t-t^{-1})(q^{-1}t - qt^{-1}) - \alpha(q^{-1/2}t + q^{1/2}t^{-1})(\sigma_1 + q^{-1}\alpha^2\sigma_3) \right] m_{0,0}(q^{-1/2}t) + (t-t^{-1})w(q^{-1}\alpha t)m_{0,0}(q^{-3/2}t) = 0,$$

which is just (7.8) with $a_5 \mapsto q^{-3/2} \alpha t, a_6 \mapsto q^{-1/2} \alpha t^{-1}$.

We know for L = 1 that the deformation coefficients can be parameterised thus

(7.79)
$$\Re_{\pm} = r_{1\pm}x + r_{0\pm}$$

(7.80)
$$\mathfrak{P}_{\pm} = p_{\pm}.$$

Furthermore we know from the large x expansions (6.32-6.36) what the leading terms are, however in contrast to those of the spectral coefficients these are related to the three-term recurrence coefficients a_n, γ_n in a nontrivial way.

Corollary 7.2. The leading order terms of the elements of the matrix B_n^* are given by

(7.81)
$$r_{1+} = 4\alpha q^{-1/2} H_n \frac{t\gamma_n(v_+)}{\gamma_{n-1}(v_-)}$$

(7.82)
$$r_{1-} = 4\alpha q^{-1/2} H_n \frac{\gamma_n(v_-)}{t\gamma_{n-1}(v_+)}$$

(7.83)
$$p_{+} = 4\alpha q^{-1/2} H_n \left[\frac{t\gamma_n(v_{+})}{\gamma_n(v_{-})} - \frac{\gamma_n(v_{-})}{t\gamma_n(v_{+})} \right],$$

(7.84)
$$p_{-} = 4\alpha q^{-1/2} \frac{a_n H_{n-1}}{a_{n-1}} \left[\frac{t\gamma_{n-1}(v_+)}{\gamma_{n-1}(v_-)} - \frac{\gamma_{n-1}(v_-)}{t\gamma_{n-1}(v_+)} \right]$$

for $n \ge 1$ whilst for n = 0 one can use $\gamma_{-1} = a_0 \gamma_0$.

Up to this point we haven't exercised a choice regarding the decoupling factor H_n but one can take $H_n = \frac{1}{2}a_n$ henceforth.

Remark 7.4. We note that the eigenvalues of $B_n^*(z;t)$ as $z \to 0, \infty$ are $r_{1\pm}$. By construction B_n^* is symmetrical with respect to $z \mapsto z^{-1}$. Under the mapping interchanging the interior and exterior of the unit circle in the deformation variable $t \mapsto 1/t$ we observe (see (6.54), (6.55) and (6.56)) that $\mathfrak{P}_{\pm} \mapsto -\mathfrak{P}_{\pm}$ and $\mathfrak{R}_{\pm} \mapsto \mathfrak{R}_{\mp}$. As a consequence

(7.85)
$$B_n^*(;t)B_n^*(;t^{-1}) = \frac{4H_n^2}{a_n(v_-)a_n(v_+)}(R^2 - \Delta v^2 S^2) \mathrm{Id},$$

and therefore this mapping corresponds to a reversal of direction on the deformation lattice.

As part of an efficient methodology we will require formulae which relate the lower order terms, with respect to x, of the deformation coefficients to the leading order term.

Proposition 7.8. The trailing terms of the deformation coefficients are related to each other and to the spectral parameterisation by

(7.86)
$$p_{+} = r_{1+}a_{n}(v_{-}) - r_{1-}a_{n}(v_{+}),$$

(7.87)
$$p_{-} = r_{1+}a_n(v_{+}) - r_{1-}a_n(v_{-})$$

(7.88)
$$r_{0+} = \frac{r_{1+}}{\Delta v_2} \left[-w_2(v_+) + w_2(v_-) - 2\sqrt{\Delta}(q^{-1/2}\alpha + q^{1/2}\alpha^{-1})(\tilde{\sigma}_6 q^n t - q^{-n}t^{-1}) \right],$$

(7.89)
$$r_{0-} = \frac{r_{1-}}{\Delta v_2} \left[w_2(v_+) - w_2(v_-) - 2\sqrt{\Delta}(q^{-1/2}\alpha + q^{1/2}\alpha^{-1})(\tilde{\sigma}_6 q^n t^{-1} - q^{-n}t) \right],$$

Proof. The formulae given above are ones which involve both shifts up and down on the *u*-lattice of the spectral variables, and these are easily derived by resolving the compatibility relation (6.67) as a Laurent polynomial in z. Such a polynomial is constructed in our application by substituting the elements (7.57-7.59) and (7.79,7.80) into (6.67) and collecting terms. For the 1,2 component we find that the leading order non-zero contributions occur at $z^{\pm 7}$ and the coefficients of both these terms will vanish if (7.86) holds. In the case of the 2,1 component the $z^{\pm 7}$ terms are the leading non-trivial ones and these both vanish when (7.87) holds. The two latter relations (7.88,7.89) follow from the requirement that the coefficients of z^7 terms of the 1,1 and 2, 2 elements vanish respectively. Finally we observe that an independent way of verifying (7.86) and (7.87) is through a trivial combination of the formulae (7.81-7.84)

A vital part of our analysis consists of resolving the compatibility relations (6.70-6.73) with respect to the zeros and poles of $\chi(z; t)$.

Corollary 7.3. The residues of the compatibility relation given in Proposition 6.11 consist of the following equations:

(i) At the advanced co-ordinate $z = z_5 = \alpha t$

(7.90)
$$r_{1-}E_{u}^{-}x_{5} + r_{0-} = -\frac{\mathfrak{W}_{+}(z_{5};q^{1/2}t)}{\mathfrak{T}_{+}(z_{5};q^{1/2}t)}p_{+}$$

(7.91)
$$r_{1+}E_u^-x_5 + r_{0+} = -\frac{\mathfrak{W}_{-}(z_5;q^{1/2}t)}{\mathfrak{T}_{-}(z_5;q^{1/2}t)}p_{-},$$

(ii) at the advanced co-ordinate $z = E_u^{2+} z_6 = q^{-1} \alpha t^{-1}$

(7.92)
$$r_{1-}E_{u}^{+}x_{6} + r_{0-} = \frac{\mathfrak{W}_{-}(E_{u}^{2+}z_{6};q^{1/2}t)}{\mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)}p_{+},$$

(7.93)
$$r_{1+}E_u^+x_6 + r_{0+} = \frac{\mathfrak{W}_+(E_u^{2+}z_6;q^{1/2}t)}{\mathfrak{T}_-(E_u^{2+}z_6;q^{1/2}t)}p_-,$$

(iii) at the retarded co-ordinate $z = E_u^{2-} z_5 = q^{-1} \alpha t$

(7.94)
$$r_{1+}E_u^-x_5 + r_{0+} = -\frac{\mathfrak{W}_{-}(E_u^{-2}z_5;q^{-1/2}t)}{\mathfrak{T}_{+}(E_u^{-2}z_5;q^{-1/2}t)}p_+,$$

(7.95)
$$r_{1-}E_{u}^{-}x_{5} + r_{0-} = -\frac{\mathfrak{W}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)}{\mathfrak{T}_{-}(E_{u}^{2-}z_{5};q^{-1/2}t)}p_{-},$$

(iv) and at the retarded co-ordinate $z = z_6 = \alpha t^{-1}$

(7.96)
$$r_{1+}E_u^+x_6 + r_{0+} = \frac{\mathfrak{W}_+(z_6;q^{-1/2}t)}{\mathfrak{T}_+(z_6;q^{-1/2}t)}p_+,$$

(7.97)
$$r_{1-}E_{u}^{+}x_{6} + r_{0-} = \frac{\mathfrak{W}_{-}(z_{6};q^{-1/2}t)}{\mathfrak{T}_{-}(z_{6};q^{-1/2}t)}p_{-}.$$

All the deformation parameters are evaluated at $t, u, i.e. p_{\pm}(t), r_{1\pm}(t), r_{0\pm}(t)$.

Proof. The multiplicative factor (6.17) is given, in this case, by

(7.98)
$$\chi(z;t) = \frac{(1-q^{-1}\alpha t^{-1}z)(1-\alpha tz^{-1})}{(1-q^{-1}\alpha tz^{-1})(1-\alpha t^{-1}z)}.$$

At the zeros of χ the right-hand sides of (6.70,6.71,6.72,6.73) must vanish whereas at the poles the lefthand factors must vanish. Thus for the zero $z = z_5 = \alpha t$ applied to (6.70) we deduce (7.90). For the other zero at $z = 1/E_u^{2+}z_6 = q\alpha^{-1}t$ applied to (6.70) we make additional use of the symmetries of A_n^* and B_n^* with respect to inversion of z to derive (7.92). At the pole $z = E_u^{2-}z_5 = q^{-1}\alpha t$ the residue of (6.70) yields (7.94) whereas at the other pole $z = 1/z_6 = \alpha^{-1}t$ we have to apply the additional symmetries to the residue to arrive at (7.96). However we have further relations, which are equivalent to the four relations derived above, by specialising the spectral variable in these ways due to the fact that $(W^2 - \Delta y^2 V^2)(z = \alpha t, q^{-1}\alpha t^{-1}; q^{1/2}t) = 0$, $(W^2 - \Delta y^2 V^2)(z = q^{-1}\alpha t, \alpha t^{-1}; q^{-1/2}t) = 0$, $(R^2 - \Delta v^2 S^2)(z = q^{-1/2}\alpha t; t) = 0$ and $(R^2 - \Delta v^2 S^2)(z = q^{-1/2}\alpha t^{-1}; t) = 0$, which implies

(7.99)
$$(\mathfrak{W}_+\mathfrak{W}_- + \mathfrak{T}_+\mathfrak{T}_-)(z = \alpha t, q^{-1}\alpha t^{-1}; q^{1/2}t) = 0,$$

(7.100)
$$(\mathfrak{W}_{+}\mathfrak{W}_{-} + \mathfrak{T}_{+}\mathfrak{T}_{-})(z = q^{-1}\alpha t, \alpha t^{-1}; q^{-1/2}t) = 0,$$

(7.101)
$$(\mathfrak{R}_+\mathfrak{R}_- + \mathfrak{P}_+\mathfrak{P}_-)(z = q^{-1/2}\alpha t; t) = 0,$$

(7.102)
$$(\mathfrak{R}_{+}\mathfrak{R}_{-} + \mathfrak{P}_{+}\mathfrak{P}_{-})(z = q^{-1/2}\alpha t^{-1}; t) = 0.$$

This means that each of the identities (7.90, 7.92, 7.94, 7.96) can take four forms and the remaining relations (7.91, 7.93, 7.95, 7.97) are examples of just one of those forms. This equivalence also ensures that if the residue condition is satisfied by (6.70) then it is automatically satisfied by (6.71, 6.72, 6.73) as well.

Corollary 7.4. The following product relation holds amongst the spectral coefficients evaluated at the zeros of the spectral determinant

(7.103)
$$\frac{\mathfrak{W}_{+}(z_{5};q^{1/2}t)}{\mathfrak{T}_{+}(z_{5};q^{1/2}t)}\frac{\mathfrak{W}_{-}(E_{u}^{2-}z_{5};q^{-1/2}t)}{\mathfrak{T}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)} = \frac{\mathfrak{W}_{+}(z_{6};q^{-1/2}t)}{\mathfrak{T}_{+}(z_{6};q^{-1/2}t)}\frac{\mathfrak{W}_{-}(E_{u}^{2+}z_{6};q^{1/2}t)}{\mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)}.$$

Proof. By comparing the right-hand sides of the pairs, (7.90, 7.95), (7.92, 7.97), (7.94, 7.91) and (7.96, 7.93), we see that

(7.104)
$$\frac{\mathfrak{W}_{+}(z_{5};q^{1/2}t)}{\mathfrak{T}_{+}(z_{5};q^{1/2}t)}p_{+} = \frac{\mathfrak{W}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)}{\mathfrak{T}_{-}(E_{u}^{2-}z_{5};q^{-1/2}t)}p_{-},$$

(7.105)
$$\frac{\mathfrak{W}_{-}(E_{u}^{2+}z_{6};q^{1/2}t)}{\mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)}p_{+} = \frac{\mathfrak{W}_{-}(z_{6};q^{-1/2}t)}{\mathfrak{T}_{-}(z_{6};q^{-1/2}t)}p_{-},$$

(7.106)
$$\frac{\mathfrak{W}_{-}(E_{u}^{2-}z_{5};q^{-1/2}t)}{\mathfrak{T}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)}p_{+} = \frac{\mathfrak{W}_{-}(z_{5};q^{1/2}t)}{\mathfrak{T}_{-}(z_{5};q^{1/2}t)}p_{-},$$

(7.107)
$$\frac{\mathfrak{W}_{+}(z_{6};q^{-1/2}t)}{\mathfrak{T}_{+}(z_{6};q^{-1/2}t)}p_{+} = \frac{\mathfrak{W}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)}{\mathfrak{T}_{-}(E_{u}^{2+}z_{6};q^{1/2}t)}p_{-}$$

By forming the cross products of the first and second, or equivalently the third and fourth we can establish (7.103), assuming $p_+p_- \neq 0$. Taking cross products of the first and third, or second and fourth leads to the identities that are trivial consequences of the determinantal relation.

Definition 7.2. In conformity with the remark made immediately preceeding Corollary 6.4 let us define the variables with deformation arguments evaluated at the advanced and retarded co-ordinates through the notation $\hat{l} = l(q^{1/2}t)$, $\check{l} = l(q^{-1/2}t)$ etc.

Lemma 7.1. One solution for the component p_+ is given by

$$(7.108) \quad q^{1/2}\alpha^{-1}(t^{-1}-t)[n+\frac{1}{2}]\frac{r_{1-}\hat{a}_{n}}{p_{+}} = -\frac{qt\hat{l}}{(\hat{l}-\hat{l}^{-1})^{2}} \left[\frac{\hat{\mathfrak{z}}_{+}}{(\alpha t-\hat{l})(qt-\alpha\hat{l})} + \frac{\hat{\mathfrak{z}}_{-}}{(-\alpha+qt\hat{l})(-1+\alpha t\hat{l})}\right] \\ -\frac{1}{\alpha t}(q^{-n}+q^{n+1}t^{2}\sigma_{6}) + \frac{1}{4}w(1)\frac{1}{\hat{\lambda}-1} + \frac{1}{4}w(-1)\frac{1}{\hat{\lambda}+1}$$

A second solution for the component p_+ is

$$(7.109) \quad q^{1/2}\alpha^{-1}(t^{-1}-t)[n+\frac{1}{2}]\frac{r_{1+\check{a}_n}}{p_+} = \frac{qt\check{l}}{(\check{l}-\check{l}^{-1})^2} \left[\frac{\check{\mathfrak{z}}_+}{(q-\alpha t\check{l})(-\alpha+t\check{l})} + \frac{\check{\mathfrak{z}}_-}{(t-\alpha\check{l})(-\alpha t+q\check{l})}\right] \\ - \frac{1}{\alpha t}(q^{-n}t^2 + q^{n+1}\sigma_6) + \frac{1}{4}w(1)\frac{1}{\check{\lambda}-1} + \frac{1}{4}w(-1)\frac{1}{\check{\lambda}+1}.$$

Proof. The first relation (7.108) follows from the subtraction of (7.92) from (7.90) and simplifying. The second relation (7.109) follows from similar reasoning applied to (7.94) and (7.96). \Box

Remark 7.5. A profound difference between the expressions (7.108,7.109) and the corresponding formulae in the case of the *q*-linear spectral lattice is the presence of both $\hat{\mathfrak{z}}_{\pm}$ (or $\check{\mathfrak{z}}_{\pm}$) whereas in the latter case only one these variables in present, which significantly complicates the ensuing analysis.

Definition 7.3. We define the following sequence of co-ordinate transformations

(7.110)
$$\mathfrak{z}_{+} = l^{-2} w(l) (1 - q^{-1/2} \alpha t l^{-1}) (1 - q^{-1/2} \alpha t^{-1} l) \frac{lf - 1}{f - l},$$

(7.111)
$$\mathfrak{z}_{-} = l^2 w(l^{-1})(1 - q^{-1/2}\alpha t l)(1 - q^{-1/2}\alpha t^{-1} l^{-1}) \frac{f - l}{lf - 1},$$

where the new variable f(t) is introduced. Clearly it is apparent that under the mapping $l \mapsto l^{-1}$ we have the interchange $\mathfrak{z}_+ \leftrightarrow \mathfrak{z}_-$ and that the product satisfies the relation (7.27), i.e. $\mathfrak{z}_+\mathfrak{z}_- = \prod_{j=1}^6 (1 - q^{-1/2}a_j l^{\pm})$.

Let $\sigma_4 = q^2 s_4^2$. In addition we define the further variable $\rho(t)$ so that

(7.112)
$$\frac{r_{1+\check{a}_n}}{p_+} := \frac{1}{q^{1/2}[n+\frac{1}{2}]} \frac{2\alpha s_4 \hat{\rho} - q^{-n}t - q^n \alpha^2 s_4^2 t^{-1}}{t - t^{-1}}$$

(7.113)
$$\frac{r_{1-}\hat{a}_n}{p_+} := \frac{1}{q^{1/2}[n+\frac{1}{2}]} \frac{2\alpha s_4\hat{\rho} - q^n\alpha^2 s_4^2 t - q^{-n}t^{-1}}{t - t^{-1}}$$

Together these relations ensure that (7.86) is automatically satisfied. We further define g(t) by $2\rho := g + g^{-1}$. We shall find that f, l, g will be our primary variables.

A consequence of these definitions and the previous Lemma is the following transformation formula.

Corollary 7.5. Let us assume $f \neq 0, \infty$, $f \neq l^{\pm 1}$ and $w(f) \neq 0$. There exists an invertible mapping between $\rho(t)$ and $\lambda(t)$ with the form

(7.114)
$$-2s_4f\rho + 1 + s_4^2f^2 = \frac{w(f)}{f^2 + 1 - 2f\lambda}$$

Proof. Employing the definitions (7.113), (7.110) and (7.111) in (7.108) we compute the relation

(7.115)
$$2s_4\rho = q^{-3/2}\sigma_3 - s_4^2[l+l^{-1}] + \frac{w(l)}{(l^2-1)(l-f)} + \frac{l^3w(l^{-1})}{(l^2-1)(lf-1)},$$

where we have shifted the arguments $t \mapsto q^{-1/2}t$ of all variables. The relation (7.114) then follows by performing a partial fraction expansion of the above equation with respect to the variable l(t) (the former equation can be thought of as an expansion with respect to f(t)). We also have an alternative form of the above expression

(7.116)
$$\lambda = q^{1/2} \frac{\sigma_3}{2\sigma_4} - \frac{1}{2s_4} (g + g^{-1}) - \frac{1}{2(g^2 - 1)} \frac{w(s_4^{-1}g)}{f - s_4^{-1}g} + \frac{g^2}{2(g^2 - 1)} \frac{w(s_4^{-1}g^{-1})}{f - s_4^{-1}g^{-1}}.$$

Remark 7.6. What is surprising about this result in comparison to the situation with the q-linear lattice is that instead of a linear transformation between ρ and f we find one between ρ and λ with f being an intermediary. Given (7.115) it is clear there is no linear inversion possible for f.

Our next task is to compute expressions for the four independent evaluations of the spectral matrix elements given on the left-hand sides of (7.104-7.107) in terms of the variables introduced above. We will carry this out in stages of successive refinement.

Lemma 7.2. The ratios of the evaluated spectral matrix elements satisfy the relations (7.117)

$$\frac{\hat{a}_n}{\check{a}_n} \frac{2\alpha s_4 \hat{\rho} - q^{-n}t - q^n \alpha^2 s_4^2 t^{-1}}{2\alpha s_4 \hat{\rho} - q^n \alpha^2 s_4^2 t - q^{-n}t^{-1}} = \frac{\mathfrak{W}_+(z_6; q^{-1/2}t)}{\mathfrak{T}_+(z_6; q^{-1/2}t)} \frac{\mathfrak{T}_+(z_5; q^{1/2}t)}{\mathfrak{W}_+(z_5; q^{1/2}t)} = \frac{\mathfrak{W}_-(E_u^{2-} z_5; q^{-1/2}t)}{\mathfrak{T}_+(E_u^{2-} z_5; q^{-1/2}t)} \frac{\mathfrak{T}_+(E_u^{2+} z_6; q^{1/2}t)}{\mathfrak{W}_-(E_u^{2+} z_6; q^{1/2}t)}$$

Proof. Subtracting (7.90) from (7.92) gives

(7.118)
$$r_{1-}\left(E_{u}^{+}x_{6}-E_{u}^{-}x_{5}\right) = \left(\frac{\mathfrak{W}_{-}(E_{u}^{2+}z_{6};q^{1/2}t)}{\mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)} + \frac{\mathfrak{W}_{+}(z_{5};q^{1/2}t)}{\mathfrak{T}_{+}(z_{5};q^{1/2}t)}\right)p_{+}$$

whilst subtracting (7.94) from (7.96) gives

(7.119)
$$r_{1+}\left(E_u^+ x_6 - E_u^- x_5\right) = \left(\frac{\mathfrak{W}_+(z_6; q^{-1/2}t)}{\mathfrak{T}_+(z_6; q^{-1/2}t)} + \frac{\mathfrak{W}_-(E_u^{2-}z_5; q^{-1/2}t)}{\mathfrak{T}_+(E_u^{2-}z_5; q^{-1/2}t)}\right)p_+.$$

Taking ratios of these two gives

$$(7.120) \quad \frac{r_{1+}}{r_{1-}} = \frac{\mathfrak{T}_{+}(z_{5};q^{1/2}t)\mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)}{\mathfrak{T}_{+}(z_{6};q^{-1/2}t)\mathfrak{T}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)} \\ \times \frac{\mathfrak{T}_{+}(z_{6};q^{-1/2}t)\mathfrak{W}_{-}(E_{u}^{2-}z_{5};q^{-1/2}t) + \mathfrak{T}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)\mathfrak{W}_{+}(z_{6};q^{-1/2}t)}{\mathfrak{T}_{+}(z_{5};q^{1/2}t)\mathfrak{W}_{-}(E_{u}^{2+}z_{6};q^{1/2}t) + \mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)\mathfrak{W}_{+}(z_{5};q^{1/2}t)}$$

Likewise subtracting (7.91) from (7.93) yields

(7.121)
$$r_{1+} \left(E_u^+ x_6 - E_u^- x_5 \right) = - \left(\frac{\mathfrak{T}_+ (E_u^{2+} z_6; q^{1/2} t)}{\mathfrak{W}_- (E_u^{2+} z_6; q^{1/2} t)} + \frac{\mathfrak{T}_+ (z_5; q^{1/2} t)}{\mathfrak{W}_+ (z_5; q^{1/2} t)} \right) p_-,$$

and (7.95) from (7.97)

(7.122)
$$r_{1-}\left(E_{u}^{+}x_{6}-E_{u}^{-}x_{5}\right)=-\left(\frac{\mathfrak{T}_{+}(z_{6};q^{-1/2}t)}{\mathfrak{W}_{+}(z_{6};q^{-1/2}t)}+\frac{\mathfrak{T}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)}{\mathfrak{W}_{-}(E_{u}^{2-}z_{5};q^{-1/2}t)}\right)p_{-}.$$

Taking the ratio of these latter two equations gives

$$(7.123) \quad \frac{r_{1+}}{r_{1-}} = \frac{\mathfrak{W}_{+}(z_{6};q^{-1/2}t)\mathfrak{W}_{-}(E_{u}^{2-}z_{5};q^{-1/2}t)}{\mathfrak{W}_{+}(z_{5};q^{1/2}t)\mathfrak{W}_{-}(E_{u}^{2+}z_{6};q^{1/2}t)} \times \frac{\mathfrak{T}_{+}(z_{5};q^{1/2}t)\mathfrak{W}_{-}(E_{u}^{2+}z_{6};q^{1/2}t) + \mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)\mathfrak{W}_{+}(z_{5};q^{1/2}t)}{\mathfrak{T}_{+}(z_{6};q^{-1/2}t)\mathfrak{W}_{-}(E_{u}^{2-}z_{5};q^{-1/2}t) + \mathfrak{T}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)\mathfrak{W}_{+}(z_{6};q^{-1/2}t)}.$$

The factors appearing on the right-hand sides of these two ratios constructed have simple evaluations, the first factor being (7.124)

$$\frac{\hat{\mathfrak{T}}_{+}(z_{5};q^{1/2}t)\mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)}{\mathfrak{T}_{+}(z_{6};q^{-1/2}t)\mathfrak{T}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)} = \frac{\hat{a}_{n}^{2}}{\tilde{a}_{n}^{2}}\frac{(q^{2}t^{2}-\alpha^{2})(\alpha^{2}t^{2}-1)(\alpha t-\hat{l})(\alpha t-\hat{l}^{-1})(qt-\alpha\hat{l}^{-1})(qt-\alpha\hat{l})}{(t^{2}-\alpha^{2})(\alpha^{2}t^{2}-q^{2})(\alpha t-q\check{l})(\alpha t-q\check{l}^{-1})(t-\alpha\check{l}^{-1})(t-\alpha\check{l})}.$$

Employing this evaluation in the first ratio (7.120) yields the evaluation for the second factor

$$(7.125) \quad \frac{\mathfrak{T}_{+}(z_{6};q^{-1/2}t)\mathfrak{W}_{-}(E_{u}^{2-}z_{5};q^{-1/2}t) + \mathfrak{T}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)\mathfrak{W}_{+}(z_{6};q^{-1/2}t)}{\mathfrak{T}_{+}(z_{5};q^{1/2}t)\mathfrak{W}_{-}(E_{u}^{2+}z_{6};q^{1/2}t) + \mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)\mathfrak{W}_{+}(z_{5};q^{1/2}t)} = \frac{\check{a}_{n}}{\hat{a}_{n}}\frac{(t^{2}-\alpha^{2})(\alpha^{2}t^{2}-q^{2})}{(q^{2}t^{2}-\alpha^{2})(\alpha^{2}t^{2}-1)}\frac{(\alpha t-q\check{l})(\alpha t-q\check{l}^{-1})(t-\alpha\check{l}^{-1})(t-\alpha\check{l})}{(\alpha t-\hat{l})(\alpha t-\hat{l}^{-1})(qt-\alpha\hat{l})}\frac{2\alpha s_{4}\hat{\rho}-q^{-n}t-q^{n}\alpha^{2}s_{4}^{2}t^{-1}}{2\alpha s_{4}\hat{\rho}-q^{n}\alpha^{2}s_{4}^{2}t-q^{-n}t^{-1}}.$$

Employing this latter factor in the second ratio (7.123) we can construct two relations involving perfect squares, upon using (7.103). After taking the square roots we need to resolve the sign ambiguity. The final result is then (7.117).

Remark 7.7. In fact we can make separate evaluations of the numerator and denominator of the last ratio, which gives us

$$\begin{aligned} (7.126) \quad \mathfrak{T}_{+}(z_{5};q^{1/2}t)\mathfrak{W}_{-}(E_{u}^{2+}z_{6};q^{1/2}t) + \mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)\mathfrak{W}_{+}(z_{5};q^{1/2}t) \\ &= \frac{2\hat{a}_{n}[n+\frac{1}{2}]}{q^{3}\alpha^{5}t^{4}}(q^{2}t^{2}-\alpha^{2})(\alpha^{2}t^{2}-1)(\alpha^{2}-q)(\alpha t-\hat{l})(\alpha t-\hat{l}^{-1})(qt-\alpha\hat{l})(qt-\alpha\hat{l}^{-1})\left[2\alpha s_{4}\hat{\rho}-q^{n}\alpha^{2}s_{4}^{2}t-q^{-n}t^{-1}\right], \end{aligned}$$

and

$$\begin{aligned} (7.127) \quad & \mathfrak{T}_{+}(z_{6};q^{-1/2}t)\mathfrak{W}_{-}(E_{u}^{2-}z_{5};q^{-1/2}t) + \mathfrak{T}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)\mathfrak{W}_{+}(z_{6};q^{-1/2}t) \\ &= \frac{2\check{a}_{n}[n+\frac{1}{2}]}{q^{3}\alpha^{5}t^{4}}(t^{2}-\alpha^{2})(\alpha^{2}t^{2}-q^{2})(\alpha^{2}-q)(\alpha t-q\check{l})(\alpha t-q\check{l}^{-1})(t-\alpha\check{l})(t-\alpha\check{l}^{-1})\left[2\alpha s_{4}\hat{\rho}-q^{-n}t-q^{n}\alpha^{2}s_{4}^{2}t^{-1}\right]. \end{aligned}$$

In order to proceed any further we will require representations of the individual evaluated spectral coefficients which we give as two distinct partial-fraction expansions. We offer these without proofs as they are the outcome of straightforward although lengthy computations. Lemma 7.3. The evaluated spectral coefficients possess the rational function forms in the f, l variables

$$(7.128) \quad 2t[n][n+\frac{1}{2}]\hat{a}_n \frac{\mathfrak{W}_+(z_5;q^{1/2}t)}{\mathfrak{T}_+(z_5;q^{1/2}t)} \\ = -q^{-1}\hat{f}^{-2}(qt-\alpha\hat{f})(q^n\alpha t\tilde{\sigma}_6 - q^{-n}\hat{f})\frac{w(\hat{f})}{(\hat{f}-\hat{l})(\hat{f}-\hat{l}^{-1})} + \alpha^{-1}\tilde{\sigma}_6\hat{f}^{-1}(\alpha t - q^{-n}\hat{f})(q^n\alpha t - \hat{f})(\hat{l}+\hat{l}^{-1}) \\ + \alpha^{-1}\hat{f}^{-2}(q^n\alpha t - \hat{f})[-q^{-n}\tilde{\sigma}_6(\hat{f}^2 - 1)(q^n\alpha t + \hat{f}) - q^{-1/2}\alpha t\tilde{\sigma}_6\sigma_1\hat{f} \\ + q^{-n-5/2}\alpha^2\sigma_3\hat{f}^2 + (q^{-n+2} + q^{n-2}\alpha^4\sigma_4)\hat{f}(-q^{-n-2} + q^{-4}\alpha t\sigma_4\hat{f})],$$

$$(7.129) \quad 2t[n][n+\frac{1}{2}]\check{a}_{n}\frac{\mathfrak{W}_{+}(z_{6};q^{-1/2}t)}{\mathfrak{T}_{+}(z_{6};q^{-1/2}t)} \\ = -q^{-1}\check{f}^{-2}\frac{(t-\alpha\check{f})(\alpha t-q\check{f})(q^{n}\alpha\check{\sigma}_{6}-q^{-n}t\check{f})}{(\alpha-t\check{f})}\frac{w(\check{f})}{(\check{f}-\check{l})(\check{f}-\check{l}^{-1})} \\ + \alpha^{-1}\check{\sigma}_{6}\check{f}^{-1}(\alpha t-q^{-n+1}\check{f})(q^{n-1}\alpha t-\check{f})(\check{l}+\check{l}^{-1}) \\ - \frac{[n](q-t^{2})t^{3}}{q}\frac{(t-\alpha\check{f})}{(\alpha-t\check{f})}\frac{w(\alpha t^{-1})}{(\alpha-t\check{l})(\alpha-t\check{l}^{-1})} \\ + \alpha^{-1}t^{-1}\check{f}^{-2}(q^{n-1}\alpha t-\check{f})[-q^{-n}t\check{\sigma}_{6}(\check{f}^{2}-1)(q^{n}\alpha t+q\check{f})-q^{-n}(q-t^{2})\check{\sigma}_{6}\check{f}(q^{n}t+\alpha\check{f}) \\ + q^{-2-2n}t^{2}\check{f}(-q^{2}t+q^{n}\alpha\sigma_{4}\check{f})-q^{-1/2}\alpha t^{2}\check{\sigma}_{6}\sigma_{1}\check{f}+q^{-\frac{3}{2}-n}\alpha^{2}t\sigma_{3}\check{f}^{2}-\alpha^{2}\check{\sigma}_{6}\check{f}(t-q^{n-2}\alpha\sigma_{4}\check{f})], \end{cases}$$

$$(7.130) \quad 2t[n][n+\frac{1}{2}]\hat{a}_{n}\frac{\mathfrak{W}_{-}(E_{u}^{2+}z_{6};q^{1/2}t)}{\mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)} \\ = q^{-1/2}\hat{f}^{-2}(\alpha t - \hat{f})(q^{n+1/2}t\tilde{\sigma}_{6} - q^{-n-1/2}\alpha\hat{f})\frac{w(\hat{f})}{(\hat{f}-\hat{t})(\hat{f}-\hat{t}^{-1})} - \alpha^{-1}\tilde{\sigma}_{6}\hat{f}^{-1}(\alpha t - q^{-n}\hat{f})(q^{n}\alpha t - \hat{f})(\hat{t}+\hat{t}^{-1}) \\ + \hat{f}^{-2}(q^{n}\alpha t - \hat{f})[q^{-3-n}(q^{n}-1)\alpha\sigma_{4}\hat{f}(1+\alpha t\hat{f}) + q^{-1/2}t\tilde{\sigma}_{6}\sigma_{1}\hat{f} - q^{-n-5/2}\alpha\sigma_{3}\hat{f}^{2} \\ - (q^{n+1/2}t\tilde{\sigma}_{6} - q^{-n-1/2}\alpha\hat{f})(q^{-n-1/2} + q^{-5/2}\sigma_{4}\hat{f}^{2})], \end{cases}$$

$$(7.131) \quad 2t[n][n+\frac{1}{2}]\check{a}_{n}\frac{\mathfrak{W}_{-}(E_{u}^{2-}z_{5};q^{-1/2}t)}{\mathfrak{T}_{+}(E_{u}^{2-}z_{5};q^{-1/2}t)} \\ = q^{-1/2}\check{f}^{-2}\frac{(t-\alpha\check{f})(\alpha t-q\check{f})(q^{n+1/2}\check{\sigma}_{6}-q^{-n-1/2}\alpha t\check{f})}{(q-\alpha t\check{f})}\frac{w(\check{f})}{(\check{f}-\check{t})(\check{f}-\check{t}^{-1})} \\ -\alpha^{-1}\check{\sigma}_{6}\check{f}^{-1}(\alpha t-q^{-n+1}\check{f})(q^{n-1}\alpha t-\check{f})(\check{t}+\check{t}^{-1}) \\ +\frac{[n](q-t^{2})\alpha^{4}t^{3}}{q^{2}}\frac{(\alpha t-q\check{f})}{(q-\alpha t\check{f})}\frac{w(q\alpha^{-1}t^{-1})}{(q-\alpha t\check{t})(q-\alpha t\check{t}^{-1})} \\ +t^{-1}\check{f}^{-2}(q^{n-1}\alpha t-\check{f})[q^{1-n}\alpha^{-1}t\check{\sigma}_{6}(\check{f}^{2}-1)(q^{n-1}\alpha t+\check{f})+q^{-4-n}(q-t^{2})\sigma_{4}\check{f}(q^{n}\alpha^{3}t+q^{2}\check{f}) \\ +q^{-1/2}t^{2}\check{\sigma}_{6}\sigma_{1}\check{f}-q^{-n-3/2}\alpha t\sigma_{3}\check{f}^{2}-q^{-4-2n}\alpha(qt^{2}+q^{2n}\sigma_{4})\check{f}(-q^{2}t+q^{n}\alpha\sigma_{4}\check{f})].$$

In addition the evaluations at the advanced co-ordinate have alternative forms.

Lemma 7.4. The advanced evaluated spectral coefficients are rational functions in the g, f variables

$$(7.132) \quad 2t[n][n+\frac{1}{2}]\hat{a}_{n}\frac{\mathfrak{W}_{+}(z_{5};q^{1/2}t)}{\mathfrak{T}_{+}(z_{5};q^{1/2}t)} \\ = -q^{-n-1}\alpha s_{4}\frac{\left(q^{n}t\alpha s_{4}\hat{g}^{-1}-1\right)^{2}}{\hat{g}-\hat{g}^{-1}}\frac{w(s_{4}^{-1}\hat{g})}{\hat{f}-s_{4}^{-1}\hat{g}} + q^{-n-1}\alpha s_{4}\frac{\left(q^{n}t\alpha s_{4}\hat{g}-1\right)^{2}}{\hat{g}-\hat{g}^{-1}}\frac{w(s_{4}^{-1}\hat{g}^{-1})}{\hat{f}-s_{4}^{-1}\hat{g}^{-1}} \\ -q^{-n-1}\alpha\left(\hat{g}^{2}+\hat{g}^{-2}\right) - q^{-n-6}s_{4}^{-1}\left[-q^{7/2}\alpha\sigma_{3}+t\sigma_{4}(q^{4}-2q^{n+3}\alpha^{2}+q^{2n}\alpha^{4}\sigma_{4})\right]\left(\hat{g}+\hat{g}^{-1}\right) \\ +q^{-2n-6}\alpha^{-1}\left[q^{6}+t^{2}q^{4n}\alpha^{6}\sigma_{4}^{2}-t^{2}q^{3n+1}\alpha^{4}\sigma_{4}(q^{2}+\sigma_{4})+q^{2n+2}\alpha^{2}(t^{2}q^{2}+\alpha^{2})\sigma_{4} \\ -q^{n+4}\alpha^{2}(2q+\sigma_{2})+q^{2n+5/2}\alpha^{3}t(-q\sigma_{3}+\sigma_{1}\sigma_{4})\right],$$

and

$$(7.133) \quad 2t[n][n+\frac{1}{2}]\hat{a}_{n}\frac{\mathfrak{W}_{-}(E_{u}^{2+}z_{6};q^{1/2}t)}{\mathfrak{T}_{+}(E_{u}^{2+}z_{6};q^{1/2}t)} \\ = q^{-n-1}\alpha s_{4}\frac{\left(q^{n}t\alpha s_{4}\hat{g}^{-1}-1\right)^{2}}{\hat{g}-\hat{g}^{-1}}\frac{w(s_{4}^{-1}\hat{g})}{\hat{f}-s_{4}^{-1}\hat{g}} - q^{-n-1}\alpha s_{4}\frac{\left(q^{n}t\alpha s_{4}\hat{g}-1\right)^{2}}{\hat{g}-\hat{g}^{-1}}\frac{w(s_{4}^{-1}\hat{g}^{-1})}{\hat{f}-s_{4}^{-1}\hat{g}^{-1}} \\ + q^{-n-1}\alpha\left(\hat{g}^{2}+\hat{g}^{-2}\right) - q^{-n-15/2}\alpha s_{4}^{-1}\left[q^{5}\sigma_{3}-t\alpha\sigma_{4}(q^{9/2}-2q^{n+9/2}+q^{2n+5/2}\sigma_{4})\right]\left(\hat{g}+\hat{g}^{-1}\right) \\ + q^{-2n-11/2}\alpha\left[-q^{9/2}-q^{4n+1/2}\alpha^{2}t^{2}\sigma_{4}^{2}+q^{3n+1/2}\alpha^{2}t^{2}\sigma_{4}(q^{2}+\sigma_{4})-q^{2n+5/2}(1+\alpha^{2}t^{2})\sigma_{4} \\ + q^{n+7/2}(2q+\sigma_{2})-q^{2n+2}\alpha t(-q\sigma_{3}+\sigma_{1}\sigma_{4})\right].$$

At this stage we have accumulated enough results to deduce the dynamical equations for our system on the deformation lattice.

Proposition 7.9. The t-evolution of our system in the variables ρ , λ , f is given by a pair of coupled first order equations the first of which is

$$(7.134) \quad \frac{2qs_4\hat{\rho} - t^2\check{f}^{-1} - q^2t^{-2}s_4^2\check{f}}{2s_4\check{\rho} - \check{f}^{-1} - s_4^2\check{f}} = \frac{q(t - \alpha\check{f})(q\check{f} - \alpha t)}{(t\check{f} - \alpha)(q - \alpha t\check{f})} \\ + \frac{\alpha^4t^2(q - t^2)}{(q - \alpha^2)}\frac{(q\check{f} - \alpha t)}{(q - \alpha t\check{f})}\frac{\check{f}^2w(q\alpha^{-1}t^{-1})}{q\alpha tw(\check{f}) - (q\check{f} - \alpha t)(q - \alpha t\check{f})\check{f}[2s_4\check{\rho} - \check{f}^{-1} - s_4^2\check{f}]} \\ - \frac{qt^2(q - t^2)}{(q - \alpha^2)}\frac{(t - \alpha\check{f})}{(t\check{f} - \alpha)}\frac{\check{f}^2w(\alpha t^{-1})}{\alpha tw(\check{f}) - (t\check{f} - \alpha)(t - \alpha\check{f})\check{f}[2s_4\check{\rho} - \check{f}^{-1} - s_4^2\check{f}]}.$$

The auxiliary equation for the leading coefficient of the polynomial is

(7.135)
$$t^2 \frac{\hat{\gamma}_n^2}{\check{\gamma}_n^2} = \frac{t^2 + q^{2n}\alpha^2 s_4^2 - 2q^n \alpha t s_4 \hat{\rho}}{1 + q^{2n}\alpha^2 t^2 s_4^2 - 2q^n \alpha t s_4 \hat{\rho}}$$

Proof. The first of these equations is derived utilising the following steps. We substitute the expression for $r_{1+}\check{a}_n/p_+$ given by (7.112) and the transformation formulae for $\check{\mathfrak{z}}_{\pm}$ as given in (7.110,7.111) into the solution at the retarded co-ordinate (7.109). This yields an equation involving $\hat{\rho}$ on one hand and \check{l}, \check{f} on the other hand. Now we perform a partial fraction expansion of this with respect to \check{l} and this produces the following expression

$$(7.136) \quad 2s_4\hat{\rho} = q^{-1}t^2\check{f}^{-1} + qt^{-2}s_4^2\check{f} - \frac{(t-\alpha\check{f})(q\check{f}-\alpha t)}{\check{f}(t\check{f}-\alpha)(q-\alpha t\check{f})}\frac{w(\check{f})}{\check{f}^2 + 1 - 2\check{f}\check{\lambda}} \\ - \frac{\alpha^4t^2(q-t^2)(q\check{f}-\alpha t)}{q(q-\alpha^2)(q-\alpha t\check{f})}\frac{w(q\alpha^{-1}t^{-1})}{q^2 + \alpha^2t^2 - 2q\alpha t\check{\lambda}} + \frac{t^2(q-t^2)(t-\alpha\check{f})}{(q-\alpha^2)(t\check{f}-\alpha)}\frac{w(\alpha t^{-1})}{\alpha^2 + t^2 - 2\alpha t\check{\lambda}},$$

which is a simple function of λ . Then one substitutes for λ using the inversion of the transformation (7.114) at the retarded time and the result is (7.134). Alternatively one can prove this formula by substituting (7.129) and (7.131) into (7.127).

We employ the variable transformation (7.112) and note that the left-hand side of this expression has been evaluated in (7.81) and (7.83). Equating these two forms gives (7.135).

Remark 7.8. The right-hand side of formula (7.134) exhibits apparent poles at $\alpha = t \check{f}$ and $q = \alpha t \check{f}$. However this is not the case as the former is cancelled by opposing contributions from the first and third terms, while the latter is cancelled by contributions from the first and second term.

Clearly (7.134) is not manifestly invertible for $\check{\rho}$, however it is possible to construct a linear equation for the retarded variables by switching from ρ to λ .

Proposition 7.10. The inverse to (7.134) is given as a relation for $\check{\lambda}$, which we give in two alternative forms

$$\begin{aligned} (7.137) \quad \check{\lambda} &= \frac{(\hat{f} - q^n \alpha t)(t - q^n \alpha s_4^2 \hat{f})}{(t\hat{f} - q^n \alpha)(1 - q^n \alpha t s_4^2 \hat{f})} \left[\hat{\lambda} - \frac{1}{2}(\hat{f} + \hat{f}^{-1}) \right] + \frac{t^4 + \hat{f}^2}{2t^2 \hat{f}} \\ &+ \frac{t^2(t^2 - 1)s_4^3}{2q^{n+1/2}[n + \frac{1}{2}]} \left[-s_4^3 \frac{(\hat{f} - q^n \alpha t)}{(1 - q^n \alpha t s_4^2 \hat{f})} \frac{q^{4n} \alpha^4 w (q^{-n} \alpha^{-1} t^{-1} s_4^{-2})}{1 + q^{2n} \alpha^2 t^2 s_4^2 - 2q^n \alpha t s_4 \hat{\rho}} + s_4^{-3} \frac{(t - q^n \alpha s_4^2 \hat{f})}{(t\hat{f} - q^n \alpha)} \frac{w(q^n \alpha t^{-1})}{t^2 + q^{2n} \alpha^2 s_4^2 - 2q^n \alpha t s_4 \hat{\rho}} \right] \\ or \end{aligned}$$

01

(7.138)

$$\begin{split} \check{\lambda} &= -\frac{(\hat{g} - q^n \alpha ts_4)(t - q^n \alpha s_4 \hat{g})}{2(\hat{g}^2 - 1)(1 - q^n \alpha ts_4 \hat{g})(t\hat{g} - q^n \alpha s_4)} \frac{w(s_4^{-1}\hat{g})}{\hat{f} - s_4^{-1}\hat{g}} + \frac{\hat{g}^2(1 - q^n \alpha ts_4 \hat{g})(t\hat{g} - q^n \alpha s_4)}{2(\hat{g}^2 - 1)(\hat{g} - q^n \alpha ts_4)(t - q^n \alpha s_4 \hat{g})} \frac{w(s_4^{-1}\hat{g}^{-1})}{\hat{f} - s_4^{-1}\hat{g}^{-1}} \\ &\quad + \frac{1}{2\alpha t^3 \sigma_4} \left[q^{1/2} \alpha t\sigma_3 + q^{5/2}(t^2 - 1)\{n + \frac{1}{2}\} \right] - \frac{1}{2t^2 s_4} \left[\hat{g} + \hat{g}^{-1} \right] \\ &\quad + \frac{\alpha^2 t(t^2 - 1)s_4^3}{2q^{1/2}[n + \frac{1}{2}]} \left[q^{2n} \alpha s_4 \frac{w(q^{-n} \alpha^{-1} t^{-1} s_4^{-2})}{(1 - q^n \alpha ts_4 \hat{g})(1 - q^n \alpha ts_4 \hat{g}^{-1})} - \alpha^{-1} s_4^{-1} \frac{w(q^n \alpha t^{-1})}{(t - q^n \alpha s_4 \hat{g})(t - q^n \alpha s_4 \hat{g}^{-1})} \right]. \end{split}$$

Proof. One can solve the compatibility relation (6.67) for the (1,2) component of $A_n^*(z;q^{-1/2}t)$ and find $\mathfrak{T}_+(z;q^{-1/2}t)$

$$\begin{aligned} (7.139) \quad \mathfrak{T}_{+}(z;q^{-1/2}t) &= \frac{1}{\chi(z,t)} \frac{\check{a}_{n}\hat{a}_{n}}{4H_{n}^{2}(R+\Delta vS)(q^{1/2}z;t)(R-\Delta vS)(q^{1/2}z;t)} \\ &\times \left[\mathfrak{R}_{-}(q^{1/2}z;t)\mathfrak{R}_{-}(q^{-1/2}z;t)\mathfrak{T}_{+}(z;q^{1/2}t) + \mathfrak{P}_{+}(q^{1/2}z;t)\mathfrak{P}_{+}(q^{-1/2}z;t)\mathfrak{T}_{-}(z;q^{1/2}t) \\ &+ \mathfrak{R}_{-}(q^{1/2}z;t)\mathfrak{P}_{+}(q^{-1/2}z;t)\mathfrak{W}_{+}(z;q^{1/2}t) - \mathfrak{P}_{+}(q^{1/2}z;t)\mathfrak{R}_{-}(q^{-1/2}z;t)\mathfrak{W}_{-}(z;q^{1/2}t) \right]. \end{aligned}$$

To simplify the calculations of the spectral matrix elements we employ an alternative parameterisation to that of (7.57), (7.58), (7.59)

$$\begin{aligned} (7.140) \quad \mathfrak{W}_{+}(z;q^{1/2}t) &= -\frac{\alpha t(z^{2}-1)(qt-\alpha z)(q^{2n+1}t\tilde{\sigma}_{6}z-\alpha)}{z^{2}(1-\alpha^{2}t^{2})(q-\alpha^{2})(q^{2n+1}t^{2}\tilde{\sigma}_{6}-1)}\mathfrak{W}_{+}(\alpha t;q^{1/2}t) \\ &+ \frac{q^{2}\alpha t(z^{2}-1)(z-\alpha t)(q^{2n}\alpha t\tilde{\sigma}_{6}z-1)}{z^{2}(q^{2}t^{2}-\alpha^{2})(q-\alpha^{2})(q^{2n+1}t^{2}\tilde{\sigma}_{6}-1)}\mathfrak{W}_{-}(q^{-1}\alpha t^{-1};q^{1/2}t) \\ &+ \frac{(z+1)(z-\alpha t)(qt-\alpha z)(q^{2}-q^{2n}\alpha^{2}t^{2}\sigma_{4}z)}{2qtz^{2}(q^{2}-q^{2n}\alpha^{2}t^{2}\sigma_{4})}w(1) \\ &- \frac{(z-1)(z-\alpha t)(qt-\alpha z)(q^{2}+q^{2n}\alpha^{2}t^{2}\sigma_{4}z)}{2qtz^{2}(q^{2}-q^{2n}\alpha^{2}t^{2}\sigma_{4}z)}w(-1) \\ &+ \frac{q^{-n-3}(z^{2}-1)(z-\alpha t)(qt-\alpha z)}{\alpha t^{2}z^{3}(q^{2}-q^{2n}\alpha^{2}t^{2}\sigma_{4})}\left[q^{4}+q^{4n}\alpha^{4}t^{4}\sigma_{4}^{2}z^{2}+q^{2n+3/2}\alpha^{2}t^{2}(q\sigma_{3}z+\sigma_{1}\sigma_{4}z-q^{1/2}(1+z^{2}))\right], \end{aligned}$$

$$\begin{aligned} (7.141) \quad \mathfrak{W}_{-}(z;q^{1/2}t) &= \frac{\alpha t(z^{2}-1)(qtz-\alpha)(q^{2n+1}t\tilde{\sigma}_{6}-\alpha z)}{z^{2}(1-\alpha^{2}t^{2})(q-\alpha^{2})(q^{2n+1}t^{2}\tilde{\sigma}_{6}-1)}\mathfrak{W}_{+}(\alpha t;q^{1/2}t) \\ &+ \frac{q^{2}\alpha t(z^{2}-1)(1-\alpha tz)(z-q^{2n}\alpha t\tilde{\sigma}_{6})}{z^{2}(q^{2}t^{2}-\alpha^{2})(q-\alpha^{2})(q^{2n+1}t^{2}\tilde{\sigma}_{6}-1)}\mathfrak{W}_{-}(q^{-1}\alpha t^{-1};q^{1/2}t) \\ &+ \frac{(z+1)(1-\alpha tz)(qtz-\alpha)(q^{2}z-q^{2n}\alpha^{2}t^{2}\sigma_{4})}{2qtz^{2}(q^{2}-q^{2n}\alpha^{2}t^{2}\sigma_{4})}w(1) \\ &+ \frac{(z-1)(1-\alpha tz)(qtz-\alpha)(q^{2}z+q^{2n}\alpha^{2}t^{2}\sigma_{4})}{2qtz^{2}(q^{2}-q^{2n}\alpha^{2}t^{2}\sigma_{4})}w(-1) \\ &- \frac{q^{-n-3}(z^{2}-1)(1-\alpha tz)(qtz-\alpha)}{\alpha t^{2}z^{3}(q^{2}-q^{2n}\alpha^{2}t^{2}\sigma_{4})}\left[q^{4}z^{2}+q^{4n}\alpha^{4}t^{4}\sigma_{4}^{2}+q^{2n+3/2}\alpha^{2}t^{2}(q\sigma_{3}z+\sigma_{1}\sigma_{4}z-q^{1/2}(1+z^{2}))\right], \end{aligned}$$

and

$$\begin{aligned} (7.142) \quad \mathfrak{T}_{+}(z;q^{1/2}t) &= -\frac{\alpha t(z-z^{-1})}{(q-\alpha^{2})(qt^{2}-1)}p_{+} \\ &\times \left[\frac{(qtz-\alpha)(qtz^{-1}-\alpha)}{(\alpha^{2}t^{2}-1)}\frac{\mathfrak{W}_{+}(\alpha t;q^{1/2}t)}{r_{1,-\frac{1}{2}}(q^{-1/2}\alpha t+q^{1/2}\alpha^{-1}t^{-1})+r_{0,-}}\right. \\ &+ q^{2}\frac{(z-\alpha t)(z^{-1}-\alpha t)}{(\alpha^{2}-q^{2}t^{2})}\frac{\mathfrak{W}_{-}(q^{-1}\alpha t^{-1};q^{1/2}t)}{r_{1,-\frac{1}{2}}(q^{-1/2}\alpha t^{-1}+q^{1/2}\alpha^{-1}t)+r_{0,-}}\right],\end{aligned}$$

$$\begin{aligned} (7.143) \quad \mathfrak{T}_{-}(z;q^{1/2}t) &= \frac{\alpha t(z-z^{-1})}{(q-\alpha^{2})(qt^{2}-1)} \frac{1}{p_{+}} \\ &\times \left[\frac{(qtz-\alpha)(qtz^{-1}-\alpha)}{(\alpha^{2}t^{2}-1)} (r_{1,-\frac{1}{2}}(q^{-1/2}\alpha t+q^{1/2}\alpha^{-1}t^{-1})+r_{0,-})\mathfrak{W}_{-}(\alpha t;q^{1/2}t) \right. \\ &+ q^{2} \frac{(z-\alpha t)(z^{-1}-\alpha t)}{(\alpha^{2}-q^{2}t^{2})} (r_{1,-\frac{1}{2}}(q^{-1/2}\alpha t^{-1}+q^{1/2}\alpha^{-1}t)+r_{0,-})\mathfrak{W}_{+}(q^{-1}\alpha t^{-1};q^{1/2}t) \right]. \end{aligned}$$

This particular parameterisation implies that the numerator of the right-hand side of (7.139) manifestly contains a factor of $(z - z^{-1})(qtz - \alpha)(qtz^{-1} - \alpha)(z - \alpha t)(z^{-1} - \alpha t)$ which is present in the denominator. This ensures that the ratio is linear in x, as it must. Finding the zero of this linear polynomial then gives λ , which after further substantial manipulation and simplification yields (7.138) and (7.137).

A crucial fact enabling further progress is the following factorisation formula for a quantity that will subsequently figure prominently in certain discriminants.

Lemma 7.5. The bi-quadratic in $\hat{\rho}, \check{\lambda}$

$$(7.144) \quad 16\sigma_4(\hat{\rho}^2\check{\lambda}^2 - \hat{\rho}^2 - \check{\lambda}^2) - 8s_4(q^2 + q\sigma_2 + \sigma_4)\hat{\rho}\check{\lambda} + 8q^{1/2}s_4(q\sigma_1 + \sigma_3)\hat{\rho} \\ + 8q^{-1/2}(\sigma_1\sigma_4 + q\sigma_3)\check{\lambda} + (q - \sigma_2)^2 - 4\sigma_1\sigma_3 + 2\sigma_4 - 2q^{-1}\sigma_2\sigma_4 + q^{-2}\sigma_4^2,$$

is a perfect square which can be given in either of two ways. In the first way this is the square of

$$\begin{array}{l} (7.145) \\ qt^{2}\hat{f}^{-2} - q^{1/2}t^{2}\sigma_{1}\hat{f}^{-1} + q^{-1/2}t^{-2}\sigma_{3}\hat{f} - q^{-1}t^{-2}\sigma_{4}\hat{f}^{2} + qt^{-2} - q^{-1}t^{2}\sigma_{4} + q^{3/2}\{n + \frac{1}{2}\}\alpha^{-1}t^{-3}(t^{2} - 1)(\hat{f} + t^{4}\hat{f}^{-1}) \\ & - 2qs_{4}(t^{-2}\hat{f} - t^{2}\hat{f}^{-1})\hat{\rho} - q\frac{(1 - s_{4}^{2}\hat{f}^{2})(\hat{f} - q^{n}\alpha t)(t - q^{n}\alpha s_{4}^{2}\hat{f})}{\hat{f}^{2}(t\hat{f} - q^{n}\alpha)(1 - q^{n}\alpha ts_{4}^{2}\hat{f})} \frac{w(\hat{f})}{1 + s_{4}^{2}\hat{f}^{2} - 2s_{4}\hat{f}\hat{\rho}} \\ & + \frac{q^{1/2}\alpha t(t^{2} - 1)s_{4}^{3}}{[n + \frac{1}{2}]} \left[\frac{q^{2n}\alpha^{2}s_{4}^{3}(1 - q^{2n}\alpha^{2}t^{2}s_{4}^{2})(\hat{f} - q^{n}\alpha t)w(q^{-n}\alpha^{-1}t^{-1}s_{4}^{-2})}{(1 - q^{n}\alpha ts_{4}^{2}\hat{f})(1 + q^{2n}\alpha^{2}t^{2}s_{4}^{2} - 2q^{n}\alpha ts_{4}\hat{\rho})} \\ & + \frac{q^{-2n}\alpha^{-2}s_{4}^{-3}(t^{2} - q^{2n}\alpha^{2}s_{4}^{2})(t - q^{n}\alpha s_{4}^{2}\hat{f})w(q^{n}\alpha t^{-1})}{(t\hat{f} - q^{n}\alpha)(t^{2} + q^{2n}\alpha^{2}s_{4}^{2} - 2q^{n}\alpha ts_{4}\hat{\rho})} \right], \end{array}$$

and in the second way this is the square of

$$\begin{array}{ll} (7.146) \quad q^{1/2}\sigma_{3}t^{-2}\check{f} - q^{-1/2}\sigma_{1}t^{2}\check{f}^{-1} - (t^{2}\check{f}^{-2} + q^{2}t^{-2}s_{4}^{2})\left[\check{f}^{2} - 1 + q^{-1}(q + \alpha^{2})\alpha^{-1}t^{-1}(q - t^{2})\check{f}\right] \\ &\quad + 2\check{\lambda}(t^{2}\check{f}^{-1} - q^{2}t^{-2}s_{4}^{2}\check{f}) + \frac{q(\check{f}^{2} - 1)(q\check{f} - \alpha t)(t - \alpha\check{f})}{\check{f}^{2}(t\check{f} - \alpha)(q - \alpha t\check{f})} \frac{w(\check{f})}{\check{f}^{2} + 1 - 2\check{f}\check{\lambda}} \\ &\quad + \frac{t(q - t^{2})}{q\alpha(q - \alpha^{2})} \left[\frac{t - \alpha\check{f}}{t\check{f} - \alpha} \frac{q^{2}(t^{2} - \alpha^{2})w(\alpha t^{-1})}{\alpha^{2} + t^{2} - 2\alpha t\check{\lambda}} + \frac{q\check{f} - \alpha t}{q - \alpha t\check{f}} \frac{\alpha^{4}(q^{2} - \alpha^{2}t^{2})w(q\alpha^{-1}t^{-1})}{q^{2} + \alpha^{2}t^{2} - 2q\alpha t\check{\lambda}} \right]. \end{array}$$

Proof. In the first way we substitute (7.138) into (7.144), whereas in the second way we substitute (7.136) into (7.144). After considerable simplification we arrive at the two results.

We are now in a position to derive the evolution equations for the f-variable, firstly in the advanced direction.

Proposition 7.11. Assuming $\check{f} \neq 0$, $t\check{f} - \alpha \neq 0$, and $q - \alpha t\check{f} \neq 0$ we have the forward evolution for f

$$\begin{array}{ll} (7.147) \quad \hat{f} = \left\{ \frac{2t^2(t^2-1)s_4^2}{q^{n-1/2}[n+\frac{1}{2}]} \left[-\frac{q^{4n}\alpha^4 s_4^4 w(q^{-n}\alpha^{-1}t^{-1}s_4^{-2})}{1+q^{2n}\alpha^2 t^2 s_4^2 - 2q^n \alpha t s_4 \hat{\rho}} + \frac{w(q^n \alpha t^{-1})}{t^2 + q^{2n}\alpha^2 s_4^2 - 2q^n \alpha t s_4 \hat{\rho}} \right] \\ \quad -\sigma_2 - q^{-1}(1-2t^2)\sigma_4 - q(1-2t^{-2}) + 2\check{\lambda} \left[2qs_4\hat{\rho} - t^2\check{f}^{-1} + q^2t^{-2}s_4^2\check{f} \right] \\ \quad -q^{1/2}\sigma_3 t^{-2}\check{f} + q^{-1/2}\sigma_1 t^2\check{f}^{-1} + (t^2\check{f}^{-2} + q^2t^{-2}s_4^2) \left[\check{f}^2 - 1 + q^{-1}(q+\alpha^2)\alpha^{-1}t^{-1}(q-t^2)\check{f} \right] \\ \quad -\frac{q(\check{f}^2 - 1)(q\check{f} - \alpha t)(t-\alpha\check{f})}{\check{f}^2(t\check{f} - \alpha)(q-\alpha t\check{f})} \frac{w(\check{f})}{\check{f}^2 + 1 - 2\check{f}\check{\lambda}} \\ \quad -\frac{t(q-t^2)}{q\alpha(q-\alpha^2)} \left[\frac{t-\alpha\check{f}}{t\check{f} - \alpha} \frac{q^2(t^2 - \alpha^2)w(\alpha t^{-1})}{\alpha^2 + t^2 - 2\alpha t\check{\lambda}} + \frac{q\check{f} - \alpha t}{q-\alpha t\check{f}} \frac{\alpha^4(q^2 - \alpha^2 t^2)w(q\alpha^{-1}t^{-1})}{q^2 + \alpha^2 t^2 - 2q\alpha t\check{\lambda}} \right] \right\} \\ \quad \times \left\{ \frac{2q^{1/2}t(t^2 - 1)}{[n+\frac{1}{2}]} \left[-\frac{q^{2n}\alpha^3 s_4^6 w(q^{-n}\alpha^{-1}t^{-1}s_4^{-2})}{1 + q^{2n}\alpha^2 t^2 s_4^2 - 2q^n \alpha t s_4 \hat{\rho}} + \frac{\alpha s_4^4 w(q^n \alpha t^{-1})}{t^2 + q^{2n}\alpha^2 s_4^2 - 2q^n \alpha t s_4 \hat{\rho}} \right] \\ \quad -2q^{3/2}\alpha^{-1}t^{-3}\{n+\frac{1}{2}\}(t^2-1) - 2q^{-1}t^{-2}(q^{1/2}\sigma_3 - 2t^2\sigma_4\check{\lambda}) + 4qt^{-2}s_4\hat{\rho} \right\}^{-1} \end{array} \right\}$$

where $\hat{\rho}$ is given by (7.134).

Proof. Whilst (7.138) can be primarily viewed as a linear equation for $\hat{\lambda}$ it can also be viewed as a quadratic equation for \hat{f} , and as such, possesses a discriminant which contains, apart from explicit squared factors, the factor (7.144). By substituting for $\hat{\rho}$ using (7.136) we can employ the result (7.146), and effect a factorisation of the quadratic into linear factors. The choice of the factors can be settled by consideration of the known solution for n = 0 and it transpires that the negative branch is appropriate. This yields (7.147).

Our last task is to derive the inverse to (7.147) and complete the system of evolution equations.

Proposition 7.12. The inverse of the evolution for f is given by

$$\begin{split} (7.148) \quad \check{f} = \begin{cases} \frac{2t^2(q-t^2)}{q-\alpha^2} \left[\frac{w(\alpha t^{-1})}{\alpha^2 + t^2 - 2\alpha t\check{\lambda}} - q^{-1}\alpha^4 \frac{w(q\alpha^{-1}t^{-1})}{q^2 + \alpha^2 t^2 - 2q\alpha t\check{\lambda}} \right] \\ &+ 4s_4 \rho\check{\lambda} + 2s_4(t^{-2}\hat{f} - t^2\hat{f}^{-1})\hat{\rho} - t^2\hat{f}^{-2} + q^{-1/2}t^2\sigma_1\hat{f}^{-1} - q^{-1}\sigma_2 - q^{-3/2}t^{-2}\sigma_3\hat{f} + q^{-2}t^{-2}\sigma_4\hat{f}^2 \\ &- q^{-2}t^{-2}(-2q + t^2 - t^4)\sigma_4 - q^{-1}t^{-2}(q(1+t^2) - 2t^4) - q^{1/2}\{n + \frac{1}{2}\}\alpha^{-1}t^{-3}(t^2 - 1)(\hat{f} + t^4\hat{f}^{-1}) \\ &+ \frac{(1 - s_4^2\hat{f}^2)(\hat{f} - q^n\alpha t)(t - q^n\alpha s_4^2\hat{f})}{\hat{f}^2(t\hat{f} - q^n\alpha t)(1 - q^n\alpha ts_4^2\hat{f})} \frac{w(\hat{f})}{1 + s_4^2\hat{f}^2 - 2s_4\hat{f}\rho} \\ &- \frac{\alpha t(t^2 - 1)s_4^3}{q^{1/2}[n + \frac{1}{2}]} \left[\frac{q^{2n}\alpha^2 s_4^3(1 - q^{2n}\alpha^2 t^2 s_4^2)(\hat{f} - q^n\alpha t)w(q^{-n}\alpha^{-1}t^{-1}s_4^{-2})}{(1 - q^n\alpha ts_4^2\hat{f})(1 + q^{2n}\alpha^2 t^2 s_4^2 - 2q^n\alpha ts_4\hat{\rho})} \\ &+ \frac{q^{-2n}\alpha^{-2}s_4^{-3}(t^2 - q^{2n}\alpha^2 s_4^2)(t - q^n\alpha s_4^2\hat{f})w(q^n\alpha t^{-1})}{(t\hat{f} - q^n\alpha)(t^2 + q^{2n}\alpha^2 s_4^2 - 2q^n\alpha ts_4\hat{\rho})} \right] \\ &\times \left\{ \frac{2\alpha t(q - t^2)}{q - \alpha^2} \left[\frac{w(\alpha t^{-1})}{\alpha^2 + t^2 - 2\alpha t\check{\lambda}} - \frac{\alpha^2 w(q\alpha^{-1}t^{-1})}{q^2 + \alpha^2 t^2 - 2q\alpha t\check{\lambda}} \right] \\ &- 2q^{-1/2}t^{-2}\sigma_3 + 2q^{-2}\alpha^{-1}t^{-3}(q - t^2)(q + \alpha^2)\sigma_4 + 4s_4\hat{\rho} + 4q^{-1}t^{-2}\sigma_4\check{\lambda} \right\}^{-1} \end{split} \right\}$$

where $\check{\lambda}$ is given by (7.138).

Proof. In a similar manner to the previous proof we can solve (7.136) for \check{f} . On appearance this polynomial should be a quintic in \check{f} , however in line with Remark 7.8, we observe this contains a factor of $\check{f}(t\check{f}-\alpha)(q-\alpha t\check{f})$ and we only have to deal with a quadratic. Upon examining this quadratic we find that the discriminant contains, again, the factor (7.144) as the only manifestly square-free factor. Upon substituting for $\check{\lambda}$ using (7.138) we can now employ (7.145) and factorise the quadratic. As in the previous case consideration of the n = 0 solution resolves the sign ambiguity in favour of the negative root, yielding (7.148).

7.4. Seed Solution. We can now make contact with the earlier theory characterising the moments, as given in Subsection 7.1, through a study of the classical "seed" solutions to the coupled recurrence system given above. We now append a subscript on the variables to indicate the n-value.

Proposition 7.13. The recurrence relations (7.134) and (7.147) admit the classical "seed" solution at n = 0

(7.149)
$$f_0(t) = q^{-1/2} \alpha t,$$

(7.150)
$$2qs_4\rho_0(t) = \frac{C(q^{1/2}\alpha^{-1}t + \sigma_4q^{-1/2}\alpha t^{-1})m(t) + D(q^{3/2}\alpha^{-1}t^{-1} + \sigma_4q^{-3/2}\alpha t)m(q^{-1}t)}{Cm(t) + Dm(q^{-1}t)}$$

where m(t) is the general solution to the second-order q-difference equation

$$(7.151) \quad (t^2 - q^2)t^4 w(\alpha t^{-1})m(q^{1/2}t) \\ + \frac{\hat{D}}{\hat{C}} \left[(t^2 - q^2)t^4 w(\alpha t^{-1}) + q^3(t^2 - 1)w(q^{-1}\alpha t) - q^{-3}(q - \alpha^2)(q^2 - \sigma_4\alpha^2)(t^2 - 1)(t^2 - q)(t^2 - q^2) \right] m(q^{-1/2}t) \\ + \frac{\hat{D}\check{D}}{\hat{C}\check{C}}q^3(t^2 - 1)w(q^{-1}\alpha t)m(q^{-3/2}t) = 0.$$

Proof. It is clear that both (7.130) and (7.131) both vanish when (7.149) holds at n = 0, therefore together they satisfy (7.117). Given (7.149) we observe that two terms vanish in (7.134) and $\rho_0(t)$ satisfies the discrete Riccati equation

(7.152)
$$2s_4\hat{\rho}_0 - \alpha^{-1}t - \alpha t^{-1}s_4^2 = \frac{qt^4w(\alpha t^{-1})\left[2qs_4\check{\rho}_0 - q^2\alpha^{-1}t^{-1} - \alpha ts_4^2\right]}{\alpha t(q-t^2)(q-\alpha^2)\left[2qs_4\check{\rho}_0 - q^2\alpha^{-1}t^{-1} - \alpha ts_4^2\right] + q^4w(q^{-1}\alpha t)}.$$

Alternatively we can derive this relation by specialising (7.137) under (7.149) which yields

(7.153)
$$t^{2} + \alpha^{2} - 2\alpha t \check{\lambda}_{0} = \frac{t^{4} w(\alpha t^{-1})}{t^{2} + \alpha^{2} s_{4}^{2} - 2\alpha t s_{4} \hat{\rho}_{0}}$$

and then using (7.114) to substitute for λ_0 . Making the standard linearising transformation

(7.154)
$$2qs_4\rho_0(t) = \frac{Am(t) + Bm(q^{-1}t)}{Cm(t) + Dm(q^{-1}t)},$$

we find, in our solution for the decoupling factors, that $\hat{A} = (q\alpha^{-1}t + \sigma_4 q^{-1}\alpha t^{-1})\hat{C}$ and $\check{B} = (q^2\alpha^{-1}t^{-1} + \sigma_4 q^{-2}\alpha t)\check{D}$ and arrive at (7.151).

Remark 7.9. Our explicit initial orthogonal polynomial variables are one specialisation of the classical solutions given above. As remarked in the proof of Proposition 7.3 we have $\mu_0 = V(\lambda_0)$ and $\nu_0 = W(\lambda_0)$ at n = 0, where λ_0 is the unique zero of the right-hand side of (7.18)

(7.155)
$$2\left[\frac{1}{2}\right]\lambda_0 = -q^{-1/2}\tilde{\sigma}_1 + q^{1/2}\tilde{\sigma}_5 - \left[1\right]\frac{m_{0,+} + m_{0,-}}{m_{0,0}}$$

We easily see that $\mathfrak{z}_{\pm,0}(t) = l_0^{\pm 3} \prod_{j=1}^6 (1 - q^{-1/2} a_j l_0^{\pm 1})$ and from the transformations (7.110,7.111) we deduce that $f_0(t) = q^{-1/2} \alpha t$. The elements of the spectral matrix A_0^* are

(7.156)
$$\mathfrak{W}_{+,0}(z;t) = z^{-3}w(z)(1-q^{-1/2}\alpha tz)(1-q^{-1/2}\alpha t^{-1}z) = W + \Delta yV,$$

(7.157)
$$\mathfrak{W}_{-,0}(z;t) = z^3 w(z^{-1})(1 - q^{-1/2}\alpha t z^{-1})(1 - q^{-1/2}\alpha t^{-1} z^{-1}) = W - \Delta y V,$$

(7.158)
$$\mathfrak{T}_{+,0}(z;t) = 2a_0[\frac{1}{2}](z-z^{-1})(z-l_0)(1-l_0^{-1}z^{-1})$$

(7.159)
$$\mathfrak{T}_{-,0}(z;t) = 0,$$

whilst those of the deformation matrix B_0^* are

(7.160)
$$\mathfrak{R}_{+,0}(z;t) = -\frac{a_0\hat{\gamma}_0}{\check{a}_0\check{\gamma}_0}(1-q^{-1/2}\alpha tz)(1-q^{-1/2}\alpha tz^{-1}),$$

(7.161)
$$\mathfrak{R}_{-,0}(z;t) = -\frac{a_0\check{\gamma}_0}{\hat{a}_0\hat{\gamma}_0}(1-q^{-1/2}\alpha t^{-1}z)(1-q^{-1/2}\alpha t^{-1}z^{-1}),$$

(7.162)
$$\mathfrak{P}_{+,0}(z;t) = 2q^{-1/2}\alpha a_0 \left[t \frac{\hat{\gamma}_0}{\check{\gamma}_0} - t^{-1} \frac{\check{\gamma}_0}{\hat{\gamma}_0} \right],$$

(7.163) $\mathfrak{P}_{-,0}(z;t) = 0.$

We also note from (7.135) and (3.16,3.10) that

(7.164)
$$2\alpha s_4 \rho_0(t) = \frac{(1+q^{-1}\alpha^2 s_4^2 t^2)m_{0,0}(q^{-1}t) - (1+q\alpha^2 s_4^2 t^{-2})m_{0,0}(t)}{q^{-1/2}t m_{0,0}(q^{-1}t) - q^{1/2}t^{-1}m_{0,0}(t)}$$

which is precisely the case of (7.150) with $D = -q^{-1}t^2C$. With this solution we also deduce that the moment recurrence (7.78) for $m_{0,0}$ coincides with (7.151).

At the beginning of this section, in Remark 7.2, we noted that some explicit solutions to the moment recurrences were given in [97]. This knowledge enables the possibility of checking any aspect of the foregoing theory either symbolically or numerically to essentially unlimited precision. We wish to report that extensive checking of all the key relations for the cases n = 0 and n = 1 has been carried out employing Mathematica code and utilities where possible by exact symbolic means or if not by numerical means to a level of one part in $10^{20} \rightarrow 10^{30}$ or better at random exact values of the input parameters. These checks have also utilised a body of unreported work [96] covering the system of Laguerre-Freud recurrences under $n \mapsto n+1$ for general $n \in \mathbb{Z}_{\geq 0}$.

We now come to the question regarding the identity of this system in the Sakai scheme. Whilst not providing an explicit transformation from our parameters and variables to those appearing in the canonical coupled q-difference system [33, 86], we offer unambiguous evidence that it is one of the classical yet full parameter set cases of the $E_7^{(1)}$ q-Painlevé system as can be seen from the following inspection of their classical solutions [71, 51, 52], and the recent systematic study of "two-Casorati" determinantal forms of

their classical solutions [69]. Clearly the τ -functions of our construction, given by (3.10) and (3.11), are of the "two-Casorati" determinant form because if we choose $b = a_r, a = a_s$ with $r, s \in \{1, 2, 3, 4\}$ say and this is employed in the definition (3.9) we have a moment determinant with elements $m_{0,0}(q^j a_r, q^k a_s), j, k \in \mathbb{Z}_{\geq 0}$ where $m_{0,0}$ given by any q-constant linear combination of the two ${}_8W_7$ solutions appearing in (7.9).

Several tasks, that have arisen in the course of our study, remain unfinished and we conclude by detailing them and the issues involved. Clarification of the explicit relationship of the evolution system (7.134,7.137,7.147,7.148) with the canonical q-difference equations as say given in [51, 52] is required. It remains to complete the construction of the $n \to n + 1$ recurrence relations which occupy a special place in our approach, but figure as Schlesinger transformations in the integrable theory. In fact our framework can easily treat the Schlesinger transformations of this example (or any for that matter), and in particular the $n \to n + 1$ recurrences, or the $a_j, \alpha \mapsto qa_j, q\alpha, j = 1, \ldots, 4$ mappings. These latter transformations are manifested in our context as specialised Christoffel-Uvarov transformations, however we will postpone this undertaking.

Significant progress has been made in finding the analog of an isomonodromic system for the elliptic Painlevé courtesy of a preprint by Eric Rains [85], and in the work by Yamada [98]. The approach taken in this former work is very much in the spirit of the present study and it would be natural to expect that a limiting case of the results reported there would correspond our own. Very recently Yamada [99] has given Lax pairs for the $E_8^{(1)}$, $E_7^{(1)}$ and $E_6^{(1)}$ q-Painlevé equations by reformulating the $E_8^{(1)}$ elliptic Painlevé Lax pair system and taking limits $E_8^{(1)} \rightarrow E_7^{(1)} \rightarrow E_6^{(1)}$. One would expect expect that a gauge transformation and co-ordinate transformations would link our Lax Pair with his.

8. Acknowledgments

This research has been supported by the Australian Research Council. The author wishes to acknowledge Alphonse Magnus, whose seminal study formed the inspiration for this work and has freely offered many clear perceptions into the subject, the generous insights shared by Eric Rains, the kind interest of Mizan Rahman, the enthusiastic encouragement from Alexander Its and the support of Peter Forrester.

References

- R. Álvarez-Nodarse and J. C. Medem. q-classical polynomials and the q-Askey and Nikiforov-Uvarov tableaus. J. Comput. Appl. Math., 135(2):197–223, 2001.
- [2] D. Arinkin and A. Borodin. Moduli spaces of d-connections and difference Painlevé equations. Duke Math. J., 134(3):515-556, 2006.
- [3] D. Arinkin and A. Borodin. τ-function of discrete isomonodromy transformations and probability. Compos. Math., 145(3):747-772, 2009. math.AG:0706.3073.
- [4] R. Askey and J. Wilson. Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials. Mem. Amer. Math. Soc., 54(319):iv+55, 1985.
- [5] N. M. Atakishiyev and S. K. Suslov. On the moments of classical and related polynomials. Rev. Mexicana Fis., 34(2):147– 151, 1988.
- [6] N. M. Atakishiyev and S. K. Suslov. Difference hypergeometric functions. In Progress in approximation theory (Tampa, FL, 1990), volume 19 of Springer Ser. Comput. Math., pages 1–35. Springer, New York, 1992.
- [7] G. Bangerezako. The factorization method for the Askey-Wilson polynomials. J. Comput. Appl. Math., 107(2):219-232, 1999.
- [8] G. Bangerezako. The fourth order difference equation for the Laguerre-Hahn polynomials orthogonal on special non-uniform lattices. Ramanujan J., 5(2):167–181, 2001.
- [9] G. Bangerezako and M Foupouagnigni. Laguerre-Freud equations for the recurrence coefficients of the Laguerre-Hahn orthogonal polynomials on special non-uniform lattices. ICTP preprint 2003/119, 2003.
- [10] G. Bangerezako and M. N. Hounkonnou. The factorization method for the general second-order q-difference equation and the Laguerre-Hahn polynomials on the general q-lattice. J. Phys. A, 36(3):765–773, 2003.
- [11] G. Bangerezako and A. P. Magnus. The factorization method for the semi-classical polynomials. In Self-similar systems (Dubna, 1998), pages 295–300. Joint Inst. Nuclear Res., Dubna, 1999.
- [12] R. J. Baxter. Exactly Solved Models in Statistical Mechanics. Academic Press, London, 1982.
- [13] S. Belmehdi and A. Ronveaux. Laguerre-Freud's equations for the recurrence coefficients of semi-classical orthogonal polynomials. J. Approx. Theory, 76(3):351–368, 1994.
- [14] P. Biane. Orthogonal polynomials on the unit circle, q-Gamma weights, and discrete Painlevé equations. math.CA:0901.0947, 2009.
- [15] L. Boelen, C. Smet, and W. Van Assche. q-Discrete Painlevé equations for recurrence coefficients of modified q-Freud orthogonal polynomials. math.CA/0808.0982, 2008.

- [16] A. Borodin. Discrete gap probabilities and discrete Painlevé equations. Duke Math. J., 117(3):489–542, 2003.
- [17] A. Borodin. Isomonodromy transformations of linear systems of difference equations. Ann. of Math. (2), 160(3):1141–1182, 2004.
- [18] A. Borodin and D. Boyarchenko. Distribution of the first particle in discrete orthogonal polynomial ensembles. Comm. Math. Phys., 234(2):287–338, 2003.
- [19] Y. Chen and M. E. H. Ismail. Ladder operators and differential equations for orthogonal polynomials. J. Phys. A, 30(22):7817–7829, 1997.
- [20] Y. Chen and M. E. H. Ismail. Ladder operators for q-orthogonal polynomials. J. Math. Anal. Appl., 345(1):1–10, 2008.
- [21] Y. Chen, M. E. H. Ismail, and W. Van Assche. Tau-function constructions of the recurrence coefficients of orthogonal polynomials. Adv. in Appl. Math., 20(2):141–168, 1998.
- [22] R. S. Costas-Santos. Classical orthogonal polynomials: a general difference calculus approach. math.CA/0612097, 2006.
- [23] A. S. Fokas, A. R. It-s, and A. V. Kitaev. Discrete Painlevé equations and their appearance in quantum gravity. Comm. Math. Phys., 142(2):313–344, 1991.
- [24] A. S. Fokas, A. R. It-s, and A. V. Kitaev. The isomonodromy approach to matrix models in 2D quantum gravity. Comm. Math. Phys., 147(2):395–430, 1992.
- [25] P. J. Forrester. Log Gases and Random Matrices, volume 34 of London Mathematical Society Monograph. Princeton University Press, Princeton NJ, first edition, 2010.
- [26] P. J. Forrester and N. S. Witte. Discrete Painlevé equations, orthogonal polynomials on the unit circle, and N-recurrences for averages over U(N)—P_{III}, and P_V τ-functions. Int. Math. Res. Not., (4):160–183, 2004.
- [27] P. J. Forrester and N. S. Witte. Bi-orthogonal polynomials on the unit circle, regular semi-classical weights and integrable systems. Constr. Approx., 24(2):201–237, 2006.
- [28] M. Foupouagnigni. On difference equations for orthogonal polynomials on nonuniform lattices. J. Difference Equ. Appl., 14(2):127–174, 2008.
- [29] M. Foupouagnigni, M. N. Hounkonnou, and A. Ronveaux. Laguerre-Freud equations for the recurrence coefficients of D_{ω} semi-classical orthogonal polynomials of class one. In *Proceedings of the VIIIth Symposium on Orthogonal Polynomials and Their Applications (Seville, 1997)*, volume 99, pages 143–154, 1998.
- [30] G. Freud. Orthogonal Polynomials. Pergamon Press, Oxford, 1971.
- [31] G. Gasper and M. Rahman. Basic Hypergeometric Series, volume 96 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, second edition, 2004. With a foreword by Richard Askey.
- [32] A. Ghressi and L. Khériji. The symmetrical H_q-semiclassical orthogonal polynomials of class one. SIGMA Symmetry Integrability Geom. Methods Appl., 5:Paper 076, 22, 2009.
- [33] B. Grammaticos and A. Ramani. On a novel q-discrete analogue of the Painlevé VI equation. Phys. Lett. A, 257(5-6):288– 292, 1999.
- [34] D. P. Gupta and D. R. Masson. Solutions to the associated q-Askey-Wilson polynomial recurrence relation. In Approximation and computation (West Lafayette, IN, 1993), volume 119 of Internat. Ser. Numer. Math., pages 273–284. Birkhäuser Boston, Boston, MA, 1994.
- [35] W. Hahn. Beiträge zur Theorie der Heineschen Reihen. Die 24 Integrale der Hypergeometrischen q-Differenzengleichung. Das q-Analogon der Laplace-Transformation. Math. Nachr., 2:340–379, 1949.
- [36] W. Hahn. Über Orthogonalpolynome, die q-Differenzengleichungen genügen. Math. Nachr., 2:4–34, 1949.
- [37] W. Hahn. Uber die höheren Heineschen Reihen und eine einheitliche Theorie der sogenannten speziellen Funktionen. Math. Nachr., 3:257–294, 1950.
- [38] W. Hahn. Über uneigentliche Lösungen linearer geometrischer Differenzengleichungen. Math. Ann., 125:67–81; Berichtigung, 125, 324 (1953), 1952.
- [39] A. Iatrou and J. A. G. Roberts. Integrable mappings of the plane preserving biquadratic invariant curves. II. Nonlinearity, 15(2):459–489, 2002.
- [40] M. E. H. Ismail. The Askey-Wilson operator and summation theorems. In Mathematical analysis, wavelets, and signal processing (Cairo, 1994), volume 190 of Contemp. Math., pages 171–178. Amer. Math. Soc., Providence, RI, 1995.
- [41] M. E. H. Ismail. Functional equations and electrostatic models for orthogonal polynomials. In Random matrix models and their applications, volume 40 of Math. Sci. Res. Inst. Publ., pages 225–244. Cambridge Univ. Press, Cambridge, 2001.
- [42] M. E. H. Ismail. Difference equations and quantized discriminants for q-orthogonal polynomials. Adv. in Appl. Math., 30(3):562–589, 2003.
- [43] M. E. H. Ismail. Classical and quantum orthogonal polynomials in one variable, volume 98 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, 2005. With two chapters by Walter Van Assche, With a foreword by Richard A. Askey.
- [44] M. E. H. Ismail, I. Nikolova, and P. Simeonov. Difference equations and discriminants for discrete orthogonal polynomials. *Ramanujan J.*, 8(4):475–502 (2005), 2004.
- [45] M. E. H. Ismail and M. Rahman. The associated Askey-Wilson polynomials. Trans. Amer. Math. Soc., 328(1):201–237, 1991.
- [46] M. E. H. Ismail and P. Simeonov. q-Difference operators for orthogonal polynomials. Journal of Computational and Applied Mathematics, 233(3):749 – 761, 2009. 9th OPSFA Conference.
- [47] M. E. H. Ismail and D. Stanton. q-Taylor theorems, polynomial expansions, and interpolation of entire functions. J. Approx. Theory, 123(1):125–146, 2003.

- [48] M. E. H. Ismail and N. S. Witte. Discriminants and functional equations for polynomials orthogonal on the unit circle. J. Approx. Theory, 110(2):200–228, 2001.
- [49] A. R. It·s, A. V. Kitaev, and A. S. Fokas. Matrix models of two-dimensional quantum gravity, and isomonodromic solutions of Painlevé "discrete equations". Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI), 187(Differentsialnaya Geom. Gruppy Li i Mekh. 12):3–30, 171, 174, 1991.
- [50] M. Jimbo and H. Sakai. A q-analog of the sixth Painlevé equation. Lett. Math. Phys., 38(2):145–154, 1996.
- [51] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta, and Y. Yamada. 10 E₉ solution to the elliptic Painlevé equation. J. Phys. A, 36(17):L263–L272, 2003.
- [52] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta, and Y. Yamada. Hypergeometric solutions to the q-Painlevé equations. Int. Math. Res. Not., (47):2497–2521, 2004.
- [53] E. G. Kalnins and W. Miller, Jr. Symmetry techniques for q-series: Askey-Wilson polynomials. Rocky Mountain J. Math., 19(1):223–230, 1989. Constructive Function Theory—86 Conference (Edmonton, AB, 1986).
- [54] L. Kheriji. An introduction to the H_q -semiclassical orthogonal polynomials. Methods Appl. Anal., 10(3):387–411, 2003.
- [55] L. Khériji and P. Maroni. The H_q -classical orthogonal polynomials. Acta Appl. Math., 71(1):49–115, 2002.
- [56] R. Koekoek and R. F. Swarttouw. The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue. Technical Report 98-17, Department of Technical Mathematics and Iformatics, Delft University of Technology, Delft, The Netherlands, 1998. http://aw.twi.tudelft.nl/ koekoek/askey/.
- [57] H. T. Koelink and T. H. Koornwinder. q-special functions, a tutorial. In Deformation theory and quantum groups with applications to mathematical physics (Amherst, MA, 1990), volume 134 of Contemp. Math., pages 141–142. Amer. Math. Soc., Providence, RI, 1992.
- [58] T. H. Koornwinder. The structure relation for Askey-Wilson polynomials. J. Comput. Appl. Math., 207(2):214–226, 2007.
- [59] P. A. Lesky. Eine Charakterisierung der klassischen kontinuierlichen-diskreten- und q-Orthogonal polynome. Shaker Verlag, Aachen, 2005.
- [60] A. P. Magnus. Associated Askey-Wilson polynomials as Laguerre-Hahn orthogonal polynomials. In Orthogonal polynomials and their applications (Segovia, 1986), volume 1329 of Lecture Notes in Math., pages 261–278. Springer, Berlin, 1988.
- [61] A. P. Magnus. Painlevé equations for semi-classical recurrence coefficients. CA/9409228, 1994.
- [62] A. P. Magnus. Painlevé-type differential equations for the recurrence coefficients of semi-classical orthogonal polynomials. In Proceedings of the Fourth International Symposium on Orthogonal Polynomials and their Applications (Evian-Les-Bains, 1992), volume 57, pages 215–237, 1995.
- [63] A. P. Magnus. Special nonuniform lattice (snul) orthogonal polynomials on discrete dense sets of points. In Proceedings of the International Conference on Orthogonality, Moment Problems and Continued Fractions (Delft, 1994), volume 65, pages 253-265, 1995.
- [64] A. P. Magnus. Elliptic hypergeometric solutions to elliptic difference equations. SIGMA Symmetry Integrability Geom. Methods Appl., 5:Paper 038, 12, 2009.
- [65] F. Marcellán and L. Salto. Discrete semi-classical orthogonal polynomials. J. Differ. Equations Appl., 4(5):463–496, 1998.
- [66] P. Maroni. Une caractérisation des polynômes orthogonaux semi-classiques. C. R. Acad. Sci. Paris Sér. I Math., 301(6):269– 272, 1985.
- [67] P. Maroni and M. Mejri. The $I_{(q,\omega)}$ classical orthogonal polynomials. Appl. Numer. Math., 43(4):423–458, 2002.
- [68] P. Maroni and M. Mejri. The symmetric D_{ω} -semi-classical orthogonal polynomials of class one. Numer. Algorithms, 49(1-4):251–282, 2008.
- [69] T. Masuda. Hypergeometric τ -functions of the q-Painlevé system of type $E_7^{(1)}$. SIGMA Symmetry Integrability Geom. Methods Appl., 5:Paper 035, 30, 2009.
- [70] M. Mejri. q-extension of some symmetrical and semi-classical orthogonal polynomials of class one. Appl. Anal. Discrete Math., 3(1):78–87, 2009.
- [71] M. Murata, H. Sakai, and J. Yoneda. Riccati solutions of discrete Painlevé equations with Weyl group symmetry of type $E_s^{(1)}$. J. Math. Phys., 44(3):1396–1414, 2003.
- [72] F. W. Nijhoff. On a q-deformation of the discrete Painlevé i equation and q-orthogonal polynomials. Lett. Math. Phys., 30(4):327–336, 1994.
- [73] A. F. Nikiforov and S. K. Suslov. Systems of classical orthogonal polynomials of a discrete variable on nonuniform grids. Akad. Nauk SSSR Inst. Prikl. Mat. Preprint, (8):33, 1985.
- [74] A. F. Nikiforov and S. K. Suslov. Classical orthogonal polynomials of a discrete variable on nonuniform lattices. In Group theoretical methods in physics, Vol. I (Yurmala, 1985), pages 505–511. VNU Sci. Press, Utrecht, 1986.
- [75] A. F. Nikiforov and S. K. Suslov. Classical orthogonal polynomials of a discrete variable on nonuniform lattices. Lett. Math. Phys., 11(1):27–34, 1986.
- [76] A. F. Nikiforov, S. K. Suslov, and V. B. Uvarov. Construction of particular solutions for a difference equation of hypergeometric type. Akad. Nauk SSSR Inst. Prikl. Mat. Preprint, (142):15, 1984.
- [77] A. F. Nikiforov, S. K. Suslov, and V. B. Uvarov. Classical orthogonal polynomials of a discrete variable on nonuniform grids. Dokl. Akad. Nauk SSSR, 291(5):1056–1059, 1986.
- [78] A. F. Nikiforov, S. K. Suslov, and V. B. Uvarov. Classical orthogonal polynomials of a discrete variable. Springer Series in Computational Physics. Springer-Verlag, Berlin, 1991. Translated from the Russian.
- [79] C. M. Ormerod, N. S. Witte, and P. J. Forrester. Connection preserving deformations and q-semi-classical orthogonal polynomials. 2010.
- [80] P. I. Pastro. Orthogonal polynomials and some q-beta integrals of Ramanujan. J. Math. Anal. Appl., 112(2):517–540, 1985.

- [81] M. Rahman. An integral representation of a $10\varphi_9$ and continuous bi-orthogonal $10\varphi_9$ rational functions. Canad. J. Math., 38(3):605-618, 1986.
- [82] M. Rahman. q-Wilson functions of the second kind. SIAM J. Math. Anal., 17(5):1280-1286, 1986.
- [83] M. Rahman and S. K. Suslov. Barnes and Ramanujan-type integrals on the q-linear lattice. SIAM J. Math. Anal., 25(3):1002-1022, 1994.
- [84] M. Rahman and S. K. Suslov. The Pearson equation and the beta integrals. SIAM J. Math. Anal., 25(2):646-693, 1994.
- [85] E. Rains. The isomondromy interpretation of the elliptic Painlevé equation. I. the hypergeometric solution. 2007.
- [86] H. Sakai. Rational surfaces associated with affine root systems and geometry of the Painlevé equations. Comm. Math. Phys., 220(1):165–229, 2001.
- [87] H. Sakai. A q-analog of the Garnier system. Funkcial. Ekvac., 48(2):273–297, 2005.
- [88] H. Sakai. Lax form of the q-Painlevé equation associated with the $A_2^{(1)}$ surface. J. Phys. A, 39(39):12203–12210, 2006. [89] V. P. Spiridonov and A. S. Zhedanov. Generalized eigenvalue problem and a new family of rational functions biorthogonal on elliptic grids. In Special functions 2000: current perspective and future directions (Tempe, AZ), volume 30 of NATO Sci. Ser. II Math. Phys. Chem., pages 365–388. Kluwer Acad. Publ., Dordrecht, 2001.
- [90] V. P. Spiridonov and A. S. Zhedanov. Elliptic grids, rational functions, and the Padé interpolation. Ramanujan J., 13(1-3):285-310, 2007.
- [91] S. K. Suslov. On the theory of difference analogues of special functions of hypergeometric type. Uspekhi Mat. Nauk, 44(2(266)):185-226, 1989.
- [92] G. Szegő. Orthogonal Polynomials. Colloquium Publications 23. American Mathematical Society, Providence, Rhode Island, third edition, 1967.
- [93] W. Van Assche. Discrete Painlevé equations for recurrence coefficients of orthogonal polynomials. In Difference equations, special functions and orthogonal polynomials, pages 687-725. World Sci. Publ., Hackensack, NJ, 2007.
- [94] N. S. Witte. Bi-orthogonal systems on the unit circle, regular semi-classical weights and integrable systems. II. J. Approx. Theory, 161(2):565-616, 2009.
- [95] N. S. Witte. Biorthogonal systems on the unit circle, regular semiclassical weights, and the discrete Garnier equations. Int. Math. Res. Not. IMRN, (6):988-1025, 2009.
- [96] N. S. Witte. Deformations of the Askey-Wilson polynomials and their Bäcklund transformations. 2010.
- [97] N. S. Witte. On a family of integrals that extend the Askey-Wilson integral. 2010.
- [98] Y. Yamada. A Lax formalism for the elliptic difference Painlevé equation. SIGMA Symmetry Integrability Geom. Methods Appl., 5:Paper 042, 15, 2009.
- [99] Y. Yamada. Lax formalism for q-painleve equations with affine weyl group symmetry of type $E_n^{(1)}$. arXiv:1004.1687v1 [math-ph], 2010.

DEPARTMENT OF MATHEMATICS AND STATISTICS. UNIVERSITY OF MELBOURNE, VICTORIA 3010, AUSTRALIA E-mail address: n.witte@ms.unimelb.edu.au