# Invariance of the Kaup-Kupershmidt equation and triangular auto-Bäcklund transformations<sup>[1](#page-0-0)</sup>

Marianna Euler and Norbert Euler

Department of Engineering Sciences and Mathematics Luleå University of Technology SE-971 87 Luleå, Sweden Emails: marianna@ltu.se; norbert@ltu.se

Abstract: We report triangular auto-Bäcklund transformations for the solutions of a fifthorder evolution equation, which is a constraint for an invariance condition of the Kaup-Kupershmidt equation derived by E. G. Reyes in his paper titled "Nonlocal symmetries and the Kaup-Kupershmidt equation" [J. Math. Phys. 46, 073507, 19 pp., 2005]. These auto-Bäcklund transformations can then be applied to generate solutions of the Kaup-Kupershmidt equation. We show that triangular auto-Bäcklund transformations result from a systematic multipotentialisation of the Kupershmidt equation.

## 1 Introduction

In his paper [\[7\]](#page-7-0), Reyes reports an invariance of the Kaup-Kupershmidt equation

<span id="page-0-1"></span>
$$
q_t = q_{xxxxx} + 5qq_{xxx} + \frac{25}{2}q_xq_{xx} + 5q^2q_x, \tag{1.1}
$$

by the following

**Proposition 1:** [\[7\]](#page-7-0) The Kaup-Kupershmidt equation,  $(1.1)$ , is invariant under the transformation  $q \mapsto \bar{q}$ , in which

<span id="page-0-3"></span> $\bar{q} = q + 3(\ln B)_{rr},$  (1.2)

where the variables q and B are related by

<span id="page-0-4"></span>
$$
q(x,t) = -\frac{B_{xxx}}{B_x} + \frac{3}{4} \left(\frac{B_{xx}}{B_x}\right)^2
$$
\n(1.3)

and  $B(x, t)$  is a solution to

<span id="page-0-2"></span>
$$
B_t = B_{xxxxx} - 5\frac{B_{xx}B_{xxxxx}}{B_x} - \frac{15}{4}\frac{B_{xxx}^2}{B_x} + \frac{65}{4}\frac{B_{xxx}B_{xx}^2}{B_x^2} - \frac{135}{16}\frac{B_{xx}^4}{B_x^3}.
$$
 (1.4)

<span id="page-0-0"></span><sup>1</sup>Communicated by F. Calogero

In the current letter we report a class of auto-Bäcklund transformations, known as the triangular auto-Bäcklund transformations (or  $\triangle$ -auto-Bäcklund transformations [\[5\]](#page-7-1)), by a systematic multipotentialisation of the equation  $(1.4)$ . Triangular auto-Bäcklund transformations have been defined and demonstrated in our recent paper [\[5\]](#page-7-1). For details on the multipotentialisation procedure we refer the reader to our papers [\[3\]](#page-6-0) and [\[4\]](#page-6-1). We show that the equation,

<span id="page-1-0"></span>
$$
v_t = v_{xxxxx} - 5\frac{v_{xx}v_{xxxx}}{v_x} + 5\frac{v_{xx}^2 v_{xxx}}{v_x^2}
$$
\n(1.5)

plays a central role in the construction of solutions of the Kaup-Kupershmidt equation and consequently we find the general stationary solution of [\(1.5\)](#page-1-0) by calculating its first integrals.

We remark that both equations  $(1.4)$  and  $(1.5)$  first appeared in a paper by J. Weiss [\[9\]](#page-7-2) on the Painlev´e analysis in the form of singularity manifold constraints for the Kupershmidt equation (see eq. (3.32) in [\[9\]](#page-7-2)) and the Caudrey-Dodd-Gibbon equation (see eq.  $(3.21)$  in [\[9\]](#page-7-2)), respectively. For more details on the Painlevé analysis and singularity manifolds we refer the reader to [\[8\]](#page-7-3). Furthermore, both [\(1.4\)](#page-0-2) and [\(1.5\)](#page-1-0) are known symmetry-integrable equations and appear in [\[6\]](#page-7-4) as part of the list of fifth-order semilinear integrable evolution equations (see equations (4.2.12) and (4.2.11) in [\[6\]](#page-7-4)). The recursion operators for both equations [\(1.4\)](#page-0-2) and [\(1.5\)](#page-1-0) are given in [\[2\]](#page-6-2).

## 2 Multipotentialisations and  $\triangle$ -auto-Bäcklund transformations

Let

<span id="page-1-1"></span>
$$
u_t = F[u] \tag{2.1}
$$

denote an evolution equation in u, where  $F[u]$  denotes a given function that depends in general on x, t, u and x-derivatives of u. The procedure to potentialise  $(2.1)$  in the equation

<span id="page-1-2"></span> $v_t = G[v_x]$  (2.2)

is well known (see e.g.  $[1]$  and  $[4]$ ). Equation  $(2.2)$  is known as the potential equation of [\(2.1\)](#page-1-1). For the benefit of the reader and to establish the notation we describe this procedure briefly: The potentialisation of  $(2.1)$  in  $(2.2)$ , if it exists, is established by a conserved current  $\Phi^t[u]$  of [\(2.1\)](#page-1-1) and the relation to the potential variable v is then

$$
v_x = \Phi^t[u],\tag{2.3}
$$

where

$$
D_t \Phi^t[u] + D_x \Phi^x[u] \bigg|_{u_t = F[u]} = 0 \tag{2.4}
$$

and  $\Phi^x$  is the conserved flux of [\(2.1\)](#page-1-1). Here  $D_t$  and  $D_x$  are the total t- and x-derivatives, repectively. Conserved currents,  $\Phi^t[u]$ , for [\(2.1\)](#page-1-1) can be obtained by the relation

$$
\Lambda[u] = \hat{E}[u]\Phi^t[u],\tag{2.5}
$$

where  $\Lambda$  is an integrating factor (also called multiplier) of [\(2.1\)](#page-1-1) that can be calculated by the relation

$$
\hat{E}[u] \left( \Lambda[u]u_t - \Lambda[u]F[u] \right) = 0. \tag{2.6}
$$

Here  $E[u]$  is the Euler operator,

$$
\hat{E}[u] = \frac{\partial}{\partial u} - D_x \circ \frac{\partial}{\partial u_x} - D_t \circ \frac{\partial}{\partial u_t} + D_x^2 \circ \frac{\partial}{\partial u_{2x}} - D_x^3 \circ \frac{\partial}{\partial u_{3x}} + \cdots
$$

A multipotentialisation of [\(2.1\)](#page-1-1) exists if [\(2.2\)](#page-1-2) can also be potentialised. A  $\triangle$ -auto-Bäcklund transformation for  $(2.1)$  exists if  $(2.2)$  potentialises back into the original equa-tion [\(2.1\)](#page-1-1). In fact we have define three different types of  $\triangle$ -auto-Bäcklund transformations. Details are in [\[5\]](#page-7-1), where several examples of  $\triangle$ -auto-Bäcklund transformations are given for some third-order and fifth-order evolution equations, as well as for systems in  $(1+1)$ dimensions and higher dimensional evolution equations.

Our starting point is the Kupershmidt equation

<span id="page-2-1"></span>
$$
K_t = K_{xxxxx} + \lambda \left( K_x K_{xxx} + K_{xx}^2 \right) - \frac{\lambda^2}{5} \left( K^2 K_{xxx} + 5K_x K_{xx} + K_x^3 \right) + \frac{\lambda^4}{125} K^4 K_x \tag{2.7}
$$

which potentialises under

$$
U_x = K \tag{2.8}
$$

in the first potential Kupershmidt equation,

<span id="page-2-0"></span>
$$
U_t = U_{xxxxx} + \lambda U_{xx} U_{xxx} - \frac{\lambda^2}{5} \left( U_x^2 U_{xxx} + U_x U_{xx}^2 \right) + \left( \frac{\lambda}{5} \right)^4 U_x^5, \tag{2.9}
$$

and, in turn again potentialises under

$$
u_x = -\frac{5}{2\lambda} \exp\left(-\frac{2\lambda}{5}U\right) \tag{2.10}
$$

in a second-potential Kupershmidt equation,

<span id="page-2-2"></span>
$$
u_t = u_{xxxxx} - \frac{5u_{xx}u_{xxxx}}{u_x} - \frac{15u_{xxx}^2}{4u_x} + \frac{65u_{xx}^2 u_{xxx}}{u_x^2} - \frac{135u_{xx}^4}{16u_x^3}.
$$
 (2.11)

In addition, the first potential Kupershmidt equation [\(2.9\)](#page-2-0) also potentialises in

<span id="page-2-3"></span>
$$
v_t = v_{xxxxx} - 5\frac{v_{xx}v_{xxxx}}{v_x} + 5\frac{v_{xx}^2 v_{xxx}}{v_x^2}
$$
\n(2.12)

under

$$
v_x = \frac{5}{\lambda} \exp\left(\frac{\lambda}{5}U\right). \tag{2.13}
$$

A connection between the Kupershmidt equation, [\(2.7\)](#page-2-1), and the equation [\(2.11\)](#page-2-2) is given by the differential substitution

<span id="page-3-0"></span>
$$
K(x,t) = -\frac{5}{2\lambda} \frac{u_{xx}}{u_x},
$$
\n(2.14)

which is obtained by combining  $U_x = K$  and  $u_x = -5/(2\lambda) \exp(-2U/(5\lambda))$ .

Furthermore, we can connect the first potential Kupershmidt equation, [\(2.9\)](#page-2-0), and the Kaup-Kupershmidt equation

$$
Q_t = Q_{xxxx} + \mu Q Q_{xxx} + \frac{5\mu}{2} Q_x Q_{xx} + \frac{\mu^2}{5} Q^2 Q_x \tag{2.15}
$$

for arbitrary constant  $\mu \neq 0$ , by the differential substitution (given in [\[6\]](#page-7-4) for the special case,  $\mu = 5$  and  $\lambda = 5$ )

$$
Q = \left(\frac{2\lambda}{\mu}\right)U_{xx} - \left(\frac{\lambda}{5\mu}\right)U_x^2.
$$
\n(2.16)

We now focus our attention on equation [\(2.11\)](#page-2-2). For this equation the most general second-order integrating factors are

$$
\Lambda_1[u] = \alpha u_x^{-5/2} u_{xx}, \qquad \Lambda_2[u] = \alpha \left( u u_x^{-5/2} u_{xx} - 2 u_x^{-1/2} \right), \tag{2.17}
$$

which leads to two potentialisations in [\(2.12\)](#page-2-3) with  $\alpha = -3/4$ , namely

$$
v_{1,x} = u_x^{-1/2}, \qquad v_{2,x} = uu_x^{-1/2}, \tag{2.18}
$$

respectively. See the Diagram below. Furthermore, the most general second-order integrating factors of equation [\(2.12\)](#page-2-3) are

$$
\Lambda_n[v] = v^n \frac{v_{xx}}{v_x^2} - \frac{n}{2} v^{n-1} \frac{1}{v_x^2}, \quad n = 0, 1, 2, 3, 4. \tag{2.19}
$$

The integrating factor  $\Lambda_0$  and  $\Lambda_4$  lead to the potentialisation of equation [\(2.12\)](#page-2-3) in equation  $(2.11)$ , namely

$$
u_{3,x} = v_x^{-2}, \qquad u_{4,x} = v^4 v_x^{-2}, \tag{2.20}
$$

respectively. See the Diagram below.

#### Diagram:



$$
4\,
$$

Combining the above potentialisations for the equation  $(2.11)$  and the equation  $(2.12)$ , leads to the following  $\triangle$ -auto-Bäcklund transformations for these equations:

Proposition 2: The equation  $(2.11)$ , viz.

$$
u_t = u_{xxxxx} - 5 \frac{u_{xx} u_{xxxx}}{u_x} - \frac{15}{4} \frac{u_{xxx}^2}{u_x} + \frac{65}{4} \frac{u_{xxx} u_{xx}^2}{u_x^2} - \frac{135}{16} \frac{u_{xx}^4}{u_x^3}
$$

admits the following  $\triangle$ -auto-Bäcklund transformations:

<span id="page-4-0"></span>
$$
u_{j+1,xx} = u_{j+1,x} \left[ \frac{u_{j,xx}}{u_{j,x}} - 2 \frac{u_{j,x}}{u_j} \right] + 4 u_{j+1,x}^{3/4} \left[ \frac{u_j^{1/2}}{u_{j,x}^{1/4}} \right],
$$
\n(2.21a)

<span id="page-4-1"></span>
$$
u_{j+1,xx} = u_{j+1,x} \left[ \frac{u_{j,xx}}{u_{j,x}} \right] + 4 u_{j+1,x}^{3/4} \left[ \frac{1}{u_{j,x}^{1/4}} \right],
$$
\n(2.21b)

where  $u_j$  and  $u_{j+1}$  satisfy [\(2.11\)](#page-2-2) for all natural numbers j.

Proposition 3: The equation [\(2.12\)](#page-2-3), viz.

$$
v_t = v_{xxxxx} - 5\frac{v_{xx}v_{xxxx}}{v_x} + 5\frac{v_{xx}^2 v_{xxx}}{v_x^2},
$$

admits the following  $\triangle$ -auto-Bäcklund transformations:

$$
v_{j+1,xx} = v_{j+1,x} \left[ \frac{v_{j,xx}}{v_{j,x}} \right] - \frac{1}{v_{j,x}},
$$
\n(2.22a)

$$
v_{j+1,xx} = v_{j+1,x} \left[ \frac{v_{j,xx}}{v_{j,x}} \right] + \frac{1}{v_{j,x}},\tag{2.22b}
$$

$$
v_{j+1,xx} = v_{j+1,x} \left[ \frac{v_{j,xx}}{v_{j,x}} - 2 \frac{v_{j,x}}{v_j} \right] + \frac{v_j^2}{v_{j,x}},
$$
\n(2.22c)

where  $v_j$  and  $v_{j+1}$  satisfy [\(2.12\)](#page-2-3) for all natural numbers j.

Remark 1: We note that in addition to the transformations given by Proposition 2 and Proposition 3, both [\(2.11\)](#page-2-2) and [\(2.12\)](#page-2-3) admit the symmetry transformation

$$
u_{j+1} = -\frac{1}{u_j} + \text{constant}, \qquad v_{j+1} = -\frac{1}{v_j} + \text{constant}
$$
 (2.23)

which are included in the Diagram: combine  $v_x = u_{1,x}^{-1/2}$  with  $v_x = u_2 u_{2,x}^{-1/2}$ , and  $u_x = v_{1,x}^{-2}$ with  $u_x = v_2^4 v_{2,x}^{-2}$ , respectively. Note that these relations were also identified by Weiss (see relation  $(3.37)$  in  $[9]$ ).

**Remark 2:** The  $\triangle$ -auto-Bäcklund transformations [\(2.21a\)](#page-4-0) and [\(2.21b\)](#page-4-1) given in Proposition 2 can be linearised, as those are in the form of first-order Bernoulli equations under the substitution  $u_{j+1,x} = w_{j+1}.$ 

## 3 Solutions

In the above Diagram it is clear that equation [\(2.12\)](#page-2-3), viz.

$$
v_t = v_{xxxxx} - \frac{5v_{xx}v_{xxxx}}{v_x} + \frac{5v_{xx}^2 v_{xxx}}{v_x^2},
$$

plays a central role in the connection between the Kupershmidt-, the Kaup-Kupershmidt and equation [\(2.11\)](#page-2-2).

We now find the most general stationary solution for equation  $(2.12)$  by calculating the first integrals and the general solution of the ordinary differential equation

<span id="page-5-0"></span>
$$
v_{xxxxx} - \frac{5v_{xx}v_{xxxx}}{v_x} + \frac{5v_{xx}^2 v_{xxx}}{v_x^2} = 0.
$$
\n(3.1)

We find that equation [\(3.1\)](#page-5-0) admits five second-order integrating factors,  $\{\Lambda_0, \Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4\}$ :

$$
\Lambda_n(x, v, v_x, v_{xx}) = a_n \frac{v_{xx}}{v_x^2} - \frac{a'_n}{2} \frac{1}{v_x^2},\tag{3.2}
$$

where

$$
a_n(v) = v^n
$$
 with  $n = 0, 1, 2, 3, 4.$ 

Here the primes denote derivatives with respect to  $v$ . The corresponding five first integrals are then

$$
I_n = \left( a_n \frac{v_{xx}}{v_x^4} - \frac{1}{2} \frac{a'_n}{v_x^2} \right) v_{xxxx} - \frac{1}{2} a_n \frac{v_{xxx}^2}{v_x^4} + \left( \frac{1}{2} a'_n \frac{v_{xx}}{v_x^3} - a_n \frac{v_{xx}^2}{v_x^5} + \frac{1}{2} \frac{a''_n}{v_x} \right) v_{xxx}
$$

$$
- \frac{1}{2} a''_n v_{xx} + \frac{1}{4} a_n^{(4)} v_x^2, \qquad n = 0, 1, 2, 3, 4.
$$

Combining  $I_0$ ,  $I_1$ ,  $I_2$ ,  $I_3$  and  $I_4$ , we obtain

$$
v_x = (I_0 v^4 - 4I_1 v^3 + 6I_2 v^2 - 4I_3 v + I_4)^{1/2},
$$

with the condition

$$
I_0I_4 - 4I_1I_3 + 3I_2^2 = 0.
$$

So the general solution of [\(3.1\)](#page-5-0) is given by the quadrature

<span id="page-5-1"></span>
$$
\int \left( I_0 v^4 - 4I_1 v^3 + 6I_2 v^2 - 4I_3 v + I_4 \right)^{-1/2} dv = x + C,
$$
\n(3.3)

where C is a constant of integration. Since the quadrature  $(3.3)$  contains five independent free constants, it represents the most general stationary solution of [\(2.12\)](#page-2-3). This solution can now be used to generate nonstationary solutions for  $(2.12)$  by the  $\triangle$ -auto-Bäcklund transformations of Proposition 3 and, by applying the transformations in the above Diagram, solutions of  $(2.11)$  can be constructed. Then by Proposition 1, the so constructed solutions of [\(2.11\)](#page-2-2) lead to solutions of the Kaup-Kupershmidt equation, as well as solutions of the Kupershmidt equation by the differential substitution [\(2.14\)](#page-3-0).

For example, by [\(3.3\)](#page-5-1) with  $I_0 = I_1 = I_2 = I_4 = C = 0$  and  $I_3 = -2$  we obtain the solution

 $v(x,t) = 2x^2$ 

for [\(2.12\)](#page-2-3) and hence the solution

$$
u(x,t) = 16x
$$

for [\(2.11\)](#page-2-2). Applying now, for example, the  $\triangle$ -auto-Bäcklund transformation [\(2.21a\)](#page-4-0) leads to the solution

$$
u(x,t) = \frac{1}{7}x^7 - \frac{4}{5}c_1x^5 + 2c_1^2x^3 - 4c_1^3x - c_1^4x^{-1} - 576c_1t + c_2,
$$

where  $c_1$  and  $c_2$  are arbitrary constants. Appplying again [\(2.21a\)](#page-4-0), setting  $c_1 = c_2 = 0$  for simplicity, we obtain the following solution for  $(2.11)$ :

$$
u(x,t) = \frac{1}{5^4 \cdot 7 \cdot 91} x^{13} + \frac{36}{5^3 \cdot 7} x^8 t + \frac{2^7 \cdot 3^4}{5^2} x^3 t^2 - 2^{12} \cdot 3^8 \cdot 7 x^{-7} t^4 - \frac{2^{10} \cdot 3^6 \cdot 7}{5} x^{-2} t^3.
$$

Proposition 1 can now be applied with the above solutions of  $(2.11)$  (put  $u(x,t) \equiv B(x,t)$  in [\(1.2\)](#page-0-3) and [\(1.3\)](#page-0-4)) to gain solutions for the Kaup-Kupershmidt equation [\(1.1\)](#page-0-1). Furthermore, solutions of the Kupershmidt equation, [\(2.7\)](#page-2-1), are then given by the differential substitution  $(2.14).$  $(2.14).$ 

## Acknowledgements

We thank the anonymous referees for their constructive and critical suggestions which led to the improvement of this letter, and the Wenner-Grenn Foundations for financial support.

## <span id="page-6-3"></span>References

- [1] Anco C S and Bluman G W, Direct construction method for conservation laws of partial differential equations Part II: General treatment, Euro. Jnl. of Applied Mathematics 13, 567–585, 2002.
- <span id="page-6-2"></span>[2] Euler M and Euler N, A class of semilinear fifth-order evolution equations: Recursion operators and multipotentialisations, J. Nonlinear Math. Phys. 18 Suppl. 1, 61–75, 2011.
- <span id="page-6-0"></span>[3] Euler N and Euler M, On nonlocal symmetries, nonlocal conservation laws and nonlocal transformations of evolution equations: Two linearisable hierarchies, J. Nonlinear Math. Phys. 16, 489–504, 2009.
- <span id="page-6-1"></span>[4] Euler N and Euler M, Multipotentialisation and iterating-solution formulae: The Krichever-Novikov equation, J. Nonlinear Math. Phys. 16 Suppl. 1, 93–106, 2009.
- <span id="page-7-4"></span><span id="page-7-1"></span>[5] Euler N and Euler M, The converse problem for the multipotentialisation of evolution equations and systems J. Nonlinear Math. Phys. 18 Suppl. 1, 77–105, 2011.
- [6] Mikhailov A V, Shabat A B and Sokolov V V, The symmetry approach to classification of integrable equations, in What is Integrability?, Editor Zhakarov E V, Springer, 115–184, 1991
- <span id="page-7-3"></span><span id="page-7-0"></span>[7] Reyes E G, Nonlocal symmetries and the Kaup-Kupershmidt equation J. Math. Phys. 46 073507, 19 pp., 2005.
- <span id="page-7-2"></span>[8] Steeb W-H and Euler N, Nonlinear Evolution Equations and Painlevé Test, World Scientific, 1988.
- [9] Weiss J, On classes of integrable systems and the Painlevé property J. Math. Phys. 25, 13–24, 1984.