

PERIODIC APPROXIMATIONS OF IRRATIONAL PSEUDO-ROTATIONS USING PSEUDOHOLOMORPHIC CURVES

BARNEY BRAMHAM

ABSTRACT. We prove that every C^∞ -smooth, area preserving diffeomorphism of the closed 2-disk having not more than one periodic point is the uniform limit of periodic C^∞ -smooth diffeomorphisms. In particular every smooth irrational pseudo-rotation can be C^0 -approximated by integrable systems. This partially answers a long standing question of A. Katok regarding zero entropy Hamiltonian systems in low dimensions. Our approach uses pseudoholomorphic curve techniques from symplectic geometry.

CONTENTS

1. Introduction	1
2. Irrational pseudo-rotations	6
3. Finite energy foliations	7
4. Proof of theorem 1	11
5. Calculation of the ω -energy	14
6. Compactness	15
7. Construction of the finite energy foliations	21
Appendix A. Proof of proposition 2	30
Appendix B. Nondegeneracy	33
References	35

1. INTRODUCTION

1.1. **The main result.** In this paper we prove a statement which is a significant first step towards answering the following general question of Katok.

*In low dimensions is every conservative dynamical system
with zero topological entropy a limit of integrable systems?*

This is stated as problem 1 in [26], but relates also to the much older [1]. Low dimensions means maps on surfaces or flows on 3-dimensional manifolds. Arguably one of the main obstacles to answering this question affirmatively is the existence of ergodic components of positive measure.

Date: April 23, 2012.

Indeed, ergodic maps (with respect to Lebesgue measure) exhibit strongly different dynamical behavior to integrable ones. In particular, almost every point is the initial condition for a dense orbit.

Using finite energy foliations by pseudoholomorphic curves we obtain a result of this nature for the class of area preserving diffeomorphisms of the 2-disk known as irrational pseudo-rotations. Irrational pseudo-rotations are of particular interest with regard to this question because they include all known ergodic disk maps with zero topological entropy.

For each $t \in \mathbb{R}$ let $R_{2\pi t} : D \rightarrow D$ denote the rigid rotation through angle $2\pi t$ on the disk $D = \{(x, y) \mid x^2 + y^2 \leq 1\}$. Let $\text{Diff}_+^\infty(D)$ and $\text{Diff}^\infty(D, \omega_0)$ be the spaces of C^∞ -smooth diffeomorphisms of the disk which preserve orientation and the area form $\omega_0 = dx \wedge dy$ respectively.

Theorem 1 (Main result). *Suppose $\varphi \in \text{Diff}^\infty(D, \omega_0)$ fixes the origin and has no other periodic points. Then it is the C^0 -limit of a sequence of maps of the form $\varphi_k = g_k^{-1} R_{2\pi p_k/q_k} g_k$, for a sequence of conjugating maps $g_k \in \text{Diff}_+^\infty(D)$ which fix the origin, and a sequence of rationals p_k/q_k converging to an irrational number.*

Elements $\varphi \in \text{Diff}^\infty(D, \omega_0)$ having precisely one periodic point are also known as (smooth) *irrational pseudo-rotations*. See definition 3.

Concerning the convergence of the approximation maps in theorem 1 it is natural to ask whether C^0 -convergence is so weak as to allow “almost anything” to be obtained in the limit? A more natural topology to consider Katok’s question, as discussed in [26], is at least a $C^{1,\varepsilon}$ -topology, $\varepsilon > 0$, in which the topological entropy is lower semi-continuous. The author would therefore like to thank Patrice Le Calvez for pointing out the following and the idea of its proof using work of Franks. A slightly more general statement is proven in appendix A.

Proposition 2. *Suppose $\varphi \in \text{Diff}^\infty(D, \omega_0)$ is the C^0 -limit of a sequence of maps of the form $\varphi_k = g_k^{-1} R_{2\pi p_k/q_k} g_k$, for a sequence of conjugating maps $g_k \in \text{Diff}_+^\infty(D)$ fixing the origin, and a sequence of rationals p_k/q_k converging to an irrational number. Then φ necessarily has precisely one periodic point. In particular it is an irrational pseudo-rotation.*

Thus C^0 -convergence is still strong enough to guarantee that the limit object is an irrational pseudo-rotation. In particular that it has zero entropy. (Due, for example, to Katok’s theorem for $C^{1,\varepsilon}$ surface diffeomorphisms [25] that bounds entropy from above by the exponential growth rate of periodic points.)

The existence of ergodic disk maps with zero entropy was established back in 1970 by Anosov and Katok [1]. Previous to their constructions it was even an open question in the non-conservative setting; Shnirelman 1930 [31] found a (non-area preserving) diffeomorphism of the disk with a dense orbit, further discussion of which can be found in [10].

It is interesting to compare the Anosov-Katok construction to the statement of theorem 1. They construct an ergodic map $\varphi : D \rightarrow D$ as the C^∞ -limit of a sequence of maps $\varphi_k : D \rightarrow D$ which are inductively constructed with the following form. For each $k \in \mathbb{N}$ there exists $(p_k, q_k) \in \mathbb{Z} \times \mathbb{N}$ relatively prime, and $g_k \in \text{Diff}^\infty(D, \omega_0)$, also fixing the origin, so that

$$(1) \quad \varphi_k = g_k^{-1} \circ R_{2\pi p_k/q_k} \circ g_k.$$

The maps g_k are arranged so that the orbits of φ_k increasingly spread out over the disk as $k \rightarrow \infty$. Consequently, the sequence $\{g_k\}$ blows up in every C^r topology. But by iteratively choosing $q_{k+1} - q_k$ sufficiently large depending on the size of $\|g_k\|_{C^k}$, the C^k norm of φ_k can be controlled. A limiting subsequence converges to a map φ with the desired “pathological” behavior such as a dense orbit, or ergodicity, or even weak mixing. More details of this method, other results and questions, are in Fayad-Katok [10]. See also Fayad-Saprykina [11].

In some sense then, theorem 1 reverses the limiting process just described above. However our conclusions are in two respects weaker than a word for word converse to the Anosov-Katok construction. Firstly, the convergence $\varphi_k \rightarrow \varphi$ in (1) is in the C^∞ -sense. Secondly, each φ_k in (1) preserves the standard area form. We do not show this for the approximation maps in theorem 1. This raises natural questions for further investigation.

A remark on integrability: The notion of integrability for a map on a surface that appears to be referred to in [26] is that the map should admit a “first integral”, that is, a continuous or smooth real valued function on the surface that is not constant on any open set but is constant on the orbits of the given map. It is obviously in this sense that each of our approximation maps in theorem 1 is integrable. A natural question is whether approximation maps can be found that are integrable in the Liouville-Arnold sense. This would follow if they were area preserving.

1.2. Idea of the proof. Pick a closed loop of Hamiltonians $H_t : D \rightarrow D$, over $t \in \mathbb{R}/\mathbb{Z}$, which generate a symplectic isotopy whose time-one map is φ . Denote the 1-periodic path of Hamiltonian vector fields on the disk by X_{H_t} . For each $n \in \mathbb{N}$ equip $Z_n = \mathbb{R}/n\mathbb{Z} \times D$ with coordinates (τ, z) . Then the vector field $R_n(\tau, z) = \partial_\tau + X_{H_t}(z)$ defines a flow on Z_n with time-one map φ and first return map φ^n .

Consider the 4-manifold $W_n := \mathbb{R} \times Z_n$ with the unique almost complex structure satisfying

$$\begin{cases} J_n \partial_{\mathbb{R}} = R_n \\ J_n|_{TD} = i \end{cases}$$

where $\partial_{\mathbb{R}}$ is the vector field dual to the \mathbb{R} -coordinate on W_n . Then (W_n, J_n) is a so called cylindrical, symmetric, almost complex manifold. That is, it is compatible in a precise way with the necessary symplectic structures for the compactness framework from symplectic field theory [4] to apply to J_n -holomorphic curves.

In section 7 we adapt techniques developed by Hofer, Wysocki, and Zehnder [19, 20, 21, 23, 24] to construct finite energy foliations of (W_n, J_n) . That is, a foliation \mathcal{F} by embedded surfaces that are the images of finite energy J_n -holomorphic curves, and where the set \mathcal{F} is invariant under translations in the \mathbb{R} -direction on W_n .




FIGURE 1. A leaf F in a finite energy foliation \mathcal{F} of $W_n = \mathbb{R} \times Z_n$ intersects the hypersurface $\{0\} \times Z_n$ in a closed 1-dimensional curve. We use this to define a disk map $\varphi_{\mathcal{F}}$.

In this setting a finite energy foliation \mathcal{F} of (W_n, J_n) can be used to define a disk map $\varphi_{\mathcal{F}} : D \rightarrow D$ in the following fairly natural way: identify the mapping torus Z_n with the hypersurface $\{0\} \times Z_n \subset W_n$. Then each leaf $F \in \mathcal{F}$ is either disjoint from Z_n or intersects it transversally in an embedded closed curve as in figure 1. The closed curve will then intersect each disk slice $\{\tau\} \times D$ in Z_n transversally and precisely once (each such disk is also J_n -holomorphic). If the curve intersects the disk $\{0\} \times D$ at a point $(0, \xi)$ and intersects the disk $\{1\} \times D$ at a point $(1, \xi')$, then we set $\varphi_{\mathcal{F}}(\xi) = \xi'$.

For each $n \in \mathbb{N}$ there is also a “trivial” \mathbb{R} -invariant foliation of (W_n, J_n) by J_n -holomorphic curves we could refer to as the *vertical* foliation $\mathcal{F}^{\text{vert}}(W_n, J_n)$. The leaves of the vertical foliation are all of the form $\mathbb{R} \times \gamma(\mathbb{R})$ for each trajectory $\gamma : \mathbb{R} \rightarrow Z_n$ of the vector field R_n . Usually this foliation is of little interest due to its “instability” and each leaf besides one has infinite so called λ -energy. However if we use the vertical foliation to define a disk map as above, we obtain the map φ of interest. The leaves of the vertical foliation are characterized by having vanishing so called ω -energy.

It turns out that there exists a sequence of finite energy foliations \mathcal{F}_n of (W_n, J_n) with the following property: each leaf in \mathcal{F}_n has ω -energy zero or

$$\{n\alpha\} \in [0, 1)$$

where $\alpha \in \mathbb{R}$ can be identified with the rotation number of the circle map $\varphi|_{\partial D}$. In particular α is irrational. $\{x\}$ denotes the fractional part of x .

Restricting to a subsequence n_j for which $\{n_j\alpha\} \rightarrow 0$ as $j \rightarrow \infty$, the leaves in \mathcal{F}_{n_j} converge in some sense to the vertical foliation $\mathcal{F}^{\text{vert}}$. Correspondingly the disk maps $\varphi_{\mathcal{F}_{n_j}}$ converge pointwise to the disk map $\varphi_{\mathcal{F}^{\text{vert}}} = \varphi$.

By being slightly more careful one obtains uniform convergence. Finally, each foliation \mathcal{F}_n has a certain symmetry that ensures that the induced map $\varphi_{\mathcal{F}_n}$ has n -th iterate $(\varphi_{\mathcal{F}_n})^n = \text{id}_D$. This is equivalent to $\varphi_{\mathcal{F}_n}$ being conjugate to a rigid rotation through an angle $2\pi p/n$ for some $p \in \{0, 1, \dots, n-1\}$.

Remark 1. We only use that φ is an irrational pseudo-rotation at two points in this argument. First, to find the nice expression for the ω -energy of the leaves. Secondly to show that the maps $\varphi_{\mathcal{F}_n}$ are roots of unity. Indeed the finite energy foliations themselves exist for generic area preserving disk maps, not just pseudo-rotations. This will be shown in [5].

1.3. Results in the literature of related interest. Using quite different techniques, Le Calvez proved in 2004, see Theorem 1.9 in [29], that every minimal C^1 -diffeomorphism of the 2-torus that is homotopic to the identity can be C^0 -approximated by periodic C^1 -diffeomorphisms. Recall that a diffeomorphism is minimal if every point is the initial condition for a dense orbit. Thus, in this result also strongly non-integrable maps are approximated by, in some sense, integrable ones.

An interesting result about irrational pseudo-rotations in the class of homeomorphisms of the open and closed annulus homotopic to the identity, was obtained by Béguin-Crovisier-LeRoux-Patou [2] and Béguin-Crovisier-LeRoux [3]. Stated for maps on the closed disk this is as follows. Let φ be an orientation preserving, measure preserving, homeomorphism of the disk with a single periodic point and boundary rotation number $\alpha \in \mathbb{R}/\mathbb{Z}$. Then the rigid rotation $R_{2\pi\alpha}$ is the C^0 -limit of maps (not necessarily area preserving) conjugate to φ . The authors of these papers note that one does not know from their approach that φ is in the closure of the set of maps conjugate to $R_{2\pi\alpha}$.

1.4. Acknowledgements. I would particularly like to thank Helmut Hofer for his interest, encouragement, and many valuable discussions and suggestions. Also for constructive comments on an earlier version of this paper. I thank Patrice LeCalvez for proposition 2, and Richard Siefring and Chris Wendl for many helpful discussions about pseudoholomorphic curves. I also wish to thank Anatole Katok for his interest.

This work is based upon work supported by the National Science Foundation under agreement No. DMS-0635607. Any opinions, findings and conclusions or recommendations in this material are those of the author and do not necessarily reflect the views of the National Science Foundation.

2. IRRATIONAL PSEUDO-ROTATIONS

Definition 3. A (smooth) *irrational, pseudo-rotation* is a C^∞ -diffeomorphism $\varphi : D \rightarrow D$ of the closed 2-disk D with the following properties: (1) φ preserves the volume form $dx \wedge dy$. (2) $\varphi(0) = 0$. (3) φ has no periodic points in $D \setminus \{0\}$.

There are equivalent definitions which admit generalizations to rational pseudo-rotations which we will not need. See for example [30] and [2].

If $\varphi : D \rightarrow D$ is an irrational pseudo-rotation, then the restriction of φ to the boundary is an orientation preserving circle diffeomorphism without periodic points. It therefore has irrational rotation number on the boundary.

More precisely, let $\pi : \mathbb{R} \rightarrow \partial D$ be the projection map $x \mapsto e^{2\pi i x}$. Then for any lift $f : \mathbb{R} \rightarrow \mathbb{R}$ of $\varphi|_{\partial D}$ via π , the limit

$$(2) \quad \tau(f) := \lim_{n \rightarrow \infty} \frac{f^n(x) - x}{n} \in \mathbb{R}$$

exists and is independent of x , see for example [27], and is called the translation number of f . Furthermore, the element $[\tau(f)] \in \mathbb{R}/\mathbb{Z}$ in the quotient space, is even independent of the choice of lift f , and is called the rotation number of $\varphi|_{\partial D}$.

Definition 4. Let $\varphi : D \rightarrow D$ be an irrational pseudo-rotation. Then we define the *rotation number* of φ to be the value on the circle

$$\text{Rot}(\varphi) := [\tau(f)] \in \mathbb{R}/\mathbb{Z}$$

for any lift $f : \mathbb{R} \rightarrow \mathbb{R}$ of the restriction $\varphi : \partial D \rightarrow \partial D$.

A preferred homotopy $\{\varphi_t\}_{t \in [0,1]}$ from $\varphi_0 = \text{id}_{\partial D}$ to $\varphi_1 = \varphi|_{\partial D}$ gives us a preferred lift of $\text{id}_{\partial D}$. Namely the terminal map of the unique lift to a homotopy in the universal covering space which begins at $\text{id}_{\mathbb{R}}$. In particular, any Hamiltonian generating φ as its time-one map, restricts to a homotopy on the boundary of the disk from $\text{id}_{\partial D}$ to $\varphi|_{\partial D}$ and thus determines a canonical lift of the latter. Using this we define:

Definition 5. Let $\varphi : D \rightarrow D$ be an irrational pseudo-rotation. Let $H_t \in C^\infty(D, \mathbb{R})$ be a path of Hamiltonians on $(D, \omega_0 = dx \wedge dy)$ generating φ as its time-one map. Then we define the *rotation number of φ with respect to H* to be the real number

$$\text{Rot}(\varphi; H) := \tau(f) \in \mathbb{R}$$

where $f : \mathbb{R} \rightarrow \mathbb{R}$ is the canonical lift of $\varphi : \partial D \rightarrow \partial D$ determined by H .

If φ is an irrational pseudo-rotation, then the unique periodic point is non-degenerate in the sense that for all $n \in \mathbb{N}$, the linearization $D\varphi^{(n)}(0)$ does not have eigenvalue 1. The proof is a well known application of the Poincaré-Birkhoff fixed point theorem. See appendix B.

3. FINITE ENERGY FOLIATIONS

For any $H \in C^\infty(\mathbb{R}/\mathbb{Z} \times D, \mathbb{R})$ with $H_t := H(t, \cdot)$ constant on the boundary of D for each $t \in \mathbb{R}/\mathbb{Z}$, the smooth time-dependent vector field $X_H(t, \cdot) := X_{H_t}$ on D defined by

$$\omega_0(X_{H_t}(z), \cdot) = -dH_t(z)$$

for all $z = (x, y) \in D$ is tangent to ∂D and therefore generates a 1-parameter family of diffeomorphisms $\phi^t : D \rightarrow D$ over $t \in \mathbb{R}$. Using that the disk is simply connected it is well known that one may find an H for any element $\varphi \in \text{Diff}^\infty(D, \omega_0)$ so that $\varphi = \phi^1$. Then H is said to generate φ .

From now on let $\varphi : D \rightarrow D$ be a fixed irrational pseudo-rotation. Unless stated otherwise $H \in C^\infty(\mathbb{R}/\mathbb{Z} \times D, \mathbb{R})$ is a 1-periodic time-dependent Hamiltonian generating φ .

Remark 2. By precomposing H with a suitable closed loop in $\text{Diff}^\infty(D, \omega_0)$ based at the identity, we may assume that the unique 1-periodic orbit of X_{H_t} corresponding to the fixed point $0 \in D$ of φ is the constant trajectory $t \mapsto 0 \in D$ for all $t \in \mathbb{R}$. This is not necessary, but makes the proof of theorem 8 slightly easier.

Define a smooth vector field R_H on the solid torus $Z := \mathbb{R}/\mathbb{Z} \times D$ by

$$(3) \quad R_H(\tau, z) := \partial_\tau + X_H(\tau, z)$$

for all $\tau \in \mathbb{R}/\mathbb{Z}, z \in D$. The first return map on $\{0\} \times D$ is canonically identified with the pseudo-rotation φ .

For each $n \in \mathbb{N}$ let Z_n be the 3-manifold-with-boundary $\mathbb{R}/n\mathbb{Z} \times D$, and R_n the vector field on Z_n that projects down to R_H under the natural projection $\pi_n : Z_n \rightarrow Z$. Clearly the first return map of the flow generated by R_n is the n -th iterate $\varphi^n : D \rightarrow D$. We will refer to the pair (Z_n, R_n) as *the mapping torus of length- n associated to H* . It will also be useful to denote by $Z_\infty := \mathbb{R} \times D$ the universal covering of each Z_n .

All the dynamical information on (Z_n, R_n) can be captured by an almost complex structure on the 4-manifold $\mathbb{R} \times Z_n$ as follows. For each $n \in \mathbb{N} \cup \{\infty\}$ define J_n on $\mathbb{R} \times Z_n$ by

$$(4) \quad \begin{cases} J_n(a, \tau, z) \partial_{\mathbb{R}} = R_n \\ J_n(a, \tau, z)|_{T_z D} = i \end{cases}$$

for all $(a, \tau, z) \in \mathbb{R} \times Z_n$. Here, $\partial_{\mathbb{R}}$ is the vector field dual to the \mathbb{R} -coordinate on $\mathbb{R} \times Z_n$, and i denotes the constant almost complex structure on the disk

coming from the standard integrable complex structure on \mathbb{C}^* . In other words $i\partial_x = \partial_y$ and $i\partial_y = -\partial_x$. Observe that J_n is independent of the \mathbb{R} -coordinate on $\mathbb{R} \times Z_n$, referred to as \mathbb{R} -invariance. This idea of coupling a suitable conservative vector field in an odd-dimensional manifold with the \mathbb{R} -direction in the product 4-manifold by an almost complex structure is due to Hofer [17].

There is a 1-parameter family of 2-tori

$$L_c := \{c\} \times \partial Z_n$$

for $c \in \mathbb{R}$, that fill the boundary $\mathbb{R} \times \partial Z_n$ of the 4-manifold. Each L_n is totally real with respect to the almost complex structure J_n , that is

$$TL_c \oplus J_n T(L_c)$$

is the full 4-dimensional tangent space at each point of L_c . These will form the boundary conditions for our pseudoholomorphic curves with boundary.

Let us describe the J_n -holomorphic half infinite cylinders with totally real boundary conditions that we are interested in. Let $\mathbb{R}^+ = [0, \infty)$ and $\mathbb{R}^- = (-\infty, 0]$. For $n \in \mathbb{N}$ we consider maps $\tilde{u} = (a, \tau, z) \in C^\infty(\mathbb{R}^\pm \times \mathbb{R}/n\mathbb{Z}, \mathbb{R} \times Z_n = \mathbb{R} \times \mathbb{R}/n\mathbb{Z} \times D)$ for which there exists $c \in \mathbb{R}$ such that

$$(5) \quad \begin{cases} \partial_s \tilde{u}(s, t) + J_n(\tilde{u}(s, t)) \partial_t \tilde{u}(s, t) = 0 & \text{for all } (s, t) \in \mathbb{R}^\pm \times \mathbb{R}/n\mathbb{Z} \\ \tilde{u}(0, t) \in L_c & \text{for all } t \in \mathbb{R}/n\mathbb{Z} \\ \tau(0, \cdot) : \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R}/n\mathbb{Z} & \text{has degree } +1, \end{cases}$$

having so called finite total energy, which we define in a moment.

This setting is a special case of that described in [4]. In particular $(\mathbb{R} \times Z_n, J_n)$ is a cylindrical symmetric almost complex manifold, and the almost complex structure J_n is compatible with the stable Hamiltonian structure (ω_n, λ_n) on Z_n given by

$$(6) \quad \begin{cases} \omega_n = dx \wedge dy + d\tau \wedge dH \\ \lambda_n = d\tau \end{cases}$$

in coordinates $(\tau, (x, y))$ on $\mathbb{R}/n\mathbb{Z} \times D$. Recall that this means that $\lambda_n \wedge \omega_n > 0$ and $\ker(\omega_n) \subset \ker(d\lambda_n)$, see for example [4] or [8]. The compactness theory in [4] leads us to consider the following two quantities for a solution to (5) which we will refer to as the ω -energy, the λ -energy, and the sum of them as the total energy. In our context the λ -energy of a solution $\tilde{u} = (a, \tau, z) \in \mathbb{R} \times \mathbb{R}/n\mathbb{Z} \times D$ to (5) is the quantity

$$(7) \quad E_\lambda(\tilde{u}) := \sup_{\psi \in \mathcal{C}} \int_{\mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z}} \tilde{u}^* \left(\psi(a) da \wedge d\tau \right) \in [0, +\infty]$$

*For convenience we use this “non-generic” almost complex structure, although it is not necessary. The pseudoholomorphic curves we encounter are either orbit cylinders or embedded with genus zero, one boundary component, and Fredholm index 2. Such curves are automatically regular.

where \mathcal{C} is the set of smooth functions $\psi : \mathbb{R} \rightarrow [0, \infty)$ for which $\int_{\mathbb{R}} \psi(s) ds = 1$. The second energy, that which in the more general context of [4] is called the ω -energy, is

$$(8) \quad E_{\omega}(\tilde{u}) := \int_{\mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z}} \tilde{u}^* \omega_n \in [0, +\infty].$$

In section 6 we will prove the following.

Lemma 6. *Let $n \in \mathbb{N}$. Suppose $\tilde{u} \in C^{\infty}(\mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z}, \mathbb{R} \times Z_n)$ is a solution to (5) with $E_{\lambda}(\tilde{u}) < \infty$. Then there exists $(s_0, t_0) \in \mathbb{R} \times \mathbb{R}/n\mathbb{Z}$ so that*

$$(9) \quad \tilde{u}(s + s_0, t + t_0) = (s, t, z(s, t))$$

for all $(s, t) \in [-s_0, \infty) \times \mathbb{R}/n\mathbb{Z}$, where $z \in C^{\infty}([-s_0, \infty) \times \mathbb{R}/n\mathbb{Z}, D)$ satisfies the Floer equation

$$(10) \quad \partial_s z(s, t) + i \left(\partial_t z(s, t) - X_H(t, z(s, t)) \right) = 0$$

for all $(s, t) \in [-s_0, \infty) \times \mathbb{R}/n\mathbb{Z}$.

This is a converse to ‘‘Gromov’s trick’’ [15].

It follows that if a solution $\tilde{u} = (a, \tau, z)$ to (5) has finite λ -energy then $E_{\lambda}(\tilde{u}) = n$, and the ω -energy of \tilde{u} is equal to the Floer energy of z ;

$$E_{\omega}(\tilde{u}) = \frac{1}{2} \int_{s=0}^{+\infty} \int_{t=0}^n |\partial_s z(s, t)|^2 + |\partial_t z(s, t) - X_H(t, z(s, t))|^2 ds dt.$$

Definition 7. For a solution $\tilde{u} = (a, \tau, z) \in C^{\infty}(\mathbb{R}^{\pm} \times \mathbb{R}/n\mathbb{Z}, \mathbb{R} \times Z_n)$ to (5) we refer to the degree of the circle map

$$z(0, \cdot) : \mathbb{R}/n\mathbb{Z} \rightarrow \partial D$$

as the *boundary index* of \tilde{u} .

Theorem 8. *Let $H \in C^{\infty}(\mathbb{R}/\mathbb{Z} \times D, \mathbb{R})$ be a Hamiltonian generating an irrational pseudo-rotation φ . Let $(Z_1, R_1), (Z_2, R_2), \dots$ be the corresponding sequence of mapping tori. For each $n \in \mathbb{N}$ let $\gamma_n : \mathbb{R}/n\mathbb{Z} \rightarrow Z_n$ be the unique n -periodic orbit of R_n for which $\gamma_n(0) \in \{0\} \times D$. Assume H was chosen so that $\gamma(t) = (t, 0)$ for all $t \in \mathbb{R}/\mathbb{Z}$ (see remark 2). Let $\alpha := \text{Rot}(\varphi; H) \in \mathbb{R}$, which is necessarily irrational.*

Then for each $n \in \mathbb{N}$ there exist two foliations $\mathcal{F}_n^+, \mathcal{F}_n^-$ of $\mathbb{R} \times Z_n$ by smoothly embedded surfaces, with the following properties:

- **Cylinder leaf:** The cylinder $C_n := \mathbb{R} \times \gamma_n(\mathbb{R}/n\mathbb{Z}) \subset \mathbb{R} \times Z_n$ is a leaf in both \mathcal{F}_n^+ and \mathcal{F}_n^- .
- **Pseudo-holomorphic:** If $F \in \mathcal{F}_n^+$ (resp. $F \in \mathcal{F}_n^-$) is not C_n , then F is parameterized by a solution \tilde{u} to (5) with J_n as in (4), with $E_{\lambda}(\tilde{u}) + E_{\omega}(\tilde{u}) < \infty$ and boundary index $\lfloor n\alpha \rfloor$ (resp. $\lceil n\alpha \rceil$).
- **\mathbb{R} -invariance:** If $F \in \mathcal{F}_n^+$ (resp. $F \in \mathcal{F}_n^-$) is a leaf and $c \in \mathbb{R}$, the set $F + c := \{(a + c, \tau, z) \mid (a, \tau, z) \in F\}$ is also a leaf in \mathcal{F}_n^+ (resp. in \mathcal{F}_n^-).

- **Uniqueness:** \mathcal{F}_n^+ and \mathcal{F}_n^- are uniquely determined by the above properties.
- **Smooth foliation:** \mathcal{F}_n^+ and \mathcal{F}_n^- are C^∞ -smooth foliations at each point on the complement of C_n .

The proof is postponed to section 7. This is a special case of a much more general result to appear in [5].

Remark 3. For each leaf $F \in \mathcal{F}_n^+$ (resp. $F \in \mathcal{F}_n^-$) that is not the cylinder, any parameterization \tilde{u} satisfying (5) has domain $\mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z}$ (resp. $\mathbb{R}^- \times \mathbb{R}/n\mathbb{Z}$). Hence the superscripts in \mathcal{F}_n^\pm . In either case, as the unique n -periodic orbit γ_n is non-degenerate the finite energy of \tilde{u} implies that the Z_n component $u_\pm(s, \cdot) : \mathbb{R}/n\mathbb{Z} \rightarrow Z_n$ converges to γ_n uniformly in $C^\infty(\mathbb{R}/n\mathbb{Z}, Z_n)$ as $s \rightarrow +\infty$ (resp. $s \rightarrow -\infty$). This can be seen in two ways. Either as a consequence of the compactness results in [17] applied to \tilde{u} (as generalized in [18, 4]); or, via lemma 6, the original work of Floer [12] applied to the disk component z .

Remark 4. Let $n \in \mathbb{N}$. The foliations \mathcal{F}_n^\pm of $\mathbb{R} \times Z_n$ can be visualized as follows. Under the projection map $\pi : \mathbb{R} \times Z_n \rightarrow Z_n$, \mathcal{F}_n^+ and \mathcal{F}_n^- project to smooth foliations of $Z_n \setminus \{\gamma_n\}$ by smoothly embedded surfaces diffeomorphic to $\mathbb{R}/n\mathbb{Z} \times (0, 1]$. The vector field $R_n = \partial_\tau + X_{H_\tau}$ on Z_n is transverse to these leaves coming from \mathcal{F}_n^+ and \mathcal{F}_n^- in opposite directions. A typical transverse disk slice to either of these looks something like in figure 2.




FIGURE 2.

The following formula will be crucial to our application. Recall that $\alpha := \text{Rot}(\varphi; H) \in \mathbb{R}$. In particular α is irrational.

Lemma 9. *Let $n \in \mathbb{N}$. For any half cylinder leaf $F \in \mathcal{F}_n^+$,*

$$E_\omega(F) = \{n\alpha\}\pi$$

where $\{\cdot\}$ denotes the fractional part of a real number.

This is proven in section 5.

4. PROOF OF THEOREM 1

We use the finite energy foliations of theorem 8 to define new disk maps.

Definition 10. For each $n \in \mathbb{N}$ define $\varphi_n : D \rightarrow D$ as follows. For $\xi \in D$ there is a unique leaf $F \in \mathcal{F}_n^+$ containing $(0, 0, \xi) \in \mathbb{R} \times \mathbb{R}/n\mathbb{Z} \times D$. Define $\varphi_n(\xi) = \xi'$ where $\xi' \in D$ is unique such that $(0, 0, \xi') \in F$.

Remark 5. φ_n is well defined: by lemma 6 if a leaf $F \in \mathcal{F}_n^+$ intersects the hypersurface

$$\hat{Z}_n := \{0\} \times Z_n$$

then it does so transversally, and for each $\tau \in \mathbb{R}/n\mathbb{Z}$ it will intersect the disk slice $\{\tau\} \times D \subset \hat{Z}_n$ in a unique point.

Remark 6. We could as easily define maps in terms of the foliations \mathcal{F}_n^- .

Remark 7. The inverse map φ_n^{-1} exists and can be defined similarly in terms of \mathcal{F}_n^+ .

Lemma 11. *Each map $\varphi_n : D \rightarrow D$ is n -periodic, that is $(\varphi_n)^n = \text{id}_D$.*

Proof. We could define n many disk maps using \mathcal{F}_n^+ ; say $\varphi_{n,i}$ for $i = 0, 1, \dots, n-1$, by requiring that $\varphi_{n,i}$ takes the point $\xi \in D$ to $\xi' \in D$ if $(0, i, \xi)$ and $(0, i+1, \xi')$ lie on the same leaf in \mathcal{F}_n^+ . Since each leaf in the foliation closes up after going once around in the $\mathbb{R}/n\mathbb{Z}$ direction, it follows that the composition $\varphi_{n,n-1} \circ \dots \circ \varphi_{n,0}$ is the identity map.

We now exploit a symmetry in \mathcal{F}_n^+ to see that each of the maps $\varphi_{n,i}$ is equal to φ_n . Consider the \mathbb{Z}_n action on $\mathbb{R} \times Z_n$ generated by the deck transformation

$$\begin{aligned} \mathcal{T} : \mathbb{R} \times Z_n &\rightarrow \mathbb{R} \times Z_n \\ \mathcal{T}(a, \tau, z) &= (a, \tau - 1, z) \end{aligned}$$

which preserves the almost complex structure; $\mathcal{T}^*J_n = J_n$. From the uniqueness part of theorem 8 we conclude that the foliation $\mathcal{T}(\mathcal{F}_n^+) := \{\mathcal{T}(F) \mid F \in \mathcal{F}_n^+\}$ is equal to \mathcal{F}_n^+ . Hence $\varphi_{n,i} = \varphi_n$ for each i . ■

Note that $\varphi_n(0) = 0$ as the cylinder C_n passes through the center of the disk slices $\{0\} \times D$ and $\{1\} \times D$ in $\hat{Z}_n = \{0\} \times Z_n$.

Lemma 12. *$\varphi_n : D \rightarrow D$ is C^∞ -smooth on $D \setminus \{0\}$.*

Proof. Recall that by theorem 8 \mathcal{F}_n^+ is a C^∞ -smooth foliation on the complement of the leaf C_n . From lemma 6 the leaves of \mathcal{F}_n^+ are uniformly transverse to the hypersurface $\hat{Z}_n = \{0\} \times Z_n$, and within this to all the disk slices, in particular to $\{0\} \times D$ and $\{1\} \times D$. Now consider any point $\xi \in D \setminus \{0\}$. Then $\xi' := \varphi_n(\xi) \in D \setminus \{0\}$. Therefore there exist local smooth foliation charts about $(0, 0, \xi)$ and $(0, 1, \xi')$. In these charts the map φ_n from its definition has a smooth expression. ■

Lemma 13. *$\varphi_n : D \rightarrow D$ is continuous at the origin.*

Proof. Let $\xi_j \in D \setminus \{0\}$ be a sequence of points converging to $0 \in D$, and $p_j := (0, 0, \xi_j) \in \mathbb{R} \times Z_n$. By lemma 21 there exists a sequence of parameterizations \tilde{u}_j of the unique leaf $F_j \in \mathcal{F}_n^+$ containing p_j that converges in the C_{loc}^∞ -topology to C_n . By lemma 6 we may choose each \tilde{u}_j to take the form $\tilde{u}_j : [S_j, \infty) \times \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R} \times Z_n$ for some $S_j \in \mathbb{R}$, with

$$\tilde{u}_j(s, t) = (s, t, z_j(s, t))$$

for some sequence z_j . It follows that the sequence of points

$$F_j \cap \{(0, 1)\} \times D = (0, 1, z_j(0, 1))$$

converges to $(0, 1, 0)$. Which means that $\varphi_n(\xi_j) = z_j(0, 1) \rightarrow 0$ as $j \rightarrow \infty$. \blacksquare

Fix a subsequence $\{n_j\}_{j \in \mathbb{N}}$ for which the sequence of fractional parts

$$(11) \quad \lim_{j \rightarrow \infty} \{n_j \alpha\} = 0.$$

From lemma (9) this implies that the ω -energies of the leaves in the foliations $\mathcal{F}_{n_j}^+$ tends to zero, as $j \rightarrow \infty$, uniformly over all leaves.

For maps $f, g : D \rightarrow D$, define $d_{C^0}(f, g)$ using the linear structure and Euclidean norm on \mathbb{R}^2 by

$$d_{C^0}(f, g) := \sup_{\xi \in D} |f(\xi) - g(\xi)|.$$

Proposition 14. *The subsequence φ_{n_j} converges to the pseudo-rotation φ in the following sense:*

$$d_{C^0}(\varphi_{n_j}, \varphi) + d_{C^0}(\varphi_{n_j}^{-1}, \varphi^{-1}) \rightarrow 0$$

as $j \rightarrow \infty$.

Proof. We show that $d_{C^0}(\varphi_{n_j}, \varphi) \rightarrow 0$ as $j \rightarrow \infty$, as the same argument will work for the inverses.

Arguing indirectly, there exists a sequence of points $\xi_j \in D$ and $\delta > 0$ such that $|\varphi_{n_j}(\xi_j) - \varphi(\xi_j)| \geq \delta$ for all $j \in \mathbb{N}$, where $|\cdot|$ is the Euclidean norm on \mathbb{R}^2 . Restricting to a subsequence we may assume that $\xi_j \rightarrow \xi$ for some $\xi \in D$, and $\delta \leq |\varphi_{n_j}(\xi_j) - \varphi(\xi_j)| \leq |\varphi_{n_j}(\xi_j) - \varphi(\xi)| + |\varphi(\xi_j) - \varphi(\xi)|$ for all $j \in \mathbb{N}$. Therefore as φ is continuous,

$$(12) \quad \frac{1}{2}\delta \leq |\varphi_{n_j}(\xi_j) - \varphi(\xi)|$$

for all j sufficiently large.

For each $j \in \mathbb{N}$ let $F_j \in \mathcal{F}_{n_j}^+$ be the unique leaf containing the point $(0, 0, \xi_j) \in \mathbb{R} \times \mathbb{R}/n_j\mathbb{Z} \times D$. Let us assume that each F_j is a half cylinder, otherwise the argument is even easier. There exists a solution \tilde{u}_j to (5) parameterizing F_j . After a holomorphic reparameterization we may assume that

$$\begin{aligned} \tilde{u}_j : [S_j, \infty) \times \mathbb{R}/n_j\mathbb{Z} &\rightarrow \mathbb{R} \times Z_{n_j} \\ \tilde{u}_j(0, 0) &= (0, 0, \xi_j) \end{aligned}$$

for some $S_j \leq 0$. For each j , $E_\lambda(\tilde{u}_j) < \infty$, so by lemma 6, \tilde{u}_j takes the form

$$(13) \quad \tilde{u}_j(s, t) = (s, t, z_j(s, t))$$

some $z_j : [S_j, \infty) \times \mathbb{R}/n_j\mathbb{Z} \rightarrow D$. Moreover, the sequence $\{\tilde{u}_j\}_{j \in \mathbb{N}}$ satisfies all the criterion for the compactness result theorem 25. In particular

$$\lim_{j \rightarrow \infty} E_\omega(\tilde{u}_j) = 0$$

due to our choice of subsequence satisfying (11). We conclude that the sequence \tilde{u}_j converges in the following sense: for each j let $\bar{u}_j : [S_j, \infty) \times \mathbb{R} \rightarrow \mathbb{R} \times Z_\infty$ be the unique lift of \tilde{u}_j to the universal covering, satisfying

$$(14) \quad \bar{u}_j(0, 0) = (0, 0, \xi_j).$$

After restricting to a further subsequence we can assume that $\bar{u}_j \rightarrow \bar{u}_\infty$ in the $C_{\text{loc}}^\infty(\mathbb{R}^2, \mathbb{R} \times Z_\infty)$ topology, that is, uniformly on compact sets, where \bar{u}_∞ takes the form

$$\bar{u}_\infty(s, t) = (s, t, \gamma(t))$$

for some $\gamma \in C^\infty(\mathbb{R}, D)$ solving $\dot{\gamma}(t) = X_H(t, \gamma(t))$ for all $t \in \mathbb{R}$. From (14) we have $\bar{u}_\infty(0, 0) = (0, 0, \xi)$. We conclude that for all j ,

$$(0, 1, \varphi_{n_j}(\xi_j)) = \bar{u}_j(0, 1).$$

The right hand side converges to

$$\begin{aligned} \bar{u}_\infty(0, 1) &= (0, 1, \gamma(1)) \\ &= (0, 1, \varphi(\gamma(0))) \\ &= (0, 1, \varphi(\xi)). \end{aligned}$$

This contradicts (12) and we are done. ■

Combining the results of this section we have proven the following statement which is almost theorem 1.

Theorem 15. *Suppose $\varphi \in \text{Diff}^\infty(D, \omega_0)$ fixes the origin and has no other periodic points. Then there exists a sequence of maps $\varphi_j \in \text{Homeo}_+(D) \cap \text{Diff}^\infty(D \setminus \{0\})$ over $j \in \mathbb{N}$, with the following properties. For each $j \in \mathbb{N}$, $\varphi_j(0) = 0$, there exists $n_j \in \mathbb{N}$ such that $\varphi_j^{n_j} = \text{id}_D$, and $d_{C^0}(\varphi_j, \varphi) + d_{C^0}(\varphi_j^{-1}, \varphi^{-1}) \rightarrow 0$ as $j \rightarrow \infty$.*

There are presumably nicer ways to go from this conclusion to the final statement. For example using changes of coordinates from the pseudoholomorphic curves themselves. This will presumably follow from a more serious analysis of the asymptotic properties of the curves.

Proof of theorem 1. It is a classical result [7, 9, 28] that if $f \in \text{Homeo}^+(D)$ satisfies $f^n = \text{id}_D$ for some $n \in \mathbb{N}$, then there exists $g \in \text{Homeo}^+(D)$ and $q \in \{0, 1, \dots, n-1\}$ so that

$$g \circ f \circ g^{-1} = R_{2\pi q/n}.$$

If $q \neq 0$ then g must fix the origin, and if $q = 0$ then $f = \text{id}_D$ anyway, so we may assume g fixes the origin. Applying this to each φ_j we find $g_j \in \text{Homeo}^+(D)$, fixing the origin, and $p_j \in \mathbb{Z}$ such that

$$\varphi_j = g_j^{-1} \circ R_{2\pi p_j/n_j} \circ g_j.$$

Now we replace g_j by a C^0 -close smooth approximation. More precisely, let \hat{g}_j be a sequence in $\text{Diff}^\infty(D)$, each fixing the origin, with $d_{C^0}(\hat{g}_j, g_j) + d_{C^0}(\hat{g}_j^{-1}, g_j^{-1}) \rightarrow 0$ as $j \rightarrow \infty$. Then the maps $\hat{\varphi}_j := \hat{g}_j^{-1} \circ R_{2\pi p_j/n_j} \circ \hat{g}_j$ are C^∞ -diffeomorphisms which converge in the C^0 -sense to the irrational pseudo-rotation φ . The maps $\hat{\varphi}_j$ satisfy the conditions of theorem 1. \blacksquare

5. CALCULATION OF THE ω -ENERGY

Choose a 1-form λ_0 on the disk so that $d\lambda_0 = \omega_0 = dx \wedge dy$. For each $n \in \mathbb{N}$ define the action functional $\mathbf{A}_n : C^\infty(\mathbb{R}/n\mathbb{Z}, Z_n) \rightarrow \mathbb{R}$ (associated to λ_0) by

$$(15) \quad \mathbf{A}_n(\sigma) := \int_{\mathbb{R}/n\mathbb{Z}} \sigma^* \lambda_0 - \int_0^n H(\sigma(t)) dt.$$

We may rewrite this as

$$(16) \quad \mathbf{A}_n(\sigma) := \int_{\mathbb{R}/n\mathbb{Z}} \sigma^* \eta_n$$

where $\eta_n := \lambda_0 - Hd\tau$ is a primitive of ω_n the 2-form used to define the ω -energy. Note that η_n restricts to a closed 1-form on ∂Z_n since R_n is tangent to ∂Z_n and $d\eta_n(R_n, \cdot) = \omega_n(R_n, \cdot) = 0$. Hence \mathbf{A}_n restricted to $C^\infty(\mathbb{R}/n\mathbb{Z}, \partial Z_n)$ descends to a map on homology.

Lemma 16. *For each $n \in \mathbb{N}$*

$$(17) \quad \mathbf{A}_n(1_{\mathbb{R}/n\mathbb{Z}}) + \lfloor n\alpha \rfloor \mathbf{A}_n(1_{\partial D}) \leq \mathbf{A}_n(\gamma_n) \leq \mathbf{A}_n(1_{\mathbb{R}/n\mathbb{Z}}) + \lceil n\alpha \rceil \mathbf{A}_n(1_{\partial D}).$$

Proof. Let $\tilde{u}_\pm = (a_\pm, u_\pm) : \mathbb{R}^\pm \times \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R} \times Z_n$ be parameterizations of leaves $F^\pm \in \mathcal{F}_n^\pm$ respectively which satisfy (5). In either case $u_\pm(s, \cdot)$ converges uniformly in $C^\infty(\mathbb{R}/n\mathbb{Z}, Z_n)$ to a parameterization $\gamma_n(\text{const}_\pm + \cdot)$ as $s \rightarrow \pm\infty$ respectively. Applying Stokes theorem,

$$E_\omega(\tilde{u}_+) = \int_{\mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z}} u_+^* \omega_n = \mathbf{A}_n(\gamma_n) - \mathbf{A}_n(u_+(0, \cdot)),$$

and

$$E_\omega(\tilde{u}_-) = \int_{\mathbb{R}^- \times \mathbb{R}/n\mathbb{Z}} u_-^* \omega_n = \mathbf{A}_n(u_-(0, \cdot)) - \mathbf{A}_n(\gamma_n).$$

Therefore, as the energies are non-negative,

$$\mathbf{A}_n(u_+(0, \cdot)) \leq \mathbf{A}_n(\gamma_n) \leq \mathbf{A}_n(u_-(0, \cdot)).$$

We observed that the action \mathbf{A}_n of a closed loop in ∂Z_n depends only on its homology class. From theorem 8 $u_+(0, \cdot) : \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R}/n\mathbb{Z} \times \partial D$ and

$u_-(0, \cdot) : \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R}/n\mathbb{Z} \times \partial D$ are homologous to $t \mapsto (t, e^{2\pi i(\lfloor n\alpha \rfloor/n)t})$ and $t \mapsto (t, e^{2\pi i(\lceil n\alpha \rceil)t})$ respectively. Therefore we get the inequalities in (17). ■

Corollary 17. *The unique 1-periodic orbit $\gamma_1 : \mathbb{R}/\mathbb{Z} \rightarrow Z_1$ has action*

$$(18) \quad \mathbf{A}_1(\gamma_1) = \mathbf{A}_1(1_{\mathbb{R}/\mathbb{Z}}) + \alpha \mathbf{A}_1(1_{\partial D})$$

where $1_{\mathbb{R}/\mathbb{Z}}$ and $1_{\partial D}$ are the closed loops in $\partial Z = \mathbb{R}/\mathbb{Z} \times \partial D$ given by $t \mapsto (t, \text{pt})$ and $t \mapsto (\text{pt}, t)$ respectively.

Proof. From the definition of \mathbf{A}_n ,

$$\begin{aligned} \mathbf{A}_n(\gamma_n) &= n\mathbf{A}_1(\gamma_1) \\ \mathbf{A}_n(1_{\mathbb{R}/n\mathbb{Z}}) &= n\mathbf{A}_1(1_{\mathbb{R}/\mathbb{Z}}) \\ \mathbf{A}_n(1_{\partial D}) &= \mathbf{A}_1(1_{\partial D}) \end{aligned}$$

for all $n \in \mathbb{N}$. Substituting these into the inequalities in lemma (16), and dividing through by n and letting $n \rightarrow +\infty$ gives (18). ■

Lemma 18. *Let $n \in \mathbb{N}$. Every leaf $F \in \mathcal{F}_n^+$ with boundary has ω -energy*

$$E_\omega(F) = \{n\alpha\}\pi$$

where $\{\cdot\}$ applied to any real number denotes its fractional part.

Proof. By Stokes theorem as in the last lemma,

$$E_\omega(F) = \mathbf{A}_n(\gamma_n) - \mathbf{A}_n(u(0, \cdot))$$

where $\tilde{u} = (a, u) : \mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R} \times Z_n$ is a parameterization of F . Using corollary 17 and that $u(0, \cdot) : \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R}/n\mathbb{Z} \times \partial D$ is homologous to $t \mapsto (t, e^{2\pi i(\lfloor n\alpha \rfloor/n)t})$, this becomes

$$\begin{aligned} E_\omega(F) &= n\left(\mathbf{A}_1(1_{\mathbb{R}/\mathbb{Z}}) + \alpha \mathbf{A}_1(1_{\partial D})\right) - \\ &\quad \left(\mathbf{A}_n(1_{\mathbb{R}/n\mathbb{Z}}) + \lfloor n\alpha \rfloor \mathbf{A}_n(1_{\partial D})\right) \\ &= \left(n\mathbf{A}_1(1_{\mathbb{R}/\mathbb{Z}}) + n\alpha \mathbf{A}_1(1_{\partial D})\right) - \\ &\quad \left(n\mathbf{A}_1(1_{\mathbb{R}/\mathbb{Z}}) + \lfloor n\alpha \rfloor \mathbf{A}_1(1_{\partial D})\right) \\ &= (n\alpha - \lfloor n\alpha \rfloor) \mathbf{A}_1(1_{\partial D}) \\ &= \{n\alpha\} \cdot \int_D dx \wedge dy. \end{aligned}$$

■

6. COMPACTNESS

Norms, such as $\|\nabla \tilde{u}_n\|_{L^\infty}$, are implicitly with respect to the Riemannian metric $dx^2 + dy^2 + d\tau^2 + da^2$ on $\mathbb{R} \times Z_n$, where (x, y) are the standard Euclidean coordinates on the disk, τ is the ‘‘coordinate’’ on $\mathbb{R}/n\mathbb{Z}$ and a is the \mathbb{R} -coordinate. This metric is J_n -invariant.

6.1. Compactness when n is uniformly bounded. We wish to prove lemma 6. This will follow from the next two lemmas.

Lemma 19 (The Floer equation from the Cauchy-Riemann equations). *Let $\tilde{u} = (a, \tau, z) : \mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R} \times \mathbb{R}/n\mathbb{Z} \times D$ be a solution to (5) for which $\|\nabla \tilde{u}\|_{L^\infty} < \infty$. Then there exist $(a_0, \tau_0) \in \mathbb{R} \times \mathbb{R}/n\mathbb{Z}$ such that*

$$(19) \quad \begin{cases} a(s, t) = s + a_0 \\ \tau(s, t) = t + \tau_0 \end{cases}$$

for all $(s, t) \in \mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z}$, and moreover $z : \mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z} \rightarrow D$ satisfies the following Floer equation:

$$(20) \quad \partial_s z(s, t) + i \left(\partial_t z(s, t) - X_H(t + \tau_0, z(s, t)) \right) = 0$$

for all $(s, t) \in \mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z}$.

Proof. Writing out in coordinates what it means for \tilde{u} to satisfy (5) gives us

$$(21) \quad (a_s - \tau_t) \partial_a + (a_t + \tau_s) \partial_\tau + \left(a_t X_H(\tau, z) + z_s + i(z_t - \tau_t X_H(\tau, z)) \right) = 0.$$

The boundary condition on \tilde{u} in (5) implies $a_t(0, t) = 0$ for all $t \in \mathbb{R}/n\mathbb{Z}$. From (21),

$$(22) \quad \begin{aligned} a_s(s, t) &= \tau_t(s, t) \\ a_t(s, t) &= -\tau_s(s, t) \end{aligned}$$

for all $(s, t) \in \mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z}$. In particular both functions a, τ lift to harmonic functions on the upper half plane with gradient bounded in L^∞ . The boundary conditions on a allow a smooth extension by reflection to the whole plane, still with gradient in L^∞ , and therefore by Liouville the partial derivatives of a are constant. So there exists $b, c, a_0 \in \mathbb{R}$ so that $a(s, t) = cs + bt + a_0$ for all $(s, t) \in \mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z}$. Putting this into (22) there exists $\tau_0 \in \mathbb{R}$ so that $\tau(s, t) = ct - bs + \tau_0$ for all $(s, t) \in \mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z}$. The n -periodicity of a in the t variable implies $b = 0$. Also $\tau(s, t + n) = \tau(s, t) + n$ for all $(s, t) \in \mathbb{R}^+ \times \mathbb{R}$ because we are assuming that $\deg(\tilde{u}) = 1$. Therefore $c = 1$. This proves (6.2).

From (21) we also have $a_t X_H(\tau, z) + z_s + i(z_t - \tau_t X_H(\tau, z)) = 0$. But we have shown that $\tau_t \equiv 1$ and $a_t \equiv 0$. Substituting these in we obtain

$$z_s + i(z_t - X_H(t + \tau_0, z)) = 0$$

as required. ■

The next statement says that we can use the above relation between the Cauchy-Riemann and Floer equations if (and only if) the λ -energy is finite.

Lemma 20. *Let $\tilde{u} = (a, \tau, z)$ be a solution to (5). Then $E_\lambda(\tilde{u}) < \infty$ implies $\|\nabla a\|_{L^\infty} < \infty$ (equivalently $\|\nabla \tau\|_{L^\infty} < \infty$).*

Proof. The equations (6.2) in the last lemma did not require the gradient bounds, and so the map $f : \mathbb{R}^+ \times \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R} \times \mathbb{R}/n\mathbb{Z}$ given by $f(s, t) := (a(s, t), \tau(s, t))$ in terms of the a and τ components of \tilde{u} , is holomorphic. Also $f(0, t) \in \{c\} \times \mathbb{R}/n\mathbb{Z}$ for all $t \in \mathbb{R}/n\mathbb{Z}$.

Arguing indirectly suppose that the gradient of a is unbounded. Then the gradient must blow up along a sequence of points that leaves every compact subset of the domain, in particular does not converge to the boundary. Therefore standard rescaling arguments applied to f yield a holomorphic plane $g : \mathbb{C} \rightarrow \mathbb{R} \times \mathbb{R}/n\mathbb{Z}$ with the following properties:

$$\begin{aligned} |\nabla g(0)| &= 1 \\ |\nabla g(\xi)| &\leq 2 \quad \text{for all } \xi \in \mathbb{C} \\ E_\lambda(g) &< \infty. \end{aligned}$$

Indeed, it is easily checked that $E_\lambda(g) \leq E_\lambda(\tilde{u})$. The first two properties imply that g has constant, non-zero, gradient from Liouville's theorem. But this implies the contradiction $E_\lambda(g) = +\infty$. ■

In our proof that the maps φ_n are continuous we used the following.

Lemma 21. *For each fixed $n \in \mathbb{N}$ \mathcal{F}_n^+ is compact in the following sense. Suppose that $F_j \in \mathcal{F}_n^+$ is a sequence of leaves over $j \in \mathbb{N}$, and $p_j \in F_j$ is a sequence of points. Suppose that $p_j \rightarrow p$ some $p \in \mathbb{R} \times Z_n$. Then there exists a sequence of parameterizations of F_{j_k} which converge in a C_{loc}^∞ -sense to a parameterization of the unique leaf in \mathcal{F}_n^+ containing p .*

Proof. This is a standard property of finite energy foliations from positivity of intersections, used many times in [23] as the λ -energy and ω -energy are uniformly bounded in j . ■

6.2. Compactness as $n \rightarrow \infty$. In our proof of convergence of the disk maps φ_n in proposition 14, we used a compactness statement for a sequence of J_n -holomorphic maps $\tilde{u}_n : [S_n, \infty) \times \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R} \times Z_n$ for which

$$\begin{aligned} E_\lambda(\tilde{u}_n) &= n \rightarrow +\infty \\ E_\omega(\tilde{u}_n) &= \{n\alpha\}\pi \end{aligned}$$

for some irrational real number α . Hence the total energy $E(\tilde{u}_n) = E_\lambda(\tilde{u}_n) + E_\omega(\tilde{u}_n)$ diverges to $+\infty$. In general, for a sequence of maps $\{\tilde{u}_n\}$ for which the total energy is unbounded one cannot expect uniform bounds on the gradient in L^∞ . However if the λ -energy grows at most linearly with n , and the ω -energy is bounded then indeed uniform bounds on $\|\nabla \tilde{u}_n\|_{L^\infty}$ can be achieved. (Actually much weaker assumptions suffice, but we will not need to explore these here.) Our arguments will be further simplified since we restricted to a subsequence for which the ω -energy of the sequence decays to zero.

Consider a sequence $\{\tilde{u}_n\}_{n \in \mathbb{N}}$ of smooth J_n -holomorphic maps, and numbers $c_n, S_n \in \mathbb{R}$ with $S_n \leq 0$, satisfying for each n ,

$$(23) \quad \begin{cases} \tilde{u}_n = (a_n, \tau_n, z_n) : [S_n, \infty) \times \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R} \times Z_n, \\ \partial_s \tilde{u}_n(s, t) + J_n(\tilde{u}_n(s, t)) \partial_t \tilde{u}_n(s, t) = 0 \\ \tilde{u}_n(S_n, t) \in L_{c_n} \\ \tau_n : (S_n, \cdot) : \mathbb{R}/n\mathbb{Z} \rightarrow \mathbb{R}/n\mathbb{Z} \quad \text{has degree 1} \end{cases}$$

for all $(s, t) \in [S_n, \infty) \times \mathbb{R}/n\mathbb{Z}$.

Proposition 22. *Suppose that $E_\lambda(\tilde{u}_n) < \infty$ for each $n \in \mathbb{N}$, and $\lim_{n \rightarrow \infty} E_\omega(\tilde{u}_n) = 0$. Then there exists $C \in (0, \infty)$ such that*

$$\|\nabla \tilde{u}_n\|_{L^\infty([S_n, \infty) \times \mathbb{R}/n\mathbb{Z})} \leq C$$

for all $n \in \mathbb{N}$.

Note that we do not assume uniform bounds on the λ -energy.

Proof. Since $E_\lambda(\tilde{u}_n) < \infty$ for each n , lemma 20 implies $\|\nabla a_n\|_{L^\infty} < \infty$ (for each n). Therefore, since also each τ_n has degree 1, lemma 19 applies so

$$\begin{cases} a_n(s, t) = s + a_n \\ \tau_n(s, t) = t + \tau_n \end{cases}$$

for all $(s, t) \in [S_n, \infty) \times \mathbb{R}/n\mathbb{Z}$, some constants $(a_n, \tau_n) \in [S_n, \infty) \times \mathbb{R}/n\mathbb{Z}$. Thus

$$\|\nabla a_n\|_{L^\infty} \leq 1 \quad \text{and} \quad \|\nabla \tau_n\|_{L^\infty} \leq 1$$

for all $n \in \mathbb{N}$.

It therefore remains to show that the gradients of the z_n are uniformly bounded. Arguing indirectly we find a sequence $\xi_n \in [S_n, \infty) \times \mathbb{R}/n\mathbb{Z}$ for which $|\nabla \tilde{u}_n(\xi_n)| \geq |\nabla z_n(\xi_n)| \rightarrow \infty$ as $n \rightarrow \infty$. A standard rescaling argument produces a J_∞ -holomorphic plane or half plane in $\mathbb{R} \times Z_\infty$. That is, a map $\tilde{v} : \mathbb{C} \rightarrow \mathbb{R} \times Z_\infty$, or $\tilde{v} : \mathbb{H} \rightarrow \mathbb{R} \times Z_\infty$ with totally real boundary conditions $\tilde{v}(\partial\mathbb{H}) \subset \{c\} \times \partial Z_\infty$ for some $c \in \mathbb{R}$. In either case, as a result of the rescaling process, $\tilde{v} = (a, \tau, z)$ has the following properties:

$$\begin{aligned} \nabla a &\equiv 0 \\ \nabla \tau &\equiv 0 \\ |\nabla \tilde{v}(0)| &> 0, \\ E_\omega(\tilde{v}) &= 0. \end{aligned}$$

The first two properties are because of the uniform bounds $\|\nabla a_j\|_{L^\infty} \leq 1$ and $\|\nabla \tau_j\|_{L^\infty} \leq 1$ respectively even before rescaling, so that rescaling ‘‘kills’’ these terms in the limit. The vanishing ω -energy is because $E_\omega(\tilde{v}) \leq \lim_{n \rightarrow \infty} E_\omega(\tilde{u}_n) = 0$ by Fatou’s lemma. Thus there exist constants $a_0, \tau_0 \in \mathbb{R}$ such that

$$\tilde{v}(s, t) = (a_0, \tau_0, z(s, t)) \in \mathbb{R} \times \mathbb{R} \times D$$

for all $(s, t) \in \mathbb{C}$ (resp. all $(s, t) \in \mathbb{H}$). That \tilde{v} is J_∞ -holomorphic translates into $z : \mathbb{C} \rightarrow D$ or $z : \mathbb{H} \rightarrow D$ satisfying the equation $a_t X_H(\tau, z) + z_s + i(z_t - \tau_t X_H(\tau, z)) = 0$, see (21). So a and τ constant implies $z_s + iz_t = 0$. (We could alternatively have just rescaled the sequence $\{z_n\}$ as in Floer theory, to get the same conclusion.) Therefore,

$$0 = E_\omega(\tilde{v}) = \int \tilde{v}^* \omega = \int \frac{1}{2} (|z_s|^2 + |z_t|^2) ds dt$$

and so z is also constant. Thus we have shown that \tilde{v} is constant, contradicting $|\nabla \tilde{v}(0)| > 0$. \blacksquare

Now standard arguments can convert these uniform bounds on the gradient in C^0 to uniform C^k -bounds on the gradient for all $k \in \mathbb{N}$. The key result is the following local statement which is proven using the $W^{k,p}$ -elliptic estimates for the linear Cauchy-Riemann operator.

For each $r \geq 0$ let $D_r := \{(x, y) \in \mathbb{R}^2 | x^2 + y^2 \leq r\}$.

Theorem 23 (Hofer[17]). *Let J be a fixed, smooth, almost complex structure on \mathbb{R}^{2d} , $d \in \mathbb{N}$. Let $C \in (0, \infty)$. Consider the set of maps $\mathcal{B}(J, C) \subset C^\infty(D_1, \mathbb{R}^{2d})$, consisting of all f satisfying,*

$$\begin{aligned} \partial_s f + J(f) \partial_t f &= 0 \\ |f(0)| &\leq 2 \\ \|\nabla f\|_{C^0(D_1)} &< C. \end{aligned}$$

Then for all $r \in (0, 1)$ there exists a sequence $c_k \in (0, \infty)$ over $k \in \mathbb{N}$, such that for all $f \in \mathcal{B}(J, C)$,

$$\|\nabla f\|_{C^k(D_r)} < c_k$$

for all $k \in \mathbb{N}$.

Replacing D_1, D_r for half disks D_1^+, D_r^+ , where $D_r^+ := \{(x, y) \in \mathbb{R}^2 | x^2 + y^2 \leq r, y \geq 0\}$, the same statement holds for maps that take the boundary points $[-1, 1] \times \{0\}$ into a smooth path of J -totally real subspaces in \mathbb{R}^{2d} .

Corollary 24. *Suppose $c_0 \in (0, \infty)$. Then there exists a sequence $c_k \in (0, \infty)$ over $k \in \mathbb{N}$, with the following property. If $\{\tilde{u}_n\}_{n \in \mathbb{N}}$ is a sequence of solutions to (23) such that $\sup_{n \in \mathbb{N}} \|\nabla \tilde{u}_n\|_{L^\infty(\{S_n, \infty\} \times \mathbb{R}/n\mathbb{Z})} \leq c_0$, then*

$$(24) \quad \|\nabla \tilde{u}_n\|_{C^k(\{S_n, \infty\} \times \mathbb{R}/n\mathbb{Z})} < c_k$$

for all $k \in \mathbb{N}$ and $n \in \mathbb{N}$.

Proof. This follows easily from the local result, theorem 23, using that the almost complex structures J_n satisfy: (1) they are invariant under the \mathbb{R} and \mathbb{Z}_n actions on $\mathbb{R} \times \mathbb{R}/n\mathbb{Z} \times D$, and (2) they each lift to the same almost complex structure J_∞ on the universal covering $\mathbb{R} \times \mathbb{R} \times D$. \blacksquare

Combining the last three statements we can prove the following.

Theorem 25. *Let $\bar{u}_n : [S_n, \infty) \times \mathbb{R} \rightarrow \mathbb{R} \times Z_\infty$ be a sequence of J_∞ -holomorphic maps over $n \in \mathbb{N}$, where each \bar{u}_n is a lift of a solution \tilde{u}_n to (23). Suppose that $\bar{u}_n(0, 0)$ is uniformly bounded in n and that $E_\lambda(\tilde{u}_n) < \infty$ for each $n \in \mathbb{N}$, and that $\lim_{n \rightarrow \infty} E_\omega(\tilde{u}_n) = 0$. Then there exists a subsequence $\{\tilde{u}_{n_j}\}_{j \in \mathbb{N}}$ such that \bar{u}_{n_j} converges in $C_{\text{loc}}^\infty(\mathbb{C}, \mathbb{R} \times Z_\infty)$ to a J_∞ -holomorphic map \tilde{u}_∞ having domain either $\Sigma = \mathbb{C}$ or $\Sigma = [S, \infty) \times \mathbb{R} \subset \mathbb{C}$ for some $S \in (-\infty, 0]$. Moreover \tilde{u}_∞ takes the following form: there exist constants $a_0, \tau_0 \in \mathbb{R}$ such that*

$$(25) \quad \begin{aligned} \tilde{u}_\infty : \Sigma &\rightarrow \mathbb{R} \times \mathbb{R} \times D \\ \tilde{u}_\infty(s, t) &= (s + a_0, t + \tau_0, \gamma(t)) \end{aligned}$$

for all $(s, t) \in \Sigma$, where $\gamma \in C^\infty(\mathbb{R}, D)$ satisfies

$$(26) \quad \dot{\gamma}(t) = X_{H_{t+\tau_0}}(\gamma(t))$$

for all $t \in \mathbb{R}$.

Proof. Taking a subsequence we may assume that $\tilde{u}_{n_j}(0, 0)$ converges, and that S_{n_j} converges to some $S \in (-\infty, 0] \cup \{-\infty\}$. From proposition 22 and corollary 24 we obtain uniform bounds on $\|\nabla \bar{u}_n\|_{C^k}$ for each $k \in \mathbb{N}$, and therefore also C^0 -bounds on \bar{u}_n on compact subsets, uniform in n . Repeated use of the Arzela-Ascoli theorem yields a subsequence converging uniformly with all derivatives on each compact subset of \mathbb{C} to a smooth map $\tilde{u}_\infty : \Sigma \rightarrow \mathbb{R} \times Z_\infty$ where $\Sigma = [S, \infty) \times \mathbb{R}$ if S is finite and $\Sigma = \mathbb{C}$ otherwise. From lemma 19 each map in the sequence \bar{u}_{n_j} takes the form

$$\bar{u}_{n_j}(s, t) = (s + a_j, t + \tau_j, z_j(s, t))$$

for constants $a_j, \tau_j \in \mathbb{R}$, with $z_j : [S_{n_j}, \infty) \times \mathbb{R} \rightarrow D$ satisfying

$$\partial_s z_j(s, t) + i \left(\partial_t z_j(s, t) - X_H(t + \tau_j, z_j(s, t)) \right) = 0$$

for all $(s, t) \in [S_{n_j}, \infty) \times \mathbb{R}$. Therefore \tilde{u}_∞ takes the form

$$\tilde{u}_\infty(s, t) = (s + a_0, t + \tau_0, z_\infty(s, t))$$

for constants $a_\infty, \tau_\infty \in \mathbb{R}$ and some $z_\infty : \Sigma \rightarrow D$ satisfying

$$\partial_s z_\infty + i \left(\partial_t z_\infty - X_H(t + \tau_\infty, z_\infty) \right) = 0.$$

Let $\omega_\infty := dx \wedge dy + d\tau \wedge dH$ on Z_∞ . Then

$$0 \leq \int_{\mathbb{R}^2} \tilde{u}_\infty^* \omega_\infty \leq \lim_{j \rightarrow \infty} E_\omega(\tilde{u}_{n_j}) = 0.$$

Thus

$$\begin{aligned} \frac{1}{2} \iint \left| \frac{\partial z_\infty}{\partial s}(s, t) \right|^2 + \left| \frac{\partial z_\infty}{\partial t}(s, t) - X_H(t + \tau_\infty, z_\infty(s, t)) \right|^2 ds dt = \\ \int_{\mathbb{R}^2} \tilde{u}_\infty^* \omega_\infty = 0. \end{aligned}$$

Hence $z_\infty(s, t) = \gamma(t)$ for some solution $\gamma : \mathbb{R} \rightarrow D$ to (26). \blacksquare

7. CONSTRUCTION OF THE FINITE ENERGY FOLIATIONS

In this final section we give a terse proof of theorem 8. The approach is along standard lines, the only part that has some small surprise is due to the presense of the boundary of the almost complex manifold. A more general construction will appear in [5].

We will assume more familiarity with terminology from [4] than elsewhere in this article, and with the homotopy invariant generalized intersection number for punctured pseudoholomorphic curves in [32].

We will prove the statement of theorem 8 for $n = 1$ only, as the proof of the general case is the same.* We begin then by recalling the statement of theorem 8 when $n = 1$.

Theorem 26. *Let $H \in C^\infty(\mathbb{R}/\mathbb{Z} \times D, \mathbb{R})$ be a Hamiltonian generating an irrational pseudo-rotation φ . Let $(Z = \mathbb{R}/\mathbb{Z} \times D, R)$ be the corresponding Hamiltonian mapping torus. Let $\gamma : \mathbb{R}/\mathbb{Z} \rightarrow Z$ be the unique 1-periodic orbit of R for which $\gamma(0) \in \{0\} \times D$. Assume H was chosen so that $\gamma(t) = (t, 0)$ for all $t \in \mathbb{R}/\mathbb{Z}$. Let $\alpha := \text{Rot}(\varphi; H) \in \mathbb{R}$, which is necessarily irrational.*

Then there exist two foliations $\mathcal{F}^+, \mathcal{F}^-$ of $\mathbb{R} \times Z$ by smoothly embedded surfaces with the following properties:

- **Cylinder leaf:** *The cylinder $C := \mathbb{R} \times \gamma(\mathbb{R}/\mathbb{Z}) \subset \mathbb{R} \times Z$ is a leaf in both \mathcal{F}^+ and \mathcal{F}^- .*
- **Pseudo-holomorphic:** *If $F \in \mathcal{F}^+$ (resp. $F \in \mathcal{F}^-$) is not C , then F is parameterized by a solution \tilde{u} to (5) with respect to J_n as in (4), with $E_\lambda(\tilde{u}) + E_\omega(\tilde{u}) < \infty$ and boundary index $[\alpha]$ (resp. $[\alpha]$).*
- **\mathbb{R} -invariance:** *If $F \in \mathcal{F}^+$ (resp. $F \in \mathcal{F}^-$) is a leaf and $c \in \mathbb{R}$, the set $F + c := \{(a + c, \tau, z) \mid (a, \tau, z) \in F\}$ is also a leaf in \mathcal{F}^+ (resp. in \mathcal{F}^-).*
- **Uniqueness:** *\mathcal{F}^+ and \mathcal{F}^- are uniquely determined by the above properties.*
- **Smooth foliation:** *\mathcal{F}^+ and \mathcal{F}^- are C^∞ -smooth foliations at each point on the complement of C .*

First we explain the construction of \mathcal{F}^+ , and then how the approach to constructing \mathcal{F}^- differs.

7.1. Construction of \mathcal{F}^+ when $\varphi|_{\partial D}$ is conjugate to a rigid rotation.

In this section suppose that there exists $g \in \text{Diff}_+^\infty(\partial D)$ so that

$$(27) \quad \sigma \circ \varphi|_{\partial D} \circ \sigma^{-1} = R_{2\pi\alpha}$$

where $R_{2\pi\alpha} : \partial D \rightarrow \partial D$ is the rigid rotation $z \mapsto ze^{2\pi i\alpha}$.

Let $H_-, H_+ \in C^\infty(Z, \mathbb{R})$ be as follows. $H_- = H$ where H is as in the statement of theorem 26. In particular the corresponding closed loop of

*For maps more general than irrational pseudo-rotations the proof for $n > 1$ is a little more involved as the almost complex structure has additional symmetry which makes transversality less obvious. But for irrational pseudo-rotations automatic transversality suffices and the same proof works for all $n \geq 1$.

Hamiltonians on the disk $H_-^t := H_-(t, \cdot) : D \rightarrow \mathbb{R}$ has time-one map φ , and $\text{Rot}(\varphi, H_-) = \alpha \in \mathbb{R}$. Define H_+ by

$$(28) \quad H_+(\tau, z) := \pi\alpha|z|^2 + C$$

for some constant $C \leq 0$ chosen so that $\max H_+ < \min H_-$. Then the two pairs of differential forms $\mathcal{H}_\pm = (\omega_\pm, \lambda_\pm)$, on Z , given by

$$\omega_\pm = dx \wedge dy + d\tau \wedge dH_\pm \quad \lambda_\pm = d\tau$$

define stable Hamiltonian structures on Z . Let R_+ and R_- denote the associated (stable Hamiltonian) Reeb vector fields, defined by $\omega_\pm(R_\pm, \cdot) = 0$ and $\lambda_\pm(R_\pm) = 1$. These are found to be

$$R_\pm(\tau, z) = \partial_\tau + X_{H_\pm^\tau}(z)$$

where $X_{H_\pm^\tau}(z)$ is the Hamiltonian vector field associated to $H_\pm^\tau = H_\pm(\tau, \cdot) : D \rightarrow \mathbb{R}$ and the symplectic form $\omega_0 = dx \wedge dy$. Choose a function $H \in C^\infty(\mathbb{R} \times Z, \mathbb{R})$ interpolating between H_- and H_+ so that

$$\begin{cases} H(a, m) = H_+(m) & \text{for all } a \geq 1 \\ \partial_a H(a, m) < 0 & \text{for } -1 < a < 1 \\ H(a, m) = H_-(m) & \text{for all } a \leq -1. \end{cases}$$

For example a good choice is $H(a, m) = \chi(a)H_+(m) + (1 - \chi(a))H_-(m)$ for some $\chi \in C^\infty(\mathbb{R}, [0, 1])$ with $\chi \equiv 0$ on $(-\infty, -1]$ and $\chi \equiv 1$ on $[1, \infty)$ and such that $\chi'(a) > 0$ for all $a \in (-1, 1)$.

Define an almost complex structure \hat{J} on $W := \mathbb{R} \times Z$ by

$$\begin{cases} \hat{J}(a, \tau, z)\partial_{\mathbb{R}} = \partial_\tau + X_{H_a^\tau}(z) \\ \hat{J}|_{TD} = i \end{cases}$$

where for $(a, \tau) \in \mathbb{R} \times \mathbb{R}/\mathbb{Z}$, $X_{H_a^\tau}$ is the Hamiltonian vector field on $(D, \omega_0 = dx \wedge dy)$ for the Hamiltonian function $H_a^\tau := H(a, \tau, \cdot) : D \rightarrow \mathbb{R}$. Then (W, \hat{J}) is an almost complex manifold with cylindrical ends $E_+ = [1, \infty) \times Z$ and $E_- = (-\infty, -1] \times Z$, adjusted to the stable Hamiltonian structures $\mathcal{H}_\pm = (\omega_\pm, \lambda_\pm)$ on these ends, while on the region $(-1, 1) \times Z$, \hat{J} tames the symplectic 2-form

$$\Omega := dx \wedge dy + d\tau \wedge dH.$$

Moreover, the cylinder $C := \mathbb{R} \times \gamma(\mathbb{R}/\mathbb{Z})$ is a \hat{J} -holomorphic curve, even though \hat{J} is not everywhere \mathbb{R} -invariant. This is because the unique 1-periodic orbits of R_+ and R_- are the same as parameterized closed loops, and the expression for H in terms of H_-, H_+, χ .

Finally, we will show in section 7.6 that the boundary of W , that is $\mathbb{R} \times \partial Z$, is filled by a set \mathcal{S} of immersed \hat{J} -holomorphic planes which, in the ends $E_\pm \cap (\mathbb{R} \times \partial Z)$ coincide with the product of the \mathbb{R} -component and a Reeb trajectory of R_\pm .

We proceed to construct \mathcal{F}^+ in four steps.

Step 1: The almost complex structure \hat{J} on W satisfies $\hat{J}|_{E_+} = J_+|_{E_+}$ where J_+ is the cylindrical almost complex structure

$$(29) \quad \begin{cases} J_+ \partial_{\mathbb{R}} = \partial_{\tau} + 2\pi\alpha \partial_{\theta} \\ J_+|_{TD} = i \end{cases}$$

on $\mathbb{R} \times Z$, in standard polar coordinates (r, θ) on the disk. For each $c \in \mathbb{R}$ and $z \in \partial D$, the map

$$(30) \quad \begin{aligned} \tilde{u}_{c,z} : \mathbb{R}^+ \times \mathbb{R}/\mathbb{Z} &\rightarrow \mathbb{R} \times Z \\ \tilde{u}_{c,z}(s, t) &= (s + c, t, ze^{2\pi([\alpha]-\alpha)s} e^{2\pi i[\alpha]t}) \end{aligned}$$

is J_+ -holomorphic. The combined images of these maps along with the cylinder $C := \{(a, \tau, 0, 0) \mid a \in \mathbb{R}, \tau \in \mathbb{R}/\mathbb{Z}\}$ defines an \mathbb{R} -invariant finite energy foliation for $(\mathbb{R} \times Z, J_+)$ with boundary index $[\alpha]$. This is the model foliation from which we homotope.

Step 2: We return to the manifold $(W = \mathbb{R} \times Z, \hat{J})$ with cylindrical ends. Let \mathcal{M} denote the moduli space of all finite energy \hat{J} -holomorphic curves $F \subset W$ which admit a \hat{J} -holomorphic parameterization by a map $\tilde{u} = (a, \tau, z) \in C^\infty(\mathbb{R}^+ \times \mathbb{R}/\mathbb{Z}, \mathbb{R} \times Z)$ satisfying

$$\begin{cases} \tilde{u}(0, \cdot) \in L_c \text{ and } \tilde{u} \text{ meets } \mathbb{R} \times \partial Z \text{ transversely} \\ \tau(0, \cdot) : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}/\mathbb{Z} \text{ has degree } +1 \\ z(0, \cdot) : \mathbb{R}/\mathbb{Z} \rightarrow \partial D \text{ has degree } [\alpha] \end{cases}$$

for any $c \in \mathbb{R}$, and equip \mathcal{M} with the topology coming from $C_{\text{loc}}^\infty \cap C^0([0, \infty) \times \mathbb{R}/\mathbb{Z}, W)$ convergence. \mathcal{M} is non-empty as it contains the image of each curve $\tilde{u}_{c,z}$ from (30) provided we choose $c \geq 1$, as $\hat{J} = J_+$ on the positive end $E_+ = [1, \infty) \times Z$.

Each curve $F \in \mathcal{M}$ is embedded, and any two curves $F_1, F_2 \in \mathcal{M}$ are equal or disjoint. Indeed, any two curves in \mathcal{M} are homotopic through half cylinders with so called asymptotically cylindrical ends (in the sense of [32]). It therefore suffices to know that there exists a single curve $F_0 \in \mathcal{M}$ satisfying

$$F_0 \cdot F_0 = 0 \quad \text{and} \quad F_0 \cdot C = 0$$

where C is the cylinder above. This is easily verified for the explicit curves in (30). Then homotopy invariance of \cdot implies that for any $F \in \mathcal{M}$ we have $F \cdot F = 0$ and $F \cdot C = 0$. These imply F is embedded by the adjunction formula in [32]. Finally, for any two curves $F_1, F_2 \in \mathcal{M}$ we have $F_1 \cdot F_2 = F_1 \cdot F_1 = 0$ and so either $F_1 = F_2$ or $F_1 \cap F_2 = \emptyset$.

As a solution to the Cauchy-Riemann equations with “free” boundary conditions (meaning that $c \in \mathbb{R}$ above is not fixed) each curve in \mathcal{M} is Fredholm with index 2. Standard automatic transversality arguments apply as for example in [22, ?] and so each curve is Fredholm regular despite the non-generic choice of almost complex structure.

Lemma 27. *The set $\mathcal{E} := \{w \in W \mid w \in F \text{ for some } F \in \mathcal{M}\}$ is an open and closed subset of $W \setminus C$.*

Proof. Openness: as each curve F in \mathcal{M} is regular and transverse to the boundary of W , an implicit function theorem as in [22] applies (or more precisely a version in [33]), and all curves in \mathcal{M} sufficiently close to F correspond to sections of the kernel of the linearized Cauchy-Riemann operator at F . Non-trivial elements of this kernel have no zeros. This is enough to show that there is an open neighborhood of F filled by curves coming from the implicit function theorem which by construction lie in \mathcal{M} .

Closedness: there is a uniform bound on the total energy of all curves in \mathcal{M} . Thus for any sequence $F_k \in \mathcal{M}$, see the discussion above, symplectic field theory compactness as in [4] applies. We have curves with boundary, which is not treated by [4], but this is not a problem due to the filling \mathcal{S} . Suppose that $p_k \in \mathbb{R} \times Z$ is a sequence of points in \mathcal{E} , with $p_k \rightarrow p_\infty$ some point $p_\infty \notin C$. Let $F_k \in \mathcal{M}$ be a curve containing p_k . Viewing these as marked points on the curves F_k we take a limit in the SFT sense. The unique stable limiting building \overline{F} has a component F_∞ with a marked point, this component must contain p_∞ . It is easy to see that the component F_∞ must be a half cylinder in \mathcal{M} provided it is not the cylinder C (see section 7.7 for a brief justification of this). But F_∞ cannot equal C as $p_\infty \notin C$. ■

We conclude that there is a unique curve in \mathcal{M} going through each point in $W \setminus C$.

Step 3: Let $p_k \in W \setminus C$ be any sequence of points which converge to a point $p_\infty \in C$ on the distinguished cylinder. From the previous step there exists then a sequence of curves $F_k \in \mathcal{M}$ with $p_k \in F_k$. Applying the SFT compactness theorem to this sequence, where p_k is viewed as the image of a marked point of F_k , we get convergence to a stable nodal holomorphic building \overline{F} say, with a single marked point which corresponds to p_∞ and which lies on the distinguished middle level of \overline{F} . It follows that the middle level of \overline{F} is simply the cylinder C as p_∞ cannot be an isolated intersection point with C . Therefore the building \overline{F} must have non-empty lower levels. These lower levels are holomorphic curves in the cylindrical manifold $(\mathbb{R} \times Z, J_-)$ corresponding to the negative end of (W, \hat{J}) , which was modelled on the pseudo-rotation φ . As the building is stable and has no marked points in the lower levels, there is precisely one lower level in \overline{F} and it must correspond to a half cylinder with positive puncture asymptotic to the unique simply covered periodic orbit γ_- of the Reeb flow of \mathcal{H}_- .

Denote this half cylinder by G_0 . Intersection considerations show that since $C \cdot F_k = 0$ and $F_k \cdot F_k = 0$ for all k , and these curves only break at elliptic orbits, we must have

$$G_0 \cdot G_0 = 0 \quad \text{and} \quad G_0 \cdot C_- = 0$$

where C_- is the unique orbit cylinder in $(\mathbb{R} \times Z, J_-)$. It follows as in step 2 that the half cylinder G_0 is embedded.

Step 4: The embedded half cylinder G_0 in the cylindrical manifold $(\mathbb{R} \times Z, J_-)$ found at the end of the last step is automatically Fredholm regular and has Fredholm index 2. We may now argue in the manner of step 2, and conclude G_0 lies in a non-empty 2-dimensional moduli space of half cylinders \mathcal{M}_- which fill an open and closed subset of $(\mathbb{R} \times Z) \setminus C_-$. Each $G \in \mathcal{M}_-$ is homotopic to G_0 through curves with asymptotically cylindrical ends, and we can conclude that

$$G \cdot G = 0 \quad \text{and} \quad G \cdot C_- = 0$$

holds for all $G \in \mathcal{M}_-$. This implies that:

- (1) Each $G \in \mathcal{M}_-$ is embedded.
- (2) Each pair of curves $G_1, G_2 \in \mathcal{M}_-$ is either equal or disjoint.

The second of these also implies that if $G \in \mathcal{M}_-$ and $c \in \mathbb{R} \setminus \{0\}$, then the curve translated in the \mathbb{R} -direction $G + c := \{(a + c, m) \mid (a, m) \in G\}$ (which is in \mathcal{M}_- as J_- is \mathbb{R} -invariant) is disjoint from G because G and $G + c$ cannot be equal as their boundaries are in different totally real submanifolds.

7.2. Construction of \mathcal{F}^+ without boundary restrictions. This completes the construction for any irrational pseudo-rotation $\varphi : D \rightarrow D$ which restricts to a circle diffeomorphism $\varphi|_{\partial D}$ that is smoothly conjugate to a rigid rotation. By a deep result of Herman the set of such pseudo-rotations is dense amongst all irrational pseudo-rotations, and so a further limiting step can remove the boundary restrictions. We explain this now.

Say that an irrational number α belongs to \mathcal{D} if it satisfies the following Diophantine condition: there exists $n \geq 2$ and $C \in (0, \infty)$ such that for all $(p, q) \in \mathbb{Z} \times \mathbb{N}$

$$\left| \alpha - \frac{p}{q} \right| \geq C \frac{1}{q^n}.$$

It is easy to show that \mathcal{D} is dense in \mathbb{R} (indeed of full measure). Denote by $\text{Diff}_{\mathcal{D}}(\partial D)$ those orientation preserving C^∞ -smooth circle diffeomorphisms that have rotation number in \mathcal{D} .

Theorem 28 (Herman [16]). *If $f \in \text{Diff}_{\mathcal{D}}(\partial D)$ then there exists $\sigma \in \text{Diff}_+^\infty(\partial D)$ such that $\sigma^{-1}f\sigma = R$ where $R : \partial D \rightarrow \partial D$ is a rigid rotation.*

Lemma 29. *$\text{Diff}_{\mathcal{D}}(\partial D)$ is dense in $\text{Diff}_+^\infty(\partial D)$ with the C^∞ -topology.*

Proof. Fix any $f \in \text{Diff}_{\mathcal{D}}(\partial D)$. Consider the continuous path $f_t := R_{2\pi t} \circ f \in \text{Diff}_+^\infty(\partial D)$ over $t \in [0, 1/2]$. The rotation numbers $\text{Rot}(f_t)$ vary continuously with t . Moreover, there is a continuous family of lifts $\tilde{f}_t : \mathbb{R} \rightarrow \mathbb{R}$ with the monotonicity property that for $t > 0$, $\tilde{f}_t(x) > \tilde{f}_0(x)$ for all $x \in \mathbb{R}$. Therefore, since $\text{Rot}(f_0)$ is *irrational*, it follows that for all $t \in (0, 1/2]$, $\text{Rot}(f_t) > \text{Rot}(f_0)$. See for example proposition 11.1.9 in [27]. As \mathcal{D} is dense in \mathbb{R}/\mathbb{Z} , we find a sequence $t_j \in (0, 1/2]$ converging to zero such that $\text{Rot}(f_{t_j}) \in \mathcal{D}$. ■

Now suppose that $\varphi \in \text{Diff}^\infty(D, \omega_0)$ is any smooth irrational pseudo-rotation. Let $H \in C^\infty(\mathbb{R}/\mathbb{Z} \times D, \mathbb{R})$ be a Hamiltonian with time-one map φ . Using the lemma one can find smooth perturbations $H_j \in C^\infty(\mathbb{R}/\mathbb{Z} \times D, \mathbb{R})$ of H near the boundary $\mathbb{R}/\mathbb{Z} \times \partial D$, so that

$$\begin{aligned} H_j &\rightarrow H \text{ in } C^\infty \\ \text{Rot}(\varphi_j|_{\partial D}) &\in \mathcal{D} \end{aligned}$$

for all j , where φ_j is the time-one map of H_j . For j sufficiently large φ_j has no fixed points besides the origin. (It may not be a pseudo-rotation but that does not matter.) By Herman's theorem $\varphi_j|_{\partial D}$ is smoothly conjugate to a rigid rotation. Therefore, by steps 2 to 4 in section 7.1 we can find a finite energy foliation \mathcal{F}_{H_j} of the cylindrical almost complex manifold $(\mathbb{R} \times Z, J_{H_j})$.

The almost complex structures J_{H_j} converge uniformly to J_H as $j \rightarrow \infty$, and we may pass to a limit of foliations and obtain a finite energy foliation for $(\mathbb{R} \times Z, J_H)$. For example one can take a limit of a single sequence of half cylinder leaves $F_j \in \mathcal{F}_{H_j}$ and obtain a single J_H -holomorphic half cylinder F_∞ disjoint from the orbit cylinder C and with vanishing self intersection number. The moduli space containing F_∞ fills up the complement of C by embedded curves that are pairwise identical or disjoint as we require. This completes the construction of \mathcal{F}^+ in theorem 26.

7.3. Uniqueness. That the foliation \mathcal{F}^+ is uniquely determined by the other properties in theorem 26 is more or less immediate from the intersection theory because there is only one 1-periodic orbit and all curves with boundary have the same boundary index.

7.4. Smoothness. The proof that \mathcal{F}^+ is a smooth foliation on $(\mathbb{R} \times Z) \setminus C$ is the same as that used in Hofer-Wysocki-Zehnder [23], based on their implicit function theorem in [22].

By smoothness of \mathcal{F}^+ at a point $p \in \mathbb{R} \times Z$ we mean that there exists a smooth local diffeomorphism $\sigma : U \rightarrow V \subset \mathbb{R}^2 \times \mathbb{R}^2$ on a neighborhood of p onto an open convex subset $V \subset \mathbb{R}^4$, such that each leaf $F \in \mathcal{F}^+$ having non-empty intersection with U is mapped under σ onto a subset of the form $V \cap (\mathbb{R}^2 \times \{c\})$ for some $c \in \mathbb{R}^2$.

The point is that each leaf $F \in \mathcal{F}^+$, with $F \neq C$, is embedded, meets the boundary of $\mathbb{R} \times Z$ transversally in some totally real submanifold, has Fredholm index 2, and is Fredholm regular. An implicit function theorem as generalized in [33] applied to F gives a neighborhood of F in $(\mathbb{R} \times Z, J)$ filled by J -holomorphic curves, given by exponentiating sections of a normal bundle over F . (With respect to a metric for which each L_c , and $\mathbb{R} \times \partial Z$, are totally geodesic.) These curves from the implicit function theorem form a C^∞ -smooth local foliation $\tilde{\mathcal{F}}$ of a neighborhood of F in $\mathbb{R} \times Z$. Intersection theory considerations show that each leaf in $\tilde{\mathcal{F}}$ is also a leaf in the global foliation \mathcal{F}^+ . So \mathcal{F}^+ is also smooth at points on F .

7.5. Constructing \mathcal{F}^- . The construction of \mathcal{F}^- is along exactly the same lines as for \mathcal{F}^+ , but we start with a different model foliation in step 1. Indeed, in place of the curves in (30), each of which is bounded from below, we use curves of the form

$$(31) \quad \begin{aligned} v_{c,z} : \mathbb{R}^- \times \mathbb{R}/\mathbb{Z} &\rightarrow \mathbb{R} \times Z \\ v_{c,z}(s, t) &= (s + c, t, ze^{2\pi([\alpha] - \alpha)s} e^{2\pi i[\alpha]t}) \end{aligned}$$

over $c \in \mathbb{R}$ and $z \in \partial D$. (These maps are also pseudoholomorphic with respect to the almost complex structure in (29).) Each $v_{c,z}$ has image bounded from above, so we have to insert one of them into the negative end of our almost complex manifold ($W = \mathbb{R} \times Z, \hat{J}$). So the main difference is that from the start we reverse the roles of H_- and H_+ , this time picking a positive constant C in (28) so that $\min H_- > \max H_+$ still holds. Then the remaining steps are exactly analogous, and in the final foliation the curves with boundary have boundary index $[\alpha]$ instead, as the curves in (31) do.

7.6. Foliating the boundary. We need to justify the existence of the foliation \mathcal{S} of the boundary of $\mathbb{R} \times Z$ that was used in section 7.1.

For each $a \in \mathbb{R}$, consider for a fixed value of a the resulting time-dependent Hamiltonian on the disk H_a given by

$$H_a^t := H(a, t, \cdot) : D \rightarrow \mathbb{R}$$

over $t \in \mathbb{R}/\mathbb{Z}$. By modifying H_a on any arbitrarily small neighborhood of the boundary of the disk we may arrange that the time-one map of the path of generated Hamiltonian disk maps is any prescribed orientation preserving diffeomorphism on the boundary of the disk. By extension, given any smooth path $a \mapsto f_a \in \text{Diff}_+^\infty(\partial D)$ over $a \in \mathbb{R}$, satisfying

$$(32) \quad f_a = \begin{cases} R_{2\pi\alpha}|_{\partial D} & \text{if } a \geq 1 \\ \varphi|_{\partial D} & \text{if } a \leq -1, \end{cases}$$

the function $H : \mathbb{R} \times Z \rightarrow \mathbb{R}$ may be modified on any small neighborhood of $[-1, 1] \times \partial Z$ so that for each $a \in \mathbb{R}$ the time-one map of the modified Hamiltonian H_a now coincides with f_a on the boundary of the disk.

Suppose that we can find a smooth path $a \mapsto f_a \in \text{Diff}_+^\infty(\partial D)$ satisfying (32), and which additionally has the property that each f_a is smoothly conjugate to the rigid rotation $R_{2\pi\alpha}|_{\partial D}$. More precisely, suppose that we find a smooth map $g \in C^\infty(\mathbb{R} \times \partial D, \partial D)$, so that for each $a \in \mathbb{R}$ the map $g_a := g(a, \cdot)$ is an element of $\text{Diff}_+^\infty(\partial D)$, and with the property that the path $f_a := g_a R_{2\pi\alpha} g_a^{-1}$ satisfies (32). Then we may modify $H \in C^\infty(\mathbb{R} \times Z, \mathbb{R})$ near $[-1, 1] \times \partial Z$ so that for each $a \in \mathbb{R}$ the time-one map of the time-dependent Hamiltonian $H_a := H(a, \cdot)$ equals f_a on the boundary of the disk. For each $a \in \mathbb{R}$ let

$$\begin{aligned} \phi_a : \mathbb{R} \times \partial Z &\rightarrow \partial Z \\ (t, z) &\mapsto \phi_a^t(z) \end{aligned}$$

denote the 1-parameter family of maps generated by $X_{H_{a,t}}$ on ∂D . So in particular $\phi_a^1 = f_a$ for all $a \in \mathbb{R}$. Now for each $z \in \partial D$,

$$S_z := \left\{ (a, t, g_a(\phi_a^t(z))) \in \mathbb{R} \times \mathbb{R}/\mathbb{Z} \times \partial D \mid a \in \mathbb{R}, t \in \mathbb{R} \right\}$$

is an immersed surface in $\mathbb{R} \times \partial Z$, and the union

$$\mathcal{S} := \bigcup_{z \in \partial D} S_z$$

is a foliation of $\mathbb{R} \times \partial Z$. As α is irrational each S_z is dense in $\mathbb{R} \times \partial Z$. However, the relation

$$(33) \quad f_a := g_a R_{2\pi\alpha} g_a^{-1}$$

for all $a \in \mathbb{R}$ enables us to find a C^∞ -smooth almost complex structure J' on $\mathbb{R} \times Z$, prescribed at points on $\mathbb{R} \times \partial Z$ so that each S_z has J' -invariant tangent bundle. Indeed, differentiating the expression $(a, t, g_a(\phi_a^t(z)))$ in a gives a vector field V_1 say, while differentiating it in t results in a vector field V_2 . Both are non-vanishing and transverse as we will see, so we can set $J'V_1 = V_2$. That V_1 is indeed a well defined vector field uses (33). Moreover, one finds that:

$$V_2 = \partial_\tau + X_{H_a^\tau},$$

while

$$V_1 = \partial_a + V_3,$$

for some V_3 on $\mathbb{R} \times \partial Z$ that has no ∂_a component, and tends to zero in C^0 as $\|\partial_a f_a\|_{C^0}$ tends to zero. We can arrange that $\|\partial_a f_a\|_{C^0}$ is as small as we wish by “slowing everything down”, that is, replacing the interval $[-1, 1] \times Z$ by $[-N, N] \times Z$ for sufficiently large $N > 0$. Then, from these expressions for V_1, V_2 we see that J' extends to an almost complex structure J'' on $\mathbb{R} \times Z$ with the following properties if $\|\partial_a f_a\|_{C^0}$ is sufficiently small:

- (1) J'' coincides with the almost complex structure J outside of a small neighborhood of $[-N + 1, N - 1] \times \partial Z$.
- (2) Each surface $L_c := \{c\} \times \partial Z$ is totally real with respect to J'' .
- (3) J'' is tamed by the symplectic form Ω on $(-N, N) \times Z$.

And finally of course \mathcal{S} is a J'' -holomorphic filling of the boundary $\mathbb{R} \times \partial Z$.

The only remaining question is when the relation (33) can be arranged for all $a \in \mathbb{R}$. But this holds if and only if the circle maps $\varphi|_{\partial D}$ and $R_{2\pi\alpha}|_{\partial D}$ are conjugate by an orientation preserving C^∞ -smooth diffeomorphism. Necessity is obvious, let us show sufficiency. Suppose that there exists $g \in \text{Diff}_+^\infty(\partial D)$ such that $\varphi|_{\partial D} = gR_{2\pi\alpha}|_{\partial D}g^{-1}$. Since g has degree +1 it is smoothly isotopic to the identity and we may find a smooth isotopy $g_a \in \text{Diff}_+^\infty(\partial D)$ over $a \in \mathbb{R}$, satisfying $g_a = \text{id}$ for all $a \geq 1$, and $g_a = g$ for all $a \leq -1$. Thus the smooth path $f_a \in \text{Diff}_+^\infty(\partial D)$ over $a \in \mathbb{R}$ defined by $f_a := g_a R_{2\pi\alpha} g_a^{-1}$ satisfies

$$f_a = \begin{cases} R_{2\pi\alpha} & \text{if } a \geq 1 \\ \varphi|_{\partial D} & \text{if } a \leq -1 \end{cases}$$

and therefore has the properties we require.

7.7. Compactness. In section 7.1 we applied, a number of times, the compactness theory in [4] that was developed for symplectic field theory, to sequences of punctured pseudoholomorphic curves in $(W = \mathbb{R} \times Z, \hat{J})$.

Although $\partial W = \mathbb{R} \times \partial Z$ is non-empty, we arranged in the last section that it can be foliated by immersed \hat{J} -holomorphic curves without boundary, in a reasonably nice way (meaning that they lift to properly embedded curves on the universal covering of ∂W).

Consider a sequence F_k of \hat{J} -holomorphic curves in (W, \hat{J}) which are half cylinders with totally real boundary conditions $\partial F_k \subset L_{c_k}$ for some $c_k \in \mathbb{R}$. Suppose that for each k the generalized intersection number, in the sense of [32], with the \hat{J} -holomorphic cylinder C is zero; $F_k \cdot C = 0$ for all k . Furthermore, suppose that each F_k meets the boundary of W transversely. Then we used the following several times:

Proposition 30. *If each F_k has at most one marked point, and the sequence F_k converges to a stable nodal holomorphic building \overline{F} in the sense of [4], then \overline{F} has the following form:*

- *At most one component in each level.*
- *No nodal points or closed curves.*
- *One component with boundary, which is a half cylinder (meaning a disk with an interior puncture) with the same sign puncture as that of each F_k , k large, and meeting ∂W transversally along the boundary of the curve.*
- *All other components of \overline{F} , if there are any, are cylinders with punctures of opposite sign.*
- *Interior points of \overline{F} are disjoint from ∂W .*

Proof. \overline{F} has genus zero in the generalized sense of [4] as each F_k has genus zero. So each component of \overline{F} has genus zero also.

There are no planes as there are no contractible periodic orbits.

There are also no disk components in \overline{F} . To see this we argue indirectly. Suppose that \overline{F} has a disk component D . Then ∂D must be a closed loop in one of the totally real surfaces $L_c \simeq \partial Z$ say. As ∂D is contractible in W it must lie in a homology class $m1_{\partial D} \in H_1(\partial Z)$ for some $m \in \mathbb{Z}$. The topological count of intersections between D and the cylinder C is m . The building is stable with at most one marked point implies that the component D is not just a point. Therefore it has non-zero energy, which implies that $m \neq 0$. Therefore D and C are distinct holomorphic curves with interior intersections. Therefore F_k and C intersect for large k , contradicting that they are infact disjoint for all k .

There are no closed curves in \overline{F} as Ω is exact. Also there are no nodes, as these would lead to a component asymptotic to a contractible periodic orbit which is impossible.

It follows that the limit building has one boundary component, and so has a unique component with non-empty boundary, and that this is a half cylinder. Due to the orientations of the periodic orbits in the ends this half cylinder cannot have a different sign puncture to each F_k .

The presence of the foliation \mathcal{S} of ∂W prevents interior intersections between components of \overline{F} and ∂W , and prevents the boundary of \overline{F} becoming tangent anywhere to ∂W . \blacksquare

APPENDIX A. PROOF OF PROPOSITION 2

In this appendix we prove the following statement, which implies proposition 2. The idea of the proof was explained to be by Patrice LeCalvez.

We write $R_\theta : D \rightarrow D$ to denote the rigid rotation $z \mapsto e^{i\theta}z$ through angle $\theta \in \mathbb{R}$.

Proposition 31. *Consider a sequence $\varphi_k \in \text{Homeo}_+(D)$ converging in the C^0 -topology to $\varphi \in \text{Homeo}_+(D)$, where all maps fix $0 \in D$. Under the following additional assumptions it follows that φ has no periodic points in $D \setminus \{0\}$.*

- (1) *For each $k \in \mathbb{N}$, there exists $g_k \in \text{Homeo}_+(D)$, fixing the origin, such that $\varphi_k = g_k^{-1} R_{2\pi\theta_k} g_k$, some $\theta_k \in \mathbb{R}$.*
- (2) *$\theta_k \rightarrow \theta$ as $k \rightarrow \infty$, where θ is irrational.*

To prove this we need to recall the notions of positively and negatively returning disks due to Franks [13]. Let $\mathring{D} := D \setminus \partial D$. Denote by $A := \mathring{D} \setminus \{0\}$ and $\tilde{\mathbb{A}} := (0, 1) \times \mathbb{R}$ the open annulus and its universal covering via the covering map

$$\begin{aligned} \pi : (0, 1) \times \mathbb{R} &\rightarrow \mathring{D} \setminus \{0\} \\ (x, y) &\mapsto xe^{2\pi iy}. \end{aligned}$$

Let $T : \tilde{\mathbb{A}} \rightarrow \tilde{\mathbb{A}}$ be the deck transformation $T(x, y) = (x, y + 1)$. By an open disk $U \subset \tilde{\mathbb{A}}$ is meant an open subset homeomorphic to \mathring{D} with the subspace topology.

Definition 32. Let $f : \mathbb{A} \rightarrow \mathbb{A}$ be a homeomorphism homotopic to the identity, and $\tilde{f} : \tilde{\mathbb{A}} \rightarrow \tilde{\mathbb{A}}$ a lift via π . Consider an open disk $U \subset \tilde{\mathbb{A}}$ for which

$$(34) \quad \tilde{f}(U) \cap U = \emptyset.$$

If there exist integers $n > 0$, $k \neq 0$, such that

$$(35) \quad \tilde{f}^n(U) \cap T^k U \neq \emptyset$$

then U is called a *positively*, resp. *negatively*, *returning disk* for \tilde{f} if $k > 0$, resp. $k < 0$.

Remark 8. Consider a disk $U \subset \tilde{\mathbb{A}}$ satisfying (34) and (35). If moreover the closure of the disk satisfies (34), that is, $\tilde{f}(\bar{U}) \cap \bar{U} = \emptyset$, then for any

sufficiently C^0 -small perturbation of \tilde{f} , U will satisfy (34) and (35) for the perturbed map also.

The key result for us is the following strong generalization of the Poincaré-Birkhoff fixed point theorem, theorem 2.1 in [13].

Theorem 33 (Franks). *Let $f : \mathbb{A} \rightarrow \mathbb{A}$ be a homeomorphism of the open annulus homotopic to the identity, and for which every point is non-wandering. If there exists a lift $\tilde{f} : \tilde{\mathbb{A}} \rightarrow \tilde{\mathbb{A}}$ having a positively returning disk which is a lift of a disk in \mathbb{A} , and a negatively returning disk that is a lift of a disk in \mathbb{A} , then f has a fixed point.*

Recall that a point $x \in \mathbb{A}$ is *non-wandering* for f if for every open neighborhood U of x there exists $n > 0$ such that $f^n(U) \cap U \neq \emptyset$. It is easy to prove that every point is non-wandering for a homeomorphism f if f preserves a finite measure that is positive on open sets.

We now use theorem 33 to prove the proposition.

Proof of proposition 31. Arguing indirectly, suppose that φ has a periodic point $z_0 \in D \setminus \{0\}$. Then we will show that some iterate of φ , let us call it φ^n , has a lift via the covering map

$$\begin{aligned} \pi : (0, 1] \times \mathbb{R} &\rightarrow D \setminus \{0\} \\ (x, y) &\mapsto xe^{2\pi iy} \end{aligned}$$

to a map of the half closed infinite strip $\tilde{\varphi}^n : (0, 1] \times \mathbb{R} \rightarrow (0, 1] \times \mathbb{R}$, having disks $U_-, U_+ \subset (0, 1) \times \mathbb{R}$ which satisfy the following:

- (1) U_+ is a positively returning disk for $\tilde{\varphi}^n$. U_- is a negatively returning disk for $\tilde{\varphi}^n$.
- (2) The closures satisfy $\tilde{f}(\bar{U}_+) \cap \bar{U}_+ = \emptyset$ and $\tilde{f}(\bar{U}_-) \cap \bar{U}_- = \emptyset$.
- (3) U_+ and U_- are lifts of disks in $\mathring{D} \setminus \{0\}$.

Suppose for a moment that we have established this. Then since $\varphi_k : D \rightarrow D$ converges uniformly in the C^0 -topology to φ , there exists a sequence of lifts

$$\tilde{\varphi}_k : (0, 1] \times \mathbb{R} \rightarrow (0, 1] \times \mathbb{R}$$

of φ_k which converges uniformly in the C^0 -topology to $\tilde{\varphi}$. Hence by remark 8 there exists $K \in \mathbb{N}$ such that for all $k \geq K$, U_+ is a positively returning disk for $\tilde{\varphi}_k^n$, and U_- is a negatively returning disk for $\tilde{\varphi}_k^n$.

Recall that U_+ and U_- lie in the interior $(0, 1) \times \mathbb{R}$, and both are lifts of disks in $\mathring{D} \setminus \{0\}$. Therefore we may apply Franks' theorem, theorem 33, to each map of the open annulus $\varphi_k^n = (\varphi_k)^n : \mathring{D} \setminus \{0\} \rightarrow \mathring{D} \setminus \{0\}$ for $k \geq K$. Indeed, being conjugate to a rigid rotation, every point is non-wandering for φ_k^n . We conclude that for all $k \geq K$, φ_k^n has a fixed point in $D \setminus \{0\}$. But φ_k^n is conjugate to the rigid rotation $R_{2\pi n\theta_k}$. So $n\theta_k \in \mathbb{Z}$ for all $k \geq K$, which contradicts the convergence of the sequence θ_k to an irrational number.

With this contradiction we will be done, and so it remains to establish that some iterate of φ has a lift for which we can find disks $U_+, U_- \subset (0, 1) \times \mathbb{R}$ satisfying conditions (1), (2), and (3) listed above.

We are assuming that φ has a periodic point besides $0 \in D$. From this it is not hard to see that for some $n > 0$ sufficiently large there exists a lift $\tilde{\varphi}^n : (0, 1] \times \mathbb{R} \rightarrow (0, 1] \times \mathbb{R}$ of φ^n such that one of the following two possibilities occurs: (1) there is a fixed point $z \in \mathring{D} \setminus \{0\}$ of φ^n with a lift $(x, y) \in (0, 1) \times \mathbb{R}$, such that $\tilde{\varphi}^n(x, y) = (x, y - 1)$, and for all $y \in \mathbb{R}$, $\tilde{\varphi}^n(1, y) = (1, y')$ where $y' > y + 1$. Or (2), there is a fixed point $z \in \mathring{D} \setminus \{0\}$ of φ^n with a lift $(x, y) \in (0, 1) \times \mathbb{R}$, such that $\tilde{\varphi}^n(x, y) = (x, y + 1)$, and for all $y \in \mathbb{R}$, $\tilde{\varphi}^n(1, y) = (1, y')$ where $y' < y - 1$.

Let us consider the first situation, as the second is dealt with similarly. Any sufficiently small disk neighborhood $U_- \subset (0, 1) \times \mathbb{R}$ of (x, y) , satisfies $\tilde{\varphi}^n(\bar{U}_-) \cap \bar{U}_- = \emptyset$ and is a lift of a disk in $\mathring{D} \setminus \{0\}$. Moreover $\tilde{\varphi}^n(U_-) \cap T^{-1}(U_-)$ is non-empty as it contains the point $(x, y - 1)$. Thus U_- is a negatively returning disk for $\tilde{\varphi}^n$ satisfying conditions (1), (2), and (3) above.

It remains to find a suitable positively returning disk for $\tilde{\varphi}^n$. Recall that for all $y \in \mathbb{R}$,

$$\tilde{\varphi}^n(1, y) = (1, y')$$

where $y' > y + 1$. As ∂D is a compact invariant set for φ^n , there exists a point $z_1 \in \partial D$ that is non-wandering for φ^n (e.g. take any point in the ω -limit set of an orbit). Let $(1, y) \in (0, 1] \times \mathbb{R}$ be a lift of z_1 . Then for every sufficiently small closed neighborhood $I \subset \partial \tilde{\mathbb{A}} := \{1\} \times \mathbb{R}$ of $(1, y)$ there exists $m > 0$ and $k > 0$ such that

$$(36) \quad \tilde{\varphi}^n(I) \cap I = \emptyset$$

$$(37) \quad \tilde{\varphi}^{nm}(I) \cap T^k(I) \neq \emptyset.$$

Taking I sufficiently small we may also assume that it is a lift of an interval in ∂D , that is, a closed, connected, non-empty, simply connected set.

Consider the open neighborhoods of I in $\tilde{\mathbb{A}}$ of the form

$$V := \{z \in (0, 1] \times \mathbb{R} \mid d(z, I) < \varepsilon\}$$

for $\varepsilon > 0$. Clearly,

$$(38) \quad \tilde{\varphi}^{nm}(V) \cap T^k(V) \neq \emptyset$$

from (37) and is open in $(0, 1] \times \mathbb{R}$. And for all $\varepsilon > 0$ sufficiently small we have

$$(39) \quad \tilde{\varphi}^n(\bar{V}) \cap \bar{V} = \emptyset$$

from (36). Set $U_+ := V \cap \tilde{\mathbb{A}} = V \cap ((0, 1) \times \mathbb{R})$. From (38) we get

$$\tilde{\varphi}^{nm}(U_+) \cap T^k(U_+) \neq \emptyset.$$

From (39)

$$\tilde{\varphi}^n(\bar{U}_+) \cap \bar{U}_+ = \emptyset$$

as $\bar{U}_+ = \bar{V}$. Moreover, for $\varepsilon > 0$ sufficiently small U_+ is a disk in $\tilde{\mathbb{A}}$ and is a lift of a disk in $\mathring{D} \setminus \{0\}$. Thus U_+ is a positively returning disk for $\tilde{\varphi}^n$ satisfying the required conditions (1), (2), and (3). \blacksquare

APPENDIX B. NONDEGENERACY

Let $\varphi : D \rightarrow D$ be an irrational pseudo-rotation with rotation number $[\alpha] \in \mathbb{R}/\mathbb{Z}$. In this appendix we prove the following.

Lemma 34. *The linearization $D\varphi(0)$ has eigenvalues $\{e^{2\pi i\alpha}, e^{-2\pi i\alpha}\}$.*

We begin with:

Lemma 35. *Let $\pi : (0, 1] \times \partial D \rightarrow D \setminus \{0\}$ be the C^∞ -diffeomorphism $\pi(t, x) := \sqrt{t}x$. Then the diffeomorphism $\pi^{-1} \circ \varphi \circ \pi$ on the half open annulus $(0, 1] \times \partial D$ preserves the area form $dt \wedge d\theta$ and has an extension to a homeomorphism on the closed annulus $\hat{\varphi} : [0, 1] \times \partial D \rightarrow [0, 1] \times \partial D$ given by*

$$\hat{\varphi}(0, x) = \left(0, \frac{D\varphi(0)[x]}{|D\varphi(0)[x]|} \right).$$

Proof. First, $\pi^*(dx \wedge dy) = \pi^*(rdr \wedge d\theta) = \sqrt{t}d(\sqrt{t}) \wedge d(\theta) = (1/2)dt \wedge d\theta$. So $\hat{\varphi}$ preserves the area form $dt \wedge d\theta$.

It remains to justify that the extension is continuous. Define $\Phi \in C^\infty(D, \mathfrak{gl}(\mathbb{R}^2))$ by

$$\Phi(z)[v] = \int_0^1 D\varphi(tz)[v]dt$$

for all $z \in D$ and $v \in \mathbb{R}^2$. As $\varphi(0) = 0$, by the fundamental theorem of calculus

$$(40) \quad \varphi(z) = \Phi(z)[z]$$

for all $z \in D$. Hence the continuous map $[0, 1] \times \partial D \rightarrow \mathbb{R}^2$ given by

$$(t, x) \mapsto \Phi(\sqrt{t}x)[x]$$

is nowhere vanishing; on $t > 0$ because of relation (40) and that φ maps only the origin to the origin; at $t = 0$ because $\Phi(0) = D\varphi(0)$ has no kernel.

A calculation shows that for all $t > 0$ and $x \in \partial D$,

$$(41) \quad \hat{\varphi}(t, x) = \left(|\varphi(\sqrt{t}x)|^2, \frac{\varphi(\sqrt{t}x)}{|\varphi(\sqrt{t}x)|} \right).$$

Using (40) we have, for $t > 0$ and $x \in \partial D$,

$$(42) \quad \frac{\varphi(\sqrt{t}x)}{|\varphi(\sqrt{t}x)|} = \frac{\Phi(\sqrt{t}x)[x]}{|\Phi(\sqrt{t}x)[x]|}.$$

As we observed, the denominator in the right hand side is well defined, continuous, and nowhere vanishing, even at $t = 0$. Thus the right hand

side of (42) defines a continuous extension of the left hand side, and at $t = 0$ takes the value

$$\frac{D\varphi(0)[x]}{|D\varphi(0)[x]|}.$$

Thus the right hand side of (41) extends continuously to $t = 0$, and the extended map satisfies

$$\hat{\varphi}(0, x) = \left(0, \frac{D\varphi(0)[x]}{|D\varphi(0)[x]|} \right).$$

This proves lemma 35. ■

A real linear map $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defines a circle map by

$$(43) \quad \begin{aligned} f_T : \partial D &\rightarrow \partial D \\ v &\mapsto \frac{Tv}{|Tv|}. \end{aligned}$$

Let us record some elementary properties of this correspondence:

- (1) If T is an orientation preserving isomorphism then f_T is an orientation preserving homeomorphism.
- (2) If T_1 and T_2 are two linear maps then $f_{T_1 \circ T_2} = f_{T_1} \circ f_{T_2}$, and moreover $f_{\text{id}} = \text{id}_{S^1}$.
- (3) T has a positive real eigenvalue if and only if f_T has a fixed point if and only if $\text{rot}(f_T) = 0$. Therefore T has a negative real eigenvalue implies $\text{rot}(f_T) \in \{0, 1/2\}$.

Proof of lemma 34. By lemma 35 we may conjugate the map $\varphi : D \setminus \{0\} \rightarrow D \setminus \{0\}$ via a smooth orientation preserving change of coordinates $\pi : (0, 1] \times \partial D \rightarrow D \setminus \{0\}$ to obtain an area preserving map on the half open annulus that extends to a homeomorphism on the closed annulus

$$\hat{\varphi} : [0, 1] \times \partial D \rightarrow [0, 1] \times \partial D.$$

Moreover, on the “new” boundary $\{0\} \times \partial D$ the extended map is the circle map induced by the differential of φ at the removed fixed point. That is,

$$\hat{\varphi}(0, x) = \left(0, \frac{D\varphi(0)[x]}{|D\varphi(0)[x]|} \right)$$

for all $x \in \partial D$. The annulus map $\hat{\varphi}$ preserves Lebesgue measure and is homotopic to the identity and so the Poincaré-Birkhoff fixed point theorem applies. That is, since φ has no periodic points on $D \setminus \{0\}$ the map $\hat{\varphi}$ has no periodic points in $(0, 1] \times \partial D$, so the restriction of $\hat{\varphi}$ to its two boundary components give circle homeomorphisms with equal rotation numbers. In terms of φ this is

$$\text{rot}(f_{D\varphi(0)}) = \text{rot}(\varphi|_{\partial D})$$

where $f_{D\varphi(0)}$ is the circle map determined by the linear map $T = D\varphi(0) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ as described in (43) above. As $\varphi|_{\partial D}$ has no periodic points, its rotation number

$$[\alpha] := \text{rot}(f_{D\varphi(0)}) \in \mathbb{R}/\mathbb{Z}$$

is irrational. Therefore, $\text{rot}(f_{D\varphi(0)})$ is irrational and so $D\varphi(0)$ cannot have any real eigenvalues from the discussion immediately following (43) above. Hence, being symplectic, the eigenvalues of $D\varphi(0)$ must be of the form

$$(44) \quad \sigma(D\varphi(0)) = \{e^{2\pi i\theta}, e^{-2\pi i\theta}\}$$

for some $\theta \notin (\mathbb{Z} \cup 1/2\mathbb{Z})$. From the real canonical form, $D\varphi(0)$ is therefore similar to the rotation through $2\pi\theta$ and also to the rotation through $-2\pi\theta$. Since these two rotations are themselves conjugate to each other via an orientation reversing linear map, we conclude that there always exists an *orientation preserving* real linear map $P : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that

$$P^{-1}D\varphi(0)P = R_{2\pi\theta} \quad \text{or} \quad P^{-1}D\varphi(0)P = R_{-2\pi\theta}.$$

Therefore, $[\alpha] = \text{rot}(f_{D\varphi(0)}) = \text{rot}(f_{R_{\pm 2\pi\theta}}) = \pm[\theta]$. Meaning that $[\alpha] = [\theta]$ or $[\alpha] = -[\theta]$. So by (44) the eigenvalues of $D\varphi(0)$ are $\{e^{2\pi i\alpha}, e^{-2\pi i\alpha}\}$. ■

REFERENCES

- [1] D. V. Anosov, A. B. Katok, New examples in smooth ergodic theory. Ergodic diffeomorphisms. Trans. Moscow Math. Soc. 23 (1970), 1-35
- [2] F. Béguin, S. Crovisier, F. Le Roux, A Patou, Pseudo-rotations of the closed annulus: variation on a theorem of J. Kwapisz. Nonlinearity 17 (2004), 1427-1453.
- [3] F. Béguin, S. Crovisier, F. Le Roux, Pseudo-rotations of the open annulus. Bull. Braz. Math. Soc. (N.S.) 37 (2006), no. 2, 275-306.
- [4] F. Bourgeois, Y. Eliashberg, H. Hofer, K. Wysocki and E. Zehnder, Compactness results in symplectic field theory. Geom. Topol. 7 (2003), 799-888 (electronic).
- [5] B. Bramham, Finite energy foliations with a chosen binding orbit. In preparation.
- [6] B. Bramham, H. Hofer, First steps towards a symplectic dynamics. To appear in Surveys in Differential Geometry, Vol. 17 (2012).
- [7] L. E. J. Brouwer, Über die periodischen Transformationen der Kugel. Math. Ann. 80 (1919), no. 1, 39-41.
- [8] K. Cieliebak, E. Volkov, First steps in stable Hamiltonian topology. arXiv:1003.5084.
- [9] S. Eilenberg, Sur les transformations periodiques de la surface de sphere, Fund. Math. 22 (1934) 228-244.
- [10] B. Fayad, A. Katok, Constructions in elliptic dynamics. Ergodic Theory Dynam. Systems 24 (2004), no. 5, 1477-1520.
- [11] B. Fayad, M. Saprykina, Weak mixing disc and annulus diffeomorphisms with arbitrary Liouville rotation number on the boundary. Ann. Sci. École Norm. Sup. (4) 38 (2005), 339-364.
- [12] A. Floer, The unregularized gradient flow of the symplectic action. Comm. Pure Appl. Math. 41 (1988), no. 6, 775-813.
- [13] J. Franks, Generalizations of the Poincar-Birkhoff theorem. Ann. of Math. (2) 128 (1988), no. 1, 139-151.
- [14] J. Franks, Geodesics on S^2 and periodic points of annulus homeomorphisms. Invent. Math. 108 (1992), no. 2, 403-418.
- [15] M. Gromov, Pseudo-holomorphic curves in symplectic manifolds, Invent. Math. 82 (1985) 307-347

- [16] M. Herman, Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations. *Inst. Hautes Études Sci. Publ. Math.* No. 49 (1979), 5-233.
- [17] H. Hofer, Pseudoholomorphic curves in symplectizations with applications to the Weinstein conjecture in dimension three. *Invent. Math.* 114 (1993), 515-563.
- [18] H. Hofer, K. Wysocki, E. Zehnder, Properties of pseudoholomorphic curves in symplectisations. I. Asymptotics. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 13 (1996), no. 3, 337-379.
- [19] H. Hofer, K. Wysocki, E. Zehnder, A characterisation of the tight three-sphere, *Duke Math. J.* 81 (1) (1995) 159-226.
- [20] H. Hofer, K. Wysocki, E. Zehnder, The dynamics on three-dimensional strictly convex energy surfaces. *Ann. of Math. (2)* 148 (1998), no. 1, 197-289.
- [21] H. Hofer, K. Wysocki, E. Zehnder, A characterization of the tight 3-sphere. II. *Comm. Pure Appl. Math.* 52 (1999), no. 9, 1139-1177.
- [22] H. Hofer, K. Wysocki, E. Zehnder, Properties of pseudoholomorphic curves in symplectizations. III. Fredholm theory. *Topics in nonlinear analysis*, 381-475, *Progr. Nonlinear Differential Equations Appl.*, 35, Birkhäuser, Basel, 1999.
- [23] H. Hofer, K. Wysocki, E. Zehnder, Finite energy foliations of tight three-spheres and Hamiltonian dynamics, *Ann. Math. (2)* 157 (1) (2003) 125-255.
- [24] H. Hofer, K. Wysocki, E. Zehnder, Pseudoholomorphic curves and dynamics in three dimensions. *Handbook of dynamical systems*, Vol. 1A, 1129-1188, North-Holland, Amsterdam, 2002.
- [25] A. Katok, Lyapunov exponents, entropy and periodic orbits for diffeomorphisms. *Inst. Hautes Études Sci. Publ. Math.* No. 51 (1980), 137-173.
- [26] A. Katok, Open problems in elliptic dynamics, www.math.psu.edu/katok_a/problems.html
- [27] A. Katok, B. Hasselblatt, Introduction to the modern theory of dynamical systems. With a supplementary chapter by Katok and Leonardo Mendoza. *Encyclopedia of Mathematics and its Applications*, 54. Cambridge University Press, Cambridge, 1995. xviii+802 pp.
- [28] B. Kerekjarto, Sur la structure des transformations topologiques des surfaces en elles-mêmes, *Enseign. Math.* 35 (1936) 297-316
- [29] P. Le Calvez, Ensembles invariants non enlacs des difféomorphismes du tore et de l'anneau. (Unlinked invariant sets of torus and annulus diffeomorphisms). *Invent. Math.* 155 (2004), no. 3, 561-603.
- [30] P. Le Calvez, From Brouwer theory to the study of homeomorphisms of surfaces. *International Congress of Mathematicians. Vol. III*, 77-98, Eur. Math. Soc., Zürich, 2006.
- [31] L. G. Shnirelman, An example of a transformation of the plane (Russian). *Proc. Don Polytechnic Inst (Novocherkassk)* 14 (Science section, Fis-math. part) (1930), 64-74.
- [32] R. Siefring, Intersection theory of punctured pseudoholomorphic curves. *Geom. Topol.* 15 (2011), no. 4, 2351-2457.
- [33] C. Wendl, Punctured holomorphic curves with boundary in 3-manifolds: Fredholm theory and embeddedness, in preparation.

BARNEY BRAMHAM, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NJ 08540

E-mail address: bramham@ias.edu