

# BOOLEAN NESTED CANALIZING FUNCTIONS: A COMPREHENSIVE ANALYSIS

YUAN LI<sup>1\*</sup>, JOHN O. ADEYEYE<sup>2\*</sup>, DAVID MURRUGARRA<sup>3\*\*</sup>, BORIS AGUILAR<sup>4</sup>, REINHARD LAUBENBACHER<sup>5\*\*</sup>

**ABSTRACT.** Boolean network models of molecular regulatory networks have been used successfully in computational systems biology. The Boolean functions that appear in published models tend to have special properties, in particular the property of being nested canalizing, a property inspired by the concept of canalization in evolutionary biology. It has been shown that networks comprised of nested canalizing functions have dynamic properties that make them suitable for modeling molecular regulatory networks, namely a small number of (large) attractors, as well as relatively short limit cycles.

This paper contains a detailed analysis of this class of functions, based on a novel normal form as polynomial functions over the Boolean field. The concept of layer is introduced that stratifies variables into different classes depending on their level of dominance. Using this layer concept a closed form formula is derived for the number of nested canalizing functions with a given number of variables. Additional metrics analyzed include Hamming weight, the activity number of any variable, and the average sensitivity of the function. It is also shown that the average sensitivity of any nested canalizing function is between 0 and 2. This provides a rationale for why nested canalizing functions are stable, since a random Boolean function has average sensitivity  $\frac{n}{2}$ . The paper also contains experimental evidence that the layer number is an important factor in network stability.

## 1. INTRODUCTION

Canalizing Boolean functions were introduced by S. Kauffman [20] as appropriate rules in Boolean network models of gene regulatory networks. More recently, a subclass of these functions, so-called *nested canalizing* functions (NCF) was introduced [21] and studied from the point of view of stability properties of network dynamics. A multi-state version of such functions have been introduced in [32, 33], where it was shown that networks whose dynamics is controlled by nested canalizing functions have similar stability properties, namely large attractor basins and short limit cycles. An analysis

---

*Key words and phrases.* Boolean function, nested canalizing function, layer number, extended monomial, multinomial coefficient, dynamical system, Hamming weight, activity, average sensitivity.

\* Supported by an award from the US DoD # W911NF-11-10166. \*\* Supported by NSF Grant CMMI-0908201.

of published models, both Boolean and multi-state, of molecular regulatory networks revealed that the large majority of regulatory rules in them is canalizing, with most of these in fact nested canalizing [14, 22, 34, 32]. Thus, nested canalizing rules and the properties of networks governed by them are important to study because of their relevance in systems biology. Furthermore, they are also important in computational science. In [16] it was shown that the class of nested canalizing Boolean functions is identical to the class of so-called unate cascade Boolean functions, which has been studied extensively in engineering and computer science. It was shown, for instance, in [8] that this class has the property that it corresponds exactly to the class of Boolean function with corresponding binary decision diagrams of shortest average path length. Thus, a more detailed mathematical study of nested canalizing functions has applications to problems in engineering as well.

In this paper, we carry out such a detailed study for the case of Boolean nested canalizing functions, obtaining a more explicit characterization than the one obtained in [17]. We introduce a new concept, the *layer number*, leading to a finer classification and, in particular, an explicit formula for the number of nested canalizing functions. This provides a closed form solution to the recursive formula derived in [17], which may be of independent mathematical interest. We also study standard properties of Boolean functions, such as variable activity and average sensitivity. In particular, we obtain a formula for the average sensitivity of a nested canalizing function with  $n$  variables, and show that it lies between  $\frac{n}{2^{n-1}}$  as a lower bound and 2 as upper bound (independent of  $n$ ), which is much smaller than  $\frac{n}{2}$ , the average sensitivity of a random Boolean function. This can be interpreted as providing a theoretical justification for why Boolean networks with nested canalizing rules are stable. We also find a formula for the Hamming weight (the number of 1's in its truth table) of a nested canalizing function. Finally, we conjecture that a nested canalizing function with  $n$  variables has maximal average sensitivity if it has the maximal layer number  $n - 1$ . Based on this result, we conjecture the tight upper bound  $\frac{4}{3}$  for this value. The paper is organized as follows. We first review existing results on nested canalizing functions and networks, after which we introduce some definitions and notation. The subsequent sections contain the main results of the paper.

## 2. BACKGROUND

In [31] it was shown that the dynamics of a Boolean network which operates according to canalizing rules is robust with regard to small perturbations. In [19], an exact formula was derived for the number of canalizing functions. In [29], the definition of canalizing functions was generalized to any finite fields  $\mathbb{F}_q$ , where  $q$  is a power of a prime. Both exact formulas and asymptotics for the number of the generalized canalizing functions were obtained.

Nested canalizing functions were introduced in [21]. One important characteristic of (nested) canalizing functions is that they exhibit a stabilizing effect on the dynamics of a system. That is, small perturbations of an initial state should not grow in time and must eventually end up in the same attractor of the initial state. The stability is typically measured using so-called Derrida plots which monitor the Hamming distance between a random initial state and its perturbed state as both evolve over time. If the Hamming distance decreases over time, the system is considered stable. The slope of the Derrida curve is used as a numerical measure of stability. Roughly speaking, the phase space of a stable system has few components and the limit cycle of each component is short.

In [22], the authors studied the dynamics of nested canalizing Boolean networks over a variety of dependency graphs. That is, for a given random graph on  $n$  nodes, where the in-degree of each node is chosen at random between 0 and  $k$ , for  $k \leq n$ , a nested canalizing function is assigned to each node in the in-degree variables of that node. The dynamics of these networks was then analyzed and the stability measured using Derrida plots. It was shown there that nested canalizing networks are remarkably stable regardless of the in-degree distribution and that the stability increases as the average number of inputs of each node increases.

Most published molecular networks are given in the form of a wiring diagram, or dependency graph, constructed from experiments and prior published knowledge. However, for most of the molecular species in the network, little knowledge, if any, could be deduced about their regulatory rules, for instance in the gene transcription networks in yeast [15] and *E. Coli* [3]. Each one of these networks contains more than 1000 genes. Kauffman et. al [21] investigated the effect of the topology of a sub-network of the yeast transcriptional network where many of the transcriptional rules are not known. They generated ensembles of different models where all models have the same dependency graph. Their heuristic results imply that the dynamics of those models which used only nested canalizing functions were far more stable than the randomly generated models. Since it is already established that the yeast transcriptional network is stable, this suggests that the unknown interaction rules are very likely nested canalizing functions. In a recent article [2], a transcriptional network of yeast, with 3459 genes as well as the transcriptional networks of *E. Coli* (1481 genes) and *B. subtilis* (840 genes) have been analyzed in a similar fashion, with similar findings.

The notion of sensitivity was introduced in [12]. The sensitivity of a Boolean function of a variable  $x$  is defined as the number of Hamming neighbors of  $x$  on which the function value is different than on  $x$ . The average sensitivity of the function is then computed by taking the average value of the sensitivities of the function on all the input  $x$ . Although the definition is straightforward, the sensitivity measure is understood only for a few classes of functions. For monotone functions, Ilya Shmulevich [38] derived asymptotic formulas for a typical monotone Boolean functions. Recently,

S. Zhang [44] found a formula for the average sensitivity of any monotone Boolean function, and derived a tight bound. In [39], I. Shmulevich and S. A. Kauffman obtained the average sensitivity of Boolean functions with only one canalizing variable.

### 3. DEFINITIONS AND NOTATION

Let  $\mathbb{F} = \mathbb{F}_2$  be the field with 2 elements, and let  $f : \mathbb{F}^n \rightarrow \mathbb{F}$ . It is well known [30] that  $f$  can be expressed as a polynomial, called the algebraic normal form (ANF) of  $f$ :

$$f(x_1, \dots, x_n) = \bigoplus_{\substack{0 \leq k_i \leq 1, \\ i=1, \dots, n}} a_{k_1 \dots k_n} x_1^{k_1} \dots x_n^{k_n}$$

where each coefficient  $a_{k_1 \dots k_n} \in \mathbb{F}$ . The number  $k_1 + k_2 + \dots + k_n$  is the multivariate degree of the term  $a_{k_1 k_2 \dots k_n} x_1^{k_1} x_2^{k_2} \dots x_n^{k_n}$  with nonzero coefficient  $a_{k_1 k_2 \dots k_n}$ . The greatest degree of all the terms of  $f$  is called its algebraic degree, denoted by  $\deg(f)$ . The symbol  $\bigoplus$  stands for addition modulo 2, whereas the symbol  $+$  will be reserved for addition of real numbers.

**Definition 3.1.** *The function  $f(x_1, x_2, \dots, x_n)$  is essential in the variable  $x_i$  if there exist  $r, s \in \mathbb{F}$  and  $x_1^*, \dots, x_{i-1}^*, x_{i+1}^*, \dots, x_n^*$  such that*

$$f(x_1^*, \dots, x_{i-1}^*, r, x_{i+1}^*, \dots, x_n^*) \neq f(x_1^*, \dots, x_{i-1}^*, s, x_{i+1}^*, \dots, x_n^*).$$

**Definition 3.2.** *A function  $f(x_1, x_2, \dots, x_n)$  is  $\langle i : a : b \rangle$  canalizing if  $f(x_1, \dots, x_{i-1}, a, x_{i+1}, \dots, x_n) = b$ , for all  $x_j, j \neq i$ , where  $i \in \{1, \dots, n\}$ ,  $a, b \in \mathbb{F}$ .*

This definition is reminiscent of the concept of ‘‘canalization’’ introduced by the geneticist C. H. Waddington [42] to represent the ability of a genotype to produce the same phenotype in the presence of environmental variability.

**Definition 3.3.** *Let  $f$  be a Boolean function in  $n$  variables. Let  $\sigma$  be a permutation of the set  $\{1, 2, \dots, n\}$ . The function  $f$  is a nested canalizing function in the variable order  $x_{\sigma(1)}, \dots, x_{\sigma(n)}$  with canalizing input values  $a_1, \dots, a_n$  and canalized values  $b_1, \dots, b_n$ , if it can be represented in the*

$$\text{form } f(x_1, \dots, x_n) = \begin{cases} b_1 & x_{\sigma(1)} = a_1, \\ b_2 & x_{\sigma(1)} = \bar{a}_1, x_{\sigma(2)} = a_2, \\ b_3 & x_{\sigma(1)} = \bar{a}_1, x_{\sigma(2)} = \bar{a}_2, x_{\sigma(3)} = a_3, \\ \dots & \\ b_n & x_{\sigma(1)} = \bar{a}_1, \dots, x_{\sigma(n-1)} = \bar{a}_{n-1}, x_{\sigma(n)} = a_n, \\ \bar{b}_n & x_{\sigma(1)} = \bar{a}_1, \dots, x_{\sigma(n-1)} = \bar{a}_{n-1}, x_{\sigma(n)} = \bar{a}_n. \end{cases}$$

Here,  $\bar{a} = a \oplus 1$ .

*The function  $f$  is nested canalizing if  $f$  is nested canalizing in the variable order  $x_{\sigma(1)}, \dots, x_{\sigma(n)}$  for some permutation  $\sigma$ .*

Let  $\alpha = (a_1, a_2, \dots, a_n)$  and  $\beta = (b_1, b_2, \dots, b_n)$ . We say that  $f$  is  $\{\sigma : \alpha : \beta\}$  NCF if it is NCF in the variable order  $x_{\sigma(1)}, \dots, x_{\sigma(n)}$  with canalizing input values  $\alpha = (a_1, \dots, a_n)$  and canalized values  $\beta = (b_1, \dots, b_n)$ .

Given a vector  $\alpha = (a_1, a_2, \dots, a_n)$ , we define  $\alpha^{i_1, \dots, i_k} = (a_1, \dots, \overline{a_{i_1}}, \dots, \overline{a_{i_k}}, \dots, a_n)$ . Then, from the above definition, we immediately have the following result.

**Proposition 3.4.** *The function  $f$  is  $\{\sigma : \alpha : \beta\}$  NCF if and only if  $f$  is  $\{\sigma : \alpha^n : \beta^n\}$  NCF.*

**Example 3.5.** *The function  $f(x_1, x_2, x_3) = x_1(x_2 \oplus 1)x_3 \oplus 1$  is  $\{(1, 2, 3) : (0, 1, 0) : (1, 1, 1)\}$  NCF. Actually, one can check that this function is nested canalizing in any variable order.*

**Example 3.6.** *Let  $f(x_1, x_2, x_3) = (x_1 \oplus 1)(x_2(x_3 \oplus 1) \oplus 1) \oplus 1$ . This function is  $\{(1, 2, 3) : (1, 0, 1) : (1, 0, 0)\}$  NCF. It is also  $\{(1, 3, 2) : (1, 1, 1) : (1, 0, 1)\}$  NCF. One can check that this function can be nested canalizing in only two variable orders, namely  $(x_1, x_2, x_3)$  and  $(x_1, x_3, x_2)$ .*

From the above definitions, we know that if a function is NCF, all the variables appearing in it must be essential. However, a constant function  $b$  can be  $\langle i : a : b \rangle$  NCF for any  $i$  and  $a$ .

#### 4. A DETAILED CATEGORIZATION OF NESTED CANALIZING FUNCTIONS

As we will see, in a nested canalizing function, some variables are more dominant than others. We will classify all the variables of an NCF into different levels according to the extent of their dominance.

**Definition 4.1.** [18] *A function  $M(x_1, \dots, x_n)$  is an extended monomial of essential variables  $x_1, \dots, x_n$  if*

$$M(x_1, \dots, x_n) = (x_1 \oplus a_1) \cdots (x_n \oplus a_n),$$

where  $a_i \in \mathbb{F}_2$  for  $i \in \{0, \dots, n\}$ .

We now rewrite Theorem 3.1 in [17] with more information.

**Lemma 4.2.** *The function  $f(x_1, x_2, \dots, x_n)$  is  $\langle i : a : b \rangle$  canalizing if and only if  $f(x_1, x_2, \dots, x_n) = (x_i \oplus a)Q(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \oplus b$  for some polynomial  $Q$ .*

*Proof.* Using the algebraic normal form of  $f$ , we rewrite it as

$$f = x_i g_1(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \oplus g_0(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n). \text{ Hence, } \\ f(X) = f(x_1, x_2, \dots, x_n) = (x_i \oplus a)g_1(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \oplus a g_1(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \oplus \\ g_0(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n).$$

$$\text{Let } g_1(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) = Q(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$$

$$\text{and } r(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) =$$

$$a g_1(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \oplus g_0(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n).$$

$$\text{Then } f(X) = f(x_1, \dots, x_n) =$$

$(x_i \oplus a)Q(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \oplus r(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$ . Since  $f(X)$  is  $\langle i : a : b \rangle$  canalizing, we get that

$$f(X) = f(x_1, \dots, x_{i-1}, a, x_{i+1}, \dots, x_n) = b$$

for any  $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n$ , i.e.,  $r(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) = b$  for any  $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n$ . So  $r(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$  must be the constant  $b$ . This shows necessity, and sufficiency is obvious.  $\square$

**Remark 4.3.** *We have the following observations.*

- (1) *When we contrast this lemma to the first part of Theorem 3.1 in [17], it is important to note that  $x_i$  is not essential in  $Q$ .*
- (2) *In [29], there is a general version of this result over any finite field.*
- (3) *In the above lemma, if  $f$  is constant, then  $Q = 0$ .*

From Definition 3.3, we have the following result.

**Proposition 4.4.** *Let  $f(x_1, \dots, x_n)$  be  $\{\sigma : \alpha : \beta\}$  NCF, i.e.,  $f$  is NCF in the variable order  $x_{\sigma(1)}, \dots, x_{\sigma(n)}$  with canalizing input values  $\alpha = (a_1, \dots, a_n)$  and canalized output values  $\beta = (b_1, \dots, b_n)$ . For  $1 \leq k \leq n-1$ , let  $x_{\sigma(1)} = \overline{a_1}, \dots, x_{\sigma(k)} = \overline{a_k}$ . Then the function  $f(x_1, \dots, \overline{a_1}, \dots, \overline{a_k}, \dots, x_n)$  is  $\{\sigma^* : \alpha^* : \beta^*\}$  NCF on the remaining variables, where  $\sigma^* = x_{\sigma(k+1)}, \dots, x_{\sigma(n)}$ ,  $\alpha^* = (a_{k+1}, \dots, a_n)$  and  $\beta^* = (b_{k+1}, \dots, b_n)$ .*

**Definition 4.5.** *Let  $f(x_1, \dots, x_n)$  be NCF. We call a variable  $x_i$  a most dominant variable of  $f$  if there is a variable order  $\alpha = (x_i, \dots)$  such that  $f$  is NCF with in this variable order.*

In Example 3.5, all three variables are most dominant. In Example 3.6, only  $x_1$  is a most dominant variable. We have:

**Theorem 4.1.** *Given an NCF  $f(x_1, \dots, x_n)$ , all variables are most dominant if and only if  $f = M(x_1, \dots, x_n) \oplus b$ , where  $M$  is an extended monomial, i.e.,  $M = (x_1 \oplus a_1)(x_2 \oplus a_2) \cdots (x_n \oplus a_n)$ .*

*Proof.* If  $x_1$  is most dominant, from Lemma 4.2, we know there exist  $a_1$  and  $b$  such that  $f(x_1, x_2, \dots, x_n) = (x_1 \oplus a_1)Q(x_2, \dots, x_n) \oplus b$ , i.e.,  $(x_1 \oplus a_1)|(f \oplus b)$ . Now, if  $x_2$  is also most dominant, then we have  $a_2$  and  $b'$  such that  $f(x_1, a_2, x_3, \dots, x_n) = b'$  for any  $x_1, x_3, \dots, x_n$ . Specifically, if  $x_1 = a_1$ , we get  $f(a_1, a_2, x_3, \dots, x_n) = b = b'$ . Hence, we also get  $(x_2 \oplus a_2)|(f \oplus b) = (x_1 \oplus a_1)Q(x_2, \dots, x_n)$ . Since  $x_1 \oplus a_1$  and  $x_2 \oplus a_2$  are coprime, we get  $(x_2 \oplus a_2)|Q(x_2, \dots, x_n)$ , hence,  $f(x_1, x_2, \dots, x_n) = (x_1 \oplus a_1)(x_2 \oplus a_2)Q'(x_3, \dots, x_n) \oplus b$ . With the induction principle, the necessity is proved. The sufficiency is evident.  $\square$

We are now ready to prove the main result of this section.

**Theorem 4.2.** *Given  $n \geq 2$ , the function  $f(x_1, \dots, x_n)$  is nested canalizing if and only if it can be uniquely written as*

$$f(x_1, \dots, x_n) = M_1(M_2(\cdots (M_{r-1}(M_r \oplus 1) \oplus 1) \cdots) \oplus 1) \oplus b, \quad (4.1)$$

where each  $M_i$  is an extended monomial in a set of disjoint variables. More precisely,  $M_i = \prod_{j=1}^{k_i} (x_{i_j} \oplus a_{i_j})$ ,  $i = 1, \dots, r$ ,  $k_i \geq 1$  for  $i = 1, \dots, r-1$ ,  $k_r \geq 2$ ,  $k_1 + \cdots + k_r = n$ ,  $a_{i_j} \in \mathbb{F}_2$ ,  $\{i_j | j = 1, \dots, k_i, i = 1, \dots, r\} = \{1, \dots, n\}$ .

*Proof.* We use induction on  $n$ . When  $n = 2$ , there are 16 Boolean functions, 8 of which are NCFs. Namely

$$(x_1 \oplus a_1)(x_2 \oplus a_2) \oplus c = M_1 \oplus 1 \oplus b,$$

where  $b = 1 \oplus c$  and  $M_1 = (x_1 \oplus a_1)(x_2 \oplus a_2)$ .

If  $(x_1 \oplus a_1)(x_2 \oplus a_2) \oplus c = (x_1 \oplus a_1')(x_2 \oplus a_2)' \oplus c'$ , then, by equating coefficients, we immediately obtain  $a_1 = a_1'$ ,  $a_2 = a_2'$  and  $c = c'$ . So uniqueness holds. We have proved that equation 4.1 is true for  $n = 2$ , where  $r = 1$ .

Assume now that Equation 4.1 is true for any nested canalizing function which has at most  $n - 1$  essential variables. Consider a NCF  $f(x_1, \dots, x_n)$ . Suppose  $x_{\sigma(1)}, \dots, x_{\sigma(k_1)}$  are all most dominant canalizing variables of  $f$ , for  $1 \leq k_1 \leq n$ .

*Case 1:*  $k_1 = n$ . Then, by Theorem 4.1, the conclusion is true with  $r = 1$ .

*Case 2:*  $k_1 < n$ . Then, with the same arguments as in Theorem 4.1, we get that  $f = M_1 g \oplus b$ , where  $M_1 = (x_{\sigma(1)} \oplus a_{\sigma(1)}) \cdots (x_{\sigma(k_1)} \oplus a_{\sigma(k_1)})$ . Let  $x_{\sigma(1)} = \overline{a_{\sigma(1)}}, \dots, x_{\sigma(k_1)} = \overline{a_{\sigma(k_1)}}$  in  $f$ , then  $f = g \oplus b$  (hence,  $g$ ) will also be nested canalizing in the remaining variables, by Proposition 4.4.

Since  $g$  has  $n - k_1 \leq n - 1$  variables, by the induction assumption, we get that  $g = M_2(M_3(\cdots(M_{r-1}(M_r \oplus 1) \oplus 1) \cdots) \oplus 1) \oplus b_1$ . It follows that  $b_1$  must be 1. Otherwise, all the variables in  $M_2$  will also be most dominant variables of  $f$ . This completes the proof.  $\square$

Because each NCF can be uniquely written in the form 4.1 and the number  $r$  is uniquely determined by  $f$ , we can make the following definition.

**Definition 4.6.** *For an NCF  $f$ , written in the form 4.1, the number  $r$  will be called its layer number. Essential variables of  $M_1$  will be called most dominant variables (canalizing variables), and are part of the first layer of  $f$ . Essential variables of  $M_2$  will be called second most dominant variables and are part of the second layer; etc.*

The function in Example 3.5 has layer number 1 and the function in Example 3.6 has layer number 2.

**Remark 4.7.** *We make some remarks on Theorem 4.2.*

- (1) *It is impossible that  $k_r = 1$ . Otherwise,  $M_r \oplus 1$  will be a factor of  $M_{r-1}$ , which means that the layer number is  $r - 1$ . Hence,  $k_r \geq 2$ .*
- (2) *If variable  $x_i$  is in the first layer, and  $x_i \oplus a_i$  is a factor of  $M_i$ , then this NCF is  $\langle i : a_i : b \rangle$  canalizing. We simply say that  $x_i$  is a canalizing variable of this NCF.*

From the previous examples, we know that an NCF can have different variable orders, but only the variables in the same layer can be reordered. More precisely, we have the following result.

**Corollary 4.8.** *If  $\sigma$  and  $\eta$  are two permutations on  $\{1, \dots, n\}$ , and  $f$  is both  $\{\sigma : \alpha : \beta\}$  NCF and  $\{\eta : \alpha' : \beta'\}$  NCF, then we have*

$$\{x_{\sigma(1)}, \dots, x_{\sigma(k_1)}\} = \{x_{\eta(1)}, \dots, x_{\eta(k_1)}\}, \dots,$$

and

$$\{x_{\sigma(k_1+\dots+k_{r-1}+1)}, \dots, x_{\sigma(n)}\} = \{x_{\eta(k_1+\dots+k_{r-1}+1)}, \dots, x_{\eta(n)}\}.$$

We also have  $\beta = \beta' = (b_1, \dots, b_n)$ , and

$$b_1 = \dots = b_{k_1}, \dots, b_{k_1+\dots+k_{r-1}+1} = \dots = b_n.$$

Furthermore,

$$b_1 \neq b_{k_1+1} \neq b_{k_1+k_2+1} \neq \dots \neq b_{k_1+\dots+k_{r+1}+1}.$$

*Proof.* These results all follow from the expression for  $f$  in Theorem 4.2.  $\square$

From this corollary we can determine the layer number of any NCF by its canalized value. For example, if  $f$  is NCF with canalized value  $(1, 1, 1, 0, 0, 0, 1, 0, 0, 1, 1)$  ( $n = 11$ ), then its layer number is 5.

Let  $\text{NCF}(n, r)$  denote the set of all nested canalizing functions in  $n$  variables with layer number  $r$ , and let  $\text{NCF}(n)$  denote the set of all nested canalizing functions in  $n$  variables.

**Corollary 4.9.** *For  $n \geq 2$ ,*

$$|\text{NCF}(n, r)| = 2^{n+1} \sum_{\substack{k_1+\dots+k_r=n \\ k_i \geq 1, i=1, \dots, r-1, k_r \geq 2}} \binom{n}{k_1, \dots, k_{r-1}},$$

and

$$|\text{NCF}(n)| = 2^{n+1} \sum_{r=1}^{n-1} \sum_{\substack{k_1+\dots+k_r=n \\ k_i \geq 1, i=1, \dots, r-1, k_r \geq 2}} \binom{n}{k_1, \dots, k_{r-1}}$$

where the multinomial coefficient  $\binom{n}{k_1, \dots, k_{r-1}}$  is equal to  $\frac{n!}{k_1! \dots k_{r-1}!}$ .

*Proof.* It follows from Equation 4.1, that for each choice of  $k_1, \dots, k_r$ , with  $k_1 + \dots + k_r = n$ ,  $k_i \geq 1$ ,  $i = 1, \dots, r-1$  and  $k_r \geq 2$ , there are  $2^{k_j} \binom{n-k_1-\dots-k_{j-1}}{k_j}$  ways to form  $M_j$ ,  $j = 1, \dots, r$ .

Note that we have two choices for  $b$ . Hence,

$$\begin{aligned} |\text{NCF}(n, r)| &= 2 \sum_{\substack{k_1+\dots+k_r=n \\ k_i \geq 1, i=1, \dots, r-1, k_r \geq 2}} 2^{k_1+\dots+k_r} \\ &\quad \binom{n}{k_1} \binom{n-k_1}{k_2} \dots \binom{n-k_1-\dots-k_{r-1}}{k_r} \\ &= 2^{n+1} \sum_{\substack{k_1+\dots+k_r=n \\ k_i \geq 1, i=1, \dots, r-1, k_r \geq 2}} \end{aligned}$$



$$\begin{aligned}
 & \frac{n!}{(k_1)!(n-k_1)!} \frac{(n-k_1)!}{(k_2)!(n-k_1-k_2)!} \cdots \\
 & \frac{(n-k_1-\cdots-k_{r-1})!}{k_r!(n-k_1-\cdots-k_r)!} \\
 & = 2^{n+1} \sum_{\substack{k_1+\cdots+k_r=n \\ k_i \geq 1, i=1, \dots, r-1, k_r \geq 2}} \frac{n!}{k_1!k_2!\cdots k_r!} = \\
 & = 2^{n+1} \sum_{\substack{k_1+\cdots+k_r=n \\ k_i \geq 1, i=1, \dots, r-1, k_r \geq 2}} \binom{n}{k_1, \dots, k_{r-1}}.
 \end{aligned}$$

Since  $\text{NCF}(n) = \bigcup_{r=1}^{n-1} \text{NCF}(n, r)$  and  $\text{NCF}(n, i) \cap \text{NCF}(n, j) = \phi$  when  $i \neq j$ , we get the formula for  $|\text{NCF}(n)|$ .  $\square$

As examples, one can check that  $|\text{NCF}(2)| = 8$ ,  $|\text{NCF}(3)| = 64$ ,  $|\text{NCF}(4)| = 736$ ,  $|\text{NCF}(5)| = 10624$ ,  $\dots$ . These results are consistent with those in [4, 37]. By equating our formula to the recursive relation in [4, 37], we have

**Corollary 4.10.** *The solution of the nonlinear recursive sequence*

$$a_2 = 8, a_n = \sum_{r=2}^{n-1} \binom{n}{r-1} 2^{r-1} a_{n-r+1} + 2^{n+1}, n \geq 3$$

is

$$a_n = 2^{n+1} \sum_{r=1}^{n-1} \sum_{\substack{k_1+\cdots+k_r=n \\ k_i \geq 1, i=1, \dots, r-1, k_r \geq 2}} \binom{n}{k_1, \dots, k_{r-1}}.$$

## 5. HAMMING WEIGHT, ACTIVITY, AND AVERAGE SENSITIVITY

A Boolean function is called *balanced* if it takes on the value 1 on exactly half the states (and 0 on the other half). In other words, its Hamming weight (the number of 1's in its truth table) is  $2^{n-1}$ , where  $n$  is the number of variables. There are  $\binom{2^n}{2^{n-1}}$  balanced Boolean functions. It is easy to show that a Boolean function with canalizing variables is not balanced, i.e., is biased, actually, very biased. For example, the two constant functions are trivially canalizing, and they are also the most biased functions. Extended monomial functions are the second most biased since for any of them, only one value is nonzero. But biased functions may have no canalizing variables. For example,  $f(x_1, x_2, x_3) = x_1x_2x_3 \oplus x_1x_2 \oplus x_1x_3 \oplus x_2x_3$  is biased but without canalizing variables.

In Boolean functions, some variables have greater influence over the output of the function than other variables. To formalize this, a concept called *activity* was introduced. Let  $x = (x_1, \dots, x_n)$ , and

$$\frac{\partial f(x)}{\partial x_i} = f(x_1, \dots, x_i \oplus 1, \dots, x_n) \oplus f(x_1, \dots, x_i, \dots, x_n).$$

The *activity* of variable  $x_i$  is defined as

$$\lambda_i^f = \frac{1}{2^n} \sum_{(x_1, \dots, x_n) \in \mathbb{F}_2^n} \frac{\partial f(x_1, \dots, x_n)}{\partial x_i} \quad (5.1)$$

Note that the above definition can also be written as follows:

$$\lambda_i^f = \frac{1}{2^{n-1}} \sum_{(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \in \mathbb{F}_2^{n-1}} (f(x_1, \dots, \overset{i}{0}, \dots, x_n) \oplus f(x_1, \dots, \overset{i}{1}, \dots, x_n)) \quad (5.2)$$

The activity of any variable in a constant function is 0. For an affine function  $f(x_1, \dots, x_n) = x_1 \oplus \dots \oplus x_n \oplus b$ ,  $\lambda_i^f = 1$  for any  $i$ . It is clear, for any  $f$  and  $i$ , that we have  $0 \leq \lambda_i^f \leq 1$ .

Another important quantity is the sensitivity of a Boolean function, which measures how sensitive the output of the function is if the input changes (This was introduced in [12]). The sensitivity  $s^f(x_1, \dots, x_n)$  of  $f$  on the input  $(x_1, \dots, x_n)$  is defined as the number of Hamming neighbors of  $(x_1, \dots, x_n)$  (that is, all states that have Hamming distance 1) on which the function value is different from  $f(x_1, \dots, x_n)$ . That is,

$$s^f(x_1, \dots, x_n) = |\{i | f(x_1, \dots, \overset{i}{0}, \dots, x_n) \neq f(x_1, \dots, \overset{i}{1}, \dots, x_n), i = 1, \dots, n\}|.$$

Obviously,  $s^f(x_1, \dots, x_n) = \sum_{i=1}^n \frac{\partial f(x_1, \dots, x_n)}{\partial x_i}$ .

The average sensitivity of a function  $f$  is defined as

$$s^f = E[s^f(x_1, \dots, x_n)] = \frac{1}{2^n} \sum_{(x_1, \dots, x_n) \in \mathbb{F}_2^n} s^f(x_1, \dots, x_n) = \sum_{i=1}^n \lambda_i^f.$$

It is clear that  $0 \leq s^f \leq n$ . The concept of average sensitivity of a Boolean function is one of the most studied concepts in the analysis of Boolean functions, and has received a lot attention recently [1, 5, 6, 7, 10, 11, 23, 25, 27, 36, 38, 39, 40, 41, 43]. Bernasconi [5] has shown that a random Boolean function has average sensitivity  $\frac{n}{2}$ . This means the average value of the average sensitivities of all Boolean functions in  $n$  variables is  $\frac{n}{2}$ . In [39], Shmulevich and Kauffman calculated the activity of all the variables of a Boolean function with exactly one canalizing variable and unbiased input for the other variables. Adding all the activities, the average sensitivity of a Boolean function was also obtained.

First, we have

**Lemma 5.1.** *We have the equality*

$$(x_1 \oplus a_1) \cdots (x_k \oplus a_k) = \begin{cases} 1, & \text{if } (x_1, \dots, x_k) = (\overline{a_1}, \dots, \overline{a_k}) \\ 0, & \text{otherwise.} \end{cases}$$

That is, only one value is equal to 1 and all the other  $2^k - 1$  values are 0.

**Theorem 5.1.** For  $n \geq 2$ , let  $f_1 = M_1$ ,  $f_r = M_1(M_2(\cdots(M_{r-1}(M_r \oplus 1) \oplus 1)\cdots) \oplus 1)$ ,  $r \geq 2$ , where  $M_i$  is same as in Theorem 4.2. Then the Hamming weight of  $f_r$  is

$$W(f_r) = \sum_{j=1}^r (-1)^{j-1} 2^{n-\sum_{i=1}^j k_i} \quad (5.3)$$

The Hamming weight of  $f_r \oplus 1$  is

$$W(f_r \oplus 1) = \sum_{j=0}^r (-1)^j 2^{n-\sum_{i=1}^j k_i}, \quad (5.4)$$

where  $\sum_{i=1}^0 k_i$  is to be interpreted as 0.

*Proof.* First, consider the Hamming weight of  $f_r$ . When  $r = 1$ , we know the result is true by Lemma 5.1. When  $r > 1$ , we consider two cases:

*Case A:*  $r = 2t + 1$  is odd. Then all the states on which  $f$  evaluates to 1 will be divided into the following disjoint groups:

- Group  $j$  for  $j = 1, \dots, t$ :  $M_1 = 1, M_2 = 1, \dots, M_{2j-1} = 1, M_{2j} = 0$ ;
- Group  $t + 1$ :  $M_1 = 1, M_2 = 1, \dots, M_{2t} = 1, M_{2t+1} = M_r = 1$ .

In Group  $j$ , the number of states is  $(2^{k_{2j}} - 1)2^{n-k_1-\cdots-k_{2j}} = 2^{n-k_1-\cdots-k_{2j-1}-2^{n-k_1-\cdots-k_{2j}}}$ .

In Group  $t + 1$ , the number of vectors is  $2^{n-k_1-\cdots-k_r} = 1$ .

Adding all of them together, we get Equation 5.3.

*Case B:*  $r = 2t$  is even.

The proof in this case is similar, and we omit it.

Because  $|\{(x_1, \dots, x_n) | f(x_1, \dots, x_n) = 0\}| + |\{(x_1, \dots, x_n) | f(x_1, \dots, x_n) = 1\}| = 2^n$ , we know that the Hamming weight of  $f_r \oplus 1$  is equal to

$$\begin{aligned} W(f_r \oplus 1) &= 2^n - W(f_r) = 2^n - \sum_{j=1}^r (-1)^{j-1} 2^{n-\sum_{i=1}^j k_i} \\ &= \sum_{j=0}^r (-1)^j 2^{n-\sum_{i=1}^j k_i}, \end{aligned}$$

where  $\sum_{i=1}^0 k_i$  should be interpreted as 0. □

In the following, we will calculate the activities of the variables of any NCF. Let  $f$  be an NCF, written in the form of Theorem 4.2. Without loss of generality, we can assume that  $M_1 = (x_1 \oplus a_1)(x_2 \oplus a_2) \cdots (x_{k_1} \oplus a_{k_1})$  and  $m_1 = (x_1 \oplus a_1) \cdots (x_{i-1} \oplus a_{i-1})(x_{i+1} \oplus a_{i+1}) \cdots (x_{k_1} \oplus a_{k_1})$ , i.e.,  $M_1 = (x_i \oplus a_i)m_1$ .

If  $r = 1$ , i.e.,  $k_1 = n$ , then

$$\lambda_i^f = \frac{1}{2^{n-1}} \sum_{(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \in \mathbb{F}_2^{n-1}}$$

$$\begin{aligned}
& (f(x_1, \dots, \overset{i}{0}, \dots, x_n) \oplus f(x_1, \dots, \overset{i}{1}, \dots, x_n)) \\
&= \frac{1}{2^{n-1}} \sum_{(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \in \mathbb{F}_2^{n-1}} m_1 = \\
& \frac{1}{2^{n-1}} W(m_1) = \frac{1}{2^{n-1}}.
\end{aligned}$$

by Lemma 5.1.

If  $1 < r \leq n - 1$ , then consider the activity of  $x_i$  in the first layer, i.e.,  $1 \leq i \leq k_1$ . We have

$$\begin{aligned}
\lambda_i^f &= \frac{1}{2^{n-1}} \sum_{(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \in \mathbb{F}_2^{n-1}} \\
& (f(x_1, \dots, \overset{i}{0}, \dots, x_n) \oplus f(x_1, \dots, \overset{i}{1}, \dots, x_n)) \\
&= \frac{1}{2^{n-1}} \sum_{(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \in \mathbb{F}_2^{n-1}} \\
& m_1(M_2(\cdots (M_{r-1}(M_r \oplus 1) \oplus 1) \cdots) \oplus 1) \\
&= \frac{1}{2^{n-1}} W(m_1(M_2(\cdots (M_{r-1}(M_r \oplus 1) \oplus 1) \cdots) \oplus 1)). \\
&= \begin{cases} \frac{1}{2^{n-1}} \sum_{j=1}^r (-1)^{j-1} 2^{n-1 - (\sum_{i=1}^j k_i - 1)}, & \text{if } k_1 > 1 \\ \frac{1}{2^{n-1}} \sum_{j=0}^{r-1} (-1)^j 2^{n-1 - \sum_{i=1}^j k_{i+1}}, & \text{if } k_1 = 1. \end{cases} \\
&= \begin{cases} \frac{1}{2^{n-1}} \sum_{j=1}^r (-1)^{j-1} 2^{n - \sum_{i=1}^j k_i}, & \text{if } k_1 > 1 \\ \frac{1}{2^{n-1}} \sum_{j=0}^{r-1} (-1)^j 2^{n - \sum_{i=0}^j k_{i+1}}, & \text{if } k_1 = 1. \end{cases} \\
&= \begin{cases} \frac{1}{2^{n-1}} \sum_{j=1}^r (-1)^{j-1} 2^{n - \sum_{i=1}^j k_i}, & \text{if } k_1 > 1 \\ \frac{1}{2^{n-1}} \sum_{j=0}^{r-1} (-1)^j 2^{n - \sum_{i=1}^{j+1} k_i}, & \text{if } k_1 = 1. \end{cases} \\
&= \begin{cases} \frac{1}{2^{n-1}} \sum_{j=1}^r (-1)^{j-1} 2^{n - \sum_{i=1}^j k_i}, & \text{if } k_1 > 1 \\ \frac{1}{2^{n-1}} \sum_{j=1}^r (-1)^{j-1} 2^{n - \sum_{i=1}^j k_i}, & \text{if } k_1 = 1. \end{cases} \\
&= \frac{1}{2^{n-1}} \sum_{j=1}^r (-1)^{j-1} 2^{n - \sum_{i=1}^j k_i} \text{ by Theorem 5.1. Note, in the above, } k_1 = 1 \text{ means } m_1 = 1, \text{ so we used Equation 5.4 with layer number } r - 1, \text{ and the first layer is } M_2 \text{ for } n - 1 \text{ variable functions.}
\end{aligned}$$

Now let's consider the variables in the  $l$ -th layer, i.e.,  $x_i$  is an essential variable of  $M_l$ ,  $2 \leq l \leq r - 1$ . We have  $M_l = (x_i + a_i)m_l$  and

$$\begin{aligned}
\lambda_i^f &= \frac{1}{2^{n-1}} \sum_{(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \in \mathbb{F}_2^{n-1}} \\
& (f(x_1, \dots, \overset{i}{0}, \dots, x_n) \oplus f(x_1, \dots, \overset{i}{1}, \dots, x_n)) \\
&= \frac{1}{2^{n-1}} \sum_{(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \in \mathbb{F}_2^{n-1}} \\
& M_1 \cdots M_{l-1} m_l (M_{l+1}(\cdots (M_r \oplus 1) \cdots) \oplus 1).
\end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2^{n-1}} \sum_{j=1}^{r-l+1} (-1)^{j-1} 2^{n-1-((k_1+\dots+k_{l-1})+k_{l+1}+\dots+k_{j+l-1})} = \\
 &\quad \frac{1}{2^{n-1}} \sum_{j=1}^{r-l+1} (-1)^{j-1} 2^{n-\sum_{i=1}^{j+l-1} k_i}
 \end{aligned}$$

by Equation 5.3 in Theorem 5.1. Note,  $M_1 \cdots M_{l-1} m_l$  is the first layer,  $M_{l+1}$  is the second layer, etc.

Let  $x_i$  be the variable in the last layer  $M_r$ , then we have

$$= \frac{1}{2^{n-1}} \sum_{(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \in \mathbb{F}_2^{n-1}} M_1 M_2 \cdots M_{r-1} m_r = \frac{1}{2^{n-1}}$$

by Lemma 5.1. Variables in the same layer have the same activities, so we use  $A_l^f$  to stand for the activity number of each variable in the  $l$ th layer  $M_l$ ,  $1 \leq l \leq r$ . We find that the formula of  $A_l^f$  for  $2 \leq l \leq r-1$  is also true when  $l = r$  or  $r = 1$ . The next theorem summarizes these.

**Theorem 5.2.** *Let  $f$  be a NCF and written as in the Theorem 4.2, then the activity of each variable in the  $l$ th layer,  $1 \leq l \leq r$ , is*

$$A_l^f = \frac{1}{2^{n-1}} \sum_{j=1}^{r-l+1} (-1)^{j-1} 2^{n-\sum_{i=1}^{j+l-1} k_i} \quad (5.5)$$

The average sensitivity of  $f$  is

$$s^f = \sum_{l=1}^r k_l A_l^f = \frac{1}{2^{n-1}} \sum_{l=1}^r k_l \sum_{j=1}^{r-l+1} (-1)^{j-1} 2^{n-\sum_{i=1}^{j+l-1} k_i} \quad (5.6)$$

Next, we analyze the formulas in Theorem 5.2.

**Corollary 5.2.** *If  $n \geq 3$ , then  $A_1^f > A_2^f > \cdots > A_r^f$  and  $\frac{n}{2^{n-1}} \leq s^f < 2 - \frac{1}{2^{n-2}}$ .*

*Proof.*

$$\begin{aligned}
 A_l^f &= \frac{1}{2^{n-1}} \sum_{j=1}^{r-l+1} (-1)^{j-1} 2^{n-\sum_{i=1}^{j+l-1} k_i} \\
 &= \frac{1}{2^{n-1}} (2^{n-k_1-\dots-k_l} - 2^{n-k_1-\dots-k_{l+1}} + \dots + (-1)^{r-l})
 \end{aligned}$$

Since the sum is an alternating decreasing sequence and  $k_{l+1} \geq 1$ , we have

$$\begin{aligned}
 \frac{1}{2^{n-1}} (2^{n-k_1-\dots-k_{l-1}}) &\leq \frac{1}{2^{n-1}} (2^{n-k_1-\dots-k_l} - 2^{n-k_1-\dots-k_{l+1}}) \\
 &< A_l^f < \frac{1}{2^{n-1}} (2^{n-k_1-\dots-k_l})
 \end{aligned}$$

Hence,

$$A_{l+1}^f < \frac{1}{2^{n-1}} (2^{n-k_1-\dots-k_{l+1}})$$

$$\leq \frac{1}{2^{n-1}}(2^{n-k_1-\dots-k_{l-1}}) < A_l^f.$$

We have

$$\begin{aligned} k_1 A_1^f &= \frac{k_1}{2^{n-1}}(2^{n-k_1} - 2^{n-k_1-k_2} + \dots (-1)^{r-1}) \\ k_2 A_2^f &= \frac{k_2}{2^{n-1}}(2^{n-k_1-k_2} - 2^{n-k_1-k_2-k_3} + \dots (-1)^{r-2}) \\ &\dots\dots\dots \\ k_l A_l^f &= \frac{k_l}{2^{n-1}}(2^{n-k_1-\dots-k_l} - 2^{n-k_1-\dots-k_l-k_{l+1}} - \dots (-1)^{r-l}) \\ &\dots \\ k_r A_r^f &= \frac{k_r}{2^{n-1}} \end{aligned}$$

Hence,  $s^f = \sum_{l=1}^r k_l A_l^f \geq \frac{k_1}{2^{n-1}} + \frac{k_2}{2^{n-1}} + \dots + \frac{k_r}{2^{n-1}} = \frac{n}{2^{n-1}}$ , so we know that the NCFs with layer number 1 have minimal average sensitivity. On the other hand,

$$s^f = \sum_{l=1}^r k_l A_l^f < \frac{k_1}{2^{n-1}} 2^{n-k_1} + \frac{k_2}{2^{n-1}} 2^{n-k_1-k_2} + \dots + \frac{k_l}{2^{n-1}} 2^{n-k_1-\dots-k_l} + \dots + \frac{k_r}{2^{n-1}} = U(k_1, \dots, k_r),$$

where  $k_1 + \dots + k_r = n$ ,  $k_i \geq 1$ ,  $i = 1, \dots, r-1$  and  $k_r \geq 2$ . We will find the maximal value of  $U(k_1, \dots, k_r)$  in the following.

First, we claim  $k_r = 2$  if  $U(k_1, \dots, k_r)$  reaches its maximal value. Because, if  $k_r$  is increased by 1, and the last term makes  $\frac{1}{2^{n-1}}$  more contributions to  $U(k_1, \dots, k_r)$ , then there exists  $l$  such that  $k_l$  will be decreased by 1 ( $k_1 + \dots + k_r = n$ ), hence

$$\frac{k_l}{2^{n-1}} 2^{n-k_1-\dots-k_l}$$

will be decreased more than  $\frac{1}{2^{n-1}}$ . Now, it is obvious that  $\frac{k_1}{2^{n-1}} 2^{n-k_1}$  attains its maximal value only when  $k_1 = 1$  or 2, but  $k_1 = 1$  will be the choice since it also make all the other terms greater. Likewise,  $\frac{k_2}{2^{n-1}} 2^{n-k_1-k_2}$  attains its maximal value when  $k_1 = k_2 = 1$  or  $k_1 = 1$  and  $k_2 = 2$ ; again,  $k_2 = 1$  is the best choice to make all the other terms greater. In general, if  $k_1 = \dots = k_{l-1} = 1$ , then  $\frac{k_l}{2^{n-1}} 2^{n-k_1-\dots-k_l}$  attains its maximal value when  $k_l = 1$ , where  $1 \leq l \leq r-1$ . In summary, we have showed that  $U(k_1, \dots, k_r)$  reaches its maximal value when  $r = n-1$ ,  $k_1 = \dots = k_{n-2} = 1$ ,  $k_{n-1} = 2$  and

$$\text{Max}[U(k_1, \dots, k_r) = U(1, \dots, 1, 2)] = \frac{1}{2^{n-1}}(2^{n-1} + 2^{n-2} + \dots + 2^2 + 2) = 2 - \frac{1}{2^{n-2}}.$$

□

**Remark 5.3.** *We know that the average sensitivity is bounded by constants for any NCF with any number of variables. Since the minimal value approaches 0 and the maximal value of  $U(k_1, \dots, k_r)$  approaches 2 as  $n \rightarrow \infty$ . Hence,  $0 < s^f < 2$  for any NCF with an arbitrary number of variables.*

In the following, we evaluate Equation 5.6 for some parameters  $k_1, \dots, k_r$ .

- Lemma 5.4.** (1) If  $r = n - 1, k_1 = \dots = k_{n-2} = 1, k_{n-1} = 2$ , then  $s^f = \frac{4}{3} - \frac{3+(-1)^n}{3 \times 2^n}$ .
- (2) Given  $n \geq 4, r = n - 2, k_1 = \dots = k_{n-3} = 1, k_{n-2} = 3$ , then  $s^f = \frac{4}{3} - \frac{9+5(-1)^{n-1}}{3 \times 2^n}$ .
- (3) If  $n$  is even and  $n \geq 6, r = \frac{n}{2}, k_1 = 1, k_2 = \dots = k_{\frac{n}{2}-1} = 2, k_{\frac{n}{2}} = 3$ , then  $s^f = \frac{4}{3} - \frac{4}{3 \times 2^n}$ . Hence, these three cardinalities are equal if  $n$  is even.

*Proof.* When  $r = n - 1$ , then  $k_1 = \dots = k_{n-2} = 1, k_{n-1} = 2$  by Equation 5.6. We have

$$\begin{aligned} s^f &= \sum_{l=1}^r k_l A_l^f = \frac{1}{2^{n-1}} \sum_{l=1}^{n-1} k_l \sum_{j=1}^{n-l} (-1)^{j-1} 2^{n-\sum_{i=1}^{j+l-1} k_i} \\ &= \frac{1}{2^{n-1}} \sum_{l=1}^{n-1} k_l \left( \sum_{j=1}^{n-l-1} (-1)^{j-1} 2^{n-j-l+1} + (-1)^{n-l-1} \right) \\ &= \frac{1}{2^{n-1}} \sum_{l=1}^{n-1} k_l \left( \frac{1}{3} 2^{n-l+1} + \frac{1}{3} (-1)^{n-l} \right) \\ &= \frac{1}{2^{n-1}} \left( \sum_{l=1}^{n-2} \left( \frac{1}{3} 2^{n-l+1} + \frac{1}{3} (-1)^{n-l} \right) + 2 \right) = \frac{4}{3} - \frac{3+(-1)^n}{3 \times 2^n} \end{aligned}$$

The other two formulas are also routine applications of Equation 5.6.  $\square$

Based on our numerical calculations, Lemma 5.4, and the proof of Corollary 5.2, we can make the following conjecture.

**Conjecture 5.5.** The maximal value of  $s^f$  is  $s^f = \frac{4}{3} - \frac{3+(-1)^n}{3 \times 2^n}$ . It will be reached if the NCF has the maximal layer number  $n - 1$ , i.e., if  $r = n - 1, k_1 = \dots = k_{n-2} = 1, k_{n-1} = 2$ . When  $n$  is even, this maximal value is also reached by an NCF with parameters  $n \geq 4, r = n - 2, k_1 = \dots = k_{n-3} = 1, k_{n-2} = 3$  or  $n \geq 6, r = \frac{n}{2}, k_1 = 1, k_2 = \dots = k_{\frac{n}{2}-1} = 2$  and  $k_{\frac{n}{2}} = 3$ .

**Remark 5.6.** When  $n = 6$ , the NCF with  $k_1 = 1, k_2 = 2, k_3 = 1$ , and  $k_4 = 2$  also has the maximal average sensitivity  $\frac{21}{16}$ . But this can not be generalized. If the above conjecture is true, then we have  $0 < s^f < \frac{4}{3}$  for any NCF with an arbitrary number of variables. In other words, both 0 and  $\frac{4}{3}$  are uniform tight bounds for any NCF.

## 6. SIMULATIONS OF NETWORK DYNAMICS

The sensitivity of functions [39] has been shown to be a good indicator of the stability of dynamical networks constructed. We have generated random networks controlled by nested canalizing functions with fixed layer number  $m$ , where  $m = 1, \dots, n - 1$ . For our simulations we followed a similar approach as in [39]. Starting with a Boolean network  $F$ , we sample pairs of random states  $x$  and  $y$  for  $F$ . Let  $H(t)$  be the Hamming distance of  $x$  and




FIGURE 1. Derrida plots for different layer numbers. Here,  $n$  is the number of nodes and  $k$  is the connectivity. For each layer number, a Derrida plot was generated from 4096 random networks. The  $x$ -axis represents Hamming distance of pairs of states, and the  $y$ -axis represents Hamming distance of their images.

$y$ , i.e.  $H(t)$  is the number of bits in which  $x$  and  $y$  differ, and let  $H(t+1)$  be the Hamming distance of  $F(x)$  and  $F(y)$ , i.e. the Hamming distance of the successor states of  $x$  and  $y$ . A Derrida curve [13] is a plot of  $H(t+1)$  against  $H(t)$  for all possible Hamming distances. Figure 1 shows Derrida plots for different layer numbers. Each Derrida curve was generated from 4096 random networks by taking the average Hamming distances.

As can be observed from Figure 1, networks made up of nested canalizing functions of fixed layer number equal to 1 are significantly more stable than networks constructed from nested canalizing functions with higher layer numbers. It should be noted that the class of NCFs with layer number equal to 1 is the same family as the AND-NOT networks [45] or the class of extended monomials. Similarly, networks made up of nested canalizing functions with fixed layer number equal to 2 are more stable than networks constructed from nested canalizing functions with higher layer numbers, and so on. Figure 1 shows that as the layer number increases networks becomes less stable. This matches our results in Section 5.



## 7. CONCLUSION

Nested canalizing Boolean functions were inspired by structural and dynamic features of biological networks. In this study we took a careful look at the computational properties of Boolean functions and of Boolean networks constructed from them. The main tool for our analysis is a particular polynomial normal form of the Boolean functions in question. In particular, we introduced a new invariant for nested canalizing functions, their layer number. Using it, we obtain an explicit formula of the number of nested canalizing functions, which improves on the known recursive formula. Based on the polynomial form, we also obtain a formula for the Hamming weight of an NCF. The activity number of each variable of a NCF is also provided with an explicit formula. An important result we obtain is that the average sensitivity of any NCF is less than 2, which provides a theoretical argument why nested canalizing functions exhibit robust dynamic properties. This leads us to conjecture that the tight upper bound for the average sensitivity of any NCF is  $\frac{4}{3}$ .

It should be noticed that all the variables in the first layer of an NCF are canalizing variables (the most dominant), all the variables in the second layer are the second most dominant, etc. The fact that networks with low layer number are more stable than those with high layer number could be a consequence of this observation, since lower layer number means more dominant variables. The most extreme examples are: when the layer number is equal to 1, we have proved that such a function has minimal average sensitivity, therefore we also conjecture that the function with layer number equal to  $n - 1$  (the maximal layer number) has maximal average sensitivity. Hence, it should be reasonable that the corresponding networks show similar behavior.

## REFERENCES

- [1] Kazuyuki Amano, "Tight bounds on the average sensitivity of k-CNF," *Theory of Computing*, Vol 7 (2011), pp. 45-48.
- [2] E. Balleza, E. R. Alvarez-Buylla, A. Chaos, S. Kauffman, I. Shmulevich, and M. Aldana, "Critical dynamics in genetic regulatory networks: Examples from four kingdoms," *PLoS ONE*, 3 (2008), p. e2456
- [3] C. Barrett, C. Herring, J. Reed, and B. Palsson, "The global transcriptional regulatory network for metabolism in *Escherichia coli* exhibits few dominant functional states," *Proc Natl Acad Sci USA*, 102 (2005), pp. 19103-19108.
- [4] E. A. Bender, J. T. Butler, "Asymptotic approximations for the number of fanout-tree functions," *IEEE Trans. Comput.* 27 (12) (1978) 1180-1183.
- [5] A. Bernasconi, "Mathematical techniques for the analysis of Boolean functions," Ph.D. thesis, Dipartimento di Informatica, Universita di Pisa (March, 1998).
- [6] A. Bernasconi, "Sensitivity vs. block sensitivity (an average-case study)" *Information processing letters* 59 (1996) 151-157.
- [7] Ravi B. Boppana, "The average sensitivity of bounded-depth circuits" *Information processing letters* 63 (1997) 257-261.
- [8] J. T. Butler, T. Sasao, and M. Matsuura, "Average path length of binary decision diagrams," *IEEE Transactions on Computers*, 54 (2005), pp. 1041-1053.

- [9] David Canright, Sugata Gangopadhyay, Subhamoy Maitra, Pantelimon Stănică, “Laced Boolean functions and subset sum problems in finite fields,” *Discrete applied mathematics*, 159 (2011), pp. 1059-1069.
- [10] Shijian Chen and Yiguang Hong, “Control of random Boolean networks via average sensitivity of Boolean functions,” *Chin. Phys. B* Vol. 20, No 3 (2011) 036401.
- [11] Demetres Christofides, “Influences of Monotone Boolean Functions,” Preprint 2009.
- [12] S. A. Cook, C. Dwork, R. Reischuk, “Upper and lower time bounds for parallel random access machines without simultaneous writes,” *SIAM J. Comput*, 15 (1986), pp. 87-89.
- [13] B. Derrida and Y. Pomeau, “Random networks of automata: a simple annealed approximation”, *Europhys. Lett.*, 1:45-49, 1986.
- [14] S. E. Harris, B. K. Sawhill, A. Wuensche, and S. Kauffman, “A model of transcriptional regulatory networks based on biases in the observed regulation rules,” *Complex*, 7 (2002), pp. 2340.
- [15] M. Herrgard, B. Lee, V. Portnoy, and B. Palsson, “Integrated analysis of regulatory and metabolic networks reveals novel regulatory mechanisms in *saccharomyces cerevisiae*,” *Genome Res*, 16 (2006), pp. 627635.
- [16] A. Jarrah, R. Laubenbacher, and A. Veliz-Cuba, “A polynomial framework for modeling and analyzing logical models.” In Preparation, 2008.
- [17] A. Jarrah, B. Ropasa and R. Laubenbacher, “Nested canalizing, Unate Cascade, and Polynomial Functions”, *Physica D* 233 (2007), pp. 167-174.
- [18] Winfried Just, “The steady state system problem is NP-hard even for monotone quadratic Boolean dynamical systems” Preprint, 2006
- [19] Winfried Just, Ilya Shmulevich, John Konvalina, “The number and probability of canalizing functions”, *Physica D* 197 (2004), pp. 211-221.
- [20] S. A. Kauffman, “The Origins of Order: Self-Organization and Selection in Evolution”, *Oxford University Press, New York, Oxford* (1993).
- [21] S. A. Kauffman, C. Peterson, B. Samuelsson, C. Troein, “Random Boolean Network Models and the Yeast Transcription Network”, *Proc. Natl. Acad. Sci* 100 (25) (2003), pp. 14796-14799.
- [22] S. A. Kauffman, C. Peterson, B. Samuelsson, and C. Troein, “Genetic networks with canalizing Boolean rules are always stable,” , *PNAS*, 101 (2004), pp. 1710217107.
- [23] N. Keller and H. Pilpel, “Linear transformations on monotone functions on the discrete cube,” *Discrete Math.* 309 (2009), 4210-4214.
- [24] R. Laubenbacher and B. Pareigis, “Equivalence relations on finite dynamical systems,” , *Advances in applied mathematics* 26 (2001), pp. 237-251.
- [25] W. Liu, H. Lähdedmäki, Edward R. Dougherty and I. Shmulevich, “Inference of Boolean Networks Using Sensitivity Regularization,” , *EURASIP Journal of Bioinformatics and System Biology* Volume 2008, Article ID 780541, 12 pages.
- [26] Lori Layne, “Biologically Relevant Classes of Boolean Functions”, *Ph.D Thesis, Clemson University* (2011).
- [27] Jiyou Li, “On the average sensitivity of the weighted sum function,” arXiv:1108.3198v2 [cs.IT] 18 Aug 2011.
- [28] Yuan Li, “Results on Rotation Symmetric Polynomials Over  $GF(p)$ ”, *Information Sciences* 178 (2008), pp. 280-286.
- [29] Yuan Li, David Murrugarra, John O Adeyeye and Reinhard Laubenbacher “Multi-State Canalizing Functions over Finite Fields ”, <http://arxiv.org/pdf/1110.6481v1.pdf>
- [30] R. Lidl and H. Niederreiter, “Finite Fields”, *Cambridge University Press, New York* (1977).
- [31] A. A. Moreira and L. A. Amaral, “Canalizing Kauffman networks: Nonergodicity and its effect on their critical behavior,” , *Phys. Rev. Lett.* 94 (21) (2005), 218702.

- [32] D. Murrugarra and R. Laubenbacher, “Regulatory patterns in molecular interaction networks,” (2011), *Journal of Theoretical Biology*, 288, 66-72.
- [33] D. Murrugarra and R. Laubenbacher, “The number of multistate nested canalizing functions,” (2012), *Physica D: Nonlinear Phenomena*, 241, 921-938.
- [34] S. Nikolajewaa, M. Friedela, and T. Wilhelm, “Boolean networks with biologically relevant rules show ordered behaviorstar, open,” *Biosystems*, 90 (2007), pp. 4047.
- [35] N. Nisan, “CREW PRAMs and decision tree,” *SIAM J. Comput.*, 20 (6) (1991), PP. 999-1070.
- [36] Xiaoning Qian and Edward R. Dougherty, “A comparative study on sensitivits of Boolean networks,” 978-1-61284-792-4/10 2011 IEEE.
- [37] T. Sasao, K. Kinoshita, “On the number of fanout-tree functions and unate cascade functions,” *IEEE Trans. Comput.* 28 (1) (1979) 66-72.
- [38] Ilya Shmulevich, “Average sensitivity of typical monotone Boolean functions,” *arXiv:math/0507030v1 [math. Co]* 1 July 2005.
- [39] Ilya Shmulevich and Stuart A. Kauffman, “Activities and sensitivities in Boolean network models,” *Physical Review Letters*, Vol 93, Number 4 (2004), 048701.
- [40] Igor. E. Shparlinski, “Bounds on the Fourier coefficients of the weighted sum function,” *Information Processing Letters*. 103 (2007), 83-87.
- [41] Steffen Schober and Martin Bossert, “Analysis of random Boolean networks using the average sensitivity,” *arXiv: 0704.0197v1 [nlin.CG]* 2 Apr 2007.
- [42] C. H. Waddington, “Canalisation of development and the inheritance of acquired characters,” *Nature*, 150 (1942), pp. 563564.
- [43] Madars Virza, “Sensitivity versus block sensitivity of Boolean functions,” *arXiv:1008.0521v2 [cs.CC]*8 Dec 2010.
- [44] Shengyu Zhang, “Note on the average sensitivity of monotone Boolean functions,” *Preprint* 2011.
- [45] A. Veliz-Cuba, K. Buschur, R. Hamershock, A. Kniss, E. Wolff, and R. Laubenbacher, “AND-NOT decomposition of boolean network models”, *Preprint* 2012.

<sup>1</sup>DEPARTMENT OF MATHEMATICS, WINSTON-SALEM STATE UNIVERSITY, NC 27110, USA, EMAIL: LIYU@WSSU.EDU, <sup>2</sup>DEPARTMENT OF MATHEMATICS, WINSTON-SALEM STATE UNIVERSITY, NC 27110, USA, EMAIL: ADEYEY EJ@WSSU.EDU, <sup>3</sup> BIOINFORMATICS INSTITUTE, VIRGINIA TECH, BLACKSBURG, VA 24061, USA, EMAIL: DAVIDMUR@VT.EDU, <sup>4</sup>DEPARTMENT OF COMPUTER SCIENCE, VIRGINIA TECH BLACKSBURG, VA 24061-0123, USA, EMAIL: BAGUILAR@VT.EDU, <sup>5</sup> BIOINFORMATICS INSTITUTE, VIRGINIA TECH, BLACKSBURG, VA 24061, USA, EMAIL: REINHARD@VBI.VT.EDU