R.R. Kamalian^a*

^aInstitute for Informatics and Automation Problems, National Academy of Sciences of RA, 0014 Yerevan, Republic of Armenia

A proper edge t-coloring of a graph G is a coloring of edges of G with colors 1, 2, ..., t such that each of t colors is used, and adjacent edges are colored differently. The set of colors of edges incident with a vertex x of G is called a spectrum of x. A proper edge t-coloring of a graph G is interval for its vertex x if the spectrum of x is an interval of integers. A proper edge t-coloring of a graph G is persistent-interval for its vertex x if the spectrum of x is an interval of integers beginning from the color 1.

For graphs G from some classes of graphs, we obtain estimates for the possible number of vertices for which a proper edge t-coloring of G can be interval or persistent-interval.

1. Introduction

We consider undirected, simple, finite, connected graphs. For a graph G, we denote by V(G) and E(G) the sets of its vertices and edges, respectively. For any $x \in V(G)$, $d_G(x)$ denotes the degree of the vertex x in G. For a graph G, we denote by $\Delta(G)$ the maximum degree of a vertex of G. A function $\varphi: E(G) \to \{1, 2, ..., t\}$ is called a proper edge t-coloring of a graph G if each of t colors is used, and adjacent edges are colored differently. The set of all proper edge t-colorings of G is denoted by $\alpha(G, t)$. The minimum value of t for which there exists a proper edge t-coloring of a graph G is called a chromatic index [22] of G and is denoted by $\chi'(G)$. Let us also define the set $\alpha(G)$ of all proper edge colorings of the graph G

$$\alpha(G) \equiv \bigcup_{t=\chi'(G)}^{|E(G)|} \alpha(G,t).$$

If G is a graph, $\varphi \in \alpha(G)$, $x \in V(G)$, then the set of colors of edges of G incident with x is called a spectrum of the vertex x in the coloring φ of the graph G and is denoted by $S_G(x,\varphi)$.

An arbitrary nonempty subset of consecutive integers is called an interval. An interval with the minimum element p and the maximum element q is denoted by [p,q]. An interval D is called a h-interval if |D| = h.

^{*}email: rrkamalian@yahoo.com

For any real number ξ , we denote by $\lfloor \xi \rfloor$ ($\lceil \xi \rceil$) the maximum (minimum) integer which is less (greater) than or equal to ξ .

If G is a graph, $\varphi \in \alpha(G)$, and $x \in V(G)$, then we say that φ is interval (persistent-interval) for x if $S_G(x,\varphi)$ is a $d_G(x)$ -interval (a $d_G(x)$ -interval with 1 as its minimum element). For an arbitrary graph G and any $\varphi \in \alpha(G)$, we denote by $f_{G,i}(\varphi)(f_{G,pi}(\varphi))$ the number of vertices of the graph G for which φ is interval (persistent-interval). For any graph G, let us [17] set

$$\eta_i(G) \equiv \max_{\varphi \in \alpha(G)} f_{G,i}(\varphi), \quad \eta_{pi}(G) \equiv \max_{\varphi \in \alpha(G)} f_{G,pi}(\varphi).$$

A bipartite graph G with bipartition (X,Y) is called (a,b)-biregular, if $d_G(x)=a$ for any vertex $x \in X$, and $d_G(y)=b$ for any vertex $y \in Y$.

The terms and concepts that we do not define can be found in [23].

It is clear that if for any graph G $\eta_{pi}(G) = |V(G)|$, then $\chi'(G) = \Delta(G)$. For a regular graph G, these two conditions are equivalent: $\eta_{pi}(G) = |V(G)|$ iff $\chi'(G) = \Delta(G)$. It is known [15, 19] that for a regular graph G, the problem of deciding whether $\chi'(G) = \Delta(G)$ or not is NP-complete. It means that for a regular graph G, the problem of deciding whether $\eta_{pi}(G) = |V(G)|$ or not is also NP-complete. For any tree G, some necessary and sufficient condition for $\eta_{pi}(G) = |V(G)|$ was obtained in [8]. In this paper, for an arbitrary regular graph G, we obtain a lower bound for the parameter $\eta_{pi}(G)$.

If G is a graph, $R_0 \subseteq V(G)$, and the coloring $\varphi \in \alpha(G)$ is interval (persistent-interval) for any $x \in R_0$, then we say that φ is interval (persistent-interval) on R_0 .

 $\varphi \in \alpha(G)$ is called an interval coloring of a graph G if φ is interval on V(G).

We define the set \mathfrak{N} as the set of all graphs for which there is an interval coloring. Clearly, for any graph $G, G \in \mathfrak{N}$ if and only if $\eta_i(G) = |V(G)|$.

The notion of an interval coloring was introduced in [6]. In [6, 16, 7] it is shown that if $G \in \mathfrak{N}$, then $\chi'(G) = \Delta(G)$. For a regular graph G, these two conditions are equivalent: $G \in \mathfrak{N}$ iff $\chi'(G) = \Delta(G)$ [6, 16, 7]. Consequently, for a regular graph G, four conditions are equivalent: $G \in \mathfrak{N}$, $\chi'(G) = \Delta(G)$, $\eta_i(G) = |V(G)|$, $\eta_{pi}(G) = |V(G)|$. It means that for any regular graph G,

- 1. the problem of deciding whether or not G has an interval coloring is NP-complete,
- 2. the problem of deciding whether $\eta_i(G) = |V(G)|$ or not is NP-complete.

In this paper, for an arbitrary regular graph G, we obtain a lower bound for the parameter $\eta_i(G)$.

We also obtain some results for bipartite graphs. The complexity of the problem of existence of an interval coloring for bipartite graphs is investigated in [3, 9, 21]. In [16] it is shown that for a bipartite graph G with bipartition (X,Y) and $\Delta(G)=3$ the problem of existence of a proper edge 3-coloring which is persistent-interval on $X \cup Y$ (or even only on Y [6, 16]) is NP-complete.

Suppose that G is an arbitrary bipartite graph with bipartition (X, Y). Then $\eta_i(G) \ge \max\{|X|, |Y|\}$.

Suppose that G is a bipartite graph with bipartition (X, Y) for which there exists a coloring $\varphi \in \alpha(G)$ persistent-interval on Y. Then $\eta_{pi}(G) \geq 1 + |Y|$.

Some attention is devoted to (a, b)-biregular bipartite graphs [4, 14, 13, 18] in the case b = a + 1.

We show that if G is a (k-1,k)-biregular bipartite graph, $k \geq 4$, then

$$\eta_i(G) \ge \frac{k-1}{2k-1} \cdot |V(G)| + \left\lceil \frac{k}{\left\lceil \frac{k}{2} \right\rceil \cdot (2k-1)} \cdot |V(G)| \right\rceil.$$

We show that if G is a (k-1,k)-biregular bipartite graph, $k \geq 3$, then

$$\eta_{pi}(G) \ge \frac{k}{2k-1} \cdot |V(G)|.$$

2. Results

Theorem 1 [17] If G is a regular graph with $\chi'(G) = 1 + \Delta(G)$, then

$$\eta_{pi}(G) \ge \left\lceil \frac{|V(G)|}{1 + \Delta(G)} \right\rceil.$$

Proof. Suppose that $\beta \in \alpha(G, 1 + \Delta(G))$. For any $j \in [1, 1 + \Delta(G)]$, define

$$V_{G,\beta,j} \equiv \{x \in V(G)/j \notin S_G(x,\beta)\}.$$

For arbitrary integers j', j'', where $1 \le j' < j'' \le 1 + \Delta(G)$, we have

$$V_{G,\beta,j'} \cap V_{G,\beta,j''} = \emptyset$$

and

$$\bigcup_{j=1}^{1+\Delta(G)} V_{G,\beta,j} = V(G).$$

Hence, there exists $j_0 \in [1, 1 + \Delta(G)]$ for which

$$|V_{G,\beta,j_0}| \ge \left\lceil \frac{|V(G)|}{1 + \Delta(G)} \right\rceil.$$

Set $R_0 \equiv V_{G,\beta,j_0}$.

Case 1 $1j_0 = 1 + \Delta(G)$.

Clearly, β is persistent-interval on R_0 .

Case 2 $2 j_0 \in [1, \Delta(G)].$

Define a function $\varphi: E(G) \to [1, 1 + \Delta(G)]$. For any $e \in E(G)$, set:

$$\varphi(e) \equiv \begin{cases} \beta(e), & \text{if } \beta(e) \notin \{j_0, 1 + \Delta(G)\} \\ j_0, & \text{if } \beta(e) = 1 + \Delta(G) \\ 1 + \Delta(G), & \text{if } \beta(e) = j_0. \end{cases}$$

It is not difficult to see that $\varphi \in \alpha(G, 1 + \Delta(G))$ and φ is persistent-interval on R_0 .

Corollary 1 [17] If G is a cubic graph, then there exists a coloring from $\alpha(G, \chi'(G))$ which is persistent-interval for at least $\lceil \frac{|V(G)|}{4} \rceil$ vertices of G.

Theorem 2 [17] If G is a regular graph with $\chi'(G) = 1 + \Delta(G)$, then

$$\eta_i(G) \ge \left\lceil \frac{|V(G)|}{\left\lceil \frac{1+\Delta(G)}{2} \right\rceil} \right\rceil.$$

Proof. Suppose that $\beta \in \alpha(G, 1 + \Delta(G))$. For any $j \in [1, 1 + \Delta(G)]$, define

$$V_{G,\beta,j} \equiv \{x \in V(G)/j \notin S_G(x,\beta)\}.$$

For arbitrary integers j', j'', where $1 \le j' < j'' \le 1 + \Delta(G)$, we have

$$V_{G,\beta,j'} \cap V_{G,\beta,j''} = \emptyset$$

and

$$\bigcup_{j=1}^{L+\Delta(G)} V_{G,\beta,j} = V(G).$$

For any $i \in [1, \lceil \frac{1+\Delta(G)}{2} \rceil]$, let us define the subset $V(G, \beta, i)$ of the set V(G) as follows:

$$V(G,\beta,i) \equiv \begin{cases} V_{G,\beta,2i-1} \cup V_{G,\beta,2i}, & \text{if } \Delta(G) \text{ is odd and } i \in [1,\frac{1+\Delta(G)}{2}] \\ & \text{or } \Delta(G) \text{ is even and } i \in [1,\frac{\Delta(G)}{2}], \\ V_{G,\beta,1+\Delta(G)}, & \text{if } \Delta(G) \text{ is even and } i = 1 + \frac{\Delta(G)}{2}. \end{cases}$$

For arbitrary integers i', i'', where $1 \le i' < i'' \le \left\lceil \frac{1 + \Delta(G)}{2} \right\rceil$, we have

$$V(G, \beta, i') \cap V(G, \beta, i'') = \emptyset$$

and

$$\bigcup_{i=1}^{\left[\frac{1+\Delta(G)}{2}\right]} V(G,\beta,i) = V(G).$$

Hence, there exists $i_0 \in \left[1, \left\lceil \frac{1+\Delta(G)}{2} \right\rceil \right]$ for which

$$|V(G, \beta, i_0)| \ge \left\lceil \frac{|V(G)|}{\left\lceil \frac{1+\Delta(G)}{2} \right\rceil} \right\rceil.$$

Set $R_0 \equiv V(G, \beta, i_0)$.

Case 3
$$1 i_0 = \left\lceil \frac{1 + \Delta(G)}{2} \right\rceil$$
.

Case 4 1.a $\Delta(G)$ is even. Clearly, β is interval on R_0 .

Case 5 1.b $\Delta(G)$ is odd.

Define a function $\varphi: E(G) \to [1, 1 + \Delta(G)]$. For any $e \in E(G)$, set:

$$\varphi(e) \equiv \begin{cases} (\beta(e) + 1)(\operatorname{mod}(1 + \Delta(G))), & \text{if } \beta(e) \neq \Delta(G), \\ 1 + \Delta(G), & \text{if } \beta(e) = \Delta(G). \end{cases}$$

It is not difficult to see that $\varphi \in \alpha(G, 1 + \Delta(G))$ and φ is interval on R_0 .

Case 6 2 $1 \le i_0 \le \lceil \frac{\Delta(G)-1}{2} \rceil$. Define a function $\varphi : E(G) \to [1, 1 + \Delta(G)]$. For any $e \in E(G)$, set:

$$\varphi(e) \equiv \begin{cases} (\beta(e) + 2 + \Delta(G) - 2i_0) (\bmod(1 + \Delta(G))), & \text{if } \beta(e) \neq 2i_0 - 1, \\ 1 + \Delta(G), & \text{if } \beta(e) = 2i_0 - 1. \end{cases}$$

It is not difficult to see that $\varphi \in \alpha(G, 1 + \Delta(G))$ and φ is interval on R_0 .

Corollary 2 [17] If G is a cubic graph, then there exists a coloring from $\alpha(G, \chi'(G))$ which is interval for at least $\frac{|V(G)|}{2}$ vertices of G.

Theorem 3 [6, 16, 7] Let G be a bipartite graph with bipartition (X,Y). Then there exists a coloring $\varphi \in \alpha(G, |E(G)|)$ which is interval on X.

Corollary 3 Let G be a bipartite graph with bipartition (X,Y). Then $\eta_i(G) \geq \max\{|X|,|Y|\}$.

Theorem 4 [1, 6, 7] Let G be a bipartite graph with bipartition (X,Y) where $d_G(x) < 0$ $d_G(y)$ for each edge $(x,y) \in E(G)$ with $x \in X$ and $y \in Y$. Then there exists a coloring $\varphi_0 \in \alpha(G, \Delta(G))$ which is persistent-interval on Y.

Theorem 5 Suppose G is a bipartite graph with bipartition (X,Y), and there exists a coloring $\varphi_0 \in \alpha(G, \Delta(G))$ which is persistent-interval on Y. Then, for an arbitrary vertex $x_0 \in X$, there exists $\psi \in \alpha(G, \Delta(G))$ which is persistent-interval on $\{x_0\} \cup Y$.

Proof. Case 7 1 $S_G(x_0, \varphi_0) = [1, d_G(x_0)]$. In this case ψ is φ_0 .

Case 8 2 $S_G(x_0, \varphi_0) \neq [1, d_G(x_0)].$

Clearly, $[1, d_G(x_0)] \setminus S_G(x_0, \varphi_0) \neq \emptyset$, $S_G(x_0, \varphi_0) \setminus [1, d_G(x_0)] \neq \emptyset$. Since $|S_G(x_0, \varphi_0)| = 0$ $|[1,d_G(x_0)]|=d_G(x_0)$, there exists $\nu_0\in[1,d_G(x_0)]$ satisfying the condition $|[1,d_G(x_0)]\setminus S_G(x_0,\varphi_0)|=$ $|S_G(x_0,\varphi_0)\backslash[1,d_G(x_0)]|=\nu_0.$

Now let us construct the sequence $\Theta_0, \Theta_1, \ldots, \Theta_{\nu_0}$ of proper edge $\Delta(G)$ -colorings of the graph G, where for any $i \in [0, \nu_0]$, Θ_i is persistent-interval on Y.

Set $\Theta_0 \equiv \varphi_0$.

Suppose that for some $k \in [0, \nu_0 - 1]$, the subsequence $\Theta_0, \Theta_1, \ldots, \Theta_k$ is already constructed.

Let

$$t_k \equiv \max(S_G(x_0, \Theta_k) \setminus [1, d_G(x_0)]),$$

$$s_k \equiv \min([1, d_G(x_0)] \setminus S_G(x_0, \Theta_k)).$$

Clearly, $t_k > s_k$. Consider the path P(k) in the graph G of maximum length with the initial vertex x_0 whose edges are alternatively colored by the colors t_k and s_k . Let Θ_{k+1} is obtained from Θ_k by interchanging the two colors t_k and s_k along P(k).

It is not difficult to see that Θ_{ν_0} is persistent-interval on $\{x_0\} \cup Y$. Set $\psi \equiv \Theta_{\nu_0}$.

Corollary 4 Let G be a bipartite graph with bipartition (X,Y) where $d_G(x) \leq d_G(y)$ for each edge $(x,y) \in E(G)$ with $x \in X$ and $y \in Y$. Let x_0 be an arbitrary vertex of X. Then there exists a coloring $\varphi_0 \in \alpha(G,\Delta(G))$ which is persistent-interval on $\{x_0\} \cup Y$.

Corollary 5 [17] Let G be a bipartite graph with bipartition (X, Y) where $d_G(x) \leq d_G(y)$ for each edge $(x, y) \in E(G)$ with $x \in X$ and $y \in Y$. Then $\eta_{pi}(G) \geq 1 + |Y|$.

Remark 1 Notice that the complete bipartite graph $K_{n+1,n}$ for an arbitrary positive integer n satisfies the conditions of Corollary 5. Is is not difficult to see that $\eta_{pi}(K_{n+1,n}) = 1 + n$. It means that the bound obtained in Corollary 5 is sharp since in this case |Y| = n.

Remark 2 Let G be a bipartite (k-1,k)-biregular graph with bipartition (X,Y), where $k \geq 3$. Then the numbers $\frac{|X|}{k}$, $\frac{|Y|}{k-1}$, and $\frac{|V(G)|}{2k-1}$ are integer. It follows from the equalities gcd(k-1,k)=1 and $|E(G)|=|X|\cdot (k-1)=|Y|\cdot k$.

Theorem 6 [17] Let G be a bipartite (k-1,k)-biregular graph, where $k \geq 4$. Then

$$\eta_i(G) \ge \frac{k-1}{2k-1} \cdot |V(G)| + \left\lceil \frac{k}{\left\lceil \frac{k}{2} \right\rceil \cdot (2k-1)} \cdot |V(G)| \right\rceil.$$

Proof. Suppose that (X,Y) is a bipartition of G. Clearly, $\chi'(G) = \Delta(G) = k$. Suppose that $\beta \in \alpha(G,k)$. For any $j \in [1,k]$, define:

$$V_{G,\beta,j} \equiv \{x \in X/j \not\in S_G(x,\beta)\}.$$

For arbitrary integers j', j'', where $1 \le j' < j'' \le k$, we have

$$V_{G,\beta,j'} \cap V_{G,\beta,j''} = \emptyset$$

and

$$\bigcup_{j=1}^{k} V_{G,\beta,j} = X.$$

For any $i \in [1, \lceil \frac{k}{2} \rceil]$, let us define the subset $V(G, \beta, i)$ of the set X as follows:

$$V(G,\beta,i) \equiv \begin{cases} V_{G,\beta,2i-1} \cup V_{G,\beta,2i}, & \text{if } k \text{ is odd and } i \in [1,\frac{k-1}{2}] \\ & \text{or } k \text{ is even and } i \in [1,\frac{k}{2}], \\ V_{G,\beta,k}, & \text{if } k \text{ is odd and } i = \frac{1+k}{2}. \end{cases}$$

For arbitrary integers i', i'', where $1 \le i' < i'' \le \lceil \frac{k}{2} \rceil$, we have

$$V(G,\beta,i')\cap V(G,\beta,i'')=\emptyset$$

and

$$\bigcup_{i=1}^{\left[\frac{k}{2}\right]} V(G,\beta,i) = X.$$

Hence, there exists $i_0 \in \left[1, \left\lceil \frac{k}{2} \right\rceil \right]$ for which

$$|V(G, \beta, i_0)| \ge \left\lceil \frac{|X|}{\left\lceil \frac{k}{2} \right\rceil} \right\rceil.$$

Set $R_0 \equiv Y \cup V(G, \beta, i_0)$.

It is not difficult to verify that

$$|R_0| \ge \frac{k-1}{2k-1} \cdot |V(G)| + \left\lceil \frac{k}{\left\lceil \frac{k}{2} \right\rceil \cdot (2k-1)} \cdot |V(G)| \right\rceil.$$

Case 9 1 $i_0 = \lceil \frac{k}{2} \rceil$.

Case 10 1.a k is odd.

Clearly, β is interval on R_0 .

Case 11 1.b k is even.

Define a function $\varphi: E(G) \to [1,k]$. For any $e \in E(G)$, set:

$$\varphi(e) \equiv \left\{ \begin{array}{ll} (\beta(e)+1)(\bmod k), & \text{if } \beta(e) \neq k-1, \\ k, & \text{if } \beta(e) = k-1. \end{array} \right.$$

It is not difficult to see that $\varphi \in \alpha(G, k)$ and φ is interval on R_0 .

Case 12 $2 i_0 \in [1, \lceil \frac{k}{2} \rceil - 1]$. Define a function $\varphi : E(G) \to [1, k]$. For any $e \in E(G)$, set:

$$\varphi(e) \equiv \begin{cases} (\beta(e) + 1 + k - 2i_0) \pmod{k}, & \text{if } \beta(e) \neq 2i_0 - 1, \\ k, & \text{if } \beta(e) = 2i_0 - 1. \end{cases}$$

It is not difficult to see that $\varphi \in \alpha(G,k)$ and φ is interval on R_0 .

Corollary 6 [17] Let G be a bipartite (k-1,k)-biregular graph, where k is even and $k \geq 4$. Then

$$\eta_i(G) \ge \frac{k+1}{2k-1} \cdot |V(G)|.$$

Corollary 7 [17] Let G be a bipartite (3,4)-biregular graph. Then there exists a coloring from $\alpha(G,4)$ which is interval for at least $\frac{5}{7}|V(G)|$ vertices of G.

Remark 3 For an arbitrary bipartite graph G with $\Delta(G) \leq 3$, there exists an interval coloring of G [12, 10, 11]. Consequently, if G is a bipartite (2,3)-biregular graph, then $\eta_i(G) = |V(G)|.$

Remark 4 Some sufficient conditions for existence of an interval coloring of a (3,4)biregular bipartite graph were obtained in [2, 5, 20].

Theorem 7 [17] Let G be a bipartite (k-1,k)-biregular graph, where $k \geq 3$. Then

$$\eta_{pi}(G) \ge \frac{k}{2k-1} \cdot |V(G)|.$$

Proof. Suppose that (X,Y) is a bipartition of G. Clearly, $\chi'(G) = \Delta(G) = k$. Suppose that $\beta \in \alpha(G,k)$.

For any $j \in [1, k]$, define:

$$V_{G,\beta,j} \equiv \{x \in X/j \notin S_G(x,\beta)\}.$$

For arbitrary integers j', j'', where $1 \le j' < j'' \le k$, we have

$$V_{G,\beta,j'} \cap V_{G,\beta,j''} = \emptyset$$

and

$$\bigcup_{j=1}^{k} V_{G,\beta,j} = X.$$

Hence, there exists $j_0 \in [1, k]$ for which

$$|V_{G,\beta,j_0}| \ge \frac{|X|}{k}.$$

Set $R_0 \equiv Y \cup V_{G,\beta,j_0}$.

It is not difficult to verify that

$$|R_0| \ge \frac{k}{2k-1} \cdot |V(G)|.$$

Case 13 $1 j_0 = k$.

Clearly, β is persistent-interval on R_0 .

Case 14 $2 j_0 \in [1, k-1].$

Define a function $\varphi: E(G) \to [1, k]$. For any $e \in E(G)$, set:

$$\varphi(e) \equiv \begin{cases} \beta(e), & \text{if } \beta(e) \notin \{j_0, k\} \\ j_0, & \text{if } \beta(e) = k \\ k, & \text{if } \beta(e) = j_0. \end{cases}$$

It is not difficult to see that $\varphi \in \alpha(G, k)$ and φ is persistent-interval on R_0 .

Corollary 8 [17] Let G be a bipartite (3,4)-biregular graph. Then there exists a coloring from $\alpha(G,4)$ which is persistent-interval for at least $\frac{4}{7}|V(G)|$ vertices of G.

3. Acknowledgment

The author thanks professors A.S. Asratian and P.A. Petrosyan for their attention to this work.

REFERENCES

- 1. A.S. Asratian, *Investigation of some mathematical model of Scheduling Theory*, Doctoral dissertation, Moscow University, 1980 (in Russian).
- 2. A.S. Asratian, C.J. Casselgren, A sufficient condition for interval edge colorings of (4,3)-biregular bipartite graphs, Research report LiTH-MAT-R-2006-07, Linköping University, 2006.
- 3. A.S. Asratian, C.J. Casselgren, Some results on interval edge colorings of (α, β) -biregular bipartite graphs, Research report LiTH-MAT-R-2006-09, Linköping University, 2006.
- 4. A.S. Asratian, C.J. Casselgren, On interval edge colorings of (α, β) -biregular bipartite graphs, Discrete Math 307 (2007), pp. 1951–1956.
- 5. A.S. Asratian, C.J. Casselgren, J. Vandenbussche, D.B. West, *Proper path-factors and interval edge-coloring of* (3,4)-biregular bigraphs, J. of Graph Theory 61 (2009), pp. 88–97.
- 6. A.S. Asratian, R.R. Kamalian, *Interval colorings of edges of a multigraph*, Appl. Math. 5 (1987), Yerevan State University, pp. 25–34 (in Russian).
- 7. A.S. Asratian, R.R. Kamalian, *Investigation of interval edge-colorings of graphs*, Journal of Combinatorial Theory. Series B 62 (1994), no.1, pp. 34–43.
- 8. Y. Caro, J. Schönheim, Generalized 1-factorization of trees, Discrete Math 33 (1981), pp. 319–321.
- 9. K. Giaro, The complexity of consecutive Δ -coloring of bipartite graphs: 4 is easy, 5 is hard. Ars Combin. 47(1997), 287–298.
- 10. K. Giaro, Compact Task Scheduling on Dedicated Processors with no Waiting Periods, Ph.D. Thesis, Technical University of Gdańsk, ETI Faculty, Gdańsk (1999) (in Polish).
- 11. K. Giaro, M. Kubale, M. Małafiejski, Compact scheduling in open shop with zero-one time operations, INFOR 37 (1999), pp. 37–47.
- 12. H. Hansen, *Scheduling with minimum waiting periods*, Master Thesis, Odense University, Odense, Denmark, 1992 (in Danish).
- 13. D. Hanson, C.O.M. Loten, A lower bound for Interval colouring bi-regular bipartite graphs, Bulletin of the ICA 18 (1996), pp. 69–74.
- 14. D. Hanson, C.O.M. Loten, B. Toft, On interval colourings of bi-regular bipartite graphs, Ars Combin. 50 (1998), pp. 23–32.
- 15. I. Holyer, The NP-completeness of edge-coloring, SIAM J. Comput. 10 (1981), pp. 718–720.
- 16. R.R. Kamalian, *Interval Edge Colorings of Graphs*, Doctoral dissertation, the Institute of Mathematics of the Siberian Branch of the Academy of Sciences of USSR, Novosibirsk, 1990 (in Russian).
- 17. R.R. Kamalian, On a number of vertices with an interval spectrum in proper edge colorings of some graphs. Research report LiTH-MAT-R-2011/03-SE, Linköping University, 2011.
- 18. A.V. Kostochka, Unpublished manuscript, 1995.
- 19. D. Leven and Z. Galil, NP-completeness of finding the chromatic index of regular graphs, J. Algorithms 4 (1983), pp. 35–44.
- $20. \ A.V. \ Pyatkin, \ Interval \ coloring \ of \ (3,4)-biregular \ bipartite \ graphs \ having \ large \ cubic$

- subgraphs, J. of Graph Theory 47 (2004), pp. 122–128.
- 21. S.V. Sevast'janov, Interval colorability of the edges of a bipartite graph, Metody Diskret. Analiza 50(1990), 61–72 (in Russian).
- 22. V.G. Vizing, The chromatic index of a multigraph, Kibernetika 3 (1965), pp. 29–39.
- 23. D.B. West, Introduction to Graph Theory, Prentice-Hall, New Jersey, 1996.