

Formulation of the cell MAPK signaling pathway by applying Shannon entropy

Tatsuaki Tsuruyama M.D, Ph.D

*Department of Molecular Pathology, Graduate School of Medicine, Kyoto University,
Yoshida Konoe-cho, Sakyo-ku, Kyoto 606-8501, Japan*

Correspondence to: Tatsuaki Tsuruyama, Email: tsuruyam@kuhp.kyoto-u.ac.jp

Phone; +81-75-753-4473

Running title: Formulation of biological information

Key words: biological information, signal transduction, Michaelis-Menten coefficient, Shannon entropy.

Abbreviations used in this paper: ATP, adenosine triphosphate; MAPK, mitogen-activated protein kinase.

Abstract

Biological signal transmission and transduction independent of individual biological systems have not been sufficiently established. As a result, a theoretical framework of biological information is yet to be determined. Here, we report on an attempt to formulate the transmission of biological information by an ATP-binding enzyme reaction on the basis of a Michaelis-Menten reaction system by applying Shannon entropy. This formulation allows us to set preliminary equations for the transmission rate of biological information and signal information on the basis of the substrate concentration and to identify the core step of the mitogen-activated protein kinase pathway in the signaling cascade during the transmission. Evaluation of biological information by calculation from a thermodynamic viewpoint is expected.

Introduction

Cellular information required for biological cellular activity is transmitted by the intracellular signaling system. This system is caused by protein association reactions triggered by cell stimulations through ligand binding, following a change in the extracellular environment. The reaction cascade is mediated by phosphorylation of a protein. The phosphorylated protein further phosphorylates another protein via an association reaction with another protein. The final phosphorylated protein translocates into the nucleus and controls the expression of genes by binding to a specific region of DNA, such as the promoter region. The mitogen-activated protein kinase (MAPK) pathway is a highly conserved signaling cascade of three protein kinases involved in various biological processes. Although the biological responses associated with MAPK signaling highly vary, there is a basic structure to the MAPK cascade. The cascade always involves MAPK kinase kinase (MAPKKK), MAPK kinase (MAPKK), and a MAPK. MAPKKK activates MAPKK by phosphorylation at two conserved serine residues, while MAPKK activates MAPK by phosphorylation at conserved threonine and tyrosine residues. The cascade relays signals from the plasma membrane to the targets in the cytoplasm and nucleus (1). In these cases, protein phosphorylation is commonly employed for biological signal transmission.

In this paper, we will consider the general formulation of biological cellular signaling information *in vivo*. The present study is a significance attempt to depict the essence of biological reactions, and there have been only a few of such studies so far. The formulation of cellular information proposed illustrates a unique feature of the cell signaling system.

Results and Discussion

Formulation of biological Shannon entropy (information)


We consider a simple system that includes two states of the biological chemical species, c^* and c . Here we introduce the Shannon entropy I ,

$$I \approx -\left(\frac{c}{C}\right)\log_2\left(\frac{c}{C}\right) - \left(\frac{c^*}{C}\right)\log_2\left(\frac{c^*}{C}\right) \quad [1]$$

where, c and c^* also denote the concentrations of the biological chemical species, respectively (2). Meanwhile, C denotes the total concentration of the species.

$$c + c^* = C \quad [2]$$

Here, we consider a simple chemical enzymatic reaction for the formulation of biological signaling information



where, k_1 and k_2 denote the kinetic coefficients of the forward reaction and k_{-1} denotes the kinetic coefficient of the reversed reaction. S is the enzyme that has the ability to bind c . Assuming the concentration of the active species is constant and using the Michaelis-Menten coefficient K_M , c^* is given as

$$c^* = \frac{C[S]}{K_M + [S]} \quad [4]$$

where,

$$K_M = \frac{k_{-1} + k_{+2}}{k_{+1}} \quad [5]$$

Subsequently, substituting [4] and [5] into [1] gives

$$I = -\frac{[S]}{K_M + [S]} \log_2 \frac{[S]}{K_M + [S]} - \frac{K_M}{K_M + [S]} \log_2 \frac{K_M}{K_M + [S]} \quad [6]$$

Therefore, differentiating [6] gives

$$dI \approx -\log_2 \frac{[S]}{K_M + [S]} d \frac{[S]}{K_M + [S]} - \log_2 \frac{K_M}{K_M + [S]} d \frac{K_M}{K_M + [S]} \quad [7]$$

As shown in a real cellular signal cascade such as the MAPK cascade, since $[S]$ is sufficiently smaller than K_M , we can obtain the following equation with sufficient accuracy:

$$I \approx -\frac{[S]}{K_M} \left(\log_2 \frac{[S]}{K_M} \right) \quad [6']$$

In addition, the following is obtained:

$$dI \approx -\frac{1}{K_M} \log_2 \frac{[S]}{K_M} d[S] \quad [7']$$

This shows that the Shannon entropy I is defined by the logarithm of the substrate

concentration $[S]$. In the next section, we will apply this formula to real signal transduction and transmission in biological cell reaction systems.

Cascade of reactions carrying biological information

In the current study, we consider a reaction cascade in which the substrate is adenosine triphosphate, ATP, which is one of the mostly employed signal transduction molecule in the cellular reaction. This substrate is involved in the phosphorylation of various enzyme, in which ATP is added to tyrosine and other amino acid residues that have a hydroxyl group. In the phosphorylation reaction of a protein, ATP forms a complex with the protein substrate S , $[S-ATP]$. Because ATP is involved in a variety of reactions, we represent the complex in the j -th reaction with the protein S_j as $[S_j-ATP]$.



Here, * marks activation of the enzyme S_i .

Assuming the concentration of the active species is constant and using the Michaelis-Menten coefficient K_{Mj} , $[S_j-ATP]$ is given as

$$[S_j - ATP] = \frac{[ATP]_0 [S_j]}{K_{Mj} + [S_j]} \quad [9]$$

where, $[ATP]_0$ represents the total concentration of ATP. We set the total concentration of ATP in a cell system as

$$[ATP] + \sum_j [S_j - ATP] = [ATP]_0 \quad [10]$$

Using a simple calculation of [9] and [10], the following is obtained:

$$[ATP] = \frac{[ATP]_0}{1 + \sum_j [S_j] / K_{Mj}} \quad [11]$$

According to [6], the total entropy I of the reaction cascade consisting of n reactions is given by

$$I = -\frac{[ATP]}{[ATP]_0} \log_2 \frac{[ATP]}{[ATP]_0} - \sum_{j=1}^n \frac{[S_j - ATP]}{[ATP]_0} \log_2 \frac{[S_j - ATP]}{[ATP]_0} \quad [12]$$


Substituting [9] into [12] gives


$$\begin{aligned} I &= -\frac{1}{1 + \sum_j \frac{[S_j]}{K_{Mj}}} \log_2 \frac{1}{1 + \sum_j \frac{[S_j]}{K_{Mj}}} - \sum_{j=1}^n \frac{[S_j]}{K_{Mj} + [S_j]} \log_2 \frac{[S_j]}{K_{Mj} + [S_j]} \\ &\approx -\frac{1}{1 + \sum_j \frac{[S_j]}{K_{Mj}}} \log_2 \frac{1}{1 + \sum_j \frac{[S_j]}{K_{Mj}}} - \sum_{j=1}^n \frac{[S_j]}{K_{Mj}} \log_2 \frac{[S_j]}{K_{Mj}} \end{aligned} \quad [13]$$

In the above, $[S]$ is sufficiently smaller than K_M .

Actual estimation of a MAPK cascade

For simplicity, we summarize the MAPK signaling cascade as follows





Here, K_{M1} , K_{M2} , and K_{M3} , denote the Michaelis-Menten coefficients of [14]-[16], respectively. In [14]-[16], the amino acid residues, tyrosine or serine, are phosphorylated. “-P” indicates the phosphorylation.

In this cascade, the total Shannon entropy is given by

$$I \approx -\frac{1}{1 + \sum_{j=1}^3 \frac{[S_j]}{K_{Mj}}} \log_2 \frac{1}{1 + \sum_{j=1}^3 \frac{[S_j]}{K_{Mj}}} - \sum_{j=1}^3 \frac{[S_j]}{K_{Mj}} \log_2 \frac{[S_j]}{K_{Mj}} \quad [17]$$

where, S_1 , S_2 , and S_3 are MAPKKK, MAPKK, and MAPK, respectively. By inputting the actual values of the parameter and coefficients from a previous report (3), we obtain the total entropy (Table 1). Here, we introduce partial entropy as follows:

$$I_1 = \frac{\partial I}{\partial [\text{MAPKKK}]}, I_2 = \frac{\partial I}{\partial [\text{MAPKK}]}, I_3 = \frac{\partial I}{\partial [\text{MAPK}]} \quad [18]$$

By inputting the actual values of the parameter and coefficients from a previous report (3), we obtain the sum of I_1 , I_2 , and I_3 (Table 1). As mentioned above for the approximation in [6'] and [7'], the substrate enzyme concentration is significantly smaller than K_M . This indicates that the first step, phosphorylation of MAPKKK, is

dominant with respect to entropy. In fact, when the enzyme concentrations are sufficient small, partial entropies are roughly estimated as follows:

$$I_1 = \frac{\partial I}{\partial[\text{MAPKKK}]} \sim -\frac{\log_2([\text{MAPKKK}]/K_{M1})}{K_{M1}} > 0 \quad [19]$$

$$I_2 = \frac{\partial I}{\partial[\text{MAPKK}]} \sim -\frac{\log_2([\text{MAPKK}]/K_{M2})}{K_{M2}} > 0 \quad [20]$$

$$I_3 = \frac{\partial I}{\partial[\text{MAPK}]} \sim -\frac{\log_2([\text{MAPK}]/K_{M3})}{K_{M3}} > 0 \quad [21]$$

As mentioned above, molecular biology lacks a generalization method that estimates the inner cellular information transmission. As a result, the term *biological information* appears to be a metaphor or has been used without consciousness of the definition of information. However, the transmitted information is based on a protein association reaction, that is, an actual physical chemical reaction cascade. From this perspective, it was not difficult to formulate equations illustrating the cascade. It is not known yet to what extent this formulation will be applicable to estimation of the actual transmission of biological information; however, the formulation of biological information provides us a way to quantify the cell phenomenon. We hope that this method will become a tool of theoretical molecular biology.

Reference

1. Shannon, Claude E. (1951) Prediction and entropy of printed English, *The Bell System Technical Journal* 30: 50-64.
2. Menten, L. & Michaelis, M.I. (1913) Die Kinetik der Invertinwirkung, *Biochem Z* 49: 333-369.
3. Huang CY & Ferrell JE, Jr. (1996) Ultrasensitivity in the mitogen-activated protein kinase cascade. *Proc Natl Acad Sci USA* 93(19): 10078-10083.
4. Yoon J, Deisboeck TS. Investigating differential dynamics of the MAPK signaling cascade using a multi-parametric global sensitivity analysis. *PLoS One*. 2009;4:e4560.
5. Angeli D, De Leenheer P, Sontag E. Graph-theoretic characterizations of monotonicity of chemical networks in reaction coordinates. *J Math Biol*. 2010;61:581-616.
6. Blossey R, Bodart JF, Devys A, et al. Signal propagation of the MAPK cascade in *Xenopus* oocytes: role of bistability and ultrasensitivity for a mixed problem. *Journal of Mathematical Biology*. 2012;64:1-39.
7. Babu CVS, Song EJ, Yoo YS. Modeling and simulation in signal transduction pathways: a systems biology approach. *Biochimie*. 2006;88:277-283.
8. Imamura H, Nhat KPH, Togawa H, et al. Visualization of ATP levels inside single living cells with fluorescence resonance energy transfer-based genetically encoded indicators. *P Natl Acad Sci USA*. 2009; 106: 15651-15656.

Table 1. Predicted Shannon entropy in the MAPK pathway

Signaling molecule	Concentration of the substrate enzyme		
	K _m	Minimum (nM)	Maximum (nM)
MAPKKK→MAPKKK-P	60	0.0006	0.015
MAPKK→MAPKK-P	60	0.24	6
MAPK→MAPK-P	60	0.24	6
<i>I</i> =Sum of the entropy		0.075	0.851
Partial entropy (bit)			
<i>I</i> ₁		0.276	0.189
<i>I</i> ₂		0.131	0.0414
<i>I</i> ₃		0.131	0.0414

K_M is the Michaelis-Menten coefficient; concentration of the substrate enzyme is that of MAPKKK in the first step, MAPKK in the second step, and MAPK in the third step.

The unit of K_M and concentration is nM.