# COACTIONS AND SKEW PRODUCTS OF TOPOLOGICAL GRAPHS

S. KALISZEWSKI AND JOHN QUIGG

ABSTRACT. We show that the  $C^*$ -algebra of a skew-product topological graph  $E \times_{\kappa} G$  is isomorphic to the crossed product of  $C^*(E)$ by a coaction of the locally compact group G.

## 1. INTRODUCTION

In [4, Theorem 2.4] we proved that if E is a directed graph and  $\kappa$  is a function from the edges of E to a discrete group G, then the graph algebra  $C^*(E \times_{\kappa} G)$  of the skew-product graph is a crossed product of  $C^*(E)$  by a coaction of G. This was later generalized to homogeneous spaces G/H in [2, Theorem 3.4], and to higher-rank graphs in [9, Theorem 7.1]. In this paper we generalize the result to topological graphs and locally compact groups. More precisely, we prove in Theorem 3.1 that if  $\kappa: E \to G$  is a continuous function (that is, a *cocycle*), then there exists a coaction  $\varepsilon$  of G on  $C^*(E)$  such that

$$C^*(E \times_{\kappa} G) \cong C^*(E) \times_{\varepsilon} G.$$

We give two distinct approaches to the coaction: in Section 3 we obtain the coaction indirectly, via an application of Landstad duality, and in Section 4 we construct the coaction directly, applying techniques developed in [6]. We thank Iain Raeburn for helpful conversations concerning this direct approach.

In Section 2 we record our conventions for topological graphs,  $C^*$ correspondences, skew products, multiplier modules, and functoriality of Cuntz-Pimsner algebras. In an appendix we develop a few tools that we need for dealing with certain bimodule multipliers in terms of function spaces.

Date: December 12, 2012.

<sup>2000</sup> Mathematics Subject Classification. Primary 46L05; Secondary 46L55. Key words and phrases. topological graph, coaction, skew product.

#### KALISZEWSKI AND QUIGG

### 2. Preliminaries

In general, we refer to [11] (see also [8]) for topological graphs, and to [11, 3] (see also [7]) for  $C^*$ -correspondences, except we make a few minor, self-explanatory modifications. Thus, a topological graph E comprises locally compact Hausdorff spaces  $E^1$ ,  $E^0$  and maps  $s, r: E^1 \to E^0$ with s a local homeomorphism and r continuous. Let  $A = C_0(E^0)$ , and let X = X(E) be the associated A-correspondence, which is the completion of  $C_c(E^1)$  with operations defined for  $f \in A$  and  $\xi, \eta \in C_c(E^1)$ by

$$f \cdot \xi(e) = f(r(e))\xi(e)$$
  
$$\xi \cdot f(e) = \xi(e)f(s(e))$$
  
$$\langle \xi, \eta \rangle(v) = \sum_{s(e)=v} \overline{\xi(e)}\eta(e)$$

Throughout this paper we will also write  $A' = C_0(E^1)$ , so that Xcan be regarded as an A' - A correspondence as well as an Acorrespondence. Recall from [8] that the left A'-module multiplication is nondegenerate in the sense that  $A' \cdot X = X$ , and is determined by the homomorphism  $\pi_E \colon A' \to \mathcal{L}(X)$  given by  $(\pi_E(f)\xi)(e) = f(e)\xi(e)$ for  $f \in A'$  and  $\xi \in C_c(E^1)$ , and the (nondegenerate) left A-module multiplication  $\varphi_A \colon A \to \mathcal{L}(X)$  is then given by  $\varphi_A(f) = \pi_E(1 \circ r)$  for  $f \in A$ .

We denote by  $(k_X, k_A)$ :  $(X, A) \to C^*(E) = \mathcal{O}_X$  the universal Cuntz-Pimsner covariant representation, and for any Cuntz-Pimsner covariant representation  $(\psi, \pi)$  of (X, A) in a  $C^*$ -algebra B we denote by  $\psi^{(1)} \colon \mathcal{K}(X) \to B$  the associated homomorphism<sup>2</sup> determined by  $\psi^{(1)}(\theta_{\xi,\eta}) = \psi(\xi)\psi(\eta)^*$ , and by  $\psi \times \pi \colon C^*(E) \to B$  the unique homomorphism satisfying

$$(\psi \times \pi) \circ k_X = \psi$$
 and  $(\psi \times \pi) \circ k_A = \pi$ .

Note that in [3], correspondences were called right-Hilbert bimodules, and nondegeneracy was built into the definition. All our correspondences will in fact be nondegenerate, so we can freely apply the results from [3].

For skew products of topological graphs, we use a slight variation of the definition in [1]: the main difference is that we use the same notational conventions as those in [11] for skew products of discrete directed graphs. Thus, a *cocycle* of a locally compact group G on a

1

<sup>&</sup>lt;sup>2</sup>and here we use the notation of [1]; Raeburn would write  $(\psi, \pi)^{(1)}$ 

topological graph E is a continuous map  $\kappa \colon E^1 \to G$ , and the *skew* product is the topological graph  $E \times_{\kappa} G$  with

$$(E \times_{\kappa} G)^{i} = E^{i} \times G \quad (i = 0, 1),$$
  
 
$$r(e, t) = (r(e), \kappa(e)t), \quad \text{and} \quad s(e, t) = (s(e), t).$$

Our conventions for multipliers of correspondences are taken primarily from [3, Chapter 1], but also see [7]. If  $(\pi, \psi, \tau)$ :  $(A, X, B) \rightarrow$ (M(C), M(Y), M(D)) is a correspondence homomorphism, then there is a unique homomorphism  $\psi^{(1)} \colon \mathcal{K}(X) \to M(\mathcal{K}(Y)) = \mathcal{L}(Y)$  such that  $\psi^{(1)}(\theta_{\xi,\eta}) = \psi(\xi)\psi(\eta)^*$  for  $\xi, \eta \in X$ . (For this result in the stated level of generality, in particular with no nondegeneracy assumption on  $(\psi, \pi)$ , see [7, Lemma 2.1].) If  $(\psi, \pi)$  happens to be nondegenerate, then so is  $\psi^{(1)}$ , and hence  $\psi^{(1)}$  extends uniquely to a homomorphism  $\overline{\psi^{(1)}} \colon \mathcal{L}(X) \to \mathcal{L}(Y)$ .

A correspondence homomorphism  $(\psi, \pi) \colon (X, A) \to (M(Y), M(B))$ is defined in [7] to be *Cuntz-Pimsner covariant* if

- (i)  $\psi(X) \subset M_B(Y)$ ,
- (ii)  $\pi: A \to M(B)$  is nondegenerate,
- (iii)  $\pi(J_X) \subset M(B; J_Y)$ , and
- (iv) the diagram

commutes,

where, for an ideal I of a  $C^*$ -algebra C,

$$M(C;I) := \{ m \in M(C) : mC \cup Cm \subset I \}.$$

By [7, Corollary 3.6], for each Cuntz-Pimsner covariant homomorphism  $(\psi, \pi)$ , there is a unique homomorphism  $\mathcal{O}_{\psi,\pi}$  making the diagram



commute. Moreover,  $\mathcal{O}_{\psi,\pi}$  is nondegenerate, and is injective if  $\pi$  is.

Our conventions for coactions on correspondences mainly follow [3], but see also [6].

#### KALISZEWSKI AND QUIGG

## 3. Indirect Approach

In this section we apply Landstad duality to give an indirect approach to the following result:

**Theorem 3.1.** If  $\kappa: E^1 \to G$  is a cocycle on a topological graph E, then there is a coaction  $\varepsilon$  of G on  $C^*(E)$  such that

$$C^*(E \times_{\kappa} G) \cong C^*(E) \times_{\varepsilon} G.$$

Throughout the rest of this paper, in addition to  $A = C_0(E^0)$  and X = X(E), we will also use the following abbreviations:

- $F = E \times_{\kappa} G;$
- $Y = X(E \times_{\kappa} G);$
- $B = C_0((E \times_{\kappa} G)^0).$

*Proof.* To apply Landstad duality [10, Theorem 3.3] (stated in more modern form in [5, Theorem 4.1]), we need the following ingredients: an action  $\alpha: G \to \operatorname{Aut} C^*(F)$ , a  $\operatorname{rt} - \alpha$  equivariant nondegenerate homomorphism  $\mu: C_0(G) \to M(C^*(F))$  (where "rt" is action of G on  $C_0(G)$ by right translation), and an injective nondegenerate homomorphism

$$\Pi \colon C^*(E) \to M(C^*(F))$$

whose image coincides with Rieffel's generalized fixed-point algebra  $C^*(F)^{\alpha}$ . Note that in [5],  $C^*(F)^{\alpha}$  would be written as  $Fix(C^*(F), \alpha, \mu)$ .

Since G acts on the right of the skew-product topological graph F via right translation in the second coordinate, by [1, Proposition 5.4 and discussion preceding Remark 5.3] we have an action  $\beta = (\beta^1, \beta^0) \colon G \to$ Aut Y such that

$$\beta_t^1(\xi)(e,r) = \xi(e,rt) \quad \text{for } \xi \in Y$$
  
$$\beta_t^0(g)(v,r) = g(v,rt) \quad \text{for } g \in B,$$

which in turn gives an action on  $C^*(F)$  such that

$$\alpha_t \circ k_Y = k_Y \circ \beta_t^1$$
$$\alpha_t \circ k_B = k_B \circ \beta_t^0.$$

Since  $F^0 = E^0 \times G$ , we have

$$B = C_0(F^0) = C_0(E^0) \otimes C_0(G) = A \otimes C_0(G),$$

so we can define a nondegenerate homomorphism  $\mu: C_0(G) \to M(C^*(F))$  by

$$\mu(g) = \overline{k_B}(1_{M(A)} \otimes g),$$

and then it is routine to verify that  $\mu$  is  $rt - \alpha$  equivariant.

Finally, since the action of G on F is free and proper, the proof of [1, Theorem 5.6] constructs an isomorphism

$$\Pi \colon C^*(E) \xrightarrow{\cong} C^*(F)^{\alpha},$$

and then the result follows from Landstad duality.

## 4. A DIRECT APPROACH TO THE COACTION

As in Section 3, we suppose we are given a cocycle  $\kappa \colon E^1 \to G$  of a locally compact group G on a topological graph E, and we continue to write  $A = C_0(E^0)$ ,  $A' = C_0(E^1)$ , X = X(E),  $F = E \times_{\kappa} G$ , Y = X(F), and  $B = C_0(F^0)$ .

Recall that the canonical embedding  $G \hookrightarrow M(C^*(G))$  is identified with a unitary element  $w_G$  of  $M(C_0(G) \otimes C^*(G))$ . Similarly, we may identify  $\kappa$  with a unitary element of

$$C_b(E^1, M^{\beta}(C^*(G))) = M(A' \otimes C^*(G)),$$

where  $M^{\beta}(C^*(G))$  denotes the multiplier algebra  $M(C^*(G))$  with the strict topology.

Define a nondegenerate homomorphism  $\kappa^* \colon C_0(G) \to M(A')$  by  $\kappa^*(f) = f \circ \kappa$ , and a nondegenerate homomorphism  $\nu \colon C_0(G) \to \mathcal{L}(X)$  by

$$\nu = \pi_E \circ \kappa^*,$$

where  $\pi_E \colon A' \to \mathcal{L}(X)$  is the homomorphism given on  $C_c(E^1)$  by pointwise multiplication.

**Proposition 4.1.** With the above notation, there is a coaction ( $\sigma$ , id<sub>A</sub> $\otimes$  1) of G on (X, A) defined by

$$\sigma(\xi) = v \cdot (\xi \otimes 1),$$

where

$$v = \overline{\nu \otimes \mathrm{id}}(w_G) \in \mathcal{L}(X \otimes C^*(G)),$$

and moreover there is a coaction  $\zeta$  of G on  $C^*(E)$  such that

$$\zeta \circ k_X = \overline{k_X \otimes \mathrm{id}} \circ \sigma$$
$$\zeta \circ k_A = k_A \otimes 1.$$

*Proof.* This follows from [6, Corollaries 3.4–3.5], because  $\nu \colon C_0(G) \to \mathcal{L}(X)$  commutes with  $\varphi_A$ .

It will be convenient for us to find an equivalent expression for the coaction  $\sigma$ . Note that we may regard X as an A' - A correspondence,

and hence  $X \otimes C^*(G)$  as an  $(A' \otimes C^*(G)) - (A \otimes C^*(G))$  correspondence. Thus we can write

$$\sigma(\xi) = \overline{\kappa^* \otimes \mathrm{id}}(w_G) \cdot (\xi \otimes 1).$$

However, we can go further: by construction the unitary element  $\overline{\kappa^* \otimes \mathrm{id}}(w_G)$  of  $M(A' \otimes C^*(G))$  coincides with the function in  $C_b(E^1, M^\beta(C^*(G)))$  whose value at an edge e is

$$\overline{\kappa^* \otimes \mathrm{id}}(w_G)(e) = w_G(\kappa(e)) = \kappa(e);$$

thus we can write

$$\sigma(\xi) = \kappa \cdot (\xi \otimes 1).$$

In Theorem 3.1 we used Landstad duality to show that  $C^*(F)$  is isomorphic to the crossed product of  $C^*(E)$  by a coaction  $\varepsilon$  of G; on the other hand, in Proposition 4.1 we directly constructed a coaction  $\zeta$ of G on  $C^*(E)$ . To show that also  $C^*(E) \times_{\zeta} G \cong C^*(F)$ , we now show that in fact the coactions  $\varepsilon$  and  $\zeta$  coincide. Since the mechanism behind Landstand duality is that  $\varepsilon$  is pulled back along  $\Pi^{-1}$  from the inner coaction  $\delta^{\mu}$  on  $C^*(F)$ , this is accomplished by the following:

**Proposition 4.2.** Let  $\zeta$  be the coaction on  $C^*(E)$  from Proposition 4.1, and let  $\Pi: C^*(E) \to M(C^*(F))$  and  $\mu: C_0(G) \to M(C^*(F))$  be as in the proof of Theorem 3.1. Then  $\Pi$  is  $\zeta - \delta^{\mu}$  equivariant, and hence  $\zeta$ coincides with the coaction  $\varepsilon$  from Theorem 3.1.

Proof. It is equivalent to show that  $(\Pi, \mu)$ :  $(C^*(E), C_0(G)) \rightarrow M(C^*(F))$  is a covariant representation for the coaction  $\zeta$ , and for this we will apply [6, Corollary 4.3].

We will need to know how the homomorphism  $\Pi$  from [1] can be described using the techniques of [7]: [1, Proof of Theorem 5.6] constructs a correspondence homomorphism

 $(\psi, \pi) \colon (X, A) \to (M_B(Y), M(B)),$ 

although the notation in [1] is substantially different<sup>3</sup>. In the terminology of [7, Definition 3.1], [1, Proof of Theorem 5.6] shows that  $(\psi, \pi)$ is Cuntz-Pimsner covariant, so that by [7, Corollary 3.6] there is a nondegenerate homomorphism  $\mathcal{O}_{\psi,\pi}$  making the diagram



<sup>&</sup>lt;sup>3</sup>The roles of E, X, A and F, Y, B are interchanged, and what we call  $(\psi, \pi)$  here was written as  $(\mu, \nu)$  in [1].

6

commute; the homomorphism  $\Pi$  from [1] coincides with  $\mathcal{O}_{\psi,\pi}$ . Thus, by [6, Corollary 4.3] it suffices to show that

$$(\psi, \pi, \mu) \colon (X, A, C_0(G)) \to (M_B(Y), M(B))$$

is covariant for  $(\sigma, \mathrm{id}_A \otimes 1)$ , in the sense of [6, Definition 2.9]. Thus we must show that

- (i)  $(\pi, \mu)$  is covariant for  $(A, \mathrm{id}_A \otimes 1)$ , and
- (ii)  $\overline{\psi \otimes \mathrm{id}} \circ \sigma(\xi) = \overline{\mu \otimes \mathrm{id}}(w_G) \cdot (\psi(\xi) \otimes 1) \cdot \overline{\mu \otimes \mathrm{id}}(w_G)^*$  for all  $\xi \in X.$

Condition (i) is immediate because  $\pi$  and  $\mu$  commute. Next, we rewrite (ii) in an equivalent form:

(ii)' 
$$\overline{\psi \otimes \mathrm{id}}(\sigma(\xi)) \cdot \overline{\mu \otimes \mathrm{id}}(w_G) = \overline{\mu \otimes \mathrm{id}}(w_G) \cdot (\psi(\xi) \otimes 1)$$
 for all  $\xi \in X$ 

To proceed further, notice that the maps  $\psi$ ,  $\pi$ , and  $\mu$  from [1] take a particularly simple form in our present context:

- $\psi = \operatorname{id}_X \otimes 1_{M(C_0(G))};$
- $\pi = \operatorname{id}_A \otimes 1_{M(C_0(G))};$
- $\mu = 1_{M(A)} \otimes \operatorname{id}_{C_0(G)}$ .

(We should explain our notation in the above expression for  $\psi$ : it follows from the definitions that, as a Hilbert  $(A \otimes C_0(G))$ -module, Y coincides with the external tensor product  $X \otimes C_0(G)$  (where  $C_0(G)$ ) is regarded as a Hilbert module over itself in the canonical way). One just has to keep in mind that Y does not coincide with  $X \otimes C_0(G)$  as a *B*-correspondence — the left *B*-module multiplication is twisted by the cocycle  $\kappa$ .) Thus, for  $\xi \in X$  we can write:

- $\psi(\xi) = \xi \otimes 1;$
- $\overline{\psi \otimes \mathrm{id}}(\sigma(\xi)) = \sigma(\xi)_{13} = \kappa_{13} \cdot (\xi \otimes 1 \otimes 1);$
- $\overline{\mu \otimes \mathrm{id}}(w_G) = 1 \otimes w_G.$

Since both sides of (ii)' are adjointable Hilbert-module maps from  $B \otimes C^*(G)$  to  $Y \otimes C^*(G)$ , and  $A \odot C_c(G)$  is dense in B, it suffices to check that the two sides of (ii)' take equal values on elementary tensors of the form  $f \otimes g \otimes a$ , with  $f \in A$ ,  $g \in C_c(G)$ , and  $a \in C^*(G)$ . Evaluating the right-hand side of (ii)' gives

(4.1) 
$$((1 \otimes w_G) \cdot (\xi \otimes 1 \otimes 1)) \cdot (f \otimes g \otimes a)$$
$$= (1 \otimes w_G) \cdot ((\xi \otimes 1 \otimes 1) \cdot (f \otimes g \otimes a))$$
$$= (1 \otimes w_G) \cdot (\xi \cdot f \otimes g \otimes a).$$

Now we must use the function-space techniques from Appendix A. We have  $1 \otimes w_G \in C_b(F^0, M^\beta(C^*(G)))$ , with value t at  $(v, t) \in F^0$ , and

 $\xi \cdot f \otimes g \otimes a \in C_c(F^1, C^*(G))$ , so by Corollary A.3 we can evaluate the last quantity in (4.1) at  $(e, t) \in F^1$ , giving

$$(1 \otimes w_G) (r(e,t)) (\xi \cdot f \otimes g \otimes a)(e,t)$$
  
=  $(1 \otimes w_G) (r(e), \kappa(e)t) (\xi \cdot f)(e)g(t)a$   
=  $\kappa(e)t\xi(e)f(s(e))g(t)a$   
=  $\xi(e)f(s(e))g(t)\kappa(e)ta.$ 

We proceed similarly with the left-hand side of (ii)':

(4.2) 
$$\begin{pmatrix} \kappa_{13} \cdot (\xi \otimes 1 \otimes 1) \cdot (1 \otimes w_G) \end{pmatrix} \cdot (f \otimes g \otimes a) \\ = \kappa_{13} \cdot (\xi \cdot f \otimes w_G(g \otimes a))$$

Now,  $\kappa_{13} \in C_b(F^1, M^\beta(C^*(G)))$ , with value  $\kappa(e)$  at (e, t), and  $\xi \cdot f \otimes w_G(g \otimes a) \in C_c(F^1, C^*(G))$  because  $\xi \cdot f \in C_c(E^1)$  and  $w_G(g \otimes a) \in C_c(G, C^*(G))$ , so by Corollary A.3 we can evaluate the right-hand side of (4.2) at  $(e, t) \in F^1$ , giving

$$\begin{pmatrix} \kappa_{13} \cdot (\xi \cdot f \otimes w_G(g \otimes a)) \end{pmatrix} (e,t) \\ = \kappa_{13}(e,t) (\xi \cdot f \otimes w_G(g \otimes a)) (e,t) \\ = \kappa(e) (\xi \cdot f(e) (w_G(g \otimes a)) (t) \\ = \kappa(e) \xi(e) f(s(e)) w_G(t) (g \otimes a) (t) \\ = \kappa(e) \xi(e) f(s(e)) tg(t) a \\ = \xi(e) f(s(e)) g(t) \kappa(e) ta. \end{cases}$$

Therefore we have verified (ii)', and this finishes the proof.

## APPENDIX A. FUNCTIONS AND MULTIPLIERS

In Section 4 we need to compute with bimodule multipliers in terms of functions. If T is a locally compact Hausdorff space and C is a  $C^*$ -algebra, we will use without comment the following identifications (see, e.g., [12] or [3, Appendix C]):

- $C_0(T,C) = C_0(T) \otimes C;$
- $M(C_0(T) \otimes C) = C_b(T, M^\beta(C)),$

where we write  $M^{\beta}(C)$  to denote M(C) with the strict topology. Note that since the action of  $C_b(T, M^{\beta}(C))$  by multipliers on  $C_0(T, C)$  is via pointwise multiplication, it preserves  $C_c(T, C)$ .

We will need to use functions as multipliers on certain  $C^*$ correspondences; since this theory is not easily available in the
literature, we give details for all the results we need. However, we

make no attempt to construct a general theory — rather, we do only enough to establish Corollary A.3, which we need in Section 4.

For a topological graph E, we write (as in the rest of the paper)  $A = C_0(E^0), X = X(E)$ , and  $A' = C_0(E^1)$ . We will regard  $X \otimes C$ both as an  $(A' \otimes C) - (A \otimes C)$  correspondence and as an  $(A \otimes C)$ correspondence.

The following lemma is routine:

**Lemma A.1.**  $C_c(E^1, C)$  embeds densely in the  $(A' \otimes C) - (A \otimes C)$ correspondence  $X \otimes C$  in the following way: if  $\xi, \eta \in C_c(E^1, C) \subset X \otimes C$ ,  $f \in C_c(E^1, C) \subset A' \otimes C$ , and  $g \in C_c(E^0, C) \subset A \otimes C$ , then  $f \cdot \xi$  and  $\xi \cdot g$  are the elements of  $C_c(E^1, C)$  given by

(A.1) 
$$(f \cdot \xi)(e) = f(e)\xi(e)$$

(A.2) 
$$(\xi \cdot g)(e) = \xi(e) \cdot g(s(e)),$$

and  $\langle \xi, \eta \rangle$  is the element of  $C_c(E^0, C) \subset A \otimes C$  given by

(A.3) 
$$\langle \xi, \eta \rangle(v) = \sum_{s(e)=v} \xi(e)^* \eta(e).$$

Moreover,  $g \cdot \xi$  is the element of  $C_c(E^1, C)$  given by

(A.4) 
$$(g \cdot \xi)(e) = g(r(e))\xi(e).$$

*Proof.* First of all, (A.2)–(A.3) make  $C_c(E^1, C)$  into a pre-Hilbert  $C_c(E^0, C)$ -module (where the latter is regarded as a dense \*-subalgebra of  $C_0(E^0, C) = A \otimes C$ ). The only non-obvious property of pre-Hilbert modules is that (A.3) does give an element of  $C_c(E^0, C)$ , but this can be proved by an argument similar to those used in [8, Lemma 1.5].

Observe that the Hilbert-module norm on  $C_c(E^1, C)$  is given by

(A.5) 
$$\|\xi\| = \sup_{v \in E^0} \left\| \sum_{s(e)=v} \xi(e)^* \xi(e) \right\|^{1/2}$$

which is larger than the uniform norm. In particular, for  $e \in E^1$  the evaluation map  $\xi \mapsto \xi(e)$  from  $C_c(E^1, C)$  to C is bounded from the Hilbert-module norm to the norm of C.

Computing with elementary tensors of the form  $\xi \otimes c$  for  $\xi \in C_e(E^1)$ and  $c \in C$ , it is now routine to verify that the completion of the pre-Hilbert module  $C_c(E^1, C)$  is isomorphic to the external tensor product  $X \otimes C$  of the Hilbert A-module X and the Hilbert C-module C.

Now regarding  $X \otimes C$  as an  $(A' \otimes C) - (A \otimes C)$  correspondence, (A.1) is obviously true on elementary tensors, hence for  $f, \xi \in C_c(E^1) \odot C$ , and therefore as stated by density of  $C_c(E^1) \odot C$  in  $C_c(E^1, C)$  and by continuity of evaluation. Finally, (A.4) follows from (A.1).

**Lemma A.2.** Let  $K \subset E^1$  be compact. On the subspace

$$C_K(E^1, C) := \{\xi \in C_c(E^1, C) : \operatorname{supp} \xi \subset K\}$$

of  $X \otimes C$ , the Hilbert-module norm and the uniform norm are equivalent. Consequently,  $C_K(E^1, C)$  is norm-closed in  $X \otimes C$ .

*Proof.* By (A.5), the uniform norm on  $C_K(E^1, C)$  is smaller than the Hilbert-module norm from  $X \otimes C$ . Thus it suffices to show that the Hilbert-module norm is bounded above by a multiple of the uniform norm. Let  $\xi \in C_K(E^1, C)$ . Using compactness of K and local homeomorphicity of s, it is easy to verify that the cardinalities of the intersections  $K \cap s^{-1}(v)$  for  $v \in E^0$  are bounded above by some nonnegative integer d. Then for any  $v \in E^0$  we have

$$\left\|\sum_{s(e)=v} \xi(e)^* \xi(e)\right\| \le \sum_{s(e)=v} \|\xi(e)\|^2 \le d \|\xi\|_u^2,$$

where  $\|\xi\|_u$  denotes the uniform norm of  $\xi$ , and the result follows.  $\Box$ 

Since  $X \otimes C$  is a nondegenerate  $(A' \otimes C) - (A \otimes C)$  correspondence, the left module action of  $A' \otimes C$  extends canonically to the multiplier algebra  $M(A' \otimes C) = C_b(E^1, M^\beta(C))$  (and similarly for the left module action of  $A \otimes C$ ). The following corollary allows us to compute this extended left module action on generators:

**Corollary A.3.** If  $m \in C_b(E^1, M^{\beta}(C))$  and  $\xi \in C_c(E^1, C) \subset X \otimes C$ , then the element  $m \cdot \xi$  of  $X \otimes C$  lies in  $C_c(E^1, C)$ , and

(A.6)  $(m \cdot \xi)(e) = m(e)\xi(e) \text{ for } e \in E^1.$ 

If  $n \in C_b(E^0, M^\beta(C))$  then both  $n \cdot \xi$  and  $\xi \cdot n$  lie in  $C_c(E^1, C)$ , and  $(n \cdot \xi)(e) = n(r(e))\xi(e);$  $(\xi \cdot n)(e) = \xi(e)n(s(e)).$ 

*Proof.* Choose a net  $\{m_i\}$  in  $C_c(E^1, C)$  converging strictly to m in  $M(C_0(E^1, C))$ . The conclusion holds for each  $m_i \cdot \xi$ , by Lemma A.1. Let  $K = \operatorname{supp} \xi$ , a compact subset of  $E^1$ . Then  $m_i \cdot \xi \in C_K(E^1, C)$  for all i. Since  $m_i \cdot \xi \to m \cdot \xi$  in the norm of  $X \otimes C$ , we have  $m \cdot \xi \in C_K(E^1, C)$ , by Lemma A.2.

For each  $e \in E^1$ , by norm-continuity of evaluation on  $C_c(E^1, C)$  we have

$$(m \cdot \xi)(e) = \lim_{i} (m_i \cdot \xi)(e) = \lim_{i} m_i(e)\xi(e).$$

Moreover, for any  $a \in C$ , we can choose  $f \in C_0(E^1, C)$  such that f(e) = a and compute:

$$m_i(e)a = m_i(e)f(e) = (m_i f)(e) \to (mf)(e) = m(e)a.$$

Thus evaluation is strictly continuous on  $C_b(E^1, M^{\beta}(C))$ ; in particular,

$$\lim m_i(e)\xi(e) = m(e)\xi(e),$$

which establishes (A.6).

The statement for  $n \cdot \xi$  follows by composing with the range map  $r: E^1 \to E^0$ , and the statement for  $\xi \cdot n$  is proved similarly to the above argument for  $m \cdot \xi$ .

## References

- V. Deaconu, A. Kumjian, and J. Quigg, Group actions on topological graphs, Ergodic Theory Dynam. Systems **32** (2012), 1527–1566.
- [2] K. Deicke, D. Pask, and I. Raeburn, Coverings of directed graphs and crossed products of C<sup>\*</sup>-algebras by coactions of homogeneous spaces, Internat. J. Math. 14 (2003), no. 7, 773–789.
- [3] S. Echterhoff, S. Kaliszewski, J. Quigg, and I. Raeburn, A Categorical Approach to Imprimitivity Theorems for C\*-Dynamical Systems, vol. 180, Mem. Amer. Math. Soc., no. 850, American Mathematical Society, Providence, RI, 2006.
- [4] S. Kaliszewski, J. Quigg, and I. Raeburn, Skew products and crossed products by coactions, J. Operator Theory 46 (2001), 411–433.
- [5] \_\_\_\_\_, Proper actions, fixed-point algebras and naturality in nonabelian duality, J. Funct. Anal. 254 (2008), 2949–2968.
- [6] S. Kaliszewski, J. Quigg, and D. Robertson, Coactions on Cuntz-Pimsner algebras, arXiv:1204.5822.
- [7] \_\_\_\_\_, Functoriality of Cuntz-Pimsner correspondence maps, arXiv:1204.5820.
- [8] T. Katsura, A class of C\*-algebras generalizing both graph algebras and homeomorphism C\*-algebras. I. Fundamental results, Trans. Amer. Math. Soc. 356 (2004), no. 11, 4287–4322.
- [9] D. Pask, J. Quigg, and I. Raeburn, *Coverings of k-graphs*, J. Algebra 289 (2005), no. 1, 161–191.
- [10] J.C. Quigg, Landstad duality for C<sup>\*</sup>-coactions, Math. Scand. **71** (1992), 277– 294.
- [11] I. Raeburn, Graph algebras, CBMS Regional Conference Series in Mathematics, vol. 103, Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2005.
- [12] I. Raeburn and D. P. Williams, Morita equivalence and continuous-trace C<sup>\*</sup>algebras, Math. Surveys and Monographs, vol. 60, American Mathematical Society, Providence, RI, 1998.

(S. Kaliszewski) SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, ARIZONA STATE UNIVERSITY, TEMPE, ARIZONA 85287 *E-mail address*: kaliszewski@asu.edu

(John Quigg) SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, ARI-ZONA STATE UNIVERSITY, TEMPE, ARIZONA 85287 *E-mail address*: quigg@asu.edu