

Regularity of the Gurtin-Pipkin equation

S. A. Ivanov*[†]

Abstract

We study regularity of the solution θ to the Gurtin-Pipkin integral-differential equation of the first order in time. In particular, we prove that the 'perturbation' part, namely, the difference of θ and the solution to the corresponding wave equation is smoother than θ .

1 Introduction

In several fields of physics such as heat transfer with finite propagation speed [2], systems with thermal memory [5], viscoelasticity problems [3], and acoustic waves in composite media [1], the integro-differential equations arise. We consider the equation of the first order in time

$$\theta_t(x, t) = \int_0^t k(t-s)\theta_{xx}(x, s) ds + f(x, t), \quad x \in (0, \pi), \quad t > 0, \quad (1)$$

with the Dirichlet boundary conditions and with the initial data $\theta(0, x) = \xi(x)$.

In the case $k(t) = Const = \alpha^2$ the equation (1) is, in a fact, an integrated wave equation. Indeed, differentiate (1) gives

$$\theta_{tt} = \alpha^2\theta_{xx} + f_t(x, t), \quad \theta(x, 0) = \xi, \quad \theta_t(x, 0) = 0. \quad (2)$$

Thus, the wave equation is a special case of (1) and we will compare general regularity results with the regularity of the solutions to the wave equation.

*St. Petersburg Institute of Terrestrial Magnetism, Ionosphere and Radio Wave Propagation. sergei.a.ivanov@mail.ru

[†]The work was supported by Russian Foundation for Basic Research, RFBR Project 11-01-00790a and RFBR Project 11-01-00667a.

2 Fourier method and the Laplace transform

First, apply the Fourier method: we set $\varphi_n = \sqrt{\frac{2}{\pi}} \sin nx$ and expand the solution, the RHS, and the initial data in series in φ_n

$$\theta(x, t) = \sum_1^{\infty} \theta_n(t) \varphi_n(x), \quad \xi(x) = \sum_1^{\infty} \xi_n \varphi_n(x), \quad f(x, t) = \sum_1^{\infty} f_n(t) \varphi_n(x).$$

The components θ_n satisfy ordinary integral-differential equations

$$\dot{\theta}_n(t) = -n^2 \int_0^t k(t-s) \theta_n(s) ds + f_n(t), \quad t > 0, \quad \theta_n(0) = \xi_n. \quad (3)$$

Note that the solutions to this integral-differential is unique and continuous, what we can see by t -integration of the equation from 0 to t . Indeed, we obtain a Volterra integral equation: with $\varphi = \int_0^t f(t) dt$, we have

$$\theta_n(t) - \xi_n = -n^2 \int_0^t d\tau \int_0^{\tau} k(\tau-s) \theta_n(s) ds + \varphi(t).$$

Change order of integrations

$$\theta_n(t) = -n^2 \int_0^t ds \theta_n(s) \int_s^t d\tau k(\tau-s) + \varphi(t) + \xi_n = -n^2 \int_0^t q(t-s) \theta_n(s) ds + \varphi(t) + \xi_n \quad (4)$$

with

$$q(s) = \int_0^s k(y) dy.$$

Introduce the scale \mathcal{H}_s , s is real, of the Hilbert spaces $\mathcal{H}_s = \text{Dom}(A^{s/2})$, where the operator A is $-d^2/dx^2$ with the Dirichlet boundary conditions at 0 and at π . A space \mathcal{H}_s is a subspace of the Sobolev space H^s and may be described in terms of the Fourier coefficients. Let the space l_s be the space of sequences $\{c_n\}$ such that

$$\sum_1^{\infty} |c_n|^2 n^{2s} < \infty.$$

Then

$$\mathcal{H}_s = \left\{ u(x) = \sum_1^{\infty} u_n \varphi_n(x) \mid \{u_n\} \in l_s \right\}.$$

Consider also the space $\mathfrak{H}_{s,\varepsilon}$ of functions $g(x,t) = \sum_1^\infty f_n(t)\varphi_n(x)$ with the norm

$$\|g\|_{\mathfrak{H}_{s,\varepsilon}}^2 = \sum_1^\infty n^{2s} \|e^{-2\varepsilon t} f_n\|_{L^2(0,\infty)}^2.$$

Definition 1 *The function $\theta(x,t) = \sum_1^\infty \theta_n(t)\varphi_n(x)$ is a solution to (1) in $\mathfrak{H}_{s,\varepsilon}$ if the functions θ_n , satisfy the integral equation (4) and $\theta \in \mathfrak{H}_{s,\varepsilon}$ with $s \in \mathbb{R}$.*

Let H_ε^2 denote the Hardy space in the right half plane $\Re z > \varepsilon$. The Paley-Wiener theorem says that

$$\|F\|_{H_\varepsilon^2}^2 = \int_0^\infty e^{-2\varepsilon t} |f(t)|^2 dt.$$

Here and in what follows will denote the Laplace image by the capital characters. Applying the Laplace Transform to (3) we find

$$z\Theta_n(z) - \xi_n = -n^2 K(z)\Theta_n(z) + F_n(z)$$

or

$$\Theta_n(z) = \frac{\xi_n + F_n(z)}{z + n^2 K(z)}. \quad (5)$$

Denote the denominators in (5) by $G_n(z)$. The set Λ of all zeros of $G_n(z)$ is called the *spectrum* of the equation (1).

Regularity of the Gurtin-Pipkin type equation is studied in [4] for several spatial variables, where under assumption that $k(t)$ is twice continuously differentiable it was shown, in particular, that $\Theta(x,t) \in C([0,T]; L^2(0,T))$. Regularity of strong solutions has been studied in several works of V. Vlasov with the coauthors, see, e.g., [8] and the Sec.5 below. The spectrum of the equation is studied in [9],[8].

Let us describe the regularity of the solutions to the the wave equation (2). Let $Q_T = (0, \pi) \times (0, T)$.

Proposition 2 *The solution to the (2) satisfy the following estimates: (i) Let $f = 0$. Then the Dalambert solution gives*

$$\|\theta\|_{\mathfrak{H}_{s,\varepsilon}} \prec \|\xi\|_{\mathcal{H}_s}, \quad \|\partial_t \theta\|_{\mathfrak{H}_{s,\varepsilon}} \prec \|\xi\|_{\mathcal{H}_{s+1}}, \quad (6)$$

(ii) let $\xi = 0$. Then, see [10] the (generalized) solutions satisfy

$$\|\partial_t \theta\|_{L^2(Q_T)} + \|\partial_x \theta\|_{L^2(Q_T)} \prec \|\xi\|_{\mathcal{H}_1} + \|f_t\|_{L^2(Q_T)}. \quad (7)$$

If $k(t) = \alpha^2 e^{-bt}$ (and $K(z) = \alpha^2/(z + b)$), then differentiation gives a damped wave equation

$$\theta_{tt} = \alpha^2 \theta_{xx} - b\theta_t. \quad (8)$$

By θ^0 we denote the solution to this equation with the initial data

$$\theta(\cdot, 0) = \xi, \quad \theta_t(\cdot, 0) = 0. \quad (9)$$

This will be an unperturbed equation, see the Sect. 4.

In application, see, e.g., [1], the kernels $k(t)$ is a series of exponentials

$$k(t) = \sum_1^{\infty} a_k e^{-b_k t}, \quad a_k \geq 0, \quad 0 \leq b_1 < b_2 < \dots < b_k < \dots$$

We can consider the following smoothness conditions

$$\sum_1^{\infty} \frac{a_k}{b_k} < \infty, \quad (10)$$

$$\alpha^2 = \sum_1^{\infty} a_k < \infty, \quad (11)$$

or

$$\beta = \sum_1^{\infty} a_k b_k < \infty, \quad (12)$$

or

$$\gamma = \sum_1^{\infty} a_k b_k^2 < \infty. \quad (13)$$

Remark 3 *These conditions maybe written as*

$$\begin{aligned} k &\in L^1(0, \infty), \\ k &\in C[0, \infty), \quad k, k' \in L^1(0, \infty), \\ k' &\in C[0, \infty), \quad k, k', k'' \in L^1(0, \infty), \\ k'' &\in C[0, \infty), \quad k, k', k'', k''' \in L^1(0, \infty). \end{aligned}$$

Write the asymptotic of $K(z)$. The Laplace image of $k(t)$ is

$$K(z) = \sum_1^{\infty} \frac{a_k}{z + b_k}, \quad k(0) = \alpha = \sum_1^{\infty} a_k.$$

Without loss of generality we can set $\alpha = 1$ if α is finite.

Proposition 4 *Let for a $\delta > 0$*

$$|\arg z| < \pi - \delta,$$

Then for large z

(i) under (10)

$$K(z) = o(1),$$

(ii) under (11)

$$K(z) = \frac{1}{z} + o\left(\frac{1}{z}\right),$$

(iii) under (12)

$$K(z) = \frac{1}{z} - \frac{\beta}{z^2} + o\left(\frac{1}{z^2}\right),$$

(iv) under (13)

$$K(z) = \frac{1}{z} - \frac{\beta}{z^2} + \frac{\gamma}{z^3} + o\left(\frac{1}{z^3}\right). \quad (14)$$

The statement of this proposition follows from known results about Cauchy transform of a measure.

3 Regularity of the solution in the spatial variable

Here we prove the results about the regularity with respect to the x -variable, i.e., in terms of \mathcal{H}_s spaces.

Theorem 5 *Let (11) be true, $\{\xi_n\} \in \ell_s$ and $f \in L^2(0, \infty; \mathcal{H}_s)$. Then for any $\varepsilon > 0$ the solution θ to (1) satisfy*

$$\|\theta\|_{\mathfrak{H}_{s,\varepsilon}}^2 \prec \|\xi\|_s^2 + \|f\|_{\mathfrak{H}_{s,\varepsilon}}^2. \quad (15)$$

PROOF:

Lemma 6 *The following estimates are fulfilled*

$$|z/G_n(z)| \prec 1, \quad \Re z > \varepsilon \quad (16)$$

$$\|1/G_n\|_{L^2(\varepsilon - i\infty, \varepsilon + i\infty)} \prec 1. \quad (17)$$

The lemma implies by (5)

$$\int_0^\infty |e^{-\varepsilon t} \theta_n(t)|^2 \prec |\xi_n|^2 + \|e^{-\varepsilon t} f_n\|_{L^2(0, \infty)}^2$$

and then (15).

Proof of the lemma. Set for the simplicity $\varepsilon = 1$ Then for $z = 1 + iy$ and $\gamma_k = 1 + b_k$ we obtain

$$G_n(z) = (1 + iy) + n^2 \sum_1^{\infty} \frac{a_k \gamma_k}{\gamma_k^2 + y^2} - iyn^2 \sum_1^{\infty} \frac{a_k}{\gamma_k^2 + y^2}$$

Therefore

$$|G_n(z)|^2 \geq \left(1 + n^2 \frac{a_1 \gamma_1}{\gamma_1^2 + y^2}\right)^2 + y^2 \left(1 - n^2 \sum_1^{\infty} \frac{a_k}{\gamma_k^2 + y^2}\right)^2$$

Setting

$$s(y) = \sum_1^{\infty} \frac{a_k}{\gamma_k^2 + y^2}.$$

we have

$$|G_n(z)|^2 \succ \left(1 + n^2 \frac{1}{1 + y^2}\right)^2 + y^2 (1 - n^2 s(y))^2.$$

This gives (16).

Divide $[0, \infty)$ into three intervals

$$I_1 = [0, n/2], I_2 = [n/2, 3n/2], I_3 = [3n/2, /iy].$$

Write

$$\|1/G_n\|_{L^2(\varepsilon - i\infty, \varepsilon + i\infty)}^2 = \int_{-\infty}^{\infty} \frac{dy}{|G_n(y)|^2} = 2 \left[\int_0^{n/2} + \int_{n/2}^{2n} + \int_{2n}^{\infty} \right] \frac{dy}{|G_n(y)|^2} = 2(J_1 + J_2 + J_3).$$

1. Estimates on $I_1 = [0, n/2]$.

Evidently, $s(y)$ decreases and then on $[0, n/2]$ we have $s(n/2) < s(y) < s(0)$. Further, the series

$$n^2 s(y) = \sum_1^{\infty} \frac{a_k n^2}{\gamma_k^2 + n^2/4}$$

has the majorant $4 \sum_1^{\infty} a_k = 4\alpha = 4$ and the terms of this series approaches to $4a_k$. Then

$$n^2 s(n/2) \rightarrow 4.$$

Take $n > n_0$ such that $n^2 s(n/2) \geq 2$ for $n/2 \geq n_0$. We obtain $n^2 s(y) > n^2 s(n/2) \geq 2$ and

$$(n^2 s(n/2) - 1)^2 \geq 1.$$

This gives

$$|G_n(z)|^2 \geq 1 + y^2 (n^2 s(n/2) - 1)^2 \geq 1 + y^2.$$

Estimate J_1 .

$$J_1 \leq \int_0^{n/2} \frac{1}{1+y^2} < 1. \quad (18)$$

2. Estimate J_2 .

For $n/2 \leq y \leq 2n$ we have

$$|G_n(z)| > 1 + n^2(1 - n^2s(y)),$$

Consider the increasing variable $\xi = 1 - n^2s(y)$. Then

$$\begin{aligned} \xi' = -n^2s' &= n^2 \sum_1^\infty \frac{2a_k y}{(\gamma_k^2 + y^2)^2} \asymp n^2 \sum_1^\infty \frac{a_k n}{(\gamma_k^2 + n^2)^2} \\ &\asymp \frac{1}{n} \sum_1^\infty \frac{a_k}{(\gamma_k^2/n^2 + 1)^2} \asymp \frac{1}{n}. \end{aligned}$$

Indeed,

$$\frac{a_1}{(\gamma_1^2/n^2 + 1)^2} \leq \sum_1^\infty \frac{a_k}{(\gamma_k^2/n^2 + 1)^2} \leq \sum_1^\infty a_k.$$

Now for J_2 we have " $nd\xi \asymp dy$ " and

$$J_2 > \int_{n/2}^{2n} \frac{dy}{1 + n^2(1 - n^2s(y))^2} = \int_{\xi(n/2)}^{\xi(2n)} \frac{nd\xi}{1 + n^2\xi^2} < \infty.$$

3. Estimate J_3 .

For $y \geq 2n$ we have

$$n^2s(y) \leq n^2s(2n) = \sum_1^\infty \frac{a_k n^2}{\gamma_k^2 + 4n^2} \leq \frac{1}{4} \sum_1^\infty a_k = \frac{1}{4}.$$

Now

$$|G_m(z)| \geq 1 + y^2(1 - n^2s(y))^2 \geq 1 + y^2 \frac{9}{16},$$

and

$$J_3 < \int_{2n}^\infty \frac{dy}{1 + y^2} < 1.$$

The theorem is proved.

Remark 7 For the case $k(t) = 1$, and $f = 0$, i.e., for the wave equation we have

$$\Theta_n(z) = \frac{\xi_n}{z + n^2/z}, \quad \theta_n(t) = \xi_n \cos nt.$$

We see that $\theta_n \notin L^2(0, \infty)$ and $e^{-\epsilon t} \theta_n \in L^2(0, \infty)$. In this sense Theorem 5 is sharp.

Theorem 8 Let (11) is true, $\{\xi_n\} \in \ell_s$ and $f \in L^2(0, \infty; \mathcal{H}_s)$. Then $\theta(x, t)$ is an \mathcal{H}_s valued continuous function:

$$\|\theta(t) - \theta(t + t_0)\|_{\mathfrak{H}_{s, \varepsilon}} \rightarrow 0,$$

as $t_0 \rightarrow t_0$.

Proof. The solutions θ_n are continuous and the series in θ_n has a majorant.

4 GP as a perturbation to the wave equation

Let us find regularity of the 'perturbation' $\theta - \theta^0$ of the solution to the wave equation. Recall that θ^0 is the solution to the problem (8),(9). Let $f(x, t) = 0$ for simplicity and set

$$K_0(z) = \frac{1}{z + \beta}, \quad G_n^0(z) = z + n^2 K_0(z), \quad D_n(z) = \frac{1}{G_n(z)} - \frac{1}{G_n^0(z)}.$$

The solution to (1) has the form

$$\Theta_n(z) = \frac{1}{G_n^0(z)} \xi_n + D_n(z) \xi_n = \Theta_n^0(z) + D_n(z) \xi_n.$$

Theorem 9 Let (13) is true and $f = 0$. Then for $s < 9/2$

$$\|z^s D_n\|_{L^2(i\mathbb{R})} \prec n^{s-1} |\xi_n|.$$

PROOF:

If $\beta = 0$ the theorem is trivial: $D_n = 0$. Thus, we can assume $\beta \neq 0$ and then integrate the functions on the imaginary axis.

Lemma 10 For $z = iy$, $y \rightarrow \infty$ and $\beta \neq 0$ we have

$$|G_n(z)|^2 \asymp |G_n^0(z)|^2 \asymp \frac{1}{y^4} \left[(y^2(y^2 - n^2))^2 + n^4 \right] =: Q(y).$$

Proof of the lemma. (14) implies

$$\begin{aligned} G_n(iy) &= iy + n^2 \left(\frac{1}{iy} - \frac{\beta}{y^2} + o\left(\frac{1}{y^2}\right) \right) \\ &= \frac{i}{y^2} [y^3 - n^2 y + o(1)n^2] - \frac{n^2}{y^2} [\beta + o(1)]. \end{aligned}$$

And the same is true for G_n^0 . From here

$$|G_n(y)|^2 = \frac{1}{y^4} [y^3 - n^2 y + o(1)n^2]^2 + \frac{n^4}{y^2} [\beta + o(1)]^2 \asymp \frac{1}{y^4} [y(y^2 - n^2) + o(1)n^2]^2 + \frac{n^4}{y^2}.$$

Now

$$\frac{|G_n(y)|^2}{Q(y)} = \frac{[y(y^2 - n^2) + o(1)n^2]^2 + n^4}{(y^2(y^2 - n^2))^2 + n^4}$$

Use the elementary inequalities

$$(a + qb)^2 + b^2 \asymp a^2 + b^2$$

with, say, $q < 1/2$. Setting

$$a = y(y^2 - n^2), \quad b = n^2$$

we complete the proof of the lemma.

Return to the proof of the theorem.

$$\|z^s D_n\|_{L^2(i\mathbb{R})}^2 = \int_{-\infty}^{\infty} n^4 \frac{y^{2s} |K(iy) - K_0(iy)|^2}{|iy + n^2 K(iy)|^2 |iy + n^2 K_0(iy)|^2} dy.$$

Use Lemma 10 to estimate the denominator and the condition (13) to estimate the numerator. By (14) we have $K(iy) - K_0(iy) = O(1/z^3)$ and obtain

$$\|z^s D_n\|_{L^2(i\mathbb{R})}^2 \prec n^4 \int_0^{\infty} \frac{y^{2s+2}}{[y^2(y^2 - n^2)^2 + n^4]^2} dy. \quad (19)$$

For $s < 9/2$ this integral converges. The main contribution gives the interval of the length $O(n)$ centered at $y = n$. Estimate it. Set

$$J = \int_{n/2}^{3n/2} \frac{y^{2s+2}}{[y^2(y^2 - n^2)^2 + n^4]^2} dy.$$

(this integral enter (19) with the factor n^4).

$$\begin{aligned} J &\asymp n^{2s-2} \int_{n/2}^{3n/2} \frac{dy}{[(y^2 - n^2)^2 + n^2]^2} \asymp n^{2s-2} \int_{n/2}^{3n/2} \frac{dy}{[n^2(y - n)^2 + n^2]^2} \\ &\asymp n^{2s-6} \int_{n/2}^{3n/2} \frac{dy}{[(y - n)^2 + 1]^2} \leq n^{2s-6} \int_{n/2}^{3n/2} \frac{dy}{[t^2 + 1]^2} = Cn^{2s-6}. \end{aligned}$$

Thus, the interval of $(n/2, 3n/2)$ gives the contribution n^{2s-2} . The theorem is proved.

This theorem give, of course, information about regularity of $\theta - \theta^0$, for example

Corollary 11

$$e^{-\varepsilon t}[\theta(x, t) - \theta_0(x, t)] \in L^2(0, \infty; \mathcal{H}_{s+1}) \cap W_2^1(0, \infty; \mathcal{H}_s)$$

5 Estimate of the derivative with respect to time

Before now we have considered a weak solution. Here we present results about a strong solutions, as, e.g., in [8]. By the definition for the strong solution the equation (1) can be considered as an equality of elements of L^2 spaces.

Proposition 12

Let (12) is fulfilled, $f_t \in \mathfrak{H}_{1,\gamma}$, $f(x, 0) = 0$, and $\xi \in \mathcal{H}_2$. Then for any $\varepsilon > \gamma$

$$\|e^{-\varepsilon t} \partial_t \theta\|_{L^2(0,\infty;L^2(0,\pi))}^2 + \|e^{-\varepsilon t} \theta\|_{L^2((0,\infty);\mathcal{H}_2)} \prec \|e^{-\varepsilon t} \partial_t f\|_{L^2((0,\infty),\mathcal{H}_1)} + \|\xi\|_{\mathcal{H}_2}.$$

In the right hand side we see, roughly speaking, the L^2 norm of $\partial_t \partial_x f$. We can slightly strengthen this estimate with the L^2 norm of $\partial_t f$ in the right hand side. It is a sharp result in the sense that it close to the estimate (7)

Theorem 13

Let (11) is fulfilled, $f_t \in \mathfrak{H}_{0,\gamma}$, $f(x, 0) = 0$, and $\xi \in \mathcal{H}_s$. Then

$$\|e^{-\varepsilon t} \partial_t \theta\|_{L^2(0,\infty;L^2(0,\pi))}^2 + \|e^{-\varepsilon t} \theta\|_{L^2((0,\infty);\mathcal{H}_2)} \prec \|e^{-\varepsilon t} \partial_t f\|_{L^2((0,\infty),\mathcal{H}_0)} + \|\xi\|_{\mathcal{H}_2}.$$

PROOF:

(i) Let $F = 0$. Then

$$\mathcal{L}[\theta'_n] = z\mathcal{L}[\theta_n] - \theta_n(0) = \left(\frac{z}{z + n^2 K(z)} - 1 \right) \xi_n = -\frac{n^2 K(z)}{z + n^2 K(z)} \xi_n.$$

$K(z)$ decreases in infinity,

$$|K(1 + iy)| \prec \frac{1}{1 + |y|},$$

Therefore we can use

$$|\mathcal{L}\theta'_n| \prec n \frac{|K|}{|G_n|} |\xi_n| \prec n |\xi_n|,$$

what gives by (17) the estimate

$$\|e^{-\varepsilon t} \theta_t\| \prec \|\xi\|_{H_{s+2}}. \quad (20)$$

(ii). Let $\xi = 0$. Then, taking into account $f(x, 0) = 0$, we obtain

$$\mathcal{L}(\theta'_n) = \frac{zF_n(z)}{z + n^2 K(z)}.$$

By (16) we have

$$\|\mathcal{L}(\theta'_n)\| \prec \|zF_n(z)\|.$$

Then

$$\|e^{-\gamma t}\theta'_n\| \prec \|e^{-\gamma t}f'_n\|_{L^2(0,\infty)}.$$

and

$$\|e^{-\gamma t}\theta_t\| \prec \|e^{-\gamma t}f'\|_{L^2(0,\infty;\mathcal{H}_s)}. \quad (21)$$

The theorem is proved.

Acknowledgements

The author grateful to Prof. V. V. Vlasov for the fruitful discussions.

References

- [1] A. A. Gavrikov, S.A. Ivanov, D.Yu. Knyazkov, V.A. Samarain , A.S. Shamaev, V. V. Vlasov, *Spectral properties of composite medaia*, Contemporary Problems of Mathematic and Mechanic, v.1, 2009, 142-159 (Russian).
- [2] M. E. Gurtin, A. C. Pipkin *A general theory of heat conduction with finite wave speeds*. Archive for Rational Mechanics and Analysis 1968; 32:113-126.
- [3] C. M. Dafermos, *Asymptotic stability in viscoelasticity*, Arch. Rational Mech. Anal., 37 (1970), 297-308.
- [4] L. Pandolfi, *The controllability of the Gurtin-Pipkin equation: a cosine operator approach*. Appl. Math. Optim. 52 (2005), no. 2, 143–165.
- [5] F.M. Vegni. *Dissipativity of a conserved phase field systems with memory*, Discrete and contiouns dynamical systems , Volume 9, Number 4, July 2003.
- [6] V. V. Vlasov, J. Wu, *Solvability and Spectral Analysis of Abstract Hyperbolic Equations with Delay*. Funct. Differ. Equ. 16 (2009), no. 4, 751-768.
- [7] Ivanov S.A., 'Wave type' spectrum of the Gurtin-Pipkin equation of the second order, arXiv; arxiv.org/abs/1002.2831, 8 p.
- [8] Vlasov V.V., Rautian N.A., Shamaev A.S., *Spectral analysis and correct solvability of abstract integrodifferential equations arising in thermophysics and acoustics*, Journal of Mathematical Sciences April 2013, Volume 190, Issue 1, pp 34-65.
- [9] Ivanov S.A., EremenkoA., *Spectra of the Gurtin-Pipkin type equations*, SIAM J. Math. Anal. 43, pp. 2296-2306.

- [10] V. Mikhailov, *Partial Differential Equations* Mir Publishers, Moscow, Russia, 1978.