

UDC 512

© 2000

T. R. Seifullin

Koszul complexes of embedded systems of polynomials and duality

(Presented by Corresponding Member of the NAS of Ukraine A. A. Letichevsky)

The author considers the dependence of Koszul complexes and dependence of dual Koszul complexes of two systems of polynomials, when one system is a part of other system, in connection with the duality in a Koszul complex established by author earlier. Whence, the dependence of Koszul complexes and dependence of dual Koszul complexes follow when one system is linearly expressed through other system. The obtained results are used in the proof of homotopic equivalence, formulated earlier by the author, of a Koszul complex and a dual Koszul complex, what happens under certain conditions.

In the present paper it is establishing the connection between Koszul complexes of embedded systems of polynomials under duality, that established in the author's paper [3]. On the basis of this it is given another proof of the connection Koszul complexes of systems of polynomials, connected by linear dependence, than has been earlier obtained by author [4]. The obtained result implies theorem 4 in [3]. By novelty of definitions and notations, for their exposition it is require a large volume, for this reason for all necessary definitions and notations we refer the reader to works [3] and [4], and also other author's works.

Let $a = (a_1, \dots, a_n)$ be a collection of elements, then we will write $|a| = n$.

Let $(\mathbf{C}; \partial)$ be a complex, and $c', c'' \in \mathbf{C}$, we will write $c' \stackrel{\partial}{\simeq} c''$ if $\exists c \in \mathbf{C}$ such that $c' - c'' = \partial[c]$.

Let \mathbf{R} be a commutative ring with unity 1 and zero 0, if $x = (x_1, \dots, x_n)$ be a collection of variables, then $\mathbf{R}[x]$ denotes a ring of polynomials in commuting variables x with coefficients in \mathbf{R} . Let \mathbf{A} be a commutative ring with unity 1 and zero 0, if $\hat{f} = (\hat{f}_1, \dots, \hat{f}_s)$ be a collection of Grassmann anticommuting variables, then $\Lambda_{\mathbf{A}}(\hat{f})$ denotes Grassmann algebra in variables \hat{f} over \mathbf{A} .

Let $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_m)$ be commuting variables; $\hat{f} = (\hat{f}_1, \dots, \hat{f}_s)$ and $\hat{g} = (\hat{g}_1, \dots, \hat{g}_t)$ be Grassmann anticommuting variables. Denote by

$$\begin{aligned} \mathbf{1}_{(x, \hat{f})}(a(y), \hat{g}b(y)) &= \mathbf{1}_{(x_1, \dots, x_n, \hat{f}_1, \dots, \hat{f}_s)}(a_1(y), \dots, a_n(y), \hat{g}b_1(y), \dots, \hat{g}b_s(y)) = \\ &= \exp(a(y)x_* + \hat{g}b(y)\hat{f}_*) = \exp\left(\sum_k a_k(y)x_*^k + \sum_{i,j} \hat{g}_j b_i^j(y)\hat{f}_*^i\right) \end{aligned}$$

linear over \mathbf{R} homomorphism $\Lambda_{\mathbf{R}[x]}(\hat{f}) \rightarrow \Lambda_{\mathbf{R}[y]}(\hat{g})$, such that $1 \mapsto 1$, $\forall k = 1, n : x_k \mapsto a_k(y)$, $\forall i = 1, s : \hat{f}_i \mapsto \sum_j \hat{g}_j b_i^j(y)$, and we call it an exponent.

Let $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$; $\partial : \hat{u} \mapsto (x - y) = (x_1 - y_1, \dots, x_n - y_n)$; $F(x) \in \mathbf{R}[x]$; denote by $\hat{u}\nabla F(x, y) = \sum_{k=1}^n \hat{u}_k \nabla^k F(x, y)$ such element $\in \mathbf{C}(x, y, \hat{u})$, that

$$\partial[\hat{u}\nabla F(x, y)] = \partial\left[\sum_{k=1}^n \hat{u}_k \nabla^k F(x, y)\right] = \sum_{k=1}^n (x_k - y_k) \nabla^k F(x, y) = F(x) - F(y).$$

Lemma 1. Let \mathbf{A} be a commutative ring with unity 1 and zero 0; $\hat{f} = (\hat{f}_1, \dots, \hat{f}_s)$, $\hat{g} = (\hat{g}_1, \dots, \hat{g}_t)$ be collections of anticommuting variables, $\forall k = 1, n : \forall i = 1, s : a_k^i \in \mathbf{A}$ and $\forall k = 1, n : \forall j = 1, t : b_k^j \in \mathbf{A}$, then

$$\frac{1}{\hat{f}} \det \|\hat{f}_*\| \det \|\hat{f}a + \hat{g}b\| = \det \begin{vmatrix} a & \hat{f}_* \\ \hat{g}b & 0 \end{vmatrix}.$$

Proof.

$$\begin{aligned} \frac{1}{\hat{f}} \det \|\hat{f}_*\| \det \|\hat{f}a + \hat{g}b\| &= \frac{1}{\hat{f}'} \det \|\hat{f}'_*\| \det \|\hat{f}'a + \hat{g}b\| \exp(\hat{f}'\hat{f}_*) = \\ &= \frac{1}{\hat{f}'} \frac{1}{\hat{g}'} \exp(\hat{g}\hat{g}'_*) \det \|\hat{f}'_*\| \cdot \det \|\hat{f}'a + \hat{g}'b\| \exp(\hat{f}'\hat{f}_*) = \det \begin{vmatrix} a & \hat{f}_* \\ \hat{g}b & 0 \end{vmatrix}. \end{aligned}$$

Lemma 2. Let \mathbf{A} be a commutative ring with unity 1 and zero 0; $\hat{f} = (\hat{f}_1, \dots, \hat{f}_s)$, $\hat{g} = (\hat{g}_1, \dots, \hat{g}_t)$ be collections of anticommuting variables, $\forall i = 1, s : \forall j = 1, t : b_i^j \in \mathbf{A}$, then:

$$1) \frac{1}{\hat{f}} \det \|\hat{f}_*\| \det \|\hat{f} - \hat{g}b\| = \det \|\hat{g}b\hat{f}_*\| = \exp(\hat{g}b\hat{f}_*);$$

$$2) \top_{\widehat{f}} \det \|\widehat{f}_*\| \det \|\widehat{f} - \widehat{g}b\| = (\widehat{g})^0.$$

Proof 1. By virtue of lemma 1

$$\begin{aligned} \perp_{\widehat{f}} \det \|\widehat{f}_*\| \det \|\widehat{f} - \widehat{g}b\| &= \det \left\| \begin{array}{c|c} E_s & \widehat{f}_* \\ \hline -\widehat{g}b & 0 \end{array} \right\| = \det \left\| \begin{array}{c|c} E_s & \widehat{0}_* \\ \hline -\widehat{g}b & \widehat{g}b\widehat{f}_* \end{array} \right\| \\ &\quad \parallel \quad \parallel \\ &= \det \left\| \begin{array}{c|c} E_s & \widehat{f}_* \\ \hline \widehat{0} & \widehat{g}b\widehat{f}_* \end{array} \right\| = \det \left\| \begin{array}{c|c} E_s & \widehat{0}_* \\ \hline \widehat{0} & \widehat{g}b\widehat{f}_* \end{array} \right\| = \det \|\widehat{g}b\widehat{f}_*\|, \end{aligned}$$

where $\forall i, i' = 1, s : (E_s)_{i'}^i = \widehat{f}_*^i \cdot \widehat{f}_{i'} = (i = i')$.

Proof 2. By virtue of 1) of this lemma

$$\top_{\widehat{f}} \det \|\widehat{f}_*\| \det \|\widehat{f} - \widehat{g}b\| = \top_{\widehat{f}} (\perp_{\widehat{f}} \det \|\widehat{f}_*\| \det \|\widehat{f} - \widehat{g}b\|) (\widehat{f})^0 = \top_{\widehat{f}} \exp(\widehat{g}b\widehat{f}_*) (\widehat{f})^0 = (\widehat{g})^0.$$

Lemma 3. Let \mathbf{R} be a commutative ring with unity 1 and zero 0; $x = (x_1, \dots, x_n)$, $f(x) = (f_1(x), \dots, f_s(x)) \in \mathbf{R}[x]^s$, $\partial : \widehat{f}_x \mapsto f(x)$, then

$$\partial \left[(x)^0 \det \|\widehat{f}_*^x\| \right] = - (f(x) \widehat{f}_*^x) \det \|\widehat{f}_*^x\| = 0.$$

Theorem 1. Let \mathbf{R} be a commutative ring with unity 1 and zero 0; $x = (x_1, \dots, x_n)$, $f(x) = (f_1(x), \dots, f_s(x)) \in \mathbf{R}[x]^s$, $F(x) = (F_1(x), \dots, F_t(x)) \in \mathbf{R}[x]^t$; $\partial : \widehat{f}_x, \widehat{f}'_x \mapsto f(x)$, $\widehat{F}_x, \widehat{F}'_x \mapsto F(x)$, then

$$\partial \left[x^0 \det \|\widehat{F}'_*\| \exp(\widehat{f}_x \widehat{f}'_x) \right] = 0 \quad \& \quad \partial \left[x^0 (\widehat{F}_x)^0 \exp(\widehat{f}_x \widehat{f}'_x) \right] = 0$$

and, hence, the maps

$$\mathbf{C}(x, \widehat{f}_x, \widehat{F}_x) \ni c(x, \widehat{f}_x, \widehat{F}_x) \mapsto \top_{\widehat{f}'_x \widehat{F}'_x} \top_{x^0} \det \|\widehat{F}'_*\| \exp(\widehat{f}_x \widehat{f}'_x) c(x, \widehat{f}'_x, \widehat{F}'_x) =$$

$$= \top_{\widehat{F}_x} \det \|\widehat{F}_*^x\| c(x, \widehat{f}_x, \widehat{F}_x) \in \mathbf{C}(x, \widehat{f}_x),$$

$$\mathbf{C}(x_*, \widehat{f}_*^x, \widehat{F}_*^x) \ni c(x_*, \widehat{f}_*^x) \mapsto \perp_x \top_{\widehat{f}'_x} \top_{x^0} \det \|\widehat{F}_*^x\| \exp(\widehat{f}'_x \widehat{f}_*^x) c(x_*, \widehat{f}'_x) =$$

$$= \det \|\widehat{F}_*^x\| c(x_*, \widehat{f}_*^x) \in \mathbf{C}(x_*, \widehat{f}_*^x, \widehat{F}_*^x),$$

$$\mathbf{C}(x, \widehat{f}_x) \ni c(x, \widehat{f}_x) \mapsto \top_{\widehat{f}'_x} x^0 (\widehat{F}_x)^0 \exp(\widehat{f}_x \widehat{f}'_x) c(x, \widehat{f}'_x) =$$

$$= (\widehat{F}_x)^0 c(x, \widehat{f}_x) \in \mathbf{C}(x, \widehat{f}_x, \widehat{F}_x),$$

$$\mathbf{C}(x_*, \widehat{f}_*^x, \widehat{F}_*^x) \ni c(x_*, \widehat{f}_*^x, \widehat{F}_*^x) \mapsto \perp_x \top_{\widehat{f}'_x} \top_{\widehat{F}'_x} x^0 (\widehat{F}'_x)^0 \exp(\widehat{f}'_x \widehat{f}_*^x) c(x_*, \widehat{f}'_x, \widehat{F}'_x) =$$

$$= \top_{\widehat{F}_x} (\widehat{F}_x)^0 c(x_*, \widehat{f}_*^x, \widehat{F}_*^x) \in \mathbf{C}(x_*, \widehat{f}_*^x)$$

are complex morphisms.

Proof. The equality $\partial \left[x^0 \det \|\widehat{F}'_*\| \exp(\widehat{f}_x \widehat{f}'_x) \right] = 0$ holds, since

$\partial \left[x^0 \det \|\widehat{F}'_x\| \right] = 0$ (lemma 3) and $\partial \left[x^0 \exp(\widehat{f}_x \widehat{f}'_x) \right] = 0$, and collections of adjoint variables of factors has empty intersection.

The equality $\partial \left[x^0 (\widehat{F}_x)^0 \exp(\widehat{f}_x \widehat{f}'_x) \right] = 0$ holds, since $\partial \left[x^0 (\widehat{F}_x)^0 \right] = 0$ and $\partial \left[x^0 \exp(\widehat{f}_x \widehat{f}'_x) \right] = 0$.

Theorem 2. Let \mathbf{R} be a commutative ring with unity 1 and zero 0; $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n)$, $f(x) = (f_1(x), \dots, f_s(x)) \in \mathbf{R}[x]^s$, $F(x) = (F_1(x), \dots, F_t(x)) \in \mathbf{R}[x]^t$; $\partial : \widehat{f}_x, \widehat{f}'_x \mapsto f(x)$, $\widehat{F}_x, \widehat{F}'_x \mapsto F(x)$, $\widehat{f}_y, \widehat{f}'_y \mapsto f(y)$, $\widehat{F}_y, \widehat{F}'_y \mapsto F(y)$, $\widehat{u} \mapsto (x - y) = (x_1 - y_1, \dots, x_n - y_n)$, then

$$\begin{aligned} 1) \quad & \prod_{\widehat{f}_y} \prod_{\widehat{F}_y} \det \left\| \begin{array}{cc} \nabla F(x, y) & \nabla f(x, y) \\ \widehat{F}_x - \widehat{F}_y & \widehat{f}_x - \widehat{f}_y \end{array} \right\| y^0 \det \|\widehat{f}_y^y\| \exp(\widehat{F}'_y \widehat{F}_y^y) = \\ & = \prod_{\widehat{F}'_x} x^0 (\widehat{f}_x)^0 \exp(\widehat{F}_x \widehat{F}'_x) \det \left\| \begin{array}{cc} \nabla F(x, y) & \\ \widehat{F}'_x - \widehat{F}'_y & \end{array} \right\|, \\ 2) \quad & \prod_{\widehat{f}_y} \det \left\| \begin{array}{cc} \nabla f(x, y) & \\ \widehat{f}_x - \widehat{f}_y & \end{array} \right\| y^0 (\widehat{F}'_y)^0 \exp(\widehat{f}'_y \widehat{f}_y^y) = \\ & = \prod_{\widehat{f}'_x} \prod_{\widehat{F}'_x} (-1)^{|F||x|} x^0 \det \|\widehat{F}'_x\| \exp(\widehat{f}_x \widehat{f}'_x) \det \left\| \begin{array}{cc} \nabla F(x, y) & \nabla f(x, y) \\ \widehat{F}'_x - \widehat{F}'_y & \widehat{f}'_x - \widehat{f}'_y \end{array} \right\|, \end{aligned}$$

hence,

$$\begin{aligned} 1') \quad & \prod_y \prod_{\widehat{f}_y} \prod_{\widehat{F}_y} \det \left\| \begin{array}{cc} \nabla F(x, y) & \nabla f(x, y) \\ \widehat{F}_x - \widehat{F}_y & \widehat{f}_x - \widehat{f}_y \end{array} \right\| \left(\det \|\widehat{f}_y^y\| c(y_*, \widehat{F}_y^y) \right) = \\ & = (\widehat{f}_x)^0 \left(\prod_y \prod_{\widehat{F}_y} \det \left\| \begin{array}{cc} \nabla F(x, y) & \\ \widehat{F}_x - \widehat{F}_y & \end{array} \right\| c(y_*, \widehat{F}_y^y) \right), \\ 2') \quad & \prod_y \prod_{\widehat{f}_y} \det \left\| \begin{array}{cc} \nabla f(x, y) & \\ \widehat{f}_x - \widehat{f}_y & \end{array} \right\| \left(\prod_{\widehat{F}_y} (\widehat{F}_y)^0 c(y_*, \widehat{f}_y^y, \widehat{F}_y^y) \right) = \\ & = \prod_{\widehat{F}_x} (-1)^{|F||x|} \det \|\widehat{F}_x\| \left(\prod_y \prod_{\widehat{f}_y} \prod_{\widehat{F}_y} \det \left\| \begin{array}{cc} \nabla F(x, y) & \nabla f(x, y) \\ \widehat{F}_x - \widehat{F}_y & \widehat{f}_x - \widehat{f}_y \end{array} \right\| c(y_*, \widehat{f}_y^y, \widehat{F}_y^y) \right). \end{aligned}$$

Proof 1. We have

$$\begin{aligned} & \prod_{\widehat{f}_y} \prod_{\widehat{F}_y} \det \left\| \begin{array}{cc} \nabla F(x, y) & \nabla f(x, y) \\ \widehat{F}_x - \widehat{F}_y & \widehat{f}_x - \widehat{f}_y \end{array} \right\| y^0 \det \|\widehat{f}_y^y\| \exp(\widehat{F}'_y \widehat{F}_y^y) = \\ & = \prod_{\widehat{f}_y} \det \left\| \begin{array}{cc} \nabla F(x, y) & \nabla f(x, y) \\ \widehat{F}_x - \widehat{F}'_y & \widehat{f}_x - \widehat{f}_y \end{array} \right\| \det \|\widehat{f}_y^y\| = \\ & = \prod_{\widehat{f}_y} \prod_{\widehat{u}} \det \|\widehat{u}_*\| \det \|\widehat{F}_x - \widehat{F}'_y - \widehat{u} \nabla F(x, y)\| \det \|\widehat{f}_x - \widehat{f}_y - \widehat{u} \nabla f(x, y)\| \det \|\widehat{f}_y^y\| = \\ & = \prod_{\widehat{u}} \det \|\widehat{u}_*\| \det \|\widehat{F}_x - \widehat{F}'_y - \widehat{u} \nabla F(x, y)\| \left(\prod_{\widehat{f}_y} \det \|\widehat{f}_x - \widehat{f}_y - \widehat{u} \nabla f(x, y)\| \det \|\widehat{f}_y^y\| \right) = \\ & = \prod_{\widehat{u}} \det \|\widehat{u}_*\| \det \|\widehat{F}_x - \widehat{F}'_y - \widehat{u} \nabla F(x, y)\| \left(\prod_{\widehat{f}_y} \det \|\widehat{f}_y^y\| \det \|\widehat{f}_x - \widehat{f}_y + \widehat{u} \nabla f(x, y)\| \right) = \end{aligned}$$

(by virtue of 2) of lemma 2)

$$\begin{aligned}
&= \underset{\hat{u}}{\top} \det \| -\hat{u}_* \| \det \| \hat{F}_x - \hat{F}'_y - \hat{u} \nabla F(x, y) \| \left((\hat{u})^0 (\hat{f}_x)^0 \right) = \\
&= \det \left\| \frac{\nabla F(x, y)}{\hat{F}_x - \hat{F}'_y} \right\| (\hat{f}_x)^0 = \underset{\hat{F}'_x}{\top} x^0 (\hat{f}_x)^0 \exp(\hat{F}_x \hat{F}'_x) \det \left\| \frac{\nabla F(x, y)}{\hat{F}'_x - \hat{F}'_y} \right\|
\end{aligned}$$

Proof 2. We have

$$\begin{aligned}
&\underset{\hat{f}'_x \hat{F}'_x}{\top} \underset{\hat{F}'_x}{\top} x^0 \det \| \hat{F}'_x \| \exp(\hat{f}_x \hat{F}'_x) \det \left\| \frac{\nabla F(x, y)}{\hat{F}'_x - \hat{F}'_y} \quad \frac{\nabla f(x, y)}{\hat{f}'_x - \hat{f}'_y} \right\| = \\
&= \underset{\hat{F}'_x}{\top} \det \| \hat{F}'_x \| \det \left\| \frac{\nabla F(x, y)}{\hat{F}'_x - \hat{F}'_y} \quad \frac{\nabla f(x, y)}{\hat{f}'_x - \hat{f}'_y} \right\| = \\
&= \underset{\hat{F}'_x}{\top} \det \| \hat{F}'_x \| \underset{\hat{u}}{\top} \det \| -\hat{u}_* \| \det \| \hat{F}'_x - \hat{F}'_y - \hat{u} \nabla F(x, y) \| \det \| \hat{f}'_x - \hat{f}'_y - \hat{u} \nabla f(x, y) \| = \\
&= (-1)^{|F||x|} \cdot \underset{\hat{u}}{\top} \det \| -\hat{u}_* \| \left(\underset{\hat{F}'_x}{\top} \det \| \hat{F}'_x \| \det \| \hat{F}'_x - \hat{F}'_y - \hat{u} \nabla F(x, y) \| \right) \det \| \hat{f}'_x - \hat{f}'_y - \hat{u} \nabla f(x, y) \| =
\end{aligned}$$

(by virtue of 2) of lemma 2)

$$\begin{aligned}
&= (-1)^{|F||x|} \underset{\hat{u}}{\top} \det \| -\hat{u}_* \| \left((\hat{F}'_y)^0 (\hat{u})^0 \right) \det \| \hat{f}'_x - \hat{f}'_y - \hat{u} \nabla f(x, y) \| = \\
&= (-1)^{|F||x|} (\hat{F}'_y)^0 \det \left\| \frac{\nabla f(x, y)}{\hat{f}'_x - \hat{f}'_y} \right\| = (-1)^{|F||x|} \underset{\hat{f}'_y}{\top} \det \left\| \frac{\nabla f(x, y)}{\hat{f}'_x - \hat{f}'_y} \right\| y^0 (\hat{F}'_y)^0 \exp(\hat{f}'_y \hat{F}'_y).
\end{aligned}$$

Theorem 3. Let \mathbf{R} be a commutative ring with unity 1 and zero 0; $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n)$, $f(x) = (f_1(x), \dots, f_s(x)) \in \mathbf{R}[x]^s$, $F(x) = (F_1(x), \dots, F_t(x)) \in \mathbf{R}[x]^t$, $\forall j = 1, t : F_j(x) = \sum_{i=1}^s f_i(x) G_j^i(x)$; $\partial : \hat{f}_x \mapsto f(x)$, $\hat{F}_x \mapsto F(x)$, $\hat{f}_y \mapsto f(y)$, $\hat{F}_y \mapsto F(y)$, then

$$\begin{aligned}
&\underset{\hat{F}_x}{\top} \exp(\hat{f}_x G(x) \hat{F}_x^x) \det \left\| \frac{\nabla F(x, y)}{\hat{F}_x - \hat{F}_y} \right\| = \det \left\| \frac{\nabla F(x, y)}{\hat{f}_x G(x) - \hat{F}_y} \right\| \underset{\partial}{\simeq} \\
&\underset{\hat{f}_y}{\top} \det \left\| \frac{\nabla f(x, y)}{\hat{f}_x - \hat{f}_y} \right\| \det \left\| \begin{array}{c} G(y) \quad \hat{f}_y^y \\ -\hat{F}_y \quad 0 \end{array} \right\|;
\end{aligned}$$

moreover, $\partial \left[\det \left\| \frac{\nabla F(x, y)}{\hat{f}_x G(x) - \hat{F}_y} \right\| \right] = 0$, $\partial \left[\det \left\| \begin{array}{c} G(y) \quad \hat{f}_y^y \\ -\hat{F}_y \quad 0 \end{array} \right\| \right] = 0$.

In particular, for $L(x_*, \hat{F}_*^x) \in \mathbf{C}(x_*, \hat{F}_*^x)$, if $\partial \left[L(x_*, \hat{F}_*^x) \right] = 0$, then

$$\partial \left[\underset{x \hat{F}_x}{\perp} \underset{\hat{F}_x}{\top} \det \left\| \begin{array}{c} G(x) \quad \hat{f}_x^x \\ -\hat{F}_x \quad 0 \end{array} \right\| L(x_*, \hat{F}_*^x) \right] = 0$$

and

$$\begin{aligned} & \underset{\widehat{F}_x}{\top} \exp(\widehat{f}_x G(x) \widehat{F}_*^x) \underset{y \widehat{F}_y}{\top} \top \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \right\| L(y_*, \widehat{F}_*^y) = \underset{y \widehat{F}_y}{\top} \top \det \left\| \frac{\nabla F(x, y)}{\widehat{f}_x G(x) - \widehat{F}_y} \right\| L(y_*, \widehat{F}_*^y) \stackrel{\partial}{\simeq} \\ & \stackrel{\partial}{\simeq} \underset{y \widehat{f}_y}{\top} \top \det \left\| \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| \left(\underset{y \widehat{F}_y}{\perp} \top \det \left\| \begin{array}{c} G(y) \quad \widehat{f}_*^y \\ -\widehat{F}_y \quad 0 \end{array} \right\| L(y_*, \widehat{F}_*^y) \right). \end{aligned}$$

Proof. Since $\forall j = 1, t : \partial \left[-\widehat{F}_{j,x} + \widehat{f}_x G_j(x) \right] = -F_j(x) + f(x) G_j(x) = 0$, then

$$\partial \left[\det \left\| -\widehat{F}_x + \widehat{f}_x G(x) \right\| \right] = \partial \left[\prod_{j=1, t} (-\widehat{F}_{j,x} + \widehat{f}_x G_j(x)) \right] = 0.$$

Let $z = (z_1, \dots, z_n)$, $\partial : \widehat{f}_z \mapsto f(z)$, $\widehat{F}_z \mapsto F(z)$. By virtue of theorem ([3])

$$\begin{aligned} & \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \quad \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| \exp(xz_* + \widehat{f}_x \widehat{f}_*^z + \widehat{F}_x \widehat{F}_*^z) \stackrel{\partial}{\simeq} \\ & \stackrel{\partial}{\simeq} \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \quad \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| \exp(yz_* + \widehat{f}_y \widehat{f}_*^z + \widehat{F}_y \widehat{F}_*^z). \end{aligned}$$

Hence, by $\partial \left[\det \left\| -\widehat{F}_z + \widehat{f}_z G(z) \right\| \right] = 0$ we have

$$\begin{aligned} & \det \left\| \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \quad \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \right\| \det \left\| -\widehat{F}_x + \widehat{f}_x G(x) \right\| = \\ & = \underset{z \widehat{f}_z \widehat{F}_z}{\top} \top \top \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \quad \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| \exp(xz_* + \widehat{f}_x \widehat{f}_*^z + \widehat{F}_x \widehat{F}_*^z) \det \left\| -\widehat{F}_z + \widehat{f}_z G(z) \right\| \stackrel{\partial}{\simeq} \\ & \stackrel{\partial}{\simeq} \underset{z \widehat{f}_z \widehat{F}_z}{\top} \top \top \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \quad \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| \exp(yz_* + \widehat{f}_y \widehat{f}_*^z + \widehat{F}_y \widehat{F}_*^z) \det \left\| -\widehat{F}_z + \widehat{f}_z G(z) \right\| = \\ & = \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \quad \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| \det \left\| -\widehat{F}_y + \widehat{f}_y G(y) \right\|. \end{aligned}$$

Then by $\partial \left[x^0 \det \left\| \widehat{F}_*^x \right\| \right] = 0$ and $\partial \left[y^0 \det \left\| \widehat{f}_*^y \right\| \right] = 0$ (lemma 3) it holds

$$\begin{aligned} & \underset{\widehat{F}_x \widehat{f}_y}{\top} \top x^0 \det \left\| \widehat{F}_*^x \right\| \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \quad \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| y^0 \det \left\| \widehat{f}_*^y \right\| \det \left\| -\widehat{F}_x + \widehat{f}_x G(x) \right\| \stackrel{\partial}{\simeq} \\ & \stackrel{\partial}{\simeq} \underset{\widehat{F}_x \widehat{f}_y}{\top} \top x^0 \det \left\| \widehat{F}_*^x \right\| \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \quad \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| y^0 \det \left\| \widehat{f}_*^y \right\| \det \left\| -\widehat{F}_y + \widehat{f}_y G(y) \right\|. \end{aligned}$$

Consider the left part of the equality

$$\begin{aligned} & \underset{\widehat{F}_x \widehat{f}_y}{\top} \top x^0 \det \left\| \widehat{F}_*^x \right\| \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \quad \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| y^0 \det \left\| \widehat{f}_*^y \right\| \det \left\| -\widehat{F}_x + \widehat{f}_x G(x) \right\| = \\ & = (-1)^{|F|(|F|-|x|)+|F||F|} \cdot \underset{\widehat{F}_x}{\top} \left(\underset{\widehat{F}_x}{\perp} \det \left\| \widehat{F}_*^x \right\| \det \left\| \widehat{F}_x - \widehat{f}_x G(x) \right\| \right) \left(\underset{\widehat{f}_y}{\top} \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \quad \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| \det \left\| \widehat{f}_*^y \right\| \right) = \end{aligned}$$

(by virtue of 1) of theorem 2 and 1) of lemma 2)

$$\begin{aligned}
&= (-1)^{|F||x|} \underset{\widehat{F}_x}{\top} \exp(\widehat{f}_x G(x) \widehat{F}_*^x) \left((\widehat{f}_x)^0 \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \right\| \right) = \\
&= (-1)^{|F||x|} \underset{\widehat{F}_x}{\top} \det \|\widehat{f}_x G(x) \widehat{F}_*^x\| \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \right\|,
\end{aligned}$$

consider the right part of the equality

$$\begin{aligned}
&\underset{\widehat{F}_x}{\top} \underset{\widehat{f}_y}{\top} x^0 \det \|\widehat{F}_*^x\| \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \quad \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| y^0 \det \|\widehat{f}_*^y\| \det \|\widehat{F}_y + \widehat{f}_y G(y)\| = \\
&= \underset{\widehat{f}_y}{\top} \left(\underset{\widehat{F}_x}{\top} \det \|\widehat{F}_*^x\| \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \quad \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| \right) \left(\underset{\widehat{f}_y}{\perp} \det \|\widehat{f}_*^y\| \det \|\widehat{F}_y + \widehat{f}_y G(y)\| \right) =
\end{aligned}$$

(by virtue of 2) of theorem 2 and lemma 1)

$$\begin{aligned}
&= \underset{\widehat{f}_y}{\top} (-1)^{|F||x|} \left(\det \left\| \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| (\widehat{F}_y)^0 \right) \left(\det \left\| \begin{array}{cc} G(y) & \widehat{f}_*^y \\ -\widehat{F}_y & 0 \end{array} \right\| \right) = \\
&= \underset{\widehat{f}_y}{\top} (-1)^{|F||x|} \det \left\| \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| \det \left\| \begin{array}{cc} G(y) & \widehat{f}_*^y \\ -\widehat{F}_y & 0 \end{array} \right\|.
\end{aligned}$$

Hence,

$$\underset{\widehat{f}_y}{\top} \det \left\| \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| \det \left\| \begin{array}{cc} G(y) & \widehat{f}_*^y \\ -\widehat{F}_y & 0 \end{array} \right\| \underset{\widehat{F}_x}{\simeq} \underset{\widehat{F}_x}{\top} \exp(\widehat{f}_x G(x) \widehat{F}_*^x) \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \right\|.$$

Let us prove theorem 4 in [3]:

Theorem 4. Let \mathbf{R} be a commutative ring with unity 1 and zero 0; $x = (x_1, \dots, x_n)$, $f(x) = (f_1(x), \dots, f_s(x)) \in \mathbf{R}[x]^s$; $\partial : \widehat{f}_x \mapsto f(x)$, $\widehat{f}_y \mapsto f(y)$. If the ring $\mathbf{H}_0(x, \widehat{f}_x) \simeq \mathbf{R}[x]/(f(x))_x$ is a finitely generated module over \mathbf{R} , then

$$\exists e(x_*, \widehat{f}_*^x) \in \mathbf{Z}(x_*, \widehat{f}_*^x) : \underset{y}{\top} \underset{\widehat{f}_y}{\top} \det \left\| \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| e(y_*, \widehat{f}_*^y) \underset{\widehat{F}_x}{\simeq} x^0 (\widehat{f}_x)^0 = 1.$$

Proof. From commutative algebra it is known, that if the ring $\mathbf{R}[x]/(f(x))_x$ is a finitely generated module over \mathbf{R} , then for any polynomial $h(x) \in \mathbf{R}[x]$ there exists a monic, i. e. with coefficient 1 of the greatest degree of variable, polynomial $T(h) = (h)^d + a_{d-1}(h)^{d-1} + \dots + a_0(h)^0 \in \mathbf{R}[h]$ such that $T(h(x)) \in (f(x))_x$.

This is proved as follows: we take the characteristic polynomial of the matrix of the multiplication operator by $h(x)$ on $\mathbf{R}[x]/(f(x))_x$ in some finite system of generators of $\mathbf{R}[x]/(f(x))_x$ as a module over \mathbf{R} , this is a monic polynomial, a root of which is $h(x)$.

Then $\forall j = 1, n$: there exist a monic polynomial $T_j(x_j) \in \mathbf{R}[x_j]$ such that $T_j(x_j) \in (f(x))_x$, hence, $T_j(x_j) = \sum_{i=1}^s f_i(x) G_j^i(x)$. Let $\partial : \widehat{T}_{j,x_j} \mapsto T_j(x_j)$, $\widehat{T}_{j,y_j} \mapsto T_j(y_j)$, then there exists $L_j(x_*^j, \widehat{T}_*^{j,x_j}) = l_j(x_*^j) \mathbf{1}_{\widehat{T}_{j,x_j}}(0) \in \mathbf{C}(x_*^j, \widehat{T}_*^{j,x_j})$ such that

$$\partial \left[L_j(x_*^j, \widehat{T}_*^{j,x_j}) \right] = \underset{x_j}{\perp} T_j(x_j) l_j(x_*^j) \widehat{T}_*^{j,x_j} = 0$$

and

$$\prod_{y_j} \prod_{\widehat{T}_{y_j}} \det \left\| \frac{\nabla T_j(x_j, y_j)}{\widehat{T}_{j, x_j} - \widehat{T}_{j, y_j}} \right\| L_j(y_j^j, \widehat{T}_{*}^{j, y_j}) = \prod_{y_j} \det \|\nabla T_j(x_j, y_j)\| l_j(y_j^j) (\widehat{T}_{j, x_j})^0 = (x_j)^0 (\widehat{T}_{j, x_j})^0.$$

Indeed, the condition $\perp_{x_j} T_j(x_j) l_j(x_*^j) = 0$ is equivalent to the following condition:

$$\begin{aligned} \forall \delta \geq 0 : \prod_{x_j} (x_j)^\delta T_j(x_j) l_j(x_*^j) &= \\ &= \prod_{x_j} (x_j)^\delta \left((x_j)^{d_j} + a_{j, d_j-1} (x_j)^{d_j-1} + \dots + a_{j, 0} (x_j)^0 \right) l_j(x_*^j) = \\ &= l_j(x_*^j) \cdot (x_j)^{d_j+\delta} + a_{j, d_j-1} (l_j(x_*^j) \cdot (x_j)^{d_j+\delta-1}) + \dots + a_{j, 0} (l_j(x_*^j) \cdot (x_j)^\delta) = 0, \end{aligned}$$

then any values of $(\lambda_0, \dots, \lambda_{d_j-1}) \in \mathbf{R}^{d_j}$ uniquely determine $l_j(x_*^j)$ such that $\forall \delta = 0, d_j - 1 : l_j(x_*^j) \cdot (x_j)^\delta = \lambda_\delta$ and $\perp_{x_j} T_j(x_j) l_j(x_*^j) = 0$. If $l_j(x_*^j)$ such that $\forall \delta = 0, d_j - 2 : l_j(x_*^j) \cdot (x_j)^\delta = 0$ and $l_j(x_*^j) \cdot (x_j)^{d_j-1} = 1$, then $\prod_{y_j} \det \|\nabla T_j(x_j, y_j)\| l_j(y_j^j) = (x_j)^0$.

Let $F(x) = (F_1(x), \dots, F_n(x))$, where $\forall j = 1, n : F_j(x) = T_j(x_j) = \sum_{i=1}^s f_i(x) G_j^i(x)$, and let $\partial : \widehat{F}_x \mapsto F(x), \widehat{F}_y \mapsto F(y)$, then $L(x_*, \widehat{F}_*^x) = \prod_{j=1, n} L_j(x_*^j, \widehat{F}_*^{j, x}) = l(x_*) \mathbf{1}_{\widehat{F}_x}(0)$ such that $\partial [L(x_*, \widehat{F}_*^x)] = 0$, since $\forall j = 1, n : \partial [L_j(x_*^j, \widehat{F}_*^{j, x})] = 0$, and

$$\prod_y \prod_{\widehat{F}_y} \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \right\| L(y_*, \widehat{F}_*^y) = \left(\prod_y \det \|\nabla F(x, y)\| l(y_*) \right) (\widehat{F}_x)^0 = (x)^0 (\widehat{F}_x)^0,$$

since

$$\begin{aligned} \prod_y \det \|\nabla F(x, y)\| l(y_*) &= \prod_{y_1} \dots \prod_{y_n} \left(\prod_{j=1, n} \det \|\nabla T_j(x_j, y_j)\| \right) \left(\prod_{j=1, n} l_j(y_j^j) \right) = \\ &= \prod_{j=1, n} \left(\prod_{y_j} \det \|\nabla T_j(x_j, y_j)\| l_j(y_j^j) \right) = \prod_{j=1, n} (x_j)^0 = (x)^0. \end{aligned}$$

Denote by

$$\begin{aligned} e(x_*, \widehat{f}_*^x) &= \perp_x \prod_{\widehat{F}_x} \det \left\| \begin{array}{c} G(x) \quad \widehat{f}_*^x \\ -\widehat{F}_x \quad 0 \end{array} \right\| L(x_*, \widehat{F}_*^x) = \\ &= \perp_x \prod_{\widehat{F}_x} \det \left\| \begin{array}{c} G(x) \quad \widehat{f}_*^x \\ -\widehat{F}_x \quad 0 \end{array} \right\| l(x_*) \mathbf{1}_{\widehat{F}_x}(0) = \perp_x \det \|G(x) \quad \widehat{f}_*^x\| l(x_*), \end{aligned}$$

then by virtue of theorem 3 it hold: $\partial [e(x_*, \widehat{f}_*^x)] = 0$ and

$$\prod_y \prod_{\widehat{f}_y} \det \left\| \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| e(y_*, \widehat{f}_*^y) = \prod_y \prod_{\widehat{f}_y} \det \left\| \frac{\nabla f(x, y)}{\widehat{f}_x - \widehat{f}_y} \right\| \left(\prod_y \prod_{\widehat{F}_y} \det \left\| \begin{array}{c} G(y) \quad \widehat{f}_*^y \\ -\widehat{F}_y \quad 0 \end{array} \right\| L(y_*, \widehat{F}_*^y) \right) \stackrel{\partial}{\simeq}$$

$$\begin{aligned} &\stackrel{\partial}{\simeq} \top_{\widehat{F}_x} \exp(\widehat{f}_x G(x) \widehat{F}_*^x) \top_y \top_{\widehat{F}_y} \det \left\| \frac{\nabla F(x, y)}{\widehat{F}_x - \widehat{F}_y} \right\| L(y_*, \widehat{F}_*^y) = \\ &= \top_{\widehat{F}_x} \exp(\widehat{f}_x G(x) \widehat{F}_*^x) (x)^0 (\widehat{F}_x)^0 = (x)^0 (\widehat{f}_x)^0. \end{aligned}$$

1. *Seifullin, T. R.* Root functionals and root polynomials of a system of polynomials. (Russian) *Dopov. Nats. Akad. Nauk Ukraïni* – 1995, – no. 5, 5–8.
2. *Seifullin, T. R.* Root functionals and root relations of a system of polynomials. (Russian) *Dopov. Nats. Akad. Nauk Ukraïni* – 1995, – no. 6, 7–10.
3. *Seifullin, T. R.* Homology of the Koszul complex of a system of polynomial equations. (Russian) *Dopov. Nats. Akad. Nauk Ukr. Mat. Prirodozn. Tekh. Nauki* 1997, no. 9, 43–49. [arXiv:1205.0472](#) (English).
4. *Seifullin, T. R.* Koszul complexes of systems of polynomials connected by linear dependence. (Russian) *Some problems in contemporary mathematics* (Russian), 326–349, *Pr. Inst. Mat. Nats. Akad. Nauk Ukr. Mat. Zastos.*, 25, Natsional. Akad. Nauk Ukraïni, Inst. Mat., Kiev, 1998.

V. M. Glushkov Institute of Cybernetics of the NAS of Ukraine, Kiev

Received 23.02.99

E-mail: timur_sf@mail.ru