

## ON CURVES CONTAINED IN CONVEX SUBSETS OF THE PLANE

DON COPPERSMITH, GYOZO NAGY, AND SASHA RAVSKY

ABSTRACT. If  $K' \subset K$  are convex bodies of the plane then the perimeter of  $K'$  is not greater than the perimeter of  $K$ . We obtain the following generalization of this fact. Let  $K$  be a convex compact body of the plane with the perimeter  $p$  and the diameter  $d$  and  $r > 1$  be an integer. Let  $s$  be the smallest number such that for any curve of length greater than  $s$  contained in  $K$  there is a straight line intersecting the curve at least in  $r + 1$  different points. Then  $s = rp/2$  if  $r$  is even and  $s = (r - 1)p/2 + d$  if  $r$  is odd.

Let  $K$  be a compact subset of the plane. Recall that the set  $K$  is called *convex* if for each pair of points  $x, y \in K$  the segment  $[x, y]$  connecting these points belongs to  $K$ . The set  $K$  is called *strictly convex* if for each pair of points  $x, y \in K$  the interval  $(x, y) = [x, y] \setminus \{x, y\}$  belongs to the interior  $\text{int } K$  of the set  $K$ . Equivalently, the set  $K$  is strictly convex if its boundary  $dK$  contains no segments of nonzero length.

Let  $I = [0; 1]$  be the unit segment. A *curve* is a continuous map  $\varphi : I \rightarrow \mathbb{R}^2$ . Let  $0 = a_0 < \dots < a_n = 1$  be a sequence. Let  $[\varphi]$  be the broken line with the sequence of vertices  $\varphi(a_0), \dots, \varphi(a_n)$  that is the union of the segments  $\bigcup_{i=0}^{n-1} [\varphi_i; \varphi_{i+1}]$ . By  $l(a_0, \dots, a_n)$  we denote the length of the broken line  $[\varphi]$ . We say that the curve  $\varphi$  has *length*  $l(\varphi)$  if for each  $\varepsilon > 0$  there is  $\delta > 0$  such that  $|l(\varphi) - l(a_0, \dots, a_n)| < \varepsilon$  for any sequence  $0 = a_0 < \dots < a_n = 1$  with  $a_{i+1} - a_i < \delta$  for each  $0 \leq i \leq n - 1$ .

The convex subset  $K$  of the plane is called a *convex body* if  $\text{int } K \neq \emptyset$ . If  $K$  is a convex body then the boundary  $dK$  of  $K$  is an image of a curve and there exists the length  $l(dK)$  which is equal to the upper bound of the perimeters of convex polygons<sup>1</sup> inscribed in  $K$  [Ale, p. 373],[YB, p.28]. The length of the boundary we shall call the *perimeter* of the body  $K$  and shall denote as  $p(K)$ .

It is well known that  $p(K') \leq p(K)$  for every pair of convex bodies  $K$  and  $K'$  with  $K' \subset K$  ([Ale, p. 373],[YB, p.28]). This means that if the image of a curve  $\varphi : I \rightarrow K$  is a boundary of a convex body then  $l(\varphi) \leq p(K)$ . A slightly weaker claim is: if the image of a curve  $\varphi : I \rightarrow K$  is a boundary of a strictly convex body then  $l(\varphi) \leq p(K)$ . The boundary  $dK'$  of the strictly convex body  $K'$  has the property that every line intersects  $dK'$  in at most two points. We are going to generalize the claim showing that  $l(\varphi) \leq p(K)$  for every curve  $\varphi : I \rightarrow K$  with every line intersecting the image of  $\varphi$  at most in two points. This result is closely related with the following proposition.

**Proposition 1.** *Let  $S^1$  be a circle and  $\varphi : S^1 \rightarrow \mathbb{R}^2$  be a homeomorphic embedding. Then the following conditions are equivalent:*

- (1)  $\varphi(S^1)$  is a boundary of a strictly convex body.
- (2) Every line intersects the image of  $\varphi$  at most in two points.
- (3) Every line intersects the image of  $\varphi$  at most in three points.

1991 *Mathematics Subject Classification.* 52A10, 52A38.

*Key words and phrases.* convex body, strictly convex body.

<sup>1</sup>We can consider a convex polygon as the convex hull of a broken line

*Proof.* The implication (1)  $\Rightarrow$  (2) follows from the definition of the strictly convex body and the implication (2)  $\Rightarrow$  (3) is obvious. Now we show the implication (3)  $\Rightarrow$  (1). Put  $K = \text{conv } \varphi(S^1)$ . Since  $\varphi$  is a homeomorphism then  $K$  is not a segment and hence  $K$  is a convex compact body. Let  $x \in S^1$  be an arbitrary point. Let  $C \subset \mathbb{R}^2$  be the circle of radius 1 with the center at the point  $\varphi(x)$  and  $p : \mathbb{R}^2 \setminus \{\varphi(x)\} \rightarrow C$  be the radial projection. Since the map  $p$  is continuous the set  $p\varphi(S^1 \setminus \{x\})$  is connected and therefore an arc. Suppose that the length of  $p\varphi(S^1)$  is greater than  $\pi$ . Then there exist points  $x_1, x_2, x_3 \in S^1 \setminus \{x\}$  such that the point  $\varphi(x)$  lies in the interior of the triangle with the vertices  $p\varphi(x_1), p\varphi(x_2)$  and  $p\varphi(x_3)$ . Then the point  $\varphi(x)$  lies in the interior of the triangle with the vertices  $\varphi(x_1), \varphi(x_2)$  and  $\varphi(x_3)$ . Fix an orientation on the circle  $S^1$ . After a re-enumerating we may suppose that the points lie on the circle  $S^1$  in the order  $x, x_1, x_2, x_3$ . Let  $l \subset \mathbb{R}^2$  be a line separating the points  $\varphi(x)$  and  $\varphi(x_2)$  from the points  $\varphi(x_1)$  and  $\varphi(x_3)$ . Since the images under the map  $\varphi$  of the oriented arcs  $(x, x_1), (x_1, x_2), (x_2, x_3)$  and  $(x_3, x)$  are linearly connected then each of the images contain a point from the line  $l$ . Obtained contradiction shows that the length of the arc  $p\varphi(S^1 \setminus \{x\})$  is not greater than  $\pi$ .

Therefore there is a line  $l$  going through the point  $\varphi(x)$  such that one of the open half planes created by  $l$  contains no points of the set  $p\varphi(S^1 \setminus \{x\})$  and hence no points of the set  $\varphi(S^1)$ . This implies that the point  $\varphi(x)$  is a boundary point of the set  $K$ .

Hence  $\varphi(S^1) \subset dK$ . Since  $dK$  is homeomorphic to a circle [Ale, p. 372] and  $\varphi(S^1) \subset dK$  is homeomorphic to a circle the  $\varphi(S^1)$  must coincide with  $dK$ . The set  $K$  is strictly convex since its boundary  $\varphi(S^1)$  contains no segments of nonzero length.  $\square$

The main result of the paper is the following

**Theorem 1.** *Let  $K \subset \mathbb{R}^2$  be a convex compact body with perimeter  $p$ , diameter  $d$  and let  $r > 1$  be an integer. Let  $s$  be the smallest number such that for any curve  $\varphi \subset K$  of length greater than  $s$  there is a line intersecting the curve  $\varphi$  at least in  $r + 1$  different points. Then*

$$s = \begin{cases} rp/2, & \text{if } r \text{ is even} \\ (r-1)p/2 + d, & \text{if } r \text{ is odd.} \end{cases}$$

*Proof.* Fix the compact body  $K$  and the number  $r$ . Let  $p$  be the perimeter of  $K$  and  $d$  be the diameter of  $K$ .

*The upper bound.* Put  $s = rp/2$  if  $r$  is even and  $s = (r-1)p/2 + d$  if  $r$  is odd. Let  $\varphi : I \rightarrow K$  be a curve of length greater than  $s$ . By the definition of the length we can choose numbers  $0 = a_0 < \dots < a_n = 1$  such that the length of the broken line with the sequence of vertices  $\varphi(a_0), \dots, \varphi(a_n)$  is greater than  $s$ . Let  $l_1, \dots, l_n$  be the lengths of the segments of the broken line and  $\alpha_1, \dots, \alpha_n$  be the angles with these segments and the  $Ox$  axis. For an angle  $\alpha \in [0; 2\pi]$ , by  $l(\alpha)$  we denote the sum of the lengths of the projections of the segments  $l_i$  onto the line  $L_\alpha = \{(x, y) : (x, y) = t(\cos \alpha, \sin \alpha)\}$ . Thus  $l(\alpha) = \sum l_i |\cos(\alpha - \alpha_i)|$ . By  $k(\alpha)$  we denote the length of the projection of  $K$  onto the line  $L_\alpha$ . By Cauchy formula [Had, 6.1.5]

$$\int_0^{2\pi} k(\alpha) d\alpha = 2p.$$

At first suppose that  $r$  is even. To the rest of the proof it suffice to show that there is an angle  $\alpha'$  such that  $l(\alpha') > rk(\alpha')$ . Suppose the contrary. Then

$$\begin{aligned} 2rp &= r \int_0^{2\pi} k(\alpha) d\alpha \geq \int_0^{2\pi} l(\alpha) d\alpha = \int_0^{2\pi} \sum l_i |\cos(\alpha - \alpha_i)| d\alpha = \\ &= \sum \int_0^{2\pi} l_i |\cos(\alpha - \alpha_i)| d\alpha = \sum \int_0^{2\pi} l_i |\cos(\beta)| d\beta = \sum 4l_i > 4s = 2rp, \end{aligned}$$

the contradiction.


Now suppose that  $r$  is odd. Add to the broken line the segment connecting its ends. Let the length of the segment be  $l_0$  and  $\alpha_0$  be the angle with the segment and the  $Ox$  axis. To the rest of the proof it suffice to show that there is an angle  $\alpha'$  such that  $l(\alpha') > rl_0 |\cos(\alpha' - \alpha_0)| + (r-1)(k(\alpha') - l_0 |\cos(\alpha' - \alpha_0)|) = (r-1)k(\alpha') + l_0 |\cos(\alpha' - \alpha_0)|$ . That is because each line not intersecting  $l_0$  and the vertices of the broken line intersects the broken line an even number of times.

Suppose the contrary. Then

$$\begin{aligned} 2(r-1)p + 4l_0 &= \int_0^{2\pi} (r-1)k(\alpha) + l_0 |\cos(\alpha - \alpha_0)| d\alpha \geq \int_0^{2\pi} l(\alpha) d\alpha = \int_0^{2\pi} \sum l_i |\cos(\alpha - \alpha_i)| d\alpha = \\ &= \sum \int_0^{2\pi} l_i |\cos(\alpha - \alpha_i)| d\alpha = \sum \int_0^{2\pi} l_i |\cos(\beta)| d\beta = \sum 4l_i > 4s = 2(r-1)p + 4d, \end{aligned}$$

which yields the contradiction since  $l_0 \leq d$ .  $\square$

*The lower bound.* The idea of the proof is the following. Put  $n = \lfloor r/2 \rfloor$ . If  $r$  is even, let the curve go  $n$  times around the perimeter, curving slightly to avoid any straight lines but remaining convex. If  $r$  is odd, let the curve go almost  $n$  times around the perimeter, then down the diameter, again always curving slightly. The reason we go “almost”  $n$  times around the perimeter is to ensure that any straight line intersecting this “curved diameter” twice, will intersect the perimeter-curve at most  $r-2$  times rather than  $r-1$ . The picture illustrates the construction when  $K$  is a square and  $r=5$ .



Now more precisely. Let  $\varepsilon > 0$ . At first we suppose that  $r$  is even. There is a curve  $\varphi_1 : I \rightarrow K$  such that  $\varphi_1(0) = \varphi_1(1)$ ,  $|l(\varphi_1) - p| < \varepsilon$  and  $\varphi_1(I)$  is the boundary of a

strictly convex body. Choose a number  $\delta_1 > 0$  such that  $l(\varphi_1([1 - \delta_1; 1])) < \varepsilon$ . Similarly, there is a curve  $\varphi_2 : I \rightarrow K$  such that  $\varphi_2(0) = \varphi_2(1)$ ,  $|l(\varphi_1) - l(\varphi_2)| < \varepsilon$ ,  $\varphi_2(I)$  is the boundary of a strictly convex body,  $\varphi_2(0) = \varphi_1(1 - \delta)$  and  $\varphi_2((0; 1)) \subset \text{int conv } \varphi_1(I)$ . Choose a number  $\delta_2 > 0$  such that  $l(\varphi_2([1 - \delta_2; 1])) < \varepsilon$ . Similarly to the previous we can construct the curve  $\varphi_3 : I \rightarrow K$  and so on. From the curves  $\varphi_1, \dots, \varphi_n$  it is easily to construct the curve  $\varphi$  intersecting every straight line in at most  $r$  points.

Now suppose that  $r$  is odd. To construct the curve  $\varphi$  we shall proceed similarly to the case of even  $n$ . By induction we can construct the curves  $\varphi_1, \dots, \varphi_n$  such that

- (i)  $|\text{diam}(\varphi_1(I)) - \text{diam}(K)| < \varepsilon$ ,
- (ii)  $|\text{diam}(\varphi_i(I)) - \text{diam}(\varphi_{i-1}(I))| < \varepsilon$  for each  $i$ ,
- (iii) there is a number  $\alpha_i$  such that  $|\text{diam}(\varphi_i(I)) - |\varphi_i(0) - \varphi_i(\alpha_i)|| < \varepsilon$  for each  $i$ .

Now choose a number  $\delta_n > 0$  such that  $|\text{diam}(\varphi_n(I)) - |\varphi_n(1 - \delta_n) - \varphi_n(\alpha_n)|| < \varepsilon$  and  $l(\varphi_n([1 - \delta_n; 1])) < \varepsilon$ . There exists a curve  $\varphi_{n+1} : I \rightarrow K$  such that

- (i)  $\varphi_{n+1}(0) = \varphi_n(1 - \delta_n)$  and  $\varphi_{n+1}(1) = \varphi_n(\alpha_n)$ ,
- (ii)  $\varphi_{n+1}((0; 1))$  lies in the interior of the triangle with the vertices  $\varphi_n(0)$ ,  $\varphi_n(\alpha_n)$  and  $\varphi_n(1 - \delta_n)$ ,
- (iii)  $\varphi_{n+1}(I) \cup [\varphi_n(1 - \delta_n); \varphi_n(\alpha_n)]$  is the boundary of a convex body,
- (iv) no three points of the set  $\varphi_{n+1}(I)$  lie on a straight line.

From the curves  $\varphi_1, \dots, \varphi_{n+1}$  it is easy to construct the curve  $\varphi$  intersecting every straight line in at most  $r$  points.  $\square$

#### REFERENCES

- [Ale] A.D. Aleksandrov. *Inner geometry of convex surfaces*, OGIZ, 1948. (in Russian)
- [Gash] S.B. Gashkov. Problem M932 // *Kvant* – 1985. – V. 11. – P.36-37. (in Russian)
- [Had] H. Hadwiger. *Vorlesungen über inhalt, oberfläche und isoperimetrie*, Springer-Verlag, 1957. (in Russian, M.:Nauka, 1966)
- [YB] I.M. Yaglom, V.G. Boltyanskiy. *Convex figures*, GITTL, 1951. (in Russian)

*E-mail address:* [dcopper@us.ibm.com](mailto:dcopper@us.ibm.com); [gyozo.nagy@raiffeisen.hu](mailto:gyozo.nagy@raiffeisen.hu); [oravsky@mail.ru](mailto:oravsky@mail.ru)

DEPARTMENT OF MATHEMATICAL SCIENCES, IBM THOMAS J WATSON RESEARCH CENTER, YORKTOWN HEIGHTS NY 10598, USA; TECHNICAL CENTER, RAIFFEISEN HUNGARY, BUDAPEST KESMARK 11-13, 1158, HUNGARY; DEPARTMENT OF MATHEMATICS, IVAN FRANKO LVIV NATIONAL UNIVERSITY, UNIVERSYTETSKA 1, LVIV, 79000, UKRAINE