

SOME COMPANIONS OF PERTURBED OSTROWSKI TYPE INEQUALITIES FOR FUNCTIONS WHOSE SECOND DERIVATIVES ARE BOUNDED AND APPLICATIONS

WENJUN LIU AND YITING ZHU

ABSTRACT. In this paper we establish some companions of perturbed Ostrowski type integral inequalities for functions whose second derivatives are bounded. Some applications to composite quadrature rules, and to probability density functions are also given.

1. INTRODUCTION

In 1938, Ostrowski [21] established the following interesting integral inequality for differentiable mappings with bounded derivatives:

Theorem 1.1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable mapping on (a, b) whose derivative is bounded on (a, b) and denote $\|f'\|_\infty = \sup_{t \in (a, b)} |f'(t)| < \infty$. Then for all $x \in [a, b]$ we have*

$$(1.1) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty.$$

The constant $\frac{1}{4}$ is sharp in the sense that it can not be replaced by a smaller one.

In [13], Gussab and Schmeisser proved the following companion of Ostrowski's inequality:

Theorem 1.2. *Let $f : [a, b] \rightarrow \mathbb{R}$ be satisfying the Lipschitz condition, i.e., $|f(t) - f(s)| \leq M|t - s|$. Then for all $x \in [a, \frac{a+b}{2}]$ we have*

$$(1.2) \quad \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{8} + 2 \left(\frac{x - \frac{3a+b}{4}}{b-a} \right)^2 \right] (b-a)M.$$

The constant $\frac{1}{8}$ is sharp in the sense that it can not be replaced by a smaller one. In (1.2), the point $x = \frac{3a+b}{4}$ gives the best estimator.

Motivated by [13], Dragomir [8] proved some companions of Ostrowski's inequality, as follows:

Theorem 1.3. *Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous mapping on $[a, b]$. Then the following inequalities*

$$\left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \begin{cases} \left[\frac{1}{8} + 2 \left(\frac{x - \frac{3a+b}{4}}{b-a} \right)^2 \right] (b-a) \|f'\|_\infty, & f' \in L^\infty[a, b], \\ \frac{2^{1/q}}{(q+1)^{1/q}} \left[\left(\frac{x-a}{b-a} \right)^{q+1} + \left(\frac{\frac{a+b}{2}-x}{b-a} \right)^{q+1} \right]^{1/q} (b-a)^{1/q} \|f'\|_p, & p > 1, \frac{1}{p} + \frac{1}{q} = 1 \text{ and } f' \in L^p[a, b], \\ \left[\frac{1}{4} + \left| \frac{x - \frac{3a+b}{4}}{b-a} \right| \right] \|f'\|_1, & f' \in L^1[a, b] \end{cases}$$

hold for all $x \in [a, \frac{a+b}{2}]$.

2010 Mathematics Subject Classification. 26D15, 41A55, 41A80, 65C50.

Key words and phrases. perturbed Ostrowski type inequality; differentiable mapping; composite quadrature rule; probability density function.

Recently, Alomari [1] studied the companion of Ostrowski inequality (1.2) for differentiable bounded mappings. In [19], Liu established some companions of an Ostrowski type integral inequality for functions whose first derivatives are absolutely continuous and second derivatives belong to L^p ($1 \leq p \leq \infty$) spaces.

Theorem 1.4. *Let $f : [a, b] \rightarrow \mathbb{R}$ be such that f' is absolutely continuous on $[a, b]$ and $f'' \in L^\infty[a, b]$. Then for all $x \in [a, \frac{a+b}{2}]$ we have*

$$(1.3) \quad \left| \frac{f(x) + f(a+b-x)}{2} - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{96} + \frac{1}{2} \frac{\left(x - \frac{3a+b}{4}\right)^2}{(b-a)^2} \right] (b-a)^2 \|f''\|_\infty.$$

The constant $\frac{1}{96}$ is sharp in the sense that it can not be replaced by a smaller one.

For other related results, the reader may refer to [2, 3, 4, 7, 9, 11, 14, 15, 16, 17, 18, 22, 23, 24, 25, 26, 27, 28] and the references therein.

The main aim of this paper is to establish some companions of perturbed Ostrowski type integral inequalities for functions whose second derivatives are bounded (Theorem 2.1-2.5). Some applications to composite quadrature rules, and to probability density functions are also given.

2. MAIN RESULTS

To prove our main results, we need the following lemmas.

Lemma 2.1. [19] *Let $f : [a, b] \rightarrow \mathbb{R}$ be such that f' is absolutely continuous on $[a, b]$. Denote by $K(x, t) : [a, b] \rightarrow \mathbb{R}$ the kernel given by*

$$(2.1) \quad K(x, t) = \begin{cases} \frac{1}{2}(t-a)^2, & t \in [a, x], \\ \frac{1}{2}\left(t - \frac{a+b}{2}\right)^2, & t \in (x, a+b-x], \\ \frac{1}{2}(t-b)^2, & t \in (a+b-x, b], \end{cases}$$

then the identity

$$(2.2) \quad \frac{1}{b-a} \int_a^b K(x, t) f''(t) dt = \frac{1}{b-a} \int_a^b f(t) dt - \frac{f(x) + f(a+b-x)}{2} + \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2}$$

holds.

Lemma 2.2. [12, Grüss inequality] *Let $f, g : [a, b] \rightarrow \mathbb{R}$ be two integrable functions such that $\phi \leq f(t) \leq \Phi$ and $\gamma \leq g(t) \leq \Gamma$ for all $t \in [a, b]$, where ϕ, Φ, γ and Γ are constants. Then we have*

$$(2.3) \quad \left| \frac{1}{b-a} \int_a^b f(t)g(t) dt - \frac{1}{b-a} \int_a^b f(t) dt \cdot \frac{1}{b-a} \int_a^b g(t) dt \right| \leq \frac{1}{4} (\Phi - \phi)(\Gamma - \gamma).$$

Theorem 2.1. Let $f : [a, b] \rightarrow \mathbb{R}$ be such that f' is absolutely continuous on $[a, b]$. If $f'' \in L^1[a, b]$ and $\gamma \leq f''(x) \leq \Gamma, \forall x \in [a, b]$, then for all $x \in [a, \frac{a+b}{2}]$ we have

$$(2.4) \quad \begin{aligned} & \left| \frac{f(x) + f(a+b-x)}{2} - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \right. \\ & \quad \left. + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{1}{8} (\Gamma - \gamma) \left[\frac{b-a}{4} + \left| x - \frac{3a+b}{4} \right| \right]^2. \end{aligned}$$

Proof. It is clear that for all $t \in [a, b]$ and $x \in [a, \frac{a+b}{2}]$, we have

$$\begin{aligned} 0 \leq K(x, t) & \leq \max \left\{ \frac{1}{2} (x-a)^2, \frac{1}{2} \left(\frac{a+b}{2} - x \right)^2 \right\} \\ & = \frac{1}{4} \left\{ \left[(x-a)^2 + \left(\frac{a+b}{2} - x \right)^2 \right] + \left| (x-a)^2 - \left(\frac{a+b}{2} - x \right)^2 \right| \right\} \\ & = \frac{1}{2} \left[\left(x - \frac{3a+b}{4} \right)^2 + \frac{(b-a)^2}{16} + \frac{b-a}{2} \left| x - \frac{3a+b}{4} \right| \right] \\ & = \frac{1}{2} \left[\frac{b-a}{4} + \left| x - \frac{3a+b}{4} \right| \right]^2. \end{aligned}$$

Applying Lemma 2.2 to the functions $K(x, \cdot)$ and $f''(\cdot)$, we get

$$(2.5) \quad \begin{aligned} & \left| \frac{1}{b-a} \int_a^b K(x, t) f''(t) dt - \frac{1}{b-a} \int_a^b K(x, t) dt \cdot \frac{1}{b-a} \int_a^b f''(t) dt \right| \\ & \leq \frac{1}{8} (\Gamma - \gamma) \left[\frac{b-a}{4} + \left| x - \frac{3a+b}{4} \right| \right]^2 \end{aligned}$$

for all $x \in [a, \frac{a+b}{2}]$. By a simple calculation, we obtain

$$(2.6) \quad \frac{1}{b-a} \int_a^b f''(t) dt = \frac{f'(b) - f'(a)}{b-a}$$

and

$$(2.7) \quad \frac{1}{b-a} \int_a^b K(x, t) dt = \frac{1}{2} \left(x - \frac{3a+b}{4} \right)^2 + \frac{(b-a)^2}{96}.$$

Combining (2.2), (2.5)-(2.7), we obtain (2.4) as required. \square

Corollary 2.1. In the inequality (2.4), choose

(1) $x = \frac{3a+b}{4}$, we get

$$(2.8) \quad \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} + \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{96} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{128} (\Gamma - \gamma) (b-a)^2.$$

(2) $x = a$, we get

$$\left| \frac{f(a) + f(b)}{2} - \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{12} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{32} (\Gamma - \gamma) (b-a)^2,$$

which is better than [5, Corollary 2.3] since a smaller estimator is given here.

(3) $x = \frac{a+b}{2}$, we get

$$\left| f\left(\frac{a+b}{2}\right) + \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{24} - \frac{1}{b-a} \int_a^b f(t)dt \right| \leq \frac{1}{32}(\Gamma - \gamma)(b-a)^2,$$

which is the inequality given in [5, Corollay 2.2].

Corollary 2.2. Let f as in Theorem 2.1. Additionally, if f is symmetric about $x = \frac{a+b}{2}$, i.e., $f(a+b-x) = f(x)$, then we have

$$\begin{aligned} & \left| f(x) - \left(x - \frac{3a+b}{4}\right) f'(x) + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t)dt \right| \\ & \leq \frac{1}{8}(\Gamma - \gamma) \left[\frac{b-a}{4} + \left| x - \frac{3a+b}{4} \right| \right]^2 \end{aligned}$$

for all $x \in [a, \frac{a+b}{2}]$.

Theorem 2.2. Let $f : [a, b] \rightarrow \mathbb{R}$ be such that f' is absolutely continuous on $[a, b]$. If $f'' \in L^1[a, b]$ and $\gamma \leq f''(x) \leq \Gamma, \forall x \in [a, b]$, then for all $x \in [a, \frac{a+b}{2}]$ we have

$$\begin{aligned} & \left| \frac{f(x) + f(a+b-x)}{2} - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \right. \\ & \quad \left. + \frac{\Gamma + \gamma}{2} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t)dt \right| \\ (2.9) \quad & \leq \frac{\Gamma - \gamma}{2} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right]. \end{aligned}$$

Proof. From (2.2) and (2.7), we have

$$\begin{aligned} & \frac{1}{b-a} \int_a^b K(x, t)[f''(t) - C]dt \\ & = \frac{1}{b-a} \int_a^b f(t)dt - \frac{f(x) + f(a+b-x)}{2} \\ & \quad + \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} - C \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right]. \end{aligned}$$

Let $C = \frac{\Gamma + \gamma}{2}$, we get

$$\begin{aligned} & \left| \frac{f(x) + f(a+b-x)}{2} - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \right. \\ & \quad \left. + \frac{\Gamma + \gamma}{2} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t)dt \right| \\ (2.10) \quad & \leq \max_{t \in [a, b]} |f''(t) - C| \frac{1}{b-a} \int_a^b |K(x, t)|dt. \end{aligned}$$

We also have

$$(2.11) \quad \max_{t \in [a, b]} |f''(t) - C| \leq \frac{\Gamma - \gamma}{2}$$

and

$$(2.12) \quad \frac{1}{b-a} \int_a^b |K(x, t)|dt = \frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96}.$$

Therefore, from (2.10)-(2.12), we obtain the desired inequality (2.9). \square

Corollary 2.3. *In the inequality (2.9), choose*

(1) $x = \frac{3a+b}{4}$, we get

$$(2.13) \quad \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} + \frac{\Gamma + \gamma}{2} \frac{(b-a)^2}{96} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{192} (\Gamma - \gamma) (b-a)^2.$$

(2) $x = a$, we get

$$\left| \frac{f(a) + f(b)}{2} - \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{8} + \frac{\Gamma + \gamma}{2} \frac{(b-a)^2}{24} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{48} (\Gamma - \gamma) (b-a)^2.$$

(3) $x = \frac{a+b}{2}$, we get

$$\left| f\left(\frac{a+b}{2}\right) + \frac{\Gamma + \gamma}{2} \frac{(b-a)^2}{24} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{48} (\Gamma - \gamma) (b-a)^2.$$

Corollary 2.4. *Let f as in Theorem 2.2. Additionally, if f is symmetric about $x = \frac{a+b}{2}$, then we have*

$$\begin{aligned} & \left| f(x) - \left(x - \frac{3a+b}{4}\right) f'(x) + \frac{\Gamma + \gamma}{2} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{\Gamma - \gamma}{2} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right], \end{aligned}$$

for all $x \in [a, \frac{a+b}{2}]$.

Theorem 2.3. *Let $f : [a, b] \rightarrow \mathbb{R}$ be such that f' is absolutely continuous on $[a, b]$. If $f'' \in L^1[a, b]$ and $\gamma \leq f''(x) \leq \Gamma, \forall x \in [a, b]$, then for all $x \in [a, \frac{a+b}{2}]$ we have*

$$(2.14) \quad \begin{aligned} & \left| \frac{f(x) + f(a+b-x)}{2} - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \right. \\ & \quad \left. + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq (S - \gamma) \left[\frac{(b-a)^2}{48} + \frac{b-a}{4} \left| x - \frac{3a+b}{4} \right| \right] \end{aligned}$$

and

$$(2.15) \quad \begin{aligned} & \left| \frac{f(x) + f(a+b-x)}{2} - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \right. \\ & \quad \left. + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq (\Gamma - S) \left[\frac{(b-a)^2}{48} + \frac{b-a}{4} \left| x - \frac{3a+b}{4} \right| \right], \end{aligned}$$

where $S = (f'(b) - f'(a))/(b-a)$.

Proof. From (2.2), (2.6), (2.7), it follows that

$$(2.16) \quad \begin{aligned} & \frac{1}{b-a} \int_a^b K(x,t) f''(t) dt - \frac{1}{(b-a)^2} \int_a^b K(x,t) dt \int_a^b f''(t) dt \\ &= \frac{1}{b-a} \int_a^b f(t) dt - \frac{f(x) + f(a+b-x)}{2} + \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \\ & \quad - \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right]. \end{aligned}$$

We denote

$$(2.17) \quad R_n(x) = \frac{1}{b-a} \int_a^b K(x,t) f''(t) dt - \frac{1}{(b-a)^2} \int_a^b K(x,t) dt \int_a^b f''(t) dt.$$

If $C \in \mathbb{R}$ is an arbitrary constant, then we have

$$(2.18) \quad R_n(x) = \frac{1}{b-a} \int_a^b (f''(t) - C) \left[K(x,t) - \frac{1}{b-a} \int_a^b K(x,s) ds \right] dt.$$

Furthermore, we have

$$(2.19) \quad |R_n(x)| \leq \frac{1}{b-a} \max_{t \in [a,b]} \left| K(x,t) - \frac{1}{b-a} \int_a^b K(x,s) ds \right| \int_a^b |f''(t) - C| dt.$$

To compute

$$(2.20) \quad \begin{aligned} & \max_{t \in [a,b]} \left| K(x,t) - \frac{1}{b-a} \int_a^b K(x,s) ds \right| \\ &= \max \left\{ \left| \frac{1}{2}(x-a)^2 - \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] \right|, \right. \\ & \quad \left. \left| \frac{1}{2} \left(\frac{a+b}{2} - x\right)^2 - \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] \right|, \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] \right\} \\ &= \max \left\{ \frac{b-a}{24} |6x - 5a - b|, \frac{b-a}{12} |3x - 2a - b|, \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] \right\}, \end{aligned}$$

we denote

$$y_1 = \frac{b-a}{24} |6x - 5a - b|, \quad y_2 = \frac{b-a}{12} |3x - 2a - b|, \quad y_3 = \frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96}.$$

A direct computation gives that

$$(2.21) \quad \begin{cases} y_2 \geq \max\{y_1, y_3\}, & x \in \left[a, \frac{3a+b}{4} \right], \\ y_1 > \max\{y_2, y_3\}, & x \in \left(\frac{3a+b}{4}, \frac{a+b}{2} \right]. \end{cases}$$

Therefore, we get

$$(2.22) \quad \max_{t \in [a,b]} \left| K(x,t) - \frac{1}{b-a} \int_a^b K(x,s) ds \right| = \max\{y_1, y_2\} = \left[\frac{(b-a)^2}{48} + \frac{b-a}{4} \left| x - \frac{3a+b}{4} \right| \right].$$

We also have

$$(2.23) \quad \int_a^b |f''(t) - \gamma| dt = (S - \gamma)(b-a)$$

and

$$(2.24) \quad \int_a^b |f''(t) - \Gamma| dt = (\Gamma - S)(b - a).$$

Therefore, we obtain (2.14) and (2.15) by using (2.16) – (2.19), (2.22) – (2.24) and choosing $C = \gamma$ and $C = \Gamma$ in (2.19), respectively. \square

Corollary 2.5. *Under the assumptions of Theorem 2.3, choose*

(1) $x = \frac{3a+b}{4}$, we have

$$(2.25) \quad \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} + \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{96} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{48} (S - \gamma)(b - a)^2,$$

$$(2.26) \quad \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} + \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{96} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{48} (\Gamma - S)(b - a)^2.$$

(2) $x = a$, we have

$$\left| \frac{f(a) + f(b)}{2} - \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{12} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{12} (S - \gamma)(b - a)^2,$$

$$\left| \frac{f(a) + f(b)}{2} - \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{12} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{12} (\Gamma - S)(b - a)^2.$$

(3) $x = \frac{a+b}{2}$, we have

$$\left| f\left(\frac{a+b}{2}\right) + \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{24} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{12} (S - \gamma)(b - a)^2,$$

$$\left| f\left(\frac{a+b}{2}\right) + \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{24} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{12} (\Gamma - S)(b - a)^3.$$

Corollary 2.6. *Let f as in Theorem 2.3. Additionally, if f is symmetric about $x = \frac{a+b}{2}$, then for all $x \in [a, \frac{a+b}{2}]$ we have*

$$\begin{aligned} & \left| f(x) - \left(x - \frac{3a+b}{4}\right) f'(x) + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq (S - \gamma) \left[\frac{(b-a)^2}{48} + \frac{b-a}{4} \left| x - \frac{3a+b}{4} \right| \right] \end{aligned}$$

and

$$\begin{aligned} & \left| f(x) - \left(x - \frac{3a+b}{4}\right) f'(x) + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq (\Gamma - S) \left[\frac{(b-a)^2}{48} + \frac{b-a}{4} \left| x - \frac{3a+b}{4} \right| \right]. \end{aligned}$$

Theorem 2.4. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a thrice continuously differentiable mapping in (a, b) with $f''' \in L^2[a, b]$. Then for all $x \in [a, \frac{a+b}{2}]$ we have*

$$\begin{aligned}
& \left| \frac{f(x) + f(a+b-x)}{2} - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \right. \\
& \quad \left. + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\
(2.27) \quad & \leq \frac{1}{\pi} \|f'''\|_2 \left\{ \frac{1}{320} (a+b-2x)^5 + \frac{1}{10} (x-a)^5 - (b-a) \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right]^2 \right\}^{1/2}.
\end{aligned}$$

Proof. Let $R_n(x)$ be defined by (2.17). From (2.16), we get

$$\begin{aligned}
(2.28) \quad R_n(x) &= \frac{1}{b-a} \int_a^b f(t) dt - \frac{f(x) + f(a+b-x)}{2} + \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \\
&\quad - \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right].
\end{aligned}$$

If we choose $C = f''((a+b)/2)$ in (2.18) and use the Cauchy inequality, then we get

$$\begin{aligned}
(2.29) \quad & |R_n(x)| \\
& \leq \frac{1}{b-a} \int_a^b \left| f''(t) - f''\left(\frac{a+b}{2}\right) \right| \left| K(x,t) - \frac{1}{b-a} \int_a^b K(x,s) ds \right| dt \\
& \leq \frac{1}{b-a} \left[\int_a^b \left(f''(t) - f''\left(\frac{a+b}{2}\right) \right)^2 dt \right]^{1/2} \left[\int_a^b \left(K(x,t) - \frac{1}{b-a} \int_a^b K(x,s) ds \right)^2 dt \right]^{1/2}.
\end{aligned}$$

We can use the Diaz-Metcalf inequality to get

$$\int_a^b \left(f''(t) - f''\left(\frac{a+b}{2}\right) \right)^2 dt \leq \frac{(b-a)^2}{\pi^2} \|f'''\|_2^2.$$

We also have

$$\begin{aligned}
(2.30) \quad & \int_a^b \left(K(x,t) - \frac{1}{b-a} \int_a^b K(x,s) ds \right)^2 dt \\
& = \int_a^b K(x,t)^2 dt - (b-a) \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right]^2 \\
& = \frac{1}{320} (a+b-2x)^5 + \frac{1}{10} (x-a)^5 - (b-a) \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right]^2.
\end{aligned}$$

Therefore, using the above relations, we obtain (2.27). \square

Corollary 2.7. *Under the assumptions of Theorem 2.4, choose*

(1) $x = \frac{3a+b}{4}$, we have

$$(2.31) \quad \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} + \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{96} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)^{5/2}}{48\pi\sqrt{5}} \|f'''\|_2.$$

(2) $x = a$, we have

$$\left| \frac{f(a) + f(b)}{2} - \frac{f'(b) - f'(a)}{b-a} \frac{(b-a)^2}{12} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)^{5/2}}{12\pi\sqrt{5}} \|f'''\|_2.$$

(3) $x = \frac{a+b}{2}$, we have

$$\left| f\left(\frac{a+b}{2}\right) + \frac{f'(b) - f'(a)(b-a)^2}{b-a} \frac{1}{24} - \frac{1}{b-a} \int_a^b f(t)dt \right| \leq \frac{(b-a)^{5/2}}{12\pi\sqrt{5}} \|f'''\|_2.$$

Corollary 2.8. Let f as in Theorem 2.3. Additionally, if f is symmetric about $x = \frac{a+b}{2}$, i.e., $f(a+b-x) = f(x)$, then for all $x \in [a, \frac{a+b}{2}]$ we have

$$\begin{aligned} & \left| f(x) - \left(x - \frac{3a+b}{4}\right) f'(x) + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t)dt \right| \\ & \leq \frac{1}{\pi} \|f'''\|_2 \left\{ \frac{1}{320}(a+b-2x)^5 + \frac{1}{10}(x-a)^5 - (b-a) \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right]^2 \right\}^{1/2}. \end{aligned}$$

Theorem 2.5. Let $f : [a, b] \rightarrow \mathbb{R}$ be such that f' is absolutely continuous on $[a, b]$ with $f'' \in L^2[a, b]$. Then for all $x \in [a, \frac{a+b}{2}]$ we have

$$\begin{aligned} & \left| \frac{f(x) + f(a+b-x)}{2} - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \right. \\ & \quad \left. + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t)dt \right| \\ (2.32) \quad & \leq \frac{\sqrt{\sigma(f'')}}{b-a} \left\{ \frac{1}{320}(a+b-2x)^5 + \frac{1}{10}(x-a)^5 - (b-a) \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right]^2 \right\}^{1/2}, \end{aligned}$$

where $\sigma(f'')$ is defined by

$$(2.33) \quad \sigma(f'') = \|f''\|_2^2 - \frac{(f'(b) - f'(a))^2}{b-a} = \|f''\|_2^2 - S^2(b-a)$$

and S is defined in Theorem 2.3.

Proof. Let $R_n(x)$ be defined by (2.17). If we choose $C = \frac{1}{b-a} \int_a^b f''(s)ds$ in (2.18) and use the Cauchy inequality and (2.30), then we get

$$\begin{aligned} & |R_n(x)| \\ & \leq \frac{1}{b-a} \int_a^b \left| f''(t) - \frac{1}{b-a} \int_a^b f''(s)ds \right| \left| K(x, t) - \frac{1}{b-a} \int_a^b K(x, s)ds \right| dt \\ & \leq \frac{1}{b-a} \left[\int_a^b \left(f''(t) - \frac{1}{b-a} \int_a^b f''(s)ds \right)^2 dt \right]^{1/2} \left[\int_a^b \left(K(x, t) - \frac{1}{b-a} \int_a^b K(x, s)ds \right)^2 dt \right]^{1/2} \\ & \leq \frac{\sqrt{\sigma(f'')}}{b-a} \left\{ \frac{1}{320}(a+b-2x)^5 + \frac{1}{10}(x-a)^5 - (b-a) \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right]^2 \right\}^{1/2}. \end{aligned}$$

□

Corollary 2.9. Under the assumptions of Theorem 2.5, choose

(1) $x = \frac{3a+b}{4}$, we have

$$(2.34) \quad \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} + \frac{f'(b) - f'(a)(b-a)^2}{b-a} \frac{1}{96} - \frac{1}{b-a} \int_a^b f(t)dt \right| \leq \frac{(b-a)^{3/2}}{48\sqrt{5}} \sqrt{\sigma(f'')}.$$

(2) $x = a$, we have

$$\left| \frac{f(a) + f(b)}{2} - \frac{f'(b) - f'(a)(b-a)^2}{b-a} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)^{3/2}}{12\sqrt{5}} \sqrt{\sigma(f'')}.$$

(3) $x = \frac{a+b}{2}$, we have

$$\left| f\left(\frac{a+b}{2}\right) + \frac{f'(b) - f'(a)(b-a)^2}{b-a} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)^{3/2}}{12\sqrt{5}} \sqrt{\sigma(f'')}.$$

Corollary 2.10. *Let f as in Theorem 2.3. Additionally, if f is symmetric about $x = \frac{a+b}{2}$, then for all $x \in [a, \frac{a+b}{2}]$ we have*

$$\begin{aligned} & \left| f(x) - \left(x - \frac{3a+b}{4}\right) f'(x) + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{\sqrt{\sigma(f'')}}{b-a} \left\{ \frac{1}{320} (a+b-2x)^5 + \frac{1}{10} (x-a)^5 - (b-a) \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right]^2 \right\}^{1/2}. \end{aligned}$$

3. APPLICATION TO COMPOSITE QUADRATURE RULES

Let $I_n : a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ be a partition of the interval $[a, b]$ and $h_i = x_{i+1} - x_i$ ($i = 0, 1, 2, \dots, n-1$).

Consider the perturbed composite quadrature rules

$$(3.1) \quad Q_n^1(I_n, f) = \frac{1}{2} \sum_{i=0}^{n-1} \left[f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i + \sum_{i=0}^{n-1} \frac{f'(x_{i+1}) - f'(x_i)}{96} h_i^2$$

and

$$(3.2) \quad Q_n^2(I_n, f) = \frac{1}{2} \sum_{i=0}^{n-1} \left[f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i + \frac{\Gamma + \gamma}{192} \sum_{i=0}^{n-1} h_i^3.$$

The following result holds.

Theorem 3.1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be such that f' is absolutely continuous on $[a, b]$. If $f'' \in L^1[a, b]$ and $\gamma \leq f''(x) \leq \Gamma, \forall x \in [a, b]$, then for all $x \in [a, \frac{a+b}{2}]$ we have*

$$\int_a^b f(t) dt = Q_n^1(I_n, f) + R_n^1(I_n, f),$$

where $Q_n^1(I_n, f)$ is defined by formula (3.1), and the remainder $R_n^1(I_n, f)$ satisfies the estimate

$$(3.3) \quad |R_n^1(I_n, f)| \leq \frac{\Gamma - \gamma}{128} \sum_{i=0}^{n-1} h_i^3.$$

Proof. Applying inequality (2.8) to the intervals $[x_i, x_{i+1}]$, then we get

$$\begin{aligned} & \left| \int_{x_i}^{x_{i+1}} f(t) dt - \frac{1}{2} \left[f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i - \frac{f'(x_{i+1}) - f'(x_i)}{96} h_i^2 \right| \\ & \leq \frac{\Gamma - \gamma}{128} h_i^3 \end{aligned}$$

for $i = 0, 1, 2, \dots, n-1$. Now summing over i from 0 to $n-1$ and using the triangle inequality, we get (3.3). \square

Theorem 3.2. Let $f : [a, b] \rightarrow \mathbb{R}$ be such that f' is absolutely continuous on $[a, b]$. If $f'' \in L^1[a, b]$ and $\gamma \leq f''(x) \leq \Gamma, \forall x \in [a, b]$, then for all $x \in [a, \frac{a+b}{2}]$ we have

$$\int_a^b f(t)dt = Q_n^2(I_n, f) + R_n^2(I_n, f),$$

where $Q_n^2(I_n, f)$ is defined by formula (3.2), and the remainder $R_n^2(I_n, f)$ satisfies the estimate

$$(3.4) \quad |R_n^2(I_n, f)| \leq \frac{\Gamma - \gamma}{192} \sum_{i=0}^{n-1} h_i^3.$$

Proof. Applying inequality (2.13) to the intervals $[x_i, x_{i+1}]$, then we get

$$\begin{aligned} & \left| \int_{x_i}^{x_{i+1}} f(t)dt - \frac{1}{2} \left[f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i - \frac{\Gamma + \gamma}{192} h_i^3 \right| \\ & \leq \frac{\Gamma - \gamma}{192} h_i^3 \end{aligned}$$

for $i = 0, 1, 2, \dots, n-1$. Now summing over i from 0 to $n-1$ and using the triangle inequality, we get (3.4). \square

Theorem 3.3. Let $f : [a, b] \rightarrow \mathbb{R}$ be such that f' is absolutely continuous on $[a, b]$. If $f'' \in L^1[a, b]$ and $\gamma \leq f''(x) \leq \Gamma, \forall x \in [a, b]$, then for all $x \in [a, \frac{a+b}{2}]$ we have

$$\int_a^b f(t)dt = Q_n^1(I_n, f) + R_n^1(I_n, f),$$

where $Q_n^1(I_n, f)$ is defined by formula (3.1), and the remainder $R_n^1(I_n, f)$ satisfies the estimate

$$(3.5) \quad |R_n^1(I_n, f)| \leq \frac{1}{48} (S - \gamma) \sum_{i=0}^{n-1} h_i^3$$

and

$$(3.6) \quad |R_n^1(I_n, f)| \leq \frac{1}{48} (S - \gamma) \sum_{i=0}^{n-1} h_i^3.$$

Proof. Applying inequality (2.25) and (2.26) to the intervals $[x_i, x_{i+1}]$, then we get

$$\begin{aligned} & \left| \int_{x_i}^{x_{i+1}} f(t)dt - \frac{1}{2} \left[f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i - \frac{f'(x_{i+1}) - f'(x_i)}{96} h_i^2 \right| \\ & \leq \frac{1}{48} (S - \gamma) h_i^3 \end{aligned}$$

and

$$\begin{aligned} & \left| \int_{x_i}^{x_{i+1}} f(t)dt - \frac{1}{2} \left[f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i - \frac{f'(x_{i+1}) - f'(x_i)}{96} h_i^2 \right| \\ & \leq \frac{1}{48} (\Gamma - S) h_i^3, \end{aligned}$$

for $i = 0, 1, 2, \dots, n-1$. Now summing over i from 0 to $n-1$ and using the triangle inequality, we get (3.5) and (3.6). \square

Theorem 3.4. Let $f : [a, b] \rightarrow \mathbb{R}$ be a thrice continuously differentiable mapping in (a, b) with $f''' \in L^2[a, b]$. Then for all $x \in [a, \frac{a+b}{2}]$ we have

$$\int_a^b f(t)dt = Q_n^1(I_n, f) + R_n^1(I_n, f),$$

where $Q_n^1(I_n, f)$ is defined by formula (3.1), and the remainder $R_n^1(I_n, f)$ satisfies the estimate

$$(3.7) \quad |R_n^1(I_n, f)| \leq \frac{\|f'''\|_2}{48\pi\sqrt{5}} \sum_{i=0}^{n-1} h_i^{7/2}.$$

Proof. Applying inequality (2.31) to the intervals $[x_i, x_{i+1}]$, then we get

$$\begin{aligned} & \left| \int_{x_i}^{x_{i+1}} f(t)dt - \frac{1}{2} \left[f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i - \frac{f'(x_{i+1}) - f'(x_i)}{96} h_i^2 \right| \\ & \leq \frac{h_i^{7/2}}{48\pi\sqrt{5}} \|f'''\|_2, \end{aligned}$$

for $i = 0, 1, 2, \dots, n-1$. Now summing over i from 0 to $n-1$ and using the triangle inequality, we get (3.7). \square

Theorem 3.5. Let $f : [a, b] \rightarrow \mathbb{R}$ be such that f' is absolutely continuous on $[a, b]$ with $f'' \in L^2[a, b]$. Then for all $x \in [a, \frac{a+b}{2}]$ we have

$$\int_a^b f(t)dt = Q_n^1(I_n, f) + R_n^1(I_n, f),$$

where $Q_n^1(I_n, f)$ is defined by formula (3.1), and the remainder $R_n^1(I_n, f)$ satisfies the estimate

$$(3.8) \quad |R_n^1(I_n, f)| \leq \frac{\sqrt{\sigma(f'')}}{48\sqrt{5}} \sum_{i=0}^{n-1} h_i^{5/2}.$$

Proof. Applying inequality (2.34) to the intervals $[x_i, x_{i+1}]$, then we get

$$\begin{aligned} & \left| \int_{x_i}^{x_{i+1}} f(t)dt - \frac{1}{2} \left[f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i - \frac{f'(x_{i+1}) - f'(x_i)}{96} h_i^2 \right| \\ & \leq \frac{h_i^{5/2}}{48\sqrt{5}} \sqrt{\sigma(f'')}, \end{aligned}$$

for $i = 0, 1, 2, \dots, n-1$. Now summing over i from 0 to $n-1$ and using the triangle inequality, we get (3.8). \square

In order to compare the error between these two methods and composite trapezoidal formula, we apply the above methods and composite trapezoidal formula to specific examples. For this purpose consider the following Table:

TABLE 1. Numerical results

$f(x)$	n	$[a, b]$	$\int_a^b f(x)dx$	T_n	Error of T_n	Q_n^1	Q_n^2	Error of Q_n^1 and Q_n^2
$\cos x - x$	20	$[0, \frac{\pi}{2}]$	-0.233701	-0.234215	5.14E-4	-0.233636	-0.233636	6.5E-5
$e^{2x} \cos(e^x)$	20	$[0, 1]$	-1.176887	-1.181466	4.579E-3	-1.176316	-1.176316	5.71E-4
$\frac{1}{x^4+4x^2+3}$	10	$[0, 1]$	0.241549	0.241393	1.56E-4	0.241569	0.241569	2E-5
$\tan x + x$	20	$[0, \frac{\pi}{4}]$	0.654999	0.655127	1.28E-4	0.654983	0.654983	7E-6
$\ln(x^2 + 1)$	20	$[-1, 1]$	0.527887	0.529554	1.667E-3	0.527679	0.527679	2.08E-4

4. APPLICATION TO PROBABILITY DENSITY FUNCTIONS

Now, let X be a random variable taking values in the finite interval $[a, b]$, with the probability density function $f : [a, b] \rightarrow [0, 1]$ and with the cumulative distribution function

$$F(x) = \Pr(X \leq x) = \int_a^x f(t)dt.$$

The following results hold:

Theorem 4.1. *With the assumptions of Theorem 2.1, we have*

$$(4.1) \quad \begin{aligned} & \left| \frac{1}{2}[F(x) + F(a+b-x)] - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \right. \\ & \left. + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{b - E(X)}{b-a} \right| \\ & \leq \frac{1}{8}(\Gamma - \gamma) \left[\frac{b-a}{4} + \left|x - \frac{3a+b}{4}\right| \right]^2. \end{aligned}$$

for all $x \in [a, \frac{a+b}{2}]$, where $E(X)$ is the expectation of X .

Proof. By (2.4) on choosing $f = F$ and taking into account

$$E(X) = \int_a^b t dF(t) = b - \int_a^b F(t)dt,$$

we obtain (4.1). □

Corollary 4.1. *Under the assumptions of Theorem 4.1 with $x = \frac{3a+b}{4}$, we have*

$$(4.2) \quad \left| \frac{1}{2} \left[F\left(\frac{3a+b}{4}\right) + F\left(\frac{a+3b}{4}\right) \right] + \frac{b-a}{96} [f'(b) - f'(a)] - \frac{b - E(x)}{b-a} \right| \leq \frac{\Gamma - \gamma}{128} (b-a)^2.$$

Theorem 4.2. *With the assumptions of Theorem 2.2, we have*

$$(4.3) \quad \begin{aligned} & \left| \frac{1}{2}[F(x) + F(a+b-x)] - \left(x - \frac{3a+b}{4}\right) \frac{f(x) - f(a+b-x)}{2} \right. \\ & \left. + \frac{\Gamma + \gamma}{2} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{b - E(X)}{b-a} \right| \\ & \leq \frac{\Gamma - \gamma}{2} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] \end{aligned}$$

for all $x \in [a, \frac{a+b}{2}]$, where $E(X)$ is the expectation of X .

Proof. By (2.9) on choosing $f = F$ and taking into account

$$E(X) = \int_a^b t dF(t) = b - \int_a^b F(t)dt,$$

we obtain (4.3). □

Corollary 4.2. *Under the assumptions of Theorem 4.2 with $x = \frac{3a+b}{4}$, we have*

$$(4.4) \quad \left| \frac{1}{2} \left[F\left(\frac{3a+b}{4}\right) + F\left(\frac{a+3b}{4}\right) \right] + \frac{\Gamma + \gamma}{192} (b-a)^2 - \frac{b - E(x)}{b-a} \right| \leq \frac{\Gamma - \gamma}{192} (b-a)^2.$$

Theorem 4.3. *With the assumptions of Theorem 2.3, we have*

$$\begin{aligned}
& \left| \frac{1}{2}[F(x) + F(a+b-x)] - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \right. \\
& \quad \left. + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{b-E(X)}{b-a} \right| \\
(4.5) \quad & \leq (S-\gamma) \left[\frac{(b-a)^2}{48} + \frac{b-a}{4} \left| x - \frac{3a+b}{4} \right| \right]
\end{aligned}$$

and

$$\begin{aligned}
& \left| \frac{1}{2}[F(x) + F(a+b-x)] - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \right. \\
& \quad \left. + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{b-E(X)}{b-a} \right| \\
(4.6) \quad & \leq (\Gamma-S) \left[\frac{(b-a)^2}{48} + \frac{b-a}{4} \left| x - \frac{3a+b}{4} \right| \right]
\end{aligned}$$

for all $x \in [a, \frac{a+b}{2}]$, where $E(X)$ is the expectation of X .

Proof. By (2.14) and (2.15) on choosing $f = F$ and taking into account

$$E(X) = \int_a^b t dF(t) = b - \int_a^b F(t) dt,$$

we obtain (4.5) and (4.6). □

Corollary 4.3. *Under the assumptions of Theorem 4.3 with $x = \frac{3a+b}{4}$, we have*

$$(4.7) \quad \left| \frac{1}{2} \left[F\left(\frac{3a+b}{4}\right) + F\left(\frac{a+3b}{4}\right) \right] + \frac{b-a}{96} [f'(b) - f'(a)] - \frac{b-E(x)}{b-a} \right| \leq \frac{1}{48} (S-\gamma) (b-a)^2$$

and

$$(4.8) \quad \left| \frac{1}{2} \left[F\left(\frac{3a+b}{4}\right) + F\left(\frac{a+3b}{4}\right) \right] + \frac{b-a}{96} [f'(b) - f'(a)] - \frac{b-E(x)}{b-a} \right| \leq \frac{1}{48} (\Gamma-S) (b-a)^2.$$

Theorem 4.4. *With the assumptions of Theorem 2.4, we have*

$$\begin{aligned}
& \left| \frac{1}{2}[F(x) + F(a+b-x)] - \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} \right. \\
& \quad \left. + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right] - \frac{b-E(X)}{b-a} \right| \\
(4.9) \quad & \leq \frac{1}{\pi} \|f'''\|_2 \left\{ \frac{1}{320} (a+b-2x)^5 + \frac{1}{10} (x-a)^5 - (b-a) \left[\frac{1}{2} \left(x - \frac{3a+b}{4}\right)^2 + \frac{(b-a)^2}{96} \right]^2 \right\}^{1/2}
\end{aligned}$$

for all $x \in [a, \frac{a+b}{2}]$, where $E(X)$ is the expectation of X .

Proof. By (2.27) on choosing $f = F$ and taking into account

$$E(X) = \int_a^b t dF(t) = b - \int_a^b F(t) dt,$$

we obtain (4.9). □

Corollary 4.4. Under the assumptions of Theorem 4.4 with $x = \frac{3a+b}{4}$, we have

$$(4.10) \quad \left| \frac{1}{2} \left[F \left(\frac{3a+b}{4} \right) + F \left(\frac{a+3b}{4} \right) \right] + \frac{b-a}{96} [f'(b) - f'(a)] - \frac{b-E(x)}{b-a} \right| \leq \frac{(b-a)^{5/2}}{48\pi\sqrt{5}} \|f'''\|_2.$$

Theorem 4.5. With the assumptions of Theorem 2.5, we have

$$(4.11) \quad \left| \frac{1}{2} [F(x) + F(a+b-x)] - \left(x - \frac{3a+b}{4} \right) \frac{f'(x) - f'(a+b-x)}{2} + \frac{f'(b) - f'(a)}{b-a} \left[\frac{1}{2} \left(x - \frac{3a+b}{4} \right)^2 + \frac{(b-a)^2}{96} \right] - \frac{b-E(X)}{b-a} \right| \leq \frac{\sqrt{\sigma(f'')}}{b-a} \left\{ \frac{1}{320} (a+b-2x)^5 + \frac{1}{10} (x-a)^5 - (b-a) \left[\frac{1}{2} \left(x - \frac{3a+b}{4} \right)^2 + \frac{(b-a)^2}{96} \right]^2 \right\}^{1/2}$$

for all $x \in [a, \frac{a+b}{2}]$, where $E(X)$ is the expectation of X .

Proof. By (2.32) on choosing $f = F$ and taking into account

$$E(X) = \int_a^b t dF(t) = b - \int_a^b F(t) dt,$$

we obtain (4.11). □

Corollary 4.5. Under the assumptions of Theorem 4.5 with $x = \frac{3a+b}{4}$, we have

$$(4.12) \quad \left| \frac{1}{2} \left[F \left(\frac{3a+b}{4} \right) + F \left(\frac{a+3b}{4} \right) \right] + \frac{b-a}{96} [f'(b) - f'(a)] - \frac{b-E(x)}{b-a} \right| \leq \frac{(b-a)^{3/2}}{48\sqrt{5}} \sqrt{\sigma(f'')}.$$

Acknowledgments. This work was partly supported by the National Natural Science Foundation of China (Grant No. 40975002) and the Natural Science Foundation of the Jiangsu Higher Education Institutions (Grant No. 09KJB110005).

REFERENCES

- [1] M. W. Alomari, A companion of Ostrowski's inequality with applications, *Transylv. J. Math. Mech.* **3** (2011), no. 1, 9–14.
- [2] M. W. Alomari, A companion of Ostrowski's inequality for mappings whose first derivatives are bounded and applications in numerical integration, *RGMA Res. Rep. Coll.*, 14 (2011) article 57.
- [3] M. W. Alomari, A generalization of companion inequality of Ostrowski's type for mappings whose first derivatives are bounded and applications in numerical integration, *RGMA Res. Rep. Coll.*, 14 (2011) article 58.
- [4] N. S. Barnett, S. S. Dragomir and I. Gomm, A companion for the Ostrowski and the generalised trapezoid inequalities, *Math. Comput. Modelling* **50** (2009), no. 1-2, 179–187.
- [5] P. Cerone, S. S. Dragomir, J. Roulmliotis, An inequality of Ostrowski type for mappings whose second derivatives are bounded and applications, *East Asian Math. J.* **15** (1999), no 1–9.
- [6] S. S. Dragomir, A companions of Ostrowski's inequality for functions of bounded variation and applications, *RGMA Res. Rep. Coll.*, 5 (2002), Supp., article 28.
- [7] S. S. Dragomir, Ostrowski type inequalities for functions defined on linear spaces and applications for semi-inner products, *J. Concr. Appl. Math.* **3** (2005), no. 1, 91–103.
- [8] S. S. Dragomir, Some companions of Ostrowski's inequality for absolutely continuous functions and applications, *Bull. Korean Math. Soc.* **42** (2005), no. 2, 213–230.
- [9] S. S. Dragomir, Ostrowski's type inequalities for continuous functions of selfadjoint operators on Hilbert spaces: a survey of recent results, *Ann. Funct. Anal.* **2** (2011), no. 1, 139–205.
- [10] S. S. Dragomir and A. Sofo, An integral inequality for twice differentiable mappings and applications, *Tamkang J. Math.* **31** (2000), no. 4, 257–266.
- [11] J. Duoandikoetxea, A unified approach to several inequalities involving functions and derivatives, *Czechoslovak Math. J.* **51(126)** (2001), no. 2, 363–376.
- [12] G. Grüss, Über das Maximum des absoluten Betrages von $\frac{1}{b-a} \int_a^b f(x)g(x) dx - \frac{1}{(b-a)^2} \int_a^b f(x) dx \int_a^b g(x) dx$, *Math. Z.* **39** (1935), no. 1, 215–226.

- [13] A. Guessab and G. Schmeisser, Sharp integral inequalities of the Hermite-Hadamard type, *J. Approx. Theory* **115** (2002), no. 2, 260–288.
- [14] V. N. Huy and Q. -A. Ngô, New bounds for the Ostrowski-like type inequalities, *Bull. Korean Math. Soc.* **48** (2011), no. 1, 95–104.
- [15] W. J. Liu, Several error inequalities for a quadrature formula with a parameter and applications, *Comput. Math. Appl.* **56** (2008), no. 7, 1766–1772.
- [16] W. J. Liu, Some weighted integral inequalities with a parameter and applications, *Acta Appl. Math.* **109** (2010), no. 2, 389–400.
- [17] W. J. Liu, Q. L. Xue and S. F. Wang, New generalization of perturbed Ostrowski type inequalities and applications, *J. Appl. Math. Comput.* **32** (2010), no. 1, 157–169.
- [18] Z. Liu, Note on a paper by N. Ujević, *Appl. Math. Lett.* **20** (2007), no. 6, 659–663.
- [19] Z. Liu, Some companions of an Ostrowski type inequality and applications, *JIPAM. J. Inequal. Pure Appl. Math.* **10** (2009), no. 2, Article 52, 12 pp.
- [20] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Inequalities involving functions and their integrals and derivatives*, Mathematics and its Applications (East European Series), 53, Kluwer Acad. Publ., Dordrecht, 1991.
- [21] A. Ostrowski, Über die Absolutabweichung einer differentiierebaren Funktion von ihrem Integralmittelwert, *Comment. Math. Helv.* **10** (1937), no. 1, 226–227.
- [22] M. Z. Sarikaya, On the Ostrowski type integral inequality, *Acta Math. Univ. Comenian. (N.S.)* **79** (2010), no. 1, 129–134.
- [23] M. Z. Sarikaya, New weighted Ostrowski and Čebyšev type inequalities on time scales, *Comput. Math. Appl.* **60** (2010), no. 5, 1510–1514.
- [24] E. Set and M. Z. Sarikaya, On the generalization of Ostrowski and Grüss type discrete inequalities, *Comput. Math. Appl.* **62** (2011), no. 1, 455–461.
- [25] K.-L. Tseng, S.-R. Hwang and S. S. Dragomir, Generalizations of weighted Ostrowski type inequalities for mappings of bounded variation and their applications, *Comput. Math. Appl.* **55** (2008), no. 8, 1785–1793.
- [26] K.-L. Tseng, S.-R. Hwang, G.-S. Yang and Y.-M. Chou, Weighted Ostrowski integral inequality for mappings of bounded variation, *Taiwanese J. Math.* **15** (2011), no. 2, 573–585.
- [27] N. Ujević, New bounds for the first inequality of Ostrowski-Grüss type and applications, *Comput. Math. Appl.* **46** (2003), no. 2-3, 421–427.
- [28] S. W. Vong, A note on some Ostrowski-like type inequalities, *Comput. Math. Appl.* **62** (2011), no. 1, 532–535.

(W. J. Liu) COLLEGE OF MATHEMATICS AND STATISTICS, NANJING UNIVERSITY OF INFORMATION SCIENCE AND TECHNOLOGY, NANJING 210044, CHINA
E-mail address: wjliu@nuist.edu.cn

(Y. T. Zhu) COLLEGE OF MATHEMATICS AND STATISTICS, NANJING UNIVERSITY OF INFORMATION SCIENCE AND TECHNOLOGY, NANJING 210044, CHINA