

SEMIORTHOGONAL DECOMPOSITIONS FOR TWISTED GRASSMANNIANS

SANGHOON BAEK

ABSTRACT. In this article, we present semiorthogonal decompositions for twisted forms of grassmannians.

1. INTRODUCTION

In [5] Orlov gave the semiorthogonal decompositions for projective, grassmann, and flag bundles, which generalize the full exceptional collections on the corresponding varieties by Beilinson [1] and Kapranov [4].

In the case of projective bundles, Bernardara [2] extended the semiorthogonal decomposition to the twisted forms. In this paper, we present, in a similar way, semiorthogonal decompositions for twisted forms of grassmannians. The proof of the main theorem uses the study of K -theory of twisted grassmannians by Levine-Srinivas-Weyman [6] and Panin [7].

Notation and Conventions In this paper, a partition means a nonincreasing sequence of numbers. Given a partition $\alpha = (\alpha_i)$, the conjugate of α , written by α^* , is the partition (α_j^*) , where $\alpha_j^* = |\{i \mid \alpha_i \geq j\}|$. For a scheme X over Y , we denote by $\Delta(X/Y)$ the diagonal of $X \times_Y X$. We abbreviate $\mathcal{F} \boxtimes \mathcal{G} = \pi_1^* \mathcal{F} \otimes \pi_2^* \mathcal{G}$ for sheaves \mathcal{F}, \mathcal{G} over X , where π_i is the projection of $X \times_Y X$. For a scheme X , we denoted by $D(X)$ the bounded derived category of coherent sheaves on X .

2. PRELIMINARIES

In this section we recall some definitions from [3]. Let F be a field. We fix a triangulated F -linear category \mathcal{T} .

Given a full subcategory \mathcal{S} of \mathcal{T} , the (right) *orthogonal complement*, denoted by \mathcal{S}^\perp , is a full subcategory whose objects are objects T in \mathcal{T} satisfying

$$\mathrm{Hom}_{\mathcal{T}}(S, T) = 0$$

for all $S \in \mathcal{S}$.

A full triangulated subcategory \mathcal{S} of \mathcal{T} is (right) *admissible* if for any $T \in \mathcal{T}$ there is a distinguished triangle

$$S \rightarrow T \rightarrow S^\perp \rightarrow S[1]$$

with $S \in \mathcal{S}$ and $S^\perp \in \mathcal{S}^\perp$, i.e. $\mathcal{T} = \langle \mathcal{S}, \mathcal{S}^\perp \rangle$.

Now, we consider a sequence $(\mathcal{S}_1, \dots, \mathcal{S}_n)$ of admissible subcategories of \mathcal{T} . This sequence is called *semiorthogonal* if

$$(1) \quad \mathcal{S}_i \subset \mathcal{S}_j^\perp$$

for $1 \leq i < j \leq n$. A semiorthogonal sequence $(\mathcal{S}_1, \dots, \mathcal{S}_n)$ of \mathcal{T} is called *semiorthogonal decomposition* if $\mathcal{T} = \langle \mathcal{S}_1, \dots, \mathcal{S}_n \rangle$.

For example, the sequence $(\mathcal{S}, \mathcal{S}^\perp)$ for an admissible full triangulated subcategory \mathcal{S} gives a semiorthogonal decomposition of \mathcal{T} . By definition, any generating sequence $(\mathcal{S}_1, \dots, \mathcal{S}_n)$ of full subcategories of \mathcal{T} satisfying (1) is a semiorthogonal decomposition of \mathcal{T} .

3. TWISTED GRASSMANNIANS

Let X be a Noetherian scheme of characteristic 0 and let \mathcal{A} be a sheaf of Azumaya algebras of rank n^2 over X . For an integer $1 \leq k < n$, a twisted grassmannian $p : \mathrm{Gr}(k, \mathcal{A}) \rightarrow X$ is defined by the representable functor from the category *Schemes*/ X of schemes over X to the category of *Sets* of sets given by $(Y \xrightarrow{\phi} X) \mapsto$ the set of sheaves of left ideals \mathcal{I} of $\phi^*\mathcal{A}$ such that $\phi^*\mathcal{A}/\mathcal{I}$ is a locally free \mathcal{O}_Y -modules of rank $n(n-k)$.

There is an étale covering $i : U \rightarrow X$ and a locally free sheaf \mathcal{E} of rank n over U with the following pullback diagram

$$\begin{array}{ccc} \mathrm{Gr}(k, \mathcal{E}nd(\mathcal{E})) & \xrightarrow{j} & \mathrm{Gr}(k, \mathcal{A}) \\ \downarrow q & & \downarrow p \\ U & \xrightarrow{i} & X, \end{array}$$

where $\mathrm{Gr}(k, \mathcal{E}nd(\mathcal{E}))$ is naturally isomorphic to $\mathrm{Gr}(k, \mathcal{E})$.

Consider the tautological exact sequence of sheaves on $\mathrm{Gr}(k, \mathcal{E})$

$$0 \rightarrow \mathcal{R} \rightarrow q^*\mathcal{E} \rightarrow \mathcal{T} \rightarrow 0,$$

where $\mathrm{rank}(\mathcal{R}) = k$. For a partition $\alpha = (\alpha_1, \dots, \alpha_k)$ with $0 \leq \alpha_i \leq n-k$, we denote by S^α the Schur functor for α . We define $S(\alpha)$ to be the full subcategory of $\mathrm{D}(\mathrm{Gr}(k, \mathcal{A}))$ generated by \mathcal{M} in $\mathrm{D}(\mathrm{Gr}(k, \mathcal{A}))$ satisfying

$$\mathcal{M}|_{\mathrm{Gr}(k, \mathcal{E})} \simeq q^*\mathcal{N} \otimes S^\alpha\mathcal{R},$$

for some $\mathcal{N} \in \mathrm{D}(U)$.

Lemma 3.1. *Let α, α' be two distinct partitions with $0 \leq \alpha_i, \alpha'_i \leq n-k$ and let $\mathcal{M} \in S(\alpha)$ and $\mathcal{M}' \in S(\alpha')$. Then $R\mathcal{H}om(\mathcal{M}, \mathcal{M}') = 0$.*

Proof. Let $\mathcal{M}|_{\mathrm{Gr}(k, \mathcal{E})} \simeq q^*\mathcal{N} \otimes S^\alpha\mathcal{R}$ and $\mathcal{M}'|_{\mathrm{Gr}(k, \mathcal{E})} \simeq q^*\mathcal{N}' \otimes S^{\alpha'}\mathcal{R}$.

By the Littlewood-Richardson rule, the partition β of an irreducible summand $S^\beta\mathcal{R}$ of $\mathcal{H}om(S^\alpha\mathcal{R}, S^{\alpha'}\mathcal{R})$ is of the form $(\beta_1, \dots, \beta_k)$ with $-(n-k) \leq \beta_i \leq n-k$. Hence, by the Borel-Bott-Weil theorem as in [4, Lemma 3.2] we have

$$(2) \quad Rq_*(\mathcal{H}om(S^\alpha\mathcal{R}, S^{\alpha'}\mathcal{R})) = 0.$$

It is enough to show the result locally. By the adjoint property of Rq_* and q^* , projection formula, and (2), we have

$$\begin{aligned} R\mathcal{H}om(\mathcal{M}|_{\mathrm{Gr}(k, \mathcal{E})}, \mathcal{M}'|_{\mathrm{Gr}(k, \mathcal{E})}) &= R\mathcal{H}om(q^*\mathcal{N}, q^*\mathcal{N}' \otimes \mathcal{H}om(S^\alpha\mathcal{R}, S^{\alpha'}\mathcal{R})) \\ &= R\mathcal{H}om(\mathcal{N}, Rq_*(q^*\mathcal{N}' \otimes \mathcal{H}om(S^\alpha\mathcal{R}, S^{\alpha'}\mathcal{R}))) \\ &= R\mathcal{H}om(\mathcal{N}, \mathcal{N}' \otimes Rq_*(\mathcal{H}om(S^\alpha\mathcal{R}, S^{\alpha'}\mathcal{R}))) \\ &= 0. \end{aligned} \quad \square$$

Theorem 3.2. *Let $(S(\alpha)|_\alpha = (\alpha_1, \dots, \alpha_k), 0 \leq \alpha_i \leq n - k)$ be a sequence of the full subcategories of $D(\mathrm{Gr}(k, \mathcal{A}))$ by the lexicographical order on α . Then this sequence gives a semiorthogonal decomposition of $D(\mathrm{Gr}(k, \mathcal{A}))$.*

Proof. By Lemma 3.1, it suffices to show that $(S(\alpha))$ generates $D(\mathrm{Gr}(k, \mathcal{A}))$. Following [6] or [7], there exist sheaves \mathcal{F}_α of right $\mathcal{A}^{|\alpha|}$ -modules and sheaves \mathcal{G}_{α^*} of left $\mathcal{A}^{|\alpha|}$ -modules such that

$$j^*\mathcal{F}_\alpha \simeq S^\alpha\mathcal{R} \otimes q^*((\mathcal{E}^*)^{\otimes|\alpha|}), \quad j^*\mathcal{G}_{\alpha^*} \simeq q^*(\mathcal{E}^{\otimes|\alpha|}) \otimes S^{\alpha^*}\mathcal{T}^*.$$

Moreover, the sequence $\mathcal{R} \boxtimes \mathcal{T}^* \rightarrow \mathcal{O}_{\mathrm{Gr}(k, \mathcal{E}) \times \mathrm{Gr}(k, \mathcal{E})} \rightarrow \mathcal{O}_{\Delta(\mathrm{Gr}(k, \mathcal{E})/U)} \rightarrow 0$ descends to the sequence $\mathcal{F}_{(1)} \boxtimes \mathcal{G}_{(1)} \rightarrow \mathcal{O}_{\mathrm{Gr}(k, \mathcal{A}) \times \mathrm{Gr}(k, \mathcal{A})} \rightarrow \mathcal{O}_{\Delta(\mathrm{Gr}(k, \mathcal{A})/X)} \rightarrow 0$. Hence, we have the Koszul resolution:

$$\begin{aligned} 0 \rightarrow \Lambda^{k(n-k)}(\mathcal{F}_{(1)} \boxtimes \mathcal{G}_{(1)}) \rightarrow \Lambda^{k(n-k)-1}(\mathcal{F}_{(1)} \boxtimes \mathcal{G}_{(1)}) \rightarrow \dots \\ \dots \rightarrow \mathcal{F}_{(1)} \boxtimes \mathcal{G}_{(1)} \rightarrow \mathcal{O}_{\mathrm{Gr}(k, \mathcal{A}) \times_X \mathrm{Gr}(k, \mathcal{A})} \rightarrow \mathcal{O}_{\Delta(\mathrm{Gr}(k, \mathcal{A})/X)} \rightarrow 0. \end{aligned}$$

Therefore, as $\Lambda^m(\mathcal{F}_{(1)} \boxtimes \mathcal{G}_{(1)}) = \bigoplus_{|\alpha|=m} \mathcal{F}_\alpha \boxtimes \mathcal{G}_{\alpha^*}$ for $1 \leq m \leq k(n-k)$, the sheaf of ideals of the diagonal embedding $\mathcal{O}_{\Delta(\mathrm{Gr}(k, \mathcal{A})/X)}$ is in the subcategory

$$\langle \pi_1^*\mathcal{F}_\alpha \otimes \pi_2^*\mathcal{G}_{\alpha^*} \mid 0 \leq |\alpha| \leq k(n-k) \rangle$$

of $D(\mathrm{Gr}(k, \mathcal{A}) \times \mathrm{Gr}(k, \mathcal{A}))$. By the projection formula, for any $\mathcal{M} \in D(\mathrm{Gr}(k, \mathcal{A}))$ we have $\mathcal{M} = R(\pi_1)_*(\pi_2^*\mathcal{M} \otimes \mathcal{O}_{\Delta(\mathrm{Gr}(k, \mathcal{A})/X)})$.

To finish the proof, it is enough to verify that

$$R(\pi_1)_*(\pi_2^*\mathcal{M} \otimes (\pi_1^*\mathcal{F}_\alpha \otimes \pi_1^*\mathcal{G}_{\alpha^*})) \in S(\alpha) :$$

since we have $R(\pi_1)_*(\pi_2^*\mathcal{M} \otimes (\pi_1^*\mathcal{F}_\alpha \otimes \pi_1^*\mathcal{G}_{\alpha^*})) = R(\pi_1)_*(\pi_2^*(\mathcal{M} \otimes \mathcal{G}_{\alpha^*})) \otimes \mathcal{F}_\alpha$, this is isomorphic to

$$q^*(Rq_*(\mathcal{M} \otimes S^{\alpha^*}\mathcal{T}^*)) \otimes S^\alpha\mathcal{R} \text{ over } \mathrm{Gr}(k, \mathcal{E}). \quad \square$$

Let $1 \leq k_1 < \dots < k_m < n$ be a sequence of integers. Given a sheaf of Azumaya algebras \mathcal{A} of rank n^2 over X , we denote by $\mathrm{Fl}(k_1, \dots, k_m, \mathcal{A})$ the functor defined by $(Y \xrightarrow{\phi} X) \mapsto$ the set of sheaves of left ideals $\mathcal{I}_1 \subset \dots \subset \mathcal{I}_m$ of $\phi^*\mathcal{A}$ such that $\phi^*\mathcal{A}/\mathcal{I}_i$ is a locally free \mathcal{O}_Y -modules of rank $n(n - k_i)$.

As in the grassmannian case, we have an étale covering $i : U \rightarrow X$ and a locally free sheaf \mathcal{E} of rank n over U with the following pullback diagram

$$\begin{array}{ccc} \mathrm{Fl}(k_1, \dots, k_m, \mathcal{E}nd(\mathcal{E})) & \xrightarrow{j} & \mathrm{Fl}(k_1, \dots, k_m, \mathcal{A}) \\ \downarrow q & & \downarrow p \\ U & \xrightarrow{i} & X, \end{array}$$

where $\mathrm{Fl}(k_1, \dots, k_m, \mathcal{E}nd(\mathcal{E}))$ is identified with $\mathrm{Fl}(k_1, \dots, k_m, \mathcal{E})$. We have the tautological flags

$$\mathcal{R}_1 \hookrightarrow \dots \hookrightarrow \mathcal{R}_m \hookrightarrow q^* \mathcal{E} \twoheadrightarrow \mathcal{T}_1 \twoheadrightarrow \dots \twoheadrightarrow \mathcal{T}_m,$$

where $\mathrm{rank}(\mathcal{R}_i) = k_i$ and $\mathrm{rank}(\mathcal{T}_i) = n - k_i$.

Let $\alpha(1), \dots, \alpha(m-1), \alpha(m)$ be partitions of the forms

$$(\alpha_1, \dots, \alpha_{k_1}), \dots, (\alpha_1, \dots, \alpha_{k_{m-1}}), (\alpha_1, \dots, \alpha_{k_m})$$

with $0 \leq \alpha_i \leq k_2 - k_1, \dots, 0 \leq \alpha_i \leq k_m - k_{m-1}, 0 \leq \alpha_i \leq n - k_m$, respectively. For $1 \leq i \leq m$, we define $S(\alpha(1), \dots, \alpha(m))$ to be the full subcategory of $\mathrm{D}(\mathrm{Fl}(k_1, \dots, k_m, \mathcal{A}))$ generated by \mathcal{M} in $\mathrm{D}(\mathrm{Fl}(k_1, \dots, k_m, \mathcal{A}))$ satisfying

$$\mathcal{M}|_{\mathrm{Fl}(k_1, \dots, k_m, \mathcal{E})} \simeq q^* \mathcal{N} \otimes S^{\alpha(1)} \mathcal{R}_1 \otimes \dots \otimes S^{\alpha(m)} \mathcal{R}_m,$$

for some $\mathcal{N} \in \mathrm{D}(U)$.

Corollary 3.3. *Let $1 \leq k_1 < \dots < k_m < n$ be a sequence of integers and let $(S(\alpha(1), \dots, \alpha(m)) \mid \text{all partitions of the form } \alpha(i), 1 \leq i \leq m)$ be a sequence of the full subcategories of $\mathrm{D}(\mathrm{Fl}(k_1, \dots, k_m, \mathcal{A}))$ in lexicographical order. Then this gives a semiorthogonal decomposition of $\mathrm{D}(\mathrm{Fl}(k_1, \dots, k_m, \mathcal{A}))$.*

Proof. We shall prove by induction on m . The case $m = 1$ follows from Theorem 3.2. Assume that the result holds for $m - 1$. There are projections

$$\mathrm{Fl}(k_1, \dots, k_m, \mathcal{E}) \xrightarrow{q_m} \dots \xrightarrow{q_2} \mathrm{Fl}(k_m, \mathcal{E}) \xrightarrow{q_1} U$$

and

$$\mathrm{Fl}(k_1, \dots, k_m, \mathcal{A}) \xrightarrow{p_m} \dots \xrightarrow{p_2} \mathrm{Fl}(k_m, \mathcal{A}) \xrightarrow{p_1} X.$$

Let $\mathcal{R}'_2 \subset (q_1 \circ \dots \circ q_{m-1})^* \mathcal{E}$ be the tautological subsheaf over $\mathrm{Fl}(k_2, \dots, k_m, \mathcal{E})$ and let \mathcal{A}' be the sheaf of Azumaya algebra over $\mathrm{Fl}(k_2, \dots, k_m, \mathcal{A})$ from $\mathcal{E}nd(\mathcal{R}'_2)$ by descent. Then, we have $\mathrm{Fl}(k_1, \dots, k_m, \mathcal{E}) = \mathrm{Gr}_{\mathrm{Fl}(k_2, \dots, k_m, \mathcal{E})}(k_1, \mathcal{R}'_2)$ and $\mathrm{Fl}(k_1, \dots, k_m, \mathcal{A}) = \mathrm{Gr}_{\mathrm{Fl}(k_2, \dots, k_m, \mathcal{A})}(k_1, \mathcal{A}')$. Now the result follows from the proofs of Lemma 3.1 and Theorem 3.2. \square

REFERENCES

- [1] A. A. Beilinson, *Coherent sheaves on \mathbf{P}^n and problems in linear algebra*, Funktsional. Anal. i Prilozhen. **12** (1978), 68–69.
- [2] M. Bernardara, *A semiorthogonal decomposition for Brauer-Severi schemes*, Math. Nachr. **282** (2009), 1406–1413.
- [3] A. Bondal, *Representations of associative algebras and coherent sheaves*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), 25–44 (Russian); English transl. in Math. USSR Izv. **34** (1990), 23–42.

- [4] M. Kapranov, *On the derived category of coherent sheaves on some homogeneous spaces*, Invent. Math. **92** (1988), 479–508.
- [5] D. Orlov, *Projective bundles, monoidal transformations, and derived categories of coherent sheaves*, Izv. Akad. Nauk SSSR Ser. Mat. **56** (1992), 852–862 (Russian); English transl. in Math. USSR Izv. **38** (1993), 133–141.
- [6] M. Levine, V. Srinivas, and J. Weyman, *K-theory of twisted Grassmannians*, K-Theory **3** (1989), 99–121.
- [7] I. A. Panin, *Algebraic K-theory of Grassmannian manifolds and their twisted forms*, Funktsional. Anal. i Prilozhen. **23** (1989), 71–72 (Russian); English transl. in Funct. Anal. Appl. **23**, 143–144.

E-mail address: baek.sanghoon@gmail.com