Nonlinear Schrödinger equations near an infinite well potential

THOMAS BARTSCH MONA PARNET

Abstract

The paper deals with standing wave solutions of the dimensionless nonlinear Schrödinger equation

$$(NLS_{\lambda}) \qquad i\Phi_t(x,t) = -\Delta_x \Phi + V_{\lambda}(x)\Phi + f(x,\Phi), \quad x \in \mathbb{R}^N, \ t \in \mathbb{R},$$

where the potential $V_{\lambda} : \mathbb{R}^N \to \mathbb{R}$ is close to an infinite well potential $V_{\infty} : \mathbb{R}^N \to \mathbb{R}$, i. e. $V_{\infty} = \infty$ on an exterior domain $\mathbb{R}^N \setminus \Omega$, $V_{\infty}|_{\Omega} \in L^{\infty}(\Omega)$, and $V_{\lambda} \to V_{\infty}$ as $\lambda \to \infty$ in a sense to be made precise. The nonlinearity may be of Gross-Pitaevskii type. A solution of (NLS_{λ}) with $\lambda = \infty$ vanishes on $\mathbb{R}^N \setminus \Omega$ and satisfies Dirichlet boundary conditions, hence it solves

$$(NLS_{\infty}) \qquad \begin{cases} i\Phi_t(x,t) = -\Delta_x \Phi + V_{\lambda}(x)\Phi + f(x,\Phi), & x \in \Omega, \ t \in \mathbb{R} \\ \Phi(x,t) = 0 & x \in \partial\Omega, \ t \in \mathbb{R}. \end{cases}$$

We investigate when a solution Φ_{∞} of the infinite well potential (NLS_{∞}) gives rise to nearby solutions Φ_{λ} of the finite well potential (NLS_{λ}) with $\lambda \gg 1$ large. Considering (NLS_{∞}) as a singular limit of (NLS_{λ}) we prove a kind of singular continuation type results.

Keywords: nonlinear Schrödinger equations, infinite well potential, deep potential well, nonlinear eigenvalue problems, singular limit, variational methods, topological methods, singular continuation

AMS subject classification: 35J20, 35J61, 35J91, 35Q55, 58E05

1 Introduction

Infinite well potentials like the infinite square well or the infinite spherical well are helpful as instructive models to describe confined particles in quantum mechanical systems. They are often used as a starting point for solving finite well problems. In this paper we investigate nonlinear Schrödinger equations, like the Gross-Pitaevskii equation, with a potential $V_{\lambda} : \mathbb{R}^N \to \mathbb{R}$ close to an infinite well potential $V_{\infty} : \mathbb{R}^N \to \mathbb{R}$. More precisely, $V_{\infty} = \infty$ on an exterior domain $\mathbb{R}^N \setminus \Omega$, and $V_{\infty}|_{\Omega} \in L^{\infty}(\Omega)$. As $\lambda \to \infty$ the potential depth of V_{λ} becomes infinite, i. e. $V_{\lambda} \to V_{\infty}$, in a sense to be made precise below. Our goal is to give rigorous proofs for the passage from the infinite well potential to the finite well potential.

We are interested in standing waves $\Phi(t,x) = e^{i\omega t}u(x)$ of the finite well nonlinear Schrödinger equation

$$(NLS_{\lambda}) \qquad i\Phi_t(x,t) = -\Delta_x \Phi(x,t) + V_{\lambda}(x)\Phi + f(x,\Phi), \quad x \in \mathbb{R}^N, \ t \in \mathbb{R},$$

where $V_{\lambda}(x) \to V_{\infty}(x)$ as $\lambda \to \infty$. For $\lambda = \infty$ a solution should vanish in $\mathbb{R}^N \setminus \Omega$ and satisfy Dirichlet boundary conditions on Ω , hence it is a solution of the singular limit problem:

$$(NLS_{\infty}) \qquad \begin{cases} i\Phi_t(x,t) = -\Delta_x \Phi + V_{\lambda}(x)\Phi + f(x,\Phi), & x \in \Omega, \ t \in \mathbb{R}, \\ \Phi(x,t) = 0 & x \in \partial\Omega, \ t \in \mathbb{R}. \end{cases}$$

The question we address in this paper is: suppose we know a solution Φ_{∞} of (NLS_{∞}) , does there exist a nearby solution Φ_{λ} of (NLS_{λ}) for λ large?

Standing wave solutions of (NLS_{λ}) correspond to solutions of the stationary nonlinear Schrödinger equation

$$(S_{\lambda}) \qquad \begin{cases} -\Delta u + V_{\lambda}(x)u = f(x, u) & \text{for } x \in \mathbb{R}^{N}; \\ u(x) \to 0 & \text{as } |x| \to \infty, \end{cases}$$

where we incorporated the term ωu generated by the ansatz into the potential without changing notation. For $\lambda = \infty$ we are similarly led to consider

$$(S_{\infty}) \qquad \qquad -\Delta u + V_{\infty}(x)u = f(x, u), \quad u \in H_0^1(\Omega),$$

as a singular limit of (S_{λ}) as $\lambda \to \infty$. The original question can now be reformulated as which solutions u_{∞} of (S_{∞}) appear as limits of solutions u_{λ} of (S_{λ}) . Solutions of (S_{∞}) can be obtained via variational or topological methods. We provide conditions on the convergence of $V_{\lambda} \to V_{\infty}$ and on f such that an isolated solution u_{∞} of (S_{∞}) which can be found by variational or topological methods gives rise to a family of solutions u_{λ} of (S_{λ}) . We include of course the generic case where u_{∞} is a nondegenerate solution of (S_{∞}) .

For the proofs we develop an abstract functional analytic approach in order to deal with the above type of singular limit problem. Our results may be thought of as being continuation results near a singular limit: For $\lambda < \infty$ we look for solutions of an equation $F_{\lambda}(u) = 0$ defined on $H^1(\mathbb{R}^N)$, whereas the limit equation $F_{\infty}(u) = 0$ is only defined for $u \in H^1_0(\Omega)$. Some of the methods we develop can also be applied to more general nonlinear eigenvalue problems that are not necessarily of variational type.

The paper is organized as follows. In Section 2 we state our main results about (S_{λ}) , and we discuss related results. Then in Section 3 we formulate the functional analytic setting which will be considered throughout the paper. Here we also state our main abstract results about solutions of nonlinear equations near a singular parameter limit. The abstract results as well as the results about (S_{λ}) will be proved in sections 4 - 6.

2 NLS near an infinite well potential

We begin with collecting our assumptions on the potentials V_{λ} . These are given in the form $V_{\lambda} = a_0 + \lambda a$, so the problem we consider is

$$(S_{\lambda}) \qquad \begin{cases} -\Delta u + (a_0(x) + \lambda a(x))u = f(x, u) & \text{for } x \in \mathbb{R}^N; \\ u(x) \to 0 & \text{as } |x| \to \infty, \end{cases}$$

and the limit problem is

$$(S_{\infty}) \qquad \qquad -\Delta u + a_0(x)u = f(x, u), \quad u \in H_0^1(\Omega).$$

The distinguishing feature is that the potential $a \in L^{\infty}_{loc}(\mathbb{R}^N)$ satisfies $a \geq 0$ and $a^{-1}(0) = \overline{\Omega}$ with $\Omega \subset \mathbb{R}^N$ nonempty, open, and bounded. Consequently, $V_{\lambda}(x) \to \infty$ as $\lambda \to \infty$ for $x \notin \overline{\Omega}$.

In order to describe the assumptions on a and a_0 we need some notation. For $x \in \mathbb{R}^N$ and r > 0 we set $B_r(x) := \{y \in \mathbb{R}^N : |y - x| < r\}$. We also set $K_r^c := \{x \in \mathbb{R}^n : |x|_{\infty} > r\}$. Let $\mu_N(-\Delta + V_{\lambda}, G)$ be the infimum of the spectrum of $-\Delta + V_{\lambda}$ on an open subset $G \subset \mathbb{R}^N$ with Neumann boundary conditions, i. e.

$$\mu_N(-\Delta+V_\lambda,G) = \inf_{\psi \in H^1(G) \setminus \{0\}} \frac{\int_G (|\nabla \psi|^2 + V_\lambda \psi^2) dx}{\|\psi\|_{L^2(G)}^2}.$$

Our basic hypotheses on the potential are:

- $(V_1) \ a_0 \in L^{\infty}_{\text{loc}}(\mathbb{R}^N)$ and ess inf $a_0 > -\infty$.
- $(V_2) \ a \in L^{\infty}_{loc}(\mathbb{R}^N), a(x) \ge 0 \text{ and } \Omega := \text{int } a^{-1}(0) \text{ is a non-empty open subset of } \mathbb{R}^N$ with Lipschitz boundary.

 (V_3) There exists a sequence $R_j \to \infty$ such that

$$\lim_{\lambda \to \infty} \liminf_{j \to \infty} \mu_N(-\Delta + V_\lambda, K_{R_j}^c) = \infty.$$

The reader can find a discussion of condition (V_3) , in particular various equivalent conditions, in [8]. Condition (V_3) holds, for instance, if a satisfies:

 (V_4) There exist M > 0 and r > 0 such that

$$\operatorname{meas}(\{x \in B_r(y) : a(x) < M\}) \to 0 \quad \text{as } |y| \to \infty$$

where meas denotes the Lebesgue measure.

 (V_3) implies that the embedding $H_0^1(\Omega) \hookrightarrow L^p(\Omega)$ is compact for $2 \le p < \infty$. Observe that (V_3) and (V_4) allow that Ω may be unbounded. For some results we require the stronger condition

 (V_5) The form domain

$$E := \left\{ u \in H^1(\mathbb{R}^N) : \int_{\mathbb{R}^N} a_0 u^2 < \infty, \ \int_{\mathbb{R}^N} a u^2 < \infty \right\}$$

embeds compactly into $L^2(\mathbb{R}^N)$.

This holds, for instance, if $a_0(x) \to \infty$ or $a(x) \to \infty$ as $|x| \to \infty$, a condition usually satisfied for confining potentials. (V_5) also holds under the weaker condition

 (V_6) For any M > 0 and any r > 0 there holds:

$$\max\{x \in B_r(y) : a(x \le M\} \to 0 \text{ as } |y| \to \infty$$

A proof that (V_6) implies (V_5) can be found in [18]; see also [22].

Concerning the nonlinearity f we only require that

(f₁) f is a Carathéodory function, and there exists constants C > 0, $2 < q < p < 2^*$ such that

$$|f(x,t)| \le C(|t|^{p-1} + |t|^{q-1})$$
 for $t \in \mathbb{R}$, a. e. $x \in \mathbb{R}^N$.

This includes the model nonlinearity $f(x, u) = W(x) \cdot |u|^{p-2}u$ with 2 $and <math>W \in L^{\infty}(\mathbb{R}^N)$, which appears in the Gross-Pitaevskii equation. We define $E_{\infty} := H_0^1(\Omega)$ provided with the scalar product

$$\langle u, v \rangle := \int_{\Omega} (\nabla u \nabla v + (b + a_0) u v) \, dx$$

where $b := 1 - \text{ess inf } a_0$. As a consequence of (V_1) and (V_2) this induces a norm which is equivalent to the standard norm of $H_0^1(\Omega)$. Setting $F(x, u) := \int_0^u f(x, t) dt$, it is well known that the functional $J_\infty : E_\infty \to \mathbb{R}$ defined by

$$J_{\infty}(u) = \frac{1}{2} \int_{\Omega} (|\nabla u|^2 + a_0 u^2) \, dx - \int_{\Omega} F(x, u) \, dx$$
$$= \frac{1}{2} ||u||^2 - \int_{\Omega} \left(\frac{b}{2}u^2 + F(x, u)\right) \, dx$$

is of class C^1 , and that critical points of J_{∞} are solutions of (S_{∞}) .

Recall that the critical groups of an isolated critical point u of a functional $J : E \to \mathbb{R}$ are defined as $C_k(J, u) := H_k(J^c, J^c \setminus \{u\})$ where c := J(u). Here H_* is singular homology with coefficients in a commutative ring R with unit; typically $R = \mathbb{Z}$ or R is a field.

Now we can state our first result.

Theorem 2.1. Assume $(V_1) - (V_3)$ and (f_1) hold. Let $u_{\infty} \in E_{\infty}$ be an isolated solution of (S_{∞}) with nontrivial critical groups $C_*(J_{\infty}, u_{\infty})$. Then there exists $\Lambda \ge 1$ such that for each $\lambda \ge \Lambda$ there exists a solution $u_{\lambda} \in E$ of (S_{λ}) with $u_{\lambda} \to u_{\infty}$ in E as $\lambda \to \infty$.

Remark 2.2. If 0 does not belong to the spectrum of $-\Delta + a_0$ in $H_0^1(\Omega)$ then this holds true for $-\Delta + a_0 + \lambda a$ for λ large. Then the solutions which we obtain in Theorem 2.1 and in the theorems below decay exponentially; see [24].

If $C_*(J_{\infty}, u_{\infty}) = 0$ then the solution u_{∞} cannot be discovered using variational methods, and it can disappear under small perturbations. In our next result we strengthen the hypotheses by assuming that u_{∞} has nontrivial index. Consider the functional

$$K_{\infty}: E_{\infty} \to \mathbb{R}, \quad K_{\infty}(u) = \int_{\Omega} \left(\frac{b}{2}u^2 + F(x, u)\right) dx,$$

and define its gradient $k_{\infty} = \nabla K_{\infty} : E_{\infty} \to E_{\infty}$ with respect to the above scalar product on E_{∞} . Then k_{∞} is completely continuous because $p < 2^*$ in (f_1) . The index of u_{∞} is then defined by

$$\operatorname{ind}(k_{\infty}, u_{\infty}) := \operatorname{deg}(\operatorname{id} - k_{\infty}, B_{\delta}(u_{\infty}, E_{\infty}), 0).$$

Here deg denotes the Leray-Schauder degree, $\delta > 0$ is small so that u_{∞} is the only solution of (S_{∞}) in the δ -ball $B_{\delta}(u_{\infty}, E_{\infty})$ of u_{∞} in E_{∞} .

Theorem 2.3. Assume $(V_1), (V_2), (V_5)$ and (f_1) hold. Let $u_{\infty} \in E_{\infty}$ be an isolated solution of (S_{∞}) with nontrivial index $ind(k_{\infty}, u_{\infty})$. Then there exists a connected set

$$\mathcal{S} \subset \{(\lambda, u) \in \mathbb{R} \times E : u \text{ solves } (S_{\lambda})\} \subset \mathbb{R} \times E$$

such that S covers a parameter interval $[\Lambda, \infty)$ for some $\Lambda \ge 1$. Moreover, $u_n \to u_\infty$ for any sequence $(\lambda_n, u_n) \in E$ with $\lambda_n \to \infty$.

The assumption $\operatorname{ind}(k_{\infty}, u_{\infty}) \neq 0$ in Theorem 2.3 is stronger than the assumption $C_*(J_{\infty}, u_{\infty}) \neq 0$ in Theorem 2.1 because of the Poincaré-Hopf formula:

(2.1)
$$\operatorname{ind}(k_{\infty}, u_{\infty}) = \sum_{i=0}^{\infty} (-1)^{i} \operatorname{rank} C_{i}(J_{\infty}, u_{\infty}).$$

Surprisingly, the strong assumption (V_5) can be replaced by (V_3) if f satisfies

 (f'_1) f is differentiable in t, f and f_t are Carathéodory functions and there exist constants $c > 0, 2 < q < p < 2^* = 2N/(N-2)^+$ such that

$$|f_t(x,t)| \le c(|t|^{p-2} + |t|^{q-2})$$
 for $t \in \mathbb{R}$, a. e. $x \in \mathbb{R}^N$;

With this condition the functional J_{λ} is of class C^2 .

Theorem 2.4. Assume $(V_1) - (V_3)$ and (f'_1) hold. Let $u_{\infty} \in E_{\infty}$ be an isolated solution of (S_{∞}) with nontrivial index ind (k_{∞}, u_{∞}) . Then the conclusion of Theorem 2.3 holds.

For our last result about (S_{λ}) we consider the case of a nondegenerate solution u_{∞} .

Theorem 2.5. Assume $(V_1) - (V_3)$ and (f'_1) hold. Let $u_{\infty} \in E_{\infty}$ be a nondegenerate solution of (S_{∞}) . Then there exists $\Lambda \ge 1$ and a C^1 -function

 $[\Lambda, \infty) \to E, \quad u \mapsto u_{\lambda},$

such that u_{λ} is a solution of (S_{λ}) , and $u_{\lambda} \to u_{\infty}$ as $\lambda \to \infty$.

Problem (S_{λ}) has found much interest in recent years after being first considered in [10, 8]. Most papers deal with potentials being positive and bounded away from 0, i. e. inf $a_0 > 0$, exceptions being [9, 14]. The equation (S_{λ}) with asymptotically linear nonlinearity has been studied in [20, 21, 28, 29], with critical growth nonlinearity in [3, 4], with Neumann boundary conditions in exterior domains in [11]. In [9, 15, 27] multiplicity results have been obtained provided the bottom $\overline{\Omega}$ of the potential well consists of several connected components. Extensions to quasilinear problems can be found in [2], to the Schrödinger-Poisson system in [17].

In almost all earlier papers on the topic the authors made assumptions on a, a_0, f such that variational methods (e. g. the mountain pass theorem or some linking theorem) can be applied to show that (S_{λ}) has a solution u_{λ} . Then it is proved that u_{λ} converges as $\lambda \to \infty$ towards a solution u_{∞} of the limit problem (S_{∞}) . However, the limit u_{∞} has not been prescribed in these papers as we do here. A notable exception, and the only one we are aware of, where the limit has been prescribed is [27, Theorem 1.2]. There the authors considered the one-dimensional problem

(2.2)
$$-u'' + (1 + \lambda a(x))u = |u|^{p-1}u, \quad u \in H^1(\mathbb{R}),$$

with the limit problem

(2.3)
$$\begin{cases} -u'' + u = |u|^{p-1}u, \quad x \in \Omega = (a_1, b_1) \cup (a_2, b_2), \\ u(a_i) = u(b_i) = 0. \end{cases}$$

The solutions of (2.3) can be listed as $v_{i,j}$, $i, j \in \mathbb{Z}$, where $v_{\pm i,\pm j}$ are the unique solutions having |i| zeroes in (a_1, b_1) and |j| zeroes in (a_2, b_2) . The authors find solutions u_{λ} of (2.2) such that $u_{\lambda} \to v_{i,j}$ as $\lambda \to \infty$. The proof is based on ODE methods and cannot be extended to dimensions $N \ge 2$. It depends on the uniqueness of the solutions having a certain nodal structure. Observe that in the one-dimensional case the solutions $v_{i,j}$ are automatically non-degenerate, hence our Theorem 2.5 applies. Thus we improve and generalize [27, Theorem 1.2] considerably. Moreover, our proof is simpler and extends to the PDE setting.

In contrast to all earlier papers we do not require global linking type hypotheses. Our results may be considered as a local version of these earlier results. As a consequence, we can deal with solutions of (S_{∞}) which are obtained not using a global linking structure. This implies in particular to almost critical problems like

$$-\Delta u = |u|^{2^* - 2 - \varepsilon} u, \quad u \in H^1_0(\Omega),$$

where in the limit for $\varepsilon \to 0$ the problem can be reduced via the Lyapunov-Schmidt reduction method to finding critical points of a finite-dimensional limit function; see [5, 6, 7, 23, 26]. For instance, in [6] the solutions have been obtained by finding a local minimum and a local mountain pass of the reduced functional.

3 Critical points near a singular limit

Let E be a real Hilbert space with scalar product $\langle \cdot, \cdot \rangle$, and let $A : E \to E$ be a bounded self-adjoint linear operator. We require that $A \ge 0$ and that $E_{\infty} := \ker A \ne \{0\}$. Finally, let $K : E \to \mathbb{R}$ be a C^1 -function, and set $k := \nabla K : E \to E$. We are interested in finding critical points of the functional

$$J_{\lambda}: E \to \mathbb{R}, \quad J_{\lambda}(u) := \frac{1}{2} ||u||^2 + \frac{\lambda}{2} \langle Au, u \rangle - K(u)$$

for λ large. Observe that $J_{\lambda}(u)$ is independent of λ for $u \in E_{\infty}$. Moreover, for $u \in E \setminus E_{\infty}$ we have $J_{\lambda}(u) \to \infty$ as $\lambda \to \infty$. We set $K_{\infty} = K | E_{\infty}, k_{\infty} := \nabla K_{\infty} : E_{\infty} \to E_{\infty}$, and consider

$$J_{\infty}: E_{\infty} \to \mathbb{R}, \quad J_{\infty}(u) = \frac{1}{2} \|u\|^2 - K_{\infty}(u),$$

as singular limit functional. Clearly, J_{∞} is just the restriction of J_{λ} to E_{∞} .

Observe that $\langle Au, u \rangle > 0$ for $u \in E \setminus E_{\infty}$ and that

(3.1)
$$u_n \rightharpoonup u \text{ in } E, \langle Au_n, u_n \rangle \to 0 \implies Au_n \to 0, \ u \in E_{\infty}.$$

This can be seen by looking at the symmetric positive-semidefinite bilinear form $(u, v)_A := \langle Au, v \rangle$. The Schwarz inequality yields

$$||Au||^2 = (u, Au)_A \le \sqrt{(u, u)_A} \sqrt{(Au, Au)_A}.$$

Therefore $\langle Au, u \rangle = (u, u)_A = 0$ implies Au = 0. Similarly, $u_n \rightharpoonup u$, $\langle Au_n, u_n \rangle \rightarrow 0$ implies $Au_n \rightarrow 0 = Au$.

For $\lambda \geq 0$ and $u, v \in E$ we define

$$\langle u, v \rangle_{\lambda} := \langle u, v \rangle + \lambda \langle Au, v \rangle.$$

As a consequence of our hypotheses on A this is a scalar product on E, and it defines a norm $\|\cdot\|_{\lambda}$ on E which is equivalent to the given norm corresponding to $\lambda = 0$. Observe that the orthogonal complement of E_{∞} with respect to $\langle \cdot, \cdot \rangle_{\lambda}$,

$$E_{\infty}^{\perp} = \{ u \in E : \langle u, v \rangle_{\lambda} = 0 \text{ for all } v \in E_{\infty} \}$$

is independent of λ , hence the orthogonal projections $P : E \to E_{\infty}$ and $Q = \mathrm{id} - P : E \to E_{\infty}^{\perp}$ are independent of λ . We write $B_{r,\lambda}(0, E_{\infty}^{\perp}) := \{v \in E_{\infty}^{\perp} : \|v\|_{\lambda} \leq r\}$ for the ball of radius r > 0 around $0 \in E_{\infty}^{\perp}$ with respect to the norm $\|\cdot\|_{\lambda}$. For $\delta > 0$ and $\lambda > 0$ and $u \in E_{\infty}$ we define

$$B_{\delta,\lambda}(u) := B_{\delta}(u, E_{\infty}) \times B_{\delta,\lambda}(0, E_{\infty}^{\perp}) \subset E.$$

Given a bounded linear map $L: E \to E$ we write

$$||L||_{\lambda} := \sup\{||Lu||_{\lambda} : u \in E, \ ||u||_{\lambda} \le 1\}$$

for the operator norm of L with respect to $\|\cdot\|_{\lambda}$ on E.

For $\lambda > 0$ we define the nonlinear operators $k_{\lambda} = \nabla_{\lambda} K : E \to E$ and $\nabla_{\lambda} J_{\lambda} : E \to E$ by the equations

$$\langle k_{\lambda}(u), v \rangle_{\lambda} = \langle \nabla_{\lambda} K(u), v \rangle_{\lambda} = DK(u)[v] = \langle k(u), v \rangle,$$

and $\nabla_{\lambda} J_{\lambda} = id - k_{\lambda}$. Observe that

(3.2)
$$||k_{\lambda}(u)||_{\lambda} = \sup_{\|v\|_{\lambda} \le 1} \langle k_{\lambda}(u), v \rangle_{\lambda} = \sup_{\|v\|_{\lambda} \le 1} \langle k(u), v \rangle \le \sup_{\|v\| \le 1} \langle k(u), v \rangle = \|k(u)\|.$$

If K is of class C^2 near u then the derivatives of $k = \nabla K$ and of $k_{\lambda} = \nabla_{\lambda} K$ satisfy

(3.3)
$$\langle Dk_{\lambda}(u)[v], w \rangle_{\lambda} = \langle Dk(u)[v], w \rangle \text{ for } u, v, w \in E.$$

We also deduce for $\lambda \ge 0$ and $u \in E$ that

$$\begin{aligned} \|Dk_{\lambda}(u)\|_{\lambda}^{2} &= \sup_{\|v\|_{\lambda} \leq 1} \langle Dk_{\lambda}(u)[v], Dk_{\lambda}(u)[v] \rangle_{\lambda} = \sup_{\|v\|_{\lambda} \leq 1} \langle Dk(u)[v], Dk_{\lambda}(u)[v] \rangle \\ &\leq \sup_{\|v\|_{\lambda} \leq 1} \|Dk(u)\| \cdot \|v\| \cdot \|Dk_{\lambda}(u)\|_{\lambda} \cdot \|v\|_{\lambda} \leq \|Dk(u)\| \cdot \|Dk_{\lambda}(u)\|_{\lambda} \end{aligned}$$

hence,

$$||Dk_{\lambda}(u)||_{\lambda} \le ||Dk(u)||_{\lambda}$$

Similarly we obtain for $\lambda \ge 0$ and $u, v \in E$ that

(3.5)
$$||Dk_{\lambda}(u) - Dk_{\lambda}(v)||_{\lambda} \le ||Dk(u) - Dk(v)||.$$

Now we collect some hypotheses on J_{λ} which we will impose in the various results.

 (J_1) J_{∞} has an isolated critical point $u_{\infty} \in E_{\infty}$, and the critical groups of u_{∞} as a critical point of J_{∞} are nontrivial: $C_*(J_{\infty}, u_{\infty}) \neq 0$.

We fix $\delta_0 > 0$ such that u_{∞} is the only critical point of J_{∞} in $B_{\delta_0}(u_{\infty})$.

- (J₂) There exists $\lambda_0 > 0$ such that k is weakly sequentially continuous in $B_{\delta_0,\lambda_0}(u_\infty)$, i. e. if $u_n \in B_{\delta_0,\lambda_0}(u_\infty)$ and $u_n \rightharpoonup u$ then $k(u_n) \rightharpoonup k(u)$.
- $(J_3) \ B_{\delta_0,\lambda_n}(u_\infty) \ni u_n \rightharpoonup u, \ \lambda_n \to \infty \qquad \Longrightarrow \qquad k(u_n) \to k(u).$
- (J_4) There exists $\lambda_0 > 0$ such that k is completely continuous in $B_{\delta_0,\lambda_0}(u_\infty)$.

Condition (J_2) is rather harmless, also (J_3) holds under rather general assumptions on a, a_0 , and f. Both are much weaker than requiring that k is completely continuous near u_{∞} as in (J_4) . (J_2) does imply that k_{∞} is completely continuous in $B_{\delta_0}(u_{\infty}, E_{\infty})$. Therefore J_{∞} satisfies the Palais-Smale condition in $B_{\delta_0}(u_{\infty}, E_{\infty})$, i. e. any Palais-Smale sequence $u_n \in B_{\delta_0}(u_{\infty}, E_{\infty})$ for J_{∞} has a convergent subsequence.

Theorem 3.1. Suppose that (J_1) , (J_2) , and (J_3) hold. Then there exists $\Lambda \ge 0$ such that J_{λ} has a critical point u_{λ} for $\lambda \in [\Lambda, \infty)$ and such that $u_{\lambda} \to u_{\infty}$ as $\lambda \to \infty$.

Our next result is based on degree theory. Recall that the index of u_{∞} as fixed point of k_{∞} is defined as:

$$\operatorname{ind}(k_{\infty}, u_{\infty}) = \operatorname{deg}(\operatorname{id} - k_{\infty}, B_{\delta}(u_{\infty}, E_{\infty}), u_{\infty})$$

where deg denotes the Leray-Schauder degree, $0 < \delta \leq \delta_0$. This index is defined, for instance, if k_{∞} is completely continuous in $B_{\delta_0}(u_{\infty}, E_{\infty})$, hence if (J_2) or (J_3) holds, in particular if (J_4) holds.

Theorem 3.2. Suppose (J_1) and (J_4) hold. Suppose moreover that the local fixed point index of u_{∞} as a fixed point of k_{∞} is nontrivial:

$$\operatorname{ind}(k_{\infty}, u_{\infty}) = \operatorname{deg}(\operatorname{id} - k_{\infty}, U_{\varepsilon}(u_{\infty}, E_{\infty}), u_{\infty}) \neq 0.$$

Then there exists a connected set $S \subset [\Lambda, \infty) \times E$ covering the parameter interval $[\Lambda, \infty)$ for some $\Lambda \geq 1$, such that $\nabla J_{\lambda}(u) = 0$ for every $(\lambda, u) \in S$. Moreover, given a sequence $(\lambda_n, u_n) \in S$ with $\lambda_n \to \infty$ there holds $u_n \to u_\infty$.

Remark 3.3. a) Recall that under the conditions of Theorem 3.1 the local index ind (K_{∞}, u_{∞}) may be trivial. On the other hand, if the local index ind (K_{∞}, u_{∞}) is nontrivial, then the critical groups $C_*(f, u_{\infty})$ are nontrivial. This follows from the Poincaré-Hopf formula (2.1).

b) Assumption (J_4) can be replaced by any assumption assuring that there is a degree theory for the maps $id - k_{\lambda}$. In the case of (J_4) one has the Leray-Schauder degree. If k_{λ} is, for instance, A-proper in the sense of [25], the generalized degree of Petryshin can be applied; see Theorem 3.4 below and its proof in Section 5.

Surprisingly, the compactness condition can be considerable relaxed if K is C^2 near u_{∞} . We need the following condition on the differential $Dk(u_{\infty})$.

(J₅) If $u_n \rightarrow u$ and $||u_n||_{\lambda_n}$ is bounded for some sequence $\lambda_n \rightarrow \infty$, then $Dk(u_{\infty})[u_n] \rightarrow Dk(u_{\infty})[u]$.

Theorem 3.4. Suppose K is C^2 near u_{∞} , (J_1) , (J_3) , and (J_5) are satisfied. Then the conclusion of Theorem 3.2 holds true.

Under the assumptions of Theorem 3.4 k_{λ} need not be compact, so we cannot work with the Leray-Schauder degree. Instead we will be able to use the degree for β -condensing maps where where β is the ball measure of noncompactness; see [13].

Finally we state a result in the nondegenerate setting.

Theorem 3.5. Suppose that K is C^2 near u_{∞} , that u_{∞} is a nondegenerate critical point of J_{∞} , and that (J_5) is satisfied. Then there exists $\Lambda \ge 0$ and a C^1 -map $[\Lambda, \infty) \to E$, $\lambda \mapsto u_{\lambda}$, such that u_{λ} is the unique critical point of J_{λ} near u_{∞} for $\lambda \in [\Lambda, \infty)$. Moreover, $u_{\lambda} \to u_{\infty}$ as $\lambda \to \infty$.

4 Nontrivial critical groups

We first prove Theorem 3.1. Consider the isolated critical point $u_{\infty} \in E_{\infty}$ of J_{∞} with nontrivial critical groups. Let (W, W_{-}) be a Gromoll-Meyer pair for u_{∞} in $B_{\delta_0}(u_{\infty}, E_{\infty})$. This means that:

- (GM_1) $W \subset \operatorname{int} B_{\delta_0}(u_{\infty}, E_{\infty})$ is a closed neighborhood of u_{∞} in E_{∞} containing no other critical point of J_{∞} .
- (GM_2) There exist C^1 -functions $g_i : U \cap E_{\infty} \to \mathbb{R}$, $i = 1, \ldots, l$, having 0 as regular value, such that $W = \bigcap_{i=1}^l g_i^0$ and $\partial W = W \cap \bigcup_{i=1}^l g_i^{-1}(0)$.
- (GM_3) ∇J_{∞} is transversal to each $g_i^{-1}(0)$; more precisely, for some $\alpha > 0$:

$$\langle \nabla J_{\infty}(u), \nabla g_i(u) \rangle \leq -3\alpha < 0 \quad \text{for } u \in \partial W \cap g_i^{-1}(0), \ i = 1, \dots, j,$$

and

$$\langle \nabla J_{\infty}(u), \nabla g_i(u) \rangle \ge 3\alpha > 0 \quad \text{for } u \in \partial W \cap g_i^{-1}(0), \ i = j+1, \dots, l.$$

 (GM_4) The exit set

$$W_{-} = \left\{ u \in \partial W : \langle \nabla J_{\infty}(u), \nabla g_{i}(u) \rangle < 0 \text{ if } u \in g_{i}^{-1}(0), i = 1, ..., l \right\}$$
$$= \bigcup_{i=1}^{j} g_{i}^{-1}(0)$$

consists of those $x \in \partial W$ where $-\nabla J_{\infty}$ points outside of W.

A construction of a Gromoll-Meyer pair can be found in [12, p. 49], where l = 3, j = 1. Using a pseudo-gradient vector field for J_{∞} it is standard to show that

(4.1)
$$C_*(J_{\infty}, u_{\infty}) = H_*(J^c, J^c \setminus \{u_{\infty}\}) \cong H_*(W, W_-).$$

This uses that J_{∞} satisfies the Palais-Smale condition in $B_{\delta_0}(u_{\infty}, E_{\infty})$.

For $\delta > 0$ we set

$$W_{\delta,\lambda} := W \times B_{\delta,\lambda}(0, E_{\infty}^{\perp}).$$

Lemma 4.1. If (J_3) holds then for every $\varepsilon > 0$ there exists $\Lambda > 0$ such that

$$\sup_{u \in W_{\delta_0,\lambda}} \|k_{\lambda}(u) - k_{\infty}(Pu)\|_{\lambda} \le \varepsilon \quad \text{for all } \lambda \ge \Lambda$$

Proof. Arguing by contradiction, suppose there exist $\varepsilon > 0$, $\lambda_n \to \infty$, $u_n \in W_{\delta_0,\lambda_n}$ with

$$||k_{\lambda_n}(u_n) - k_{\infty}(Pu_n)||_{\lambda_n} \ge \varepsilon$$

Then the sequence $(u_n)_n$ is bounded, and $\langle Au_n, u_n \rangle \to 0$, hence (3.1) applies and yields u_n , $Pu_n \to u \in E_{\infty}$ along a subsequence. Now (J_3) implies $k(u_n) \to k(u)$ and $k(Pu_n) \to k(u)$. Setting $v_n := k_{\lambda_n}(u_n) - k_{\infty}(Pu_n)$ we see that $||v_n||_{\lambda_n}$ is bounded uniformly in n as a consequence of (3.2). Applying (3.1) again shows that $v_n, Pv_n \to v \in E_{\infty}$ along a subsequence. This in turn implies

$$\varepsilon^{2} \leq \|v_{n}\|_{\lambda_{n}}^{2} = \langle k_{\lambda_{n}}(u_{n}), v_{n} \rangle_{\lambda_{n}} - \langle k_{\infty}(Pu_{n}), v_{n} \rangle$$
$$= \langle k(u_{n}), v_{n} \rangle - \langle k(Pu_{n}), Pv_{n} \rangle \rightarrow \langle k(u), v \rangle - \langle k(u), v \rangle = 0$$

which is absurd.

Lemma 4.2. For all $0 < \delta \leq \delta_0$ there exists $\Lambda_{\delta} > 0$ such that for $\lambda \geq \Lambda_{\delta}$ and $v \in B_{\delta,\lambda}(0, E_{\infty}^{\perp})$, there holds:

$$\langle P\nabla_{\lambda}J_{\lambda}(u+v), \nabla g_{i}(u) \rangle \leq -2\alpha \quad \text{for } u \in \partial W \cap g_{i}^{-1}(0), \ i = 1, \dots, j,$$

and

$$\langle P \nabla_{\lambda} J_{\lambda}(u+v), \nabla g_i(u) \rangle \ge 2\alpha \quad \text{for } u \in \partial W \cap g_i^{-1}(0), \ i = j+1, \dots, l,$$

and

$$\langle Q \nabla_{\lambda} J_{\lambda}(u+v), v \rangle_{\lambda} \geq \delta^2/2 \quad \text{for } u \in W, \ v \in B_{\delta,\lambda}(0, E_{\infty}^{\perp}).$$

Proof. We may assume that

$$m := \max_{i=1,...,l} \sup_{u \in g_i^{-1}(0) \cap W} \|\nabla g_i(u)\| < \infty.$$

According to Lemma 4.1, for $0 < \delta \leq \delta_0$ there exists $\Lambda_{\delta} > 0$ such that

$$||k_{\lambda}(u) - k_{\infty}(Pu)||_{\lambda} \le \min\{\alpha/m, \delta/2\} \quad \text{if } \lambda \ge \Lambda_{\delta}, \ u \in W_{\delta_{0}, \lambda}.$$

Consequently, we obtain for $\lambda \ge \Lambda_{\delta}$, i = 1, ..., j, $u \in W \cap g_i^{-1}(0)$ and $v \in B_{\delta,\lambda}(0, E_{\infty}^{\perp})$ that

$$\langle P\nabla_{\lambda}J_{\lambda}(u+v), \nabla g_{i}(u) \rangle$$

$$\leq \langle \nabla J_{\infty}(u), \nabla g_{i}(u) \rangle + \|P\nabla_{\lambda}J_{\lambda}(u+v) - \nabla J_{\infty}(u)\| \cdot \|\nabla g_{i}(u)\|$$

$$= \langle \nabla J_{\infty}(u), \nabla g_{i}(u) \rangle + \|Pk_{\lambda}(u+v) - k_{\infty}(u)\| \cdot \|\nabla g_{i}(u)\|$$

$$\leq -3\alpha + \frac{\alpha}{m}m = -2\alpha.$$

Similarly we obtain for $\lambda \geq \Lambda_{\delta}$, i = j + 1, ..., l, $u \in W \cap g_i^{-1}(0)$ and $v \in B_{\delta,\lambda}(0, E_{\infty}^{\perp})$ that

$$\langle P\nabla_{\lambda}J_{\lambda}(u+v), \nabla g_i(u)\rangle \geq 2\alpha.$$

Finally, for $\lambda \geq \Lambda_{\delta}$, $u \in W$ and $v \in B_{\delta,\lambda}(0, E_{\infty}^{\perp})$ there holds:

$$\begin{split} \langle \nabla_{\lambda} Q J_{\lambda}(u+v), v \rangle_{\lambda} &= \|v\|_{\lambda}^{2} - \langle k_{\lambda}(u+v) - k_{\infty}(u), v \rangle_{\lambda} \geq \|v\|_{\lambda}^{2} - \|\langle k_{\lambda}(u+v) - k_{\infty}(u)\|_{\lambda} \|v\|_{\lambda} \\ &\geq \delta^{2} - \frac{\delta}{2}\delta = \frac{\delta^{2}}{2}. \end{split}$$

 \Box

Lemma 4.2 implies that for $\delta > 0$ and $\lambda \ge \Lambda_{\delta}$, the set

$$\left(W_{\delta,\lambda}, W_{-} \times B_{\delta,\lambda}(0, E_{\infty}^{\perp})\right) = \left(W \times B_{\delta,\lambda}(0, E_{\infty}^{\perp}), W_{-} \times B_{\delta,\lambda}(0, E_{\infty}^{\perp})\right)$$

is a regular index pair for pseudo-gradient flows of J_{λ} in the sense of Conley index theory.

Lemma 4.3. J_{λ} has a critical point $u_{\lambda} \in W_{\delta,\lambda}$ if $0 < \delta \leq \delta_0$ and $\lambda \geq \Lambda_{\delta}$.

Proof. If J_{λ} does not have a critical point in $W_{\delta,\lambda}$ then there exists a pseudo-gradient vector field V for J_{λ} in $W_{\delta,\lambda}$ such that the inequalities in Lemma 4.2 hold with V instead of $\nabla_{\lambda} J_{\lambda}$, α instead of 2α , and $\delta^2/4$ instead of $\delta^2/2$. Moreover,

(4.2)
$$\inf_{u \in W_{\delta,\lambda}} \|\nabla_{\lambda} J_{\lambda}(u)\|_{\lambda} > 0,$$

because if $u_n \in W_{\delta,\lambda}$ satisfies $\nabla_{\lambda} J_{\lambda}(u_n) = u_n - k_{\lambda}(u_n) \to 0$, then $u_n \rightharpoonup u \in W_{\delta,\lambda}$ along a subsequence, hence $k_{\lambda}(u_n) \rightharpoonup k_{\lambda}(u)$ as a consequence of (J_2) . This implies that $u \in W_{\delta,\lambda}$ is a critical point of J_{λ} . Observe that we do not prove strong convergence here, hence we do not prove the Palais-Smale condition in $W_{\delta,\lambda}$.

Now (4.2) implies that the flow associated to -V provides a deformation of $W_{\delta,\lambda}$ to $W_- \times B_{\delta,\lambda}(0, E_{\infty}^{\perp})$. This in turn implies

$$H_*\left(W_{\delta,\lambda}, W_- \times B_{\delta,\lambda}(0, E_\infty^{\perp})\right) \cong 0$$

in contradiction with

$$H_*\left(W_{\delta,\lambda}, W_- \times B_{\delta,\lambda}(0, E_\infty^{\perp})\right) \cong H_*(W, W_-) \cong C_*(J_\infty, u_\infty) \neq 0.$$

Proof of 3.1. The existence of a critical point $u_{\lambda} \in W \times B_{\delta,\lambda}(0, E_{\infty}^{\perp})$ of J_{λ} for $\lambda \geq \Lambda$ has been stated in Lemma 4.3. Clearly, $Qu_{\lambda} \to 0$ and $\nabla J_{\infty}(Pu_{\lambda}) \to 0$ as $\lambda \to \infty$. It follows that $u_{\lambda} \to u_{\infty} \in E_{\infty}$ because u_{∞} is the only critical point of J_{∞} in W.

Proof of 2.1. In order to apply Theorem 3.1 we set

$$E := \left\{ u \in H^1(\mathbb{R}^N) : \int_{\mathbb{R}^N} a_0 u^2 < \infty, \ \int_{\mathbb{R}^N} a u^2 < \infty \right\}$$

provided with the scalar product

$$\langle u, v \rangle := \int_{\mathbb{R}^N} (\nabla u, \nabla v + (b + a_0 + a)uv) dx$$

Here $b = 1 - ess \inf a_0$ is defined as in Section 2. The operator $A : E \to E$ is defined by the equation

$$\langle Au, v \rangle := \int_{\mathbb{R}^N} auv dx,$$

and the functional $K:E\to \mathbb{R}$ by

$$K(u) = -\int_{\mathbb{R}^N} \left(\frac{b}{2}u^2 + F(x,u)\right) \, dx.$$

A is a self-adjoint, positive semidefinite, and bounded linear operator. The kernel E_{∞} of A consists of all $u \in E$ such that u = 0 a. e. in $\mathbb{R}^N \setminus \Omega$, hence $E_{\infty} = H_0^1(\Omega)$. This uses that the boundary of Ω is Lipschitz.

Solutions of (S_{λ}) are obtained as critical points of the \mathcal{C}^1 -functional

$$J_{\lambda}(u) = \frac{1}{2} \int_{\mathbb{R}^{N}} (|\nabla u|^{2} + (a_{0} + \lambda a)u^{2} dx - \int_{\mathbb{R}^{N}} F(x, u) dx$$

= $\frac{1}{2} (||u||^{2} + (\lambda - 1)\langle Au, u \rangle) - K(u).$

Observe that λ has to be replaced by $\lambda - 1$ because $\|\cdot\|_{\lambda}$ contains the summand $\langle Au, u \rangle$. Since a = 0 on Ω , we see that J_{∞} is simply the restriction of J_{λ} to E_{∞} . It remains to prove the conditions (J_2) and (J_3) . In fact, (J_2) is an easy consequence of (f_1) because E imbeds into $L^p(\mathbb{R}^N)$ for $2 \le p \le 2^*$. In order to see (J_3) , consider sequences $\lambda_n \to \infty$ and $B_{\delta_0,\lambda_n} \ni u_n \rightharpoonup u$. Then $u_n \to u$ strongly in $L^p(\mathbb{R}^N)$ for $2 by [8, Lemma 4.2.]. And <math>u_n \to u$ strongly in $L^2(\mathbb{R}^N)$ follows from (V_3) . This implies $k(u_n) \to k(u)$ in E because of the subcritical growth of f required in (f_1) .

Theorem 2.1 is now an immediate consequence of Theorem 3.1.

5 Nontrivial index

Proof of Theorem 3.2. From $||u - k_{\infty}(Pu)|| \ge ||Pu - k_{\infty}(Pu)||$ and using Lemma 4.1 we immediately deduce that there exists $\Lambda > 0$ such that for $\lambda \ge \Lambda$ and $0 < \delta \le \delta_0$ small that

(5.1)

$$0 \neq \operatorname{ind}(k_{\infty}, u_{\infty}) = \operatorname{deg}(\operatorname{id}_{E_{\infty}} - k_{\infty}, B_{\delta}(u_{\infty}, E_{\infty}), u_{\infty})$$

$$= \operatorname{deg}(\operatorname{id}_{E} - k_{\infty} \circ P, B_{\delta,\lambda}(u_{\infty}), u_{\infty})$$

$$= \operatorname{deg}(\operatorname{id}_{E} - k_{\lambda}, B_{\delta,\lambda}(u_{\infty}), u_{\infty}).$$

Since k is completely continuous in $B_{\delta_0,\lambda_0}(u_\infty)$ so is k_λ , hence the above degree is defined Using (5.1), a standard continuation argument (see [1], for instance) shows that there exists a connected set $S \subset [\Lambda, \infty) \times B_{\delta,\lambda}(u_\infty) \subset [\Lambda, \infty) \times E$ covering the parameter inverval $[\Lambda, \infty)$, such that $\nabla J_\lambda(u) = u - k_\lambda(u) = 0$ for every $(\lambda, u) \in S$. Given a sequence $(\lambda_n, u_n) \in S$ with $\lambda_n \to \infty$, using (5.1) and Lemma 4.1 once more, we deduce that $||u_n - u_\infty||_\lambda \to 0$.

Proof of Theorem 2.3. This follows from Theorem 3.2 as Theorem 2.1 follows from Theorem 3.1. We only need to observe that k is completely continuous as a consequence of (V_5) and (f_1) , in particular (J_4) is satisfied.

For the proof of Theorem 3.4 we need the following lemma.

Lemma 5.1. Suppose (J_5) is satisfied. Then

$$||Dk_{\lambda}(u_{\infty}) - Dk_{\infty}(u_{\infty}) \circ P||_{\lambda} \to 0 \text{ as } \lambda \to \infty.$$

Proof. Arguing by contradiction, suppose that there exist sequences $\lambda_n \to \infty$, $u_n \in E$ with $||u_n||_{\lambda_n} = 1$, and

$$||Dk_{\lambda_n}(u_{\infty})[u_n] - Dk_{\infty}(u_{\infty})[Pu_n]||_{\lambda} \ge \varepsilon > 0.$$

Then $u_n \rightharpoonup u$ in E along a subsequence, and $u \in E_{\infty}$ by (3.1), hence also $Pu_n \rightharpoonup u$. Setting

$$v_n := Dk_{\lambda_n}(u_\infty)[u_n] - Dk_\infty(u_\infty)[Pu_n]$$

and using (3.4) we see that

$$\|v_n\|_{\lambda_n} \le \|Dk_{\lambda_n}(u_{\infty})[u_n]\|_{\lambda_n} + \|Dk(u_{\infty})[Pu_n]\| \le \|Dk(u_{\infty})\| + \|Dk(u_{\infty})\|$$

is bounded uniformly in n. We deduce, again by (3.1), that $v_n \rightarrow v$ in E along a subsequence, and that $v \in E_{\infty}$, hence also $Pv_n \rightarrow v$. Using condition (J_5) we obtain a contradiction:

$$\varepsilon^{2} \leq ||v_{n}||_{\lambda_{n}}^{2} = \langle Dk_{\lambda_{n}}(u_{\infty})[u_{n}], v_{n} \rangle_{\lambda_{n}} - \langle Dk_{\infty}(u_{\infty})[Pu_{n}], Pv_{n} \rangle$$

= $\langle Dk(u_{\infty})[u_{n}], v_{n} \rangle - \langle Dk(u_{\infty})[Pu_{n}], Pv_{n} \rangle$
 $\rightarrow \langle Dk(u_{\infty})[u], v \rangle - \langle Dk(u_{\infty})[u], v \rangle = 0.$

Proof of Theorem 3.4.. Let β_{λ} be the ball measure of non-compactness in E, i. e. for a subset $A \subset E$

 $\beta_{\lambda}(A) = \inf\{r > 0 : A \text{ can be covered by finitely many } \| \cdot \|_{\lambda} \text{-balls of radius } r\}.$

We claim that k_{λ} is a strict β_{λ} -set contraction in a neighborhood of u_{∞} if λ is large. We refer to [13] for properties of this class of maps and the construction of a degree theory. It is sufficient to show that

(5.2)
$$\beta_{\lambda}(k_{\lambda}(A)) \leq \frac{1}{2}\beta_{\lambda}(A)$$
 for $A \subset B_{\delta,\lambda}(u_{\infty})$ if λ is large and δ is small.

For (5.2) it suffices to prove that $k_{\lambda} - k_{\infty} \circ P$ is $\|\cdot\|_{\lambda}$ -Lipschitz continuous with Lipschitz constant $\frac{1}{2}$ because $k_{\infty} \circ P$ is completely continuous as a consequence of (J_3) , and because the sum of a completely continuous map and a Lipschitz map with Lipschitz constant $\frac{1}{2}$ satisfies (5.2). Now the Lipschitz continuity of $k_{\lambda} - k_{\infty} \circ P$ follows easily from:

$$\begin{aligned} \|k_{\lambda}(u) - k_{\infty}(Pu) - (k_{\lambda}(v) - k_{\infty}(Pv))\|_{\lambda} \\ &\leq \|k_{\lambda}(u) - k_{\lambda}(v) - Dk_{\lambda}(u_{\infty})[u - v])\|_{\lambda} \\ &+ \|Dk_{\lambda}(u_{\infty})[u - v] - Dk_{\infty}(u_{\infty})[Pu - Pv]\|_{\lambda} \\ &+ \|k_{\infty}(Pu) - k_{\infty}(Pv) - Dk_{\infty}(u_{\infty})[Pu - Pv]\|_{\lambda} \\ &\leq \sup_{w \in B_{\delta,\lambda}(u_{\infty})} \|Dk_{\lambda}(w)[u - v] - Dk_{\lambda}(u_{\infty})[u - v]\|_{\lambda} \\ &+ \|Dk_{\lambda}(u_{\infty}) - Dk_{\infty}(u_{\infty}) \circ P\|_{\lambda}\|u - v\|_{\lambda} \\ &+ \sup_{w \in B_{\delta}(u_{\infty}, E_{\infty})} \|Dk_{\infty}(w)[u - v] - Dk_{\infty}(u_{\infty})[u - v]\| \end{aligned}$$

Now $\sup_{w \in B_{\delta,\lambda}(u_{\infty})} \|Dk_{\lambda}(w) - Dk_{\lambda}(u_{\infty})\|_{\lambda}$ and $\sup_{w \in B_{\delta}(u_{\infty}, E_{\infty})} \|Dk_{\infty}(w) - Dk_{\infty}(u_{\infty})\|_{\lambda}$ can be made arbitrarily small by making $\delta > 0$ small. And $\|Dk_{\lambda}(u_{\infty}) - Dk_{\infty}(u_{\infty})\|_{\lambda} \circ P$

can be made arbitrarily small as $\lambda \to \infty$ as a consequence of Lemma 5.1.

Since k_{λ} is a strict β_{λ} -set contraction in a neighborhood of u_{∞} for λ large, we may argue as in the proof of Theorem 3.2 to conclude the proof of Theorem 3.4.

Proof of Theorem 2.4. Observe that (f'_1) implies that

$$K: E \to \mathbb{R}, \quad K(u) = \int_{\mathbb{R}^N} \left(\frac{b}{2}u^2 + F(x, u)\right) dx,$$

is of class C^2 . It remains to prove (J_3) and (J_5) . In fact, the proof of (J_3) proceeds as in the proof of Theorem 2.1. In order to see (J_5) consider a sequence $u_n \in E$ such that $u_n \rightharpoonup u$ and $||u_n||_{\lambda_n}$ is bounded for some sequence $\lambda_n \rightarrow \infty$, so that $u \in E_{\infty} = H_0^1(\Omega)$. Now assumption (V_3) yields a sequence $R_j \rightarrow \infty$ such that

$$\lim_{n \to \infty} \liminf_{j \to \infty} \frac{\|u_n - u\|_{\lambda_n}^2}{\int_{K_{R_i}^c} |u_n - u|^2} \to \infty,$$

which implies

$$\int_{K_{R_j}^c} |u_n - u|^2 \to 0 \quad \text{as } j, n \to \infty.$$

Since $u_n \to u$ in $L^2_{loc}(\mathbb{R}^N)$ we deduce that $u_n \to u$ in $L^2(\mathbb{R}^N)$. This implies that

$$\begin{aligned} |\langle Dk(u_{\infty})[u_n] - Dk(u_{\infty})[u], v\rangle| &= \left| \int_{\mathbb{R}^N} (b + f'(u_{\infty}))(u_n - u)v \, dx \right| \\ &\leq c ||u_n - u||_{L^2(\Omega)} ||v|| \end{aligned}$$

hence $Dk(u_{\infty})[u_n] \to Dk(u_{\infty})[u]$ in E.

Now Theorem 2.4 follows from Theorem 3.4.

6 The nondegenerate case

In this section we use the notation $f_{\lambda} = id_E - k_{\lambda} : E \to E$. The proof of Theorem 3.5 is an immediate consequence of the following proposition.

Proposition 6.1. For $\delta > 0$ small there exists $\Lambda_{\delta} \ge 1$ such that the map

$$g_{\lambda}: B_{\delta,\lambda}(u_{\infty}) \to B_{\delta,\lambda}(u_{\infty}), \quad g_{\lambda}(u):=u - (\mathrm{id}_E - Dk_{\infty}(u_{\infty}) \circ P)^{-1} \circ f_{\lambda}(u),$$

is well defined and a contraction for $\lambda \geq \Lambda_{\delta}$.

Proof of Theorem 3.5. According to Proposition 6.1 there exists $\delta_0 > 0$ such that for $0 < \delta \leq \delta_0$ and $\lambda \geq \Lambda_{\delta}$, the Banach fixed point theorem yields a unique fixed point $u_{\lambda} \in B_{\delta,\lambda}(u_{\infty})$ of g_{λ} , hence a zero of f_{λ} , i. e. a critical point of J_{λ} . The map

$$[\Lambda_{\delta_0}, \infty) \to E, \quad \lambda \mapsto u_\lambda,$$

is C^1 because f_{λ} is C^1 in λ . Finally, $||u_{\lambda} - u_{\infty}||_{\lambda} \to 0$ is also a consequence of Proposition 6.1.

The proof of Proposition 6.1 is based on the following lemmata.

Lemma 6.2. The bounded operator $L := id_E - Dk_{\infty}(u_{\infty}) \circ P : E \to E$ is an isomorphism, and $||L^{-1}||_{\lambda} \leq \alpha$ is bounded uniformly in λ .

Proof. That L is an isomorphism follows easily from the assumption that u_{∞} is a nondegenerate fixed point of k_{∞} , which means that $\mathrm{id}_{E_{\infty}} - Dk_{\infty}(u_{\infty}) : E_{\infty} \to E_{\infty}$ is an isomorphism. It is also clear that $||L^{-1}||_{\lambda} \leq \max\{1, ||u_{\infty} - Dk_{\infty}(u_{\infty})^{-1}||\}$ because the norms on E_{∞} do not depend on λ .

Lemma 6.3. $||f_{\lambda}(u_{\infty})||_{\lambda} \to 0 \text{ as } \lambda \to \infty.$

Proof. Arguing by contradiction, suppose there exist $\varepsilon > 0$ and $\lambda_n \to \infty$ such that $v_n := f_{\lambda_n}(u_\infty) = u_\infty - k_{\lambda_n}(u_\infty) = k_\infty(u_\infty) - k_{\lambda_n}(u_\infty)$ satisfies $||v_n||_{\lambda_n} \ge \varepsilon$. Observe that $||v_n||_{\lambda_n}$ is bounded uniformly in n as a consequence of (3.2). Now (3.1) implies $v_n, Pv_n \to v \in E_\infty$ along a subsequence. This in turn implies:

$$\varepsilon^2 \le \|v_n\|_{\lambda_n}^2 = \langle k_\infty(u_\infty), v_n \rangle - \langle k_{\lambda_n}(u_\infty), v_n \rangle_{\lambda_n} = \langle k(u_\infty), Pv_n \rangle - \langle k(u_\infty), v_n \rangle \to 0$$

which is absurd.

Proof of Proposition 6.1. By (3.5) there exists $\delta_1 > 0$ such that

(6.1)
$$\sup_{u \in B_{\delta_1,\lambda}(u_{\infty})} \|Dk_{\lambda}(u) - Dk_{\lambda}(u_{\infty})\|_{\lambda} \le \frac{1}{4\alpha} \quad \text{for all } \lambda \ge 0,$$

where $\alpha > 0$ is from Lemma 6.2. Now we fix $0 < \delta \le \delta_1$. Using Lemma 5.1 and Lemma 6.3 there exists Λ_{δ} such that

(6.2)
$$\|Dk_{\lambda}(u_{\infty}) - Dk_{\infty}(u_{\infty}) \circ P\|_{\lambda} \leq \frac{1}{4\alpha} \quad \text{for } \lambda \geq \Lambda_{\delta},$$

and

(6.3)
$$||f_{\lambda}(u_{\infty})||_{\lambda} \leq \frac{\delta}{2\alpha} \quad \text{for } \lambda \geq \Lambda_{\delta}.$$

Thus for $\lambda \geq \Lambda_{\delta}$ and $u, v \in B_{\delta,\lambda}(u_{\infty})$ there holds

(6.4)
$$\|k_{\lambda}(u) - k_{\lambda}(v) - Dk_{\lambda}(u_{\infty})(u-v)\|_{\lambda}$$
$$\leq \sup_{w \in B_{\delta,\lambda}(u_{\infty})} \|Dk_{\lambda}(w) - Dk_{\lambda}(u_{\infty})\|_{\lambda} \cdot \|u-v\|_{\lambda} \stackrel{\text{(6.1)}}{\leq} \frac{1}{4\alpha} \|u-v\|_{\lambda}.$$

Since $L = id_E - Dk_{\infty}(u_{\infty}) \circ P$ we have $g_{\lambda} = L^{-1}(k_{\lambda} - Dk_{\infty}(u_{\infty}) \circ P)$. It follows that

$$\begin{aligned} \|g_{\lambda}(u) - g_{\lambda}(v)\|_{\lambda} &\leq \alpha \|k_{\lambda}(u) - k_{\lambda}(v) - Dk_{\infty}(u_{\infty})(P(u-v))\|_{\lambda} \\ &\leq \alpha \|k_{\lambda}(u) - k_{\lambda}(v) - Dk_{\lambda}(u_{\infty})(u-v)\|_{\lambda} \\ &\quad + \alpha \|Dk_{\lambda}(u_{\infty})(u-v) - Dk_{\infty}(u_{\infty})(P(u-v))\|_{\lambda} \\ &\stackrel{(6.4)}{\leq} \alpha \frac{1}{4\alpha} \|u-v\|_{\lambda} + \alpha \|Dk_{\lambda}(u_{\infty}) - Dk_{\infty}(u_{\infty}) \circ P\|_{\lambda} \cdot \|u-v\|_{\lambda} \\ &\stackrel{(6.2)}{\leq} \frac{1}{4} \|u-v\|_{\lambda} + \frac{1}{4} \|u-v\|_{\lambda} = \frac{1}{2} \|u-v\|_{\lambda}. \end{aligned}$$

We also have

$$\|g_{\lambda}(u_{\infty}) - u_{\infty}\|_{\lambda} \le \|L^{-1}\|_{\lambda} \cdot \|f_{\lambda}(u_{\infty})\|_{\lambda} \stackrel{(6.3)}{\le} \alpha \frac{\delta}{2\alpha} = \frac{\delta}{2}$$

hence, for $u \in B_{\delta,\lambda}(u_{\infty})$ there holds:

$$\begin{aligned} \|g_{\lambda}(u) - u_{\infty}\|_{\lambda} &\leq \|g_{\lambda}(u) - g_{\lambda}(u_{\infty})\|_{\lambda} + \|g_{\lambda}(u_{\infty}) - u_{\infty}\|_{\lambda} \\ &\leq \frac{1}{2}\|u - u_{\infty}\|_{\lambda} + \frac{\delta}{2} \leq \frac{\delta}{2} + \frac{\delta}{2} = \delta. \end{aligned}$$

Therefore g_{λ} maps $B_{\delta,\lambda}(u_{\infty})$ into itself.

Proof of Theorem 2.5. As in the proof of Theorem 2.4 one sees that J_{λ} is of class C^2 and that (J_5) holds. Therefore Theorem 2.5 follows from Theorem 3.5.

References

- J. Alexander : A primer on connectivity. In: Proc. Conf. Fixed Point Theory (Sherbrooke, Que., 1980), 455âĂŞ483, Lecture Notes in Math. 886, Springer, Berlin-New York 1981.
- [2] C. O. Alves: *Existence of multi-bump solutions for a class of quasilinear problems*. Adv. Nonlin. Studies **6** (2006), 491-509.
- [3] C. O. Alves, D. C. de Morais Filho and M. A. S. Souto: *Multiplicity of positive solutions for a class of problems with critical growth in* ℝ^N. Proc. Royal Soc. Edinburgh 52 (2009), 1-21.

- [4] C. O. Alves and M. A. S. Souto: Multiplicity of positive solutions for a class of problems with exponential critical growth in ℝ². J. Diff. Equ. 244 (2008), 1502-1520.
- [5] T. Bartsch, T. d'Aprile and A. Pistoia: *Multi-bubble nodal solutions for slightly subcritical elliptic problems in domains with symmetries.* Preprint
- [6] T. Bartsch, T. d'Aprile and A. Pistoia: *On the profile of sign changing solutions of an almost critical problem in the ball.* Preprint
- [7] T. Bartsch, A. Micheletti and A. Pistoia: On the existence and the profile of nodal solutions of elliptic equations involving critical groth. Calc. Var. Part. Diff. Equ. 26 (2006), 265-282.
- [8] T. Bartsch, A. Pankov and Z.-Q.Wang: *Nonlinear Schrödinger equations with steep potential well*. Commun. Contemp. Math. **3** (2001), 549-569
- [9] T. Bartsch and Z. Tang: *Multibump solutions of nonlinear Schrödinger equations* with steep potential well and indefinite potential. Discr. Cont. Dynam. Syst. (to appear)
- [10] T. Bartsch and Z.-Q. Wang: *Existence and multiplicity results for some superlinear elliptic problems on* \mathbb{R}^N . Comm. Part. Diff. Eq. **20** (1995), 1725-1741.
- [11] J. Chabrowski and Z.-Q. Wang: *Exterior nonlinear Neumann problem*. Nonlin. Diff. Equ. Appl. 13 (2007), 683-697.
- [12] K.C. Chang: Infinite Dimensional Morse Theory and Multiple Solution Problems. Birkhäuser, Boston 1993.
- [13] K. Deimling: Nonlinear Functional Analysis. Springer-Verlag, Berlin Heidelberg 1985.
- [14] Y. Ding and A. Szulkin: Bound states for semilinear Schrödinger equations with sign-changing potential. Calc. Var. Part. Diff. Equ. 29 (2007), 397-419.
- [15] Y. Ding and K. Tanaka: *Multiplicity of positive solutions of a nonlinear Schrödinger equation*. Man. Math. **112** (2003), 109-135.
- [16] M. Furtado, E. A. B. Silva and M. S. Xavier: Multiplicity and concentration of solutions for elliptic systems with vanishing potentials. J. Diff. Equ. 249 (2010), 2377-2396.
- [17] Y. Jiang and H.-S. Zhou: Schrödinger-Poisson system with steep potential well. J. Diff. Equ. 251 (2011), 582-608.

- [18] V. Kondrat'ev and M. Shubin: Discreteness of spectrum for the Schrödinger operators on manifolds of bounded geometry. Operator theory: Advances and Applications 110 (1999), 185–226
- [19] Z. Liu, F. A. van Heerden and Z.-Q. Wang: Nodal type bound states of Schrödinger equations via invariant set and minimax methods. J. Diff. Equ. 214 (2005), 358-390.
- [20] X. Liu, Y. Huang: Sign-changing solutions for a class of nonlinear Schrödinger equations. Bull. Australian Math. Soc. **80** (2009), 294-305.
- [21] X. Liu, Y. Huang and J. Liu: Sign-changing solutions for an asymptotically linear Schrödinger equation with deepening potential well. Adv. Diff. Equ. 16 (2011), 1-30.
- [22] Molchanov, A.M.: On the discreteness of the spectrum conditions for self-adjoint differential equations of the second order. Trudy Mosk. Matem. Obshchestva 2 (1953), 169–199 (in Russian). Adv. Diff. Equ. 16 (2011), 1-30.
- [23] M. Musso and A. Pistoia: Tower of bubbles for almost critical problems in general domains. J. Math. Pure Appl. 93 (2010), 1-40.
- [24] A. Pankov: On decay of solutions to nonlinear Schrödinger equations. Proc. Amer. Math. Soc. 136 (2008), 2565-2570.
- [25] W. V. Petryshin: Generalized Topological Degree and Semilinear Equations. Cambridge University Press 1995.
- [26] A. Pistoia and T. Weth: Sign-changing bubble tower solutions in a slightly subcritical semilinear Dirichlet problem. Ann. Inst. H. Poincaré Anal. Non Lin. 24 (2007), 325-340.
- [27] Y. Sato and K. Tanaka: Sign-changing multi-bump solutions for nonlinear Schrödinger equations with steep potential wells. Trans. Amer. Math. Soc. 361 (2009), 6205-6253.
- [28] C. A. Stuart and H.-S. Zhou: Global branch of solutions for non-linear Schrödinger equations with deepening potential well. Proc. London Math. Soc. 92 (2006), 655-681.
- [29] Z. Wang and H.-S. Zhou: Positive solutions for nonlinear Schrödinger equations with deepening potential well. J. Europ. Math. Soc. 11 (2009), 545-573.
- [30] W. Zou: Sign-changing saddle point. J. Funct. Anal. 219 (2005), 433-468.

Address of the authors:

Thomas Bartsch, Mona Parnet Mathematisches Institut University of Giessen Arndtstr. 2 35392 Giessen Germany Thomas.Bartsch@math.uni-giessen.de Mona.Parnet@math.uni-giessen.de