

ÉTALE SUBQUOTIENTS OF PRIME TORSION OF ABELIAN SCHEMES

HENDRIK VERHOEK

ABSTRACT. Let A be an abelian variety over a number field K with good reduction outside a finite set of primes S . We show that if the ℓ -torsion subgroup schemes $A[\ell^n]$ lie in a certain category of group schemes, then $A[\ell^n]$ does not contain any subgroup schemes that are étale or are of multiplicative type.

CONTENTS

1. Introduction	1
2. The generic fiber of simple group schemes	2
3. Filtrations by simple group schemes	3
4. Application to abelian varieties	4
References	7

1. INTRODUCTION

Let K be a number field with ring of integers O_K and let S be a finite set of primes in O_K . Denote by O_S the ring of S -integers of K . Let ℓ be a rational prime such that none of the primes in S divides ℓ .

Definition 1.1. Let \mathcal{C} be a subcategory of the category of finite flat commutative group schemes over O_S of ℓ -power order, such that \mathcal{C} is closed under taking products, subquotients and Cartier duality.

In addition, with an eye towards our main theorem stated below, we state the following two conditions that the category \mathcal{C} might or might not satisfy. These conditions involve simple group schemes in \mathcal{C} , i.e., group schemes that have no non-trivial closed flat subgroup schemes.

Condition (1): For all simple non-étale group schemes T in \mathcal{C} and all simple étale group schemes E in \mathcal{C} , the group $\mathrm{Ext}_{\mathcal{C}}^1(T, E)$ is trivial.

Condition (2): Let F be the compositum of all $K(E)$, where E runs over all simple étale group schemes E in \mathcal{C} . Then the extension F/K is finite and the maximal abelian extension R of F , that is unramified outside S and at most tamely ramified at primes over S , is a cyclic extension.

Let A be an abelian variety over K with good reduction outside S , let \mathcal{A} denote its Néron model. Denote by $\mathcal{A}[\ell^n]$ the ℓ^n -torsion subgroup scheme of \mathcal{A} . The schemes $\mathcal{A}[\ell^n]$ are finite flat commutative group scheme over O_S . We prove:

Theorem 1.2. *Let A be an abelian variety such that $\mathcal{A}[\ell^n]$ is an object in \mathcal{C} for all $n \in \mathbf{N}$. If Conditions (1) and (2) hold for the category \mathcal{C} , then $\mathcal{A}[\ell]$ does not have subquotients that are étale or of multiplicative type.*

As an application, we prove:

Corollary 1.3. *There do not exist abelian varieties over $\mathbf{Q}(\sqrt{13})$ and $\mathbf{Q}(\sqrt{17})$ with good reduction everywhere.*

In the rest of the article we continue as follows. First we indicate how one finds simple group schemes in \mathcal{C} . Then we discuss filtrations and extensions of group schemes in \mathcal{C} and prove Theorem 1.2. The proof is divided into three steps, the same steps that can be found in [Fon85], [Sch03] and [Sch05] and that prove the non-existence or unique up to isogeny results of abelian varieties with good or semi-stable reduction at the primes in S .

- (1) Define a category \mathcal{C} that contains $\mathcal{A}[\ell^n]$ for all n
- (2) Find the simple objects in the category \mathcal{C} by using the generic fiber of objects in \mathcal{C} annihilated by ℓ and the discriminant bounds of Odlyzko to classify the generic fibers of simple objects in \mathcal{C} , and subsequently use theorems of Oort-Tate and Raynaud to determine the simple objects up to isomorphism. Verify that Condition (2) holds.
- (3) Calculate various extension groups of the objects in \mathcal{C} and verify Condition (1). If both conditions hold, then apply Theorem 1.2.

2. THE GENERIC FIBER OF SIMPLE GROUP SCHEMES

The generic fiber of a finite flat commutative group scheme J over O_S is a group scheme over K , which we denote by J_K . Since $\text{char}(K) = 0$, J_K is an étale group scheme. Therefore, the group scheme J_K is just an abelian group $J(\overline{K})$ together with the Galois action $\rho_J : G_K \rightarrow \text{Aut}(J(\overline{K}))$. We denote by $K(J)$ the field extension obtained by adjoining the \overline{K} -points of J to K . The representation ρ_J factors through a finite Galois extension $K(J)/K$. By considering the generic fiber J_K we obtain not only information about the group scheme J considered over K , but also as a scheme over O_S . It is even true that, under certain conditions (see [Ray74]), the generic fiber uniquely determines the group scheme J over O_S .

A first step to understand the category \mathcal{C} is to classify its simple objects up to isomorphism. Every simple object is annihilated by ℓ : if not, the Zariski closure of the ℓ -torsion points in the generic fiber would form a non-trivial closed flat subgroup scheme. Since by assumption \mathcal{C} is closed under taking subquotients, this subgroup scheme would again be in \mathcal{C} .

Define $T_{\mathcal{C}}$ to be the compositum of all fields $K(J)$, where the J are group schemes in \mathcal{C} that are annihilated by ℓ . We call $T_{\mathcal{C}}$ the *maximal ℓ -torsion extension* of \mathcal{C} . This extension $T_{\mathcal{C}}$ need not be finite in general. The reason that we are interested in the maximal ℓ -torsion extension of \mathcal{C} is that if $T_{\mathcal{C}}$ is finite, it enables us to find the simple objects in \mathcal{C} . Namely, the \overline{K} -points of every simple object generate an extension that is a subfield of the maximal ℓ -torsion extension of \mathcal{C} . As a side note we mention that to find $T_{\mathcal{C}}$ in practice, it is helpful that the category \mathcal{C} is closed under taking products.

Lemma 2.1. *If J is a simple finite flat commutative group scheme over O_S , then the representation $\rho_J : G_K \rightarrow \text{Aut}(J(\overline{K}))$ is irreducible.*

Proof. Suppose ρ_J admits a non-trivial G_K -stable subgroup V . Since the closure of the generic point of O_S is O_S (recall that O_S is a Dedekind ring), taking the Zariski closure of V gives a

non-trivial closed flat subgroup scheme of J . This closure is equal to J because J is simple. The generic fiber of the closure, which is equal to $J(\overline{K})$, is contained in V . \square

The generic fiber of a simple object J in \mathcal{C} is a simple $\mathbf{F}_\ell[\mathrm{Gal}(K(J)/K)]$ -module. Since simple objects are killed by ℓ , such a generic fiber is also a simple $\mathbf{F}_\ell[\mathrm{Gal}(T_\mathcal{C}/K)]$ -module. Therefore we classify all simple $\mathbf{F}_\ell[\mathrm{Gal}(T_\mathcal{C}/K)]$ -modules. If we can find a relatively large normal ℓ -subgroup H in $\mathrm{Gal}(T_\mathcal{C}/K)$, it is easier to classify irreducible submodules: the representation ρ_J factors not only through $\mathrm{Gal}(L/K)$, but also through the quotient of $\mathrm{Gal}(L/K)$ by H . This is an immediate consequence of:

Lemma 2.2. *Let J be a simple object in \mathcal{C} . Then $\mathrm{Gal}(K(J)/K)$ contains no non-trivial normal ℓ -subgroup.*

Proof. The representation ρ_J factors through $\mathrm{Gal}(K(J)/K)$. Let H be a non-trivial normal ℓ -subgroup of $\mathrm{Gal}(K(\overline{J})/K)$. Then H must act faithfully as a ℓ -group on the ℓ -group $J(\overline{K})$, but this is impossible. There are non-trivial fixed points of $J(\overline{K})$ under this action and they form a closed flat subgroup scheme of J , which must equal J since J is simple. \square

Finally, once simple $\mathbf{F}_\ell[\mathrm{Gal}(T_\mathcal{C}/K)]$ -modules have been found, the question remains if they extend to finite flat commutative group schemes over O_S . This is addressed in the work of Raynaud [Ray74] and Oort-Tate [TO70].

3. FILTRATIONS BY SIMPLE GROUP SCHEMES

In this section we discuss filtrations of group schemes in \mathcal{C} by simple subgroup schemes. These filtrations will be used to prove Theorem 1.2. Each finite flat commutative group scheme J contains a simple closed flat subgroup scheme J' . The same is true for J/J' . Continuing like this we obtain a filtration of J :

Definition 3.1. A (left) filtration of a finite flat commutative group scheme J is an ordered set $\{J_i\}_{i=1}^n$ such that

- J_1 is a simple closed flat subgroup scheme of $F_1 := J$
- for $1 < i < n$, let J_i be a simple closed flat subgroup scheme of $F_i := F_{i-1}/J_{i-1}$
- J_n is simple

We call n the length of the filtration.

We note that by using Cartier duality, we can get another (right) filtration. If A is a simple group scheme occurring in a filtration (or equivalently all filtrations) of J , we say that J admits A .

Lemma 3.2. *Let J be a group scheme in \mathcal{C} that admits the simple group scheme A . Suppose that for each simple B with $B \not\cong A$ occurring in the filtration of J , the group $\mathrm{Ext}_{\mathcal{C}}^1(A, B)$ is trivial. Then A is a closed flat subgroup scheme of J .*

Proof. Consider the short exact sequence

$$(1) \quad 0 \longrightarrow J' \longrightarrow J \longrightarrow J/J' = F_2 \longrightarrow 0,$$

where J' is simple. If $J' \cong A$ there is nothing to prove, so assume $A \not\cong J'$. We proceed by induction on the length of the filtration of J . The statement of the lemma holds for length one and two. By induction we have the following exact sequence:

$$0 \longrightarrow A \longrightarrow J/J' = F_2 \longrightarrow F_3 \longrightarrow 0.$$

The pull-back of A by J over F_2 , using (1), gives the short exact sequence

$$0 \longrightarrow J' \longrightarrow J \times_{F_2} A \longrightarrow A \longrightarrow 0.$$

The group scheme $J \times_{F_2} A$ is a closed flat subgroup scheme of J . By hypothesis, $J \times_{F_2} A \simeq A \times J'$. Hence A is a closed flat subgroup scheme of J . \square

Corollary 3.3. *Let J be a finite flat commutative group scheme in the category \mathcal{C} that admits a simple group scheme A . If for each simple B with $B \not\cong A$ occurring in the filtration of J , the group $\text{Ext}_{\mathcal{C}}^1(A, B)$ is trivial, then there exists a closed flat subgroup scheme J' of J admitting only copies of A and such that J/J' does not admit A .*

Proof. We proceed by induction on the length of the filtration of J . If the length of J is one, we are done. If the length is two, we are again done by hypothesis. Suppose the length of J is k and the statement holds if the length is at most $k - 1$. By Lemma 3.2 we can write $0 \subset A \subset J$. By induction, there exists a closed flat subgroup scheme J'' of J/A such that $(J/A)/J''$ does not admit A and J'' only admits copies of A . Then $J' := J \times_{J/A} J''$ verifies the condition of the statement. \square

The next proposition resembles the fact that for finite flat commutative group schemes over a local henselian ring, the quotient by the connected component is an étale group scheme. See for instance [CSS97, p. 138].

Proposition 3.4. *If Condition (1) holds for the category \mathcal{C} , then for any J in \mathcal{C} we have an exact sequence*

$$0 \longrightarrow J' \longrightarrow J \longrightarrow J'' \longrightarrow 0$$

such that J'' is étale and J' does not admit an étale scheme.

Proof. Let J^* be the Cartier dual of J . It suffices to show that J^* contains a subgroup scheme M of multiplicative type such that J^*/M does not admit a simple group scheme of multiplicative type. We may suppose that J^* admits a simple group scheme of multiplicative type; if not, we are done. Then by Lemma 3.2, the group scheme J^* has a simple subgroup scheme of multiplicative type.

Next, suppose that J^* has a subgroup of multiplicative type M' such that J^*/M' admits a simple group scheme of multiplicative type; if not, we are done again. Then again by Lemma 3.2, the group scheme J^*/M' has a simple subgroup scheme of multiplicative type M'' . Now $M'' \times_{J^*/M'} J^*$ is a closed flat subgroup scheme of J^* and sits inside the short exact sequence

$$0 \longrightarrow M' \longrightarrow J^* \times_{J^*/M'} M'' \longrightarrow M'' \longrightarrow 0.$$

Hence $M'' \times_{J^*/M'} J^*$ is an extension of two group schemes of multiplicative type and therefore itself of multiplicative type. Proceeding this way, we find a subgroup scheme of multiplicative type M such that J^*/M does not admit a simple group scheme of multiplicative type. \square

4. APPLICATION TO ABELIAN VARIETIES

In this section, we will prove Theorem 1.2. We first state two auxiliary lemmas:

Lemma 4.1. *Let p be a prime and G be a finite p -group such that $G/[G, G]$ is cyclic. Then G is cyclic.*

Proof. The Frattini subgroup $\text{Frat}(G)$ of G is equal to $[G, G]G^p$. The group $G/\text{Frat}(G)$ is by hypothesis a cyclic group of order p . Burnside's basis Theorem [Hal59, Theorem 12.2.1, p. 176] implies that G is cyclic. \square

Lemma 4.2. *Let G be a group and A, B, C be finite G -modules such that A and C have trivial G -action and G acts faithfully on B . Let*

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

be an exact sequence of G -modules. Let k denote the number of generators of C . Then $\#G$ divides $(\#A)^k$.

Proof. We leave the proof to the reader. □

Lemma 4.3. *Let R be the field as before in Condition (2). If Condition (2) holds for the category \mathcal{C} , then any étale object J in \mathcal{C} becomes constant over R .*

Proof. Let J be any étale group scheme in \mathcal{C} . We claim that $\Lambda = \text{Gal}(F(J_F)/F)$ is an ℓ -group. The proof proceeds by induction on the order of J . There exists an étale group scheme J' in \mathcal{C} such that we have the following short exact sequence of group schemes over the field F :

$$0 \longrightarrow J'_F \longrightarrow J_F \longrightarrow \mathbf{Z}/\ell\mathbf{Z} \longrightarrow 0.$$

By induction, $\text{Gal}(F(J'_F)/F)$ is an ℓ -group. Apply Lemma 4.2 to finish the induction and prove the claim.

We note that $\Lambda/[\Lambda, \Lambda]$ is an abelian ℓ -group and hence the fixed field of $[\Lambda, \Lambda]$ is at most tamely ramified at primes dividing the primes in S . This fixed field is contained in R , which by assumption is a cyclic extension of F . Hence also $\Lambda/[\Lambda, \Lambda]$ is cyclic and by Lemma 4.1 the group Λ is cyclic. We conclude that $F(J_F)$ is contained in R , which is exactly what we wanted to prove. □

For example, if the Hilbert class field of F is trivial and S contains only one prime that does not split in F/K , then R is a cyclic extension of F .

Proposition 4.4. *Let $q \notin S$ be a prime in O_K that is inert in R/K . Suppose that Conditions (1) and (2) hold for the category \mathcal{C} . Then for any J in \mathcal{C} having n simple étale group schemes and m simple group schemes of multiplicative type in its filtration, the following inequalities hold:*

$$|J_q(\mathbf{F}_q)| \geq \ell^n \quad \text{and} \quad |J_q^*(\mathbf{F}_q)| \geq \ell^m.$$

Proof. Let R as before. By Proposition 4.3 all étale objects in \mathcal{C} become constant over R . Let E be the étale quotient of J as in Proposition 3.4. Let \mathfrak{P} be a prime in O_R lying above q . The residue field $\mathbf{F}_{\mathfrak{P}}$ is equal to \mathbf{F}_q . Since E_R is constant, it follows that also $E_{\mathfrak{P}}$ is constant and hence that J_R has at least ℓ^n points in the fiber at \mathfrak{P} . The inequality $|J_q(\mathbf{F}_q)| \geq \ell^n$ follows. The second inequality follows by Cartier duality. □

We are now able to prove Theorem 1.2:

Proof. By contradiction, suppose that $A[\ell]$ contains k simple étale subquotients. Then for any prime q that is not in S and is inert in R/K , Proposition 4.4 says that the number of ℓ -torsion points of A in the fiber at q is at least ℓ^k . Hence $A[\ell^n]$ has at least ℓ^{kn} points in the fiber at q . This is in contradiction with the fact that $A(\mathbf{F}_q)$ is finite for n sufficiently large. Let A^{dual} be the dual abelian variety of A . For each n , the group scheme $A^{\text{dual}}[\ell^n]$ is the Cartier dual of $A[\ell^n]$. If $A[\ell]$ has subquotients of multiplicative type, then $A^{\text{dual}}[\ell]$ has étale subquotients which is impossible by the same argument given above but now applied to the abelian variety A^{dual} . □

We apply Theorem 1.2 together with the three steps described in the introduction to prove:

Theorem 4.5. *There are no non-zero abelian varieties over $\mathbf{Q}(\sqrt{13})$ with good reduction everywhere.*

Proof. We follow the steps mentioned in the introduction:

- (1) We define \mathcal{C} to be the category of finite flat commutative group schemes of 2-power order over $O = \mathbf{Z}[\frac{1+\sqrt{13}}{2}]$.
- (2) By [Fon85] we know that the root discriminant δ of the extension $T_{\mathcal{C}}/\mathbf{Q}$ satisfies $\delta < 4\sqrt{13}$. By Odlyzko's tables this implies that $[T_{\mathcal{C}} : \mathbf{Q}] < 60$. Group schemes in \mathcal{C} annihilated by 2 are isomorphic to $\mu_2, \mathbf{Z}/2\mathbf{Z}$ and the non-trivial extensions of $\mathbf{Z}/2\mathbf{Z}$ by μ_2 described in [KM85, Section 8.7, p.251] using the units -1 and $\eta = \frac{3+\sqrt{13}}{2}$. Hence $T_{\mathcal{C}}$ contains i and the square root of η :

$$\mathbf{Q}(\sqrt{13}) \subset_4 \mathbf{Q}(i, \sqrt{\eta}) \subset_{\leq 7} T_{\mathcal{C}}.$$

The extension $T_{\mathcal{C}}/\mathbf{Q}(i, \sqrt{\eta})$ is unramified outside 2 and is solvable. However, the smallest non-trivial abelian extension unramified outside 2 of $\mathbf{Q}(i, \sqrt{\eta})$ is a subfield of the ray class field of conductor π_2^6 , where π_2 is the unique prime above 2 in $\mathbf{Q}(i, \sqrt{\eta})$. This subfield violates the root discriminant bound on $T_{\mathcal{C}}$. It follows that $T_{\mathcal{C}} = \mathbf{Q}(i, \sqrt{\eta})$. By Lemma 2.2 this implies that every simple object in \mathcal{C} has rank 2.

Since 2 is inert in $\mathbf{Q}(\sqrt{13})$, this implies by [TO70, Corollary, p.21] that the simple group schemes in \mathcal{C} are μ_2 and $\mathbf{Z}/2\mathbf{Z}$. One now checks that Condition (2) is satisfied.

- (3) For this category, Now we use Theorem [Sch03, Prop. 2.6] to verify that Condition (1) is satisfied.

The 2-torsion of a non-zero abelian variety over $\mathbf{Q}(\sqrt{13})$ with good reduction everywhere is an object in \mathcal{C} , and this 2-torsion subgroup scheme must be filtered by copies of μ_2 or $\mathbf{Z}/2\mathbf{Z}$. This, however, contradicts Theorem 1.2. □

As another example, we show that:

Theorem 4.6. *There are no non-zero abelian varieties over $\mathbf{Q}(\sqrt{17})$ with good reduction everywhere.*

Proof. We follow the steps mentioned in the introduction:

- (1) Let \mathcal{C} be the category of finite flat commutative group schemes of 2-power order over $O = \mathbf{Z}[\frac{1+\sqrt{17}}{2}]$. We will see that the category \mathcal{C} does not satisfy Condition (1) of Theorem 1.2.
- (2) We find the maximal 2-torsion extension $T_{\mathcal{C}}/\mathbf{Q}(\sqrt{17})$ of \mathcal{C} . We leave it as an exercise to show that the extension $T_{\mathcal{C}}/\mathbf{Q}(\sqrt{17})$ is finite and has degree a power of 2. So we can apply Lemma 2.2. By factoring $2 = \pi\bar{\pi}$ in O we find the following simple group schemes: $\mu_2, \mathbf{Z}/2\mathbf{Z}, G_{\pi}$ and $G_{\bar{\pi}}$, where we refer to [TO70] for the meaning of G_{π} and $G_{\bar{\pi}}$.
- (3) The only simple étale group scheme is $\mathbf{Z}/2\mathbf{Z}$ and we immediately verify Condition (2) for our category \mathcal{C} . However, Condition (1) fails because $\text{Ext}_O^1(\mu_2, \mathbf{Z}/2\mathbf{Z})$ is non-trivial due to the splitting of the prime 2 in $\mathbf{Q}(\sqrt{17})/\mathbf{Q}$: A non-trivial extension is given by $G_{\pi} \times G_{\bar{\pi}}$.

Even though Condition (1) does not hold, it is true that all extensions of simple non-étale group schemes by simple étale group schemes are annihilated by 2: they are

products of G_π 's and $G_{\bar{\pi}}$'s. Using this, for any abelian variety A over $\mathbf{Q}(\sqrt{17})$ with good reduction everywhere one deduces that the rank of $A[2^n]$ (which is an object in \mathcal{C}) cannot depend on n . Hence there are no such non-zero abelian varieties. \square

We end this article by asking for which square-free integers D do there exist abelian varieties over $\mathbf{Q}(\sqrt{D})$ with good reduction everywhere?

REFERENCES

- [CSS97] Gary Cornell, Joseph H. Silverman, and Glenn Stevens, editors. *Modular forms and Fermat's last theorem*. Springer-Verlag, New York, 1997. Papers from the Instructional Conference on Number Theory and Arithmetic Geometry held at Boston University, Boston, MA, August 9–18, 1995.
- [Fon85] Jean-Marc Fontaine. Il n'y a pas de variété abélienne sur \mathbf{Z} . *Invent. Math.*, 81(3):515–538, 1985.
- [Hal59] Jr. Marshall Hall. *The theory of groups*. The Macmillan Co., New York, N.Y., 1959.
- [KM85] Nicholas M. Katz and Barry Mazur. *Arithmetic moduli of elliptic curves*, volume 108 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1985.
- [Ray74] Michel Raynaud. Schémas en groupes de type (p, \dots, p) . *Bull. Soc. Math. France*, 102:241–280, 1974.
- [Sch03] René Schoof. Abelian varieties over cyclotomic fields with good reduction everywhere. *Math. Ann.*, 325(3):413–448, 2003.
- [Sch05] René Schoof. Abelian varieties over \mathbf{Q} with bad reduction in one prime only. *Compos. Math.*, 141(4):847–868, 2005.
- [TO70] John Tate and Frans Oort. Group schemes of prime order. *Ann. Sci. École Norm. Sup. (4)*, 3:1–21, 1970.