

# An asymptotic formula of the divergent bilateral basic hypergeometric series

Takeshi MORITA\*

## Abstract

We show an asymptotic formula of the divergent bilateral basic hypergeometric series  ${}_1\psi_0(a; -; q, \cdot)$  with using the  $q$ -Borel-Laplace method. We also give the limit  $q \rightarrow 1 - 0$  of our asymptotic formula.

## 1 Introduction

In this paper, we show an asymptotic formula of the divergent bilateral basic hypergeometric series

$${}_1\psi_0(a; -; q; x) := \sum_{n \in \mathbb{Z}} (a; q)_n \left\{ (-1)^n q^{\frac{n(n-1)}{2}} \right\}^{-1} x^n.$$

Here,  $(a; q)_n$  is the  $q$ -shifted factorial;

$$(a; q)_n := \begin{cases} 1, & n = 0, \\ (1-a)(1-aq) \dots (1-aq^{n-1}), & n \geq 1, \\ [(1-aq^{-1})(1-aq^{-2}) \dots (1-aq^n)]^{-1}, & n \leq -1 \end{cases}$$

moreover,  $(a; q)_\infty := \lim_{n \rightarrow \infty} (a; q)_n$  and

$$(a_1, a_2, \dots, a_m; q)_\infty := (a_1; q)_\infty (a_2; q)_\infty \dots (a_m; q)_\infty.$$

The  $q$ -shifted factorial  $(a; q)_n$  is a  $q$ -analogue of the shifted factorial

$$(\alpha)_n = \alpha \{\alpha + 1\} \dots \{\alpha + (n - 1)\}.$$

---

\*Graduate School of Information Science and Technology, Osaka University, 1-1 Machikaneyama-machi, Toyonaka, 560-0043, Japan.

The series  ${}_1\psi_0(a; -, q, \cdot)$  is related to the bilateral basic hypergeometric series  ${}_1\psi_1(a; b; q, \cdot)$ . At first, we review the series

$${}_1\psi_1(a; b; q, z) := \sum_{n \in \mathbb{Z}} \frac{(a; q)_n}{(b; q)_n} z^n.$$

This series has Ramanujan's summation formula [3]

$${}_1\psi_1(a; b; q, z) = \frac{(q, b/a, az, q/az; q)_\infty}{(b, q/a, z, b/az; q)_\infty}, \quad |b/a| < |z| < 1, \quad (1)$$

which was first given by Ramanujan [4]. This formula is considered as an extension of the  $q$ -binomial theorem [3, 1];

$$\sum_{n \geq 0} \frac{(a; q)_n}{(q; q)_n} z^n = \frac{(az; q)_\infty}{(z; q)_\infty}, \quad |z| < 1.$$

We also regard the summation (1) as a  $q$ -analogue of the bilateral binomial theorem [6] discovered by M. E. Horn. If  $\alpha$  and  $\beta$  are complex numbers  $\Re(\beta - \alpha) > 1$  and  $z$  is a complex number with  $|z| = 1$ , then

$$\sum_{n \in \mathbb{Z}} \frac{(\alpha)_n}{(\beta)_n} z^n = \frac{(1-z)^{\beta-\alpha-1} \Gamma(1-\alpha) \Gamma(\beta)}{(-z)^{\beta-1} \Gamma(\beta-\alpha)}. \quad (2)$$

We remark that the limit  $q \rightarrow 1 - 0$  of (1) with suitable condition gives the bilateral binomial theorem (2).

The series  ${}_1\psi_1(a; b; q, z)$  satisfies the following  $q$ -difference equation

$$\left(\frac{b}{q} - az\right) u(qz) + (z-1)u(z) = 0.$$

In this paper, we study the degeneration of this equation as follows;

$$\left(\frac{1}{q} - ax\right) \tilde{u}(qx) + x\tilde{u}(x) = 0.$$

This equation has a formal solution

$$\tilde{u}(x) = {}_1\psi_0(a; -, q, x). \quad (3)$$

But  $\tilde{u}(x)$  is a divergent around the origin [3] and its properties are not clear. In section three, we show an asymptotic formula of the series (3) with using the  $q$ -Borel-Laplace transformations. The  $q$ -Borel-Laplace transformations are introduced in the study of connection problems on linear  $q$ -difference equations between the origin and the infinity.

Connection problems on  $q$ -difference equations with regular singular points are studied by G. D. Birkhoff [2]. The first example of the connection formula is given by G. N. Watson [13] in 1910 as follows;

$$\begin{aligned} {}_2\varphi_1(a, b; c; q, x) &= \frac{(b, c/a; q)_\infty (ax, q/ax; q)_\infty}{(c, b/a; q)_\infty (x, q/x; q)_\infty} {}_2\varphi_1(a, aq/c; aq/b; q, cq/abx) \\ &+ \frac{(a, c/b; q)_\infty (bx, q/bx; q)_\infty}{(c, a/b; q)_\infty (x, q/x; q)_\infty} {}_2\varphi_1(b, bq/c; bq/a; q, cq/abx). \end{aligned} \quad (4)$$

The function  ${}_2\varphi_1(a, b; c; q, x)$  is the basic hypergeometric series

$${}_2\varphi_1(a, b; c; q, x) = \sum_{n \geq 0} \frac{(a, b; q)_n}{(c; q)_n (q; q)_n} x^n,$$

which is introduced by E. Heine [5]. This series satisfies the second order linear  $q$ -difference equation

$$(c - abqx)u(q^2x) - \{c + q - (a + b)qx\}u(qx) + q(1 - x)u(x) = 0. \quad (5)$$

around the origin. On the other hand, the equation (5) has a fundamental system of solutions

$$v_1(x) = \frac{\theta(-ax)}{\theta(-x)} {}_2\varphi_1\left(a, \frac{aq}{c}; \frac{aq}{b}; q, \frac{cq}{abx}\right), \quad v_2(x) = \frac{\theta(-bx)}{\theta(-x)} {}_2\varphi_1\left(b, \frac{bq}{c}; \frac{bq}{a}; q, \frac{cq}{abx}\right)$$

around the infinity [14]. Here,  $\theta(\cdot)$  is the theta function of Jacobi (see section two). In the connection formula (4), we remark that the connection coefficients are given by  $q$ -elliptic functions.

Other connection formulae, especially connection formulae of irregular singular type  $q$ -special functions has not known for a long time. Recently, C. Zhang gives some connection formulae by two different types of  $q$ -Borel transformations and  $q$ -Laplace transformations [14, 15, 16], which are introduced by J. Sauloy[11]. We assume that  $f(x)$  is a formal power series  $f(x) = \sum_{n \geq 0} a_n x^n$ ,  $a_0 = 1$ .

**Definition 1.** For any power series  $f(x)$ , the  $q$ -Borel transformation of the first kind  $\mathcal{B}_q^+$  is

$$(\mathcal{B}_q^+ f)(\xi) := \sum_{n \geq 0} a_n q^{\frac{n(n-1)}{2}} \xi^n =: \varphi(\xi).$$

For an entire function  $\varphi$ , the  $q$ -Laplace transformation of the first kind  $\mathcal{L}_{q,\lambda}^+$  is

$$(\mathcal{L}_{q,\lambda}^+ \varphi)(x) := \sum_{n \in \mathbb{Z}} \frac{\varphi(\lambda q^n)}{\theta\left(\frac{\lambda q^n}{x}\right)}, \quad \lambda \notin q^{\mathbb{Z}}.$$

Similarly, the  $q$ -Borel transformation of the second kind  $\mathcal{B}_q^-$  is

$$(\mathcal{B}_q^- f)(\xi) := \sum_{n \geq 0} a_n q^{-\frac{n(n-1)}{2}} \xi^n =: g(\xi)$$

and the  $q$ -Laplace transformation of the second kind  $\mathcal{L}_q^-$  is

$$(\mathcal{L}_q^- g)(x) := \frac{1}{2\pi i} \int_{|\xi|=r} g(\xi) \theta\left(\frac{x}{\xi}\right) \frac{d\xi}{\xi}.$$

We remark that each  $q$ -Borel transformation is a formal inverse of each  $q$ -Laplace transformation;

$$\mathcal{L}_{q,\lambda}^+ \circ \mathcal{B}_q^+ f = f, \quad \mathcal{L}_q^- \circ \mathcal{B}_q^- f = f.$$

We can find applications of the  $q$ -Borel-Laplace method of the first kind in [14, 15, 10]. This summation method is powerful to deal with divergent type basic hypergeometric series and to study the  $q$ -Stokes phenomenon. Applications of the method of the second kind can be found in [16, 8, 9, 10]. But other examples, for example, applications to bilateral basic hypergeometric series, has not known.

In this paper, we study an asymptotic formula of the divergent bilateral basic hypergeometric series  ${}_1\psi_0(a; -; q, \cdot)$  from viewpoint of connection problems on linear  $q$ -difference equations. We show the following theorem.

**Theorem.** For any  $x \in \mathbb{C}^* \setminus [-\lambda; q]$ , we have

$$\hat{\psi}_0(a; -; \lambda; q, x) = \frac{(q; q)_\infty}{(q/a; q)_\infty} \frac{\theta(aq\lambda)}{\theta(q\lambda)} \frac{\theta(ax/\lambda)}{\theta(x/\lambda)} \frac{1}{(1/ax; q)_\infty},$$

where  $1 < |ax|$ .

Here, the function  ${}_1\hat{\psi}_0(a; -; \lambda; q, x)$  is the  $q$ -Borel-Laplace transform of the series  ${}_1\psi_0(a; -; q, x)$  and the set  $[-\lambda; q]$  is the  $q$ -spiral which is given by  $[\lambda; q] := \{\lambda q^k : k \in \mathbb{Z}\}$ . We also show the limit  $q \rightarrow 1 - 0$  of our asymptotic formula in section four. If we take a suitable limit  $q \rightarrow 1 - 0$  of the theorem above, we formally obtain the asymptotic formula of the bilateral hypergeometric series  ${}_1H_0$ .

## 2 Basic notations

In this section, we fix our notations. We assume that  $0 < |q| < 1$  and  $\sigma_q$  is the  $q$ -difference operator such that  $\sigma_q f(x) = f(qx)$ . The basic hypergeometric series is

$${}_r\varphi_s(a_1, \dots, a_r; b_1, \dots, b_s; q, x) := \sum_{n \geq 0} \frac{(a_1, \dots, a_r; q)_n}{(b_1, \dots, b_s; q)_n (q; q)_n} \left\{ (-1)^n q^{\frac{n(n-1)}{2}} \right\}^{1+s-r} x^n. \quad (6)$$

The radius of convergence  $\rho$  of the basic hypergeometric series is given by [7]

$$\rho = \begin{cases} \infty, & \text{if } r < s + 1, \\ 1, & \text{if } r = s + 1, \\ 0, & \text{if } r > s + 1. \end{cases}$$

The bilateral basic hypergeometric series is

$${}_r\psi_s(a_1, \dots, a_r; b_1, \dots, b_s; q, x) := \sum_{n \in \mathbb{Z}} \frac{(a_1, \dots, a_r; q)_n}{(b_1, \dots, b_s; q)_n} \left\{ (-1)^n q^{\frac{n(n-1)}{2}} \right\}^{s-r} x^n. \quad (7)$$

If  $s < r$ , the series (7) diverges for  $x \neq 0$  and if  $r = s$ , the series (7) converges  $|b_1 \dots b_s / a_1 \dots a_r| < |x| < 1$  (see [3] for more detail). The series (6) is a  $q$ -analogue of the generalized hypergeometric function

$${}_rF_s(\alpha_1, \dots, \alpha_r; \beta_1, \dots, \beta_s; x) := \sum_{n \geq 0} \frac{(\alpha_1, \dots, \alpha_r)_n}{(\beta_1, \dots, \beta_s)_n n!} x^n$$

and the series (7) is a  $q$ -analogue of the bilateral hypergeometric function

$${}_rH_s(\alpha_1, \dots, \alpha_r; \beta_1, \dots, \beta_s; x) := \sum_{n \in \mathbb{Z}} \frac{(\alpha_1, \dots, \alpha_r)_n}{(\beta_1, \dots, \beta_s)_n} x^n.$$

By D'Alembert's ratio test, it can be checked that  ${}_rH_r$  converges only for  $|x| = 1$  [12], provided that  $\Re(\beta_1 + \cdots + \beta_r - \alpha_1 - \cdots - \alpha_r) > 1$ .

The  $q$ -exponential function  $e_q(x)$  is

$$e_q(x) := {}_1\varphi_0(0; -; q, x) = \sum_{n \geq 0} \frac{x^n}{(q; q)_n}$$

The function  $e_q(x)$  has the infinite product representation as follows;

$$e_q(x) = \frac{1}{(x; q)_\infty}, \quad |x| < 1.$$

The limit  $q \rightarrow 1 - 0$  of  $e_q(x)$  is the exponential function [3];

$$\lim_{q \rightarrow 1-0} e_q(x(1-q)) = e^x. \quad (8)$$

The  $q$ -gamma function  $\Gamma_q(x)$  is

$$\Gamma_q(x) := \frac{(q; q)_\infty}{(q^x; q)_\infty} (1-q)^{1-x}, \quad 0 < q < 1.$$

The limit  $q \rightarrow 1 - 0$  of  $\Gamma_q(x)$  gives the gamma function [3];

$$\lim_{q \rightarrow 1-0} \Gamma_q(x) = \Gamma(x). \quad (9)$$

The theta function of Jacobi is given by

$$\theta(x) := \sum_{n \in \mathbb{Z}} q^{\frac{n(n-1)}{2}} x^n, \quad \forall x \in \mathbb{C}^*.$$

Jacobi's triple product identity is

$$\theta(x) = (q, -x, -q/x; q)_\infty$$

and the theta function satisfies the following  $q$ -difference equation

$$\theta(q^k x) = x^{-k} q^{-\frac{k(k-1)}{2}} \theta(x), \quad \forall k \in \mathbb{Z}.$$

The inversion formula is

$$\theta(x) = x\theta(1/x). \quad (10)$$

We remark that  $\theta(\lambda q^k/x) = 0$  if and only if  $x \in [-\lambda; q]$ . In our study, the following proposition [15] is useful to consider the limit  $q \rightarrow 1 - 0$  of our formula.

**Proposition 1.** For any  $x \in \mathbb{C}^*$  ( $-\pi < \arg x < \pi$ ), we have

$$\lim_{q \rightarrow 1-0} \frac{\theta(q^\alpha x)}{\theta(q^\beta x)} = x^{\beta-\alpha} \quad (11)$$

and

$$\lim_{q \rightarrow 1-0} \frac{\theta\left(\frac{q^\alpha x}{(1-q)}\right)}{\theta\left(\frac{q^\beta x}{(1-q)}\right)} (1-q)^{\beta-\alpha} = \left(\frac{1}{x}\right)^{\alpha-\beta}. \quad (12)$$

We also use the limit [3] as follows;

$$\lim_{q \rightarrow 1-0} \frac{(xq^\alpha; q)_\infty}{(x; q)_\infty} = (1-x)^{-\alpha}, \quad |x| < 1. \quad (13)$$

### 3 An asymptotic formula of the divergent series ${}_1\psi_0(a; -; q, x)$

In this section, we show an asymptotic formula of the divergent series  ${}_1\psi_0(a; -; q, x)$ . At first, we review Ramanujan's sum for  ${}_1\psi_1(a; b; q, z)$  and its property. Ramanujan gives the following sum [4]

$${}_1\psi_1(a; b; q, z) = \frac{(q, b/a, az, q/az; q)_\infty}{(b, q/a, z, b/az; q)_\infty}, \quad |b/a| < |z| < 1. \quad (14)$$

We can regard this sum as a  $q$ -analogue of the bilateral binomial theorem [6] as follows;

**Theorem.** (Horn [6]) If  $z$  is a complex number such that  $|z| = 1$ , we have

$${}_1H_1(\alpha; \beta; z) = \frac{\Gamma(\beta)\Gamma(1-\alpha)}{\Gamma(\beta-\alpha)} \frac{(1-z)^{\beta-\alpha-1}}{(-z)^{\beta-1}}, \quad (15)$$

provided that  $\Re(\beta - \alpha) > 1$ .

Let  $z$  is a complex number such that  $-\pi < \arg z < \pi$ . The summation (14) is rewritten by

$$\frac{(q, b/a, az, q/az; q)_\infty}{(b, q/a, z, b/az; q)_\infty} = \frac{(q, b/a; q)_\infty}{(b, q/a; q)_\infty} \frac{\theta(-az)}{\theta(-aqz/b)} \frac{(aqz/b; q)_\infty}{(z; q)_\infty}. \quad (16)$$

In (16), we put  $a = q^\alpha$  and  $b = q^\beta$  (with  $\Re(\beta - \alpha) > 1$ ), we obtain

$${}_1\psi_1(q^\alpha; q^\beta; q, z) = \frac{\Gamma_q(\beta)\Gamma_q(1 - \alpha)}{\Gamma_q(\beta - \alpha)} \frac{\theta(q^\alpha(-z))}{\theta(q^{\alpha+1-\beta}(-z))} \frac{(q^{\alpha+1-\beta}z; q)_\infty}{(z; q)_\infty}, \quad (17)$$

where  $|q^{\beta-\alpha}| < |z| < 1$ . Combining (9), (11) and (13), we can check out that the limit  $q \rightarrow 1 - 0$  of (17) gives the bilateral binomial theorem (15), provided that  $-\pi < \arg z < \pi$ .

The bilateral basic hypergeometric series  ${}_1\psi_1(a; b; q, z)$  satisfies the  $q$ -difference equation

$$\left(\frac{b}{q} - az\right)u(qz) + (z - 1)u(z) = 0. \quad (18)$$

We consider the degeneration of the equation (18) in the next section.

### 3.1 Local solutions of the degenerated equation

In the equation (18), we put  $z = bx$  and take the limit  $b \rightarrow \infty$ , we obtain the equation

$$\left\{ \left( \frac{1}{q} - ax \right) \sigma_q + x \right\} \tilde{u}(x) = 0. \quad (19)$$

The formal solution of (19) is

$$\tilde{u}(x) = {}_1\psi_0(a; -, q, x), \quad (20)$$

which is divergent around the origin. We consider “the basic hypergeometric type” solution around the infinity. We assume that the solution around the infinity is given by

$$\tilde{u}_\infty(x) := \frac{\theta(ax)}{\theta(x)} v_\infty(x) = \frac{\theta(ax)}{\theta(x)} \sum_{n \geq 0} v_n x^{-n}, \quad v_0 = 1.$$

By the  $q$ -difference equation of the theta function, we obtain the equation

$$\left\{ \left( \frac{1}{aq} - x \right) \sigma_q + x \right\} v_\infty(x) = 0.$$

We remark that the function  $\theta(ax)/\theta(x)$  satisfies the following  $q$ -difference equation

$$u(qx) = q^{-\alpha} u(x)$$



which is also satisfied by the function  $u(x) = x^{-\alpha}$  with  $\log_q a = \alpha$ . We can check out that the series  $v_\infty(x)$  is

$$v_\infty(x) = e_q \left( \frac{1}{ax} \right) = \frac{1}{(1/ax; q)_\infty}, \quad \left| \frac{1}{ax} \right| < 1.$$

Therefore, one of the solution of (19) around the infinity is

$$\tilde{u}_\infty(x) = \frac{\theta(ax)}{\theta(x)} e_q \left( \frac{1}{ax} \right). \quad (21)$$

In the following section, we study the relation between these solutions (20) and (21) with using the  $q$ -Borel-Laplace method of the first kind.

**Remark 1.** *We remark that “the bilateral basic hypergeometric type solution” of (19) around the infinity is given by*

$$w_\infty(x) = {}_1\psi_1 \left( 0; \frac{q}{a}; q, \frac{1}{ax} \right) = \sum_{n \in \mathbb{Z}} \frac{1}{(q/a; q)_n} \left( \frac{1}{ax} \right)^n.$$

*But this solution is not suitable for our argument. We choose and deal with the solution (21).*

## 3.2 An asymptotic formula

In this section, we show an asymptotic formula of (20) with using the  $q$ -Borel-Laplace transformations of the first kind. We show the following theorem.

**Theorem 1.** *For any  $x \in \mathbb{C}^* \setminus [-\lambda; q]$ , we have*

$${}_1\hat{\psi}_0(a; -; \lambda; q, x) = \frac{(q; q)_\infty}{(q/a; q)_\infty} \frac{\theta(aq\lambda)}{\theta(q\lambda)} \frac{\theta(ax/\lambda)}{\theta(x/\lambda)} \frac{1}{(1/ax; q)_\infty},$$

where  $1 < |ax|$ .

*Proof.* We apply the  $q$ -Borel transformation  $\mathcal{B}_q^+$  to the series  ${}_1\psi_0(a; -; q, x)$ . Then,

$$\psi(\xi) := (\mathcal{B}_q^+ {}_1\psi_0)(\xi) = {}_1\psi_1(a; 0; q, -\xi).$$

By Ramanujan’s sum (14), we obtain the infinite product representation of  $\psi(\xi)$  as follows;

$$\psi(\xi) = \frac{(q; q)_\infty}{(q/a; q)_\infty} \frac{\theta(a\xi)}{\theta(\xi)} (-q/\xi; q)_\infty.$$

We apply the  $q$ -Laplace transformation  $\mathcal{L}_{q,\lambda}^+$  to  $\psi(\xi)$ .

$$\begin{aligned}
(\mathcal{L}_{q,\lambda}^+ \psi)(x) &= \sum_{n \in \mathbb{Z}} \frac{\psi(\lambda q^n)}{\theta\left(\frac{\lambda q^n}{x}\right)} = \frac{(q; q)_\infty}{(q/a; q)_\infty} \sum_{n \in \mathbb{Z}} \frac{1}{\theta\left(\frac{\lambda q^n}{x}\right)} \frac{\theta(a\lambda q^n)}{\theta(\lambda q^n)} \left(-\frac{q}{\lambda q^n}; q\right)_\infty \\
&= \frac{(q; q)_\infty}{(q/a; q)_\infty} \frac{1}{\theta(\lambda/x)} \frac{\theta(a\lambda)}{\theta(\lambda)} \sum_{n \in \mathbb{Z}} q^{\frac{n(n-1)}{2}} \left(\frac{\lambda}{ax}\right)^n \left(-\frac{q}{\lambda q^n}; q\right)_\infty \\
&= \frac{(q; q)_\infty}{(q/a; q)_\infty} \frac{1}{\theta(\lambda/x)} \frac{\theta(a\lambda)}{\theta(\lambda)} (-q/\lambda; q)_{\infty 1} \psi_1(-\lambda; 0; q, 1/ax). \quad (22)
\end{aligned}$$

We remark that

$${}_1\psi_1(-\lambda; 0; q, 1/ax) = \frac{\theta(\lambda/ax)}{(-q/\lambda; q)_\infty (1/ax; q)_\infty}.$$

Combining (22) and (10), we obtain the conclusion.  $\square$

Theorem 1 is rewritten by

$$(\mathcal{L}_{q,\lambda}^+ \circ \mathcal{B}_q^+ \tilde{u})(x) = \frac{(q; q)_\infty}{(q/a; q)_\infty} \frac{\theta(aq\lambda)}{\theta(q\lambda)} \frac{\theta(ax/\lambda)}{\theta(x/\lambda)} \frac{\theta(x)}{\theta(ax)} \tilde{u}_\infty(x).$$

We define the function  $C(x; q)$  as follows;

$$C(x; q) := \frac{(q; q)_\infty}{(q/a; q)_\infty} \frac{\theta(aq\lambda)}{\theta(q\lambda)} \frac{\theta(ax/\lambda)}{\theta(x/\lambda)} \frac{\theta(x)}{\theta(ax)}.$$

Then,  $C(x; q)$  is single valued as a function of  $x$ .

**Corollary 1.** *The function  $C(x; q)$  is the  $q$ -elliptic function, i.e.,  $C(x; q)$  satisfies*

$$C(qx; q) = C(x; q), \quad C(e^{2\pi i} x; q) = C(x; q).$$

## 4 The limit $q \rightarrow 1 - 0$ of the asymptotic formula

In this section, we give the limit  $q \rightarrow 1 - 0$  of our asymptotic formula. We assume that  $x \in \mathbb{C}^* \setminus [-\lambda; q]$  ( $-\pi < \arg x < \pi$ ). We put  $a = q^\alpha$  and  $x \mapsto x/(1 - q)$  in theorem 1 to consider the limit. We remark that the limit of the left-hand side in theorem 1 formally gives the bilateral hypergeometric series  ${}_1H_0(\alpha; -; -x)$ . We show the following theorem.

**Theorem 2.** For any  $x \in \mathbb{C}^* \setminus [-\lambda; q]$  ( $-\pi < \arg x < \pi$ ), we have

$$\lim_{q \rightarrow 1-0} \frac{(q; q)_\infty}{(q^{1-\alpha}; q)_\infty} \frac{\theta(q^{\alpha+1}\lambda)}{\theta(q\lambda)} \frac{\theta\left(\frac{q^\alpha x}{(1-q)\lambda}\right)}{\theta\left(\frac{x}{(1-q)\lambda}\right)} \frac{1}{\left((1-q)\frac{q^{-\alpha}}{x}; q\right)_\infty} = \frac{\Gamma(1-\alpha)}{x^\alpha} e^{\frac{1}{x}},$$

where  $1 < |x|$ .

We give the proof of theorem 2.

*Proof.* The right-hand side of theorem 1 is rewritten by

$$\begin{aligned} & \frac{(q; q)_\infty}{(q^{1-\alpha}; q)_\infty} \frac{\theta(q^{\alpha+1}\lambda)}{\theta(q\lambda)} \frac{\theta\left(\frac{q^\alpha}{(1-q)} \frac{x}{\lambda}\right)}{\theta\left(\frac{1}{(1-q)} \frac{x}{\lambda}\right)} \frac{1}{\left((1-q)\frac{q^{-\alpha}}{x}; q\right)_\infty} \\ &= \Gamma_q(1-\alpha) \frac{\theta(q^{\alpha+1}\lambda)}{\theta(q\lambda)} \left\{ \frac{\theta\left(\frac{q^\alpha}{(1-q)} \frac{x}{\lambda}\right)}{\theta\left(\frac{q^0}{(1-q)} \frac{x}{\lambda}\right)} (1-q)^{-\alpha} \right\} e_q\left(\left(1-q\right)\frac{1}{q^\alpha x}\right). \end{aligned}$$

Combining (9), (11), (12) and (8), we obtain the conclusion.  $\square$

## Acknowledgement

My heartfelt appreciation goes to Professor Yousuke Ohyama who provided carefully considered feedback and valuable comments.

## References

- [1] G. E. Andrews, R. Askey and R. Roy, Special Functions, Encyclopedia of Math. Appl. 71, Cambridge Univ. Press, Cambridge, 1999.
- [2] G. D. Birkhoff, The generalized Riemann problem for linear differential equations and the allied problems for linear difference and  $q$ -difference equations, Proc. Am. Acad. Arts and Sciences, **49** (1914), 521 – 568.
- [3] G. Gasper and M. Rahman, Basic Hypergeometric Series, 2nd ed, Cambridge, 2004.

- [4] G. H. Hardy, *Ramanujan*, Cambridge University Press, Cambridge; reprinted by Chelsea, New York, 1978.
- [5] Heine. E. Untersuchungen über die Reihe..., *J. reine angew. Math.* **34**, 285-328.
- [6] M. E. Horn, Bilateral binomial theorem, SIAM Problem **03-001** (2003).
- [7] T. H. Koornwinder and R. F. Swarttouw, On  $q$ -analogues of the Fourier and Hankel transforms, *Trans. Amer. Math. Soc.* **333** (1992), 445-461.
- [8] T. Morita, A connection formula of the Hahn-Exton  $q$ -Bessel Function, *SIGMA*,**7** (2011), 115, 11pp.
- [9] T. Morita, A connection formula of the  $q$ -confluent hypergeometric function, [arXiv:1105.5770](#).
- [10] T. Morita, On local solutions of the Ramanujan equation and their connection formulae, [arXiv:1203.3404](#).
- [11] J. Sauloy, Algebraic construction of the Stokes sheaf for irregular linear  $q$ -difference equations, [arXiv:math/0409393](#)
- [12] L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966. MR0201688 (34:1570)
- [13] G. N. Watson, The continuation of functions defined by generalized hypergeometric series, *Trans. Camb. Phil. Soc.* **21** (1910), 281–299.
- [14] C. Zhang, Remarks on some basic hypergeometric series, in “Theory and Applications of Special Functions”, Springer (2005), 479–491.
- [15] C. Zhang, Sur les fonctions  $q$ -Bessel de Jackson, *J. Approx. Theory*, **122** (2003), 208–223.
- [16] C. Zhang, Une sommation discrète pour des équations aux  $q$ -différences linéaires et à coefficients, analytiques: théorie générale et exemples, in “Differential Equations and Stokes Phenomenon”, World Scientific (2002), 309–329.