

Local GW Invariants of Elliptic Multiple Fibers

Junho Lee
University of Central Florida
Orlando, FL 32816

Abstract

We use simple geometric arguments to calculate the dimension zero local Gromov-Witten invariants of elliptic multiple fibers. This completes the calculation of all dimension zero GW invariants of elliptic surfaces with $p_g > 0$.

Let X be a Kähler surface with $p_g > 0$. By the Enriques-Kodaira classification (cf. [BHPV]), its minimal model is a $K3$ or Abelian surface, a surface of general type or an elliptic surface. Each holomorphic 2-form α on X defines an almost complex structure

$$J_\alpha = (Id + JK_\alpha) J (Id + JK_\alpha). \quad (0.1)$$

Here, J is the complex structure on X and the endomorphism K_α of TX is defined by the formula $\langle u, K_\alpha v \rangle = \alpha(u, v)$ where $\langle \cdot, \cdot \rangle$ is the Kähler metric. This J_α satisfies:

Lemma 0.1 ([L]). *If f is a J_α -holomorphic map that represents a nontrivial (1,1) class then its image lies in the support of the zero divisor D_α of α and f is, in fact, holomorphic.*

The Gromov-Witten invariant $GW_{g,n}(X, A)$ is a (virtual) count of holomorphic maps representing the class A . In particular, the invariant $GW_{g,n}(X, A)$ vanishes unless A is a (1,1) class since every holomorphic map represents (1,1) class. Note that each canonical divisor D of X is a zero divisor of a holomorphic 2-form. Lemma 0.1 thus shows that the GW invariant is a sum

$$GW_{g,n}(X, A) = \sum GW_{g,n}^{loc}(D_k, A_k)$$

over the connected components D_k of the canonical divisor D of local invariants that counts the contribution of maps whose image lies in D_k (cf. [LP], [KL]). It follows that the GW invariants of minimal $K3$ or Abelian surfaces are trivial except possibly for the trivial homology class because their canonical divisors are trivial.

The local GW invariants have a universal property. If X is a minimal surface of general type with a smooth canonical divisor D then the local invariants associated with D , and hence GW invariants, are determined by the normal bundle of D — in fact, there exists a universal function of c_1^2 and c_2 that gives the GW invariants of X (cf. Section 7 of [LP]).

If $\pi : X \rightarrow C$ is a minimal elliptic surface with $p_g > 0$, after suitable deformation, we can assume X has a canonical divisor of the form

$$\sum_i n_i F^i + \sum_k (m_k - 1) F_{m_k}$$

where F^i is a regular fiber and F_{m_k} is a smooth multiple fiber of multiplicity m_k (cf. Proposition 6.1 of [LP]). In this case, the GW invariants of X are sums of universal functions, and are completely determined by the multiplicities m_k and the number

$$c_\pi = \chi(\mathcal{O}_X) - 2\chi(\mathcal{O}_C)$$

(cf. Section 6 of [LP]). In particular, the generating function for the set of all *dimension zero* GW invariants of X is given by

$$GW_X^0 = c_\pi \sum_{d>0} GW_1^{loc}(F, d) t^d + \sum_k \sum_{d>0} GW_1^{loc}(F_{m_k}, d) t_{m_k}^d \quad (0.2)$$

where the formal variables t and t_{m_k} are for the fiber class $[F]$ and the multiple fiber classes $[F_{m_k}]$ respectively; these satisfy $t_{m_k}^{m_k} = t$. The local invariants in (0.2) are counts of multiple covers of elliptic curves together with signs determined by the GW theory of 4-manifolds.

Some of the generating functions in (0.2) are known. In cases of the regular fiber F and the multiple fiber F_2 , it was proved in Section 10 of [LP] that

$$GW_1^{loc}(F, d) = -\frac{1}{d} \sigma(d) \quad \text{and} \quad GW_1^{loc}(F_2, d) = \frac{1}{d} \left(\sigma(d) - 2\sigma\left(\frac{d}{2}\right) \right) \quad (0.3)$$

where $\sigma(d) = \sum_{k|d} k$ if d is a positive integer and $\sigma(d) = 0$ otherwise. In this note we use geometric arguments to obtain the terms in (0.2) associated with fibers of higher multiplicity. Our main theorem is the following formula for the local invariants $GW_1^{loc}(F_m, d)$ for $m > 2$. This completes the calculation of all dimension zero GW invariants of all minimal elliptic surfaces with $p_g > 0$.

Main Theorem. *Let $m \geq 3$. Then*

$$GW_1^{loc}(F_m, d) = \frac{1}{d} \left(\sigma(d) - m \sigma\left(\frac{d}{m}\right) \right).$$

The contribution of each degree d cover f of elliptic curve F_m is, as a map into a 4-manifold, determined by the normal bundle N_m of F_m . In cases of $F_1 = F$ and F_2 , the almost complex structure J_α on X is generic in the sense that the linearized operator L_f (see (1.7) below) is invertible and hence the contribution of f is $(-1)^{h^0(N_m)} / |\text{Aut}(f)|$ (cf. Section 10 of [LP]). When $m \geq 3$, J_α is, in general, no longer generic. We need to perturb J_α to generic J . In Section 2, using the universal property of local invariants (see (1.3) below), we choose a local model that is convenient for our calculation. In Section 3, when L_f is not invertible, we use a lifting property of covering space to calculate the contribution of f that proves the Main Theorem. The information for dimension zero GW invariants of elliptic surfaces with $p_g > 0$ is the same as for its Seiberg-Witten invariants. We spell out the specific connection in Remark 3.4.

Acknowledgments. I am very grateful to Thomas H. Parker for helpful discussions.

1 Dimension Zero Genus One Local GW Invariants

Let X be a (not necessarily compact) elliptic Kähler surface with a holomorphic 2-form α . The 2-form α defines an almost complex structure J_α on X by the formula (0.1). Suppose that the zero divisor D_α of α has a smooth reduction $D_\alpha = (m-1)D$ where D is a regular fiber or a multiple fiber of multiplicity m for some integer $m > 1$. The adjunction formula then shows $c_1(TX)([D]) = 0$ and $c_1(N) = 0$ where N is the normal bundle of D . The moduli space

$$\overline{\mathcal{M}}_1^\alpha(X, d[D]) \tag{1.1}$$

of stable J_α -holomorphic maps from curves of genus one representing the class $d[D]$ ($d \neq 0$) carries a (virtual) fundamental class

$$[\overline{\mathcal{M}}_1^\alpha(X, d[D])]^{vir} \tag{1.2}$$

that is defined by the GW theory of 4-manifolds (cf. Section 4 of [LP]). This (virtual) fundamental class (1.2) has dimension zero since $c_1(TX)([D]) = 0$. *The dimension zero genus one local Gromov-Witten invariant of X associated with the zero divisor D_α* is then

$$GW_1^{loc}(X, D_\alpha, d) := [\overline{\mathcal{M}}_1^\alpha(X, d[D])]^{vir}.$$

This local GW invariant has the following universal property. Let X' be another elliptic Kähler surface with a holomorphic 2-form α' whose zero divisor $D_{\alpha'} = (m'-1)D'$ where D' is a regular fiber or a multiple fiber of multiplicity m' . Let N' be the normal bundle of D' . If $m = m'$ and $h^0(N) = h^0(N')$ then

$$GW_1^{loc}(X, D_\alpha, d) = GW_1^{loc}(X', D_{\alpha'}, d) \tag{1.3}$$

(cf. Section 6 of [LP]). We set

$$GW_1^{loc}(X, D_\alpha, d) = \begin{cases} GW_1^{loc}((m-1)F, d) & \text{if } D \text{ is a regular fiber} \\ GW_1^{loc}(F_m, d) & \text{if } D \text{ is a } m\text{-multiple fiber} \end{cases} \tag{1.4}$$

It was proved in Example 4.4 of [LP] that

$$GW_1^{loc}(mF, d) = m GW_1^{loc}(F, d). \tag{1.5}$$

As given in (0.2), all dimension zero GW invariants of minimal elliptic surfaces with $p_g > 0$ are sums of local invariants in (1.4).

In the below, we will give a precise description on the (virtual) fundamental class (1.2) which will be used for our calculation in Section 3. The point in the moduli space (1.1) is an equivalence class $[f, C]$ of stable maps (f, C) where two stable maps (f, C) and (f', C') are equivalent if there is a biholomorphic map $\sigma : C \rightarrow C'$ with $f' \circ \sigma = f$. By Lemma 0.1, if $d \neq 0$ every representative (f, C) of $[f, C]$ is a holomorphic d -fold covering map from C to D . Thus, if D is given by a lattice Λ in the complex plane then $[f, C]$ is determined by an index d sublattice of Λ . In particular, the moduli space (1.1) consists of $\sigma(d)$ points.

On the other hand, since the (virtual) fundamental class (1.2) is defined by the GW theory of 4-manifolds, as described in Section 3 of [IP], it is a finite sum

$$[\overline{\mathcal{M}}_1^\alpha(X, d[D])]^{vir} = \sum c([f, C]) \quad (1.6)$$

over $[f, C] \in \overline{\mathcal{M}}_1^\alpha(X, d[D])$ of the contributions $c([f, C])$ that are defined as follows. Choose a $p \in D$ and a small disk B in X with $B \cap D = \{p\}$ and, once and for all, fix a map (f, C, x) with $f(x) = p$ such that (f, C) represents $[f, C]$. Then for a generic almost complex structure J on X that is sufficiently close to J_α and tamed by the Kähler form on X , there are finitely many J -holomorphic maps (f_i, C_i, x_i) from smooth genus one curves with one marked point such that (i) $f_i(x_i) \in B$ (ii) each (f_i, C_i, x_i) is C^0 -close to (f, C, x) (in a suitable space of maps) and (iii) the index zero operator

$$L_{f_i} : \Omega^0(f_i^* N_i) \rightarrow \Omega^{0,1}(f_i^* N_i) \quad (1.7)$$

has trivial kernel (or equivalently L_{f_i} is invertible) where the operator L_{f_i} is obtained by linearizing J -holomorphic map equation (see Remark 1.1 below) and restricting to the normal bundle N_i of the image of f_i . Denote by

$$\mathcal{M}_{(f, C, x), B, J}$$

the set of such J -holomorphic maps (f_i, C_i, x_i) . Notice that for each (f_i, C_i, x_i) the preimage $f_i^{-1}(B)$ consists of $d = |\text{Aut}(f)|$ distinct points x_{ij} . Since the automorphism group of C_i acts transitively, for each x_{ij} there exists an automorphism σ_j of C_i with $\sigma_j(x_i) = x_{ij}$ such that $(f_i \circ \sigma_j, C_i, x_i)$ is also contained in the set $\mathcal{M}_{(f, C, x), B, J}$. The contribution $c([f, C])$ is thus the (weighted) sum

$$c([f, C]) = \frac{1}{d} \sum (-1)^{SF(L_{f_i})}$$

over f_i in $\mathcal{M}_{(f, C, x), B, J}$ where the sign of each f_i is given by the mod 2 spectral flow $SF(L_{f_i})$ of the invertible operator L_{f_i} . In particular, $SF(L_{f_i}) = 0$ if L_{f_i} is complex linear, namely J -linear.

Remark 1.1. The operator $D_{f_i} : \Omega^0(f_i^* TX) \rightarrow \Omega^{0,1}(f_i^* TX)$ obtained by linearizing J -holomorphic map equation at f_i is given by

$$D_{f_i}(\xi)(v) = \nabla_v \xi + J \nabla_{jv} \xi + \frac{1}{2} [(\nabla_\xi J)(df_i(jv)) - J(\nabla_\xi J)(v)] \quad (1.8)$$

where $\xi \in \Omega^0(f_i^* TX)$, $v \in TC_i$ and j is the complex structure on C_i . Here ∇ is the pull-back connection on $f_i^* TX$ of the Levi-Civita connection of the metric on X that is defined by the Kähler form and J (cf. Lemma 6.3 of [RT]).

2 Local Model

Once and for all, fix an integer $m \geq 2$ and let D denote the elliptic curve given as the complex plane (with coordinate z) modulo the lattice $\mathbb{Z} + (mi)\mathbb{Z}$. Then $S = D \times \mathbb{C}$ has an automorphism φ of order m defined by

$$\varphi(z, w) = (z + i, e^{2\pi i/m} \cdot w)$$

such that all powers φ^i are fixed-point free where w is a coordinate on \mathbb{C} . Let S_m be the quotient of S by the group $\{\varphi^i\}$ and $q : S \rightarrow S_m$ the quotient map. The map $S \rightarrow \mathbb{C} : (z, w) \rightarrow w^m$ then factors through S_m to give an elliptic fibration $S_m \rightarrow \mathbb{C}$ whose central fiber is a m -multiple fiber D_m given by the lattice $\mathbb{Z} + i\mathbb{Z}$ with torsion normal bundle N_m of order m :

$$\begin{array}{ccc} S = D \times \mathbb{C} & \xrightarrow{q} & S_m \\ & \searrow & \swarrow \\ & \mathbb{C} & \end{array}$$

$(z, w) \rightarrow w^m$

The following simple observation is a key fact for our subsequent discussions. Let $f : C \rightarrow D_m$ be a holomorphic map of degree d from an elliptic curve C that is given by a sublattice of $\mathbb{Z} + i\mathbb{Z}$ of the form

$$a\mathbb{Z} + (bi + k)\mathbb{Z} \quad \text{with} \quad d = ab, \quad 0 \leq k \leq a - 1.$$

Write $D \times \{0\} \subset S$ simply as D .

Lemma 2.1. *Let D , N_m and $f : C \rightarrow D_m$ be as above. Then,*

$$f \text{ factors through } D \iff a \mid \frac{d}{m} \iff f^*N_m = \mathcal{O}_C$$

Proof. f factors through $D \iff a\mathbb{Z} + (bi + k)\mathbb{Z}$ is a sublattice of $\mathbb{Z} + (mi)\mathbb{Z} \iff m \mid b \iff a \mid \frac{d}{m}$. This shows the first assertion. Observe that for the restriction map $g_m = q|_D : D \rightarrow D_m$,

$$g_m^*(N_m) = g_m^*([D_m]|_{D_m}) = q^*([D_m])|_D = [q^*D_m]|_D = [D]|_D = \mathcal{O}_D \quad (2.1)$$

where $[D_m]$ is the line bundle associated to the divisor D_m , $N_m = [D_m]|_{D_m}$ by adjunction, the pullback divisor $q^*D_m = D$ and again by adjunction $[D]|_D$ is the normal bundle of D that is trivial. Write as $N_m = \mathcal{O}_{D_m}(p - q)$ where

$$\int_q^p dz = \frac{k_1}{m} + i \frac{k_2}{m} \quad \text{for some} \quad 0 \leq k_1, k_2 \leq m - 1.$$

Then, by (2.1) and the Abel's Theorem, $g_m^*N_m = \mathcal{O}_D(\sum_j(p_j - q_j))$ for some p_j, q_j such that

$$\sum_j \int_{q_j}^{p_j} dz = k_1 + ik_2 \equiv 0 \pmod{\mathbb{Z} + (mi)\mathbb{Z}}.$$

Consequently, $k_2 = 0$ and $\gcd(m, k_1) = 1$ since N_m is torsion of order m . Now, again by the Abel's Theorem, $f^*N_m = \mathcal{O}_C(\sum_\ell(t_\ell - s_\ell))$ for some s_ℓ, t_ℓ such that

$$\sum_\ell \int_{s_\ell}^{t_\ell} dz = \frac{dk_1}{m} \equiv 0 \pmod{a\mathbb{Z} + (bi + k)\mathbb{Z}} \iff f^*N_m = \mathcal{O}_C$$

Therefore, $a \mid \frac{d}{m} \iff m \mid b \iff a \mid \frac{dk_1}{m} \iff f^*N_m = \mathcal{O}_C$. This shows the second assertion. \square

Remark 2.2. Since $q : S \rightarrow S_m$ is a covering map, Lemma 2.1 shows $f : C \rightarrow D_m \subset S_m$ lifts to $\tilde{f} : C \rightarrow D \subset S$ if and only if $f^*N_m = \mathcal{O}_C$. On the other hand, the Kähler form $-\frac{i}{2}(dz \wedge d\bar{z} + dw \wedge d\bar{w})$ on \mathbb{C}^2 descends to a Kähler form $\tilde{\omega}$ on S that is φ -invariant, so $\tilde{\omega}$ also descends to a Kähler form ω on S_m such that $q^*\omega = \tilde{\omega}$.

3 Calculation

Let $q : (S, D) \rightarrow (S_m, D_m)$ be as in Section 2. Fix a holomorphic 2-form

$$\alpha = w^{m-1} dw \wedge dz$$

on S whose zero divisor is $(m-1)D$ and let J_α denote the almost complex structure on S defined by the formula (0.1). The 2-form α is φ -invariant, so it descends to a holomorphic 2-form α_m on S_m whose zero divisor is $(m-1)D_m$. We denote by $J_m = J_{\alpha_m}$ the almost complex structure on S_m defined by the 2-form α_m . Since D_m is a multiple fiber of multiplicity m ,

$$GW_1^{loc}(F_m, d) = [\overline{\mathcal{M}}_1^{\alpha_m}(S_m, d[D_m])]^{vir}$$

where the right-hand side is given by the sum of contributions as in (1.6). In order to calculate them, we decompose the moduli space $\overline{\mathcal{M}}_1^{\alpha_m}(S_m, d[D_m])$ as a disjoint union

$$\overline{\mathcal{M}}_1^{\alpha_m}(S_m, d[D_m]) = \mathcal{M}_{m,d}^+ \amalg \mathcal{M}_{m,d}^- \quad \text{where} \quad \begin{cases} [f, C] \in \mathcal{M}_{m,d}^+ & \text{if } h^0(f^*N_m) = 0 \\ [f, C] \in \mathcal{M}_{m,d}^- & \text{if } h^0(f^*N_m) = 1 \end{cases}$$

It then follows from Lemma 2.1 that

$$\# \mathcal{M}_{m,d}^+ = \sigma(d) - \sigma\left(\frac{d}{m}\right) \quad \text{and} \quad \# \mathcal{M}_{m,d}^- = \sigma\left(\frac{d}{m}\right) \quad (3.1)$$

where $\#A$ is the cardinality of a set A .

We first calculate the contribution $c([f, C])$ of $[f, C]$ in $\mathcal{M}_{m,d}^+$. In the below, we always assume $m \geq 3$ and $m|d$.

Lemma 3.1. *If $[f, C] \in \mathcal{M}_{m,d}^+$ then $c([f, C]) = \frac{1}{d}$.*

Proof. The linearized operator L_f has the form $L_f = \overline{\partial}_f + R_m$ where $\overline{\partial}_f$ is the usual $\overline{\partial}$ -operator on f^*N_m and the zeroth order term R_m is given by

$$R_m(\xi) = -\nabla_\xi K_{\alpha_m} \circ J_{\alpha_m} \circ df \quad \text{for} \quad \xi \in \Omega^0(f^*N_m)$$

(cf. Section 8 of [LP]). But, $R_m \equiv 0$ since α_m (and hence K_{α_m}) vanishes of order $m-1 \geq 2$ along D_m . Consequently, $\dim \ker L_f = 2h^0(f^*N_m) = 0$, so L_f is invertible with $SF(L_f) = 0$. Now, the proof follows from the fact $f : C \rightarrow D_m$ has degree d . \square

Let $[f, C] \in \mathcal{M}_{m,d}^-$. The proof of Lemma 3.1 shows $L_f = \overline{\partial}_f$ is not invertible. In this case, we will use the m -fold covering map $q : S \rightarrow S_m$ to calculate the contribution $c([f, C])$. Observe that by Lemma 2.1 the map

$$\mathcal{M}_{m,d}^- \rightarrow \overline{\mathcal{M}}_1^\alpha(S, \frac{d}{m}[D]) \quad \text{defined by} \quad [f, C] \rightarrow [\tilde{f}, C]$$

is one-to-one and onto where \tilde{f} is a lift of f .

Lemma 3.2. *If $[f, C] \in \mathcal{M}_{m,d}^-$ then $c([f, C]) = \frac{1}{m} c([\tilde{f}, C])$.*

Proof. Let $B = \{0\} \times \Delta \subset S$ where Δ is a small disk around 0 in \mathbb{C} and $B_m = q(B)$ and fix a map (f, C, x) with $f(x) \in B_m$ such that (f, C) represents $[f, C]$. Since the restriction map $q|_B : B \rightarrow B_m$ is one-to-one, Lemma 2.1 shows that (f, C, x) uniquely lifts to a J_α -holomorphic map (\tilde{f}, C, x) with $\tilde{f}(x) \in B$ such that (\tilde{f}, C) represents $[\tilde{f}, C]$ in $\overline{\mathcal{M}}_1^\alpha(S, \frac{d}{m}[D])$.

Let ω and $\tilde{\omega}$ be the Kähler forms as in Remark 2.2 and choose a generic ω -tamed almost complex structure J on S_m that is close to J_m . Then, we have

- J lifts to an $\tilde{\omega}$ -tamed almost complex structure \tilde{J} on S close to J_α such that $dq \circ \tilde{J} = J \circ dq$,
- each f_i in $\mathcal{M}_{(f, C, x), B_m, J}$ is homotopic to f since f_i is C^0 -close to f , so (f_i, C_i, x_i) also uniquely lifts to \tilde{J} -holomorphic maps (\tilde{f}_i, C_i, x_i) with $\tilde{f}_i(x_i) \in B$ such that (\tilde{f}_i, C_i, x_i) is C^0 -close to (\tilde{f}, C, x) .

The pair (ω, J) defines a metric g on S_m whose lift $\tilde{g} = q^*g$ is the same metric defined by the pair $(\tilde{\omega}, \tilde{J})$. Let ∇ and $\tilde{\nabla}$ respectively denote the pull-back connections on $f_i^*TS_m$ and \tilde{f}_i^*TS of the Levi-Civita connection of g and \tilde{g} . The differential dq then induces a bundle isomorphism $dq : \tilde{f}_i^*TS \rightarrow \tilde{f}_i^*q^*TS_m = f_i^*TS_m$ such that $dq \circ \tilde{\nabla} = \nabla \circ dq$ (see [W] page 138) and hence by the formula (1.8) we have

$$dq \circ D_{\tilde{f}_i} = D_{f_i} \circ dq \quad (3.2)$$

The differential dq also induces a bundle isomorphism $dq_i : \tilde{f}_i^*\tilde{N}_i \rightarrow f_i^*N_i$ and restricting the equation (3.2) to $\tilde{f}_i^*\tilde{N}_i$ and $f_i^*N_i$ gives

$$dq_i \circ L_{\tilde{f}_i} = L_{f_i} \circ dq_i$$

where \tilde{N}_i and N_i are normal bundles of $\text{Im}(\tilde{f}_i)$ and $\text{Im}(f_i)$ respectively. Therefore, $L_{\tilde{f}_i}$ is also invertible and hence there is one-to-one correspondence

$$\mathcal{M}_{(f, C, x), B_m, J} \rightarrow \mathcal{M}_{(\tilde{f}, C, x), B, \tilde{J}} \quad \text{given by} \quad (f_i, C_i, x_i) \rightarrow (\tilde{f}_i, C_i, x_i).$$

Let \tilde{L}_t be a path of first order elliptic operators from an invertible \tilde{J} -linear operator \tilde{L}_0 to $\tilde{L}_1 = L_{\tilde{f}_i}$ with all \tilde{L}_t invertible except at finitely many t_k . Then, $dq_i \circ \tilde{L}_t \circ (dq_i)^{-1}$ is also a path from invertible J -linear operator to L_{f_i} such that

$$SF(L_{\tilde{f}_i}) \equiv \sum_k \dim \ker \tilde{L}_{t_k} = \sum_k \dim \ker dq_i \circ \tilde{L}_{t_k} \circ (dq_i)^{-1} \equiv SF(L_{f_i}) \pmod{2}.$$

Now, noting $\deg(f) = d$ and $\deg(\tilde{f}) = \frac{d}{m}$, we have

$$c([f, C]) = \frac{1}{d} \sum_{f_i} (-1)^{SF(L_{f_i})} = \frac{1}{d} \sum_{\tilde{f}_i} (-1)^{SF(L_{\tilde{f}_i})} = \frac{1}{m} c([\tilde{f}, C]). \quad \square$$

We are now ready to prove the Main Theorem in the introduction.

Proof of the Main Theorem : It follows from Lemma 3.1, Lemma 3.2 and (3.1) that

$$\begin{aligned} GW_1^{loc}(F_m, d) &= \sum_{[f, C] \in \mathcal{M}_{m, d}^+} c([f, C]) + \sum_{[f, C] \in \mathcal{M}_{m, d}^-} c([f, C]) \\ &= \frac{1}{d} \left(\sigma(d) - \sigma\left(\frac{d}{m}\right) \right) + \frac{1}{m} [\overline{\mathcal{M}}_1^\alpha(S, \frac{d}{m}[D])]^{vir}. \end{aligned} \quad (3.3)$$

Since the 2-form α on S has the zero divisor $(m-1)D$, so by (1.5) and (0.3) we have

$$[\overline{\mathcal{M}}_1^\alpha(S, \frac{d}{m}[D])]^{vir} = GW_1^{loc}((m-1)F, \frac{d}{m}) = -(m-1) \frac{m}{d} \sigma\left(\frac{d}{m}\right). \quad (3.4)$$

Now, the proof follows from (3.3) and (3.4). \square

Remark 3.3. One can also use the above argument to compute $GW_1^{loc}(F_2, d)$, replacing the ‘‘Taubes type’’ argument used in [LP]. Specifically, for each $f \in \overline{\mathcal{M}}_1^{\alpha_2}(S_2, d[D_2])$ the linearized operator L_f is invertible with $SF(L_f) \equiv h^0(f^*N_2) \pmod{2}$ (cf. Proposition 9.2 of [LP]). Thus, by (3.1) we have

$$GW_1^{loc}(F_2, d) = [\overline{\mathcal{M}}_1^{\alpha_2}(S_2, d[D_2])]^{vir} = \frac{1}{d} \left(\sigma(d) - \sigma\left(\frac{d}{2}\right) \right) - \frac{1}{d} \sigma\left(\frac{d}{2}\right) = \frac{1}{d} \left(\sigma(d) - 2\sigma\left(\frac{d}{2}\right) \right).$$

Remark 3.4. Ionel and Parker [IP] showed how GW invariants for the class A of a symplectic 4-manifold X are related with the Taubes’ Gromov invariants $Gr_X(A)$ [T] that count embedded (not necessarily connected) curves in X representing the class A . They used a particular function $F(t)$ that satisfies

$$\prod_d F(t^d)^{-\frac{1}{d}\sigma(d)} = (1-t)$$

to relate Taubes’ counting of multiple covers of embedded tori with the dimension zero genus one GW invariants. Let X be a minimal elliptic surface with $p_g > 0$. In this case, any GW invariant constrained to pass through generic points vanishes (cf. Corollary 3.4 of [LP]). So, by (0.2), (0.3) and the Main Theorem, the relation between two set of invariants (Theorem 4.5 of [IP]) yields

$$\begin{aligned} \sum_A Gr_X(A) t_A &= \prod_{d,k} F(t^d)^{c_\pi GW_1^{loc}(F,d)} F(t_{m_k}^d)^{GW_1^{loc}(F_{m_k},d)} \\ &= (1-t)^{c_\pi} \prod_k (1 + t_{m_k} + \dots + t_{m_k}^{m_k-1}). \end{aligned}$$

This also gives the well-known Seiberg-Witten invariants SW of X (cf. [FM], [B], [FS]) due to the famous Taubes’ theorem $SW = Gr$.

References

- [B] R. Brussee, *The canonical class and the C^∞ properties of Kähler surfaces*, New York J. Math. **2** (1996), 103–146.
- [BHPV] W. Barth, K. Hulek, C. Peters, and A. Van de Ven, *Compact complex surfaces*, second ed., Springer-Verlag, Berlin Heidelberg, 2004.
- [FM] R. Friedman and J. Morgan, *Obstruction bundles, semiregularity, and Seiberg-Witten invariants*, Comm. Anal. Geom. **7** (1999), no. 3, 451–495.

- [FS] R. Fintushel and R. Stern, *Rational blowdowns of smooth 4-manifolds*, J. Diff. Geom. **46** (1997), 181-235.
- [IP] E. Ionel and T.H. Parker, *The Gromov invariants of Ruan-Tian and Taubes*, Math. Res. Lett. **4** (1997), no. 4, 521-532.
- [KL] Y-H. Kiem and J. Li, *Low degree GW invariants of spin surfaces*, Pure Appl. Math. Q. **7** (2011), no. 4, 1449-1476.
- [L] J. Lee, *Family Gromov-Witten Invariants for Kähler Surfaces*, Duke Math. J. **123** (2004), no. 1, 209-233.
- [LP] J. Lee and T.H. Parker, *A Structure Theorem for the Gromov-Witten Invariants of Kähler Surfaces*, J. Diff. Geom. **77** (2007), 483-513.
- [RT] Y. Ruan and G. Tian, *A mathematical theory of cohomology*, J. Differential Geom. **42** (1995), 259-367.
- [T] C. Taubes, *Seiberg Witten and Gromov invariants for symplectic 4-manifolds*, Edited by Richard Wentworth. First International Press Lecture Series, 2. International Press, Somerville, MA, 2000.
- [W] G. Walschap, *Metric structures in differential geometry*, Graduate Texts in Mathematics, 224. Springer-Verlag, New York, 2004.