

THE BRAIDINGS IN THE MAPPING CLASS GROUPS OF SURFACES

YONGJIN SONG

ABSTRACT. The disjoint union of mapping class groups of surfaces forms a braided monoidal category \mathcal{M} , as the disjoint union of the braid groups \mathcal{B} does. We give a concrete, and geometric meaning of the braiding $\beta_{r,s}$ in \mathcal{M} . Moreover, we find a set of elements in the mapping class groups which correspond to the standard generators of the braid groups. Using this, we obtain an obvious map $\phi : B_g \rightarrow \Gamma_{g,1}$. We show that this map ϕ is injective and nongeometric in the sense of Wajnryb. Since this map extends to a braided monoidal functor $\Phi : \mathcal{B} \rightarrow \mathcal{M}$, the integral homology homomorphism induced by ϕ is trivial in the stable range.

1. INTRODUCTION

Let $\Gamma_{g,1}$ be the mapping class group of the surface $S_{g,1}$, the compact orientable surface of genus g with one boundary component obtained by deleting a disk from the closed surface S_g . Let $\mathcal{M} = \coprod_{g \geq 0} \Gamma_{g,1}$ be the disjoint union of $\Gamma_{g,1}$'s, then it is regarded as the category whose objects are nonnegative integers. Here, $\text{hom}_{\mathcal{M}}(g, g) = \Gamma_{g,1}$ and there is no morphism between distinct integers. The monoid structure, called F-product in \mathcal{M} , is induced by the pair-of-pants connected sum. The braid structure of \mathcal{M} was given in [8] in terms of the actions of braidings on the fundamental group of $S_{g,1}$. Recall that $\Gamma_{g,1}$ may be identified with the subgroup of the automorphism group of $\pi_1 S_{g,1}$ that consists of the automorphisms fixing the fundamental relator $R = [y_1, x_1] \cdots [y_g, x_g]$. Here $\pi_1 S_{g,1}$ is a free group on $2g$ generators $x_1, y_1, x_2, y_2, \dots, x_g, y_g$ and the fundamental relator represents a loop along the boundary of $S_{g,1}$. The F-product on \mathcal{M} may be identified with the operation taking the free product of automorphisms.

The action of the braiding $\beta_{r,s} : r \otimes s \rightarrow s \otimes r$, which is an element of $\Gamma_{r+s,1}$, on the free group on $x_1, y_1, x_2, y_2, \dots, x_{r+s}, y_{r+s}$ was given in [8]. The braiding could be also expressed as a product of the standard Dehn twists, the generators of $\Gamma_{g,1}$.

The disjoint union of braid groups $\mathcal{B} = \coprod_{g \geq 0} B_g$ forms a braided monoidal category. Let $\sigma_1, \dots, \sigma_{g-1}$ be the standard generators of B_g , where σ_i crosses the i -th and $i+1$ -st strings. The monoid structure is induced by the juxtaposition and the braiding $(\sigma_{r,s} : r \otimes s \rightarrow s \otimes r) \in B_{r+s}$ is the crossing of the front r strings and the rear s strings. We can express $\sigma_{r,s}$ in terms of σ_i 's as follows:

$$\sigma_{r,s} = (\sigma_r \sigma_{r-1} \cdots \sigma_1)(\sigma_{r+1} \sigma_r \cdots \sigma_2) \cdots (\sigma_{r+s-1} \sigma_{r+s-2} \cdots \sigma_s).$$

In this paper, we give a concrete, and geometric meaning of the braiding $\beta_{r,s}$ in the braided monoidal category \mathcal{M} . Moreover, we find a set of elements $\beta_1, \dots, \beta_{g-1}$

2010 *Mathematics Subject Classification.* Primary , 55R37, 18D10; Secondary 57M50, , 55P48.

Key words and phrases. braid group, mapping class group, Dehn twists, braided monoidal category, double loop space, plus construction.

of $\Gamma_{g,1}$ which are analogous to the standard generators of B_g . Using these generators we can define an obvious map $\phi : B_g \rightarrow \Gamma_{g,1}, \sigma_i \mapsto \beta_i$. An embedding of a braid group into a mapping class group is said to be *geometric* by Wajnryb ([14], [15]) if it maps the standard generators of the braid group to Dehn twists. We show that the map $\phi : B_g \rightarrow \Gamma_{g,1}$ is injective and is, moreover, *nongeometric*.

The natural embedding $\phi : B_g \rightarrow \Gamma_{g,1}$ can be extended to a braided monoidal functor $\Phi : \mathcal{B} \rightarrow \mathcal{M}$. Namely, Φ preserves the braidings. As a consequence we show that $\phi : B_g \rightarrow \Gamma_{g,1}$ induces a map of double loop spaces $B\phi^+ : BB_\infty^+ \rightarrow B\Gamma_\infty^+$ and hence the homology homomorphism $\phi_* : H_*(B_\infty; \mathbb{Z}) \rightarrow H_*(\Gamma_\infty; \mathbb{Z})$ induced by ϕ is trivial, where $B_\infty = \varinjlim B_g$ and $\Gamma_\infty = \varinjlim \Gamma_{g,1}$.

The classifying space of a braided monoidal category naturally gives rise to a double loop space ([5]). More precisely speaking, the group completion of the classifying space of a braided monoidal category is homotopy equivalent to a double loop space. For $\mathcal{B} = \coprod_{g \geq 0} B_g$, we have

$$\overline{B\mathcal{B}} = \Omega B\left(\coprod_{g \geq 0} BB_g\right) \simeq \mathbb{Z}_+ \times BB_\infty^+ \simeq \Omega^2 S^2,$$

where $\overline{B\mathcal{B}}$ means the group completion of the classifying space of \mathcal{B} , and $+$ means the Quillen's plus construction.

The braided monoidal functor $\Phi : \mathcal{B} \rightarrow \mathcal{M}$ induces a map of double loop spaces $\overline{B\mathcal{B}} \rightarrow \overline{B\mathcal{M}}$. In other words, $B\phi^+ : BB_\infty^+ \rightarrow B\Gamma_\infty^+$ is a map of double loop spaces. Since every map $\Psi : BB_\infty^+ \rightarrow B\Gamma_\infty^+$ of double loop spaces is null-homotopic ([10], Lemma 5.3), so is $B\phi^+$. This proves that the homomorphism $\phi_* : H_*(B_\infty; \mathbb{Z}) \rightarrow H_*(\Gamma_\infty; \mathbb{Z})$ is trivial, since the plus construction does not change the homology.


When we first think about the geometric meaning of the braiding $\beta_{r,s}$, since it interchanges the front r genus holes and rear s genus holes of $S_{r+s,1}$, we think that it would be a 180° rotation around the boundary of the surface. Here we consider the boundary as the waist of the surface with two arms; the left arm is the front r genus holes and the right arm is the rear s genus holes. However, just a 180° rotation around the waist is not sufficient. We also need half-twists of two arms. Here, it is important to determine the directions of these half-twists. There are four choices of directions; two choices on the waist and two choices on the arms. By analyzing the simplest case, $\beta_{1,1}$ for $S_{2,1}$, we show that the braiding $\beta_{r,s}$ is the composite of two half-twists; the *reverse* half Dehn twist along the boundary and half Dehn twists along the shoulders of two arms. The key part of this paper is to figure out a correct geometric meaning of $\beta_{1,1}$ as a self-homeomorphism of $S_{2,1}$ and extend this to the general case, $\beta_{r,s}$ in $\Gamma_{r+s,1}$.

2. BRAIDED MONOIDAL CATEGORY

Let $S_{g,1}$ be a compact connected orientable surface of genus g with one boundary component. The mapping class group $\Gamma_{g,1}$ is the group of isotopy classes of orientation preserving self-homeomorphisms of $S_{g,1}$ fixing the boundary of $S_{g,1}$ pointwise. $\Gamma_{g,1}$ is generated by the standard Dehn twists $a_1, \dots, a_g, b_1, \dots, b_g, w_1, \dots, w_{g-1}$ (Figure 1).

Definition 2.1. A *braided monoidal category* is a monoidal category $(\mathcal{C}, \otimes, I)$ together with a family of natural commutativity isomorphisms

$$\beta_{A,B} : A \otimes B \rightarrow B \otimes A$$

FIGURE 1. Standard Dehn twists of $\Gamma_{g,1}$

called *braidings*, natural in both variables, such that for all objects A, B, C in \mathcal{C} ,

- (a) $\beta_{A,I} = \beta_{I,A} = 1_A$
- (b) $\beta_{A \otimes B, C} = (\beta_{A,C} \otimes 1_B) \circ (1_A \circ \beta_{B,C})$
- (c) $\beta_{A, B \otimes C} = (1_B \otimes \beta_{A,C}) \circ (\beta_{A,B} \otimes 1_C)$.

The equations in (b), (c) imply the Yang-Baxter equation:

$$(1_C \otimes \beta_{A,B})(\beta_{A,C} \otimes 1_B)(1_A \otimes \beta_{B,C}) = (\beta_{B,C} \otimes 1_A)(1_B \otimes \beta_{A,C})(\beta_{A,C} \otimes 1_C).$$

The disjoint union $\mathcal{M} = \coprod_{g \geq 0} \Gamma_{g,1}$ forms a category whose objects are nonnegative integers and morphisms satisfy

$$\text{hom}(g, h) = \begin{cases} \Gamma_{g,1} & \text{if } g = h, \\ \phi & \text{if } g \neq h. \end{cases}$$

\mathcal{M} has a monoid structure: the F-product

$$\Gamma_{g,1} \times \Gamma_{h,1} \rightarrow \Gamma_{g+h,1}$$

is induced by extending two self-homeomorphisms on $S_{g,1}$ and $S_{h,1}$ to the surface $S_{g+h,1}$ obtained attaching a pair of pants (a sphere with three boundary components) to the surfaces $S_{g,1}$ and $S_{h,1}$ along the fixed boundary circles. We extend the identity map on the boundary to the whole pants.

It was shown in [8] that \mathcal{M} is a braided monoidal category whose braidings are as follows: Let $x_1, y_1, x_2, y_2, \dots, x_g, y_g$ be the generators of $\pi_1 S_{g,1}$ which are represented by loops parallel to the Dehn twist loops $a_1, b_1, a_2, b_2, \dots, a_g, b_g$, respectively, in Figure 1. $\Gamma_{g,1}$ is identified with the subgroup of the automorphism group of $\pi_1 S_{g,1}$ fixing the fundamental relator $R = [y_1, x_1][y_2, x_2] \cdots [y_g, x_g]$. The (r, s) -braiding $\beta_{r,s} : r \otimes s \rightarrow s \otimes r$, which is an element of $\Gamma_{r+s,1}$, acts on the free

group on $\{x_1, y_1, x_2, y_2, \dots, x_g, y_g\}$ as follows:

$$\begin{aligned}
 x_1 &\mapsto R_s x_{s+1} R_s^{-1}, & y_1 &\mapsto R_s y_{s+1} R_s^{-1} \\
 &\vdots & & \\
 x_r &\mapsto R_s x_{s+r} R_s^{-1}, & y_1 &\mapsto R_s y_{s+r} R_s^{-1} \\
 x_{r+1} &\mapsto x_1, & y_{r+1} &\mapsto y_1 \\
 x_{r+2} &\mapsto x_2, & y_{r+1} &\mapsto y_2 \\
 &\vdots & & \\
 x_{r+s} &\mapsto x_s, & y_{r+s} &\mapsto y_s
 \end{aligned}$$

where $R_s = [y_1, x_1] \cdots [y_s, x_s]$. We can easily check that the (r, s) -braidings fix the fundamental relator R and satisfy the braid equations (b), (c) of Definition 2.1.




FIGURE 2. The generators $x_1, y_1, \dots, x_g, y_g$ of $\pi_1 S_{g,1}$

Since the group completion of the classifying space of a braided monoidal category is homotopy equivalent to a double loop space, we have

Theorem 2.2. $\overline{BM} = \Omega B(\coprod_{g \geq 0} B\Gamma_{g,1})$ is homotopy equivalent to a double loop space. Here, \overline{BM} denotes the group completion of BM .

3. THE GEOMETRIC ANALYSIS OF BRAIDINGS

In this section we give geometric meanings of braidings in the mapping class group. By using this result we construct a natural embedding $\phi : B_g \hookrightarrow \Gamma_{g,1}$, which is nongeometric. We show that this embedding induces the trivial homology homomorphism in the stable range in the integral coefficient.

The braiding $\beta_{r,s} \in \Gamma_{r+s,1}$ can be expressed as a product of the standard Dehn twists. First, the braiding $\beta_{1,1}$ was expressed in [8] in terms of the Dehn twists a_1, a_2, b_1, b_2, w_1 :

Lemma 3.1. *The $(1,1)$ -braiding $\beta_{1,1}$ in $\Gamma_{2,1}$ equals*

$$(3.1) \quad \beta_{1,1} = (a_1 b_1 a_1)^4 (a_2 b_2 (a_1 b_1 a_1)^{-1} w_1 a_1 b_1 a_1^2 b_1)^{-3}.$$

The standard Dehn twists $a_1, \dots, a_g, b_1, \dots, b_g, w_1, \dots, w_{g-1}$ act on $\pi_1 S_{g,1}$, which is the free group generated by $\{x_1, y_1, x_2, y_2, \dots, x_g, y_g\}$, as follows:

$$\begin{aligned} a_i &: y_i \mapsto y_i x_i^{-1} \\ b_i &: x_i \mapsto x_i y_i \\ w_i &: x_i \mapsto z_i^{-1} y_{i+1} x_{i+1} y_{i+1}^{-1} \\ &: y_i \mapsto y_i z_i \\ &: y_{i+1} \mapsto z_i^{-1} y_{i+1} \end{aligned}$$

where $z_i = x_i^{-1} y_{i+1} x_{i+1} y_{i+1}^{-1}$ and these automorphisms fix the generators that do not appear in the list. By using these actions we can check that the equation given in Lemma 3.1 is correct, that is, $\beta_{1,1}$ acts on the free group on $\{x_1, y_1, x_2, y_2\}$ as

$$\begin{aligned} x_1 &\mapsto [y_1, x_1] x_2 [x_1, y_1], \quad y_1 \mapsto [y_1, x_1] y_2 [x_1, y_1] \\ x_2 &\mapsto x_1, \quad y_2 \mapsto y_1 \end{aligned}$$

as desired. Here the Dehn twists act on the right on $\pi_1 S_{g,1}$.

Now we construct the (r, s) -braiding $\beta_{r,s}$ by extending $\beta_{1,1}$. Let

$$\beta_i = (a_i b_i a_i)^4 (a_{i+1} b_{i+1} (a_i b_i a_i)^{-1} w_i a_i b_i a_i^2 b_i)^{-3}$$

for $i = 1, \dots, g-1$, then these satisfy the braid relations. The (r, s) -braiding $\beta_{r,s}$ in the group $\Gamma_{g,1}$ defined by

$$\beta_{r,s} = (\beta_r \beta_{r-1} \cdots \beta_1) (\beta_{r+1} \beta_r \cdots \beta_2) \cdots (\beta_{r+s-1} \beta_{r+s-2} \cdots \beta_s)$$

is analogous to the (r, s) -braiding in the braid group B_{r+s} . Here we may think that the composition of two braids is a stacking (from up to) down as morphisms map from top to bottom. Through simple calculations we can check that this (r, s) -braiding $\beta_{r,s}$ acts on $\pi_1 S_{g,1}$ as described in Section 2.

We now investigate the geometric meaning of $\beta_{r,s}$. We first figure out what $\beta_{1,1}$ really is as a self-homeomorphism of $S_{2,1}$.

For a simple closed curve C on $S_{g,1}$, the half Dehn twist h_C is a self-homeomorphism of $S_{g,1}$ which is 180° twist along a tubular neighborhood of C in the same direction as the usual Dehn twist, as given in Figure 3. Let h'_C denote the reverse half Dehn twist along C , namely $h'_C = h_C^{-1}$ in $\Gamma_{g,1}$.




FIGURE 3. The half Dehn twist along C

In this paper we sometimes abuse notations. For a simple closed curve C on the surface, it would be just a curve for a (half) Dehn twist, but sometimes it stands for an element of the fundamental group of the surface. For example, Let R be the closed curve along the boundary of $S_{g,1}$, then it could also mean the element $R = [y_1, x_1][y_2, x_2] \cdots [y_g, x_g] \in \pi_1 S_{g,1}$.

Let R_1, R_2 be closed curves on $S_{2,1}$ as given in Figure 4. The arrows on the curves mean the directions of them as elements of $\pi_1 S_{2,1}$.




FIGURE 4.

For the closed curve R on $S_{2,1}$, the half Dehn twist h_R and the reverse half Dehn twist h'_R are illustrated in Figure 5. We may think of them as 180° rotations of the whole surface around the axis l .




FIGURE 5. Two half Dehn twists along the boundary

We may regard h_R and h'_R as elements of the automorphism group of $\pi_1 S_{2,1} = F_{\{x_1, y_1, x_2, y_2\}}$. The geometric calculations in this section not only play key roles in analyzing the geometry of braidings in $\Gamma_{g,1}$, but also are of interest in their own right.

Proposition 3.2. *The half Dehn twist h_R acts on $\pi_1 S_{2,1}$ as follows:*

$$\begin{aligned} x_1 &\mapsto R_2^{-1} x_2^{-1}, & y_1 &\mapsto y_2^{-1} R_2 \\ x_2 &\mapsto R^{-1} x_1^{-1} R_2, & y_2 &\mapsto R_2^{-1} y_1 R. \end{aligned}$$

Note that h_R maps R_1 to R_2 , and R_2 to $R_2^{-1} R$.

Proposition 3.3. *The reverse half Dehn twist h'_R acts on $\pi_1 S_{2,1}$ as follows:*

$$\begin{aligned} x_1 &\mapsto R_1 x_2^{-1} R^{-1} = (R_1 y_2) x_2^{-1} (R_1 y_2)^{-1} \\ y_1 &\mapsto R y_2^{-1} R_1^{-1} = (R_1 y_2 x_2) y_2^{-1} (R_1 y_2 x_2)^{-1} \\ x_2 &\mapsto y_1 x_1^{-1} y_1^{-1} \\ y_2 &\mapsto R_1 y_1^{-1} = (y_1 x_1) y_1^{-1} (y_1 x_1)^{-1} \end{aligned}$$

Note that h'_R maps R_1 to RR_1^{-1} , and R_2 to R_1 .

These lemmas are obtained through series of geometric calculations of actions of half Dehn twists on $\pi_1 S_{2,1} = F_{\{x_1, y_1, x_2, y_2\}}$.

Proposition 3.4. For the half Dehn twists h_{R_1} and h_{R_2} , let $h_A = h_{R_1} \circ h_{R_2}$, called the half Dehn twists on two arms. Then h_A acts on $\pi_1 S_{g,1}$ as follows:

$$\begin{aligned} x_1 &\mapsto R_1^{-1} x_1^{-1} = (x_1 y_1) x_1^{-1} (x_1 y_1)^{-1} \\ y_1 &\mapsto y_1^{-1} R_1 = x_1 y_1^{-1} x_1^{-1} \\ x_2 &\mapsto R_2^{-1} x_2^{-1} = (x_2 y_2) x_2^{-1} (x_2 y_2)^{-1} \\ y_2 &\mapsto y_2^{-1} R_2 = x_2 y_2^{-1} x_2^{-1} \end{aligned}$$

Proposition 3.5. For the reverse half Dehn twists on two arms $h'_A = h'_{R_1} \circ h'_{R_2}$ acts on $\pi_1 S_{g,1}$ as follows:

$$\begin{aligned} x_1 &\mapsto x_1^{-1} R_1^{-1} = y_1 x_1^{-1} y_1^{-1} \\ y_1 &\mapsto R_1 y_1^{-1} = (y_1 x_1) y_1^{-1} (y_1 x_1)^{-1} \\ x_2 &\mapsto x_2^{-1} R_2^{-1} = y_2 x_2^{-1} y_2^{-1} \\ y_2 &\mapsto R_2 y_2^{-1} = (y_2 x_2) y_2^{-1} (y_2 x_2)^{-1} \end{aligned}$$

In the combination of the two half Dehn twists, the half Dehn twist along the boundary and the half Dehn twists on two arms, there are four choices of directions. For the braiding $\beta_{1,1}$, the right choice of directions is h'_R and h_A .

Theorem 3.6. The $(1,1)$ -braiding $\beta_{1,1} \in \Gamma_{2,1}$ is the product (composite) of h'_R and h_A in $\Gamma_{2,1}$, that is, $\beta_{1,1} = h'_R \circ h_A$.

Proof. Since three closed curves R, R_1, R_2 are mutually disjoint, h_R, h_{R_1}, h_{R_2} are all commutative. We have

$$\begin{aligned} x_1 &\xrightarrow{h'_R} R_1 x_2 R^{-1} \xrightarrow{h_A} R_1 x_2 R_1^{-1} \\ y_1 &\xrightarrow{h'_R} R y_2^{-1} R_1^{-1} \xrightarrow{h_A} R_1 y_2 R_1^{-1} \\ x_2 &\xrightarrow{h'_R} y_1 x^{-1} y_1^{-1} \xrightarrow{h_A} x_1 \\ y_2 &\xrightarrow{h'_R} R_1 y_1^{-1} \xrightarrow{h_A} y_1 \end{aligned}$$

as desired. \square

We have shown that $\beta_{1,1}$ is the composition of the reverse half Dehn twist along the boundary and the half Dehn twists on two arms. It is illustrated in Figure 6 and 7.





FIGURE 6. The action of $\beta_{1,1}$

FIGURE 7. The action of $\beta_{1,1}$


Recall that the (r, s) -braiding $\beta_{r,s} \in \Gamma_{r+s,1}$ equals

$$(\beta_r \beta_{r-1} \cdots \beta_1)(\beta_{r+1} \beta_r \cdots \beta_2) \cdots (\beta_{r+s-1} \beta_{r+s-2} \cdots \beta_s),$$

where $\beta_i = (a_i b_i a_i)^4 (a_{i+1} b_{i+1} (a_i b_i a_i)^{-1} w_i a_i b_i a_i^2 b_i)^{-3}$. Let $R_i = [y_i, x_i]$. Then $\beta_i \in \Gamma_{r+s,1}$ acts on $\pi_1 S_{r+s,1}$ as follows:

$$\begin{aligned} \beta_i : \quad x_i &\mapsto R_i x_{i+1} R_i^{-1} & , \quad y_i &\mapsto R_i y_{i+1} R_i^{-1} \\ x_{i+1} &\mapsto x_i & , \quad y_{i+1} &\mapsto y_i. \end{aligned}$$

Let $R_{i,j} = [y_i, x_i][y_{i+1}, x_{i+1}] \cdots [y_j, x_j]$ for $j > i$. For a closed curve $R_{i,i+1}$ on $S_{r+s,1}$ (See Figure 8) imagine that we cut out the surface along $R_{i,i+1}$, then we get a (small) surface $S_{2,1}$.

FIGURE 8. The local surface $S_{2,1}$ obtained by cutting along $R_{i,i+1}$

Then β_i is a self-homeomorphism of $S_{r+s,1}$ which is the composition of the reverse half Dehn twist along $h'_{R_{i,i+1}}$ and the half Dehn twists on two arms, that is, we have

$$\beta_i = h'_{R_{i,i+1}} \circ h_{R_i} \circ h_{R_{i+1}}.$$

We can see, by checking the actions on the fundamental group, that these β_i 's satisfy the braid relations:

$$\beta_i \beta_{i+1} \beta_i = \beta_{i+1} \beta_i \beta_{i+1}$$

$$\beta_i \beta_j = \beta_j \beta_i \text{ for } |i - j| \geq 2.$$

The (r, s) -braiding $\beta_{r,s}$ is a composition of a series of local braidings β'_i 's. It may be regarded as the composition of the reverse half Dehn twist along the boundary R of $S_{r+s,1}$ and a series of some local half Dehn twists in the upper part of the whole surface. Figure 9 shows that how $\beta_{2,3}$ acts on y_1 .




FIGURE 9. An example of action of braiding

Let $\text{Aut } F_n$ be the automorphism of the free group F_n on n generators x_1, \dots, x_n . Then there is an injection $j : B_n \rightarrow \text{Aut } F_n$, called the Artin map ([1], [12]), defined by

$$j(\sigma_i) : x_i \mapsto x_i x_{i+1} x_i^{-1}, \quad x_{i+1} \mapsto x_i$$

where $\sigma_1, \dots, \sigma_{n-1}$ are the standard generators of B_n .

There exists an obvious group homomorphism

$$\begin{aligned} \phi : B_g &\rightarrow \Gamma_{g,1} \subset \text{Aut } S_{g,1} \\ \sigma_i &\mapsto \beta_i \end{aligned}$$

where $\phi(\sigma_i) = \beta_i$ acts on $\pi_1 S_{g,1} = F_{\{x_1, y_1, x_2, y_2, \dots, x_g, y_g\}}$ as

$$\begin{aligned} x_i &\mapsto R_i x_{i+1} R_i^{-1}, & y_i &\mapsto R_i y_{i+1} R_i^{-1} \\ x_{i+1} &\mapsto x_i, & y_{i+1} &\mapsto y_i, \end{aligned}$$

where $R_i = [y_i, x_i]$. The map ϕ is injective as the Artin map $j : B_n \rightarrow \text{Aut } F_n$ is ([1], [12])


An embedding of a braid group into a mapping class group is called a geometric embedding if it takes the standard generators of braid group onto Dehn twists in the mapping class group([14]). In [15] Wajnryb raised a question whether there can be a nongeometric embedding. We here show that map $\phi : B_g \rightarrow \Gamma_{g,1}$ is a nongeometric embedding.

Lemma 3.7. *The map $\phi : B_g \rightarrow \Gamma_{g,1}$ is a nongeometric embedding.*

Proof. For a standard generator σ_i of B_g , $\phi(\sigma_i) = \beta_i$ is composite of three half Dehn twists along three disjoint simple closed curves in the local surface $S_{2,1}$. We may think that this self-homeomorphism is supported in the inner surface $S_{0,3}$ obtained by removing the outer dotted parts of the surface $S_{2,1}$ as given in Figure ???. β_i cannot be equal to a full Dehn twist along a simple closed curve in $S_{0,3}$ because there are only three simple closed curves in $S_{0,3}$ and any Dehn twist along one of those three closed curves is not equal to $\phi(\sigma_i)$. We can see this by comparing the actions of them on the fundamental group of $S_{2,1}$. \square

From the geometric construction of β'_i s and the actions of them on the fundamental group of the surface, we have the following theorem.

Theorem 3.8. *Let $\mathcal{B} = \coprod_{g \geq 0} B_g$ and $\mathcal{M} = \coprod_{g \geq 0} \Gamma_{g,1}$. The nongeometric embedding $\phi : B_g \rightarrow \Gamma_{g,1}, \sigma_i \mapsto \beta_i$ extends to a braided monoidal functor $\Phi : \mathcal{B} \rightarrow \mathcal{M}$.*

FIGURE 10. Three 180° twists along the boundary of $S_{0,3}$

The map $\phi : B_g \rightarrow \Gamma_{g,1}$ is a more natural embedding than the Harer map which is defined by

$$h(\sigma_i) = \begin{cases} b_{\frac{i+1}{2}} & \text{if } i \text{ is odd} \\ w_{\frac{i}{2}} & \text{if } i \text{ is even.} \end{cases}$$

(See Figure 1 for the definitions of b_i 's and w_i 's.), because the map h does not directly give rise to a braided monoidal functor for \mathcal{B} to \mathcal{M} . (See [9], Theorem 3.3)

The group completion of the classifying space of a braided monoidal category is naturally homotopy equivalent to a double loop space ([3], [4], [5], [2]). Therefore, Φ induces a map $\overline{B\mathcal{B}} \rightarrow \overline{B\mathcal{M}}$ of double loop spaces, where $\overline{B\mathcal{B}}$ and $\overline{B\mathcal{M}}$ denote the group completions of $B\mathcal{B}$ and $B\mathcal{M}$, respectively.

Since $\overline{B\mathcal{B}} \simeq \mathbb{Z}_+ \times BB_\infty^+$ and $\overline{B\mathcal{M}} \simeq \mathbb{Z}_+ \times B\Gamma_\infty^+$, we have

Lemma 3.9. $B\phi^+ : BB_\infty^+ \rightarrow B\Gamma_\infty^+$ is a map of double loop spaces, and is null-homotopic.

It is known ([10], Lemma 5.3) that every double loop map $f : BB_\infty^+ \rightarrow B\Gamma_\infty^+$ is null-homotopic.

Since the plus construction does not change homology groups, Lemma 3.9 implies that the homology homomorphism induced by ϕ is trivial in the stable range.

Theorem 3.10. The homomorphism $\phi_* : H_*(B_\infty; \mathbb{Z}) \rightarrow H_*(\Gamma_\infty; \mathbb{Z})$ induced by ϕ is trivial.

By the homology stability theorem ([6], [7]) we have that $\phi_* : H_i(B_g; \mathbb{Z}) \rightarrow H_i(\Gamma_{g,1}; \mathbb{Z})$ is trivial for $0 < i < \frac{g}{2}$.

ACKNOWLEDGEMENTS

This research was supported by Basic Science Research Program through the National Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2011-0004509).

REFERENCES

- [1] E. Artin, *Theorie der Zöpfe*, Abh. Math. Sem. Hambur. Univ. **4** (1926), 47–72.
- [2] C. Batenau, Z. Fiedorowicz, R. Schwänzl, R. Vogt, *Iterated monoidal categories*, Adv. in Math. **176**(2003), 277–349.

- [3] C. Berger, *Double loop spaces, braided monoidal categories and algebraic 3-type of space*, Contemporary Math. 227(2009), 49–66.
- [4] P. Carrasco, A.M. Cegarra, and A.R. Garazón, *Classifying spaces for braided monoidal categories and lax diagrams of bicategories*, arXiv0907.0930v3(2009).
- [5] Z. Fiedorowicz, *The symmetric bar construction*, Preprint, available at <http://www.math.ohio-state.edu/~fiedorowicz>.
- [6] J. Harer, *Stability of the homology of the mapping class groups of orientable surfaces*, Ann. of Math. (2) **121** (1985), no. 2, 215–249.
- [7] N. V. Ivanov, *Stabilization of the homology of Teichmüller modular groups*, Algebra i Analiz **1** (1989), no. 3, 110–126; Leningrad Math. J. **1** (1990), no. 3, 675–691.
- [8] Y. Song, *The braidings of mapping class groups and loop spaces*, Tohoku Math. J. **52** (2000), 309–319.
- [9] Y. Song, *The action of image of braiding under the harer map*, Commun. Korea math. Soc. **21** (2006), No. 2, 337–345.
- [10] Y. Song and U. Tillmann, *Braid, mapping class groups, and categorical delooping*, Math. Ann. **339** (2007), 377–393.
- [11] U. Tillmann, *On the homotopy of stable mapping class group*, Invent. Math. **130** (1997), 257–275.
- [12] U. Tillmann, *Artin’s map in stable homology*, Bull. Lond. Math. Soc. **39** (2007), no. 6, 989–992.
- [13] B. Wajnryb, *A simple representation for mapping class group of an orientable surface*, Israel Journal of Math. **45** (1983), 157–174.
- [14] B. Wajnryb, *Artin groups and geometric monodromy*, Invent. Math. **138** (1999), No. 3, 563–571.
- [15] B. Wajnryb, *A simple representation for mapping class group of an orientable surface Relations in the mapping class group*, Problems on mapping class groups and related topics, Proc. Pure Math. **74** (2006), Amer. Math. Soc., 115–120.

DEPARTMENTS OF MATHEMATICS, INHA UNIVERSITY, INCHEON 402-751, KOREA
E-mail address: yjsong@inha.ac.kr