

**POSITIVE SOLUTIONS OF NONLINEAR THREE-POINT
INTEGRAL BOUNDARY-VALUE PROBLEMS FOR
SECOND-ORDER DIFFERENTIAL EQUATIONS**

FAOUZI HADDOUCHI, SLIMANE BENAICHA

ABSTRACT. We investigate the existence of positive solutions to the nonlinear second-order three-point integral boundary value problem

$$u''(t) + a(t)f(u(t)) = 0, \quad 0 < t < T,$$

$$u(0) = \beta u(\eta), \quad u(T) = \alpha \int_0^\eta u(s) ds,$$

where $0 < \eta < T$, $0 < \alpha < \frac{2T}{\eta^2}$, $0 \leq \beta < \frac{2T - \alpha\eta^2}{\alpha\eta^2 - 2\eta + 2T}$ are given constants. We show the existence of at least one positive solution if f is either superlinear or sublinear by applying Krasnoselskii's fixed point theorem in cones.

1. INTRODUCTION

The study of the existence of solutions of multipoint boundary value problems for linear second-order ordinary differential equations was initiated by II'in and Moiseev [5]. Then Gupta [2] studied three-point boundary value problems for nonlinear second-order ordinary differential equations. Since then, nonlinear second-order three-point boundary value problems have also been studied by several authors. We refer the reader to [1, 3, 4, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21] and the references therein.

Tariboon and Sitthiwirattam [20] proved the existence of positive solutions for the three-point boundary-value problem with integral condition

$$u''(t) + a(t)f(u(t)) = 0, \quad t \in (0, 1), \quad (1.1)$$

$$u(0) = 0, \quad u(1) = \alpha \int_0^\eta u(s) ds, \quad (1.2)$$

where $0 < \eta < 1$ and $0 < \alpha < \frac{2}{\eta^2}$.

This paper is concerned with the existence of positive solutions of the equation

$$u''(t) + a(t)f(u(t)) = 0, \quad t \in (0, T), \quad (1.3)$$

with the three-point integral boundary condition

$$u(0) = \beta u(\eta), \quad u(T) = \alpha \int_0^\eta u(s) ds, \quad (1.4)$$

where $\alpha > 0$, $\beta \geq 0$, $\eta \in (0, T)$ are given constants. Clearly if $\beta = 0$ and $T = 1$, then (1.4) reduces to (1.2). The purpose of this paper is to give some results for existence

2000 *Mathematics Subject Classification.* 34B15, 34C25, 34B18.

Key words and phrases. Positive solutions; Krasnoselskii's fixed point theorem; Three-point boundary value problems; Cone.

for positive solutions to (1.3)-(1.4), assuming that $0 < \alpha < \frac{2T}{\eta^2}$, $0 \leq \beta < \frac{2T - \alpha\eta^2}{\alpha\eta^2 - 2\eta + 2T}$ and f is either superlinear or sublinear.

Our results extend and complete those obtained by J. Tariboon and T. Sitthiraththam [20]. On the other hand, we point out that the proof of the last part in the sublinear case ($f_\infty = 0$) of Theorem 3.1 in [20] is not correct since it is based on an inequality which is not true. We give a new proof, which is different from that of Theorem 3.1 in [20], and obtain an extended result.

Set

$$f_0 = \lim_{u \rightarrow 0^+} \frac{f(u)}{u}, \quad f_\infty = \lim_{u \rightarrow \infty} \frac{f(u)}{u}. \quad (1.5)$$

Then $f_0 = 0$ and $f_\infty = \infty$ correspond to the superlinear case, and $f_0 = \infty$ and $f_\infty = 0$ correspond to the sublinear case. By the positive solution of (1.3)-(1.4) we mean that function $u(t)$ is positive on $0 < t < T$ and satisfies the problem (1.3)-(1.4).

Throughout this paper, we assume the following hypotheses:

(H1) $f \in C([0, \infty), [0, \infty))$.

(H2) $a \in C([0, T], [0, \infty))$ and there exists $t_0 \in [\eta, T]$ such that $a(t_0) > 0$.

The following theorem (Krasnoselskii's fixed-point theorem), will play an important role in the proof of our main results.

Theorem 1.1 ([6]). *Let E be a Banach space, and let $K \subset E$ be a cone. Assume Ω_1, Ω_2 are open bounded subsets of E with $0 \in \Omega_1, \overline{\Omega}_1 \subset \Omega_2$, and let*

$$A : K \cap (\overline{\Omega}_2 \setminus \Omega_1) \longrightarrow K$$

be a completely continuous operator such that either

(i) $\|Au\| \leq \|u\|$, $u \in K \cap \partial\Omega_1$, and $\|Au\| \geq \|u\|$, $u \in K \cap \partial\Omega_2$; or

(ii) $\|Au\| \geq \|u\|$, $u \in K \cap \partial\Omega_1$, and $\|Au\| \leq \|u\|$, $u \in K \cap \partial\Omega_2$

hold. Then A has a fixed point in $K \cap (\overline{\Omega}_2 \setminus \Omega_1)$.

2. PRELIMINARIES

To prove the main existence results we will employ several straightforward lemmas. These lemmas are based on the linear boundary-value problem.

Lemma 2.1. *Let $\beta \neq \frac{2T - \alpha\eta^2}{\alpha\eta^2 - 2\eta + 2T}$. Then for $y \in C([0, T], \mathbb{R})$, the problem*

$$u''(t) + y(t) = 0, \quad t \in (0, T), \quad (2.1)$$

$$u(0) = \beta u(\eta), \quad u(T) = \alpha \int_0^\eta u(s) ds, \quad (2.2)$$

has a unique solution

$$\begin{aligned} u(t) = & \frac{\beta(2T - \alpha\eta^2) - 2\beta(1 - \alpha\eta)t}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^\eta (\eta - s)y(s) ds \\ & + \frac{\alpha\beta\eta - \alpha(\beta - 1)t}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^\eta (\eta - s)^2 y(s) ds \\ & + \frac{2(\beta - 1)t - 2\beta\eta}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^T (T - s)y(s) ds - \int_0^t (t - s)y(s) ds. \end{aligned}$$

Proof. From (2.1), we have

$$u(t) = u(0) + u'(0)t - \int_0^t (t-s)y(s)ds. \quad (2.3)$$

Integrating (2.3) from 0 to η , where $\eta \in (0, T)$, we have

$$\begin{aligned} \int_0^\eta u(s)ds &= u(0)\eta + u'(0)\frac{\eta^2}{2} - \int_0^\eta \left(\int_0^\tau (\tau-s)y(s)ds \right) d\tau \\ &= u(0)\eta + u'(0)\frac{\eta^2}{2} - \frac{1}{2} \int_0^\eta (\eta-s)^2 y(s)ds. \end{aligned}$$

Since

$$\begin{aligned} u(T) &= u(0) + u'(0)T - \int_0^T (T-s)y(s)ds \quad \text{and} \\ u(\eta) &= u(0) + u'(0)\eta - \int_0^\eta (\eta-s)y(s)ds. \end{aligned}$$

By (2.2), from $u(0) = \beta u(\eta)$, we have

$$(\beta - 1)u(0) + \eta\beta u'(0) = \beta \int_0^\eta (\eta-s)y(s)ds,$$

and from $u(T) = \alpha \int_0^\eta u(s)ds$, we have

$$(1 - \alpha\eta)u(0) + (T - \alpha\frac{\eta^2}{2})u'(0) = \int_0^T (T-s)y(s)ds - \frac{\alpha}{2} \int_0^\eta (\eta-s)^2 y(s)ds.$$

Therefore,

$$\begin{aligned} u(0) &= \frac{\beta(2T - \alpha\eta^2)}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^\eta (\eta-s)y(s)ds \\ &\quad - \frac{2\beta\eta}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^T (T-s)y(s)ds \\ &\quad + \frac{\alpha\beta\eta}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^\eta (\eta-s)^2 y(s)ds, \\ u'(0) &= \frac{2(\beta - 1)}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^T (T-s)y(s)ds \\ &\quad - \frac{\alpha(\beta - 1)}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^\eta (\eta-s)^2 y(s)ds \\ &\quad - \frac{2\beta(1 - \alpha\eta)}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^\eta (\eta-s)y(s)ds, \end{aligned}$$

from which it follows that

$$\begin{aligned} u(t) &= \frac{\beta(2T - \alpha\eta^2) - 2\beta(1 - \alpha\eta)t}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^\eta (\eta-s)y(s)ds \\ &\quad + \frac{\alpha\beta\eta - \alpha(\beta - 1)t}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^\eta (\eta-s)^2 y(s)ds \\ &\quad + \frac{2(\beta - 1)t - 2\beta\eta}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^T (T-s)y(s)ds - \int_0^t (t-s)y(s)ds. \end{aligned}$$

□

Lemma 2.2. *Let $0 < \alpha < \frac{2T}{\eta^2}$, $0 \leq \beta < \frac{2T - \alpha\eta^2}{\alpha\eta^2 - 2\eta + 2T}$. If $y \in C([0, T], [0, \infty))$, then the unique solution u of (2.1)-(2.2) satisfies $u(t) \geq 0$ for $t \in [0, T]$.*

Proof. From the fact that $u''(t) = -y(t) \leq 0$, we have that the graph of $u(t)$ is concave down on $(0, T)$ and

$$\int_0^\eta u(s)ds \geq \frac{\eta}{2}(u(0) + u(\eta)). \quad (2.4)$$

Combining (2.2) with (2.4), we get

$$u(T) \geq \frac{\alpha(\beta + 1)\eta}{2}u(\eta). \quad (2.5)$$

Since the graph of u is concave down, we get

$$\frac{u(\eta) - u(0)}{\eta} \geq \frac{u(T) - u(0)}{T}.$$

Combining this with (2.2) and (2.5), we obtain

$$(1 - \beta)\frac{u(\eta)}{\eta} \geq \frac{\alpha(\beta + 1)\eta - 2\beta}{2T}u(\eta).$$

If $u(0) < 0$, then $u(\eta) < 0$. It implies $\beta \geq \frac{2T - \alpha\eta^2}{\alpha\eta^2 - 2\eta + 2T}$, a contradiction to $\beta < \frac{2T - \alpha\eta^2}{\alpha\eta^2 - 2\eta + 2T}$.

If $u(T) < 0$, then $u(\eta) < 0$, and the same contradiction emerges. Thus, it is true that $u(0) \geq 0$, $u(T) \geq 0$, together with the concavity of $u(t)$, we have $u(t) \geq 0$ for $t \in [0, T]$. \square

Lemma 2.3. *Let $\alpha > \frac{2T}{\eta^2}$, $\beta \geq 0$. If $y \in C([0, T], [0, \infty))$, then the problem (2.1)-(2.2) has no positive solutions.*

Proof. Suppose that problem (2.1)-(2.2) has a positive solution u satisfying $u(t) \geq 0$, $t \in [0, T]$.

If $u(T) > 0$, then $\int_0^\eta u(s)ds > 0$. It implies

$$u(T) = \alpha \int_0^\eta u(s)ds > \frac{2T}{\eta^2} \frac{\eta}{2}(u(0) + u(\eta)) = \frac{T(\beta + 1)u(\eta)}{\eta} \geq \frac{Tu(\eta)}{\eta},$$

that is

$$\frac{u(T)}{T} > \frac{u(\eta)}{\eta},$$

which is a contradiction to the concavity of u .

If $u(T) = 0$, then $\int_0^\eta u(s)ds = 0$, this is $u(t) \equiv 0$ for all $t \in [0, \eta]$. If there exists $t_0 \in (\eta, T)$ such that $u(t_0) > 0$, then $u(0) = u(\eta) < u(t_0)$, a violation of the concavity of u . Therefore, no positive solutions exist. \square

Lemma 2.4. *Let $0 < \alpha < \frac{2T}{\eta^2}$, $0 \leq \beta < \frac{2T - \alpha\eta^2}{\alpha\eta^2 - 2\eta + 2T}$. If $y \in C([0, T], [0, \infty))$, then the unique solution u of (2.1)-(2.2) satisfies*

$$\min_{t \in [\eta, T]} u(t) \geq \gamma \|u\|, \quad \|u\| = \max_{t \in [0, T]} |u(t)|, \quad (2.6)$$

where

$$\gamma := \min \left\{ \frac{\eta}{T}, \frac{\alpha(\beta + 1)\eta^2}{2T}, \frac{\alpha(\beta + 1)\eta(T - \eta)}{2T - \alpha(\beta + 1)\eta^2} \right\} \in (0, 1). \quad (2.7)$$

Proof. We divide the proof into three cases. Set $u(t_1) = \|u\|$.

Case 1. If $\eta \leq t_1 \leq T$ and $\min_{t \in [\eta, T]} u(t) = u(\eta)$, then the concavity of u implies that

$$\frac{u(\eta)}{\eta} \geq \frac{u(t_1)}{t_1} \geq \frac{u(t_1)}{T}.$$

Thus,

$$\min_{t \in [\eta, T]} u(t) \geq \frac{\eta}{T} \|u\|.$$

Case 2. If $\eta \leq t_1 \leq T$ and $\min_{t \in [\eta, T]} u(t) = u(T)$, then (2.2), (2.4) and the concavity of u implies

$$\begin{aligned} u(T) &= \alpha \int_0^\eta u(s) ds \geq \frac{\alpha(\beta+1)\eta^2}{2} \frac{u(\eta)}{\eta} \\ &\geq \frac{\alpha(\beta+1)\eta^2}{2} \frac{u(t_1)}{t_1} \\ &\geq \frac{\alpha(\beta+1)\eta^2}{2} \frac{u(t_1)}{T}. \end{aligned}$$

This implies that

$$\min_{t \in [\eta, T]} u(t) \geq \frac{\alpha(\beta+1)\eta^2}{2T} \|u\|.$$

Case 3. If $t_1 \leq \eta < T$, then $\min_{t \in [\eta, T]} u(t) = u(T)$. Using the concavity of u and (2.2), (2.4), we obtain

$$\begin{aligned} u(t_1) &\leq u(T) + \frac{u(T) - u(\eta)}{T - \eta} (t_1 - T) \\ &\leq u(T) + \frac{u(T) - u(\eta)}{T - \eta} (0 - T) \\ &\leq u(T) \left[1 - T \frac{1 - \frac{2}{\alpha(\beta+1)\eta}}{T - \eta} \right] \\ &= \frac{2T - \alpha(\beta+1)\eta^2}{\alpha(\beta+1)\eta(T - \eta)} u(T), \end{aligned}$$

from which it follows that

$$\min_{t \in [\eta, T]} u(t) \geq \frac{\alpha(\beta+1)\eta(T - \eta)}{2T - \alpha(\beta+1)\eta^2} \|u\|.$$

Summing up, we have

$$\min_{t \in [\eta, T]} u(t) \geq \gamma \|u\|,$$

where

$$\gamma := \min \left\{ \frac{\eta}{T}, \frac{\alpha(\beta+1)\eta^2}{2T}, \frac{\alpha(\beta+1)\eta(T - \eta)}{2T - \alpha(\beta+1)\eta^2} \right\}.$$

This completes the proof. \square

3. EXISTENCE OF POSITIVE SOLUTIONS

Now we are in the position to establish the main result.

Theorem 3.1. *Assume (H1) and (H2) hold, and $0 < \alpha < \frac{2T}{\eta^2}$, $0 \leq \beta < \frac{2T - \alpha\eta^2}{\alpha\eta^2 - 2\eta + 2T}$. Then the problem (1.3)-(1.4) has at least one positive solution in the case*

- (i) $f_0 = 0$ and $f_\infty = \infty$ (superlinear), or
- (ii) $f_0 = \infty$ and $f_\infty = 0$ (sublinear).

Proof. It is known that $0 < \alpha < \frac{2T}{\eta^2}$, $0 \leq \beta < \frac{2T - \alpha\eta^2}{\alpha\eta^2 - 2\eta + 2T}$. From Lemma 2.1, u is a solution to the boundary value problem (1.3)-(1.4) if and only if u is a fixed point of operator A , where A is defined by

$$\begin{aligned} Au(t) = & \frac{\beta(2T - \alpha\eta^2) - 2\beta(1 - \alpha\eta)t}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^\eta (\eta - s)a(s)f(u(s))ds \\ & + \frac{\alpha\beta\eta - \alpha(\beta - 1)t}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^\eta (\eta - s)^2a(s)f(u(s))ds \\ & + \frac{2(\beta - 1)t - 2\beta\eta}{(\alpha\eta^2 - 2T) - \beta(2\eta - \alpha\eta^2 - 2T)} \int_0^T (T - s)a(s)f(u(s))ds \\ & - \int_0^t (t - s)a(s)f(u(s))ds. \end{aligned}$$

Denote

$$K = \left\{ u/u \in C([0, T], \mathbb{R}), u \geq 0, \min_{t \in [\eta, T]} u(t) \geq \gamma \|u\| \right\},$$

where γ is defined in (2.7). It is obvious that K is a cone in $C([0, T], \mathbb{R})$. Moreover, from Lemma 2.2 and Lemma 2.4, $AK \subset K$. It is also easy to check that $A : K \rightarrow K$ is completely continuous.

Superlinear case. $f_0 = 0$ and $f_\infty = \infty$. Since $f_0 = 0$, we may choose $H_1 > 0$ so that $f(u) \leq \epsilon u$, for $0 < u \leq H_1$, where $\epsilon > 0$ satisfies

$$\epsilon \frac{2(\beta + 1) + T^{-1}\beta\eta(\alpha\eta + 2) + \alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T T(T - s)a(s)ds \leq 1.$$

Thus, if we let

$$\Omega_1 = \{u \in C([0, T], \mathbb{R}) : \|u\| < H_1\},$$

then, for $u \in K \cap \partial\Omega_1$, we get

$$\begin{aligned}
 Au(t) &\leq \frac{2\beta(1-\alpha\eta)t - \beta(2T - \alpha\eta^2)}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta - s)a(s)f(u(s))ds \\
 &\quad + \frac{\alpha(\beta - 1)t - \alpha\beta\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta - s)^2 a(s)f(u(s))ds \\
 &\quad + \frac{2\beta\eta - 2(\beta - 1)t}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T (T - s)a(s)f(u(s))ds \\
 &\leq \frac{2\beta T + \alpha\beta\eta^2}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta - s)a(s)f(u(s))ds \\
 &\quad + \frac{\alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta - s)^2 a(s)f(u(s))ds \\
 &\quad + \frac{2\beta\eta + 2T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T (T - s)a(s)f(u(s))ds \\
 &\leq \frac{2T(\beta + 1) + \beta\eta(\alpha\eta + 2)}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T (T - s)a(s)f(u(s))ds \\
 &\quad + \frac{\alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta - s)^2 a(s)f(u(s))ds \\
 &\leq \frac{2T(\beta + 1) + \beta\eta(\alpha\eta + 2)}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T (T - s)a(s)f(u(s))ds \\
 &\quad + \frac{\alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T T(T - s)a(s)f(u(s))ds \\
 &= \frac{2(\beta + 1) + T^{-1}\beta\eta(\alpha\eta + 2) + \alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T T(T - s)a(s)f(u(s))ds \\
 &\leq \epsilon\|u\| \frac{2(\beta + 1) + T^{-1}\beta\eta(\alpha\eta + 2) + \alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T T(T - s)a(s)ds \\
 &\leq \|u\|.
 \end{aligned}$$

Thus $\|Au\| \leq \|u\|$, $u \in K \cap \partial\Omega_1$.

Further, since $f_\infty = \infty$, there exists $\widehat{H}_2 > 0$ such that $f(u) \geq \rho u$ for $u \geq \widehat{H}_2$, where $\rho > 0$ is chosen so that

$$\rho\gamma \frac{2\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_\eta^T (T - s)a(s)ds \geq 1.$$

Let $H_2 = \max\{2H_1, \frac{\widehat{H}_2}{\gamma}\}$ and $\Omega_2 = \{u \in C([0, T], \mathbb{R}) : \|u\| < H_2\}$. Then $u \in K \cap \partial\Omega_2$ implies that

$$\min_{t \in [\eta, T]} u(t) \geq \gamma\|u\| = \gamma H_2 \geq \widehat{H}_2,$$

and so,

$$\begin{aligned}
Au(\eta) &= \frac{2\beta(1-\alpha\eta)\eta - \beta(2T - \alpha\eta^2)}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta - s)a(s)f(u(s))ds \\
&\quad + \frac{\alpha(\beta - 1)\eta - \alpha\beta\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta - s)^2 a(s)f(u(s))ds \\
&\quad + \frac{2\beta\eta - 2(\beta - 1)\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T (T - s)a(s)f(u(s))ds \\
&\quad - \int_0^\eta (\eta - s)a(s)f(u(s))ds \\
&= \frac{2\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T (T - s)a(s)f(u(s))ds \\
&\quad - \frac{\alpha\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta^2 - 2\eta s + s^2)a(s)f(u(s))ds \\
&\quad - \frac{2T - \alpha\eta^2}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta - s)a(s)f(u(s))ds \\
&= \frac{2\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T (T - s)a(s)f(u(s))ds \\
&\quad + \frac{\alpha\eta^2}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta sa(s)f(u(s))ds \\
&\quad + \frac{2T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta sa(s)f(u(s))ds \\
&\quad - \frac{\alpha\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta s^2 a(s)f(u(s))ds \\
&\quad - \frac{2T\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta a(s)f(u(s))ds \\
&= \frac{2\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_\eta^T (T - s)a(s)f(u(s))ds \\
&\quad + \frac{2(T - \eta)}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta sa(s)f(u(s))ds \\
&\quad + \frac{\alpha\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta s(\eta - s)a(s)f(u(s))ds \\
&\geq \frac{2\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_\eta^T (T - s)a(s)f(u(s))ds \\
&\geq \frac{2\eta\rho}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_\eta^T (T - s)a(s)u(s)ds \\
&\geq \frac{2\eta\rho\gamma\|u\|}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_\eta^T (T - s)a(s)ds \\
&\geq \|u\|.
\end{aligned}$$

Hence, $\|Au\| \geq \|u\|$, $u \in K \cap \partial\Omega_2$. By the first part of Theorem 1.1, A has a fixed point in $K \cap (\bar{\Omega}_2 \setminus \Omega_1)$ such that $H_1 \leq \|u\| \leq H_2$. This completes the superlinear part of the theorem.

Sublinear case. $f_0 = \infty$ and $f_\infty = 0$. Since $f_0 = \infty$, choose $H_3 > 0$ such that $f(u) \geq Mu$ for $0 < u \leq H_3$, where $M > 0$ satisfies

$$M\gamma \frac{2\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_\eta^T (T-s)a(s)ds \geq 1.$$

Let $\Omega_3 = \{u \in C([0, T], \mathbb{R}) : \|u\| < H_3\}$, then for $u \in K \cap \partial\Omega_3$, we get

$$\begin{aligned} Au(\eta) &= \frac{2\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T (T-s)a(s)f(u(s))ds \\ &\quad - \frac{\alpha\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta-s)^2 a(s)f(u(s))ds \\ &\quad - \frac{2T - \alpha\eta^2}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta-s)a(s)f(u(s))ds \\ &\geq \frac{2\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_\eta^T (T-s)a(s)f(u(s))ds \\ &\geq \frac{2\eta M}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_\eta^T (T-s)a(s)u(s)ds \\ &\geq M\gamma \frac{2\eta\|u\|}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_\eta^T (T-s)a(s)ds \\ &\geq \|u\|. \end{aligned}$$

Thus $\|Au\| \geq \|u\|$, $u \in K \cap \partial\Omega_3$. Now, since $f_\infty = 0$, there exists $\hat{H}_4 > 0$ so that $f(u) \leq \lambda u$ for $u \geq \hat{H}_4$, where $\lambda > 0$ satisfies

$$\lambda \frac{2(\beta+1) + T^{-1}\beta\eta(\alpha\eta+2) + \alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T T(T-s)a(s)ds \leq 1.$$

We consider two cases:

Case (i). Suppose f is bounded, say $f(u) \leq N$ for all $u \in [0, \infty)$. Choosing $H_4 \geq \max\{2H_3, \frac{N}{\lambda}\}$. For $u \in K$ with $\|u\| = H_4$, we have

$$\begin{aligned} Au(t) &= \frac{2\beta(1 - \alpha\eta)t - \beta(2T - \alpha\eta^2)}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta-s)a(s)f(u(s))ds \\ &\quad + \frac{\alpha(\beta-1)t - \alpha\beta\eta}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^\eta (\eta-s)^2 a(s)f(u(s))ds \\ &\quad + \frac{2\beta\eta - 2(\beta-1)t}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T (T-s)a(s)f(u(s))ds \\ &\quad - \int_0^t (t-s)a(s)f(u(s))ds \\ &\leq \frac{2(\beta+1) + T^{-1}\beta\eta(\alpha\eta+2) + \alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T T(T-s)a(s)f(u(s))ds \\ &\leq N \frac{2(\beta+1) + T^{-1}\beta\eta(\alpha\eta+2) + \alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T T(T-s)a(s)ds \\ &\leq H_4\lambda \frac{2(\beta+1) + T^{-1}\beta\eta(\alpha\eta+2) + \alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T T(T-s)a(s)ds \\ &\leq H_4, \end{aligned}$$

and therefore $\|Au\| \leq \|u\|$. \square

Case (ii). If f is unbounded, then we know from $f \in C([0, \infty), [0, \infty))$ that there is H_4 : $H_4 \geq \max\{2H_3, \frac{\hat{H}_4}{\gamma}\}$ such that

$$f(u) \leq f(H_4) \text{ for } u \in [0, H_4].$$

Then for $u \in K$ and $\|u\| = H_4$, we have

$$\begin{aligned} Au(t) &\leq \frac{2(\beta + 1) + T^{-1}\beta\eta(\alpha\eta + 2) + \alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T T(T-s)a(s)f(u(s))ds \\ &\leq \frac{2(\beta + 1) + T^{-1}\beta\eta(\alpha\eta + 2) + \alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T T(T-s)a(s)f(H_4)ds \\ &\leq H_4\lambda \frac{2(\beta + 1) + T^{-1}\beta\eta(\alpha\eta + 2) + \alpha\beta T}{(2T - \alpha\eta^2) - \beta(\alpha\eta^2 - 2\eta + 2T)} \int_0^T T(T-s)a(s)ds \\ &\leq H_4 = \|u\|. \end{aligned}$$

Therefore, in either case we may set

$$\Omega_4 = \{u \in C([0, T], \mathbb{R}) : \|u\| < H_4\},$$

and for $u \in K \cap \partial\Omega_4$ we may have $\|Au\| \leq \|u\|$. By the second part of Theorem 1.1, it follows that A has a fixed point in $K \cap (\bar{\Omega}_4 \setminus \Omega_3)$ such that $H_3 \leq \|u\| \leq H_4$. This completes the sublinear part of the theorem. Therefore, the problem (1.3)-(1.4) has at least one positive solution.

4. SOME EXAMPLES

In this section, in order to illustrate our result, we consider some examples.

Example 4.1. Consider the boundary value problem

$$u''(t) + tu^p = 0, \quad 0 < t < 2, \quad (4.1)$$

$$u(0) = \frac{1}{2}u\left(\frac{3}{2}\right), \quad u(2) = \int_0^{\frac{3}{2}} u(s)ds. \quad (4.2)$$

Set $\beta = 1/2$, $\alpha = 1$, $\eta = 3/2$, $T = 2$, $a(t) = t$, $f(u) = u^p$. We can show that $0 < \alpha = 1 < 16/9 = 2T/\eta^2$, $0 < \beta = 1/2 < 7/13 = (2T - \alpha\eta^2)/(\alpha\eta^2 - 2\eta + 2T)$.

Now we consider the existence of positive solutions of the problem (4.1), (4.2) in two cases.

Case 1: $p > 1$. In this case, $f_0 = 0$, $f_\infty = \infty$ and (i) holds. Then (4.1), (4.2) has at least one positive solution.

Case 2: $p \in (0, 1)$. In this case, $f_0 = \infty$, $f_\infty = 0$ and (ii) holds. Then ((4.1), (4.2) has at least one positive solution.

Example 4.2. Consider the boundary value problem

$$u''(t) + t^2u^2 \ln(1+u) = 0, \quad 0 < t < \frac{3}{4}, \quad (4.3)$$

$$u(0) = \frac{1}{10}u\left(\frac{1}{4}\right), \quad u\left(\frac{3}{4}\right) = 20 \int_0^{\frac{1}{4}} u(s)ds. \quad (4.4)$$

Set $\beta = 1/10$, $\alpha = 20$, $\eta = 1/4$, $T = 3/4$, $a(t) = t^2$, $f(u) = u^2 \ln(1 + u)$. We can show that $0 < \alpha = 20 < 24 = 2T/\eta^2$, $0 < \beta = 1/10 < 1/9 = (2T - \alpha\eta^2)/(\alpha\eta^2 - 2\eta + 2T)$. Through a simple calculation we can get $f_0 = 0$ and $f_\infty = \infty$. Thus, by the first part of Theorem 3.1, we can get that the problem (4.3), (4.4) has at least one positive solution.

Example 4.3. Consider the boundary value problem

$$u''(t) + e^t \frac{\sin u + \ln(1 + u)}{u^2} = 0, \quad 0 < t < 1, \quad (4.5)$$

$$u(0) = u\left(\frac{1}{3}\right), \quad u(1) = 2 \int_0^{\frac{1}{3}} u(s) ds. \quad (4.6)$$

Set $\beta = 1$, $\alpha = 2$, $\eta = 1/3$, $T = 1$, $a(t) = e^t$, $f(u) = (\sin u + \ln(1 + u))/u^2$. We can show that $0 < \alpha = 2 < 18 = 2T/\eta^2$, $0 < \beta = 1 < 8/7 = (2T - \alpha\eta^2)/(\alpha\eta^2 - 2\eta + 2T)$. Through a simple calculation we can get $f_0 = \infty$ and $f_\infty = 0$. Thus, by the second part of Theorem 3.1, we can get that the problem (4.5), (4.6) has at least one positive solution.

REFERENCES

- [1] Z. Chengbo; Positive solutions for semi-positone three-point boundary value problems, J. Comput. Appl. Math. **228** (2009), 279-286.
- [2] C. P. Gupta; Solvability of a three-point nonlinear boundary value problem for a second order ordinary differential equations, J. Math. Anal. Appl. **168** (1992), 540-551.
- [3] Y. Guo, W. Ge; Positive solutions for three-point boundary value problems with dependence on the first order derivative, J. Math. Anal. Appl. **290** (2004), 291-301.
- [4] X. Han, Positive solutions for a three-point boundary value problem, Nonlinear Anal. **66** (2007), 679-688.
- [5] V. A. Il'in, E. I. Moiseev; Nonlocal boundary-value problem of the first kind for a Sturm-Liouville operator in its differential and finite difference aspects, Differ. Equ. **23** (1987), 803-810.
- [6] M. A. Krasnoselskii; Positive Solutions of Operator Equations, P. Noordhoff, Groningen, The Netherlands, 1964.
- [7] S. Liang, L. Mu; Multiplicity of positive solutions for singular three-point boundary value problems at resonance, Nonlinear Anal. **71** (2009), 2497-2505.
- [8] R. Liang, J. Peng, J. Shen; Positive solutions to a generalized second order three-point boundary value problem, Appl. Math. Comput. **196** (2008), 931-940.
- [9] J. Li, J. Shen; Multiple positive solutions for a second-order three-point boundary value problem, Appl. Math. Comput. **182** (2006), 258-268.
- [10] B. Liu; Positive solutions of a nonlinear three-point boundary value problem, Appl. Math. Comput. **132** (2002), 11-28.
- [11] B. Liu; Positive solutions of a nonlinear three-point boundary value problem, Comput. Math. Appl. **44** (2002), 201-211.
- [12] B. Liu, L. Liu, Y. Wu; Positive solutions for singular second order three-point boundary value problems, Nonlinear Anal. **66** (2007), 2756-2766.
- [13] H. Luo, Q. Ma; Positive solutions to a generalized second-order three-point boundary-value problem on time scales, Electron. J. Differen. Equat. **2005**(2005), No. 17, 1-14.
- [14] R. Ma; Multiplicity of positive solutions for second-order three-point boundary value problems, Comput. Math. Appl. **40** (2000), 193-204.
- [15] R. Ma; Positive solutions of a nonlinear three-point boundary-value problem, Electron. J. Differen. Equat. **1999**(1999), No. 34, 1-8.
- [16] R. Ma; Positive solutions for second-order three-point boundary value problems, Appl. Math. Lett. **14** (2001), 1-5.
- [17] H. Pang, M. Feng, W. Ge; Existence and monotone iteration of positive solutions for a three-point boundary value problem, Appl. Math. Lett. **21** (2008), 656-661.

- [18] Y. Sun, L. Liu, J. Zhang, R. P. Agarwal; Positive solutions of singular three-point boundary value problems for second-order differential equations, *J. Comput. Appl. Math.* **230** (2009), 738-750.
- [19] H. Sun, W. Li; Positive solutions for nonlinear three-point boundary value problems on time scales, *J. Math. Anal. Appl.* **299** (2004), 508-524.
- [20] J. Tariboon, T. Sitthiwiratttham; Positive solutions of a nonlinear three-point integral boundary value problem, *Bound. Val. Prob.* **2010** (2010), ID 519210, 11 pages, doi:10.1155/2010/519210.
- [21] X. Xu; Multiplicity results for positive solutions of some semi-positone three-point boundary value problems, *J. Math. Anal. Appl.* **291** (2004), 673-689.

FAOUZI HADDOUCHI, DEPARTMENT OF PHYSICS, UNIVERSITY OF SCIENCES AND TECHNOLOGY OF ORAN, EL MNAOUAR, BP 1505, 31000 ORAN, ALGERIA
E-mail address: fhaddouchi@gmail.com

SLIMANE BENAICHA, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ORAN, ES-SENIA, 31000 ORAN, ALGERIA
E-mail address: slimanebenaicha@yahoo.fr