

Inverse spectral problems for Dirac operators with summable matrix-valued potentials

D. V. Puyda*

Ivan Franko National University of Lviv
1 Universytetska str., Lviv, 79000, Ukraine

Abstract

We solve the direct and inverse spectral problems for Dirac operators on $(0, 1)$ with matrix-valued potentials whose entries belong to $L_p(0, 1)$, $p \in [1, \infty)$. We give a complete description of the spectral data (eigenvalues and suitably introduced norming matrices) for the operators under consideration and suggest a method for reconstructing the operator from the spectral data.

1 Introduction

The aim of the present paper is to solve the direct and inverse spectral problems for self-adjoint Dirac operators generated by the differential expressions

$$\mathfrak{t}_q := \frac{1}{i} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \frac{d}{dx} + \begin{pmatrix} 0 & q \\ q^* & 0 \end{pmatrix}$$

and some separated boundary conditions. Here q is an $r \times r$ matrix-valued function with entries belonging to $L_p(0, 1)$, $p \in [1, \infty)$, called the potential of the operator, and I is the identity $r \times r$ matrix. For such Dirac operators, we introduce the notion of the spectral data – eigenvalues and specially defined norming matrices. We then give a complete description of the class of the spectral data for the operators under consideration, show that the spectral data determine the operator uniquely and provide an efficient method for reconstructing the operator from the spectral data.

Direct and inverse spectral problems for Sturm–Liouville and Dirac operators [1, 2, 3] have been studied over the last 50 years. Already in 1966, M. Gasymov and B. Levitan suggested using the spectral function or the scattering phase for reconstructing Dirac operators on a half-line [4, 5]. Ever since, direct and inverse spectral problems for Dirac operators, including the systems of higher orders, have been considered in many papers. Among the recent investigations in that area we mention the ones by M. Malamud et al. [6, 7, 8], F. Gesztesy et al. [9, 10, 11], A. Sakhnovich [12, 13]. Problems similar to the ones considered in this paper were recently treated by D. Chelkak and E. Korotyaev for Sturm–Liouville operators with matrix-valued potentials [14, 15]. We refer the reader to the references in [6]–[15] for further results on the subject.

The direct and inverse spectral problems for Dirac operators with summable *scalar* potentials were solved in [16], where an algorithm for reconstructing the operator from two spectra or

**Email addresses:* dpuyda@gmail.com (D. V. Puyda)

from one spectrum and the norming constants was suggested. Later, using the technique that was suggested in [17] for solving the inverse spectral problems for Sturm–Liouville operators with matrix-valued potentials, the case of Dirac operators with square-integrable matrix-valued potentials was considered in [18]. Therein, a complete description of the spectral data was given and a method for reconstructing the operator from the spectral data was suggested.

In this paper we extend the results of [18] to the case of Dirac operators with summable matrix-valued potentials. However, as compared with the special case $q \in L_2$, it turns out that more general one $q \in L_p$, $p \geq 1$, meets some conceptual and technical difficulties.

In particular, the description of the spectral data in [18] involves a precise asymptotics of eigenvalues and norming matrices, which is not available for the operators with summable matrix-valued potentials. Namely, the proofs of the asymptotics in [18] were based on the results of [19, 20] that are applicable only for $p = 2$. To prove similar statements for $p \geq 1$ one may need analogues of the results of [21] covering matrix-valued functions, but to the best of the author’s knowledge these are not known yet. Furthermore, it turns out that we cannot separate the asymptotics of eigenvalues and the asymptotics of norming matrices under the present assumptions on the potential.

Instead, as opposed to the case $p = 2$, we formulate the description of the spectral data involving an auxiliary object which is the restriction of the Fourier transform of the spectral measure (see Theorem 1.1 and Definition 1.1 below and compare with Theorem 1.1 in [18]). The latter is easy to construct given the spectral data and turns out to contain all the information about the potential.

Further, as compared with the case $p = 2$, while dealing with $p \geq 1$ one also has to reconsider another condition involved in the description of the spectral data, i.e. the one on ranks of the norming matrices. Although the claim of the condition remains the same, technique of the proof that was used for $p = 2$ fails for $p \geq 1$. Instead, we introduce the approach based on the theory of Riesz bases. In particular, new our approach uses a vector analogue of well-known Kadec’s 1/4-theorem (see Appendix C), which plays an auxiliary role in this paper but may be used while solving other problems having to do with vector-valued functions.

From the practical point of view, a method for reconstructing the operator from the spectral data appears the same as for operators with square-integrable potentials and essentially consists in solving certain integral equation (see Theorem 1.4 and further comments on it). The paper is theoretical, but the results can be used in practical applications where the inverse spectral problems for Dirac operators with matrix-valued potentials arise. For instance, inverse problems for quantum graphs, which are of practical importance in nanotechnology, microelectronics, etc. (see, e.g., [22]) in some cases can be reduced to the inverse problems for operators with matrix-valued potentials. As compared with the case considered in [18], the results of the present paper allow stronger singularities of potentials which is important for applications.

The paper is organized as follows. In the reminder of this section we give a precise setting of the problem and formulate the main results. In Section 2 we provide some preliminaries: we introduce the Weyl–Titchmarsh function and establish the basic properties of the operators under consideration. In Sections 3 and 4, respectively, we solve the direct and inverse spectral problems. Several spaces used in this paper are introduced in Appendix A. In Appendix B we recall some facts on the factorization of integral operators. Finally, Appendix C is devoted to some auxiliary facts on the theory of Riesz bases.

1.1 Setting of the problem

Firstly, we introduce the space of potentials for Dirac operators considered in this paper. Let M_r denote the set of $r \times r$ matrices with complex entries, which we identify with the Banach

algebra of linear operators $\mathbb{C}^r \rightarrow \mathbb{C}^r$ endowed with the standard norm. We write $I = I_r$ for the unit element of M_r and M_r^+ for the set of all matrices $A \in M_r$ such that $A = A^* \geq 0$. We set

$$\mathfrak{Q}_p := L_p((0, 1), M_r), \quad p \in [1, \infty),$$

and endow \mathfrak{Q}_p with the norm $\|q\|_{\mathfrak{Q}_p} := \left(\int_0^1 \|q(s)\|^p ds \right)^{1/p}$, $q \in \mathfrak{Q}_p$. The space \mathfrak{Q}_p will serve as the *space of potentials* for Dirac operators under consideration.

Let $q \in \mathfrak{Q}_p$, $p \in [1, \infty)$, denote

$$\vartheta := \frac{1}{i} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \mathbf{q} := \begin{pmatrix} 0 & q \\ q^* & 0 \end{pmatrix}$$

and consider the differential expression

$$\mathbf{t}_q := \vartheta \frac{d}{dx} + \mathbf{q} \tag{1.1}$$

on the domain

$$D(\mathbf{t}_q) = \{y := (y_1, y_2)^\top \mid y_1, y_2 \in W_1^1((0, 1), \mathbb{C}^r)\},$$

where $W_1^1((0, 1), \mathbb{C}^r)$ is the Sobolev space. The object of our investigation is the *self-adjoint Dirac operator* T_q generated by the differential expression (1.1) and some separated boundary conditions:

$$T_q y = \mathbf{t}_q(y),$$

$$D(T_q) = \{y \in D(\mathbf{t}_q) \mid \mathbf{t}_q(y) \in L_2((0, 1), \mathbb{C}^{2r}), y_1(0) = y_2(0), y_1(1) = y_2(1)\}.$$

The function $q \in \mathfrak{Q}_p$ will be called the *potential* of the operator T_q .

The spectrum $\sigma(T_q)$ of the operator T_q consists of countably many isolated real eigenvalues of finite multiplicity, accumulating only at $+\infty$ and $-\infty$. We denote by $\lambda_j(q)$, $j \in \mathbb{Z}$, the pairwise distinct eigenvalues of the operator T_q labeled in increasing order so that $\lambda_0(q) \leq 0 < \lambda_1(q)$, i.e.

$$\sigma(T_q) = \{\lambda_j(q)\}_{j \in \mathbb{Z}}, \quad \lambda_0(q) \leq 0 < \lambda_1(q).$$

Further, denote by m_q the Weyl–Titchmarsh function of the operator T_q (see, e.g., [9]). The function m_q is a matrix-valued meromorphic Herglotz function (i.e. such that $\text{Im } m_q(\lambda) \geq 0$ whenever $\text{Im } \lambda > 0$), and $\{\lambda_j(q)\}_{j \in \mathbb{Z}}$ is the set of its poles. We set

$$\alpha_j(q) := - \operatorname{res}_{\lambda=\lambda_j(q)} m_q(\lambda), \quad j \in \mathbb{Z}, \tag{1.2}$$

and call $\alpha_j(q)$ the *norming matrix* of the operator T_q corresponding to the eigenvalue $\lambda_j(q)$. For every $j \in \mathbb{Z}$, the norming matrix $\alpha_j(q)$ is non-negative and multiplicity of the eigenvalue $\lambda_j(q)$ equals the rank of $\alpha_j(q)$.

The sequence $\mathbf{a}_q := ((\lambda_j(q), \alpha_j(q)))_{j \in \mathbb{Z}}$ will be called the *spectral data* of the operator T_q , and the matrix-valued measure

$$\mu_q := \sum_{j=-\infty}^{\infty} \alpha_j(q) \delta_{\lambda_j(q)} \tag{1.3}$$

will be called its *spectral measure*. Here δ_λ is the Dirac delta-measure centered at the point λ . In particular, if $q = 0$, then

$$\mu_0 = \sum_{n=-\infty}^{\infty} I \delta_{\pi n}. \tag{1.4}$$

We give a complete description of the class

$$\mathfrak{A}_p := \{\mathfrak{a}_q \mid q \in \mathfrak{Q}_p\}$$

of the spectral data for the operators under consideration, which is equivalent to the description of the class $\mathfrak{M}_p := \{\mu_q \mid q \in \mathfrak{Q}_p\}$ of the spectral measures. We then show that the spectral data of the operator T_q determine the potential q uniquely and suggest a method for reconstructing this potential from the spectral data.

1.2 Main results

We start from the description of the spectral data for the operators under consideration. In what follows, \mathfrak{a} will stand for an arbitrary sequence pretending to be the spectral data of some Dirac operator T_q , i.e. $\mathfrak{a} := ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}}$, where $(\lambda_j)_{j \in \mathbb{Z}}$ is a strictly increasing sequence of real numbers such that $\lambda_0 \leq 0 < \lambda_1$ and α_j , $j \in \mathbb{Z}$, are non-zero matrices in M_r^+ .

Given the sequence \mathfrak{a} , denote by $\mu^\mathfrak{a}$ a matrix-valued measure given by

$$\mu^\mathfrak{a} := \sum_{j=-\infty}^{\infty} \alpha_j \delta_{\lambda_j} \quad (1.5)$$

and pretending to be the spectral measure. With every measure $\mu := \mu^\mathfrak{a}$ of the form (1.5) we associate a \mathbb{C}^r -valued distribution

$$(\mu, f) := \int_{\mathbb{R}} f \, d\mu, \quad f \in \mathcal{S}^r,$$

where \mathcal{S}^r is the Schwartz space of rapidly decreasing \mathbb{C}^r -valued functions (see Appendix A). Now we introduce a kind of the Fourier transform of $\mu^\mathfrak{a}$:

Definition 1.1 *For an arbitrary measure $\mu := \mu^\mathfrak{a}$ of the form (1.5) we denote by $\widehat{\mu}$ a \mathbb{C}^r -valued distribution given by the formula*

$$(\widehat{\mu}, f) := (\mu, \widehat{f}), \quad f \in \mathcal{S}^r,$$

where

$$\widehat{f}(\lambda) := \int_{-\infty}^{\infty} e^{2i\lambda s} f(s) \, ds, \quad \lambda \in \mathbb{R}. \quad (1.6)$$

We denote by H_μ the restriction of the distribution $\widehat{\mu} - \widehat{\mu}_0$ to the interval $[-1, 1]$, i.e.

$$(H_\mu, f) := (\widehat{\mu} - \widehat{\mu}_0, f), \quad f \in \mathcal{S}^r, \quad \text{supp } f \subset [-1, 1], \quad (1.7)$$

where μ_0 given by (1.4) is the spectral measure of the free operator T_0 .

The distribution H_μ , $\mu := \mu^\mathfrak{a}$, plays an important role in establishing whether the sequence \mathfrak{a} is the spectral data of some operator T_q . Namely, we partition the real axis into pairwise disjoint intervals Δ_n , $n \in \mathbb{Z}$, by setting

$$\Delta_n := \left(\pi n - \frac{\pi}{2}, \pi n + \frac{\pi}{2} \right].$$

Then the following theorem gives a complete description of the class \mathfrak{A}_p of the spectral data for Dirac operators under consideration:

Theorem 1.1 *In order that a sequence $\mathbf{a} := ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}}$ should belong to \mathfrak{A}_p , $p \geq 1$, it is necessary and sufficient that the following conditions are satisfied:*

$$(B_1) \quad \sum_{\lambda_j \in \Delta_n} |\pi n - \lambda_j| = o(1) \text{ and } \|I - \sum_{\lambda_j \in \Delta_n} \alpha_j\| = o(1) \text{ as } |n| \rightarrow \infty;$$

$$(B_2) \quad \exists N_0 \in \mathbb{N} \quad \forall N \in \mathbb{N}: N > N_0 \Rightarrow \sum_{n=-N}^N \sum_{\lambda_j \in \Delta_n} \text{rank } \alpha_j = (2N + 1)r;$$

(B₃) *the system of functions $\{e^{i\lambda_j t} d \mid j \in \mathbb{Z}, d \in \text{Ran } \alpha_j\}$ is complete in $L_2((-1, 1), \mathbb{C}^r)$;*

(B₄) *the distribution H_μ , $\mu := \mu^{\mathbf{a}}$, belongs to $L_p((-1, 1), M_r)$.*

Remark 1.1 *Thus, in particular, the distribution H_μ , $\mu := \mu^{\mathbf{a}}$, turns out to be regular for all $\mathbf{a} \in \mathfrak{A}_p$. In this case, $H_\mu(x)$ can be formally defined by the formula*

$$H_\mu(x) = \sum_{n \in \mathbb{Z}} \left(\sum_{\lambda_j \in \Delta_n} e^{2i\lambda_j x} \alpha_j - e^{2i\pi n x} I \right), \quad x \in (-1, 1), \quad (1.8)$$

but the convergence of this series in $L_p((-1, 1), M_r)$, $p \geq 1$, is difficult to establish without knowing the precise asymptotics of eigenvalues and norming matrices. The proof for the case $p = 2$ is given in [18].

Further, it turns out that the spectral data of the operator T_q determine the potential q uniquely:

Theorem 1.2 *For all $p \geq 1$, the mapping $\mathfrak{Q}_p \ni q \mapsto \mathbf{a}_q \in \mathfrak{A}_p$ is bijective.*

This allows a possibility to reconstruct the operator from the spectral data.

As in [17, 18], we base our reconstruction algorithm on Krein's accelerant method:

Definition 1.2 *We say that a function $H \in L_1((-1, 1), M_r)$ is an accelerant if $H(-x) = H(x)^*$ a.e. on $(-1, 1)$ and for every $a \in (0, 1]$ the integral equation*

$$f(x) + \int_0^a H(x-t)f(t) dt = 0, \quad x \in (0, a),$$

has only zero solution in $L_2((0, a), \mathbb{C}^r)$. We denote by \mathfrak{H}_p , $p \geq 1$, the set of all accelerants belonging to $L_p((-1, 1), M_r)$ and endow \mathfrak{H}_p with the metric of the space $L_p((-1, 1), M_r)$.

Equivalently, it is known (see, e.g., [16]) that an arbitrary function $H \in L_p((-1, 1), M_r)$, $p \geq 1$, belongs to \mathfrak{H}_p if and only if the Krein equation

$$R(x, t) + H(x-t) + \int_0^x R(x, s)H(x-s) ds = 0, \quad (x, t) \in \Omega, \quad (1.9)$$

where $\Omega := \{(x, t) \mid 0 \leq t \leq x \leq 1\}$, is solvable in $G_p^+(M_r)$ (see Appendix A). In this case, a solution of (1.9) is unique and we denote it by $R_H(x, t)$, $(x, t) \in \Omega$.

We define the *Krein mapping* $\Theta : \mathfrak{H}_1 \rightarrow \mathfrak{Q}_1$ by the formula

$$[\Theta(H)](x) := iR_H(x, 0), \quad x \in (0, 1). \quad (1.10)$$

This mapping provides a one-to-one correspondence between accelerants $H \in \mathfrak{H}_p$ and potentials $q \in \mathfrak{Q}_p$:

Theorem 1.3 For all $p \geq 1$, the Krein mapping Θ is a homeomorphism between the metric spaces \mathfrak{H}_p and \mathfrak{Q}_p .

We use the Krein mapping to reconstruct the potential q from the spectral data of the operator T_q :

Theorem 1.4 Given an arbitrary sequence $\mathbf{a} \in \mathfrak{A}_p$, set $\mu := \mu^{\mathbf{a}}$ and $H := H_\mu$. Then $H \in \mathfrak{H}_p$ and $\mathbf{a} = \mathbf{a}_q$ for $q = \Theta(H)$.

Therefore, Theorems 1.2 and 1.4 provide an efficient method for reconstructing the Dirac operator T_q from the spectral data. Namely, given an arbitrary sequence $\mathbf{a} \in \mathfrak{A}_p$ being the spectral data of some Dirac operator T_q , construct the matrix-valued measure $\mu := \mu^{\mathbf{a}}$ by the formula (1.5); set $H := H_\mu$ by the formula (1.7) or (1.8); substitute H into the Krein equation (1.9) and find R_H ; find the potential $q = \Theta(H)$ by the formula (1.10). That the potential q is the one looked for follows from the fact that the Dirac operator T_q has the spectral data \mathbf{a} we have started with.

The method can be visualized by means of the following diagram:

$$\mathfrak{A}_p \ni \mathbf{a} \xrightarrow[s_1]{(1.5)} \mu := \mu^{\mathbf{a}} \xrightarrow[s_2]{(1.7)} H := H_\mu \xrightarrow[s_3]{(1.10)} q = \Theta(H) \in \mathfrak{Q}_p.$$

Here s_j denotes the step number j . The steps s_1 and s_2 are trivial. The basic and non-trivial step is s_3 , which requires solving the Krein equation (1.9).

Remark 1.2 One can similarly consider the Dirac operators with general separated boundary conditions. However, the description of the spectral data would be more complicated since the spectrum of the free operator T_0 (subject to $q = 0$) is more involved in this case. The author plans to consider the operators with general (not necessarily separated) boundary conditions in a forthcoming paper.

2 Preliminary results

Here we introduce the Weyl–Titchmarsh function and establish the basic properties of the operator T_q . The material of this section mainly follows [18] but forms the essential base for further considerations.

2.1 The Weyl–Titchmarsh function of the operator T_q

We start from constructing the Weyl–Titchmarsh function of the operator T_q .

Let $q \in \mathfrak{Q}_p$, $p \geq 1$. Set

$$a := \frac{1}{\sqrt{2}} \begin{pmatrix} I & -I \end{pmatrix}$$

so that the boundary conditions $y_1(0) = y_2(0)$, $y_1(1) = y_2(1)$ can be written in the form

$$ay(0) = ay(1) = 0.$$

The multiplier $\frac{1}{\sqrt{2}}$ provides the normalization $aa^* = I$.

Denote by $u_q(\cdot, \lambda) \in W_1^1((0, 1), M_{2r})$, $\lambda \in \mathbb{C}$, a matrix-valued solution of the Cauchy problem

$$\vartheta \frac{d}{dx} u + \mathbf{q}u = \lambda u, \quad u(0, \lambda) = I_{2r},$$

and set

$$\varphi_q(x, \lambda) := u_q(x, \lambda)\vartheta a^*, \quad \psi_q(x, \lambda) := u_q(x, \lambda)a^* \quad (2.1)$$

so that the $2r \times r$ matrix-valued functions $\varphi_q(x, \lambda)$ and $\psi_q(x, \lambda)$ solve the Cauchy problems

$$\vartheta \frac{d}{dx} \varphi + \mathbf{q} \varphi = \lambda \varphi, \quad \varphi(0, \lambda) = \vartheta a^*, \quad (2.2)$$

and

$$\vartheta \frac{d}{dx} \psi + \mathbf{q} \psi = \lambda \psi, \quad \psi(0, \lambda) = a^*,$$

respectively.

Next, define the $r \times r$ matrix-valued functions $s_q(\lambda)$ and $c_q(\lambda)$ by the formulae

$$s_q(\lambda) := a\varphi_q(1, \lambda), \quad c_q(\lambda) := a\psi_q(1, \lambda). \quad (2.3)$$

Then the function

$$m_q(\lambda) := -s_q(\lambda)^{-1}c_q(\lambda) \quad (2.4)$$

is called the *Weyl–Titchmarsh function* of the operator T_q .

Repeating the proofs in [18], which were done for the case of square-integrable potential, one can prove the following lemma claiming the basic properties of just introduced objects:

Lemma 2.1 (i) For every $q \in \mathfrak{Q}_p$, $p \geq 1$, there exists a unique matrix-valued function $K_q \in G_p^+(M_{2r})$ (see Appendix A) such that for all $\lambda \in \mathbb{C}$ and $x \in (0, 1)$,

$$\varphi_q(x, \lambda) = \varphi_0(x, \lambda) + \int_0^x K_q(x, s)\varphi_0(s, \lambda) ds, \quad (2.5)$$

where

$$\varphi_0(x, \lambda) = \frac{1}{\sqrt{2i}} \begin{pmatrix} e^{i\lambda x} I \\ e^{-i\lambda x} I \end{pmatrix}$$

is a solution of (2.2) in the free case $\mathbf{q} = 0$; moreover, the mapping $\mathfrak{Q}_p \ni q \mapsto K_q \in G_p^+(M_{2r})$ is continuous;

(ii) the functions $s_q(\lambda)$ and $c_q(\lambda)$ are entire and allow the representations

$$s_q(\lambda) = (\sin \lambda)I + \int_{-1}^1 e^{i\lambda t} g_q(t) dt, \quad (2.6)$$

$$c_q(\lambda) = (\cos \lambda)I + \int_{-1}^1 e^{i\lambda t} h_q(t) dt, \quad (2.7)$$

where g_q and h_q are some (depending on q) functions in $L_p((-1, 1), M_r)$; moreover, the mappings $\mathfrak{Q}_p \ni q \mapsto g_q \in L_p((-1, 1), M_r)$ and $\mathfrak{Q}_p \ni q \mapsto h_q \in L_p((-1, 1), M_r)$ are continuous;

(iii) the operator functions $\lambda \mapsto s_q(\lambda)^{-1}$ and $\lambda \mapsto m_q(\lambda)$ are meromorphic in \mathbb{C} ; moreover, $m_0(\lambda) = -(\cot \lambda)I$ and

$$\|m_q(\lambda) + (\cot \lambda)I\| = o(1) \quad (2.8)$$

as $\lambda \rightarrow \infty$ within the domain $\mathcal{O} = \{z \in \mathbb{C} \mid \forall n \in \mathbb{Z} \ |z - \pi n| > 1\}$.

Sketch of the proof. Repeating the proof in [18] and using the results of [23], one can show that there exist unique matrix-valued functions $P^+, P^- \in G_p^+(M_{2r})$ such that for all $x \in (0, 1)$ and $\lambda \in \mathbb{C}$,

$$u_q(x, \lambda) = e^{-\lambda x \vartheta} + \int_0^x P^+(x, s) e^{-\lambda(x-2s)\vartheta} ds + \int_0^x P^-(x, s) e^{\lambda(x-2s)\vartheta} ds.$$

Then, by virtue of definitions (2.1) and (2.3), straightforward calculations lead us to the representations (2.5), (2.6) and (2.7). Moreover, since the mappings $\mathfrak{Q}_p \ni q \mapsto P^\pm \in G_p^+(M_{2r})$ are continuous (see [23]), the mappings $\mathfrak{Q}_p \ni q \mapsto K_q \in G_p^+(M_{2r})$ and $\mathfrak{Q}_p \ni q \mapsto g_q, h_q \in L_p((-1, 1), M_r)$ remain continuous. Thus we obtain parts (i) and (ii) of the present lemma.

To prove part (iii), observe that by virtue of the representations (2.6), (2.7) and Lemma A.2 we have

$$\lim_{|\lambda| \rightarrow \infty} e^{-|\operatorname{Im} \lambda|} \|s_q(\lambda) - (\sin \lambda)I\| = \lim_{|\lambda| \rightarrow \infty} e^{-|\operatorname{Im} \lambda|} \|c_q(\lambda) - (\cos \lambda)I\| = 0.$$

Therefore, $s_q(\lambda)$ is invertible for all $\lambda \in \mathcal{O}$ large enough, so that m_q is meromorphic and (2.8) holds true. \square

It will follow that poles of the Weyl–Titchmarsh function m_q are eigenvalues of the Dirac operator T_q . Given also the corresponding residues of m_q , it is possible to find the potential q .

2.2 Basic properties of the operator T_q

Here we establish the basic properties of the operator T_q . We set

$$\mathbb{H} := L_2((0, 1), \mathbb{C}^r) \times L_2((0, 1), \mathbb{C}^r)$$

and denote by \mathcal{I} the identity operator in \mathbb{H} . For an arbitrary $q \in \mathfrak{Q}_p$, $p \geq 1$, and $\lambda \in \mathbb{C}$, denote by $\Phi_q(\lambda)$ the operator acting from \mathbb{C}^r to \mathbb{H} by the formula

$$[\Phi_q(\lambda)c](x) := \varphi_q(x, \lambda)c.$$

Taking into account (2.5), we obtain that for all $\lambda \in \mathbb{C}$,

$$\Phi_q(\lambda) = (\mathcal{I} + \mathcal{K}_q)\Phi_0(\lambda), \tag{2.9}$$

where \mathcal{K}_q is an integral operator in \mathbb{H} with kernel K_q (see Lemma 2.1, part (i)) and

$$[\Phi_0(\lambda)c](x) = \frac{1}{\sqrt{2i}} \begin{pmatrix} e^{i\lambda x} I \\ e^{-i\lambda x} I \end{pmatrix} c. \tag{2.10}$$

The following lemma claims basic properties of the operators $\Phi_q(\lambda)$. Particularly, the second part of the lemma is important:

Lemma 2.2 *For all $q \in \mathfrak{Q}_p$, $p \geq 1$, and $\lambda \in \mathbb{C}$,*

$$(i) \quad \ker \Phi_q(\lambda) = \{0\}, \quad \operatorname{Ran} \Phi_q^*(\lambda) = \mathbb{C}^r, \tag{2.11}$$

where $\Phi_q^*(\lambda) := [\Phi_q(\lambda)]^*$;

$$(ii) \quad \ker(T_q - \lambda \mathcal{I}) = \Phi_q(\lambda) \ker s_q(\lambda). \tag{2.12}$$

Proof. Since the operator $\mathcal{S} + \mathcal{K}_q$ is a homeomorphism of the space \mathbb{H} , part (i) easily follows from (2.9) and (2.10).

To prove part (ii), note that for all $c \in \ker s_q(\lambda)$ the function $f(x) := \varphi_q(x, \lambda)c$ verifies the equality $\vartheta f' + \mathbf{q}f = \lambda f$ and the boundary conditions $af(0) = af(1) = 0$. Conversely, the generic solution of the problem

$$\vartheta f' + \mathbf{q}f = \lambda f, \quad af(0) = 0,$$

takes the form $f(x) = \varphi_q(x, \lambda)c$, $c \in \mathbb{C}^r$, while the boundary condition $af(1) = 0$ means $c \in \ker s_q(\lambda)$. Therefore, the equality (2.12) follows. \square

Now, denote by $\lambda_j := \lambda_j(q)$, $j \in \mathbb{Z}$, the pairwise distinct eigenvalues of the operator T_q labeled in increasing order so that $\lambda_0 \leq 0 < \lambda_1$, and let $\alpha_j := \alpha_j(q)$ be the corresponding norming matrices defined by (1.2). Then the basic properties of the operator T_q are formulated in the following theorem:

Theorem 2.1 *Let $q \in \mathfrak{Q}_p$, $p \geq 1$. Then the following statements hold true:*

(i) *the operator T_q is self-adjoint;*

(ii) *the spectrum $\sigma(T_q)$ of the operator T_q consists of isolated real eigenvalues of finite multiplicity and, moreover,*

$$\sigma(T_q) = \{\lambda \in \mathbb{C} \mid \ker s_q(\lambda) \neq \{0\}\};$$

(iii) *denote by $P_{q,j}$ the orthogonal projector onto $\ker(T_q - \lambda_j \mathcal{S})$; then*

$$\sum_{j=-\infty}^{\infty} P_{q,j} = \mathcal{S}; \tag{2.13}$$

(iv) *for every $j \in \mathbb{Z}$ the norming matrix α_j is non-negative and*

$$P_{q,j} = \Phi_q(\lambda_j)\alpha_j\Phi_q^*(\lambda_j). \tag{2.14}$$

The proof of Theorem 2.1 repeats the proof in [18]. In particular, part (ii) together with (2.4) implies that eigenvalues of the operator T_q are poles of the Weyl–Titchmarsh function m_q .

By virtue of the relations (2.9), (2.13) and (2.14), the operators $\Phi_q(\lambda)$ and the function K_q from Lemma 2.1 will play an important role in this investigation.

3 Direct spectral problem

Here we prove the necessity part of Theorem 1.1: we show that for an arbitrary potential $q \in \mathfrak{Q}_p$, $p \geq 1$, the spectral data of the operator T_q satisfy the conditions (B_1) – (B_4) . Throughout this section we use the abbreviations $\lambda_j := \lambda_j(q)$ and $\alpha_j := \alpha_j(q)$ for eigenvalues and norming matrices of the operator T_q , respectively.

3.1 Condition (B_1)

In this subsection we prove the following proposition:

Proposition 3.1 For an arbitrary potential $q \in \mathfrak{Q}_p$, $p \geq 1$, the spectral data \mathbf{a}_q of the operator T_q satisfy the condition (B_1) , i.e. the following asymptotics hold true:

$$\sum_{\lambda_j \in \Delta_n} |\pi n - \lambda_j| = o(1), \quad |n| \rightarrow \infty, \quad (3.1)$$

and

$$\left\| I - \sum_{\lambda_j \in \Delta_n} \alpha_j \right\| = o(1), \quad |n| \rightarrow \infty, \quad (3.2)$$

where $\Delta_n := \left(\pi n - \frac{\pi}{2}, \pi n + \frac{\pi}{2} \right]$.

As in [18], the proof of Proposition 3.1 is based on the claim that eigenvalues of the operator T_q are zeros of certain entire function. However, as opposed to the case of square-integrable potential, we can establish only the rough asymptotics of eigenvalues and norming matrices. Thus, we start the proof of Proposition 3.1 from making the following remark:

Remark 3.1 As follows from Theorem 2.1, part (ii), the eigenvalues $(\lambda_j)_{j \in \mathbb{Z}}$ of the operator T_q are zeros of the entire function

$$\tilde{s}_q(\lambda) := \det s_q(\lambda). \quad (3.3)$$

Since the function $s_q(\lambda)$ allows a representation (2.6), repeating the proofs in [19] one can use Rouché's theorem to show that the set of zeros of the function $\tilde{s}_q(\lambda)$ can be indexed (counting multiplicities) by numbers $n \in \mathbb{Z}$ so that the corresponding sequence $(\xi_n)_{n \in \mathbb{Z}}$ has the asymptotics

$$\xi_{nr+j} = \pi n + o(1), \quad j = 1, \dots, r, \quad |n| \rightarrow \infty. \quad (3.4)$$

Further, it also follows that the set of zeros of the entire function

$$\tilde{c}_q(\lambda) := \det c_q(\lambda)$$

can be indexed (counting multiplicities) by numbers $n \in \mathbb{Z}$ so that the corresponding sequence $(\zeta_n)_{n \in \mathbb{Z}}$ has the asymptotics

$$\zeta_{nr+j} = \pi \left(n + \frac{1}{2} \right) + o(1), \quad j = 1, \dots, r, \quad |n| \rightarrow \infty. \quad (3.5)$$

Now the proof of Proposition 3.1 is straightforward:

Proof of Proposition 3.1. Since, by Remark 3.1, eigenvalues $(\lambda_j)_{j \in \mathbb{Z}}$ of the operator T_q are zeros of the entire function $\tilde{s}_q(\lambda)$, the asymptotics (3.1) directly follow from (3.4). Thus it only remains to prove (3.2).

For $n \in \mathbb{Z}$ denote

$$\beta_n := I - \sum_{\lambda_j \in \Delta_n} \alpha_j.$$

It follows from the definition (1.2) of α_j and from the asymptotics (3.1) of $(\lambda_j)_{j \in \mathbb{Z}}$ that there exists $n_0 \in \mathbb{N}$ such that for all $n \in \mathbb{Z}$, $|n| > n_0$,

$$\sum_{\lambda_j \in \Delta_n} \alpha_j = -\frac{1}{2\pi i} \oint_{|\lambda - \pi n| = 1} m_q(\lambda) \, d\lambda.$$

Therefore, for all $n \in \mathbb{Z}$, $|n| > n_0$,

$$\beta_n = \frac{1}{2\pi i} \oint_{|\lambda - \pi n| = 1} (m_q(\lambda) + (\cot \lambda)I) \, d\lambda.$$

Now taking into account (2.8), we observe that $\|\beta_n\| = o(1)$, $|n| \rightarrow \infty$, as desired. \square

3.2 Conditions (B_2) and (B_3)

In this subsection we prove that the spectral data for the operators under consideration satisfy the conditions (B_2) and (B_3) . We start from proving the condition (B_2) :

Proposition 3.2 *For an arbitrary potential $q \in \mathfrak{Q}_p$, $p \geq 1$, the spectral data \mathfrak{a}_q of the operator T_q satisfy the condition (B_2) , i.e. there exists $N_0 \in \mathbb{N}$ such that for all natural $N > N_0$,*

$$\sum_{n=-N}^N \sum_{\lambda_j \in \Delta_n} \text{rank } \alpha_j = (2N + 1)r. \quad (3.6)$$

The proof of Proposition 3.2 involves a technique based on the theory of Riesz bases. However, before starting the essential part of the proof we have to establish some auxiliary results. Namely, recalling Remark 3.1 claiming that eigenvalues $(\lambda_j)_{j \in \mathbb{Z}}$ of the operator T_q are zeros of the entire function $\tilde{s}_q(\lambda)$, we need to prove the following lemma:

Lemma 3.1 *Let n_j , $j \in \mathbb{Z}$, denote the multiplicity of zero λ_j of the function $\tilde{s}_q(\lambda)$. Then there exists $N_1 \in \mathbb{N}$ such that for all $\lambda_j \in \Delta_n$, $|n| > N_1$,*

$$n_j = \text{rank } \alpha_j. \quad (3.7)$$

Proof. Firstly, observe that by virtue of the relations (2.11), (2.12) and (2.14), we have

$$\dim \ker s_q(\lambda_j) = \text{rank } \alpha_j, \quad j \in \mathbb{Z}.$$

Since

$$s_q(\lambda) = s_q(\lambda_j) + O(\lambda - \lambda_j), \quad \lambda \rightarrow \lambda_j,$$

one can easily find that

$$n_j \geq \text{rank } \alpha_j, \quad j \in \mathbb{Z}. \quad (3.8)$$

Now let us show that there exists $N_1 \in \mathbb{N}$ such that for all $\lambda_j \in \Delta_n$, $|n| > N_1$,

$$n_j \leq \text{rank } \alpha_j. \quad (3.9)$$

Indeed, since the function $m_q(\lambda)$ is meromorphic in \mathbb{C} and has a pole of first order at point λ_j (see, e.g., [25]), we find that

$$m_q(\lambda) = \frac{\alpha_j}{\lambda - \lambda_j} + g_j(\lambda), \quad (3.10)$$

where the function $g_j(\lambda)$ is analytic in the neighborhood of λ_j . Denote by Q_j an orthogonal projector onto $\text{Ran } \alpha_j$ and set

$$\tilde{Q}_j(\lambda) := (I - Q_j) + (\lambda - \lambda_j)Q_j.$$

Then it follows from (3.10) that the function

$$\lambda \mapsto \tilde{Q}_j(\lambda)m_q(\lambda)$$

is bounded in the neighborhood of λ_j . Furthermore, observe that by virtue of the asymptotics (3.5) of zeros of $\det c_q(\lambda)$, the function $\lambda \mapsto c_q(\lambda)^{-1}$ is analytic in the neighborhood of λ_j for large values of $|\lambda_j|$.

Now, since $m_q(\lambda) = -s_q(\lambda)^{-1}c_q(\lambda)$ (see (2.4)), we find that the function

$$\tilde{Q}_j(\lambda)s_q(\lambda)^{-1} = -\tilde{Q}_j(\lambda)m_q(\lambda)c_q(\lambda)^{-1}$$

is bounded in the neighborhood of λ_j for large values of $|\lambda_j|$. Therefore, since

$$\det \tilde{Q}_j(\lambda)s_q(\lambda)^{-1} = \frac{(\lambda - \lambda_j)^{\text{rank } \alpha_j}}{\tilde{s}_q(\lambda)},$$

we observe that there exists $N_1 \in \mathbb{N}$ such that for all $\lambda_j \in \Delta_n$, $|n| > N_1$, the inequality (3.9) holds true. Together with (3.8), this proves the lemma. \square

Now, recalling the asymptotics (3.4) of zeros of $\tilde{s}_q(\lambda)$, we arrive at the following corollary which was the purpose of Lemma 3.1:

Corollary 3.1 *There exists $N_2 \in \mathbb{N}$ such that for all $n \in \mathbb{Z}$, $|n| > N_2$,*

$$\sum_{\lambda_j \in \Delta_n} \text{rank } \alpha_j = r.$$

Now we can proceed to the principal part of the proof involving the theory of Riesz bases. The approach is based on Lemma 3.2 below, which will be used in solving both the direct and inverse spectral problems. The proof of the Lemma 3.2 uses a vector analogue of well-known Kadec's 1/4-theorem which is established in Appendix C.

Thus, let $\mathbf{a} := ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}}$ be an *arbitrary* sequence where $(\lambda_j)_{j \in \mathbb{Z}}$ is a strictly increasing sequence of real numbers such that $\lambda_0 \leq 0 < \lambda_1$ and α_j , $j \in \mathbb{Z}$, are non-zero matrices in M_r^+ . Since the matrices α_j , $j \in \mathbb{Z}$, are self-adjoint and non-negative, for every $j \in \mathbb{Z}$ there exists a system of pairwise orthogonal vectors $\{v_{j,k}\}_{k=1}^{\text{rank } \alpha_j}$ in \mathbb{C}^r such that

$$\alpha_j = \sum_{k=1}^{\text{rank } \alpha_j} (\cdot | v_{j,k}) v_{j,k}. \quad (3.11)$$

Denote by ϵ_k , $k = 1, \dots, r$, a standard orthonormal basis for \mathbb{C}^r . Then the following lemma holds true:

Lemma 3.2 *Let $\mathbf{a} := ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}}$ be an arbitrary sequence satisfying the asymptotics (3.1) and (3.2), and assume that there exists $N_0 \in \mathbb{N}$ such that for all $n \in \mathbb{Z}$, $|n| > N_0$,*

$$\sum_{\lambda_j \in \Delta_n} \text{rank } \alpha_j = r. \quad (3.12)$$

Then there exists $N_1 \in \mathbb{N}$ such that for all natural $N > N_1$ the system $\mathcal{E}_N \cup \mathcal{B}_N$, where

$$\mathcal{E}_N := \left\{ \frac{1}{\sqrt{2}} e^{i\pi n t} \epsilon_s \mid n \in \mathbb{Z}, |n| \leq N, \quad s = 1, \dots, r \right\} \quad (3.13)$$

and

$$\mathcal{B}_N := \{ e^{i\lambda_j t} v_{j,k} \mid \lambda_j \in \Delta_n, |n| > N, \quad 1 \leq k \leq \text{rank } \alpha_j \}, \quad (3.14)$$

is a Riesz basis for the space $\mathcal{H} := L_2((-1, 1), \mathbb{C}^r)$ (see Appendix C).

Proof. We prove the lemma by applying Theorem C.1. As follows from the equality (3.12), for all $n \in \mathbb{Z}$, $|n| > N_0$, the system

$$V_n := \{v_{j,k} \mid \lambda_j \in \Delta_n, 1 \leq k \leq \text{rank } \alpha_j\}$$

consists of exactly r vectors. Moreover, since

$$\|I - \beta_n\| = o(1), \quad |n| \rightarrow \infty, \quad (3.15)$$

where

$$\beta_n := \sum_{\lambda_j \in \Delta_n} \alpha_j, \quad (3.16)$$

we claim that N_0 can be taken so large that for all $n \in \mathbb{Z}$, $|n| > N_0$, the system V_n forms a basis for the space \mathbb{C}^r .

For $n \in \mathbb{Z}$, $|n| > N_0$, we denote by A_n the operator acting in \mathbb{C}^r by the formula

$$A_n v_{j,k} = \lambda_j v_{j,k}, \quad v_{j,k} \in V_n, \quad \lambda_j \in \Delta_n,$$

and set B_n to be an operator transforming a standard orthonormal basis for \mathbb{C}^r into the basis V_n . It follows from the asymptotics (3.1) and (3.2) that there exists a natural $N_1 > N_0$ such that

$$\|A_n - \pi n I\| < \ln 2, \quad |n| > N_1. \quad (3.17)$$

Moreover, let us show that

$$\sup_{|n| > N_1} (\|B_n\| + \|B_n^{-1}\|) < \infty. \quad (3.18)$$

Indeed, since

$$\sum_{v_{j,k} \in V_n} (\cdot |v_{j,k}) v_{j,k} = \beta_n,$$

where β_n is self-adjoint non-negative matrix given by (3.16), we observe that for all $|n| > N_1$ the vectors $\beta_n^{-1/2} v_{j,k}$, where $v_{j,k} \in V_n$, form an orthonormal basis for \mathbb{C}^r . Therefore, $B_n = \beta_n^{1/2} U_n$, where U_n is a unitary matrix in M_r . Since, by virtue of (3.15), we have

$$\sup_{|n| > N_1} (\|\beta_n^{1/2}\| + \|\beta_n^{-1/2}\|) < \infty,$$

(3.18) follows.

Therefore, taking into account (3.17) and (3.18), we find from Theorem C.1 that for all natural $N > N_1$ the system $\mathcal{E}_N \cup \mathcal{B}_N$ forms a Riesz basis for the space \mathcal{H} , as desired. \square

Now we use Lemma 3.2 and Corollary 3.1 to prove Proposition 3.2:

Proof of Proposition 3.2. Since the norming matrices α_j , $j \in \mathbb{Z}$, are self-adjoint and non-negative, for every $j \in \mathbb{Z}$ there exists a system of pairwise orthogonal vectors $\{v_{j,k}\}_{k=1}^{\text{rank } \alpha_j}$ in \mathbb{C}^r such that (3.11) holds true.

Observe that by virtue of Proposition 3.1 and Corollary 3.1, the conditions of Lemma 3.2 are satisfied. Hence, it follows from Lemma 3.2 that there exists $N_1 \in \mathbb{N}$ such that for all natural $N > N_1$, the system $\mathcal{E}_N \cup \mathcal{B}_N$ (see (3.13) and (3.14)) is a Riesz basis for the space $\mathcal{H} := L_2((-1, 1), \mathbb{C}^r)$. Therefore, Proposition 3.2 will be proved if we show that the system

$$\mathcal{B}_0 := \{e^{i\lambda_j t} v_{j,k} \mid j \in \mathbb{Z}, 1 \leq k \leq \text{rank } \alpha_j\} \quad (3.19)$$

also forms a basis for \mathcal{H} . Indeed, since $\mathcal{B}_N \subset \mathcal{B}_0$ and

$$\mathcal{B}_0 \setminus \mathcal{B}_N = \{e^{i\lambda_j t} v_{j,k} \mid \lambda_j \in \Delta_n, |n| \leq N, 1 \leq k \leq \text{rank } \alpha_j\},$$

we then obtain that for all natural $N > N_1$, the finite systems $\mathcal{B}_0 \setminus \mathcal{B}_N$ and \mathcal{E}_N consist of the same number of elements. Obviously, this implies the claim of Proposition 3.2.

Thus let us prove that \mathcal{B}_0 is a basis for \mathcal{H} . Indeed, since the operators

$$P_{q,j} := \Phi_q(\lambda_j) \alpha_j \Phi_q^*(\lambda_j), \quad j \in \mathbb{Z},$$

form a system of pairwise orthogonal projectors in the space \mathbb{H} (see Theorem 2.1) and

$$\sum_{j=-\infty}^{\infty} P_{q,j} = \mathcal{I},$$

we find that the system

$$\mathcal{A}_q := \{\Phi_q(\lambda_j) v_{j,k} \mid j \in \mathbb{Z}, 1 \leq k \leq \text{rank } \alpha_j\}$$

is a basis for the space \mathbb{H} . Since $\Phi_q(\lambda) = (\mathcal{I} + \mathcal{K}_q) \Phi_0(\lambda)$ and the mapping $\mathcal{I} + \mathcal{K}_q$ is a homeomorphism of \mathbb{H} , we observe that the system

$$\mathcal{A}_0 := \{\Phi_0(\lambda_j) v_{j,k} \mid j \in \mathbb{Z}, 1 \leq k \leq \text{rank } \alpha_j\}$$

remains a basis for \mathbb{H} . Now introduce the unitary mapping $V : \mathcal{H} \rightarrow \mathbb{H}$ acting by the formula

$$(Vf)(t) = (f(-t), f(t))^\top, \quad t \in (0, 1),$$

and note that V maps the function $e^{i\lambda_j t} v_{j,k}$ to $\Phi_0(\lambda_j) v_{j,k}$ for all $j \in \mathbb{Z}$, $1 \leq k \leq \text{rank } \alpha_j$. Therefore, the system \mathcal{B}_0 is a basis for the space \mathcal{H} , as desired. \square

In particular, from the proof of Proposition 3.2 we also obtain the following:

Corollary 3.2 *For an arbitrary potential $q \in \mathfrak{Q}_p$, $p \geq 1$, the spectral data \mathfrak{a}_q of the operator T_q satisfy the condition (B_3) , i.e. the system*

$$\mathcal{X} := \{e^{i\lambda_j t} d \mid j \in \mathbb{Z}, d \in \text{Ran } \alpha_j\} \tag{3.20}$$

is complete in $L_2((-1, 1), \mathbb{C}^r)$.

Proof. It follows from the proof of Proposition 3.2 that the system \mathcal{B}_0 given by (3.19) is a basis for the space $\mathcal{H} := L_2((-1, 1), \mathbb{C}^r)$. Therefore, we immediately obtain that $\mathcal{X}^\perp = (\text{lin } \mathcal{B}_0)^\perp = \{0\}$, as desired. Here $\text{lin } \mathcal{B}_0$ denotes the linear span of \mathcal{B}_0 . \square

3.3 The Krein mapping. Proof of Theorem 1.3

Thus, it remains only to prove that the spectral data for the operators under consideration satisfy the condition (B_4) . In order to do this, we need to look into some properties of the Krein mapping Θ given by (1.10). In particular, here we prove Theorem 1.3 claiming that for all $p \geq 1$ the mapping Θ is a homeomorphism between the space \mathfrak{H}_p of accelerants and the space \mathfrak{Q}_p of potentials.

We start from the following remark:

Remark 3.2 It obviously follows from the results of Appendix B that for all $p \geq 1$, the mapping $H \mapsto \Theta(H) := iR_H(\cdot, 0)$ acts continuously from \mathfrak{H}_p to \mathfrak{Q}_p .

Now, for an arbitrary accelerant $H \in \mathfrak{H}_p$ denote by \mathcal{F}_H a self-adjoint integral operator in \mathbb{H} with kernel

$$F_H(x, t) := \frac{1}{2} \begin{pmatrix} H\left(\frac{x-t}{2}\right) & H\left(\frac{x+t}{2}\right) \\ H\left(-\frac{x-t}{2}\right) & H\left(-\frac{x+t}{2}\right) \end{pmatrix}, \quad 0 \leq x, t \leq 1. \quad (3.21)$$

Then the following proposition explains a natural connection between an accelerant H and the corresponding potential $q := \Theta(H)$:

Proposition 3.3 For an arbitrary accelerant $H \in \mathfrak{H}_p$,

(i) there exists a unique function $L_H \in G_p^+(M_{2r})$ such that

$$\mathcal{I} + \mathcal{F}_H = (\mathcal{I} + \mathcal{L}_H)^{-1}(\mathcal{I} + \mathcal{L}_H^*)^{-1}, \quad (3.22)$$

where $\mathcal{L}_H \in \mathcal{G}_p^+(M_{2r})$ is an integral operator in \mathbb{H} with kernel L_H ;

(ii) $L_H = K_q$ for $q = \Theta(H)$.

Proof. Part (i) of the present proposition directly follows from Lemma B.1. To prove part (ii), note that for all $p \geq 1$ the set \mathfrak{H}_p is open in $L_p((-1, 1), M_r)$, and thus the set $\mathfrak{H}_1 \cap C([-1, 1], M_r)$ of continuous accelerants is dense everywhere in \mathfrak{H}_p . Therefore, there exists a sequence $(H_n)_{n \in \mathbb{N}}$, $H_n \in \mathfrak{H}_1 \cap C([-1, 1], M_r)$, such that

$$\lim_{n \rightarrow \infty} \|H - H_n\|_{\mathfrak{H}_p} = 0.$$

Then it follows from [18] that for all $n \in \mathbb{N}$,

$$L_{H_n} = K_{q_n}, \quad q_n = \Theta(H_n). \quad (3.23)$$

Now, by virtue of Theorem B.2, we find that the mapping $\mathfrak{H}_p \ni H \mapsto L_H \in G_p^+(M_{2r})$ is continuous. From the other side, it follows from Lemma 2.1 that so is the mapping $\mathfrak{Q}_p \ni q \mapsto K_q \in G_p^+(M_{2r})$. Furthermore, as follows from Remark 3.2, the mapping Θ acts continuously from \mathfrak{H}_p to \mathfrak{Q}_p . Hence, passing to the limit in (3.23) we obtain that

$$L_H = K_q, \quad q = \Theta(H),$$

as desired. □

Now we are ready to prove Theorem 1.3. The proof also uses the results of [24] and [18]:

Proof of Theorem 1.3. As follows from Remark 3.2, the mapping Θ acts continuously from \mathfrak{H}_p to \mathfrak{Q}_p for all $p \geq 1$. Therefore, it remains to prove that Θ is invertible and that the inverse mapping Θ^{-1} acts continuously from \mathfrak{Q}_p to \mathfrak{H}_p .

Denote by \mathfrak{H}_0 the set of all accelerants that are continuous on $[-1, 1] \setminus \{0\}$ having a jump discontinuity at the origin, and set

$$\Theta_0 := \Theta|_{\mathfrak{H}_0}.$$

Then it is proved in [24] that Θ_0 maps \mathfrak{H}_0 onto $C([0, 1], M_r)$ one-to-one. Since \mathfrak{H}_0 is dense in \mathfrak{H}_p and $C([0, 1], M_r)$ is dense in \mathfrak{Q}_p for all $p \geq 1$, the present theorem will be proved if we show that Θ_0^{-1} can be extended to a continuous mapping from \mathfrak{Q}_p to \mathfrak{H}_p .

Recalling the results of [18], we find that the mapping Θ_0^{-1} can be represented in the following way. For an arbitrary $q \in C([0, 1], M_r)$ denote by \mathcal{K}_q an integral operator in \mathbb{H} with kernel K_q (see Lemma 2.1) and set

$$\mathcal{F}^q := (\mathcal{I} + \mathcal{K}_q)^{-1}(\mathcal{I} + \mathcal{K}_q^*)^{-1} - \mathcal{I}. \quad (3.24)$$

Let F^q be the kernel of \mathcal{F}^q . Since the mappings $\mathfrak{Q}_p \ni q \mapsto K_q \in G_p^+(M_{2r})$ and $\mathfrak{G}_p^+(M_{2r}) \ni \mathcal{K} \mapsto (\mathcal{I} + \mathcal{K})^{-1} - \mathcal{I} \in \mathfrak{G}_p^+(M_{2r})$ are continuous (see Lemmas 2.1 and A.1, respectively), it follows that the mapping $\mathfrak{Q}_p \ni q \mapsto F^q \in G_p(M_{2r})$ is continuous as well.

As follows from Proposition 3.3, if $q = \Theta(H)$ for some $H \in \mathfrak{H}_p$, then $F^q = F_H$. By virtue of the formula (3.21) for F_H , the inverse of the mapping $\mathfrak{H}_p \ni H \mapsto F_H \in G_p(M_{2r})$ is a restriction of certain mapping $\eta : G_p(M_{2r}) \rightarrow L_p((-1, 1), M_r)$ that can be easily written: write $F \in G_p(M_{2r})$ in the block-diagonal form

$$F = \frac{1}{2} \begin{pmatrix} F_{11} & F_{12} \\ F_{21} & F_{22} \end{pmatrix},$$

where $F_{ks} \in G_p(M_r)$, $k, s \in \{1, 2\}$, and set

$$[\eta(F)](x) := \begin{cases} F_{21}(-2x - 1, 1), & -1 \leq x \leq -\frac{1}{2}, \\ F_{11}(2x + 1, 1), & -\frac{1}{2} < x \leq 0, \\ F_{22}(-2x + 1, 1), & 0 < x \leq \frac{1}{2}, \\ F_{12}(2x - 1, 1), & \frac{1}{2} < x \leq 1. \end{cases} \quad (3.25)$$

Now define the mapping Υ acting from $C([0, 1], M_r)$ to $L_p((-1, 1), M_r)$ by the formula

$$\Upsilon(q) := \eta(F^q), \quad q \in C([0, 1], M_r).$$

Since the mappings

$$\mathfrak{Q}_p \ni q \mapsto F^q \in G_p(M_{2r})$$

and

$$\eta : G_p(M_{2r}) \rightarrow L_p((-1, 1), M_r)$$

are continuous, we obtain that Υ acts continuously from \mathfrak{Q}_p to $L_p((-1, 1), M_r)$.

It follows from [18] that

$$\Upsilon(q) = \Theta_0^{-1}(q)$$

for all $q \in C([0, 1], M_r)$. Therefore, we find that the mapping Θ_0^{-1} can be extended to a continuous one from \mathfrak{Q}_p to \mathfrak{H}_p , as desired. \square

Theorem 1.3 will find its application in the next subsection, while Proposition 3.3 will also play an important role in solving the inverse spectral problem.

3.4 Condition (B_4)

The main result of this subsection is the following theorem:

Theorem 3.1 *Let $q \in \mathfrak{Q}_p$, $p \geq 1$, set $\mu := \mu_q$ and $H := H_\mu$ (see the definitions (1.3) and (1.7), respectively). Then $H \in \mathfrak{H}_p$ and $q = \Theta(H)$.*

In particular, from Theorem 3.1 we immediately obtain the following:

Corollary 3.3 For an arbitrary potential $q \in \mathfrak{Q}_p$, $p \geq 1$, the spectral data \mathfrak{a}_q of the operator T_q satisfy the condition (B_4) , i.e. $H_\mu \in L_p((-1, 1), M_r)$, $\mu := \mu_q$.

We prove Theorem 3.1 by a limiting procedure using the results of [18]. Before starting the proof, we need to establish two auxiliary technical lemmas.

Lemma 3.3 Let $q \in \mathfrak{Q}_p$, $p \geq 1$. Assume that for all $n \in \mathbb{N}$, $q_n \in C([0, 1], M_r)$ and $\lim_{n \rightarrow \infty} \|q - q_n\|_{\mathfrak{Q}_p} = 0$. Then there exist $N_0 \in \mathbb{N}$ and $n_0 \in \mathbb{N}$ such that for any natural $N > N_0$ and $n > n_0$ the functions m_q and m_{q_n} have no poles on the circle $K_N := \{\lambda \in \mathbb{C} \mid |\lambda| = \pi N + \pi/6\}$. Moreover, for all natural $N > N_0$,

$$\lim_{n \rightarrow \infty} \sup_{\lambda \in K_N} \|m_q(\lambda) - m_{q_n}(\lambda)\| = 0. \quad (3.26)$$

Proof. It follows from Lemma 2.1 that

$$s_q(\lambda) = (\sin \lambda)I + \mathcal{F}(\lambda)g_q, \quad s_{q_n}(\lambda) = (\sin \lambda)I + \mathcal{F}(\lambda)g_{q_n}, \quad (3.27)$$

where $g_q, g_{q_n} \in L_p((-1, 1), M_r)$ and $\mathcal{F}(\lambda)$ is the operator $L_p((-1, 1), M_r) \rightarrow M_r$ acting by the formula $\mathcal{F}(\lambda)f := \int_{-1}^1 e^{i\lambda t} f(t) dt$. Moreover, since $q_n \rightarrow q$ in the metric of the space \mathfrak{Q}_p , from Lemma 2.1 we also obtain that $\lim_{n \rightarrow \infty} \|g_{q_n} - g_q\|_{L_1} = 0$.

Similarly,

$$c_q(\lambda) = (\cos \lambda)I + \mathcal{F}(\lambda)h_q, \quad c_{q_n}(\lambda) = (\cos \lambda)I + \mathcal{F}(\lambda)h_{q_n},$$

where $h_q, h_{q_n} \in L_p((-1, 1), M_r)$ and $\lim_{n \rightarrow \infty} \|h_{q_n} - h_q\|_{L_1} = 0$.

Since $\|g_{q_n} - g_q\|_{L_1} \rightarrow 0$ and $\|h_{q_n} - h_q\|_{L_1} \rightarrow 0$ as $n \rightarrow \infty$, we find that for every $N \in \mathbb{N}$,

$$\lim_{n \rightarrow \infty} \sup_{\lambda \in K_N} \|s_q(\lambda) - s_{q_n}(\lambda)\| = 0, \quad \lim_{n \rightarrow \infty} \sup_{\lambda \in K_N} \|c_q(\lambda) - c_{q_n}(\lambda)\| = 0. \quad (3.28)$$

Now let us establish some estimates for $\|s_q(\lambda)^{-1}\|$ and $\|s_{q_n}(\lambda)^{-1}\|$ as $\lambda \in K_N$.

By virtue of Lemma A.2, we obtain that there exists $N_1 \in \mathbb{N}$ such that for all natural $N > N_1$,

$$e^{-|\operatorname{Im} \lambda|} \|\mathcal{F}(\lambda)g_q\| < \frac{1}{8}, \quad \lambda \in K_N. \quad (3.29)$$

In the meantime, using the expansion of $\sin \lambda$ as an infinite product we note that $|\sin \lambda| \geq |\sin |\lambda||$, $\lambda \in \mathbb{C}$, and thus $|\sin \lambda| \geq \frac{1}{2}$ as $\lambda \in K_N$. Moreover, as $|\operatorname{Im} \lambda| > \ln 2$ we arrive at the estimate

$$|\sin \lambda| = \frac{1}{2} |e^{i\lambda} - e^{-i\lambda}| \geq \frac{1}{4} e^{|\operatorname{Im} \lambda|},$$

while as $|\operatorname{Im} \lambda| \leq \ln 2$, $\lambda \in K_N$, we have

$$|\sin \lambda| \geq \frac{1}{2} \geq \frac{1}{4} e^{|\operatorname{Im} \lambda|}.$$

Thus, $|\sin \lambda| \geq \frac{1}{4} e^{|\operatorname{Im} \lambda|}$ as $\lambda \in K_N$. Using this estimate together with (3.29), we obtain from (3.27) that

$$\|s_q(\lambda)^{-1}\| \leq \frac{|\sin \lambda|^{-1}}{1 - |\sin \lambda|^{-1} \cdot \|\mathcal{F}(\lambda)g_q\|} \leq 4, \quad \lambda \in K_N, \quad N > N_1. \quad (3.30)$$

Similarly, since $\|g_{q_n} - g_q\|_{L_1} \rightarrow 0$ as $n \rightarrow \infty$ and $\|\mathcal{F}(\lambda)(g_{q_n} - g_q)\| \leq e^{|\operatorname{Im} \lambda|} \|g_{q_n} - g_q\|_{L_1}$, we obtain that there exist $n_0 \in \mathbb{N}$ and $N_2 \in \mathbb{N}$ such that for all natural $n > n_0$ and $N > N_2$,

$$e^{-|\operatorname{Im} \lambda|} \|\mathcal{F}(\lambda)g_{q_n}\| \leq e^{-|\operatorname{Im} \lambda|} \|\mathcal{F}(\lambda)g_q\| + e^{-|\operatorname{Im} \lambda|} \|\mathcal{F}(\lambda)(g_q - g_{q_n})\| < \frac{1}{8}$$

as $\lambda \in K_N$. Therefore, for all natural $n > n_0$,

$$\|s_{q_n}(\lambda)^{-1}\| \leq \frac{|\sin \lambda|^{-1}}{1 - |\sin \lambda|^{-1} \cdot \|\mathcal{F}(\lambda)g_{q_n}\|} \leq 4, \quad \lambda \in K_N, \quad N > N_2. \quad (3.31)$$

Hence we conclude that the functions m_q and m_{q_n} , $n > n_0$, have no poles on K_N as $N > N_0 := \max\{N_1, N_2\}$. Moreover, it follows from the definitions of m_q and m_{q_n} that

$$\begin{aligned} \|m_q(\lambda) - m_{q_n}(\lambda)\| &= \|-s_q(\lambda)^{-1}c_q(\lambda) + s_{q_n}(\lambda)^{-1}c_{q_n}(\lambda)\| \\ &\leq \|s_{q_n}(\lambda)^{-1}\| \cdot \|c_q(\lambda) - c_{q_n}(\lambda)\| + \|c_q(\lambda)\| \cdot \|s_q(\lambda)^{-1} - s_{q_n}(\lambda)^{-1}\|. \end{aligned}$$

Since

$$\|s_q(\lambda)^{-1} - s_{q_n}(\lambda)^{-1}\| \leq \|s_{q_n}(\lambda)^{-1}\| \cdot \|s_q(\lambda) - s_{q_n}(\lambda)\| \cdot \|s_q(\lambda)^{-1}\|,$$

taking into account (3.28), (3.30) and (3.31) we arrive at the equality (3.26). \square

We use the preceding lemma to obtain the following auxiliary result:

Lemma 3.4 *Let $q \in \mathfrak{Q}_p$, $p \geq 1$. Assume that for all $n \in \mathbb{N}$, $q_n \in C([0, 1], M_r)$ and $\lim_{n \rightarrow \infty} \|q - q_n\|_{\mathfrak{Q}_p} = 0$. Then for all $f \in \mathcal{S}^r$ (see Appendix A) such that $\operatorname{supp} f \subset [-1, 1]$,*

$$\lim_{n \rightarrow \infty} (H_{\mu_{q_n}}, f) = (H_{\mu_q}, f),$$

where μ_{q_n} and μ_q are the spectral measures of the operators T_{q_n} and T_q , respectively.

Proof. Let $(\alpha_j, \lambda_j)_{j \in \mathbb{Z}}$ and $(\alpha_{j,n}, \lambda_{j,n})_{j \in \mathbb{Z}}$ be the spectral data of the operators T_q and T_{q_n} , respectively. Then for all $f \in \mathcal{S}^r$, $\operatorname{supp} f \subset [-1, 1]$,

$$(H_{\mu_{q_n}}, f) = \sum_{j=-\infty}^{\infty} \alpha_{j,n} \widehat{f}(\lambda_{j,n}) - \sum_{k=-\infty}^{\infty} \widehat{f}(\pi k),$$

where $\widehat{f}(\lambda)$ is defined by (1.6). Since $\widehat{f} \in \mathcal{S}^r$ and for all $n \in \mathbb{N}$ the norms $\|\alpha_{j,n}\|$, $j \in \mathbb{Z}$, are uniformly bounded (see (3.2)), it suffices to show that there exists $N_0 \in \mathbb{N}$ such that for all natural $N > N_0$,

$$\lim_{n \rightarrow \infty} \sum_{|\lambda_{j,n}| < \pi N + \frac{\pi}{6}} \alpha_{j,n} \widehat{f}(\lambda_{j,n}) = \sum_{|\lambda_j| < \pi N + \frac{\pi}{6}} \alpha_j \widehat{f}(\lambda_j). \quad (3.32)$$

Note that for fixed $N \in \mathbb{N}$ both the left and right parts of (3.32) depend only on the values of \widehat{f} on some (large enough) finite interval. Since the function $\widehat{f} \in \mathcal{S}^r$ can be uniformly approximated by ‘‘polynomials’’ from \mathcal{P}^r (see Appendix A) on any finite interval, (3.32) will be proved if we show that

$$\lim_{n \rightarrow \infty} \sum_{|\lambda_{j,n}| < \pi N + \frac{\pi}{6}} \alpha_{j,n} P(\lambda_{j,n}) = \sum_{|\lambda_j| < \pi N + \frac{\pi}{6}} \alpha_j P(\lambda_j)$$

for an arbitrary $P \in \mathcal{P}^r$.

Since the functions m_q and m_{q_n} have only poles of first order at points λ_j and $\lambda_{j,n}$, respectively, by virtue of the asymptotics (3.1) and Lemma 3.3 we find that

$$\sum_{|\lambda_{j,n}| < \pi N + \frac{\pi}{6}} \alpha_{j,n} P(\lambda_{j,n}) = \frac{1}{2\pi i} \oint_{K_N} m_{q_n}(\lambda) P(\lambda) d\lambda$$

and

$$\sum_{|\lambda_j| < \pi N + \frac{\pi}{6}} \alpha_j P(\lambda_j) = \frac{1}{2\pi i} \oint_{K_N} m_q(\lambda) P(\lambda) d\lambda$$

for large values of N , where $K_N := \{\lambda \in \mathbb{C} \mid |\lambda| = \pi N + \pi/6\}$. Hence it is enough to prove that there exists $N_0 \in \mathbb{N}$ such that for all natural $N > N_0$,

$$\lim_{n \rightarrow \infty} \frac{1}{2\pi i} \oint_{K_N} m_{q_n}(\lambda) P(\lambda) d\lambda = \frac{1}{2\pi i} \oint_{K_N} m_q(\lambda) P(\lambda) d\lambda.$$

But this claim follows directly from (3.26), and thus the proof is complete. \square

Now we can use Lemma 3.4 to prove Theorem 3.1:

Proof of Theorem 3.1. Let $q \in \mathfrak{Q}_p$, $p \geq 1$. Assume that for all $n \in \mathbb{N}$, $q_n \in C([0, 1], M_r)$ and $\lim_{n \rightarrow \infty} \|q - q_n\|_{\mathfrak{Q}_p} = 0$. Let μ_q and μ_{q_n} , $n \in \mathbb{N}$, be the spectral measures of Dirac operators T_q and T_{q_n} , respectively. Then from Lemma 3.4 we obtain that for all $f \in \mathcal{S}^r$ such that $\text{supp } f \subset [-1, 1]$,

$$\lim_{n \rightarrow \infty} (H_{\mu_{q_n}}, f) = (H_{\mu_q}, f). \quad (3.33)$$

It follows from [18] that for all $n \in \mathbb{N}$, $H_{\mu_{q_n}} \in \mathfrak{H}_2$ and that $H_{\mu_{q_n}} = \Theta^{-1}(q_n)$. Hence (3.33) reads

$$\lim_{n \rightarrow \infty} (\Theta^{-1}(q_n), f) = (H_{\mu_q}, f).$$

From the other side, since Θ^{-1} acts continuously from \mathfrak{Q}_p to \mathfrak{H}_p , we observe that for all $f \in \mathcal{S}^r$, $\text{supp } f \subset [-1, 1]$,

$$\lim_{n \rightarrow \infty} (\Theta^{-1}(q_n), f) = (\Theta^{-1}(q), f).$$

Thus we obtain that for all $f \in \mathcal{S}^r$, $\text{supp } f \subset [-1, 1]$,

$$(H_{\mu_q}, f) = (\Theta^{-1}(q), f),$$

i.e. $H_{\mu_q} = \Theta^{-1}(q)$, as desired. \square

Thus, so far we have proved that for an arbitrary potential $q \in \mathfrak{Q}_p$, $p \geq 1$, the spectral data \mathfrak{a}_q of the operator T_q satisfy the conditions (B_1) – (B_4) . This is the necessity part of Theorem 1.1. The next section is devoted to establishing its sufficiency part and solving the inverse spectral problem for the operator T_q .

4 Inverse spectral problem

In this section we finish the proof of Theorem 1.1 and prove Theorems 1.2 and 1.4, and thus solve the inverse spectral problem for the operator T_q . Namely, we show that if a sequence $\mathfrak{a} := ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}}$ satisfies the conditions (B_1) – (B_4) , then there exists a unique potential $q \in \mathfrak{Q}_p$ such that $\mathfrak{a} = \mathfrak{a}_q$. Theorem 1.4 then suggests a method how to reconstruct this potential from the spectral data.

4.1 Proof of Theorems 1.1 and 1.4

Here we prove that if a sequence \mathbf{a} satisfies the conditions (B_1) – (B_4) , $\mu := \mu^{\mathbf{a}}$ and $H := H_\mu$, then $H \in \mathfrak{H}_p$ and $\mathbf{a} = \mathbf{a}_q$ for $q = \Theta(H)$. This is the sufficiency part of Theorem 1.1 and the proof of Theorem 1.4.

We start from the following technical but nevertheless important lemma:

Lemma 4.1 *For an arbitrary sequence $\mathbf{a} := ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}}$ set $\mu := \mu^{\mathbf{a}}$ and $H := H_\mu$. If $H \in L_p((-1, 1), M_r)$, $p \geq 1$, then the following equality holds true:*

$$\mathcal{I} + \mathcal{F}_H = \text{s-lim}_{N \rightarrow \infty} \sum_{j=-N}^N \Phi_0(\lambda_j) \alpha_j \Phi_0^*(\lambda_j), \quad (4.1)$$

where $\mathcal{F}_H \in \mathcal{G}_p(M_{2r})$ is an integral operator in \mathbb{H} with kernel F_H given by (3.21) and the operators $\Phi_0(\lambda) : \mathbb{C}^r \rightarrow \mathbb{H}$ are given by (2.10).

Proof. Given an arbitrary sequence $\mathbf{a} := ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}}$, set $\mu := \mu^{\mathbf{a}}$, $H := H_\mu$, and assume that $H \in L_p((-1, 1), M_r)$. Denote by \mathcal{H} an integral operator in $L_2((0, 1), \mathbb{C}^r)$ acting by the formula

$$(\mathcal{H}f)(x) := \int_0^1 H(x-t)f(t) dt, \quad x \in (0, 1),$$

and let \mathcal{I} stand for the identity operator in $L_2((0, 1), \mathbb{C}^r)$.

Let us show that the operator $\mathcal{I} + \mathcal{H}$ is unitarily equivalent to $\mathcal{I} + \mathcal{F}_H$. Indeed, consider the unitary transformation $V : L_2((0, 1), \mathbb{C}^r) \rightarrow \mathbb{H}$ given by the formula

$$(Vf)(t) := \frac{1}{\sqrt{2}} \begin{pmatrix} f\left(\frac{1+u}{2}\right) \\ f\left(\frac{1-u}{2}\right) \end{pmatrix}, \quad f \in L_2((0, 1), \mathbb{C}^r).$$

Then a simple verification shows that

$$\mathcal{I} + \mathcal{F}_H = V(\mathcal{I} + \mathcal{H})V^{-1}.$$

Furthermore, we find that for all $j \in \mathbb{Z}$,

$$\Phi_0(\lambda_j) \alpha_j \Phi_0^*(\lambda_j) = V \Psi_0(\lambda_j) \alpha_j \Psi_0^*(\lambda_j) V^{-1},$$

where for an arbitrary $\lambda \in \mathbb{C}$ the operator $\Psi_0(\lambda)$ acts from \mathbb{C}^r to $L_2((0, 1), \mathbb{C}^r)$ by the formula

$$[\Psi_0(\lambda)c](x) := e^{2i\lambda x} c,$$

$\Psi_0^*(\lambda) := [\Psi_0(\lambda)]^*$. Hence, in order to prove (4.1) it suffices to show that

$$\mathcal{I} + \mathcal{H} = \text{s-lim}_{N \rightarrow \infty} \sum_{j=-N}^N \Psi_0(\lambda_j) \alpha_j \Psi_0^*(\lambda_j). \quad (4.2)$$

Firstly, we observe that (4.2) holds true on the set of smooth functions $f \in \mathcal{S}^r$. Namely, let $f \in \mathcal{S}^r$, $\text{supp } f \subset [0, 1]$. Then, by virtue of definitions (1.6) and (1.7), we have that for an arbitrary but fixed $x \in (0, 1)$,

$$\begin{aligned} [(\mathcal{I} + \mathcal{H})f](x) &= f(x) + \int_0^1 H(x-t)f(t) dt = f(x) + \int_{-1}^1 H(s)f(x-s) ds \\ &= \sum_{j=-\infty}^{\infty} \alpha_j \int_{-1}^1 e^{2i\lambda_j s} f(x-s) ds = \sum_{j=-\infty}^{\infty} \alpha_j \int_0^1 e^{2i\lambda_j(x-t)} f(t) dt, \end{aligned}$$

noting that the series in the last two expressions are convergent because $f \in \mathcal{S}^r$. Thus we conclude that for all $f \in \mathcal{S}^r$, $\text{supp } f \subset [0, 1]$,

$$(\mathcal{I} + \mathcal{H})f = \lim_{N \rightarrow \infty} \sum_{j=-N}^N \Psi_0(\lambda_j) \alpha_j \Psi_0^*(\lambda_j) f. \quad (4.3)$$

Since $\{f \in \mathcal{S}^r \mid \text{supp } f \subset [0, 1]\}$ is dense everywhere in $L_2((0, 1), \mathbb{C}^r)$, in order to finish the proof it remains to show that there exists the limit $\text{s-lim}_{N \rightarrow \infty} \sum_{j=-N}^N C_j$, where the operators C_j , $j \in \mathbb{Z}$, act in $L_2((0, 1), \mathbb{C}^r)$ by the formula

$$C_j := \Psi_0(\lambda_j) \alpha_j \Psi_0^*(\lambda_j).$$

To this end, note that for every $j \in \mathbb{Z}$ the operator C_j is self-adjoint and non-negative, and that (4.3) implies that for all $N \in \mathbb{N}$,

$$\sum_{j=-N}^N C_j \leq \mathcal{I} + \mathcal{H}.$$

Therefore, from well-known theorem on convergence of monotonic sequence of non-negative self-adjoint operators (see, e.g., [26, Chapter IV, §2]) we obtain that the sequence $\sum_{j=-N}^N C_j$, $N \in \mathbb{N}$, is convergent in the strong operator topology. Hence (4.2) follows, and thus the proof is complete. \square

Now, the following observation will serve as a part of Theorem 1.4:

Lemma 4.2 *Let a sequence \mathbf{a} satisfy the conditions (B_3) and (B_4) , set $\mu := \mu^{\mathbf{a}}$ and $H := H_{\mu}$. Then $H \in \mathfrak{H}_p$.*

Proof. Firstly, it follows from the condition (B_4) that $H \in L_p((-1, 1), M_r)$. Construct the function $F_H \in G_p(M_{2r})$ by formula (3.21) and let $\mathcal{F}_H \in \mathcal{G}_p(M_{2r})$ be an integral operator in \mathbb{H} with kernel F_H . Taking into account Lemma B.1, we find that the present lemma will be proved if we show that $\mathcal{I} + \mathcal{F}_H > 0$.

It follows from Lemma 4.1 that

$$\mathcal{I} + \mathcal{F}_H = \text{s-lim}_{N \rightarrow \infty} \sum_{j=-N}^N \Phi_0(\lambda_j) \alpha_j \Phi_0^*(\lambda_j).$$

Therefore, since for every $j \in \mathbb{Z}$ the operator $\Phi_0(\lambda_j) \alpha_j \Phi_0^*(\lambda_j)$ is non-negative, it follows that $\mathcal{I} + \mathcal{F}_H \geq 0$. Hence, since the operator \mathcal{F}_H is compact, we obtain that $\mathcal{I} + \mathcal{F}_H > 0$ if and only if $\ker(\mathcal{I} + \mathcal{F}_H) = \{0\}$. Now taking into account that for all $j \in \mathbb{Z}$, $\ker \Phi_0(\lambda_j) = \{0\}$, we find that

$$\ker(\mathcal{I} + \mathcal{F}_H) = \ker \left(\sum_{j=-\infty}^{\infty} \Phi_0(\lambda_j) \alpha_j \Phi_0^*(\lambda_j) \right) = \bigcap_{j=-\infty}^{\infty} \ker \alpha_j \Phi_0^*(\lambda_j) = U \mathcal{X}^{\perp},$$

where \mathcal{X} is the set from the condition (B_3) defined by (3.20) and the unitary mapping $U : L_2((-1, 1), M_r) \rightarrow \mathbb{H}$ acts by the formula

$$(Uf)(x) = \begin{pmatrix} f(x), & f(-x) \end{pmatrix}^{\top}, \quad x \in (0, 1).$$

Since, by virtue of the condition (B_3) , $\mathcal{X}^\perp = \{0\}$, we obtain that $\ker(\mathcal{I} + \mathcal{F}_H) = \{0\}$, as desired. \square

Now, given an arbitrary sequence \mathbf{a} satisfying the conditions (B_1) – (B_4) , set $\mu := \mu^\mathbf{a}$, $H := H_\mu$ and

$$q := \Theta(H).$$

For all $j \in \mathbb{Z}$ define the operators

$$\tilde{P}_{\mathbf{a},j} := \Phi_q(\lambda_j)\alpha_j\Phi_q^*(\lambda_j). \quad (4.4)$$

Remark 4.1 *By this construction, if $\mathbf{a} = \mathbf{a}_q$ for some $q \in \mathfrak{Q}_p$, then $\tilde{P}_{\mathbf{a},j} = P_{q,j}$, i.e. $\tilde{P}_{\mathbf{a},j}$ is an orthogonal projector of the operator T_q corresponding to the eigenvalue $\lambda_j(q)$ (see Theorem 2.1, parts (iii) and (iv)).*

According to (4.4), for every $j \in \mathbb{Z}$ the operator $\tilde{P}_{\mathbf{a},j}$ is self-adjoint non-negative operator of finite rank. If we show that $\{\tilde{P}_{\mathbf{a},j}\}_{j \in \mathbb{Z}}$ is a complete system of orthogonal projectors, then the same arguments as in [17, 18] will lead us to the desired equality $\mathbf{a} = \mathbf{a}_q$, $q = \Theta(H)$.

We start from proving completeness of $\{\tilde{P}_{\mathbf{a},j}\}_{j \in \mathbb{Z}}$, which is a direct consequence of Proposition 3.3:

Lemma 4.3 *Let a sequence \mathbf{a} satisfy the conditions (B_1) – (B_4) , set $\mu := \mu^\mathbf{a}$, $H := H_\mu$ and $q := \Theta(H)$. Then*

$$\text{s-lim}_{N \rightarrow \infty} \sum_{j=-N}^N \tilde{P}_{\mathbf{a},j} = \mathcal{I}. \quad (4.5)$$

Proof. Let the assumptions of the present lemma hold true. Construct the function $F_H \in G_p(M_{2r})$ by formula (3.21) and denote by $\mathcal{F}_H \in \mathcal{G}_p(M_{2r})$ an integral operator in \mathbb{H} with kernel F_H . It follows from Proposition 3.3 that

$$\mathcal{I} + \mathcal{F}_H = (\mathcal{I} + \mathcal{K}_q)^{-1}(\mathcal{I} + \mathcal{K}_q^*)^{-1}, \quad q = \Theta(H), \quad (4.6)$$

where $\mathcal{K}_q \in \mathcal{G}_p^+(M_{2r})$ is an integral operator in \mathbb{H} with kernel K_q (see Lemma 2.1). From the other side, observing that by virtue of Lemma 4.1,

$$\mathcal{I} + \mathcal{F}_H = \text{s-lim}_{N \rightarrow \infty} \sum_{j=-N}^N \Phi_0(\lambda_j)\alpha_j\Phi_0^*(\lambda_j), \quad (4.7)$$

and recalling that $\Phi_q(\lambda) = (\mathcal{I} + \mathcal{K}_q)\Phi_0(\lambda)$ (see (2.9)), by virtue of the equalities (4.6) and (4.7) we arrive at (4.5). \square

Next we prove that $\{\tilde{P}_{\mathbf{a},j}\}_{j \in \mathbb{Z}}$ is a system of pairwise orthogonal projectors. Again, as in solving the direct spectral problem, the proof uses Lemma 3.2 and the theory of Riesz bases.

Lemma 4.4 *Let a sequence \mathbf{a} satisfy the conditions (B_1) – (B_4) , set $\mu := \mu^\mathbf{a}$, $H := H_\mu$ and $q := \Theta(H)$. Then $\{\tilde{P}_{\mathbf{a},j}\}_{j \in \mathbb{Z}}$ is a system of pairwise orthogonal projectors.*

Proof. As in the proof of Proposition 3.2, for an arbitrary $j \in \mathbb{Z}$ we denote by $\{v_{j,k}\}_{k=1}^{\text{rank } \alpha_j}$ a system of pairwise orthogonal vectors in \mathbb{C}^r such that

$$\alpha_j = \sum_{k=1}^{\text{rank } \alpha_j} (\cdot | v_{j,k})v_{j,k}.$$

Then, taking into account (4.4), we obtain that

$$\tilde{P}_{a,j} = \sum_{k=1}^{\text{rank } \alpha_j} (\cdot | \mathbf{f}_{j,k}) \mathbf{f}_{j,k},$$

where $\mathbf{f}_{j,k} := \Phi_q(\lambda_j) v_{j,k}$.

The present lemma will be proved if we show that the system

$$\mathcal{A} := \{ \mathbf{f}_{j,k} \mid j \in \mathbb{Z}, \ 1 \leq k \leq \text{rank } \alpha_j \}$$

is a Riesz basis for the space \mathbb{H} . Indeed, from the equality (4.5) we then obtain that for an arbitrary function $\mathbf{f}_{s,l}$,

$$\sum_{j=-\infty}^{\infty} \sum_{k=1}^{\text{rank } \alpha_j} (\mathbf{f}_{s,l} | \mathbf{f}_{j,k}) \mathbf{f}_{j,k} = \mathbf{f}_{s,l},$$

which implies the equalities $(\mathbf{f}_{j,k} | \mathbf{f}_{j,k}) = 1$ and $(\mathbf{f}_{s,l} | \mathbf{f}_{j,k}) = 0$ as $(s,l) \neq (j,k)$. Therefore, $\tilde{P}_{a,s} \tilde{P}_{a,j} = 0$ as $s \neq j$, and thus $\{\tilde{P}_{a,j}\}_{j \in \mathbb{Z}}$ is a system of pairwise orthogonal projectors.

Hence it remains to prove that \mathcal{A} is a Riesz basis for \mathbb{H} . To this end, it suffices to show that the system

$$\mathcal{B}_0 := \{ e^{i\lambda_j t} v_{j,k} \mid j \in \mathbb{Z}, \ 1 \leq k \leq \text{rank } \alpha_j \} \quad (4.8)$$

is a Riesz basis for the space $\mathcal{H} := L_2((-1,1), \mathbb{C}^r)$. Indeed, introduce the unitary mapping $U : \mathcal{H} \rightarrow \mathbb{H}$ acting by the formula

$$(Uf)(t) = (f(-t), \ f(t))^\top, \quad t \in (0,1),$$

and note that U maps $e^{i\lambda_j x} v_{j,k}$ to $\Phi_0(\lambda_j) v_{j,k}$ for all $j \in \mathbb{Z}$, $1 \leq k \leq \text{rank } \alpha_j$. Therefore, if \mathcal{B}_0 is a Riesz basis for \mathcal{H} , then the system

$$\mathcal{A}_0 := \{ \Phi_0(\lambda_j) v_{j,k} \mid j \in \mathbb{Z}, \ 1 \leq k \leq \text{rank } \alpha_j \}$$

is a Riesz basis for \mathbb{H} . Then, since $\Phi_q(\lambda) = (\mathcal{I} + \mathcal{K}_q) \Phi_0(\lambda)$ and $\mathcal{I} + \mathcal{K}_q$ is a homeomorphism of the space \mathbb{H} , we obtain that \mathcal{A} remains a Riesz basis for \mathbb{H} , as desired.

Thus let us show that \mathcal{B}_0 is a Riesz basis for \mathcal{H} . We do this by applying Lemma 3.2 and Theorem C.2. Firstly, observe that by virtue of the conditions (B_1) and (B_2) , there exists $N_0 \in \mathbb{N}$ such that for all $n \in \mathbb{Z}$, $|n| > N_0$,

$$\sum_{\lambda_j \in \Delta_n} \text{rank } \alpha_j = r. \quad (4.9)$$

Indeed, it follows from the condition (B_1) that there exists $N_0 \in \mathbb{N}$ such that

$$\left\| I - \sum_{\lambda_j \in \Delta_n} \alpha_j \right\| < 1, \quad |n| > N_0.$$

Hence, $\sum_{\lambda_j \in \Delta_n} \text{rank } \alpha_j \geq r$ as $|n| > N_0$. Moreover, by virtue of the condition (B_2) , N_0 can be taken so large that

$$\sum_{n=-N}^N \sum_{\lambda_j \in \Delta_n} \text{rank } \alpha_j = (2N+1)r$$

for all natural $N \geq N_0$. Therefore, $\sum_{\lambda_j \in \Delta_n} \text{rank } \alpha_j + \sum_{\lambda_j \in \Delta_{-n}} \text{rank } \alpha_j = 2r$ as $|n| > N_0$, and thus we arrive at (4.9).

Now, since a sequence \mathbf{a} satisfies the condition (B_1) and for all $n \in \mathbb{Z}$, $|n| > N_0$, the equality (4.9) holds true, we find that the conditions of Lemma 3.2 are satisfied. Therefore, from Lemma 3.2 we obtain that there exists a large enough natural $N > N_0$ such that the system $\mathcal{E}_N \cup \mathcal{B}_N$, where

$$\mathcal{E}_N := \left\{ \frac{1}{\sqrt{2}} e^{i\pi n t} \epsilon_s \mid n \in \mathbb{Z}, |n| \leq N, \quad s = 1, \dots, r \right\}$$

and

$$\mathcal{B}_N := \left\{ e^{i\lambda_j t} v_{j,k} \mid \lambda_j \in \Delta_n, |n| > N, \quad 1 \leq k \leq \text{rank } \alpha_j \right\},$$

is a Riesz basis for the space \mathcal{H} .

Finally, observe that by virtue of the condition (B_3) , the system \mathcal{B}_0 is complete in \mathcal{H} . Moreover, since $\mathcal{B}_N \subset \mathcal{B}_0$ and, by virtue of the condition (B_2) , the finite systems

$$\mathcal{B}_0 \setminus \mathcal{B}_N = \left\{ e^{i\lambda_j t} v_{j,k} \mid \lambda_j \in \Delta_n, |n| \leq N, \quad 1 \leq k \leq \text{rank } \alpha_j \right\}$$

and \mathcal{E}_N consist of the same number of elements, we find that \mathcal{B}_0 is quadratically close to $\mathcal{E}_N \cup \mathcal{B}_N$. Then it follows from Theorem C.2 that \mathcal{B}_0 remains a Riesz basis for \mathcal{H} , as desired. \square

Now we are ready to prove Theorem 1.4:

Proof of Theorem 1.4. Firstly, it follows from Lemma 4.2 that if a sequence \mathbf{a} satisfies the conditions (B_1) – (B_4) , $\mu := \mu^{\mathbf{a}}$ and $H := H_{\mu}$, then $H \in \mathfrak{H}_p$. Thus, it only remains to prove that $\mathbf{a} = \mathbf{a}_q$ for $q = \Theta(H)$.

The proof of this claim repeats the technique that was suggested in [17]. Namely, as in [17, 18], we observe that it is enough to prove the embedding

$$\text{Ran } \tilde{P}_{\mathbf{a},j} \subset \ker(T_q - \lambda_j \mathcal{J}), \quad j \in \mathbb{Z}, \quad (4.10)$$

where the operators $\tilde{P}_{\mathbf{a},j}$ are given by (4.4). Indeed, taking into account completeness of $\{\tilde{P}_{\mathbf{a},j}\}_{j \in \mathbb{Z}}$, from (4.10) we immediately conclude that $\lambda_j = \lambda_j(q)$ for all $j \in \mathbb{Z}$, where $\lambda_j(q)$ are eigenvalues of T_q . Now, from this equality and from (4.10) we obtain that $P_{q,j} - \tilde{P}_{\mathbf{a},j} \geq 0$, $j \in \mathbb{Z}$, where $P_{q,j}$ are the orthogonal projectors of the operator T_q (see Theorem 2.1). However, taking into account completeness of the systems $\{\tilde{P}_{\mathbf{a},j}\}_{j \in \mathbb{Z}}$ and $\{P_{q,j}\}_{j \in \mathbb{Z}}$, we observe that $\sum_{j=-\infty}^{\infty} (P_{q,j} - \tilde{P}_{\mathbf{a},j}) = 0$, and thus $P_{q,j} - \tilde{P}_{\mathbf{a},j} = 0$ for all $j \in \mathbb{Z}$. Therefore, recalling the representation (2.14) for $P_{q,j}$, we find that

$$\Phi_q(\lambda_j) \{\alpha_j(q) - \alpha_j\} \Phi_q^*(\lambda_j) = 0, \quad j \in \mathbb{Z},$$

and taking into account (2.11) we arrive at $\alpha_j = \alpha_j(q)$. Together with $\lambda_j = \lambda_j(q)$ this means that $\mathbf{a} = \mathbf{a}_q$, as desired.

Now let us prove (4.10). Firstly, from the definition (4.4) of $\tilde{P}_{\mathbf{a},j}$ we obtain that

$$\text{Ran } \tilde{P}_{\mathbf{a},j} = \{\varphi_q(\cdot, \lambda_j) \alpha_j c \mid c \in \mathbb{C}^r\}.$$

From the other side, it follows from Lemma 2.2 that

$$\ker(T_q - \lambda_j \mathcal{J}) = \{\varphi_q(\cdot, \lambda_j) c \mid a \varphi_q(1, \lambda_j) c = 0\}.$$

Therefore, it suffices to show that $a \varphi_q(1, \lambda_j) \alpha_j = 0$. The proof of this claim is technical and literally repeats the proof in [18]. \square

Finally, the proof of Theorem 1.1 is straightforward:

Proof of Theorem 1.1. Firstly, by virtue of Propositions 3.1, 3.2 and Corollaries 3.2, 3.3 we have that for all $q \in \mathfrak{Q}_p$, $p \geq 1$, the spectral data \mathfrak{a}_q of the operator T_q satisfy the conditions (B_1) – (B_4) . This is the necessity part of Theorem 1.1. Conversely, if a sequence \mathfrak{a} satisfies the conditions (B_1) – (B_4) , $\mu := \mu^{\mathfrak{a}}$ and $H := H_\mu$, then it follows from Theorem 1.4 that $H \in \mathfrak{H}_p$ and $\mathfrak{a} = \mathfrak{a}_q$ for $q = \Theta(H)$. This is the sufficiency part of the theorem. \square

4.2 Proof of Theorem 1.2

Thus it only remains to prove Theorem 1.2 stating that the spectral data of the operator T_q determine the potential q uniquely. The proof of this claim repeats the proof in [18]:

Proof of Theorem 1.2. Evidently, the theorem will be proved if we prove the implication $\mathfrak{a}_{q_1} = \mathfrak{a}_{q_2} \Rightarrow q_1 = q_2$. Therefore, let $q_1, q_2 \in \mathfrak{Q}_p$, $p \geq 1$, and assume that $\mathfrak{a}_{q_1} = \mathfrak{a}_{q_2} =: \mathfrak{a}$. Let us show that $q_1 = q_2$. Set $H := H_\mu$, $\mu := \mu^{\mathfrak{a}}$, and construct the function $F_H \in G_p(M_{2r})$ by the formula (3.21). Denote by $\mathcal{F}_H \in \mathcal{G}_p(M_{2r})$ an integral operator in \mathbb{H} with kernel F_H . Since

$$\mathcal{I} + \mathcal{F}_H = \text{s-lim}_{N \rightarrow \infty} \sum_{j=-N}^N \Phi_0(\lambda_j) \alpha_j \Phi_0^*(\lambda_j),$$

by virtue of the equalities (2.9), (2.13) and (2.14) we find that

$$\mathcal{I} + \mathcal{F}_H = (\mathcal{I} + \mathcal{K}_{q_1})^{-1} (\mathcal{I} + \mathcal{K}_{q_1}^*)^{-1} = (\mathcal{I} + \mathcal{K}_{q_2})^{-1} (\mathcal{I} + \mathcal{K}_{q_2}^*)^{-1},$$

where \mathcal{K}_{q_j} is an integral operator in \mathbb{H} with kernel K_{q_j} , $j = 1, 2$. Since the operator $\mathcal{I} + \mathcal{F}_H$ may admit at most one factorization in $\mathcal{G}_p(M_{2r})$, we arrive at the equality

$$\mathcal{K}_{q_1} = \mathcal{K}_{q_2}.$$

Thus it remains to prove the implication $\mathcal{K}_{q_1} = \mathcal{K}_{q_2} \Rightarrow q_1 = q_2$. By virtue of (2.2) and (2.5), this can be obtained from the uniqueness theorem repeating the proof in [18]. \square

Acknowledgment

The author is extremely grateful to his supervisor Ass. Prof. Yaroslav Mykytyuk for countless discussions, valuable ideas and permanent attention to this work. The author also would like to thank Dr. Rostyslav Hryniv for valuable suggestions in preparing this manuscript.

A Spaces

Here we introduce several spaces that are used in this paper.

For an arbitrary $p \geq 1$ we denote by $G_p(M_r)$ the set of all measurable functions $K : [0, 1]^2 \rightarrow M_r$ such that for all x and t in $[0, 1]$ the functions $K(x, \cdot)$ and $K(\cdot, t)$ belong to $L_p((0, 1), M_r)$ and, moreover, the mappings

$$[0, 1] \ni x \mapsto K(x, \cdot) \in L_p((0, 1), M_r), \quad [0, 1] \ni t \mapsto K(\cdot, t) \in L_p((0, 1), M_r)$$

are continuous. The set $G_p(M_r)$ becomes a Banach space upon introducing the norm

$$\|K\|_{G_p} = \max \left\{ \max_{x \in [0, 1]} \|K(x, \cdot)\|_{L_p}, \max_{t \in [0, 1]} \|K(\cdot, t)\|_{L_p} \right\}.$$

We denote by $\mathcal{G}_p(M_r)$ the set of all integral operators in $L_2((0, 1), \mathbb{C}^r)$ with kernels $K \in G_p(M_r)$ and endow $\mathcal{G}_p(M_r)$ with the norm

$$\|\mathcal{K}\|_{\mathcal{G}_p} := \|K\|_{G_p}, \quad \mathcal{K} \in \mathcal{G}_p(M_r).$$

The space $\mathcal{G}_p(M_r)$ forms a subalgebra in the algebra \mathcal{B}_∞ of compact operators in $L_2((0, 1), \mathbb{C}^r)$.

We set

$$\Omega := \{(x, t) \mid 0 \leq t \leq x \leq 1\}, \quad \Omega^- := [0, 1]^2 \setminus \Omega,$$

and write $G_p^+(M_r)$ for the set of all functions $K \in G_p(M_r)$ such that $K(x, t) = 0$ a.e. in Ω^- , and $G_p^-(M_r)$ for set of all $K \in G_p(M_r)$ such that $K(x, t) = 0$ a.e. in Ω . By $\mathcal{G}_p^\pm(M_r)$ we denote the subalgebras of $\mathcal{G}_p(M_r)$ consisting of all operators $\mathcal{K} \in \mathcal{G}_p(M_r)$ with kernels $K \in G_p^\pm(M_r)$.

Let \mathcal{I} stand for the identity operator in $L_2((0, 1), \mathbb{C}^r)$. Then the following lemma is established in [17] for the case $p = 2$. However, its generalization to the case of an arbitrary $p \geq 1$ is straightforward:

Lemma A.1 *For all $p \geq 1$, the mapping $\mathcal{K} \mapsto (\mathcal{I} + \mathcal{K})^{-1} - \mathcal{I}$ acts continuously in $\mathcal{G}_p^+(M_r)$.*

We denote by \mathcal{P} the set of all polynomials over complex numbers and set

$$\mathcal{P}^r := \{(f_1, \dots, f_r)^\top \mid f_j \in \mathcal{P}, j = 1, \dots, r\}, \quad r \in \mathbb{N}.$$

We denote by \mathcal{S} the Schwartz space of smooth functions $f \in C^\infty(\mathbb{R})$ whose derivatives (including the function itself) decay at infinity faster than any power of $|x|^{-1}$, i.e.

$$\mathcal{S} := \{f \in C^\infty(\mathbb{R}) \mid x^\alpha D^\beta f(x) \rightarrow 0 \text{ as } |x| \rightarrow \infty, \quad \alpha, \beta \in \mathbb{N} \cup \{0\}\}.$$

Similarly, we set

$$\mathcal{S}^r := \{(f_1, \dots, f_r)^\top \mid f_j \in \mathcal{S}, j = 1, \dots, r\}, \quad r \in \mathbb{N}.$$

Also, we formulate here the following refined version of the Riemann–Lebesgue lemma that is mentioned, e.g., in [19]:

Lemma A.2 *Let $g \in L_1((-1, 1), M_r)$. Then*

$$\lim_{\mathbb{C} \ni \lambda \rightarrow \infty} e^{-|\operatorname{Im} \lambda|} \int_{-1}^1 e^{i\lambda t} g(t) dt = 0$$

in the metric of the space M_r .

B Factorization of integral operators

In this appendix we recall some facts from the theory of factorization of integral operators (see [27, 28]), which are also mentioned in [16, 17, 18]. See also [29] for the details.

We say that an operator $\mathcal{I} + \mathcal{F}$, where $\mathcal{F} \in \mathcal{G}_p(M_r)$, $p \geq 1$, admits a factorization in $\mathcal{G}_p(M_r)$ if there exist the operators $\mathcal{L}^+ \in \mathcal{G}_p^+(M_r)$ and $\mathcal{L}^- \in \mathcal{G}_p^-(M_r)$ such that

$$\mathcal{I} + \mathcal{F} = (\mathcal{I} + \mathcal{L}^+)^{-1} (\mathcal{I} + \mathcal{L}^-)^{-1}. \quad (\text{B.1})$$

If \mathcal{F} is self-adjoint, then $\mathcal{L}^- = (\mathcal{L}^+)^*$. This follows from uniqueness of \mathcal{L}^\pm (see Theorem B.1 below).

The following two theorems are established in [27, 28] for the case $p = 2$. Their generalization for the case of an arbitrary $p \geq 1$ is mentioned in [16].

Theorem B.1 *If $\mathcal{I} + \mathcal{F}$, $\mathcal{F} \in \mathcal{G}_p(M_r)$, admits a factorization in $\mathcal{G}_p(M_r)$, then the corresponding operators $\mathcal{L}^\pm = \mathcal{L}^\pm(\mathcal{F})$ in the representation (B.1) are unique. Moreover, the set of operators $\mathcal{F} \in \mathcal{G}_p(M_r)$, such that $\mathcal{I} + \mathcal{F}$ admits a factorization, is open in $\mathcal{G}_p(M_r)$ and the mappings*

$$\mathcal{G}_p(M_r) \ni \mathcal{F} \mapsto \mathcal{L}^\pm(\mathcal{F}) \in \mathcal{G}_p(M_r)$$

are continuous.

Theorem B.2 *An operator $\mathcal{I} + \mathcal{F}$, $\mathcal{F} \in \mathcal{G}_p(M_r)$, admits a factorization in $\mathcal{G}_p(M_r)$ if and only if: (A) the operators $\mathcal{I} + \chi_a \mathcal{F} \chi_a$ have trivial kernels for all $a \in [0, 1]$, where χ_a is an operator of multiplication by the indicator of the interval $[0, a]$, i.e.*

$$(\chi_a f)(x) := \begin{cases} f(x), & x \in [0, a], \\ 0, & x \in (a, 1]. \end{cases}$$

If an operator \mathcal{F} is self-adjoint, then the condition (A) is equivalent to positivity of $\mathcal{I} + \mathcal{F}$.

Now we are interested in a connection between Krein's accelerants (see Definition 1.2) and factorization of some integral operators.

Let $H \in L_p((-1, 1), M_r)$, $p \geq 1$, be an arbitrary function such that $H(-x) = H(x)^*$ for almost all $x \in (-1, 1)$. Denote by \mathcal{H} an integral operator in $L_2((0, 1), \mathbb{C}^r)$ acting by the formula

$$(\mathcal{H}f)(x) := \int_0^1 H(x-t)f(t) dt, \quad x \in (0, 1).$$

Now set

$$\mathbb{H} := L_2((0, 1), \mathbb{C}^r) \times L_2((0, 1), \mathbb{C}^r).$$

Let \mathcal{I} stand for the identity operator in $L_2((0, 1), \mathbb{C}^r)$ and \mathcal{I} for the identity operator in \mathbb{H} . Then the following lemma holds true:

Lemma B.1 *Let $H \in L_p((-1, 1), M_r)$, $H(-x) = H(x)^*$ for almost all $x \in (-1, 1)$, and let $\mathcal{F}_H \in \mathcal{G}_p(M_{2r})$ be an integral operator in \mathbb{H} with kernel $F_H \in G_p(M_{2r})$ taking the form*

$$F_H(x, t) = \frac{1}{2} \begin{pmatrix} H\left(\frac{x-t}{2}\right) & H\left(\frac{x+t}{2}\right) \\ H\left(-\frac{x+t}{2}\right) & H\left(-\frac{x-t}{2}\right) \end{pmatrix}, \quad 0 \leq x, t \leq 1.$$

Then the following statements are equivalent:

- (i) *the operator $\mathcal{I} + \mathcal{H}$ admits a factorization in $\mathcal{G}_p(M_r)$;*
- (ii) *the operator $\mathcal{I} + \mathcal{F}_H$ admits a factorization in $\mathcal{G}_p(M_{2r})$;*
- (iii) *the function H is an accelerant, i.e. $H \in \mathfrak{H}_p$;*
- (iv) *the Krein equation*

$$R(x, t) + H(x-t) + \int_0^x R(x, s)H(x-s) ds = 0, \quad (x, t) \in \Omega, \quad (\text{B.2})$$

is solvable in $G_p^+(M_r)$.

Proof. Let us establish the equivalence of (i) and (ii). Firstly, observe that both operators $\mathcal{I} + \mathcal{H}$ and $\mathcal{J} + \mathcal{F}_H$ are self-adjoint, and therefore both of them admit a factorization if and only if they are positive. Now consider the unitary transformation $V : L_2((0, 1), \mathbb{C}^r) \rightarrow \mathbb{H}$ given by the formula

$$(Vf)(t) := \frac{1}{\sqrt{2}} \begin{pmatrix} f\left(\frac{1+u}{2}\right) \\ f\left(\frac{1-u}{2}\right) \end{pmatrix}, \quad f \in L_2((0, 1), \mathbb{C}^r),$$

and verify that

$$\mathcal{J} + \mathcal{F}_H = V(\mathcal{I} + \mathcal{H})V^{-1}.$$

Therefore, the operators $\mathcal{I} + \mathcal{H}$ and $\mathcal{J} + \mathcal{F}_H$ are unitarily equivalent, and hence $\mathcal{I} + \mathcal{H} > 0$ if and only if $\mathcal{J} + \mathcal{F}_H > 0$. Thus the equivalence of (i) and (ii) follows.

The equivalence of (ii) and (iii) obviously follows from Theorem B.2 and Definition 1.2. Finally, the equivalence of (ii) and (iv) is proved in [16]. \square

For an arbitrary accelerant $H \in \mathfrak{H}_p$, a solution of the Krein equation (B.2) is unique, and we denote it by R_H . The mapping $\mathfrak{H}_p \ni H \mapsto R_H \in G_p^+(M_r)$ is continuous (see [16]).

C Riesz bases

Here we state some facts from the theory of Riesz bases.

Definition C.1 *We say that two bases $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ in a Banach space X are equivalent if there exists a bounded and boundedly invertible operator $T : X \rightarrow X$ such that $Tx_n = y_n$ for all $n \in \mathbb{N}$. A basis for a Hilbert space \mathcal{H} is called a Riesz basis if it is equivalent to some orthonormal basis for \mathcal{H} .*

We set

$$\mathcal{H} := L_2((-1, 1), \mathbb{C}^r)$$

and endow \mathcal{H} with the inner product

$$(f|g)_{\mathcal{H}} := \int_{-1}^1 (f(t)|g(t))_{\mathbb{C}^r} dt, \quad f, g \in \mathcal{H}.$$

The set \mathcal{H} with the inner product $(\cdot|\cdot)_{\mathcal{H}}$ becomes a Hilbert space. Throughout this appendix we denote by \widehat{f} the Fourier transform in \mathcal{H} given by

$$\widehat{f}(n) := \frac{1}{\sqrt{2}} \int_{-1}^1 e^{-i\pi nt} f(t) dt, \quad n \in \mathbb{Z}, \quad f \in \mathcal{H}.$$

We start from a certain analogue of well-known Kadec's 1/4-theorem (see, e.g., [30, Chapter 1]). Let $\xi := (\xi_n)_{n \in \mathbb{Z}}$ be a non-decreasing sequence of real numbers such that

$$\lim_{n \rightarrow +\infty} \xi_n = +\infty, \quad \lim_{n \rightarrow -\infty} \xi_n = -\infty,$$

and let $v := (v_n)_{n \in \mathbb{Z}}$ be a sequence of non-zero vectors in \mathbb{C}^r . Consider the system of functions

$$\mathcal{E}(\xi, v) := \{e^{i\xi_n t} v_n \mid n \in \mathbb{Z}\}$$

in the space $\widehat{\mathcal{H}}$. We are interested in finding conditions guaranteeing that the system $\mathcal{E} := \mathcal{E}(\xi, v)$ forms a Riesz basis for the space $\widehat{\mathcal{H}}$.

Definition C.2 We say that the system \mathcal{E} enjoys the condition (R_0) if for all $n \in \mathbb{Z}$ the numbers $\xi_{n,k} := \xi_{nr+k}$, $k = 1, \dots, r$, belong to the interval

$$\Delta_n := \left(\pi n - \frac{\pi}{2}, \pi n + \frac{\pi}{2} \right]$$

and the vectors $v_{n,k} := v_{nr+k}$, $k = 1, \dots, r$, form a basis for \mathbb{C}^r .

With an arbitrary system \mathcal{E} enjoying the condition (R_0) we associate two sequences $(A_n)_{n \in \mathbb{Z}}$ and $(B_n)_{n \in \mathbb{Z}}$ of bounded linear operators in \mathbb{C}^r acting by the formulae

$$A_n v_{n,k} := \xi_{n,k} v_{n,k}, \quad B_n \epsilon_k := v_{n,k}, \quad k = 1, \dots, r,$$

where $(\epsilon_k)_{k=1}^r$ is a standard orthonormal basis for \mathbb{C}^r .

Definition C.3 We say that the system \mathcal{E} enjoys the condition (R_1) if it enjoys the condition (R_0) and, moreover,

$$\sup_{n \in \mathbb{Z}} \|A_n - \pi n I\| < \ln 2$$

and

$$\sup_{n \in \mathbb{Z}} (\|B_n\| + \|B_n^{-1}\|) < \infty. \quad (\text{C.1})$$

Theorem C.1 If the system \mathcal{E} enjoys the condition (R_1) , then it forms a Riesz basis for the space \mathcal{H} .

Proof. Denote by \mathcal{E}_0 an orthonormal basis in the space \mathcal{H} given by

$$\mathcal{E}_0 := \left\{ \frac{1}{\sqrt{2}} e^{i\pi n t} \epsilon_k \mid n \in \mathbb{Z}, k = 1, \dots, r \right\}.$$

The present theorem will be proved if we show that there exists a bounded and boundedly invertible operator $S : \mathcal{H} \rightarrow \mathcal{H}$ that maps $\frac{1}{\sqrt{2}} e^{i\pi n t} \epsilon_k$ to $e^{i\xi_{n,k} t} v_{n,k}$ for all $n \in \mathbb{Z}$, $k = 1, \dots, r$.

To this end, consider the operators S and S_0 that are defined on the linear span $\text{lin } \mathcal{E}_0$ by the formulae

$$(Sf)(t) = \sum_{n=-\infty}^{\infty} e^{iA_n t} B_n \widehat{f}(n), \quad (S_0 f)(t) = \frac{1}{\sqrt{2}} \sum_{n=-\infty}^{\infty} e^{iA_n t} \widehat{f}(n),$$

$f \in \text{lin } \mathcal{E}_0$. Observe that

$$S \left(\frac{1}{\sqrt{2}} e^{i\pi n t} \epsilon_k \right) = e^{iA_n t} B_n \epsilon_k = e^{iA_n t} v_{n,k} = e^{i\xi_{n,k} t} v_{n,k}$$

for all $n \in \mathbb{Z}$, $k = 1, \dots, r$, and that the equality $S = S_0 B$ takes place, where $B : \mathcal{H} \rightarrow \mathcal{H}$ acts by the formula

$$(\widehat{Bf})(n) = \sqrt{2} B_n \widehat{f}(n), \quad n \in \mathbb{Z}.$$

Since continuity and invertibility of the operator B follows from the condition (C.1), it suffices to show that the operator

$$\mathcal{B}f := S_0 f - f, \quad f \in \text{lin } \mathcal{E}_0,$$

has the norm less than 1.

Set $\tilde{A}_n := A_n - \pi n I$, $n \in \mathbb{Z}$. Since

$$e^{iA_n t} - e^{i\pi n t I} = e^{i\pi n t} (e^{i\tilde{A}_n t} - I) = e^{i\pi n t} \sum_{k=1}^{\infty} \frac{(i\tilde{A}_n t)^k}{k!}$$

and

$$f(t) = \frac{1}{\sqrt{2}} \sum_{n=-\infty}^{\infty} e^{i\pi n t} \hat{f}(n),$$

we find that

$$\begin{aligned} \sqrt{2} \|S_0 f - f\|_{\mathcal{H}} &= \left\| \sum_{n=-\infty}^{\infty} e^{i\pi n t} (e^{i\tilde{A}_n t} - I) \hat{f}(n) \right\|_{\mathcal{H}} \\ &= \left\| \sum_{n=-\infty}^{\infty} \sum_{k=1}^{\infty} e^{i\pi n t} \frac{(i\tilde{A}_n t)^k}{k!} \hat{f}(n) \right\|_{\mathcal{H}} \leq \sum_{k=1}^{\infty} \frac{1}{k!} \left\| \sum_{n=-\infty}^{\infty} e^{i\pi n t} \tilde{A}_n^k \hat{f}(n) \right\|_{\mathcal{H}} \\ &\leq \sum_{k=1}^{\infty} \frac{\sqrt{2}}{k!} \left(\sum_{n=-\infty}^{\infty} \|\tilde{A}_n^k \hat{f}(n)\|^2 \right)^{1/2} \leq \sqrt{2} \sum_{k=1}^{\infty} \frac{\delta^k}{k!} \|f\|_{\mathcal{H}}, \end{aligned}$$

where $\delta := \sup_{n \in \mathbb{Z}} \|\tilde{A}_n\| < \ln 2$. Therefore, $\|S_0 f - f\| \leq (e^\delta - 1) \|f\|$, i.e. $\|S_0 - \mathcal{I}\| \leq e^\delta - 1 < 1$, as desired. \square

Definition C.4 Two sequences of vectors $(f_n)_{n \in \mathbb{Z}}$ and $(g_n)_{n \in \mathbb{Z}}$ in a Hilbert space are said to be quadratically close if

$$\sum_{n=-\infty}^{\infty} \|f_n - g_n\|^2 < \infty.$$

The following theorem is a simple consequence of Theorem 15 in [30, Chapter 1]:

Theorem C.2 Let $(f_n)_{n \in \mathbb{Z}}$ be a Riesz basis for \mathcal{H} . If $(g_n)_{n \in \mathbb{Z}}$ is complete and quadratically close to $(f_n)_{n \in \mathbb{Z}}$, then $(g_n)_{n \in \mathbb{Z}}$ is a Riesz basis for \mathcal{H} .

References

- [1] B. M. Levitan and I. S. Sargsjan, *Sturm–Liouville and Dirac operators* Kluwer Academic Publishers, 1991.
- [2] V. A. Marchenko, *Sturm–Liouville operators and applications* Birkhäuser, 1986.
- [3] B. Thaller, *The Dirac equation* Springer, 1992.
- [4] M. G. Gasymov and B. M. Levitan, *The inverse problem for the Dirac system* Dokl. Akad. Nauk SSSR **167** (1966), 967–70 (in Russian).
- [5] M. G. Gasymov and B. M. Levitan, *Determination of the Dirac system from the scattering phase* Dokl. Akad. Nauk SSSR **167** (1966), 1219–22 (in Russian).
- [6] M. M. Malamud, *Borg-type theorems for first-order systems on a finite interval* Funct. Anal. Appl. **33** (1999), 64–8.

- [7] M. M. Malamud, *Uniqueness questions in inverse problems for systems of differential equations on a finite interval* Trans. Moscow Math. Soc. **60** (1999), 173–224.
- [8] M. Lesch and M. Malamud, *The inverse spectral problem for first order systems on the half line* Oper. Theory Adv. Appl. **117** (2000), 199–238.
- [9] S. Clark and F. Gesztesy, *Weyl–Titchmarsh M -function asymptotics, local uniqueness results, trace formulas, and Borg-type theorems for Dirac operators* Trans. Amer. Math. Soc. **354** (2002), 3475–534.
- [10] F. Gesztesy, A. Kiselev and K. A. Makarov, *Uniqueness results for matrix-valued Schrödinger, Jacobi, and Dirac-type operators* Math. Nachr. **239/240** (2002), 103–45.
- [11] F. Gesztesy, R. Weikard and M. Zinchenko, *Initial value problems and Weyl–Titchmarsh theory for Schrödinger operators with operator-valued potentials* Preprint arXiv:1109.1613.
- [12] A. Sakhnovich, *Dirac type and canonical systems: spectral and Weyl–Titchmarsh matrix functions, direct and inverse problems* Inverse Problems **18** (2002), 331–448.
- [13] A. Sakhnovich, *Dirac type system on the axis: explicit formulae for matrix potentials with singularities and soliton-positon interactions* Inverse Problems **19** (2003), 845–54.
- [14] D. Chelkak and E. Korotyaev, *Parametrization of the isospectral set for the vector-valued Sturm–Liouville problem* J. Funct. Anal. **241** (2006), 359–73.
- [15] D. Chelkak and E. Korotyaev, *Weyl–Titchmarsh functions of vector-valued Sturm–Liouville operators on the unit interval* J. Funct. Anal. **257** (2009), 1546–88.
- [16] S. Albeverio, R. Hryniv and Ya. Mykytyuk, *Inverse spectral problems for Dirac operators with summable potentials* Russ. J. Math. Phys. **12** (2005), 406–23.
- [17] Ya. V. Mykytyuk and N. S. Trush, *Inverse spectral problems for Sturm–Liouville operators with matrix-valued potentials* Inverse Problems **26** (2010), 015009.
- [18] Ya. V. Mykytyuk and D. V. Puyda, *Inverse spectral problems for Dirac operators on a finite interval* J. Math. Anal. Appl. **386** (2012), 177–94.
- [19] N. Trush, *Asymptotics of singular values of entire matrix-valued sine-type functions* Mat. Stud. **30** (2008), 95–7.
- [20] Ya. V. Mykytyuk and N. S. Trush, *Asymptotics of zeros for some entire sine-type functions acting in a Banach algebra* Math. Bull. Shevchenko Sci. Soc. **4** (2007), 214–9.
- [21] R. O. Hryniv and Ya. V. Mykytyuk, *On zeros of some entire functions* Trans. Amer. Math. Soc. **361** (2009), 2207–23.
- [22] P. Kuchment, *Quantum graphs: an introduction and a brief survey* Analysis on graphs and its applications, Proc. Sympos. Pure Math. **77** (2008), 291–312.
- [23] N. Trush, *Solutions of the Cauchy problem for factorized Sturm–Liouville equation in a Banach algebra* Mat. Stud. **31** (2008), 75–82.
- [24] D. Alpay, I. Gohberg, M. A. Kaashoek, L. Lerer and A. L. Sakhnovich, *Krein systems and canonical systems on a finite interval: Accelerants with a jump discontinuity at the origin and continuous potentials* Integr. Equ. Oper. Theory **68(1)** (2010), 115–50.

- [25] F. Gesztesy and S. Tsekanovskii, *On matrix-valued Herglotz functions* Math. Nachr. **218** (2000), 61–138.
- [26] V. A. Sadovnichiy, *Theory of operators* Consultants Bureau, 1991.
- [27] Ya. V. Mykytyuk, *Factorization of Fredholm operators* Mat. Stud. **20(2)** (2003), 185–99 (in Ukrainian).
- [28] Ya. V. Mykytyuk, *Factorization of Fredholm operators in operator algebras* Mat. Stud. **21(1)** (2004), 87–97 (in Ukrainian).
- [29] I. Gokhberg and M. Krein, *Theory of Volterra operators in Hilbert space and its applications* Nauka, 1967 (in Russian).
- [30] R. M. Young, *An introduction to nonharmonic Fourier series* Revised 1st Edition, Academic Press, 2001.