PARTIAL REGULARITY FOR MINIMIZERS OF QUASICONVEX FUNCTIONALS WITH GENERAL GROWTH

LARS DIENING¹, DANIEL LENGELER², BIANCA STROFFOLINI³, AND ANNA VERDE⁴

ABSTRACT. We prove a partial regularity result for local minimizers of quasiconvex variational integrals with general growth. The main tool is an improved \mathcal{A} -harmonic approximation, which should be interesting also for classical growth.

1. INTRODUCTION

In this paper we study partial regularity for vector-valued minimizers $u: \Omega \to \mathbb{R}^N$ of variational integrals:

(1.1)
$$\mathcal{F}(u) := \int_{\Omega} f(\nabla u) \, dx,$$

where $\Omega \subset \mathbb{R}^n$ is a domain and $f : \mathbb{R}^{N \times n} \to \mathbb{R}$ is a continuous function.

Let us recall Morrey's notion of quasiconvexity [26]:

Definition 1.1. f is called quasiconvex if and only if

(1.2)
$$\int_{B_1} f(A + \nabla \boldsymbol{\xi}) \, dx \ge f(A)$$

holds for every $A \in \mathbb{R}^{nN}$ and every smooth $\boldsymbol{\xi} : B_1 \to \mathbb{R}^N$ with compact support in the open unit ball B_1 in \mathbb{R}^n .

By Jensen's inequality, quasiconvexity is a generalization of convexity. It was originally introduced as a notion for proving the lower semicontinuity and the existence of minimizers of variational integrals. In fact, assuming a power growth condition, quasiconvexity is proved to be a necessary and sufficient condition for the sequential weak lower semicontinuity on $W^{1,p}(\Omega.\mathbb{R}^N)$, p > 1, see [25] and [1]. For general growth condition see [20] and [31]. In the regularity issue, a stronger definition comes into play. In the fundamental paper [19] Evans considered strictly quasi-convex integrands f in the quadratic case and proved that if f is of class C^2 and has bounded second derivatives then any minimizing function \mathbf{u} is of class $C^{1,\alpha}(\Omega \setminus \Sigma)$ where Σ has *n*-dimensional Lebesgue measure zero. In [1], this result was generalized to integrands f of p-growth with $p \geq 2$ while the subquadratic growth was considered in [6].

In order to treat the general growth case, we introduce the notion of strictly $W^{1,\varphi}$ quasiconvex function, where φ is a suitable N-function, see Assumption 2.2.

¹⁹⁹¹ Mathematics Subject Classification. 35J60, 35J70, 49N60, 26B25.

Key words and phrases. quasi-convex, partial regularity, harmonic approximation, Lipschitz truncation.

Definition 1.2. The function f is strictly $W^{1,\varphi}$ - quasiconvex if and only if

$$\int_{B} f(\mathbf{Q} + \nabla \mathbf{w}) - f(\mathbf{Q}) \, dx \ge k \int_{B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{w}|) \, dx.$$

for all balls $B \subset \Omega$, all $\mathbf{Q} \in \mathbb{R}^{N \times n}$ and all $\mathbf{w} \in C_0^1(B)$, where $\varphi_a(t) \sim \varphi''(a+t) t^2$ for $a, t \geq 0$. A precise definition of φ_a is given in Section 2.

We will work with the following set of assumptions:

- (H1) $f \in C^1(\mathbb{R}^n) \cap C^2(\mathbb{R}^n \setminus \{0\}),$
- (H2) for all $\mathbf{Q} \in \mathbb{R}^{N \times n}$ it holds

$$|f(\mathbf{Q})| \le K\varphi(|\mathbf{Q}|),$$

- (H3) the function f is strictly $W^{1,\varphi}$ -quasiconvex;
- (H4) for all $\mathbf{Q} \in \mathbb{R}^{N \times n} \setminus \{\mathbf{0}\}$

$$\left| (D^2 f)(\mathbf{Q}) \right| \le c \, \varphi''(|\mathbf{Q}|)$$

(H5) the following Hölder continuity of $D^2 f$ away from 0

$$\left| D^2 f(\mathbf{Q}) - D^2 f(\mathbf{Q} + \mathbf{P}) \right| \le c \, \varphi''(|\mathbf{Q}|) |\mathbf{Q}|^{-\beta} |\mathbf{P}|^{\beta}$$

holds for all $\mathbf{P}, \mathbf{Q} \in \mathbb{R}^{N \times n}$ such that $|\mathbf{P}| \leq \frac{1}{2} |\mathbf{Q}|$.

Due to **(H2)**, \mathcal{F} is well defined on the Sobolev-Orlicz space $W^{1,\varphi}(\Omega, \mathbb{R}^N)$, see section 2. Let us observe that assumption **(H5)** has been used to show everywhere regularity of radial functionals with φ -growth, [12]. Following the argument given in [22] it is possible to prove that **(H3)** implies the following *strong Legendre-Hadamard condition*

$$(D^2 f)(\mathbf{Q})(\boldsymbol{\eta} \otimes \boldsymbol{\xi}, \boldsymbol{\eta} \otimes \boldsymbol{\xi}) \ge c \, \varphi''(|\mathbf{Q}|) |\boldsymbol{\eta}|^2 |\boldsymbol{\xi}|^2$$

for all $\eta \in \mathbb{R}^N$, $\xi \in \mathbb{R}^n$ and $\mathbf{Q} \in \mathbb{R}^{N \times n} \setminus \{\mathbf{0}\}$. Furthermore, (H3) implies that the functional

$$\mathcal{J}(t) := \int_{B} f(\mathbf{Q} + t\nabla \mathbf{w}) - f(\mathbf{Q}) - k\varphi_{|\mathbf{Q}|}(t|\nabla \mathbf{w}|) \, dx$$

attains its minimal value at t = 0. Hence $\mathcal{J}''(0) \ge 0$, that is

(1.3)
$$\int_{B} (D^{2}f)(\mathbf{Q})(\nabla \mathbf{w}, \nabla \mathbf{w}) \, dx \ge k \int_{B} \varphi_{|\mathbf{Q}|}'(0) |\nabla \mathbf{w}|^{2} \, dx \ge c \, \varphi''(|\mathbf{Q}|) \int_{B} |\nabla \mathbf{w}|^{2} \, dx.$$

As usual, the strategy for proving partial regularity consists in showing an excess decay estimate, where the *excess* function is

(1.4)
$$\Phi_s(B,\mathbf{u}) := \left(\oint_B |\mathbf{V}(\nabla \mathbf{u}) - \langle \mathbf{V}(\nabla \mathbf{u}) \rangle_B |^{2s} \, dx \right)^{\frac{1}{s}}$$

with $\mathbf{V}(\mathbf{Q}) = \sqrt{\frac{\varphi'(|\mathbf{Q}|)}{|\mathbf{Q}|}} \mathbf{Q}$ and $s \ge 1$. We write $\Phi := \Phi_1$. Note that $\Phi_{s_1}(B, \mathbf{u}) \le \Phi_{s_2}(B, \mathbf{u})$ for $1 \le s_1 \le s_2$ and $|\mathbf{V}(\mathbf{Q})|^2 \sim \varphi(|\mathbf{Q}|)$.

Our regularity theorem states:

Theorem 1.3 (Main theorem). Let **u** be a local minimizer of the quasiconvex functional (1.1), with f satisfying (H1)–(H5) and fix some $\beta \in (0,1)$. Then there exists $\delta = \delta(\beta) > 0$ such that the following holds: If

(1.5)
$$\Phi(2B,\mathbf{u}) \le \delta \oint_{2B} |\mathbf{V}(\nabla \mathbf{u})|^2 \, dx$$

for some ball $B \subset \mathbb{R}^n$ with $2B \subset \Omega$, then $\mathbf{V}(\nabla \mathbf{u})$ is β -Hölder continuous on B.

The proof of this theorem can be found at the end of Section 6. We define the set of regular points $\mathcal{R}(\mathbf{u})$ by

(1.6)
$$\mathcal{R}(\mathbf{u}) = \left\{ x_0 \in \Omega : \liminf_{r \to 0} \Phi(B(x_0, r), \mathbf{u}) = 0 \right\}.$$

As an immediate consequence of Theorem 1.3 we have:

Corollary 1.4. Let **u** be as in Theorem 1.3 and let $x_0 \in \mathcal{R}(\mathbf{u})$ with $\nabla \mathbf{u} \neq 0$. Then for every $\beta \in (0, 1)$ the function $\mathbf{V}(\nabla \mathbf{u})$ is β -Hölder continuous on a neighborhood of x_0 .

Note that the Hölder continuity of $\mathbf{V}(\nabla \mathbf{u})$ implies the Hölder continuity of $\nabla \mathbf{u}$ with a different exponent depending on φ . Consider for example the situation $\varphi(t) = t^p$ with $1 . Therefore, <math>\beta$ -Hölder continuity of $\mathbf{V}(\nabla \mathbf{u})$ implies for $p \leq 2$ that $\nabla \mathbf{u}$ is β -Hölder continuous and for p > 2 that $\nabla \mathbf{u}$ is $\beta \frac{2}{p}$ -Hölder continuous.

The proofs of the regularity results for local minimizers in [19],[1], [6], are based on a blow-up technique originally developed by De Giorgi [7] and Almgren [3], [4] in the setting of the geometric measure theory, and by Giusti and Miranda for elliptic systems, [23].

Another more recent approach for proving partial regularity for local minimizers is based on the so called \mathcal{A} -harmonic approximation method. This technique has its origin in Simon's proof of the regularity theorem [30] (see also Allard [2]). The technique has been successfully applied in the framework of the geometric measure theory, and to obtain partial-regularity results for general elliptic systems in a series of papers by Duzaar, Grotowski, Kronz, Mingione [15] [16], [17], [18]. More precisely, we consider a bilinear form on $\operatorname{Hom}(\mathbb{R}^n, \mathbb{R}^N)$ which is (strongly) elliptic in the sense of Legendre-Hadamard, i.e. if for all $\mathbf{a} \in \mathbb{R}^N$, $\mathbf{b} \in \mathbb{R}^n$ it holds

$$\mathcal{A}_{ij}^{lphaeta}a^{i}b_{lpha}a^{j}b_{eta}\geq\kappa_{\mathcal{A}}|\mathbf{a}|^{2}|\mathbf{b}|^{2}$$

for some $\kappa_{\mathcal{A}} > 0$. The method of \mathcal{A} -harmonic approximation consists in obtaining a good approximation of functions $\mathbf{u} \in W^{1,2}(B)$, which are *almost* \mathcal{A} -harmonic (in the sense of Theorem 4.1) by \mathcal{A} -harmonic functions $\mathbf{h} \in W^{1,2}(B)$, in both the L^2 -topology and in the weak topology of $W^{1,2}$. Let us recall that $\mathbf{h} \in W^{1,2}(B)$ is called \mathcal{A} -harmonic on B if

(1.7)
$$\int_{B} \mathcal{A}(D\mathbf{h}, D\boldsymbol{\eta}) \, dx = 0, \forall \boldsymbol{\eta} \in C_0^{\infty}(B)$$

holds. Here, in order to prove the result, we will follow the second approach.

As in the situations considered in the above-mentioned papers, the required approximate \mathcal{A} -harmonicity of a local minimizer $\mathbf{u} \in W^{1,\varphi}(\Omega \setminus \Sigma)$ is a consequence of the minimizing property and of the *smallness* of the *excess*.

Next, having proven the \mathcal{A} -harmonic approximation lemma and the corresponding approximate \mathcal{A} -harmonicity of the local minimizer \mathbf{u} , the other steps are quite standard. We prove a Caccioppoli-type inequality for minimizers \mathbf{u} and thus we compare \mathbf{u} with the

 \mathcal{A} -harmonic approximation **h** to obtain, via our Caccioppoli-type inequality, the desired excess decay estimate.

Thus, the main difficulty is to establish a suitable version of the \mathcal{A} -harmonic approximation lemma in this general setting. However, let us point out that our \mathcal{A} -harmonic approximation lemma differs also in the linear or p-growth situation from the classical one in [17]. Firstly, we use a direct approach based on the Lipschitz truncation technique which requires no contradiction argument. This allows for a precise control of the constants, which will only depend on the Δ_2 -condition for φ and its conjugate. In fact, we will apply the approximation lemma to the family of shifted N-functions that inherit the same Δ_2 constants of φ . Secondly, we are able to preserve the boundary values of our original function, so $\mathbf{u} - \mathbf{h}$ is a valid test function. Thirdly, we show that \mathbf{h} and \mathbf{u} are close with respect to the gradients rather than just the functions. The main tools in the proof is a Lipschitz approximation of the Sobolev functions as in [11, 5]. However, since \mathcal{A} is only strongly elliptic in the sense of Legendre-Hadamard, we will not be able to apply the Lipschitz truncation technique directly to our almost \mathcal{A} -harmonic function **u**. Instead, we need to use duality and apply the Lipschitz truncation technique to the test functions.

Let us conclude by observing that here we are able to present a unified approach for both cases: superquadratic and subquadratic growth.

2. NOTATION AND PRELIMINARY RESULTS

We use c, C as generic constants, which may change from line to line, but does not depend on the crucial quantities. Moreover we write $f \sim g$ iff there exist constants c, C > 0 such that $cf \leq g \leq Cf$. For $w \in L^1_{loc}(\mathbb{R}^n)$ and a ball $B \subset \mathbb{R}^n$ we define

(2.8)
$$\langle w \rangle_B := \oint_B w(x) \, dx := \frac{1}{|B|} \int_B w(x) \, dx,$$

where |B| is the *n*-dimensional Lebesgue measure of B. For $\lambda > 0$ we denote by λB the ball with the same center as B but λ -times the radius. For $U, \Omega \subset \mathbb{R}^n$ we write $U \subseteq \Omega$ if the closure of U is a compact subset of Ω .

The following definitions and results are standard in the context of N-functions, see for example [24, 28]. A real function $\varphi : \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ is said to be an N-function if it satisfies the following conditions: $\varphi(0) = 0$ and there exists the derivative φ' of φ . This derivative is right continuous, non-decreasing and satisfies $\varphi'(0) = 0$, $\varphi'(t) > 0$ for t > 0, and $\lim_{t\to\infty} \varphi'(t) = \infty$. Especially, φ is convex.

We say that φ satisfies the Δ_2 -condition, if there exists c > 0 such that for all $t \ge 0$ holds $\varphi(2t) \leq c \varphi(t)$. We denote the smallest possible constant by $\Delta_2(\varphi)$. Since $\varphi(t) \leq \varphi(t) \leq \varphi(t)$ $\varphi(2t)$ the Δ_2 condition is equivalent to $\varphi(2t) \sim \varphi(t)$.

By L^{φ} and $W^{1,\varphi}$ we denote the classical Orlicz and Sobolev-Orlicz spaces, i. e. $f \in L^{\varphi}$ iff $\int \varphi(|f|) dx < \infty$ and $f \in W^{1,\varphi}$ iff $f, \nabla f \in L^{\varphi}$. By $W_0^{1,\varphi}(\Omega)$ we denote the closure of $\begin{array}{l} C_0^{\infty}(\Omega) \text{ in } W^{1,\varphi}(\Omega). \\ \text{ By } (\varphi')^{-1} : \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0} \text{ we denote the function} \end{array}$

$$(\varphi')^{-1}(t) := \sup \{ s \in \mathbb{R}^{\ge 0} : \varphi'(s) \le t \}.$$

If φ' is strictly increasing then $(\varphi')^{-1}$ is the inverse function of φ' . Then $\varphi^* : \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ with

$$\varphi^*(t) := \int_0^t (\varphi')^{-1}(s) \, ds$$

is again an N-function and $(\varphi^*)'(t) = (\varphi')^{-1}(t)$ for t > 0. It is the complementary function of φ . Note that $\varphi^*(t) = \sup_{s \ge 0} (st - \varphi(s))$ and $(\varphi^*)^* = \varphi$. For all $\delta > 0$ there exists c_{δ} (only depending on $\Delta_2(\varphi, \varphi^*)$ such that for all $t, s \ge 0$ holds

(2.9)
$$t s \le \delta \varphi(t) + c_{\delta} \varphi^*(s),$$

For $\delta = 1$ we have $c_{\delta} = 1$. This inequality is called Young's inequality. For all $t \ge 0$

(2.10)
$$\frac{\frac{t}{2}\varphi'\left(\frac{t}{2}\right) \le \varphi(t) \le t\,\varphi'(t),}{\varphi\left(\frac{\varphi^*(t)}{t}\right) \le \varphi^*(t) \le \varphi\left(\frac{2\,\varphi^*(t)}{t}\right).}$$

Therefore, uniformly in $t \ge 0$

(2.11)
$$\varphi(t) \sim \varphi'(t) t, \qquad \varphi^*(\varphi'(t)) \sim \varphi(t),$$

where the constants only depend on $\Delta_2(\varphi, \varphi^*)$.

We say that a N-function ψ is of type (p_0, p_1) with $1 \le p_0 \le p_1 < \infty$, if

(2.12)
$$\psi(st) \le C \max\{s^{p_0}, s^{p_1}\}\psi(t)$$
 for all $s, t \ge 0$.

We also write $\psi \in \mathfrak{T}(p_0, p_1, C)$.

Lemma 2.1. Let ψ be an N-function with $\psi \in \Delta_2$ together with its conjugate. Then $\psi \in \mathfrak{T}(p_0, p_1, C_1)$ for some $1 < p_0 < p_1 < \infty$ and $C_1 > 0$, where p_0, p_1 and C_1 only depend on $\Delta_2(\psi, \psi^*)$. Moreover, ψ has the representation

(2.13)
$$\psi(t) = t^{p_0} (h(t))^{p_1 - p_0} \quad \text{for all } t \ge 0,$$

where h is a quasi-concave function, i.e.

$$h(\lambda t) \le C_2 \max\{1, \lambda\} h(t)$$
 for all $\lambda, t \ge 0$,

where C_2 only depends on $\Delta_2(\psi, \psi^*)$.

Proof. Let $K := \Delta_2(\psi)$ and $K_* := \max \{\Delta_2(\psi^*), 3\}$. Then $\psi^*(2t) \leq K_*\psi^*(t)$ for all $t \geq 0$ implies $\psi(t) \leq K_*\psi(2t/K_*)$ for all $t \geq 0$. Now, choose p_0, p_1 such that $1 < p_0 < p_1 < \infty$ and $K \leq 2^{p_0}$ and $(K_*/2)^{p_0} \leq K_*$. We claim that

(2.14)
$$\psi(st) \le C \max\{s^{p_0}, s^{p_1}\}\psi(t)$$
 for all $s, t \ge 0$,

where C only depends on K and K_* . Indeed, if $s \ge 1$, then choose $m \ge 0$ such that $2^m \le s \le 2^{m+1}$. Using $\psi \in \Delta_2$, we get

(2.15)
$$\psi(st) \le \psi(2^{m+1}t) \le K^{m+1}\psi(t) \le K(2^{p_1})^m\psi(t) \le Ks^{p_1}\psi(t).$$

If $s \leq 1$, then we choose $m \in \mathbb{N}_0$ such that $(K_*/2)^m s \leq 1 \leq (K_*/2)^{m+1} s$, so that

$$\psi(st) \le K_*^m \psi\left(\left(\frac{2}{K_*}\right)^m st\right) \le K_*\left(\frac{K_*}{2}\right)^{p_0(m-1)} \psi(t) \le K_* s^{p_0} \psi(t).$$

This proves (2.14).

Now, let us define

$$h(u) := \psi\left(u^{\frac{1}{p_1 - p_0}}\right) u^{-\frac{p_0}{p_1 - p_0}},$$

then ψ satisfies (2.13). It remains to show that h is quasi-concave. We estimate with (2.14)

$$h(su) \le K \,\psi\left(u^{\frac{1}{p_1 - p_0}}\right) \max\left\{s^{\frac{p_1}{p_1 - p_0}}, s^{\frac{p_0}{p_1 - p_0}}\right\} (su)^{\frac{-p_0}{p_1 - p_0}} = K\psi(u) \max\left\{s, 1\right\}$$

for all $s, u \ge 0$.

Throughout the paper we will assume that φ satisfies the following assumption.

Assumption 2.2. Let φ be an N-function such that φ is C^1 on $[0,\infty)$ and C^2 on $(0,\infty)$. Further assume that

(2.16)
$$\varphi'(t) \sim t \,\varphi''(t)$$

uniformly in t > 0. The constants in (2.16) are called the characteristics of φ .

We remark that under these assumptions $\Delta_2(\varphi, \varphi^*) < \infty$ will be automatically satisfied, where $\Delta_2(\varphi, \varphi^*)$ depends only on the characteristics of φ .

For given φ we define the associated N-function ψ by

(2.17)
$$\psi'(t) := \sqrt{\varphi'(t) t}.$$

It is shown in [8, Lemma 25] that if φ satisfies Assumption 2.2, then also φ^* , ψ , and ψ^* satisfy this assumption.

Define $\mathbf{A}, \mathbf{V} : \mathbb{R}^{N \times n} \to \mathbb{R}^{N \times n}$ in the following way:

(2.18a)
$$\mathbf{A}(\mathbf{Q}) = \varphi'(|\mathbf{Q}|) \frac{\mathbf{Q}}{|\mathbf{Q}|},$$

(2.18b)
$$\mathbf{V}(\mathbf{Q}) = \psi'(|\mathbf{Q}|) \frac{\mathbf{Q}}{|\mathbf{Q}|}$$

Another important set of tools are the shifted N-functions $\{\varphi_a\}_{a\geq 0}$ introduced in [8], see also [10, 29]. We define for $t\geq 0$

(2.19)
$$\varphi_a(t) := \int_0^t \varphi_a'(s) \, ds \quad \text{with} \quad \varphi_a'(t) := \varphi'(a+t) \frac{t}{a+t}.$$

Note that $\varphi_a(t) \sim \varphi'_a(t) t$. Moreover, for $t \geq a$ we have $\varphi_a(t) \sim \varphi(t)$ and for $t \leq a$ we have $\varphi_a(t) \sim \varphi''(a)t^2$. This implies that $\varphi_a(st) \leq c s^2 \varphi_a(t)$ for all $s \in [0,1]$, $a \geq 0$ and $t \in [0,a]$. The families $\{\varphi_a\}_{a\geq 0}$ and $\{(\varphi_a)^*\}_{a\geq 0}$ satisfy the Δ_2 -condition uniformly in $a \geq 0$.

The connection between **A**, **V** and the shifted functions of φ is best reflected in the following lemma [12, Lemma 2.4], see also [8].

Lemma 2.3. Let φ satisfy Assumption 2.2 and let A and V be defined by (2.18). Then

$$\begin{split} \left(\mathbf{A}(\mathbf{P}) - \mathbf{A}(\mathbf{Q})\right) \cdot \left(\mathbf{P} - \mathbf{Q}\right) &\sim \left|\mathbf{V}(\mathbf{P}) - \mathbf{V}(\mathbf{Q})\right|^2 \sim \varphi_{|\mathbf{P}|}(|\mathbf{P} - \mathbf{Q}|), \\ \left|\mathbf{A}(\mathbf{P}) - \mathbf{A}(\mathbf{Q})\right| &\sim \varphi'_{|\mathbf{P}|}(|\mathbf{P} - \mathbf{Q}|), \end{split}$$

uniformly in $\mathbf{P}, \mathbf{Q} \in \mathbb{R}^{N \times n}$. Moreover,

$$\mathbf{A}(\mathbf{Q}) \cdot \mathbf{Q} \sim |\mathbf{V}(\mathbf{Q})|^2 \sim \varphi(|\mathbf{Q}|),$$

6

uniformly in $\mathbf{Q} \in \mathbb{R}^{N \times n}$.

We state a generalization of Lemma 2.1 in [1] to the context of convex functions φ .

Lemma 2.4 (Lemma 20, [8]). Let φ be an N-function with $\Delta_2(\varphi, \varphi^*) < \infty$. Then uniformly for all $\mathbf{P}_0, \mathbf{P}_1 \in \mathbb{R}^{N \times n}$ with $|\mathbf{P}_0| + |\mathbf{P}_1| > 0$ holds

(2.20)
$$\int_{0}^{1} \frac{\varphi'(|\mathbf{P}_{\theta}|)}{|\mathbf{P}_{\theta}|} d\theta \sim \frac{\varphi'(|\mathbf{P}_{0}| + |\mathbf{P}_{1}|)}{|\mathbf{P}_{0}| + |\mathbf{P}_{1}|},$$

where $\mathbf{P}_{\theta} := (1 - \theta)\mathbf{P}_0 + \theta\mathbf{P}_1$. The constants only depend on $\Delta_2(\varphi, \varphi^*)$.

Note that (H5) and the previous Lemma imply that

(2.21)
$$\left| (Df)(\mathbf{Q}) - (Df)(\mathbf{P}) \right| = \left| \int_{0}^{1} (D^{2}f)(\mathbf{P} + t(\mathbf{Q} - \mathbf{P}))(\mathbf{Q} - \mathbf{P}) dt \right|$$
$$\leq c \int_{0}^{1} \varphi''(|\mathbf{P} + t(\mathbf{Q} - \mathbf{P}))|) dt |\mathbf{P} - \mathbf{Q}|$$
$$\leq c \varphi''(|\mathbf{P}| + |\mathbf{Q}|)|\mathbf{P} - \mathbf{Q}|$$
$$\leq c \varphi'_{|\mathbf{Q}|}(|\mathbf{P} - \mathbf{Q}|).$$

The following version of Sobolev-Poincaré inequality can be found in [8, Lemma 7].

Theorem 2.5 (Sobolev-Poincaré). Let φ be an N-function with $\Delta_2(\varphi, \varphi^*) < \infty$. Then there exist $0 < \alpha < 1$ and K > 0 such that the following holds. If $B \subset \mathbb{R}^n$ is some ball with radius R and $\mathbf{w} \in W^{1,\varphi}(B, \mathbb{R}^N)$, then

(2.22)
$$\int_{B} \varphi\left(\frac{|\mathbf{w} - \langle \mathbf{w} \rangle_{B}|}{R}\right) dx \leq K \left(\int_{B} \varphi^{\alpha}(|\nabla \mathbf{w}|) dx\right)^{\frac{1}{\alpha}},$$

where $\langle \mathbf{w} \rangle_B := \int_B \mathbf{w}(x) \, dx$.

3. Caccioppoli estimate

We need the following simple modification of lemma 3.1, (Chap. 5) from [21].

Lemma 3.1. Let ψ be an N-function with $\psi \in \Delta_2$, let r > 0 and $h \in L^{\psi}(B_{2r}(x_0))$. Further, let $f : [r/2, r] \to [0, \infty)$ be a bounded function such that for all $\frac{r}{2} < s < t < r$

$$f(s) \le \theta f(t) + A \int_{B_t(x_0)} \psi\left(\frac{|h(y)|}{t-s}\right) dy$$

where A > 0 and $\theta \in [0, 1)$. Then

$$f\left(\frac{r}{2}\right) \le c(\theta, \Delta_2(\psi)) A \int_{B_{2r}(x_0)} \psi\left(\frac{|h(y)|}{2r}\right) dy.$$

Proof. Since $\psi \in \Delta_2$, there exists $C_2 > 0$ and $p_1 < \infty$ (both depending only on $\Delta_2(\psi)$) such that $\psi(\lambda u) \leq C_2 \lambda^{p_1} \psi(u)$ for all $\lambda \geq 1$ and $u \geq 0$ (compare (2.15) of Lemma 2.1). This implies

$$f(t) \le \theta f(s) + A \int_{B_s(x_0)} \psi \left(\frac{|h(y)|}{2r}\right) dy \ C_2(2r)^{p_1} (s-t)^{-p_1}.$$

Now Lemma 3.1 in [21], with $\alpha := p_1$ implies

$$f\left(\frac{r}{2}\right) \le c(\theta, p_1) A \int\limits_{B_s(x_0)} \psi\left(\frac{|h(y)|}{2r}\right) dy \ C_2(2r)^{p_1} r^{-p_1},$$

which proves the claim.

Theorem 3.2. Let $\mathbf{u} \in W^{1,\varphi}_{loc}(\Omega)$ be a local minimizer of \mathcal{F} and B be a ball with radius R such that $2B \subset \subset \Omega$. Then

$$\int_{B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) \, dx \le c \int_{2B} \varphi_{|\mathbf{Q}|}\left(\frac{|\mathbf{u} - \mathbf{q}|}{R}\right) \, dx$$

for all $\mathbf{Q} \in \mathbb{R}^{N \times n}$ and all linear polynomials \mathbf{q} on \mathbb{R}^n with values in \mathbb{R}^N and $\nabla \mathbf{q} = \mathbf{Q}$, where c only depends on n, N, k, K and the characteristics of φ .

Proof. Let 0 < s < t. Further, let B_s and B_t be balls in Ω with the same center and with radius s and t, respectively. Choose $\eta \in C_0^{\infty}(B_t)$ with $\chi_{B_s} \leq \eta \leq \chi_{B_t}$ and $|\nabla \eta| \leq c/(t-s)$. Now, define $\boldsymbol{\xi} := \eta(\mathbf{u} - \mathbf{q})$ and $\mathbf{z} := (1 - \eta)(\mathbf{u} - \mathbf{q})$. Then $\nabla \boldsymbol{\xi} + \nabla \mathbf{z} = \nabla \mathbf{u} - \mathbf{Q}$. Consider

$$I := \int_{B_t} f(\mathbf{Q} + \nabla \boldsymbol{\xi}) - f(\mathbf{Q}) \, dx.$$

Then by the quasi-convexity of f, see (H3), follows

$$I \ge c \int\limits_{B_t} \varphi_{|\mathbf{Q}|}(|\nabla \boldsymbol{\xi}|) \, dx.$$

On the other hand since $\nabla \boldsymbol{\xi} + \nabla \mathbf{z} = \nabla \mathbf{u} - \mathbf{Q}$ we get

$$\begin{split} I &= \int\limits_{B_t} f(\mathbf{Q} + \nabla \boldsymbol{\xi}) - f(\mathbf{Q}) \, dx \\ &= \int\limits_{B_t} f(\mathbf{Q} + \nabla \boldsymbol{\xi}) - f(\mathbf{Q} + \nabla \boldsymbol{\xi} + \nabla \mathbf{z}) \, dx \\ &+ \int\limits_{B_t} f(\nabla \mathbf{u}) - f(\nabla \mathbf{u} - \nabla \boldsymbol{\xi}) \, dx \\ &+ \int\limits_{B_t} f(\mathbf{Q} + \nabla \mathbf{z}) - f(\mathbf{Q}) \, dx \\ &=: II + III + IV. \end{split}$$

Since **u** is a local minimizer, we know that $(III) \leq 0$. Moreover,

$$II + IV = \int_{B_t} \int_0^1 \left((Df)(\mathbf{Q} + t\nabla \mathbf{z}) - (Df)(\mathbf{Q} + \nabla \boldsymbol{\xi} - t\nabla \mathbf{z}) \right) \nabla \mathbf{z} \, dt \, dx$$
$$= \int_{B_t} \int_0^1 \left((Df)(\mathbf{Q} + t\nabla \mathbf{z}) - (Df)(\mathbf{Q}) \nabla \mathbf{z} \, dt \, dx$$
$$- \int_{B_t} \int_0^1 \left((Df)(\mathbf{Q} + \nabla \boldsymbol{\xi} - t\nabla \mathbf{z}) - (Df)(\mathbf{Q}) \right) \nabla \mathbf{z} \, dt \, dx.$$

This proves

$$|II| + |IV| \le c \int_{B_t} \int_0^1 \varphi'_{|\mathbf{Q}|}(t|\nabla \mathbf{z}|) dt |\nabla \mathbf{z}| dx$$
$$+ c \int_{B_t} \int_0^1 \varphi'_{|\mathbf{Q}|}(|\nabla \boldsymbol{\xi} - t\nabla \mathbf{z}|) dt |\nabla \mathbf{z}| dx$$

Using $\varphi'_{|\mathbf{Q}|}(|\nabla \boldsymbol{\xi} - t\nabla \mathbf{z}|) \le c \, \varphi'_{|\mathbf{Q}|}(|\nabla \boldsymbol{\xi}|) + c \, \varphi'_{|\mathbf{Q}|}(|\mathbf{z}|)$, we get

$$|II| + |IV| \le c \int_{B_t} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{z}|) \, dx + c \int_{B_t} \varphi'_{|\mathbf{Q}|}(|\nabla \boldsymbol{\xi}|) |\nabla \mathbf{z}| \, dx$$
$$\le c \int_{B_t} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{z}|) \, dx + \frac{1}{2}(I),$$

where we have used Young's inequality in the last step. Overall, we have shown the a priori estimate

(3.23)
$$\int_{B_t} \varphi_{|\mathbf{Q}|}(|\nabla \boldsymbol{\xi}|) \, dx \le c \int_{B_t} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{z}|) \, dx.$$

Note that $\nabla \mathbf{z} = (1 - \eta)(\nabla \mathbf{u} - \mathbf{Q}) - \nabla \eta(\mathbf{u} - \mathbf{q})$, which is zero outside $B_t \setminus B_s$. Hence,

$$\int_{B_t} \varphi_{|\mathbf{Q}|}(|\nabla \boldsymbol{\xi}|) \, dx \le c \int_{B_t \setminus B_s} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) \, dx + c \int_{B_t} \varphi_{|\mathbf{Q}|}\left(\frac{|\mathbf{u} - \mathbf{q}|}{t - s}\right) \, dx.$$

Since $\eta = 1$ on B_s , we get

$$\int_{B_s} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) \, dx \le c \int_{B_t \setminus B_s} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) \, dx + c \int_{B_t} \varphi_{|\mathbf{Q}|}\left(\frac{|\mathbf{u} - \mathbf{q}|}{t - s}\right) \, dx.$$

The hole-filling technique proves

$$\int_{B_s} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) \, dx \le \lambda \int_{B_t} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) \, dx + c \int_{B_t} \varphi_{|\mathbf{Q}|}\left(\frac{|\mathbf{u} - \mathbf{q}|}{t - s}\right) \, dx$$

for some $\lambda \in (0,1)$, which is independent of **Q** and **q**. Now Lemma 3.1 proves the claim.

Corollary 3.3. There exists $0 < \alpha < 1$ such that for all local minimizers $\mathbf{u} \in W^{1,\varphi}_{loc}(\Omega)$ of \mathcal{F} , all balls B with $2B \subset \subset \Omega$ and all $\mathbf{Q} \in \mathbb{R}^{N \times n}$

$$\oint_{B} |\mathbf{V}(\nabla \mathbf{u}) - \mathbf{V}(\mathbf{Q})|^2 \, dx \le c \left(\oint_{2B} |\mathbf{V}(\nabla \mathbf{u}) - \mathbf{V}(\mathbf{Q})|^{2\alpha} \, dx \right)^{\frac{1}{\alpha}}$$

Proof. Apply Theorem 3.2 with **q** such that $\langle \mathbf{u} - \mathbf{q} \rangle_{2B} = 0$. Then use Theorem 2.5 with $\mathbf{w}(x) = \mathbf{u}(x) - \mathbf{Q}x$.

Using Gehring's Lemma we deduce the following assertion.

Corollary 3.4. There exists $s_0 > 1$ such that for all local minimizers $\mathbf{u} \in W^{1,\varphi}_{loc}(\Omega)$ of \mathcal{F} , all balls B with $2B \subset \subset \Omega$ and all $\mathbf{Q} \in \mathbb{R}^{N \times n}$

$$\left(\oint_{B} |\mathbf{V}(\nabla \mathbf{u}) - \mathbf{V}(\mathbf{Q})|^{2s_0} dx\right)^{\frac{1}{s_0}} \le c \oint_{2B} |\mathbf{V}(\nabla \mathbf{u}) - \mathbf{V}(\mathbf{Q})|^2 dx.$$

4. The \mathcal{A} -harmonic approximation

In this section we present a generalization of the \mathcal{A} -harmonic approximation lemma in Orlicz spaces. Basically it says that if a function locally "almost" behaves like an \mathcal{A} -harmonic function, then it is close to an \mathcal{A} -harmonic function. The proof is based on the Lipschitz truncation technique, which goes back to Acerbi- Fusco [1] but has been refined by many others.

Orginally the closeness of the function to its \mathcal{A} -harmonic approximation was stated in terms of the L^2 -distance and later for the non-linear problems in terms of the L^p distance. Based on a refinement of the Lipschitz truncation technique [11], it has been shown in [13] that also the distance in terms of the gradients is small.

Let us consider the following elliptic system

$$-\partial_{\alpha}(\mathcal{A}_{ii}^{\alpha\beta}D_{\beta}u^{j}) = -\partial_{\alpha}H_{i}^{\alpha} \qquad \text{in } B,$$

where $\alpha, \beta = 1, \ldots, n$ and $i, j = 1, \ldots N$. We use the convention that repeated indices are summed. In short we write $-\operatorname{div}(\mathcal{A}\nabla \mathbf{u}) = -\operatorname{div} \mathbf{G}$. We assume that \mathcal{A} is constant. We say that \mathcal{A} is strongly elliptic in the sense of Legendre-Hadamard if for all $\mathbf{a} \in \mathbb{R}^N, \mathbf{b} \in \mathbb{R}^n$ holds

$$\mathcal{A}_{ij}^{\alpha\beta}a^i b_\alpha a^j b_\beta \ge \kappa_{\mathcal{A}} |\mathbf{a}|^2 |\mathbf{b}|^2$$

for some $\kappa_{\mathcal{A}} > 0$. The biggest possible constant $\kappa_{\mathcal{A}}$ is called the ellipticity constant of \mathcal{A} . By $|\mathcal{A}|$ we denote the Euclidean norm of \mathcal{A} . We say that a Sobolev function \mathbf{w} on a ball B is \mathcal{A} -harmonic, if it satisfies $-\operatorname{div}(\mathcal{A}\nabla\mathbf{w}) = 0$ in the sense of distributions.

Given a Sobolev function \mathbf{u} on a ball B we want to find an \mathcal{A} -harmonic function \mathbf{h} which is close the our function \mathbf{u} . The way to find \mathbf{h} is very simple: it will be the \mathcal{A} -harmonic function with the same boundary values as \mathbf{u} . In particular, we want to find a Sobolev function \mathbf{h} which satisfies

(4.24)
$$-\operatorname{div}(\mathcal{A}\nabla\mathbf{h}) = 0 \qquad \text{on } B$$
$$\mathbf{h} = \mathbf{u} \qquad \text{on } \partial B$$

in the sense of distributions.

Let $\mathbf{w} := \mathbf{h} - \mathbf{u}$, then (4.24) is equivalent to finding a Sobolev function \mathbf{w} which satisfies

(4.25)
$$-\operatorname{div}(\mathcal{A}\nabla\mathbf{w}) = -\operatorname{div}(\mathcal{A}\nabla\mathbf{u}) \quad \text{on } B$$
$$\mathbf{w} = \mathbf{0} \quad \text{on } \partial B$$

in the sense of distributions.

Our main approximation result is the following.

Theorem 4.1. Let $B \subset \Omega$ be a ball with radius r_B and let $\widetilde{B} \subset \Omega$ denote either Bor 2B. Let \mathcal{A} be strongly elliptic in the sense of Legendre-Hadamard. Let ψ be an Nfunction with $\Delta_2(\psi, \psi^*) < \infty$ and let s > 1. Then for every $\varepsilon > 0$, there exists $\delta > 0$ only depending on n, N, κ_A , $|\mathcal{A}|$, $\Delta_2(\psi, \psi^*)$ and s such that the following holds: let $\mathbf{u} \in W^{1,\psi}(\widetilde{B})$ be almost \mathcal{A} -harmonic on B in the sense that

(4.26)
$$\left| \oint_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla \boldsymbol{\xi} \, dx \right| \leq \delta \oint_{\widetilde{B}} |\nabla \mathbf{u}| \, dx \, \|\nabla \boldsymbol{\xi}\|_{L^{\infty}(B)}$$

for all $\boldsymbol{\xi} \in C_0^{\infty}(B)$. Then the unique solution $\mathbf{w} \in W_0^{1,\psi}(B)$ of (4.25) satisfies

$$(4.27) \quad \oint_{B} \psi\left(\frac{|\mathbf{w}|}{r_{B}}\right) dx + \oint_{B} \psi(|\nabla \mathbf{w}|) dx \le \varepsilon \left(\left(\oint_{B} \left(\psi(|\nabla \mathbf{u}|) \right)^{s} dx \right)^{\frac{1}{s}} + \oint_{\widetilde{B}} \psi(|\nabla \mathbf{u}|) dx \right).$$

The proof of this theorem can be found at the end of this section. The distinction between B and \tilde{B} on the right-hand side of (4.27) allows a finer tuning with respect to the exponents. If $B = \tilde{B}$, then only the term involving s is needed.

The following result on the solvability and uniqueness in the setting of classical Sobolev spaces $W_0^{1,q}(B,\mathbb{R}^N)$ can be found in [14, Lemma 2].

Lemma 4.2. Let $B \subset \subset \Omega$ be a ball, let \mathcal{A} be strongly elliptic in the sense of Legendre-Hadamard and let $1 < q < \infty$. Then for every $\mathbf{G} \in L^q(B, \mathbb{R}^{N \times n})$, there exists a unique weak solution $\mathbf{u} = T_{\mathcal{A}} \mathbf{G} \in W_0^{1,q}(B, \mathbb{R}^N)$ of

(4.28)
$$-\operatorname{div}(\mathcal{A}\nabla\mathbf{u}) = -\operatorname{div}\mathbf{G} \qquad on \ B,$$
$$\mathbf{u} = \mathbf{0} \qquad on \ \partial B$$

The solution operator $T_{\mathcal{A}}$ is linear and satisfies

$$\left\|\nabla T_{\mathcal{A}}\mathbf{G}\right\|_{L^{q}(B)} \le c \left\|\mathbf{G}\right\|_{L^{q}(B)}$$

where c only depends on n, N, $\kappa_{\mathcal{A}}$, $|\mathcal{A}|$ and q.

Remark 4.3. Note that our constants do not depend on the size of the ball, since the estimates involved are scaling invariant.

Let $T_{\mathcal{A}}$ be the solution operator of Lemma 4.2. Then by the uniqueness of Lemma 4.2, the operator $T_{\mathcal{A}} : L^q(B, \mathbb{R}^{N \times n}) \to W_0^{1,q}(B, \mathbb{R}^N)$ does not depend on the choice of $q \in (1, \infty)$. Therefore, $T_{\mathcal{A}}$ is uniquely defined from $\bigcup_{1 < q < \infty} L^q(B, \mathbb{R}^{N \times n})$ to $\bigcup_{1 < q < \infty} W_0^{1,q}(B, \mathbb{R}^N)$

We need to extend Lemma 4.2 to the setting of Orlicz spaces. We will do so by means of the following real interpolation theorem of Peetre [27, Theorem 5.1] which states, that whenever ψ is of the form (2.13), then L^{ψ} is an interpolation space between L^{p_0} and L^{p_1} . **Theorem 4.4.** Let ψ be an N-function with $\Delta_2(\psi, \psi^*)$ and p_0 , p_1 as in Lemma 2.1. Moreover let S be a linear, bounded operator from $L^{p_j} \to L^{p_j}$ for j = 0, 1. Then there exists K_2 , which only depends on $\Delta_2(\psi, \psi^*)$, and the operator norms of S such that

$$\|Sf\|_{\psi} \le K_2 \|f\|_{\psi}$$
$$\int \psi(|Sf|/K_2) \, d\mu \le \int \psi(|f|) \, d\mu$$

for every $f \in L^{\psi}$.

This interpolation result and Lemma 4.2 immediately imply:

Theorem 4.5. Let $B \subset \Omega$ be a ball, let \mathcal{A} be strongly elliptic in the sense of Legendre-Hadamard and ψ be an N-function with $\Delta_2(\psi, \psi^*)$. Then the solution operator $T_{\mathcal{A}}$ of Lemma 4.2 is continuous from $L^{\psi}(B, \mathbb{R}^{N \times n})$ to $W_0^{1,\psi}(B, \mathbb{R}^n)$ and

(4.29)
$$\|\nabla T_{\mathcal{A}} \mathbf{G}\|_{L^{\psi}(B)} \leq c \|\mathbf{G}\|_{L^{\psi}(B)},$$
$$\int_{B} \psi(|\nabla T_{\mathcal{A}} \mathbf{G}|) \, dx \leq c \int_{B} \psi(|\mathbf{G}|) \, dx,$$

for all $\mathbf{G} \in L^{\psi}(B, \mathbb{R}^{N \times n})$, where c only depends on n, N, $\kappa_{\mathcal{A}}$, $|\mathcal{A}|$, $\Delta_2(\psi, \psi^*)$.

Remark 4.6. Since ψ satisfies (2.12) for some $1 < p_0 < p_1 < \infty$ it follows easily that $L^{\psi}(B) \hookrightarrow L^{p_0}(B)$ for every ball $B \subset \Omega$. From this and the uniqueness in Lemma 4.2, the solution of (4.28) is also unique in $W_0^{1,\psi}(B, \mathbb{R}^N)$.

Since \mathcal{A} is only strongly elliptic in the sense of Legendre-Hadamard, we will not be able to apply the Lipschitz truncation technique directly to our almost \mathcal{A} -harmonic function **u**. Instead, we need to use duality and apply the Lipschitz truncation technique to the test functions. For this reason, we prove the following variational inequality.

Lemma 4.7. Let $B \subset \Omega$ be a ball and let \mathcal{A} be strongly elliptic in the sense of Legendre-Hadamard. Then it holds for all $\mathbf{u} \in W_0^{1,\psi}(B)$ that

(4.30a)
$$\|\nabla \mathbf{u}\|_{\psi} \sim \sup_{\substack{\boldsymbol{\xi} \in C_0^{\infty}(B) \\ \|\nabla \boldsymbol{\xi}\|_{\psi^*} \le 1^B}} \int_B \mathcal{A} \nabla \mathbf{u} \cdot \nabla \boldsymbol{\xi} \, dx,$$

(4.30b)
$$\int_{B} \psi(|\nabla \mathbf{u}|) \, dx \sim \sup_{\boldsymbol{\xi} \in C_{0}^{\infty}(B)} \left[\int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla \boldsymbol{\xi} \, dx - \int_{B} \psi^{*}(|\nabla \boldsymbol{\xi}|) \, dx \right].$$

The implicit constants only depend on $n, N, \kappa_{\mathcal{A}}, |\mathcal{A}|, \Delta_2(\psi, \psi^*)$.

Proof. We begin with the proof of (4.30a). The \gtrsim estimate is a simple consequence of Hölder's inequality, so let us concentrate on \lesssim . Since $(L^{\psi})^* \cong L^{(\psi^*)}$ (with constants bounded by 2) and $C_0^{\infty}(B)$ is dense in $L^{(\psi^*)}(\Omega)$, we have

$$\|\nabla \mathbf{u}\|_{\psi} \leq 2 \sup_{\substack{\mathbf{H} \in C_0^{\infty}(B, \mathbb{R}^{N \times n}) \\ \|\mathbf{H}\|_{\psi^*} \leq 1}} \int_B \nabla \mathbf{u} \cdot \mathbf{H} \, dx,$$

Define $\overline{\mathcal{A}}$ by $\overline{\mathcal{A}}_{ij}^{\alpha\beta} := \mathcal{A}_{ji}^{\beta\alpha}$, then $-\operatorname{div}(\overline{\mathcal{A}}\nabla\mathbf{u})$ is the formal adjoint operator of $-\operatorname{div}(\mathcal{A}\nabla\mathbf{u})$. In particular, using (4.28)

(4.31)
$$\int_{B} \nabla \mathbf{u} \cdot \mathbf{H} = \int_{B} \nabla \mathbf{u} \cdot \overline{\mathcal{A}} \nabla T_{\overline{\mathcal{A}}} \mathbf{H} \, dx$$
$$= \int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla T_{\overline{\mathcal{A}}} \mathbf{H} \, dx.$$

Hence,

$$\begin{aligned} \|\nabla \mathbf{u}\|_{\psi} &\leq 2 \sup_{\substack{\mathbf{H} \in C_{0}^{\infty}(B, \mathbb{R}^{N \times n}) \\ \|\mathbf{H}\|_{\psi^{*}} \leq 1}} \int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla T_{\overline{\mathcal{A}}} \mathbf{H} \, dx \\ &\leq 4 \sup_{\substack{\mathbf{H} \in C_{0}^{\infty}(B, \mathbb{R}^{N \times n}) \\ \|\mathbf{H}\|_{\psi^{*}} \leq 1}} \|\mathcal{A} \nabla \mathbf{u}\|_{L^{\psi}(B)} \|\nabla T_{\overline{\mathcal{A}}} \mathbf{H}\|_{\psi^{*}}. \end{aligned}$$

where we used in the last step Theorem 4.5 for $T_{\overline{\mathcal{A}}}$ and ψ^* . This proves (4.30a). Let us now prove (4.30b). The estimate \gtrsim just follows from

$$\int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla \boldsymbol{\xi} \, dx - \int_{B} \psi^{*}(|\nabla \boldsymbol{\xi}|) \, dx \leq \int_{B} \psi(|\mathcal{A}||\nabla \mathbf{u}|) \, dx$$
$$\leq c(|\mathcal{A}|) \int_{B} \psi(|\nabla \mathbf{u}|) \, dx,$$

where we used $|\mathcal{A}\nabla \mathbf{u} \cdot \nabla \boldsymbol{\xi}| \leq |\mathcal{A}| |\nabla \mathbf{u}| |\nabla \boldsymbol{\xi}|$, Young's inequality and $\psi \in \Delta_2$. We turn to \leq of (4.30b). Recall that

$$\psi^{**}(t) = \psi(t) = \sup_{u \ge 0} (ut - \psi^*(u)),$$

where the supremum is attained at $u = \psi'(t)$. Thus the choice $\mathbf{H} := \psi'(|\nabla \mathbf{u}|) \frac{\nabla \mathbf{u}}{|\nabla \mathbf{u}|}$ (with $\mathbf{H} = \mathbf{0}$ where $\nabla \mathbf{u} = \mathbf{0}$) implies

$$\int_{B} \psi(|\nabla \mathbf{u}|) \, dx \le \sup_{\mathbf{H} \in (L^{\psi^*}(B, \mathbb{R}^{N \times n}))} \left[\int_{B} \nabla \mathbf{u} \cdot \mathbf{H} \, dx - \int_{B} \psi^*(|\mathbf{H}|) \, dx \right].$$

Using $T_{\overline{\mathcal{A}}}$ we estimate with (4.31)

$$\int_{B} \psi(|\nabla \mathbf{u}|) \, dx \leq \sup_{\mathbf{H} \in L^{\psi^*}(B, \mathbb{R}^{N \times n})} \left[\int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla T_{\overline{\mathcal{A}}} \mathbf{H} \, dx - \int_{B} \psi^*(|\mathbf{H}|) \, dx \right].$$

By Theorem 4.5 there exists $c \ge 1$ such that

$$\int_{B} \psi^*(|\nabla T_{\mathcal{A}}\mathbf{H}|) \, dx \le c \, \int_{B} \psi^*(|\mathbf{H}|) \, dx.$$

This proves the following:

$$\int_{B} \psi(|\nabla \mathbf{u}|) \, dx \leq \sup_{\mathbf{H} \in (L^{\psi^*}(B, \mathbb{R}^{N \times n})} \left[\int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla T_{\overline{\mathcal{A}}} \mathbf{H} \, dx - c \int_{B} \psi^*(|\nabla T_{\overline{\mathcal{A}}} \mathbf{H}|) \, dx \right]$$
$$\leq \sup_{\boldsymbol{\xi} \in L^{\psi^*}(B, \mathbb{R}^N)} \left[\int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla \boldsymbol{\xi} \, dx - c \int_{B} \psi^*(|\nabla \boldsymbol{\xi}|) \, dx \right].$$

We replace \mathbf{u} by $c\mathbf{u}$ to get

$$\int_{B} \psi \left(c |\nabla \mathbf{u}| \right) dx \le c \sup_{\boldsymbol{\xi} \in L^{\psi^*}(B, \mathbb{R}^N)} \left[\int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla \boldsymbol{\xi} \, dx - \int_{B} \psi^*(|\nabla \boldsymbol{\xi}|) \, dx \right].$$

Now the claim follows using $\psi \in \Delta_2$ on the left-hand side and the density of $C_0^{\infty}(B, \mathbb{R}^N)$ in $L^{\psi^*}(B, \mathbb{R}^N)$ (using $\psi^* \in \Delta_2$).

Moreover, we need the following result of [13, Theorem 3.3] about *Lipschitz truncations* in Orlicz spaces.

Theorem 4.8 (Lipschitz truncation). Let $B \subset \Omega$ be a ball and let ψ be an N-function with $\Delta_2(\psi, \psi^*) < \infty$. If $\mathbf{w} \in W_0^{1,\psi}(B, \mathbb{R}^N)$, then for every $m_0 \in \mathbb{N}$ and $\gamma > 0$ there exists $\lambda \in [\gamma, 2^{m_0}\gamma]$ and $\mathbf{w}_{\lambda} \in W_0^{1,\infty}(B, \mathbb{R}^N)$ (called the Lipschitz truncation) such that

$$\|\nabla \mathbf{w}_{\lambda}\|_{\infty} \leq c \,\lambda,$$

$$\oint_{B} \psi \left(|\nabla \mathbf{w}_{\lambda}| \chi_{\{\mathbf{w}_{\lambda} \neq \mathbf{w}\}} \right) dx \leq c \,\psi(\lambda) \frac{|\{\mathbf{w}_{\lambda} \neq \mathbf{w}\}|}{|B|} \leq \frac{c}{m_{0}} \oint_{B} \psi(|\nabla \mathbf{w}|) \,dx$$

$$\oint_{B} \psi \left(|\nabla \mathbf{w}_{\lambda}| \right) dx \leq c \oint_{B} \psi \left(|\nabla \mathbf{w}| \right) dx.$$

The constant c depends only on $\Delta_2(\psi, \psi^*)$, n and N.

We are ready to prove Theorem 4.1.

Proof of Theorem 4.1. We begin with an application of Lemma 4.7:

(4.32)
$$\oint_{B} \psi(|\nabla \mathbf{u}|) \, dx \le c \sup_{\boldsymbol{\xi} \in C_{0}^{\infty}(B,\mathbb{R}^{N})} \left[\oint_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla \boldsymbol{\xi} \, dx - \oint_{B} \psi^{*}(|\nabla \boldsymbol{\xi}|) \, dx \right].$$

In the following let us fix $\boldsymbol{\xi} \in C_0^{\infty}(B)$. Choose $\gamma \geq 0$ such that

(4.33)
$$\psi^*(\gamma) = \oint_B \psi^*(|\nabla \boldsymbol{\xi}|) \, dx.$$

and let $m_0 \in \mathbb{N}$. Due to Theorem 4.8 applied to ψ^* we find $\lambda \in [\gamma, 2^{m_0}\gamma]$ and $\boldsymbol{\xi}_{\lambda} \in W_0^{1,\infty}(B)$ such that

(4.34)
$$\|\nabla \boldsymbol{\xi}_{\lambda}\|_{\infty} \le c\,\lambda,$$

(4.35)
$$\psi^*(\lambda) \frac{|\{\boldsymbol{\xi}_\lambda \neq \boldsymbol{\xi}\}|}{|B|} \le \frac{c}{m_0} \int_B \psi^*(|\nabla \boldsymbol{\xi}|) \, dx$$

(4.36)
$$\oint_{B} \psi^{*} (|\nabla \boldsymbol{\xi}_{\lambda}|) \, dx \leq c \oint_{B} \psi^{*} (|\nabla \boldsymbol{\xi}|) \, dx.$$

Let us point out that the use of the Lipschitz truncation is not a problem of the regularity of $\boldsymbol{\xi}$ as it is C_0^{∞} . It is the precise estimates above that we need.

We calculate

$$\int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla \boldsymbol{\xi} \, dx = \int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla \boldsymbol{\xi}_{\lambda} \, dx + \int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla (\boldsymbol{\xi} - \boldsymbol{\xi}_{\lambda}) \, dx =: I + II$$

Using Young's inequality and (4.36) we estimate

$$II = \int_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla (\boldsymbol{\xi} - \boldsymbol{\xi}_{\lambda}) \chi_{\{\boldsymbol{\xi} \neq \boldsymbol{\xi}_{\lambda}\}} dx$$

$$\leq c \int_{B} \psi(|\nabla \mathbf{u}| \chi_{\{\boldsymbol{\xi} \neq \boldsymbol{\xi}_{\lambda}\}}) dx + \frac{1}{2} \int_{B} \psi^{*}(|\nabla \boldsymbol{\xi}|) dx =: II_{1} + II_{2},$$

where c depends on $|\mathcal{A}|$, $\Delta_2(\psi, \psi^*)$. With Hölder's inequality we get

$$II_1 \le c \left(\int_B \left(\psi(|\nabla \mathbf{u})| \right)^s dx \right)^{\frac{1}{s}} \left(\frac{|\{\boldsymbol{\xi}_\lambda \neq \boldsymbol{\xi}\}|}{|B|} \right)^{1-\frac{1}{s}}.$$

If follows from (4.35), (4.33) and $\lambda \geq \gamma$ that

$$\frac{|\{\boldsymbol{\xi}_{\lambda} \neq \boldsymbol{\xi}\}|}{|B|} \le \frac{c\psi^{*}(\gamma)}{m_{0}\psi^{*}(\lambda)} \le \frac{c}{m_{0}}$$

Thus

$$II_1 \le c \left(\oint_B \left(\psi(|\nabla \mathbf{u})| \right)^s dx \right)^{\frac{1}{s}} \left(\frac{c}{m_0} \right)^{1-\frac{1}{s}}.$$

We choose m_0 so large such that

$$II_1 \leq \frac{\varepsilon}{2} \left(\oint_B \left(\psi(|\nabla \mathbf{u})|) \right)^s dx \right)^{\frac{1}{s}}.$$

Since **u** is almost \mathcal{A} -harmonic and $\|\nabla \boldsymbol{\xi}_{\lambda}\|_{\infty} \leq c \, \lambda \leq c \, 2^{m_0} \gamma$ we have

$$|I| \leq \delta \oint_{\widetilde{B}} |\nabla \mathbf{u}| \, dx \, \|\nabla \boldsymbol{\xi}_{\lambda}\|_{\infty} \leq \delta \oint_{\widetilde{B}} |\nabla \mathbf{u}| \, dx \, c \, 2^{m_0} \gamma.$$

We apply Young's inequality and (4.33) to get

$$|I| \leq \delta 2^{m_0} c \left(\oint_{\widetilde{B}} \psi(|\nabla \mathbf{u}|) \, dx + \psi^*(\gamma) \right)$$

$$\leq \delta 2^{m_0} c \oint_{\widetilde{B}} \psi(|\nabla \mathbf{u}|) \, dx + \delta 2^{m_0} c \oint_{B} \psi^*(|\nabla \boldsymbol{\xi}|) \, dx.$$

Now, we choose $\delta > 0$ so small such that $\delta 2^{m_0} c \leq \varepsilon/2$. Thus

$$|I| \leq \frac{\varepsilon}{2} \oint_{\widetilde{B}} \psi(|\nabla \mathbf{u}|) \, dx + \frac{1}{2} \oint_{B} \psi^*(|\nabla \boldsymbol{\xi}|) \, dx.$$

Combining the estimates for I, II and II_1 we get

$$\oint_{B} \mathcal{A} \nabla \mathbf{u} \cdot \nabla \boldsymbol{\xi} \, dx \le \varepsilon \left(\left(\oint_{B} \left(\psi(|\nabla \mathbf{u}|) \right)^{s} \, dx \right)^{\frac{1}{s}} + \oint_{\widetilde{B}} \psi(|\nabla \mathbf{u}|) \, dx \right) + \oint_{B} \psi^{*}(|\nabla \boldsymbol{\xi}|) \, dx.$$

Now taking the supremum over all $\boldsymbol{\xi} \in C_0^{\infty}(B)$ and using (4.32) we get

$$\oint_{B} \psi(|\nabla \mathbf{w}|) \, dx \le \varepsilon \left(\left(\oint_{B} \left(\psi(|\nabla \mathbf{u}|) \right)^{s} \, dx \right)^{\frac{1}{s}} + \oint_{\widetilde{B}} \psi(|\nabla \mathbf{u}|) \, dx \right).$$

The claim follows by Poincaré inequality, see Theorem 2.5.

5. Almost \mathcal{A} -harmonicity

The following result is a special case of [9, Lemma A.2].

Lemma 5.1. Let $B \subset \mathbb{R}^n$ be a ball and $\mathbf{w} \in W^{1,\varphi}(B)$. Then

$$\int_{B} |\mathbf{V}(\nabla \mathbf{w}) - \langle \mathbf{V}(\nabla \mathbf{w}) \rangle_{B}|^{2} dx \sim \int_{B} |\mathbf{V}(\nabla \mathbf{w}) - \mathbf{V}(\langle \nabla \mathbf{w} \rangle_{B})|^{2} dx.$$

The constants are independent of B and w; they only depend on the characteristics of φ .

Lemma 5.2. There exists $\delta > 0$, which only depends on the characteristics of φ , such that for every ball B with $B \subset \subset \Omega$ and every $\mathbf{u} \in W^{1,\varphi}(B)$ the estimate

(5.37)
$$\oint_{B} |\mathbf{V}(\nabla \mathbf{u}) - \langle \mathbf{V}(\nabla \mathbf{u}) \rangle_{B}|^{2} dx \leq \delta \oint_{B} |\mathbf{V}(\nabla \mathbf{u})|^{2} dx$$

implies

(5.38)
$$\oint_{B} |\mathbf{V}(\nabla \mathbf{u})|^2 \, dx \le 4 \, |\mathbf{V}(\langle \nabla \mathbf{u} \rangle_B)|^2,$$

(5.39)
$$\oint_{B} |\mathbf{V}(\nabla \mathbf{u}) - \langle \mathbf{V}(\nabla \mathbf{u}) \rangle_{B}|^{2} dx \leq 4 \,\delta \, |\mathbf{V}(\langle \nabla \mathbf{u} \rangle_{B})|^{2}.$$

Proof. It follows from (5.37) and Lemma 5.1 that

$$\begin{split} \oint_{B} |\mathbf{V}(\nabla \mathbf{u})|^{2} \, dx &\leq 2 \oint_{B} |\mathbf{V}(\nabla \mathbf{u}) - \mathbf{V}(\langle \nabla \mathbf{u} \rangle_{B})|^{2} \, dx + 2 |\mathbf{V}(\langle \nabla \mathbf{u} \rangle_{B})|^{2} \\ &\leq c \oint_{B} |\mathbf{V}(\nabla \mathbf{u}) - \langle \mathbf{V}(\nabla \mathbf{u}) \rangle_{B}|^{2} \, dx + 2 |\mathbf{V}(\langle \nabla \mathbf{u} \rangle_{B})|^{2} \\ &\leq \delta \, c \oint_{B} |\mathbf{V}(\nabla \mathbf{u})|^{2} \, dx + 2 |\mathbf{V}(\langle \nabla \mathbf{u} \rangle_{B})|^{2}. \end{split}$$

For small δ we absorb the first term of the right-hand side to get (5.38). The remaining estimate (5.39) is a combination of (5.37) and (5.38).

Lemma 5.3. Let **u** be a local minimizer of \mathcal{F} . Then for every ball B with $2B \subset \subset \Omega$ and every $\mathbf{Q} \in \mathbb{R}^{N \times n}$ it holds

$$\int_{B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) \, dx \le c \, \varphi_{|\mathbf{Q}|} \left(\int_{2B} |\nabla \mathbf{u} - \mathbf{Q}| \, dx \right).$$

Proof. From Corollary 3.3 we get

$$\oint_{B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) \, dx \le c \left(\oint_{2B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|)^{\alpha} \, dx \right)^{\frac{1}{\alpha}}.$$

We can apply then Corollary 3.4 in [9] to conclude.

Lemma 5.4. For all $\varepsilon > 0$ there exists $\delta > 0$, which only depends on ε and the characteristics of φ , such that for every ball B with $B \subset \subset \Omega$ and every $\mathbf{u} \in W^{1,\varphi}(B)$

(5.40)
$$\oint_{B} |\mathbf{V}(\nabla \mathbf{u}) - \langle \mathbf{V}(\nabla \mathbf{u}) \rangle_{B}|^{2} dx \leq \delta \oint_{B} |\mathbf{V}(\nabla \mathbf{u})|^{2} dx$$

implies

(5.41)
$$\int_{B} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{B}| \, dx \le \varepsilon \, |\langle \nabla \mathbf{u} \rangle_{B}|.$$

Proof. Let $\mathbf{Q} = \langle \nabla \mathbf{u} \rangle_B$. Then, by Jensen's inequality and Lemma 5.2 we get

$$\begin{split} \varphi_{|\langle \nabla \mathbf{u} \rangle_B|} \bigg(\oint_B |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_B | \, dx \bigg) &\leq \oint_B \varphi_{|\langle \nabla \mathbf{u} \rangle_B|} (|\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_B|) \, dx \\ &\leq c \oint_B |\mathbf{V} (\nabla \mathbf{u}) - \mathbf{V} (\langle \nabla \mathbf{u} \rangle_B)|^2 \, dx \\ &\leq \delta \, c \, |\mathbf{V} (\langle \nabla \mathbf{u} \rangle_B)|^2 \\ &\leq \delta \, c \, \varphi (|\langle \nabla \mathbf{u} \rangle_B|) \\ &\leq \delta \, c \, \varphi_{|\langle \nabla \mathbf{u} \rangle_B|} (|\langle \nabla \mathbf{u} \rangle_B|). \end{split}$$

For the last inequality we used the fact that $\varphi(a) \sim \varphi_a(a)$ for $a \geq 0$. Using the Δ_2 condition of $\varphi_{|\langle \nabla \mathbf{u} \rangle_B|}$ it follows that for every $\varepsilon > 0$ there exists a $\delta > 0$ such that

$$\oint_{B} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{B}| \, dx \le \varepsilon \, |\langle \nabla \mathbf{u} \rangle_{B}|.$$

Note that the smallness assumption in (5.40) automatically implies that $\langle \nabla \mathbf{u} \rangle_B \neq 0$ (unless $\nabla \mathbf{u} = 0$ on *B*). So the smallness assumption ensures that in some sense in the non-degenerate situation.

Lemma 5.5. For all $\varepsilon > 0$ there exists $\delta > 0$ such that for every local minimizer $\mathbf{u} \in W^{1,\varphi}_{loc}(\Omega)$ of \mathcal{F} and every ball B with $2B \subset \subset \Omega$ and

(5.42)
$$\int_{2B} |\mathbf{V}(\nabla \mathbf{u}) - \langle \mathbf{V}(\nabla \mathbf{u}) \rangle_{2B}|^2 \, dx \le \delta \int_{2B} |\mathbf{V}(\nabla \mathbf{u})|^2 \, dx$$

there holds

(5.43)
$$\left| \oint_{B} D^{2} f(\mathbf{Q}) (\nabla \mathbf{u} - \mathbf{Q}, \nabla \boldsymbol{\xi} \, dx \right| \leq \varepsilon \, \varphi''(|\mathbf{Q}|) \oint_{2B} |\nabla \mathbf{u} - \mathbf{Q}| \, dx \|\nabla \boldsymbol{\xi}\|_{\infty}.$$

for every $\boldsymbol{\xi} \in C_0^{\infty}(B)$, where $\mathbf{Q} := \langle \nabla \mathbf{u} \rangle_{2B}$. In particular, \mathbf{u} is almost \mathcal{A} -harmonic (in the sense of Theorem 4.1), with $\mathcal{A} = D^2 f(\mathbf{Q}) / \varphi''(|\mathbf{Q}|)$.

Proof. Let $\varepsilon > 0$. Without loss of generality we can assume that $\delta > 0$ is so small that the Lemmas 5.2 and 5.4 give

(5.44)
$$\oint_{2B} |\mathbf{V}(\nabla \mathbf{u})|^2 \, dx \le 4 \, |\mathbf{V}(\mathbf{Q})|^2,$$

(5.45)
$$\oint_{2B} |\nabla \mathbf{u} - \mathbf{Q}| \, dx \le \varepsilon \, |\mathbf{Q}|$$

From the last inequality we deduce

(5.46)
$$\varphi''(|\mathbf{Q}|) \left(\oint_{2B} |\nabla \mathbf{u} - \mathbf{Q}| \, dx \right)^2 \sim \varphi_{|\mathbf{Q}|} \left(\oint_{2B} |\nabla \mathbf{u} - \mathbf{Q}| \, dx \right).$$

Since the estimate (5.43) is homogeneous with respect to $\|\nabla \boldsymbol{\xi}\|_{\infty}$, it suffices to show that (5.43) holds for all $\boldsymbol{\xi} \in C_0^{\infty}(B)$ with $\|\nabla \boldsymbol{\xi}\|_{\infty} = \int_{2B} |\nabla \mathbf{u} - \mathbf{Q}| dx$. Hence, because of (5.46) it suffices to prove

(5.47)
$$\left| \oint_{B} D^{2} f(\mathbf{Q}) (\nabla \mathbf{u} - \mathbf{Q}, \nabla \boldsymbol{\xi}) \, dx \right| \leq \varepsilon \, c \, \varphi_{|\mathbf{Q}|} \left(\oint_{2B} |\nabla \mathbf{u} - \mathbf{Q}| \, dx \right)$$

for all such $\boldsymbol{\xi}$. We define

$$B^{\geq} := \left\{ x \in B : |\nabla \mathbf{u} - \mathbf{Q}| \ge \frac{1}{2} |\mathbf{Q}| \right\},\$$

$$B^{<} := \left\{ x \in B : |\nabla \mathbf{u} - \mathbf{Q}| < \frac{1}{2} |\mathbf{Q}| \right\}.$$

From the Euler-Lagrange equation we get $\int_B (Df(\nabla \mathbf{v}) - Df(\mathbf{Q})) : \nabla \boldsymbol{\xi} \, dx = 0$, and therefore

$$\begin{split} & \oint_B D^2 f(\mathbf{Q}) (\nabla \mathbf{u} - \mathbf{Q}, \nabla \boldsymbol{\xi}) \, dx \\ & = \oint_B \int_0^1 \left(D^2 f(\mathbf{Q}) - D^2 f(\mathbf{Q} + \theta(\nabla \mathbf{u} - \mathbf{Q})) \right) (\nabla \mathbf{u} - \mathbf{Q}, \nabla \boldsymbol{\xi}) \, d\theta \, dx. \end{split}$$

We split the right-hand side into the integral I over B^{\geq} and the integral II over $B^{<}$. Using **(H4)** we get

$$\begin{split} |I| &\leq c \oint_{B} \chi_{B^{\geq}} \int_{0}^{1} \left(\varphi''(|\mathbf{Q}|) + \varphi''(|\mathbf{Q} + \theta(\nabla \mathbf{u} - \mathbf{Q})|) \right) d\theta \ |\nabla \mathbf{u} - \mathbf{Q}| |\nabla \boldsymbol{\xi}| \, dx \\ &\leq c \oint_{B} \chi_{B^{\geq}} \left(\varphi''(|\mathbf{Q}|) + \varphi''(|\mathbf{Q}| + |\nabla \mathbf{u} - \mathbf{Q}|) \right) |\nabla \mathbf{u} - \mathbf{Q}| |\nabla \boldsymbol{\xi}| \, dx \\ &\leq c \oint_{B} \chi_{B^{\geq}} \left(|\nabla \mathbf{u} - \mathbf{Q}| \varphi'(|\mathbf{Q}|) + \varphi'_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) |\mathbf{Q}| \right) dx \ \frac{\|\nabla \boldsymbol{\xi}\|_{\infty}}{|\mathbf{Q}|} \\ &\leq \varepsilon c \oint_{B} \chi_{B^{\geq}} \left(|\nabla \mathbf{u} - \mathbf{Q}| \varphi'(|\mathbf{Q}|) + \varphi'_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) |\mathbf{Q}| \right) dx. \end{split}$$

We used Lemma 2.4 for the second, Assumption 2.2 for the third and (5.45) for the last estimate. Now, using $|\mathbf{Q}| \leq 2 |\nabla \mathbf{u} - \mathbf{Q}|$ on B^{\geq} and $\varphi_a(t) \sim \varphi(t)$ for $0 \leq a \leq t$ we get

$$\begin{split} |I| &\leq \varepsilon \, c \, \oint_{B} \chi_{B^{\geq}} \big(\varphi(|\nabla \mathbf{u} - \mathbf{Q}|) + \varphi_{|\mathbf{Q}|} (|\nabla \mathbf{u} - \mathbf{Q}|) \big) \, dx \\ &\leq \varepsilon \, c \, \oint_{B} \varphi_{|\mathbf{Q}|} (|\nabla \mathbf{u} - \mathbf{Q}|) \, dx. \end{split}$$

Let us estimate the modulus of II. Using (H5) and $|\nabla \mathbf{u} - \mathbf{Q}| < \frac{1}{2} |\mathbf{Q}|$ on $B^{<}$ we get

$$|II| \le c \oint_{B} \chi_{B} < \varphi''(|\mathbf{Q}|) |\mathbf{Q}|^{-\beta_1} |\nabla \mathbf{u} - \mathbf{Q}|^{1+\beta_1} |\nabla \boldsymbol{\xi}| \, dx,$$

where $\beta_1 := \min \{s_0, \beta\}$ with the constant s_0 from Corollary 3.4. Using Young's inequality we get

$$\begin{split} |II| &\leq \gamma \varphi''(|\mathbf{Q}|) \|\nabla \boldsymbol{\xi}\|_{\infty}^{2} + c_{\gamma} \int_{B} \chi_{B} \langle \varphi''(|\mathbf{Q}|) |\mathbf{Q}|^{-2\beta_{1}} |\nabla \mathbf{u} - \mathbf{Q}|^{2(1+\beta_{1})} dx \\ &\leq \gamma c \, \varphi_{|\mathbf{Q}|}(\|\nabla \boldsymbol{\xi}\|_{\infty}) + c_{\gamma}(\varphi(|\mathbf{Q}|))^{-\beta_{1}} \int_{B} \chi_{B} \langle \varphi''(|\mathbf{Q}|) |\nabla \mathbf{u} - \mathbf{Q}|^{2} \rangle^{1+\beta_{1}} dx \\ &\leq \gamma c \, \int_{2B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) \, dx + c_{\gamma}(\varphi(|\mathbf{Q}|))^{-\beta_{1}} \int_{B} \chi_{B} \langle \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) \rangle^{1+\beta_{1}} dx \\ &\leq \gamma c \, \int_{2B} |\mathbf{V}(\nabla \mathbf{u}) - \mathbf{V}(\mathbf{Q})|^{2} \, dx + c_{\gamma}(\varphi(|\mathbf{Q}|))^{-\beta_{1}} \int_{B} |\mathbf{V}(\nabla \mathbf{u}) - \mathbf{V}(\mathbf{Q})|^{2(1+\beta_{1})} \, dx. \end{split}$$

Here we used (5.46) for the second and Jensen's inequality, $\varphi''(a)t^2 \sim \varphi_a(t)$ for $0 \leq t \leq a$ and $|\nabla \mathbf{u} - \mathbf{Q}| < \frac{1}{2}|\mathbf{Q}|$ on $B^<$ for the third estimate. With the help of Corollary 3.4 we get

$$|II| \leq \gamma c \oint_{2B} |\mathbf{V}(\nabla \mathbf{u}) - \mathbf{V}(\mathbf{Q})|^2 dx + c_{\gamma}(\varphi(|\mathbf{Q}|))^{-\beta_1} \left(\oint_{2B} |\mathbf{V}(\nabla \mathbf{u}) - \mathbf{V}(\mathbf{Q})|^2 dx \right)^{1+\beta_1}.$$

L. Diening, D. Lengeler, B. Stroffolini, A. Verde

Using the assumption (5.42), Lemma 5.1 and (5.44) it follows that

$$|II| \leq \gamma c \oint_{2B} |\mathbf{V}(\nabla \mathbf{u}) - \mathbf{V}(\mathbf{Q})|^2 dx + c_{\gamma} \delta^{\beta_1} \oint_{2B} |\mathbf{V}(\nabla \mathbf{u}) - \mathbf{V}(\mathbf{Q})|^2 dx.$$

Choosing $\gamma > 0$ and then $\delta > 0$ small enough we get the assertion.

6. Excess decay estimate

In this section we will focus on the excess decay estimate. Therefore, we compare the almost harmonic solution with its harmonic approximation.

Proposition 6.1. For all $\varepsilon > 0$, there exists $\delta = \delta(\varphi, \varepsilon) > 0$ such that the following is true: if for some ball B with $2B \subset \subset \Omega$ the smallness assumption (5.42) holds true, then for every $\tau \in (0, 1]$

(6.48)
$$\Phi(\tau B, \mathbf{u}) \le c \tau^2 (1 + \varepsilon \tau^{-n-2}) \Phi(2B, \mathbf{u}),$$

where c depends only on the characteristics of φ and is independent of ε .

Proof. It suffices to consider the case $\tau \leq \frac{1}{2}$. Let s_0 be as in Corollary 3.4. Let \mathbf{q} be a linear function such that $\langle \mathbf{u} - \mathbf{q} \rangle_{2B} = 0$ and $\mathbf{Q} := \nabla \mathbf{q} = \langle \nabla \mathbf{u} \rangle_{2B}$. Define $\mathbf{z} := \mathbf{u} - \mathbf{q}$. Let \mathbf{h} be the harmonic approximation of \mathbf{z} with $\mathbf{h} = \mathbf{z}$ on ∂B . It follows from Lemma 5.5 that \mathbf{z} is almost \mathcal{A} -harmonic with $\mathcal{A} = D^2 f(\mathbf{Q}) / \varphi''(|\mathbf{Q}|)$. Thus by Theorem 4.1 for suitable $\delta = \delta(\varphi, \varepsilon)$ and by Theorem 4.1 the \mathcal{A} -harmonic approximation \mathbf{h} satisfies

$$\oint_{B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{z} - \nabla \mathbf{h}|) \, dx \le \varepsilon \left(\left(\oint_{B} \varphi_{|\mathbf{Q}|}^{s_{0}}(|\nabla \mathbf{u} - \mathbf{Q}|) \, dx \right)^{\frac{1}{s_{0}}} + \oint_{2B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{u} - \mathbf{Q}|) \, dx \right).$$

Now, it follows by Corollary 3.4 that

(6.49)
$$\int_{B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{z} - \nabla \mathbf{h}|) \, dx \le c \varepsilon \, \Phi(2B, \mathbf{u}).$$

Since $\nabla \mathbf{z} = \nabla \mathbf{u} - \mathbf{Q}$ and $\langle \nabla \mathbf{z} \rangle_{\tau B} = \langle \nabla \mathbf{u} \rangle_{\tau B} - \mathbf{Q}$, we get

$$\begin{split} \varPhi(\tau B, \mathbf{u}) &\leq c \oint_{\tau B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{z} - \langle \nabla \mathbf{z} \rangle_{\tau B}|) \, dx \\ &\leq c \oint_{\tau B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_{\tau B}|) \, dx + c \oint_{\tau B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{z} - \nabla \mathbf{h}|) \, dx \\ &=: I + II. \end{split}$$

For the second estimate we used Jensen's inequality. Using (6.49) we obtain

$$II \leq \tau^{-n} c \oint_{B} \varphi_{|\mathbf{Q}|}(|\nabla \mathbf{z} - \nabla \mathbf{h}|) \, dx \leq \tau^{-n} c \, \varepsilon \, \Phi(2B, \mathbf{u}).$$

By the interior regularity of the \mathcal{A} -harmonic function **h**, [21], and $\tau \leq \frac{1}{2}$ it holds that

$$\sup_{\tau B} |\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_{\tau B}| \le c \tau \oint_{B} |\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_{B}| \, dx.$$

This proves

$$I \le c \,\varphi_{|\mathbf{Q}|} \bigg(\tau \oint_B |\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_B | \, dx \bigg).$$

Using the estimate $\psi(st) \leq s\psi(t)$ for any $s \in [0, 1]$, $t \geq 0$ and any N-function ψ , we would get a factor τ in the estimate of I. However, to produce a factor τ^2 , we have to work differently and use the improved estimate $\varphi_a(st) \leq c s^2 \varphi_a(t)$ for all $s \in [0, 1]$, $a \geq 0$ and $t \in [0, a]$. We begin with

$$\begin{split} \oint_{B} |\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_{B} | \, dx &\leq \oint_{B} |\nabla \mathbf{z} - \langle \nabla \mathbf{z} \rangle_{B} | \, dx + 2 \oint_{B} |\nabla \mathbf{z} - \nabla \mathbf{h}| \, dx \\ &= \oint_{B} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{B} | \, dx + 2 \oint_{B} |\nabla \mathbf{z} - \nabla \mathbf{h}| \, dx, \end{split}$$

which implies

$$I \le c \,\varphi_{|\mathbf{Q}|} \left(\tau \oint_{B} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{B} | \, dx \right) + c \,\tau \varphi_{|\mathbf{Q}|} \left(\oint_{B} |\nabla \mathbf{z} - \nabla \mathbf{h}| \, dx \right).$$

Due to (5.45), we can use for the first term the improved estimate $\varphi_a(st) \leq c s^2 \varphi_a(t)$, which gives

$$I \leq c \tau^{2} \varphi_{|\mathbf{Q}|} \left(\oint_{B} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{B} | dx \right) + c \tau \varphi_{|\mathbf{Q}|} \left(\oint_{B} |\nabla \mathbf{z} - \nabla \mathbf{h}| dx \right)$$
$$\leq c \tau^{2} \oint_{B} \varphi_{|\mathbf{Q}|} \left(|\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{B} | \right) dx + c \tau \oint_{B} \varphi_{|\mathbf{Q}|} \left(|\nabla \mathbf{z} - \nabla \mathbf{h}| \right) dx.$$

Thus using (6.49) we get

$$I \le c \tau^2 \Phi(B, \mathbf{u}) + c \tau \varepsilon \Phi(2B, \mathbf{u}) \le c \left(\tau^2 + \varepsilon \tau\right) \Phi(2B, \mathbf{u}).$$

Combining the estimates for I and II we get the claim.

It follows now, by a series of standard arguments, that for any $\beta \in (0, 1)$, there exists a suitable small δ that ensures local $C^{0,\beta}$ -regularity of $\mathbf{V}(\nabla u)$, which implies Hölder continuity of the gradients as well.

Proposition 6.2 (Decay estimate). For $0 < \beta < 1$ there exists $\delta = \delta(\varphi, \beta) > 0$ such that the following is true. If for some ball $B \subset \Omega$ the smallness assumption (5.42) holds true, then

(6.50)
$$\Phi(\rho B, \mathbf{u}) \le c \, \rho^{2\beta} \Phi(2B, \mathbf{u})$$

for any $\rho \in (0,1]$, where $c = c(\varphi)$ depends only on the characteristics of φ .

Proof. Due to our assumption, we can apply Proposition 6.1 for any τ . Let $\gamma(\varepsilon, \tau) := c \tau^2 (1 + \varepsilon \tau^{-n-2})$ as in (6.48). Let us fix $\tau > 0$ and $\varepsilon > 0$, such that $\gamma(\varepsilon, \tau) \leq \min \{(\tau/2)^{2\beta}, \frac{1}{4}\}$. Let $\delta = \delta(\varphi, \varepsilon)$ chosen accordingly to Proposition 6.1 and also so small that $(1 + \tau^{-n/2})\delta^{1/2} \leq \frac{1}{2}$. By Proposition 6.1 we have

(6.51)
$$\Phi(\tau B, \mathbf{u}) \le \min\left\{(\tau/2)^{2\beta}, \frac{1}{4}\right\} \Phi(2B, \mathbf{u}).$$

We claim that the smallness assumption is inherited from 2B to τB , so that we can iterate (6.51). For this we estimate with the help of our smallness assumption

$$\begin{split} \left(\oint_{2B} |\mathbf{V}(\nabla \mathbf{u})|^2 \, dx \right)^{\frac{1}{2}} &\leq \left(\Phi(2B, \mathbf{u}) \right)^{\frac{1}{2}} + |\langle \mathbf{V}(\nabla \mathbf{v}) \rangle_{2B} - \langle \mathbf{V}(\nabla \mathbf{v}) \rangle_{\tau B}| + \left(\oint_{\tau B} |\mathbf{V}(\nabla \mathbf{v})|^2 \, dx \right)^{\frac{1}{2}} \\ &\leq \left(\Phi(2B, \mathbf{u}) \right)^{\frac{1}{2}} + \tau^{-n/2} \left(\Phi(2B, \mathbf{u}) \right)^{\frac{1}{2}} + \left(\oint_{\tau B} |\mathbf{V}(\nabla \mathbf{v})|^2 \, dx \right)^{\frac{1}{2}} \\ &\leq \left(1 + \tau^{-n/2} \right) \delta^{1/2} \left(\oint_{2B} |\mathbf{V}(\nabla \mathbf{u})|^2 \, dx \right)^{\frac{1}{2}} + \left(\oint_{\tau B} |\mathbf{V}(\nabla \mathbf{v})|^2 \, dx \right)^{\frac{1}{2}}. \end{split}$$

Using $(1 + \tau^{-n/2})\delta^{1/2} \le \frac{1}{2}$, we get

$$\oint_{2B} |\mathbf{V}(\nabla \mathbf{u})|^2 \, dx \le 4 \, \oint_{\tau B} |\mathbf{V}(\nabla \mathbf{v})|^2 \, dx.$$

Now (6.51) and the previous estimate imply

$$\Phi(\tau B, \mathbf{u}) \le \frac{1}{4} \Phi(2B, \mathbf{u}) \le \frac{1}{4} \delta \oint_{2B} |\mathbf{V}(\nabla \mathbf{u})|^2 \, dx \le \delta \oint_{\tau B} |\mathbf{V}(\nabla \mathbf{u})|^2 \, dx$$

In particular, the smallness assumption is also satisfied for τB . So by induction we get

(6.52)
$$\Phi((\tau/2)^k 2B, \mathbf{u}) \le \min\{(\tau/2)^{2\beta k}, 4^{-k}\} \Phi(2B, \mathbf{u}),$$

which is the desired claim.

Having the decay estimate, it is easy to proove our Main Theorem.

Proof of the Main Theorem 1.3. We can assume that (5.42) is satisfied with a strict inequality. By continuity, (5.42), holds for B = B(x) and all x in some neighborhood of x_0 . By Proposition 6.2 and Campanato's characterisation of Hölder continuity we deduce that $\mathbf{V}(\nabla \mathbf{u})$ is β -Hölder continuous in a neighbourhood of x_0 .

References

- E. Acerbi and N. Fusco. Semicontinuity problems in the calculus of variations. Arch. Rational Mech. Anal., 86(2):125–145, 1984.
- [2] William K. Allard. On the first variation of a varifold. Ann. of Math. (2), 95:417–491, 1972.
- [3] F. J. Almgren, Jr. Existence and regularity almost everywhere of solutions to elliptic variational problems among surfaces of varying topological type and singularity structure. Ann. of Math. (2), 87:321–391, 1968.
- [4] F. J. Almgren, Jr. Existence and regularity almost everywhere of solutions to elliptic variational problems with constraints. *Mem. Amer. Math. Soc.*, 4(165):viii+199, 1976.
- [5] D. Breit, Diening L., and Fuchs M. Solenoidal lipschitz truncation and applications in fluid mechanics. accepted, 2012.
- [6] Menita Carozza, Nicola Fusco, and Giuseppe Mingione. Partial regularity of minimizers of quasiconvex integrals with subquadratic growth. Ann. Mat. Pura Appl. (4), 175:141–164, 1998.
- [7] E. De Giorgi. Frontiere orientate di misura minima. Seminario di Matematica della Scuola Normale Superiore di Pisa, 1960-61. Editrice Tecnico Scientifica, Pisa, 1961.
- [8] L. Diening and F. Ettwein. Fractional estimates for non-differentiable elliptic systems with general growth. Forum Mathematicum, 20(3):523–556, 2008.
- [9] L. Diening, P. Kaplický, and S. Schwarzacher. BMO estimates for the p-Laplacian. Nonlinear Analysis: Theory, Methods and Applications, 2011.

- [10] L. Diening and Ch. Kreuzer. Linear convergence of an adaptive finite element method for the p-Laplacian equation. SIAM J. Numer. Anal., 46:614–638, 2008.
- [11] L. Diening, J. Málek, and M. Steinhauer. On Lipschitz truncations of Sobolev functions (with variable exponent) and their selected applications. ESAIM Control Optim. Calc. Var., 14(2):211–232, 2008.
- [12] L. Diening, B. Stroffolini, and A. Verde. Everywhere regularity of functionals with φ -growth. *Manuscripta Math.*, 129(4):449–481, 2009.
- [13] L. Diening, B. Stroffolini, and A. Verde. The φ -harmonic approximation and the regularity of φ -harmonic maps. *submitted*, 2010.
- [14] G. Dolzmann and S. Müller. Estimates for Green's matrices of elliptic systems by L^p theory. Manuscripta Math., 88(2):261–273, 1995.
- [15] F. Duzaar and J. F. Grotowski. Optimal interior partial regularity for nonlinear elliptic systems: the method of A-harmonic approximation. *Manuscripta Math.*, 103(3):267–298, 2000.
- [16] F. Duzaar, J. F. Grotowski, and M. Kronz. Regularity of almost minimizers of quasi-convex variational integrals with subquadratic growth. Ann. Mat. Pura Appl. (4), 184(4):421-448, 2005.
- [17] F. Duzaar and G. Mingione. The p-harmonic approximation and the regularity of p-harmonic maps. Calc. Var. Partial Differential Equations, 20(3):235–256, 2004.
- [18] F. Duzaar and G. Mingione. Harmonic type approximation lemmas. J. Math. Anal. Appl., 352(1):301–335, 2009.
- [19] Lawrence C. Evans. Quasiconvexity and partial regularity in the calculus of variations. Arch. Rational Mech. Anal., 95(3):227–252, 1986.
- [20] M. Focardi. Semicontinuity of vectorial functionals in Orlicz-Sobolev spaces. Rend. Istit. Mat. Univ. Trieste, 29(1-2):141-161 (1998), 1997.
- [21] Mariano Giaquinta. Multiple Integrals in the Calculus of Variations and nonlinear elliptic Systems. Lectures in mathematics. Annals of Mathematics Studies, Princeton University Press, 1982.
- [22] E. Giusti. Metodi diretti nel calcolo delle variazioni. Unione Matematica Italiana, Bologna, 1994.
- [23] E. Giusti and M. Miranda. Sulla regolarità delle soluzioni deboli di una classe di sistemi ellittici quasi-lineari. Arch. Rational Mech. Anal., 31:173–184, 1968/1969.
- [24] M. A. Krasnosel'skij and Ya. B. Rutitskij. Convex functions and Orlicz spaces. Groningen-The Netherlands: P. Noordhoff Ltd. IX, 249 p., 1961.
- [25] Paolo Marcellini. Approximation of quasiconvex functions, and lower semicontinuity of multiple integrals. Manuscripta Math., 51(1-3):1-28, 1985.
- [26] Charles B. Morrey, Jr. Quasi-convexity and the lower semicontinuity of multiple integrals. Pacific J. Math., 2:25–53, 1952.
- [27] Jaak Peetre. A new approach in interpolation spaces. Studia Math., 34:23–42, 1970.
- [28] M. M. Rao and Z. D. Ren. Theory of Orlicz spaces, volume 146 of Monographs and Textbooks in Pure and Applied Mathematics. Marcel Dekker Inc., New York, 1991.
- [29] M. Růžička and L. Diening. Non-Newtonian fluids and function spaces. In Nonlinear Analysis, Function Spaces and Applications, Proceedings of NAFSA 2006 Prague, volume 8, pages 95–144, 2007.
- [30] Leon Simon. Theorems on regularity and singularity of energy minimizing maps. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 1996. Based on lecture notes by Norbert Hungerbühler.
- [31] A. Verde and G. Zecca. Lower semicontinuity of certain quasiconvex functionals in Orlicz-Sobolev spaces. Nonlinear Anal., 71(10):4515–4524, 2009.

¹ LMU Munich, Institute of Mathematics, Theresienstr. 39, 80333 Munich, Germany. Email: Lars.diening@mathematik.uni-muenchen.de. The work was partially supported by Gnampa.

² UNIVERSITÄT REGENSBURG, FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄTSSTR. 31, 93053 RE-GENSBURG, GERMANY. EMAIL: DANIEL.LENGELER@MATHEMATIK.UNI-REGENSBURG.DE. THE WORK WAS PARTIALLY SUPPORTED BY THE EUROPEAN RESEARCH COUNCIL UNDER FP7, ADVANCED GRANT N. 226234 "ANALYTIC TECHNIQUES FOR GEOMETRIC AND FUNCTIONAL INEQUALITIES".

³ DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI NAPOLI, FEDERICO II, VIA CINTIA, 80126 NAPLES, ITALY. EMAIL: BSTROFFO@UNINA.IT. THE WORK WAS PARTIALLY SUPPORTED BY PRIN PROJECT: "CALCOLO DELLE VARIAZIONI E TEORIA GEOMETRICA DELLA MISURA".

 4 Dipartimento di Matematica, Università di Napoli, Federico II, Via Cintia, 80126 Naples, Italy. email: anverde@unina.it