

On Hall subgroups of a finite group

Wenbin Guo*

Department of Mathematics, University of Science and Technology of China,
Hefei 230026, P. R. China
E-mail: wbguo@ustc.edu.cn

Alexander N. Skiba †

Department of Mathematics, Francisk Skorina Gomel State University,
Gomel 246019, Belarus
E-mail: alexander.skiba49@gmail.com

November 5, 2018

Abstract

In the paper new criteria of existence and conjugacy of Hall subgroups of finite groups are given.

1 Introduction

Throughout this paper, all groups are finite and G always denotes a finite group. We use \mathbb{P} to denote the set of all primes, π is a non-empty subset of \mathbb{P} and $\pi' = \mathbb{P} \setminus \pi$. A natural number n is called a π -number if $(n, p) = 1$ for any prime $p \in \pi'$. If $|G|$ is a π -number, then G is said to be a π -group.

Let A and B be subgroups of G . Then A is said to permute with B if $AB = BA$. If A permutes with all (Sylow) subgroups of B , then A is called *quasinormal* (*S-quasinormal*, respectively) in B .

A group G is said to be:

- (a) an E_π -group if G has a Hall π -subgroup;
- (b) a C_π -group if G is an E_π -group and any two Hall π -subgroups of G are conjugate;

*Research of the first author is supported by a NNSF grant of China (Grant # 11071229) and Wu Wen-Tsun Key Laboratory of Mathematics, USTC, Chinese Academy of Sciences.

†Research of the second author supported by Chinese Academy of Sciences Visiting Professorship for Senior International Scientists (grant No. 2010T2J12)

Keywords: finite group, soluble group, Sylow subgroup, Hall subgroup, quasinormal subgroup, E_π -group, C_π -group, D_π -group.

Mathematics Subject Classification (2010): 20D20

(c) a D_π -group if G is an E_π -group and any π -subgroup of G is contained in some Hall π -subgroup of G .

A group G is said to be π -separable if G has a chief series

$$1 = G_0 \leq H_1 \leq \dots \leq H_{t-1} \leq H_t = G, \quad (*)$$

where each index $|H_i : H_{i-1}|$ is either a π -number or a π' -number.

The most important result of the theory of π -separable groups is the following classical result.

Theorem A (P. Hall, S.A. Čunihin). *Any π -separable group is a D_π -group.*

Our main goal here is to prove the following generalization of this theorem.

Theorem 1.1. *Suppose that G has a subgroup series*

$$1 = H_0 < H_1 \leq \dots \leq H_{t-1} \leq H_t = G,$$

where $|H_{i+1} : H_i|$ is either a π -number or a π' -number for all $i = 1, \dots, t$. Suppose also that G has a subgroup T such that $H_1T = G$ and $|G : T|$ is a π' -number. If H_i is quasinormal in T , for all $i = 1, \dots, t$, then G is a D_π -group.

The following theorem is one of the main steps in the proof of Theorem 1.1.

Theorem 1.2. *Suppose that G has a subgroup series*

$$1 = H_0 < H_1 \leq \dots \leq H_{t-1} \leq H_t = G,$$

where $|H_{i+1} : H_i|$ is either a π -number or a π' -number for all $i = 1, \dots, t$. Suppose also that G has a subgroup T such that $H_1T = G$ and $|G : T|$ is a π' -number. If H_i is S -quasinormal in T , for all $i = 1, \dots, t$, then G is a C_π -group.

Theorem 1.1 strengthens the main result in [1]. Note also that Example 1.1 in [1] shows that, under the conditions in Theorems 1.1 or 1.2 the group G is not necessary π -separable.

2 Preliminaries

Lemma 2.1. *Let N be a normal C_π -subgroup of G .*

(i) *If G/N is a C_π -group, then G is a C_π -group (S. A. Čunihin [3]).*

(ii) *If G/N is an E_π -group, then G is an E_π -group*

(iii) *If G has a nilpotent Hall π -subgroup, then G is a D_π -group ([19]).*

(iv) *If G has a Hall π -subgroup with cyclic Sylow subgroups, then G is a D_π -group (S. A. Rusakov [20]).*

Lemma 2.2. *Let N be a normal C_π -subgroup of G and N_π a Hall π -subgroup of N .*

(i) If G is a C_π -group, then G/N is a C_π -group (See [4, Lemma 9]).

(ii) If every Sylow subgroup of N_π is cyclic and G/N is a D_π -group, then G is a D_π -group (See [5] or [6, Chapter IV, Theorem 18.17]).

(iii) G is a D_π -group if and only if G/N is a D_π -group (See [7]).

Lemma 2.3 (O. Kegel [8]). *Let A and B be subgroups of G such that $G \neq AB$ and $AB^x = B^x A$, for all $x \in G$. Then G has a proper normal subgroup N such that either $A \leq N$ or $B \leq N$.*

Let A be a subgroup of G . A subgroup T is said to be a minimal supplement of A in G if $AT = G$ but $AT_0 \neq G$ for all proper subgroups T_0 of T .

The following lemma is obvious.

Lemma 2.4. *If N is normal in G and T is a minimal supplement of N in G , then $N \cap T \leq \Phi(T)$.*

Lemma 2.5 (P. Hall [10]). *Suppose that G has a Hall p' -subgroup for each prime p dividing $|G|$. Then G is soluble.*

Let A and B be subgroups of G and $\emptyset \neq X \subseteq G$. Following [9], we say that A is X -permutable (or A X -permutes) with B if $AB^x = B^x A$ for some $x \in X$.

The following lemma is also evident.

Lemma 2.6. *Let A, B, X be subgroups of G and $K \trianglelefteq G$. If A is X -permutable with B , then AK/K is XK/K -permutable with BK/K in G/K .*

Lemma 2.7 (Kegel [11]). *If a subgroup A of G permutes with all Sylow subgroups of G , then A is subnormal in G .*

Lemma 2.8 (H. Wielandt [12]). *If a π -subgroup A of G is subnormal in G , then $A \leq O_\pi(G)$.*

Lemma 2.9 (V. N. Knyagina and V. S. Monakhov [17]). *Let H, K and N be subgroups of G . If H is a Hall subgroup of G , then*

$$N \cap HK = (N \cap H)(N \cap K).$$

Lemma 2.10. *Let $U \leq B \leq G$ and $G = AB$, where the subgroup A permutes with U^b for all $b \in B$. Then A permutes with U^x for all $x \in G$.*

Proof. Since $G = AB$, $x = ab$ for some $a \in A$ and $b \in B$. Hence

$$AU^x = AU^{ab} = Aa(U^b)a^{-1} = a(U^b)a^{-1}A = U^x A.$$

Lemma 2.11 [13, Chapter I, Lemma 1.1.19]. *Let $A, B \leq G$ and $G = AB$. Then $G_p = A_p B_p$ for some $G_p \in \text{Syl}_p(G)$, $A_p \in \text{Syl}_p(A)$ and $B_p \in \text{Syl}_p(B)$.*

Lemma 2.12. *Let X be a normal C_π -subgroup of G . Suppose that G has a subgroup series*

$$1 = H_0 \leq H_1 \leq \dots \leq H_{t-1} \leq H_t = G$$

and a supplement T of H_1 in G such that H_i X -permutes with every Sylow subgroup of T for all $i = 1, \dots, t$. If each index $|H_{i+1} : H_i|$ is either a π -number or a π' -number, then G is an E_π -group.

Proof. Consider the subgroup series

$$1 = H_0X/X \leq H_1X/X \leq \dots \leq H_{t-1}X/X \leq H_tX/X = G/X \quad (1)$$

in G/X . If $T \leq X$, then $G = H_1T = XH_1$. Hence G is an E_π -group by Lemma 2.1(2). Now let $T \not\leq X$ and P/X be a Sylow p -subgroup of TX/X . Then by Lemma 2.11 there are Sylow p -subgroups T_p of T and X_p of X such that T_pX_p is a Sylow p -subgroup of TX and $T_pX_pX = T_pX = P$. Hence by Lemma 2.6, H_iX/X permutes with every Sylow subgroup of TX/X for all $i = 1, \dots, t$. On the other hand, since

$$|H_{i+1}X/X : H_iX/X| = |H_{i+1}X : H_iX| = |H_{i+1} : H_i| \cdot |X \cap H_{i+1} : X \cap H_i|,$$

every index of Series (1) is either π -number or π' -number. Hence the assertion follows from Theorem B in [1].

Lemma 2.13 (see Lemma in [14, Chapter A]). *Let H, K and N be subgroups of G . If $HK = KH$ and $HN = NH$, then $H\langle K, N \rangle = \langle K, N \rangle H$.*

3 Base proposition

Proposition 3.1. *Let X be a normal C_π -subgroup of G and A a subgroup of G such that $|G : A|$ is a π -number. Suppose that A has a Hall π -subgroup A_0 such that either A_0 is nilpotent or every Sylow subgroup of A_0 is cyclic. Suppose that A X -permutes with every Sylow p -subgroup of G for all primes $p \in \pi$ or for all primes $p \in \pi \setminus \{q\}$ for some prime q dividing $|G : A|$. Then G is a C_π -group.*

Proof. Assume that this proposition is false and let G be a counterexample of minimal order. Then $|\pi \cap \pi(G)| > 1$.

(1) G/R is a C_π -group for any non-identity normal subgroup R of G .

In order to prove this assertion, in view of the choice of G , it is enough to show that the hypothesis is still true for $(G/R, AR/R, XR/R)$. First note that $|G/R : AR/R| = |G : AR|$ is a π -number, and A_0R/R is a Hall π -subgroup of AR/R since

$$|AR/R : A_0R/R| = |AR : A_0R| = |A : A \cap A_0R| = |A : A_0(A \cap R)|.$$

On the other hand, $XR/R \simeq X/X \cap R$ is a C_π -group by Lemma 2.2 (i), and either $A_0R/R \simeq A_0/R \cap A_0$ is nilpotent or every Sylow subgroup of A_0R/R is cyclic. Finally, let P/R be a Sylow p -subgroup of G/R , where $p \in \pi \setminus \{q\}$. Then for some Sylow p -subgroup G_p we have $G_pR/R = P/R$. Hence AR/R XR/R -permutes with P/R by Lemma 2.6. Therefore the hypothesis holds for $(G/R, AR/R, XR/R)$.

(2) $X = 1$.

Indeed, if $X \neq 1$, then G/X is a C_π -group by (1). Hence G is C_π -group by Lemma 2.1 (i), a contradiction.

(3) G has a proper non-identity normal subgroup N .

Let $p \in \pi \cap \pi(G)$, where $p \neq q$. Let P be a Sylow p -subgroup of G . First assume that $AP = G$. Since $|\pi \cap \pi(G)| > 1$, there is a prime $r \in \pi \cap \pi(G)$ such that $r \neq p$, so r does not divide $|G : A|$. Let R be a Sylow r -subgroup of G . Then for any $x \in G$ we have $AR^x = R^x A$. Hence $R \leq A_G$. Since G is not a C_π -group, $A \neq G$ by Lemma 2.1 (iii)(iv). Hence $1 \neq A_G \neq G$. Now suppose that $AP \neq G$. By (2), $P^x A = AP^x$ for all $x \in G$. Hence we have (3) by Lemma 2.3.

(4) N is a C_π -group.

In view of the choice of G it is enough to prove that the hypothesis holds for (N, A_1) , where $A_1 = A \cap N$. Since $|AN : A| = |N : A \cap N|$, $|N : A_1|$ is a π -number. On the other hand, $A_0 \cap N$ is a Hall π -subgroup of N since

$$|A \cap N : A_0 \cap N| = |A_0(A \cap N) : A_0|.$$

It clear also that either $A_0 \cap N$ is nilpotent or every Sylow subgroup of $A_0 \cap N$ is cyclic. Now let N_r be any Sylow r -subgroup of N , where $r \in \pi \setminus \{q\}$. Then for some Sylow r -subgroup G_r of G we have $N_r = G_r \cap N$ and

$$N \cap G_r = (A \cap N)(N \cap G_r) = A_1 N_r = N_r A_1$$

by Lemma 2.9. Therefore the hypothesis holds for (N, A_1) .

Finally, in view of (1) and (4), G is a C_π -group by Lemma 2.1 (i), which contradicts the choice of G .

Corollary 3.2. *Let X be a normal C_π -subgroup of G and A a subgroup of G such that $|G : A|$ is a π -number. Suppose that A has a Hall π -subgroup A_0 such that either A_0 is nilpotent or every Sylow subgroup of A_0 is cyclic. Suppose that G has a subgroup T such that $|G : T|$ is a π' -number, $G = AT$ and A X -permutes with every Sylow p -subgroup of T for all primes $p \in \pi(G) \setminus \pi$, $p \neq q \in \mathbb{P}$. Then G is a $C_{\pi'}$ -group.*

Proof. Let $p \in \pi(G) \setminus \pi$, $p \neq q$ and P be a Sylow p -subgroup of G . Since $G = AT$ and $|G : T|$ is a π' -number, every Sylow p -subgroup T_p of T is a Sylow p -subgroup of G . Hence for some $x \in G$ we have $T_p = P^x$, so A permutes with P by Lemma 2.10. Therefore Corollary 3.2 follows from Proposition 3.1.

Corollary 3.3 (see Theorem 1.2 in [9]). *Let X be a normal nilpotent subgroup of G and A a Hall π -subgroup of G . Suppose $G = AT$ and A X -permutes with every subgroup of T Then G is a $C_{\pi'}$ -group.*

Corollary 3.4 (See Theorem B in [16]). *Let A be a Hall subgroup of a group G and T a minimal supplement of A in G . Suppose that A permutes with all Sylow subgroups of T and with all maximal subgroups of any Sylow subgroup of T . Then G is a $C_{\pi'}$ -group.*

Corollary 3.5 (see Theorem A* in [1]). *Let H be a Hall π -subgroup of G . Let $G = HT$ for some subgroup T of G , and q a prime. If H permutes with every Sylow p -subgroup of T for all primes $p \neq q$, then T contains a complement of H in G and any two complements of H in G are conjugate.*

Proof. It is clear that $T \cap H$ is a Hall π -subgroup of T . Moreover, if P is a Sylow subgroup of T and $HP = PH$, then $HP \cap T = P(H \cap T) = (H \cap T)P$. Hence by Proposition 3.1, T is a C_π -group. Now Corollary 3.5 follows from Proposition 3.1.

Corollary 3.6 (see [18]). *Let A be a Hall π -subgroup of G , $G = AT$ and A permutes with every subgroup of T . Then G is an $E_{\pi'}$ -group.*

4 Proof of Theorem 1.1

Theorem 1.2 is a special case (when $X = 1$) of the following theorem.

Theorem 4.1. *Let X be a normal C_π -subgroup of G . Suppose that G has a subgroup series*

$$1 = H_0 < H_1 \leq \dots \leq H_{t-1} \leq H_t = G,$$

where $|H_{i+1} : H_i|$ is either a π -number or a π' -number for all $i = 1, \dots, t$. Suppose that G has a subgroup T such that $H_1T = G$ and $|G : T|$ is a π' -number. If H_i is X -permutable with each Sylow subgroup of T , for all $i = 1, \dots, t$, then G is a C_π -group.

Proof. Suppose that this theorem is false and let G be a counterexample of minimal order. By Lemma 2.12, G has a Hall π -subgroup S . Hence some Hall π -subgroup S_1 of G is not conjugated with S . Without loss of generality, we may assume that $H_1 \neq 1$. Since $|G : T|$ is π' -number, every Sylow p -subgroup P of T , where $p \in \pi$, is also a Sylow p -subgroup of G . We proceed the proof via the following steps.

(1) G/N is a C_π -group for every non-trivial quotient G/N of G .

We consider the subgroup series

$$1 = H_0N/N \leq H_1N/N \leq \dots \leq H_{t-1}N/N \leq H_tN/N = G/N \quad (2)$$

in G/N . Then $(H_1N/N)(TN/N) = G/N$ and $|G/N : TN/N| = |G : TN|$ is a π' -number. Moreover, by Lemma 2.6, H_iN/N is XN/N -permutable with every Sylow subgroup of TN/N for all $i = 1, \dots, t$. On the other hand, since

$$|H_{i+1}N/N : H_iN/N| = |H_{i+1}N : H_iN| = |H_{i+1} : H_i| \cdot |N \cap H_{i+1} : N \cap H_i|,$$

every index of the series (2) is either π -number or π' -number. Moreover, $XN/N \simeq X/(X \cap N)$ is a C_π -group by Lemma 2.2 (i). All these show that the hypothesis holds for G/N . Hence in the case, where $N \neq 1$, G/N is a C_π -group by the choice of G .

(2) $O_{\pi'}(G) = 1 = O_\pi(G)$.

Suppose that $D = O_{\pi'}(G) \neq 1$. Then by (1), there is an element $x \in G$ such that $S_1^x D = SD$. But by the Schur-Zassenhaus theorem, S_1^x and S are conjugate in SD , which implies that S_1 and S are conjugate in G . This contradiction shows that $O_{\pi'}(G) = 1$. Analogously, one can prove that $O_{\pi}(G) = 1$.

(3) $X = 1$ (This follows from (1), Lemma 2.1 (i) and the choice of G).

(4) $T \neq G$.

Suppose that $T = G$. Then by hypothesis and (3), H_1 permutes with all Sylow subgroups of G . It follows from Lemma 2.7 that H_1 is subnormal in G . Since H_1 is either a π -group or a π' -group, $H_1 \leq O_{\pi}(G)$ or $H_1 \leq O_{\pi'}(G)$ by Lemma 2.8. It follows from (2) that $H_1 = 1$, which contradicts to our assumption about H_1 . Hence (4) holds.

(5) *The hypothesis holds for T .*

Consider the subgroup series

$$1 = H_0 \cap T \leq H_1 \cap T \leq \dots \leq H_{t-1} \cap T \leq H_t \cap T = T \quad (3)$$

of the group T . Since

$$H_{i+1} = H_i T \cap H_{i+1} = H_i(H_{i+1} \cap T)$$

we have

$$|H_{i+1} : H_i| = |H_{i+1} \cap T : H_i \cap T|,$$

for all $i = 1, \dots, t-1$ and $|H_1 \cap T : H_0 \cap T| = |H_1 \cap T|$ divides $|H_1 : 1|$, we see that every index of the series (3) is either π -number or π' -number. Now let E be a Sylow subgroup of T . By (3) and the hypothesis, $H_i E = E H_i$. Hence

$$H_i E \cap T = E(H_i \cap T) = (H_i \cap T)E.$$

This shows that the hypothesis holds for T .

(6) *T is a C_{π} -group.*

Since $T \neq G$ by (4), and the hypothesis holds for T by (5), the minimal choice of G implies that (6) holds.

(7) *T is a $E_{\pi'}$ -subgroup.*

This follows from (3), (5) and Lemma 2.12.

(8) *Let $T_{\pi'}$ be a Hall π' -subgroup of T and D a normal subgroup of G . Then $T_{\pi'} \neq 1$, and if either $H_1 \leq D$ or $T_{\pi'} \leq D$, then $D = G$.*

Suppose that $T_{\pi'} = 1$. Then H_1 is a Hall π' -subgroup of G . Therefore G is a C_{π} -group by Corollary 3.2, a contradiction. Hence $T_{\pi'} \neq 1$.

We show that the hypothesis holds for D . Consider the subgroup series

$$1 = D_0 \leq D_1 \leq \dots \leq D_{t-1} \leq D_t = D,$$

where $D_i = H_i \cap D$ for all $i = 1, \dots, t$. Let $T_0 = D \cap T$. First we show that $D_1 T_0 = D$. If $H_1 \leq D$, then

$$D = H_1(D \cap T) = H_1 T_0 = D_1 T_0.$$

Now suppose that $T_{\pi'} \leq D$. In view of (3), H_1 permutes with $T_{\pi'}$. Since $H_1 T = G$, $T \neq G$ and $|G : T|$ is a π' -number, H_1 is a π' -group. Hence $H_1 T_{\pi'}$ is a Hall π' -subgroup of G . Therefore

$$D = (D \cap H_1 T_{\pi'})(D \cap T_{\pi}) = T_{\pi'}(D \cap H_1)(D \cap T_{\pi}) = (D \cap H_1)(T_{\pi'}(D \cap T_{\pi})) = (D \cap H_1)(D \cap T) = D_1 T_0.$$

It is also clear that $|D : T_0|$ is a π' -number. Now let P be a Sylow p -subgroup of T_0 . Then for some Sylow p -subgroup T_p of T we have $P = T_p \cap D$. Hence in view of Lemma 2.9,

$$D \cap H_i T_p = (D \cap H_i)(D \cap T_p) = D_i P = P D_i.$$

Thus for any $p \in \pi$, D_i is permutable with every Sylow p -subgroup of T_0 for all $i = 1, \dots, t$. Finally, since the number

$$|D_i : D_{i-1}| = |(D \cap H_i)H_{i-1} : H_{i-1}|$$

divides $|H_i : H_{i-1}|$, each index $|D_i : D_{i-1}|$ is either a π -number or a π' -number. Therefore the hypothesis holds for D . Suppose that $D \neq G$. Then D is a C_{π} -group by the choice of G . Since either $1 \neq H_1 \leq D$ or $1 \neq T_{\pi'} \leq D$, G/D is a C_{π} -group by (1). It follows from Lemma 2.1 (i) that G is a C_{π} -group, which contradicts the choice of G . Hence, (8) holds.

Final contradiction. Since $G = H_1 T$ and H_1 permutes with all Sylow subgroups of T by (3),

$$H_1(T_{\pi'})^x = (T_{\pi'})^x H_1$$

for all $x \in G$ by Lemma 2.10. Therefore by Lemma 2.3, either $H_1^G \neq G$ or $(T_{\pi'})^G \neq G$. But in view of (8) both these cases are impossible. The contradiction completes the proof of the result.

Corollary 4.2 (see Theorem 5.1 in [1]). *Let X be a normal π -separable subgroup of G . Suppose that G has a subgroup series*

$$1 = H_0 < H_1 \leq \dots \leq H_{t-1} \leq H_t = G,$$

where $|H_{i+1} : H_i|$ is either a π -number or a π' -number for all $i = 1, \dots, t$. Suppose that G has a subgroup T such that $H_1 T = G$ and $|G : T|$ is a π' -number. If H_i is X -permutable with each subgroup of T , for all $i = 1, \dots, t$, then G is a C_{π} -group.

Theorem 1.1 is a special case (when $X = 1$) of the following theorem.

Theorem 4.3. *Let X be a normal an E_{π} -subgroup of G and X_{π} a Hall π -subgroup of X . Suppose that G has a subgroup series*

$$1 = H_0 < H_1 \leq \dots \leq H_{t-1} \leq H_t = G,$$

where $|H_{i+1} : H_i|$ is either a π -number or a π' -number for all $i = 1, \dots, t$. Suppose that G has a subgroup T such that $H_1 T = G$ and $|G : T|$ is a π' -number.

(i) Suppose that the Sylow subgroups of X_π are cyclic. If H_i is X -permutable with each cyclic subgroup H of T of prime power order, for all $i = 1, \dots, t$, then G is a D_π -group.

(ii) Suppose that X is a D_π -subgroup. If H_i is X -permutable with each cyclic subgroup H of T of prime power order, for all $i = 1, \dots, t$, then G is a D_π -group.

Proof. (i) Suppose that this assertion is false and let G be a counterexample of minimal order. In view of Theorem 4.1, G is a C_π -group. Hence there is a π -subgroup U of G such that for any Hall π -subgroup E of G we have $U \not\leq E$.

(1) G/N is a D_π -group for any non-identity normal subgroup N of G .

Consider the subgroup series

$$1 = H_0N/N \leq H_1N/N \leq \dots \leq H_{t-1}N/N \leq H_tN/N = G/N.$$

It is clear that $(H_1N/N)(TN/N) = G/N$, $|G/N : TN/N|$ is a π' -number and $|H_{i+1}N/N : H_iN/N|$ is either a π -number or a π' -number for all $i = 1, \dots, t$ (see (1) in the proof of Theorem 4.1). Now let H/N be a cyclic subgroup of TN/N of prime power order $|H/N|$. Then $H = N(H \cap T)$. Let W be a group of minimal order with the properties that $W \leq H \cap T$ and $NW = H$. If $N \cap W \not\leq \Phi(W)$, then for some maximal subgroup S of W we have $(N \cap W)S = W$. Hence $H = NW = N(N \cap W)S = NS$, a contradiction. Hence $N \cap W \leq \Phi(W)$. Since $W/W \cap N \simeq H/N$ is a cyclic group of prime power order, it follows that W is cyclic group of prime power order. Hence H_i is X -permutable with W . Thus H_iN/N is XN/N -permutable with $WN/N = H/N$ by Lemma 2.6. Therefore the hypothesis holds for G/N . But since $N \neq 1$, $|G/N| < |G|$ and so G/N is a D_π -group by the choice of G .

(2) $X = 1$.

Suppose that $X \neq 1$. Then G/X is a D_π -group by (1). Hence G is a D_π -group by Lemma 2.2 (iii), a contradiction. Thus we have (2).

(3) H_1 permutes with every subgroup of U .

Let Z be any cyclic subgroup of U of prime power order p^n . Then $p \in \pi$ and $Z \leq G_p$ for some Sylow p -subgroup G_p of G . Since $|G : T|$ is a π' -number, there is an element $x = ht$ such that $(G_p)^x \leq T$. Hence $H_1Z = ZH_1$ by Lemma 2.10, which in view of Lemma 2.13 implies that $H_1U = UH_1$.

(4) $T \neq G$ (see the proof of (4) in the proof of Theorem 4.1).

(5) T is a D_π -group.

The hypothesis holds for T (see (5) in the proof of Theorem 4.1), so in view of (4) the minimal choice of G implies that we have (5).

(6) $V = H_1U$ is a C_π -group.

Indeed, V is a group by (3), and since $T \neq G$, H_1 is a Hall π' -subgroup of V . Therefore we have (6) by (2) and Corollary 3.2.

(7) $|V : T \cap V|$ is a π' -number and $T \cap V$ is a C_π -group.

First note that $|G : T| = |V : T \cap V|$ is a π' -number. But H_1 is a Hall π' -subgroup of V . Hence $V = H_1(T \cap V)$, which implies that

$$|V : H_1| = |T \cap V : H_1 \cap T \cap V|$$

is a π -number. Hence $A = H_1 \cap T \cap V$ is a Hall π' -subgroup of $T \cap V$. Finally, if W is any π -subgroup of $T \cap V =$, then $H_1W = WH_1$ by (3). Therefore

$$AW = (H_1 \cap T \cap V)W = (H_1W \cap T \cap V) = WA.$$

Hence $T \cap V$ is a C_π -group by Proposition 3.1.

Final contradiction for (i). In view of (7), $|V : T \cap V|$ is a π' -number and $T \cap V$ is a C_π -group. Hence in view of (6), there is an element $x \in V$ such that $U^x \leq T \cap V \leq T$. But by (5), T is a D_π -group. Hence for some Hall π -subgroup T_π of T we have $U \leq T_\pi$, which is a contradiction since T_π is clearly a Hall π -subgroup of G .

(ii) See the proof of (i) and use Lemma 2.2 (iii).

5 Groups with soluble Hall π -subgroups

A group G is said to be:

- (a) an E_π^S -group (E_π^N -group) if G has a soluble (a nilpotent, respectively) Hall π -subgroup;
- (b) a C_π^S -group (C_π^N -group) if G has a soluble (nilpotent, respectively) Hall π -subgroup and G is a C_π -group;
- (c) a D_π^S -group if G is a C_π^S -group and any π -subgroup of G is contained in some Hall π -subgroup of G .

Our next results are new criteria for a group to be an E_π^S -group.

Lemma 5.1. *Let N be a normal E_π -subgroup of G and N_π a Hall π -subgroup of N . Suppose that N_π is nilpotent. If G/N is a D_π^S -group, then G is a D_π^S -group (See [4] or [6, Chapter IV, Theorem 18.15]).*

Theorem 5.2. *Let X be a normal E_π^N -subgroup of G . Suppose that G has a subgroup series*

$$1 = H_0 < H_1 \leq \dots \leq H_{t-1} \leq H_t = G,$$

where $|H_{i+1} : H_i|$ is divisible by at most one prime in π , for all $i = 1, \dots, t$. Suppose that G has a subgroup T such that $H_1T = G$ and $|G : T|$ is a π' -number. If H_i is X -permutable with each Sylow subgroup of T , for all $i = 1, \dots, t$, then G is an E_π^S -group.

Proof. Suppose that this theorem is false and let G be a counterexample of minimal order.

(1) Every non-trivial quotient G/N of G is an E_π^S -group (See (1) in the proof of Theorem 4.1).

(2) $X = 1$.

Suppose that $X \neq 1$. Then G/X of G is an E_π^S -group by (1). Let E/X be a soluble π -Hall subgroup of G/X . Then, by Lemma 5.1, E is a D_π^S -group and if U is a Hall subgroup of E , then U is a Hall subgroup of G . Hence G is an E_π^S -group, a contradiction. Thus we have (2).

(3) $T \neq G$.

Suppose that $T = G$. Then by hypothesis and (2), H_1 permutes with all Sylow subgroups of G . Hence H_1 is subnormal in G by Lemma 2.7, so

$$1 < H_1 \leq O_{\pi' \cup \{p\}}(G)$$

for some $p \in \pi$ by Lemma 2.8. Therefore $O_{\pi' \cup \{p\}}(G)$ is a non-identity normal D_π^N -group-subgroup of G , which as in the proof of (2), conducts us to the contradiction. Thus (3) holds.

Final contradiction. In view of (2), the hypothesis is true for T (see (5) in the proof of Theorem 4.1). Hence T is an E_π^S -group by (3) and the choice of G . But since $|G : T|$ is a π' -number, any Hall π -subgroup of T is a Hall π -subgroup of G as well. Hence G is an E_π^S -group, a contradiction.

The theorem is proved.

Theorem 5.3. *Let X be a normal E_π^N -subgroup of G . Suppose that G has a subgroup series*

$$1 = H_0 < H_1 \leq \dots \leq H_{t-1} \leq H_t = G,$$

where $|H_{i+1} : H_i|$ is divisible by at most one prime in π , for all $i = 1, \dots, t$. Suppose that G has a subgroup T such that $H_1 T = G$ and $|G : T|$ is a π' -number. If H_i is X -permutable with each cyclic subgroup H of T of prime power order, for all $i = 1, \dots, t$, then G is a D_π^S -group.

Proof. Suppose that this theorem is false and let G be a counterexample of minimal order. Then $|\pi \cap \pi(G)| > 1$. Without loss we may suppose that $H_1 \neq 1$ and $H_{t-1} \neq G$.

(1) Every on-trivial quotient G/N of G is a D_π^S -group (See (1) in the proof of Theorem 4.3).

(2) If N is a normal E_π^N -subgroup of G , then $N = 1$. In particular, $X = 1$.

Suppose that $N \neq 1$. Then G/N of G is a D_π^S -group by (1) and the choice of G . Hence G is a D_π^S -group by Lemma 5.1, which contradicts the choice of G . Hence we have (2).

(3) $T \neq G$ (See (3) in the proof of Theorem 5.2).

(4) The hypothesis holds for H_{t-1} and T .

Since

$$H_{t-1} = H_1(H_{t-1} \cap T)$$

and

$$|H_{t-1} : H_{t-1} \cap T| = |G : T|,$$

the hypothesis holds for H_{t-1} by (2). The second assertion of (4) may be proved as (5) in the proof of Theorem 4.1.

(5) G is a C_π^S -group.

Since $|G : H_{t-1}|$ is divisible by at most one prime in π and $|\pi \cap \pi(G)| > 1$, there is a prime $p \in \pi \cap \pi(G)$ such that for a Sylow p -subgroup P of G we have $P \leq H_{t-1}$, so in view of (2) and Lemma 2.10 we have $1 < P \leq (H_{t-1})_G$. Since $H_{t-1} \neq G$ and the hypothesis holds for H_{t-1} , H_{t-1} is a D_π^S -group by the choice of G . Hence $(H_{t-1})_G$ is a C_π^S -group, so G is a C_π^S -group by Lemma 2.1 (i) and Theorem 5.2.

(6) For any i , H_i permutes with every π -subgroup of G (see (3) in the proof of Theorem 4.3).

(7) $V = H_1U$ is a D_π^S -group for any π -subgroup U of G .

By (6), V is a subgroup of G . Moreover, V is a C_π -group by Proposition 3.1. We show that the hypothesis holds for V . Let V_π be a Hall π -subgroup of V . Then $V = H_1V_\pi$ and

$$V_i = V \cap H_i = H_1(V_\pi \cap H_i),$$

for all $i = 1, \dots, t$. Let W be any subgroup of V_π . Then

$$V_iW = (H_1(V_\pi \cap H_i))W = H_1(V_\pi \cap H_iW) = H_1W(V_\pi \cap H_i)W(H_1(V_\pi \cap H_i)) = WWV_i.$$

It is clear also that $|V_{i+1} : V_i|$ is divisible by at most one prime in π , for all $i = 0, \dots, t$. Hence the hypothesis holds for V . Suppose that $G = V$. In this case, in view of (5), we may suppose that V_π is a soluble Hall π -subgroup of G . Let L be a minimal normal subgroup of V_π . Then

$$L^G = L^{V_\pi H_1} = L^{H_1} \leq LH_1 \cap L^{H_1} = L(L^G \cap H_1)$$

and L is a q -group for some $q \in \pi$. From (2) it follows that for some prime $p \in \pi$ with $q \neq p$ we have $p \in \pi(L^G \cap H_1)$ and a Sylow p -subgroup P of $L^G \cap H_1$ is also a Sylow subgroup of L^G . For some Sylow p -subgroup G_p of G we have $P = L^G \cap G_p$, so

$$L^G \cap H_1 G_p = (L^G \cap H_1)(L^G \cap G_p) = (L^G \cap H_1)P = P(L^G \cap H_1).$$

Hence $P \leq (L^G \cap H_1)_{L^G}$. Therefore G has a non-identity subnormal subgroup $R = (L^G \cap H_1)_{L^G}$ of order divisible by at most one prime in π , which contradicts (2). Hence $V \neq G$, so V is a D_π^S -group by the choice of G .

Final contradiction. Let U be any π -subgroup of G and $V = H_1U$. Then V is a D_π^S -subgroup of G by (7), and $|G : T| = |V : V \cap T|$ is a π' -number. By (3) and (4), T is a D_π -group. Hence for some Hall π -subgroup V_π of V we have $U \leq V_\pi$, and $V_\pi \leq T_\pi$. But since $|G : T|$ is a π' -number, T_π is a Hall π -subgroup of G . Therefore G is a D_π^S -group.

Corollary 5.4. *Suppose that G has a subgroup series*

$$1 = H_0 < H_1 \leq \dots \leq H_{t-1} \leq H_t = G,$$

where $|H_{i+1} : H_i|$ is divisible by at most one prime in π , for all $i = 1, \dots, t$. Suppose also that G has a subgroup T such that $H_1T = G$ and $|G : T|$ is a π' -number. If H_i is quasinormal in T , for all $i = 1, \dots, t$, then G is a D_π^S -group.

Corollary 5.6 (see S. A. Čuniĥin [21] or [6, Chapter IV, Theorem 18.13]). *If G has a chief subgroup series*

$$1 = H_0 < H_1 \leq \dots \leq H_{t-1} \leq H_t = G,$$

where $|H_{i+1} : H_i|$ is divisible by at most one prime in π , for all $i = 1, \dots, t$, then G is a D_π^S -group.

References

- [1] W. Guo, A.N. Skiba, New criteria of existence and conjugacy of Hall subgroups of finite groups, Proc. Amer. Math. Soc., 2011, **139**, 2327-2336.
- [2] S. A. Čuniĥin, On weakening the conditions in theorems of Sylow's type, Doklady Akad. Nauk SSSR (NS), **83** (1952), 663-665.
- [3] S. A. Čuniĥin, On existence and conjugacy of subgroups of a finite group, Mat. Sbornik, , 1953, **33** (1), 111-132.
- [4] P. Hall, Theorems like Sylow's, Proc. London Math. Soc., 1956, **6** (3), 286-304.
- [5] L.A. Shemetkov, On Sylow properties of finite groups, DAN BSSR, 1972, **16** (10), 881-883.
- [6] L.A. Shemetkov, Formations of Finite Groups, Nauka, Main Editorial Board for Physical and Mathematical Literature, Moscow, 1978.
- [7] D.O. Revin, E.P. Vdovin, Hall subgroups of finite groups, Contemp. Math., 2006, **402**, 229-265.
- [8] O. H. Kegel, Produkte nilpotenter Gruppen, Arch. Math., 1961, **12**, 90-93.
- [9] Wenbin Guo, K. P. Shum and Alexander N. Skiba, X -semipermutable subgroups of finite groups, J. Algebra, 2007, **315**, 31-41.
- [10] P. Hall, A characteristic property of soluble groups, J. London Math. Soc., 1937, **12**, 188-200.
- [11] O. Kegel, Sylow-Gruppen and Subnormalteiler endlicher Gruppen, Math. Z., 1962, **78** 205–221.
- [12] H. Wielandt, Subnormal subgroups and permutation groups. Lectures given at the Ohio State University, Columbus, Ohio, 1971.
- [13] A. Ballester-Bolinches, R. Esteban-Romero, M. Asaad, Products of Finite Groups, Walter de Gruyter, Berlin, New York, 2010.
- [14] K. Doerk, T. Hawkes, Finite Soluble Groups, Walter de Gruyter, Berlin, New York, 1992.

- [15] Wenbin Guo, K.P. Shum and Alexander N. Skiba, Schur-Zassenhaus Theorem for X -permutable subgroups, Algebra Colloquium, 2008, **15**(2), 185-192.
- [16] W. Guo and A.N. Skiba, Criteria of Existence of Hall Subgroups in Non-soluble Finite Groups, Acta Math. Sinica, 2010, **26**(2), 295-304.
- [17] B.N. Knyagina, V.S. Monakhov, On π' -properties of finite group having a Hall π -subgroup, Siberian. Math. J., 2011, **52** (2), 298-309.
- [18] N. Foguel, On seminormal subgroups, J. Algebra, 1994, **165**, 633-635.
- [19] H. Wielandt, Zum Satz von Sylow, II., Math. Z., 1959, **71**, 461-462.
- [20] S.A. Rusakov, Analogous of theorem of Sylow on existence and embedding subgroups, Siberian Math. J., 1963, **4**(5), 325-342.
- [21] S. A. Čuniĥin, On π -separate groups, Doklady Akad. Nauk. SSSR, 1948, **59**, 443-445.