

# A short proof of Paouris' inequality

Radosław Adamczak <sup>\*</sup>      Rafał Łatała <sup>†</sup>  
 Alexander E. Litvak <sup>‡</sup>      Krzysztof Oleszkiewicz <sup>§</sup>  
 Alain Pajor <sup>¶</sup>      Nicole Tomczak-Jaegermann <sup>||</sup>

## Abstract

We give a short proof of a result of G. Paouris on the tail behaviour of the Euclidean norm  $|X|$  of an isotropic log-concave random vector  $X \in \mathbb{R}^n$ , stating that for every  $t \geq 1$ ,

$$\mathbb{P}(|X| \geq ct\sqrt{n}) \leq \exp(-t\sqrt{n}).$$

More precisely we show that for any log-concave random vector  $X$  and any  $p \geq 1$ ,

$$(\mathbb{E}|X|^p)^{1/p} \sim \mathbb{E}|X| + \sup_{z \in S^{n-1}} (\mathbb{E}|\langle z, X \rangle|^p)^{1/p}.$$

AMS Classification: 46B06, 46B09 (Primary), 52A23 (Secondary)

**Key Words and Phrases:** log-concave random vectors, deviation inequalities

---

<sup>\*</sup>Research partially supported by MNiSW Grant no. N N201 397437.

<sup>†</sup>Research partially supported by MNiSW Grant no. N N201 397437.

<sup>‡</sup>Research partially supported by the E.W.R. Steacie Memorial Fellowship.

<sup>§</sup>Research partially supported by MNiSW Grant no. N N201 397437.

<sup>¶</sup>Research partially supported by the ANR project ANR-08-BLAN-0311-01.

<sup>||</sup>This author holds the Canada Research Chair in Geometric Analysis.

# 1 Introduction

Let  $X$  be a random vector in the Euclidean space  $\mathbb{R}^n$  equipped with its Euclidean norm  $|\cdot|$  and its scalar product  $\langle \cdot, \cdot \rangle$ . Assume that  $X$  has a log-concave distribution (a typical example of such a distribution is a random vector uniformly distributed on a convex body). Assume further that it is centered and its covariance matrix is the identity – such a random vector will be called *isotropic*. A famous and important result of ([14], Theorem 1.1) states that

**Theorem 1.** *There exists an absolute constant  $c > 0$  such that if  $X$  is an isotropic log-concave random vector in  $\mathbb{R}^n$ , then for every  $t \geq 1$ ,*

$$\mathbb{P}(|X| \geq ct\sqrt{n}) \leq \exp(-t\sqrt{n}).$$

This result had a huge impact on the study of log-concave measures and has a lot of applications in that subject.

A Borel probability measure on  $\mathbb{R}^n$  is called log-concave if for all  $0 < \theta < 1$  and all compact sets  $A, B \subset \mathbb{R}^n$  one has

$$\mu((1 - \theta)A + \theta B) \geq \mu(A)^{1-\theta} \mu(B)^\theta.$$

We refer to [5, 6] for a general study of this class of measures. Clearly, the affine image of a log-concave probability is also log-concave. The Euclidean norm of an  $n$ -dimensional log-concave random vector has moments of all orders (see [5]). A log-concave probability is supported on some convex subset of an affine subspace where it has a density. In particular when the support of the probability generates the whole space  $\mathbb{R}^n$  (in which case we talk, in short, about full-dimensional probability) a characterization of Borell (see [5, 6]) states that the probability is absolutely continuous with respect to the Lebesgue measure and has a density which is log-concave. We say that a random vector is log-concave if its distribution is a log-concave measure.

Let  $X \in \mathbb{R}^n$ , be a random vector, denote the weak  $p$ -th moment of  $X$  by

$$\sigma_p(X) = \sup_{z \in S^{n-1}} (\mathbb{E}|\langle z, X \rangle|^p)^{1/p}.$$

The purpose of this article is to give a short proof of the following

**Theorem 2.** *For any log-concave random vector  $X \in \mathbb{R}^n$  and any  $p \geq 1$ ,*

$$(\mathbb{E}|X|^p)^{1/p} \leq C(\mathbb{E}|X| + \sigma_p(X)),$$

where  $C$  is an absolute positive constant.

This result may be deduced directly from Paouris' work in [14]. Indeed, it is a consequence of Theorem 8.2 combined with Lemma 3.9 in that paper. As formulated here, Theorem 2 first appeared in [3] (Theorem 2). Note that because trivially a converse inequality is valid (with constant 1/2), Theorem 2 states in fact an equivalence for  $(\mathbb{E}|X|^p)^{1/p}$ .

It is noteworthy that the following strengthening of Theorem 2 is still open:  $(\mathbb{E}|X|^p)^{1/p} \leq \mathbb{E}|X| + C\sigma_p(X)$ , where  $C$  is an absolute positive constant.

If  $X$  is a log-concave random vector, then so is  $\langle z, X \rangle$  for every  $z \in S^{n-1}$ . It follows that there exists an absolute constant  $C' > 0$  such that for any  $p \geq 1$ ,  $\sigma_p(X) \leq C'p\sigma_2(X)$  ([5]). (In fact one can deduce this inequality with  $C' = 1$  from [4] or from Remark 5 in [11]; see also Remark 1 following Theorem 3.1 in [2].) If moreover  $X$  is isotropic, then  $\mathbb{E}|X| \leq (\mathbb{E}|X|^2)^{1/2} = \sqrt{n}$  and  $\sigma_2(X) = 1$ ; thus

$$(\mathbb{E}|X|^p)^{1/p} \leq C(\sqrt{n} + C'p).$$

From Markov's inequality for  $p = t\sqrt{n}$ , Theorem 2 implies Theorem 1 with  $c = (C' + 1)eC$ .

Let us recall the idea underlying the proof by Paouris. Let  $X \in \mathbb{R}^n$  be an isotropic log-concave random vector. Let  $p \sim \sqrt{n}$  be an integer (for example,  $p = \lfloor \sqrt{n} \rfloor$ ). Let  $Y = PX$  where  $P$  is an orthogonal projection of rank  $p$  and let  $G \in \text{Im}P$  be a standard Gaussian vector. By rotation invariance,  $\mathbb{E}|Y|^p \sim \mathbb{E}|\langle G/\sqrt{p}, Y \rangle|^p$ . If the linear forms  $\langle z, X \rangle$  with  $|z| = 1$  had a sub-Gaussian tail behaviour, the proof would be straightforward. But in general they only obey a sub-exponential tail behaviour. The first step of the proof consists of showing that there exists some  $z$  for which  $(\mathbb{E}|\langle z, Y \rangle|^p)^{1/p}$  is in fact small, compared to  $\mathbb{E}|Y|$ . The second step uses a concentration principle to show that  $(\mathbb{E}_X|\langle z, PX \rangle|^p)^{1/p}$  is essentially constant on the sphere, for a random orthogonal projection of rank  $p \sim \sqrt{n}$ , and thus comparable to the minimum. Thus for these *good* projections, one has a good estimate of  $(\mathbb{E}|Y|^p)^{1/p}$  and the result follows by averaging over  $P$ . Our proof follows the same scheme, at least for the first step, but whereas the proof of the first step in [14] is the most technical part, our argument is very simple. Then the estimate for  $\min_{|z|=1} \mathbb{E}|\langle z, Y \rangle|^p$  brings us to a minimax problem precisely in the form answered by Gordon's inequality ([9]).

Finally we would like to note that our proof can be generalized to the case of convex measures in the sense of [5, 6]. Of course the proof is longer and more technical. We provide the details in [1].

## 2 Proof of Theorem 2

First let us notice that it is enough to prove Theorem 2 for symmetric log-concave random vectors. Indeed, let  $X$  be a log-concave random vector and let  $X'$  be an independent copy. By Jensen's inequality we have for all  $p \geq 1$ ,

$$(\mathbb{E}|X|^p)^{1/p} \leq (\mathbb{E}|X - \mathbb{E}X|^p)^{1/p} + |\mathbb{E}X| \leq (\mathbb{E}|X - X'|^p)^{1/p} + \mathbb{E}|X|.$$

On the other hand  $\mathbb{E}|X - X'| \leq 2\mathbb{E}|X|$  and for  $p \geq 1$  one has  $\sigma_p(X - X') \leq 2\sigma_p(X)$ . Since  $X - X'$  is log-concave (see [8]) and symmetric, we obtain that the symmetric case proved with a constant  $C'$  implies the non-symmetric case with the constant  $C = 2C' + 1$ .

**Lemma 3.** Let  $Y \in \mathbb{R}^q$  be a random vector. Let  $\|\cdot\|$  be a norm on  $\mathbb{R}^q$ . Then for all  $p > 0$ ,

$$\min_{|z|=1} (\mathbb{E}|\langle z, Y \rangle|^p)^{1/p} \leq \frac{(\mathbb{E}\|Y\|^p)^{1/p}}{\mathbb{E}\|Y\|} \mathbb{E}|Y|.$$

**Proof:** Let  $r$  be the largest number such that  $r\|t\| \leq |t|$  for all  $t \in \mathbb{R}^q$ . Using duality, pick  $z \in \mathbb{R}^q$  such that  $|z| = 1$  and  $\|z\|_* \leq r$  (the dual norm of  $\|\cdot\|$ ). Then  $|\langle z, t \rangle| \leq r\|t\| \leq |t|$  for all  $t \in \mathbb{R}^q$ . Therefore,  $(\mathbb{E}|\langle z, Y \rangle|^p)^{1/p} \leq r(\mathbb{E}\|Y\|^p)^{1/p}$  for any  $p > 0$ , and the proof follows from  $r\mathbb{E}\|Y\| \leq \mathbb{E}|Y|$ .  $\square$

**Lemma 4.** Let  $Y$  be a full-dimensional symmetric log-concave  $\mathbb{R}^q$ -valued random vector. Then there exists a norm  $\|\cdot\|$  on  $\mathbb{R}^q$  such that

$$(\mathbb{E}\|Y\|^q)^{1/q} \leq 500 \mathbb{E}\|Y\|.$$

**Remark.** In fact the constant 500 can be significantly improved. To keep the presentation short and transparent we omit the details.

**Proof:** From Borell's characterization  $Y$  has an even log-concave density  $g_Y$ . Thus  $g_Y(0)$  is the maximum of  $g_Y$ . Define a symmetric convex set by

$$K = \{t \in \mathbb{R}^q : g_Y(t) \geq 25^{-q} g_Y(0)\}.$$

Since clearly  $K$  has a non-empty interior, it is the unit ball of a norm which we denote by  $\|\cdot\|$ . On one hand,  $1 \geq \mathbb{P}(Y \in K) = \int_K g_Y \geq 25^{-q} g_Y(0) \text{vol}(K)$ , thus

$$\mathbb{P}(\|Y\| \leq 1/50) = \int_{K/50} g_Y \leq g_Y(0) 50^{-q} \text{vol}(K) \leq 2^{-q} \leq 1/2.$$

Therefore  $\mathbb{E}\|Y\| \geq \mathbb{P}(\|Y\| > 1/50)/50 \geq 1/100$ . On the other hand, by the log-concavity of  $g_Y$ ,

$$\forall t \in \mathbb{R}^q \setminus K \quad g_{2Y}(t) = 2^{-q} g_Y(t/2) \geq 2^{-q} g_Y(t)^{1/2} g_Y(0)^{1/2} \geq (5/2)^q g_Y(t).$$

Therefore

$$\mathbb{E}\|Y\|^q \leq 1 + \mathbb{E}(\|Y\|^q \mathbf{1}_{Y \in \mathbb{R}^q \setminus K}) \leq 1 + (2/5)^q \mathbb{E}\|2Y\|^q = 1 + (4/5)^q \mathbb{E}\|Y\|^q.$$

We conclude that  $(\mathbb{E}\|Y\|^q)^{1/q} \leq 5$  and  $(\mathbb{E}\|Y\|^q)^{1/q} / \mathbb{E}\|Y\| \leq 500$ .  $\square$

**Lemma 5.** Let  $n, q \geq 1$  be integers and  $p \geq 1$ . Let  $X$  be an  $n$ -dimensional random vector,  $G$  be a standard Gaussian vector in  $\mathbb{R}^n$  and  $\Gamma$  be an  $n \times q$  standard Gaussian matrix. Then

$$(\mathbb{E}|X|^p)^{1/p} \leq \alpha_p^{-1} \left( \mathbb{E} \min_{|t|=1} \|\Gamma t\| + (\alpha_p + \sqrt{q}) \sigma_p(X) \right),$$

where  $\|z\| = (\mathbb{E}|\langle z, X \rangle|^p)^{1/p}$  and  $\alpha_p^p$  is the  $p$ -th moment of an  $N(0, 1)$  Gaussian random variable (so that  $\lim_{p \rightarrow \infty} (\alpha_p / \sqrt{p}) = 1/\sqrt{e}$ ).

**Proof:** By rotation invariance,  $\mathbb{E}|\langle G, X \rangle|^p = \alpha_p^p \mathbb{E}|X|^p$ . Notice that

$$\sigma^2 := \sup_{\|t\|_* \leq 1} \mathbb{E}|\langle G, t \rangle|^2 = \sup_{\|t\|_* \leq 1} |t|^2 = \sigma_p^2(X),$$

where  $\|\cdot\|_*$  denotes the norm on  $\mathbb{R}^n$  dual to the norm  $\|\cdot\|$ . Denote the median of  $\|G\|$  by  $M_G$ . The classical deviation inequality for a norm of a Gaussian vector ([7], [15], see also [12], Theorem 12.2) states

$$\forall s \geq 0 \quad \mathbb{P}(\| \|G\| - M_G | \geq s) \leq 2 \int_{s/\sigma}^{\infty} \exp(-t^2/2) \frac{dt}{\sqrt{2\pi}}$$

and since  $M_G \leq \mathbb{E}\|G\|$  ([10], see also [12], Lemma 12.2) this implies

$$(\mathbb{E}|X|^p)^{1/p} = \alpha_p^{-1} (\mathbb{E}\|G\|^p)^{1/p} \leq \alpha_p^{-1} (\mathbb{E}\|G\| + \alpha_p \sigma_p(X))$$

(cf. [13], Statement 3.1).

The Gordon minimax lower bound (see [9], Theorem 2.5) states that for any norm  $\|\cdot\|$

$$\mathbb{E}\|G\| \leq \mathbb{E} \min_{|t|=1} \|\Gamma t\| + (\mathbb{E}|H|) \max_{|z|=1} \|z\| \leq \mathbb{E} \min_{|t|=1} \|\Gamma t\| + \sqrt{q} \sigma_p(X),$$

where  $H$  is a standard Gaussian vector in  $\mathbb{R}^q$ . This concludes the proof.  $\square$

**Proof of Theorem 2:** Assume that  $X$  is log-concave symmetric. We use the notation of Lemma 5 with  $q$  the integer such that  $p \leq q < p + 1$ . We first condition on  $\Gamma$ . Let  $Y = \Gamma^* X$ . Note that  $Y$  is log-concave symmetric and that

$$\|\Gamma t\| = (\mathbb{E}_X |\langle \Gamma t, X \rangle|^p)^{1/p} = (\mathbb{E}_X |\langle t, \Gamma^* X \rangle|^p)^{1/p}.$$

If  $\Gamma^* X$  is supported by a hyperplane then  $\min_{|t|=1} (\mathbb{E}_X |\langle t, \Gamma^* X \rangle|^p)^{1/p} = 0$ . Otherwise Lemma 4 applies and combined with Lemma 3 gives that

$$\min_{|t|=1} \|\Gamma t\| \leq \min_{|t|=1} (\mathbb{E}_X |\langle t, \Gamma^* X \rangle|^p)^{1/p} \leq 500 \mathbb{E}_X |\Gamma^* X|.$$

By taking expectation over  $\Gamma$  we get

$$\mathbb{E} \min_{|t|=1} \|\Gamma t\| \leq 500 \mathbb{E} |\Gamma^* X| = 500 \mathbb{E} |H| \mathbb{E} |X| \leq 500 \sqrt{q} \mathbb{E} |X|,$$

where  $H \in \mathbb{R}^q$  is a standard Gaussian vector. Applying Lemma 5 we obtain

$$(\mathbb{E}|X|^p)^{1/p} \leq 500 \alpha_p^{-1} \sqrt{q} \mathbb{E}|X| + (1 + \alpha_p^{-1} \sqrt{q}) \sigma_p(X).$$

This implies the desired result, since  $q \leq p + 1$  and hence  $\alpha_p^{-1} \sqrt{q} \leq c$  for some numerical constant  $c$  (recall that  $\lim_{p \rightarrow \infty} (\alpha_p / \sqrt{p}) = 1/\sqrt{e}$ ).  $\square$

## References

- [1] R. Adamczak, O. Guédon, R. Latała, A. E. Litvak, K. Oleszkiewicz, A. Pajor and N. Tomczak-Jaegermann, *Moment estimates for convex measures*, preprint.
- [2] R. Adamczak, R. Latała, A. E. Litvak, A. Pajor and N. Tomczak-Jaegermann, *Tail estimates for norms of sums of log-concave random vectors*, preprint, <http://arxiv.org/abs/1107.4070>.
- [3] R. Adamczak, R. Latała, A. E. Litvak, A. Pajor and N. Tomczak-Jaegermann, *Geometry of log-concave Ensembles of random matrices and approximate reconstruction*, C.R. Math. Acad. Sci. Paris, **349** (2011), 783–786.
- [4] R. E. Barlow, A. W. Marshall and F. Proschan, *Properties of probability distributions with monotone hazard rate*, Ann. Math. Statist., **34** (1963), 375–389.
- [5] C. Borell, *Convex measures on locally convex spaces*, Ark. Math., **12** (1974), 239–252.
- [6] C. Borell, *Convex set functions in  $d$ -space*, Period. Math. Hungar., **6** (1975), 111–136.
- [7] C. Borell, *The Brunn-Minkowski inequality in Gauss space*, Invent. Math., **30** (1975), 207–216.
- [8] Ju. S. Davidovic, B. I. Korenbljum and B. I. Hacet, *A certain property of logarithmically concave functions*, Soviet Math. Dokl., **10** (1969), 447–480; translation from Dokl. Akad. Nauk SSSR **185** (1969), 1215–1218.
- [9] Y. Gordon, *Some inequalities for Gaussian processes and applications*, Israel J. Math., **50** (1985), 265–289.
- [10] S. Kwapień, *A remark on the median and the expectation of convex functions of Gaussian vectors*, Probability in Banach spaces, 9 (Sandjberg, 1993), 271–272, Progr. Probab., 35, Birkhuser Boston, Boston, MA, 1994.
- [11] S. Kwapień, R. Latała and K. Oleszkiewicz, *Comparison of moments of sums of independent random variables and differential inequalities*, J. Funct. Anal., **136** (1996), 258–268.
- [12] M. A. Lifshits, *Gaussian random functions*. Mathematics and its Applications; 322. Kluwer Academic Publishers, Dordrecht, 1995.
- [13] A. E. Litvak, V. D. Milman and G. Schechtman, *Averages of norms and quasi-norms*, Math. Ann., **312** (1998), 95–124.
- [14] G. Paouris, *Concentration of mass on convex bodies*, Geom. Funct. Anal., **16** (2006), 1021–1049.

- [15] V. N. Sudakov, B. S. Cirel'son, *Extremal properties of half-spaces for spherically invariant measures*, J. Sov. Math. **9** (1978), 9–18; translation from Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov (LOMI) **41** (1974), 14–24.

Radosław Adamczak,  
Institute of Mathematics,  
University of Warsaw,  
Banacha 2, 02-097 Warszawa, Poland  
e-mail: [radamcz@mimuw.edu.pl](mailto:radamcz@mimuw.edu.pl)

Rafał Łatała,  
Institute of Mathematics,  
University of Warsaw,  
Banacha 2, 02-097 Warszawa, Poland  
e-mail: [rlatala@mimuw.edu.pl](mailto:rlatala@mimuw.edu.pl)

Alexander E. Litvak,  
Dept. of Math. and Stat. Sciences,  
University of Alberta,  
Edmonton, Alberta, Canada, T6G 2G1.  
e-mail: [alexandr@math.ualberta.ca](mailto:alexandr@math.ualberta.ca)

Krzysztof Oleszkiewicz,  
Institute of Mathematics,  
University of Warsaw,  
Banacha 2, 02-097 Warszawa, Poland  
e-mail: [koles@mimuw.edu.pl](mailto:koles@mimuw.edu.pl)

Alain Pajor,  
Université Paris-Est  
Équipe d'Analyse et Mathématiques Appliquées,  
5, boulevard Descartes, Champs sur Marne,  
77454 Marne-la-Vallée, Cedex 2, France  
e-mail: [Alain.Pajor@univ-mlv.fr](mailto:Alain.Pajor@univ-mlv.fr)

Nicole Tomczak-Jaegermann,  
Dept. of Math. and Stat. Sciences,  
University of Alberta,  
Edmonton, Alberta, Canada, T6G 2G1.  
e-mail: [nicole.tomczak@ualberta.ca](mailto:nicole.tomczak@ualberta.ca)