ON THE PLANAR ALGEBRA OF OCNEANU'S ASYMPTOTIC INCLUSION

STEPHEN CURRAN[†]

ABSTRACT. In recent joint work with V. Jones and D. Shlyakhtenko, we have given a diagrammatic description of Popa's symmetric enveloping inclusion for planar algebra subfactors. In this paper we give a diagrammatic construction of the associated Jones tower, in the case that the planar algebra is finite-depth. We then use this construction to describe the planar algebra of the symmetric enveloping inclusion, which is known to be isomorphic to the planar algebra of Ocneanu's asymptotic inclusion by a result of Popa. As an application we give a planar algebraic computation of the (reduced) fusion algebra of the asymptotic inclusion, recovering some well-known results of Ocneanu and Evans-Kawahigashi.

INTRODUCTION

Let $M_0 \subset M_1$ be a finite-index inclusion of AFD II_1 factors, and let $M_0 \subset M_1 \subset M_2 \subset \cdots \subset M_\infty$ be the associated Jones tower [12]. The asymptotic inclusion is the subfactor $M_1 \vee (M'_1 \cap M_\infty) \subset M_\infty$, which has finite index if and only if $M_0 \subset M_1$ has finite depth. This construction was introduced by Ocneanu [18], who argued that the asymptotic inclusion could be viewed as the subfactor analogue of Drinfeld's quantum double construction. This connection has since been clarified by a number of authors, including Evans-Kawahigashi [4], Izumi [10, 11] and Müger [17]. A related construction for type III factors has been developed by Longo and Rehren [16].

In [19], Popa introduced the symmetric enveloping algebra associated to a finite index inclusion $M_0 \subset M_1$ of arbitrary II_1 factors. This is the (unique) II_1 factor $M_1 \boxtimes_{e_0} M_1^{op}$ which is generated by a copy of $M_1 \otimes M_1^{op}$ and a projection e_0 which is simultaneously the Jones projection for both $M_0 \subset M_1$ and $M_0^{op} \subset M_1^{op}$. In the case that $M_0 \subset M_1$ is a finite-depth inclusion of AFD II_1 factors, Popa proved that the symmetric enveloping inclusion $M_1 \otimes M_1^{op} \subset M_1 \boxtimes_{e_0} M_1^{op}$ is conjugate to Ocneanu's asymptotic inclusion. In the infinite-depth case, Popa [21] used this inclusion to analyze a number of important analytic properties of the original subfactor $M_0 \subset M_1$.

Planar algebras were introduced by Jones [13] in the late 90s. Any finite-index subfactor gives rise to a planar algebra as its *standard invariant*. By a fundamental result of Popa [20], any planar algebra satisfying suitable positivity conditions arises in this way. With inspiration from random matrix theory and Voiculescu's free probability, Guionnet, Jones and Shlyakhtenko [9] have recently given a diagrammatic construction of a tower of subfactors $M_0 \subset M_1 \subset \cdots$, starting from a planar algebra \mathcal{P} . In subsequent work [8], they proved that if \mathcal{P} is finite-depth then M_k is an interpolated free group factor $L\mathbb{F}_{r_k}$, where r_k is computed in terms of the index $\delta^2 = [M_1 : M_0]$ and the global index $I = [M_1 \boxtimes_{e_0} M_1^{op} : M_1 \otimes M_1^{op}]$. In joint work with Jones and Shlyakhtenko [1], we have given a diagrammatic construction of the symmetric enveloping algebra associated to these planar algebra subfactors (see also [22]). As an application of our construction, we computed a certain free entropy dimension type quantity which provides some intuition for the formula for r_k .

In this paper, we give a diagrammatic construction of the Jones tower of the symmetric enveloping inclusion for planar algebra subfactors, in the case that \mathcal{P} is finite-depth. We then use this construction to derive a complete description of the planar algebra of the asymptotic inclusion in terms of the original planar algebra \mathcal{P} . As an application, we compute the fusion rules for bimodules arising from the asymptotic inclusion, recovering some well-known results of Ocneanu and

 $[\]dagger:$ Research supported by an NSF postdoctoral fellowship, NSF grant DMS-0900776 and DARPA Award 0011-11-0001.

STEPHEN CURRAN[†]

Evans-Kawahigashi [4]. In particular, we show that the (reduced) fusion algebra of $M_1 \boxtimes_{e_0} M_1^{op}$ bimodules is described in terms of the affine category [6], which is closely related to Ocneanu's tube algebra [4] and Jones' annular category [15]. The relationship between the Drinfeld center of a fusion category and the annular/affine category is well-known in the TQFT community (see e.g. [5]). In the planar algebra setting, it was very recently shown by Das, Ghosh and Gupta [3] that the category of affine Hilbert representations of a finite-depth planar algebra \mathcal{P} is equivalent to the Drinfeld center of the fusion category associated to \mathcal{P} .

The paper is organized as follows. In Section 1 we briefly recall the constructions from [9], [1]. In Section 2 we study two canonical elements appearing in any finite-depth planar algebra, and establish a number of useful "skein" relations which these satisfy. In Section 3 we construct the Jones tower for the symmetric enveloping inclusion and compute the higher relative commutants. Section 4 contains our main result: a description of the planar algebra of the asymptotic inclusion. In Section 5 we use this description to compute the fusion rules for the asymptotic inclusion.

Acknowledgements. I am grateful to Dimitri Shlyakhtenko for suggesting this problem, and for many useful discussions while completing this project. I would also like to thank Dietmar Bisch and Vaughan Jones for several helpful conversations.

1. BACKGROUND AND PRELIMINARIES

In this section we briefly recall the constructions from [9], [1].

Planar algebra subfactors: Let $\mathcal{P} = (P_k)_{k\geq 0}$ be a subfactor planar algebra. For $n, k \geq 0$ let $P_{n,k}$ be a copy of P_{n+k} . Elements of $P_{n,k}$ will be represented by diagrams



where the thick lines to the left and right represent k strings, and the thick line at top represents 2n strings. We will typically suppress the marked point \star , and take the convention that it occurs at the top-left corner which is adjacent to an unshaded region.

Define a product $\wedge_k : P_{n,k} \times P_{m,k} \to P_{n+m,k}$ by

$$x \wedge_k y =$$

The involution $\dagger: P_{n,k} \to P_{n,k}$ is given by

$$\mathbf{x}^{\dagger} = \mathbf{x}^{\star} \mathbf{x}^{\star}$$

The Voiculescu trace $\tau_k : P_{n,k} \to \mathbb{C}$ is defined by

where $\sum TL$ denotes the sum over all loopless Temperley-Lieb diagram with 2*n* boundary points. Let $Gr_k(\mathcal{P}) = \bigoplus_{n\geq 0} P_{n,k}$, and observe that the formulas above give $Gr_k(\mathcal{P})$ the structure of a graded *-algebra with trace τ_k . The unit of $Gr_k(\mathcal{P})$ is the element of $P_{0,k}$ consisting of k parallel lines. Let \mathbf{e}_k denote the following element of $P_{0,k+2}$, $k \ge 0$:

$$\mathbf{e}_k =$$

Note that there are natural inclusions of $Gr_k(\mathcal{P})$ into $Gr_{k+1}(\mathcal{P})$ defined by



The main result of [9] is the following:

Theorem 1.1. For $k \geq 0$, the Voiculescu trace τ_k is a faithful tracial state on $Gr_k(\mathcal{P})$, and its GNS completion is a II_1 factor M_k as long as $\delta > 1$. The inclusions $Gr_k(\mathcal{P}) \subset Gr_{k+1}(\mathcal{P})$ extend to $M_k \subset M_{k+1}$, and (M_{k+1}, \mathbf{e}_k) is the Jones tower of $M_0 \subset M_1$. Moreover, the planar algebra of $M_0 \subset M_1$ is isomorphic to \mathcal{P} .

Popa's symmetric enveloping algebra: For integers k, s, t with s + t + 2k = n, let $V_k(s, t)$ be a copy of P_n . Elements of $V_k(s, t)$ will be represented by diagrams of the form



where there are 2s parallel strings at the top, 2t strings at the bottom and 2k at either side. As above, we will use the convention that the marked point occurs at the upper left corner, which is adjacent to a unshaded region.

Define a product $\wedge: V_k(s,t) \times V_k(s',t') \to V_k(s+s',t+t')$ by

$$x \wedge y =$$

The adjoint $\dagger: V_k(s,t) \to V_k(s,t)$ is defined by

$$x^{\dagger}$$
 = x^{*}

Define $\tau_k \boxtimes \tau_k : V_k(s, t) \to \mathbb{C}$ by

$$(\tau_k \boxtimes \tau_k)(x) = \begin{array}{c} \underbrace{\sum TL} \\ x \\ \underbrace{\sum TL} \end{array}$$

Define

$$Gr_k \boxtimes Gr_k^{op} = \bigoplus_{s,t \ge 0} V_k(s,t).$$

Note that there is a natural anti-automorphism $y \mapsto y^{op}$ of $Gr_k(\mathcal{P}) \boxtimes Gr_k(\mathcal{P})$, determined by

Observe also that $Gr_k(\mathcal{P}) \boxtimes Gr_k(\mathcal{P})$ contains a copy of $Gr_k(\mathcal{P}) \otimes Gr_k(\mathcal{P})^{op}$ as follows:

$$x \otimes y^{op} \mapsto \underbrace{\begin{bmatrix} \star & & \\ & &$$

Let $\mathbf{f}_k \in Gr_k \boxtimes Gr_k$ be the projection

$$\mathbf{f}_k = \delta^{-1} \cdot \mathbf{b}_k^{k-1}$$

Note that \mathbf{f}_k implements the Jones projections for both $M_{k-1} \subset M_k$ and $M_{k-1}^{op} \subset M_k^{op}$.

We have proved the following result in [1]:

Theorem 1.2. For $k \ge 0$, $\tau_k \boxtimes \tau_k$ is a faithful, tracial state on $Gr_k(\mathcal{P}) \boxtimes Gr_k(\mathcal{P})$, and its GNS completion is a II_1 factor $M_k \boxtimes M_k$ as long as $\delta > 1$. The inclusion $Gr_k(\mathcal{P}) \otimes Gr_k(\mathcal{P}) \subset Gr_k \boxtimes Gr_k$ extends to $M_k \otimes M_k \subset M_k \boxtimes M_k$. Moreover, for $k \ge 1$ we have that $M_k \otimes M_k^{op} \subset M_k \boxtimes M_k^{op}$ is isomorphic to Popa's symmetric enveloping inclusion $M_k \otimes M_k^{op} \subset M_k \boxtimes f_{k-1} M_k$.

2. Skein relations

In this section we introduce two canonical elements p and q appearing in any finite-depth planar algebra, and describe a number of diagrammatic relations which they satisfy.

Throughout the paper $\mathcal{P} = (P_k)_{k\geq 0}$ will be a subfactor planar algebra. Let Γ denote the principal graph of \mathcal{P} (see e.g. [7]), and let Γ_+ denote the collection of even vertices. Associated to each $v \in \Gamma_+$ is an irreducible $M_0 - M_0$ bimodule X_v . We let * denote the distinguished vertex, corresponding to the bimodule $X_* = L^2(M_0)$.

We will assume that Γ_+ is finite (i.e. \mathcal{P} is finite-depth), and throughout the paper we fix k such that $d(*, v) \leq 2k$ for all $v \in \Gamma_+$. It follows that we have

$$_{M_0}L^2(M_k)_{M_0}\simeq \bigoplus_{v\in\Gamma_+} X_v\otimes\mathcal{H}_v,$$

where \mathcal{H}_v are auxiliary finite-dimensional Hilbert spaces, whose dimensions we denote by n_v . It follows that

$${}_{M_0 \otimes M_0^{op}} L^2(M_k \otimes M_k^{op})_{M_0 \otimes M_0^{op}} \simeq \bigoplus_{v, w \in \Gamma_+} X_v \otimes \overline{X_w} \otimes (\mathcal{H}_v \otimes \overline{\mathcal{H}_w}),$$

where $\overline{X_w}$ is the contragredient bimodule and $\overline{\mathcal{H}_w}$ is the conjugate Hilbert space. Let $1_v \in \mathcal{H}_v \otimes \overline{\mathcal{H}_v}$ be the unit under the natural identification with $\operatorname{Hom}_{\mathbb{C}}(\mathcal{H}_v)$, and define p_v to be the projection from $L^2(M_k \otimes M_k^{op})$ onto $X_v \otimes \overline{X_v} \otimes 1_v$. Then define

$$p = \sum_{v \in \Gamma_+} p_v.$$

Recall that P_{2k} can be identified with $\operatorname{Hom}_{M_0,M_0}(L^2(M_k))$, in particular the central projections of P_{2k} are indexed by $v \in \Gamma_+$. By an abuse of notation we will use v to denote the central projection corresponding to the vertex v. Let $(\mu_v)_{v\in\Gamma}$ be the Perron-Frobenius eigenvector of the adjacency matrix for Γ , normalized by $\mu(*) = 1$. Then the trace of a minimal projection in the central component of P_{2k} corresponding to $v \in \Gamma_+$ is $\operatorname{tr}_{P_{2k}}(v) = \delta^{-2k} \mu_v$. Note that if $\{e_{ij}(v) : 1 \leq i, j \leq n_v\}$ are matrix units for the central component of P_{2k} corresponding to $v \in \Gamma_+$, then we have

$$p_v = \frac{1}{n_v} \sum_{1 \le i,j \le n_v} e_{ij}(v) \otimes e_{ji}(v)^{op}.$$

Graphically, we will represent p_v using a Sweedler type convention as follows:

$$p_v = \begin{array}{c} & & \\$$

The projection p_v satisfies the following key skein relation:

Lemma 2.1. Let $x, y \in P_{2k}$ and $v \in \Gamma_+$, then we have

Proof. The left hand side is equal to

$$\frac{\delta^{4k}}{n_v} \sum_{1 \le i,j \le n_v} \operatorname{tr}(e_{ij}(v)x) \cdot \operatorname{tr}(ye_{ji}(v)) = \frac{\delta^{4k}}{n_v} \cdot \delta^{-2k} \mu_v \cdot \operatorname{tr}(xy \cdot v),$$

which is equal to the right hand side.


We will now derive a number of 'skein' relations for the projection p. First we recall the following well known relation:

Lemma 2.2. For $x, y \in P_k$ we have


$$\begin{array}{c} x \\ \hline v \\ y \\ \end{array} = \delta_{v,*} \quad \begin{array}{c} x \\ \hline y \\ \end{array}$$

Proposition 2.3. *p* satisfies the following skein relations:


(1) Trace: $tr_{P_{2k}}(p) = \delta^{-4k}I$, i.e.




(2) Rotational invariance:



(3) Capping:



(4) Multiplication: We have



Proof. For (1), we have


$$\operatorname{tr}_{P_{2k}}(p) = \sum_{v \in \Gamma_+} n_v^{-1} \sum_{1 \le i,j \le n_v} \operatorname{tr}_{P_{2k}(v)}(e_{ij}(v))^2 = \sum_{v \in \Gamma_+} \delta^{-4k} \mu_v^2 = \delta^{-4k} I.$$

(2) follows easily from Lemma 2.1, and the fact that



where \overline{v} is the vertex corresponding to the conjugate bimodule $\overline{X_v}$.

By Lemma 2.1 we have




for any $x, y \in P_k$. (3) then follows from Lemma 2.2.

For (4), the left hand side is equal to

$$\sum_{v \in \Gamma_{+}} \frac{1}{n_{v}} \sum_{1 \le i, j \le n_{v}} e_{ij}(v) e_{ji}(v) = \sum_{v \in \Gamma_{+}} v = 1.$$

We now introduce an element $q \in P_{3k}$, which will be central to our constructions.

Proposition 2.4. Fix $v, w, z \in \Gamma_+$. Then there is a unique element $q_{v,w,z} \in P_{3k} \otimes P_{3k}^{op}$, which we represent in Sweedler notation as $q_{v,w,z} = q_{v,w,z}^{(1)} \otimes q_{v,w,z}^{(2)}$, with the property that for any $x, y \in P_{3k}$ we have




Proof. With the obvious unital embedding of $P_{3k} \otimes P_{3k}^{op}$ into P_{6k} , we have

Define

$$q = \sum_{v,w,z \in \Gamma_+} q_{v,w,z}.$$

Proposition 2.5. *q* satisfies the following skein relations:

(1) Compatibility with p:




(2) Rotation:


(3) Capping: If $x, y \in P_k$ then

$$\begin{array}{c} \begin{pmatrix} x \\ q^{(1)} \\ \hline q^{(2)} \\ \hline y \\ \hline y \\ \hline \end{array} = I^{-1/2} n_*^{-1/2} \cdot \begin{array}{c} & & \\ & &$$

(4) Double capping:




(5) Connecting:



Proof. (1) and (2) follow easily from Proposition 2.4. (3) follows from Proposition 2.4 and Lemma 2.1. (4) follows from (2) and (3), and then (5) clearly follows from (4). \Box

The finite-depth assumption has not been essential so far, but the following consequence will be crucial in the remainder of the paper.

Proposition 2.6. Let $m \ge k$, then P_{2m} is spanned by elements of the form




for $x, y \in P_{m+k}$.

Proof. Since the depth of \mathcal{P} is less than or equal to 2k, we have that P_{2k+1} is equal to the basic construction of $P_{2k-1} \subset P_{2k}$. So P_{2k+1} is generated by P_{2k} and the Jones projection e_{2k-1} . Iterating, we see that P_{2m} is generated by P_{2k} and $e_{2k-1}, \ldots, e_{2m-2}$, from which the result easily follows. \Box

Proposition 2.7. Fix $w, z \in \Gamma_+$, then we have


$$\sum_{v\in\Gamma_+} \left(\frac{\mu_v}{n_v}\right)^{1/2} \cdot \left(\begin{array}{c} \star & q_{v,w,z}^{(1)} \\ q_{v,w,z}^{(2)} \\ q_{v,w,z}^{(2)} \\ \star \end{array}\right) = I^{-1/2} \left(\frac{\mu_w \mu_z}{n_w n_z}\right)^{1/2} \cdot (w) \star (z) \star$$

Proof. By Proposition 2.6, it suffices to show that



for any $x, y \in P_{3k}$.

Now by Lemma 2.1, the left hand side of the above equation is equal to




By Proposition 2.4 this is equal to

$$\sum_{v\in\Gamma_+} \left(\frac{\mu_v}{n_v}\right)^{1/2} \cdot \frac{n_v}{\mu_v} \cdot I^{-1/2} \cdot \left(\frac{\mu_v \mu_w \mu_z}{n_v n_w n_z}\right)^{1/2} \xrightarrow{*}_{p_v^{(1)}} (w) \in \mathbb{R}$$

The result then follows from (4) of Proposition 2.3.

Corollary 2.8. Fix $v_1, v_2, z_1, z_2 \in \Gamma_+$. Then if $x, y \in P_{4k}$ we have




Proof. By Proposition 2.4, the left hand side of the equation in the statement of the proposition is equal to


$$\sum_{z \in \Gamma_{+}} I^{-1/2} \left(\frac{\mu_{v_{1}} \mu_{w_{1}} \mu_{z}}{n_{v_{1}} n_{w_{1}} n_{z}} \right)^{1/2} \cdot \underbrace{ \begin{array}{c} x \\ (v_{1}) \\ (v_{1}) \\ (w_{1}) \\$$

and the result then follows from Proposition 2.7.

Corollary 2.9. Let $q_1 = q_1^{(1)} \otimes q_1^{(2)}$ and $q_2 = q_2^{(1)} \otimes q_2^{(2)}$ be copies of q. Then




Proof. It follows from Corollary 2.8 and Proposition 2.7 that




The result then follows from (2) of Proposition 2.5.

Corollary 2.10. Let q_1, q_2 be copies of q. Then we have



Proof. By Corollary 2.9 and (3) of Proposition 2.5, we have



The result follows by applying (4) of Proposition 2.5 and (1) of Proposition 2.3.


3. The Jones tower

In this section we compute the Jones tower for the inclusion $M_0 \otimes M_0^{op} \subset M_0 \boxtimes M_0^{op}$. Our construction is easily modified to obtain the Jones tower for the symmetric enveloping inclusion $M_1 \otimes M_1^{op} \subset M_1 \boxtimes M_1^{op}$. However, we prefer to work with $M_0 \boxtimes M_0^{op}$ as it simplifies the diagrams. As discussed in Remark 4.7 in the following section, this is sufficient to compute the planar algebra of the asymptotic inclusion.


As above, we will assume that \mathcal{P} is a finite-depth planar algebra and that $depth(\mathcal{P}) \leq 2k$.

Fix $n \ge 0$. Recall from Section 1 that we have a graphical representation of $P_{s,(2n+1)k} \otimes P_{t,(2n+1)k}$ as the span of elements of the form


for $x \in P_{s,(2n+1)k}$ and $y \in P_{t,(2n+1)k}$. For $s,t \ge 0$ let $\mathcal{V}_{2n+1}(s,t)$ be the subspace of $P_{s,(2n+1)k} \otimes P_{t,(2n+1)k}$ spanned by



for $x \in P_{s,(2n+1)k}$, $y \in P_{t,(2n+1)k}$, where $p_n = p_n^{(1)} \otimes p_n^{(2)}$ is defined recursively as follows: p_0 is the empty diagram, and for $n \ge 1$ we have



Note that $\mathcal{V}_{2n+1}(s,t)$ is closed under the involution on $P_{s,(2n+1)k} \otimes P_{t,(2n+1)k}$. Fix $t_1, t_2, s_1, s_2 \ge 0$ and define




for $x_i \otimes y_i \in P_{s_i,(2n+1)k} \otimes P_{t_i,(2n+1)k}$. Extend this map linearly and then restrict to the subspace $\mathcal{V}_{2n+1}(s_1, t_1) \otimes \mathcal{V}_{2n+1}(s_2, t_2)$ to obtain a "twisted" multiplication

$$\star_q: \mathcal{V}_{2n+1}(s_1, t_1) \otimes \mathcal{V}_{2n+1}(s_2, t_2) \to \mathcal{V}_{2n+1}(s_1 + s_2, t_1 + t_2).$$

We note that in the rest of the paper we will define linear maps on $\mathcal{V}_{2n+1}(s,t)$ by simply describing their action on tensors $x \otimes y$, then implicitly applying this extension/restriction procedure.

Let $\varphi_{2n+1}: \mathcal{V}_{2n+1}(s,t) \to \mathbb{C}$ be the linear functional determined by




for $x \in P_{s,(2n+1)k}$ and $y \in P_{t,(2n+1)k}$, where the second equality follows from Proposition 2.5 (4) and Proposition 2.3 (1).


Define

$$\mathcal{A}_{2n+1} = \bigoplus_{s,t \ge 0} \mathcal{V}_{2n+1}(s,t).$$

Finally set



Now define $\Psi_n: \mathcal{V}_{2n+1}(s,t) \to V_{2nk}(s,t)$ to be the linear map determined by



Proposition 3.1. Ψ_n is an isomorphism of $(\mathcal{A}_{2n+1}, \star_q)$ onto $p_n(Gr_{2nk} \boxtimes Gr_{2nk}^{op})p_n$. We have $\Psi_n(1_{\mathcal{A}_{2n+1}}) = p_n$, and

$$\delta^{4nk} I^{-n}(\tau_{2nk} \boxtimes \tau_{2nk}) \circ \Psi_n = \varphi_{2n+1}.$$

Proof. The fact that Ψ_0 is injective was proved in [1] (note that our Ψ_0 was denoted by Ψ_k there, and had a different scaling factor). The argument given there extends easily to Ψ_n for n > 0. It is clear that Ψ_n respects the involutions on \mathcal{A}_{2n+1} and $\operatorname{Gr}_{2nk} \boxtimes \operatorname{Gr}_{2nk}^{op}$. That $\Psi_n(1_{\mathcal{A}_{2n+1}}) = p_n$ and that $\delta^{4nk} I^{-n}(\tau_{2nk} \boxtimes \tau_{2nk}) \circ \Psi_n = \varphi_{2n+1}$ are clear from the definitions and Lemma 2.2. Surjectivity follows from Lemma 2.6.

It remains only to show that Ψ_n is a homomorphism. We have

$$\Psi_{n}((x_{1} \otimes y_{1}) \star_{q} (x_{2} \otimes y_{2})) = I^{-1/2} \sum_{v \in \Gamma_{+}} \left(\frac{\mu_{v}}{n_{v}}\right)^{1/2} \qquad \underbrace{v}_{1} \underbrace{q^{(1)}}_{y_{1}} \underbrace{v}_{2nk} \underbrace{q^{(2)}}_{y_{2}} \underbrace{y_{1}}_{y_{2}} \underbrace{q^{(2)}}_{y_{2}} \underbrace{y_{2}}_{y_{2}} \underbrace{y_{1}}_{y_{2}} \underbrace{q^{(2)}}_{y_{2}} \underbrace{y_{2}}_{y_{2}} \underbrace{y_{2}} \underbrace{y_{$$

which by Proposition 2.7 is equal to

This shows that Ψ_n is a homomorphism, which completes the proof.

The following theorem is now immediate.

Theorem 3.2. For $n \ge 0$, $(\mathcal{A}_{2n+1}, \star_q)$ is a unital, associative *-algebra. Moreover, φ_{2n+1} is a faithful, tracial state on \mathcal{A}_{2n+1} , and Ψ_n extends to an isomorphism of the GNS completion \mathcal{M}_{2n+1} onto $p_n(\mathcal{M}_{2nk} \boxtimes \mathcal{M}_{2nk}^{op})p_n$. In particular, \mathcal{M}_1 is isomorphic to $\mathcal{M}_0 \boxtimes \mathcal{M}_0^{op}$.

Now for $n \ge 0$, define $\mathcal{A}_{2n} = p_n(Gr_{2nk} \otimes Gr_{2nk}^{op})p_n$ and let φ_{2n} be the renormalized trace

$$\varphi_{2n} = \delta^{4nk} I^{-n} \cdot (\tau_{2nk} \otimes \tau_{2nk})|_{\mathcal{A}_{2n}}$$


Let \mathcal{M}_{2n} denote the GNS completion of \mathcal{A}_{2n} , which is naturally isomorphic to $p_n(M_{2nk} \otimes M_{2nk}^{op})p_n$. Define inclusions $i_{2n} : \mathcal{A}_{2n} \hookrightarrow \mathcal{A}_{2n+1}$ by

$$i_{2n}(x \otimes y) = \underbrace{\begin{array}{c} & & \\$$

Proposition 3.3. For $n \ge 0$, i_{2n} extends to a unital, trace-preserving inclusion of $(\mathcal{M}_{2n}, \varphi_{2n})$ into $(\mathcal{M}_{2n+1}, \varphi_{2n+1})$. Moreover, the following diagram commutes:

Proof. It is clear that $\Psi_n \circ i_{2n}$ is the inclusion of $p_n(Gr_{2nk} \otimes Gr_{2nk}^{op})p_n$ into $p_n(Gr_{2nk} \boxtimes Gr_{2nk}^{op})p_n$, from which the result follows.

We now compute the φ_{2n+1} -preserving conditional expectation from \mathcal{M}_{2n+1} onto \mathcal{M}_{2n} . Define $E_{\mathcal{M}_{2n}}: \mathcal{V}_{2n+1}(s,t) \to P_{s,2nk} \otimes P_{t,2nk}$ by



Proposition 3.4. $E_{\mathcal{M}_{2n}}$ extends to the unique φ_{2n+1} -preserving conditional expectation $\mathcal{M}_{2n+1} \rightarrow \mathcal{M}_{2n}$.

Proof. We have

$$\langle i_{2n}(a \otimes b), x \otimes y \rangle = I^{-\frac{2n+1}{2}} n_*^{-1/2} \cdot \underbrace{ \left\{ \begin{array}{c} \sum TL \\ a^* \\ \end{array} \right\}_{x}}^{\left\{ \sum TL \\ \end{array}} = \langle a \otimes b, E_{\mathcal{M}_{2n}}(x \otimes y) \rangle$$

from which the result follows.

Now define inclusions $i_{2n+1} : \mathcal{A}_{2n+1} \to \mathcal{A}_{2n+2}$ by

$$i_{2n+1}(x \otimes y) = \begin{array}{c} & & \\$$

Proposition 3.5. i_{2n+1} extends to a unital, trace preserving embedding of \mathcal{M}_{2n+1} into \mathcal{M}_{2n+2} . Moreover, the following diagram commutes:

where the arrow on the bottom row is the obvious inclusion.

Proof. This follows from combining (3) and (4) of Proposition 2.5 and (1) of Proposition 2.3. \Box

Now define $E_{\mathcal{M}_{2n+1}}: \mathcal{A}_{2n+2} \to \mathcal{A}_{2n+1}$ by

$$E_{\mathcal{M}_{2n+1}}(x \otimes y) = \overbrace{\begin{array}{c} \\ q^{(1)} \\ q^{(2)} \\ y \end{array}}^{x}$$


Proposition 3.6. $E_{\mathcal{M}_{2n+1}}$ extends to the unique φ_{2n+2} -preserving conditional expectation of \mathcal{M}_{2n+2} onto \mathcal{M}_{2n+1} .

Proof. Draw the picture and apply Proposition 2.5 (5) to see that

$$\langle a \otimes b, E_{\mathcal{M}_{2n+1}}(x \otimes y) \rangle = \langle i_{2n+1}(a \otimes b), x \otimes y \rangle,$$


from which the result follows.

Now define $\mathbf{e}_{2n} \in \mathcal{A}_{2n+2}$ by



where the second equality follows from Proposition 2.5 (4).

Also define $\mathbf{e}_{2n+1} \in \mathcal{A}_{2n+3}$ by



Theorem 3.7. $(\mathcal{M}_n, I^{-1/2}\mathbf{e}_n)$ is the Jones tower for the subfactor $\mathcal{M}_0 \subset \mathcal{M}_1$.

Proof. It follows from Propositions 3.4 and 3.6 that \mathbf{e}_n is the Jones projection for $\mathcal{M}_n \subset \mathcal{M}_{n+1}$. So it remains to show that \mathcal{M}_{n+2} is the basic construction for $\mathcal{M}_n \subset \mathcal{M}_{n+1}$ for all $n \ge 0$.

We first prove that \mathcal{M}_{2n+2} is the basic construction of $\mathcal{M}_{2n} \subset \mathcal{M}_{2n+1}$. Indeed, by a standard result in subfactor theory we have that the subalgebra $\langle \mathcal{M}_{2n+1}, \mathbf{e}_{2n} \rangle$ of \mathcal{M}_{2n+2} which is generated by \mathcal{M}_{2n+1} and \mathbf{e}_{2n} is isomorphic to the basic construction of $\mathcal{M}_{2n} \subset \mathcal{M}_{2n+1}$. Note that by Proposition 3.3 we have

$$[\mathcal{M}_{2n+1}:\mathcal{M}_{2n}] = [p_n(M_{2nk} \boxtimes M_{2nk}^{op})p_n: p_n(M_{2nk} \otimes M_{2nk}^{op})p_n]$$
$$= [M_{2nk} \boxtimes M_{2nk}: M_{2nk} \otimes M_{2nk}^{op}] = I.$$

It follows that $[\langle \mathcal{M}_{2n+1}, \mathbf{e}_{2n} \rangle : \mathcal{M}_{2n}] = I^2$. By Proposition 3.5 we have

$$\begin{aligned} [\mathcal{M}_{2n+2}:\mathcal{M}_{2n}] &= [p_{n+1}(M_{2(n+1)k} \otimes M_{2(n+1)k}^{op})p_{n+1}:p_n(M_{2nk} \otimes M_{2nk}^{op})p_n] \\ &= \left(\frac{(\tau_{2(n+1)k} \otimes \tau_{2(n+1)k})(p_{n+1})}{(\tau_{2nk} \otimes \tau_{2nk})(p_n)}\right)^2 \cdot [M_{2(n+1)k} \otimes M_{2(n+1)k}:M_{2nk} \otimes M_{2nk}] \\ &= \delta^{-8k} I^2 \cdot \delta^{8k} \\ &= [\langle \mathcal{M}_{2n+1}, \mathbf{e}_{2n} \rangle: \mathcal{M}_{2n}], \end{aligned}$$

which implies that $\mathcal{M}_{2n+2} = \langle \mathcal{M}_{2n+1}, \mathbf{e}_{2n} \rangle$ is the Jones basic construction as claimed.

It now follows that \mathcal{M}_{2n+3} is the basic construction for $\mathcal{M}_{2n+1} \subset \mathcal{M}_{2n+2}$. Indeed, we have $\langle M_{2n+2}, \mathbf{e}_{2n+1} \rangle \subset \mathcal{M}_{2n+3}$. But since we have shown that \mathcal{M}_{2n+4} is the basic construction of $\mathcal{M}_{2n+2} \subset \mathcal{M}_{2n+3}$, we have


$$[\mathcal{M}_{2n+3}:\mathcal{M}_{2n+2}] = [\mathcal{M}_{2n+4}:\mathcal{M}_{2n+2}]^{1/2} = I = [\langle \mathcal{M}_{2n+2}, \mathbf{e}_{2n+1} \rangle : \mathcal{M}_{2n+2}],$$

so that $\mathcal{M}_{2n+3} = \langle \mathcal{M}_{2n+2}, \mathbf{e}_{2n+1} \rangle$ as desired.


It follows that the planar algebra $\mathcal{P}(\mathcal{M}_0 \subset \mathcal{M}_1)$ is isomorphic to $\mathcal{P}(\mathcal{M}_0 \otimes \mathcal{M}_0^{op} \subset \mathcal{M}_0 \boxtimes \mathcal{M}_0^{op})$. We will now compute the higher relative commutants $\mathcal{M}'_i \cap \mathcal{M}_n$ for i = 0, 1, we compute the action of planar tangles in the next section.

Proposition 3.8. The higher relative commutants are as follows:


(1) For $n \ge 0$, $\mathcal{M}'_0 \cap \mathcal{M}_{2n}$ is spanned by elements of the form:




(2) For $n \ge 0$, $\mathcal{M}'_0 \cap \mathcal{M}_{2n+1}$ is spanned by elements of the form:



(3) For $n \ge 1$, $\mathcal{M}'_1 \cap \mathcal{M}_{2n}$ is spanned by elements of the form:




(4) For $n \ge 1$, $\mathcal{M}'_1 \cap \mathcal{M}_{2n+1}$ is spanned by elements of the form:




Proof. Since $\mathcal{M}_0 \subset \mathcal{M}_{2n}$ is identified with the inclusion $M_0 \otimes M_0^{op} \subset p_n(M_{2nk} \otimes M_{2nk}^{op})p_n$, we have $\mathcal{M}'_0 \cap \mathcal{M}_{2n} = p_n(P_{2nk} \otimes P_{2nk}^{op})p_n$. This proves (1). For (3), suppose that $x \in \mathcal{M}'_1 \cap \mathcal{M}_{2n}$. By (1), $x \in p_n(P_{2nk} \otimes P_{2nk}^{op})p_n$. Since x commutes with

 \mathcal{M}_1 , we have



where we are using Sweedler notation $x = x^{(1)} \otimes x^{(2)}$. It follows that




where we have applied Corollary 2.10 for the first equality. Applying Corollary 2.9 shows that x is of the desired form, which proves (3).


Finally, since $\mathcal{M}'_i \cap \mathcal{M}_{2n+1} = E_{\mathcal{M}_{2n+1}}(\mathcal{M}'_i \cap \mathcal{M}_{2n+2})$ for i = 0, 1, (2) and (4) then follow from (1) and (3) by Proposition 3.6.

As a corollary we can compute the conditional expectation $E_{\mathcal{M}'_1}(x)$ for x in $\mathcal{M}'_0 \cap \mathcal{M}_n$. The proof is a straightforward diagrammatic argument and is left to the reader.

Corollary 3.9. If $x = x^{(1)} \otimes x^{(2)} \in \mathcal{M}'_0 \cap \mathcal{M}_{2n}$, then



Likewise, if $x = x^{(1)} \otimes x^{(2)} \in \mathcal{M}'_0 \cap \mathcal{M}_{2n+1}$ then



4. The planar algebra

In this section we construct the planar algebra $Sym(\mathcal{P})$ of the subfactor $\mathcal{M}_0 \subset \mathcal{M}_1$. By Theorem 3.2, this is isomorphic to the planar algebra of $M_0 \otimes M_0^{op} \subset M_0 \boxtimes M_0^{op}$. We explain how to recover the planar algebra of the asymptotic inclusion in Remark 4.7 at the end of this section.

It will be convenient to express this in terms of the planar algebra of the inclusion $M_0 \otimes M_0^{op} \subset M_k \otimes M_k^{op}$, which we will now describe.

First we construct the planar algebra of $M_0^{op} \subset M_1^{op}$. For $n \ge 1$ let R_n be the following tangle:



where there are n strings on the top and bottom of the input box, and the region adjacent to the left side of the outer box is unshaded. Let R_n^* be the same tangle, but with the region adjacent to

the left side of the inner box unshaded. Define P_n^{rev} to be $P_{n,+}$ if n is even and $P_{n,-}$ if n is odd. Note that Z_{R_n} maps P_n^{rev} onto P_n . Let T be a tangle with 2n marked points on the boundary of its outer disc, and with input discs D_1, \ldots, D_m such that D_i has $2n_i$ marked points on its boundary. We then define


$$Z_T^{rev}: P_{n_1}^{rev} \otimes \cdots \otimes P_{n_m}^{rev} \to P_n^{rev}$$

by

$$Z_T^{rev}(x_1 \otimes \cdots \otimes x_m) = Z_{R_n^*}(Z_T(Z_{R_{n_1}}(x_1) \otimes \cdots \otimes Z_{R_{n_m}}(x_m))).$$

It is not hard to see that $\mathcal{P}^{rev} = (P_n^{rev})_{n\geq 0}$ is the planar algebra for $M_0^{op} \subset M_1^{op}$, see [13]. Now the planar algebra of $M_0 \otimes M_0^{op} \subset M_1 \otimes M_1^{op}$ is the tensor product $\mathcal{P} \otimes \mathcal{P}^{rev}$. We refer to [13] for details, but note that in particular we have $(\mathcal{P} \otimes \mathcal{P}^{rev})_n = P_n \otimes P_n^{rev}$. Finally, the planar algebra of $M_0 \otimes M_0^{op} \subset M_k \otimes M_k^{op}$ is the *k*-cabling $\mathcal{C}_k(\mathcal{P} \otimes \mathcal{P}^{rev})$ of the planar algebra of $M_0 \otimes M_0^{op} \subset M_1 \otimes M_1^{op}$. Again we will refer to [13] for the definition, but note that $(\mathcal{C}_k(\mathcal{P} \otimes \mathcal{P}^{rev}))_n = P_{nk} \otimes P_{nk}^{rev}$. In particular, the elements *p* and *q* of Section 2 are now elements of the 2 and 2 bey grave of the planar algebra. the 2- and 3-box spaces of this planar algebra, respectively. Let us now interpret some of the skein relations from that section in terms of this planar algebra.

Proposition 4.1. The following skein relations hold in $C_k(\mathcal{P} \otimes \mathcal{P}^{rev})$:



(5) For any $x \in (\mathcal{C}_k(\mathcal{P} \otimes \mathcal{P}^{rev}))_2$ we have

$$\begin{array}{c} \star \\ \hline q = q \\ \star \end{array} = \begin{array}{c} p \\ p \\ \star \end{array}$$


Because of (1) we will usually not label the marked interval for p and q, but assume that it occurs at one of the corners. Note that (5) follows from combining (4) and (5) of Proposition 2.5. The following important relation follows from (3) and (4) above.

Lemma 4.2. For $n \geq 3$ we have




where there are n q's appearing on the left hand side of the equation, and n-2 on the right hand side.

Proof. By applying relation (4) to the second and third q from left, we see that the left hand side is equal to




Now if n = 3 then the statement follows from relation (3) above, otherwise it follows from induction on n.

Define $Sym(\mathcal{P})_n \subset \mathcal{C}_k(\mathcal{P} \otimes \mathcal{P}^{rev})_n$ to be the range of the partially labelled tangle



By Proposition 3.8, we have an identification of $Sym(\mathcal{P})_n$ with $\mathcal{M}'_0 \cap \mathcal{M}_n$ (as vector spaces).

For $n \ge 1$ define partially labelled tangles F_n and C_n by



where the number of q's in each tangle is n. By relation (2) above, the linear maps associated to F_n and C_n restrict to maps $Z_{F_n} : Sym(\mathcal{P})_n \to Sym(\mathcal{P})_{2n}$ and $Z_{C_n} : Sym(\mathcal{P})_{2n} \to Sym(\mathcal{P})_n$. Note that $Z_{C_n} \circ Z_{F_n}$ restricts to the identity on $Sym(\mathcal{P})_n$ by (3) of Proposition 4.1.

Now let T be a planar tangle with 2n marked points on the boundary of its outer disc, and with input discs D_1, \ldots, D_m such that D_i has $2n_i$ marked points on its boundary. We define a "spin factor" (cf. [14]) $\sigma(T)$ as follows. First isotope T so that the output and input rectangles each have as many marked points on the top as on the bottom (with no marked points on the sides), and such that the left side is the distinguished interval. $\sigma(T)$ is then defined to be the product over all local extrema of all the strings of $I^{\pm 1/4}$, where the sign is taken to be positive if the convex region adjacent to the extreme point is unshaded, and negative if it is shaded. If n = 0, we also multiply by I^{-1} if the region adjacent to the boundary is shaded.

Let T be the tangle obtained by doubling the strings of T. We then define a linear map

$$Z_T^{Sym}: Sym(\mathcal{P})_{n_1} \otimes \cdots \otimes Sym(\mathcal{P})_{n_m} \to Sym(\mathcal{P})_n$$

by

$$Z_T^{Sym}(x_1 \otimes \cdots \otimes x_m) = \sigma(T) \cdot Z_{C_n}(Z_{\widetilde{T}}(Z_{F_{n_1}}(x_1) \otimes \cdots \otimes Z_{F_{n_m}}(x_m))).$$


In other words, Z_T^{Sym} is equal to $\sigma(T) \cdot Z_{f(T)}$, where f(T) is the partially labelled tangle obtained from T by doubling the number of strings and composing with the tangles C_n and F_{n_i} in the appropriate manner.

Let us now compute this action explicitly for several important classes of tangles.


Example 4.3. In $Sym(\mathcal{P})$, we have the following:

(1) Identity: If n is even we have


Likewise if n is odd we have



(2) Multiplication: Let $x, y \in Sym(\mathcal{P})_n$. If n is even we have




Likewise, if n is odd we have




Note that we have applied Lemma 4.2 here.

(3) Inclusions: Let $x \in Sym(\mathcal{P})_n$. If n is even we have



If n is odd we have




Again we have applied Lemma 4.2 in this case.

(4) Jones projections: If n is even we have


$$\overset{\bullet}{\overset{\bullet}}_{\overset{\bullet}{\overset{\bullet}}} = I^{-1/2} \cdot \underbrace{\overset{\bullet}{\overset{\bullet}}_{\overset{\bullet}{\overset{\bullet}}}}_{\overset{\bullet}{\overset{\bullet}}} = I^{1/2} \cdot \mathbf{e}_{n}$$

Likewise, if n is odd we have




where we have applied Lemma 4.2.

(5) Conditional expectation onto \mathcal{M}'_1 : Let $x \in Sym(\mathcal{P})_n$. If n is even we have



where we have applied Lemma 4.2. Likewise, if n is odd we have



In both cases we are applying Corollary 3.9.

(6) Trace: Let $x \in Sym(\mathcal{P})_n$. If n is even we have

$$\overset{\star}{\underbrace{x}}_{\widehat{n}} \mapsto \underbrace{\begin{pmatrix} p \\ \vdots \\ p \\ \vdots \\ \vdots \\ \vdots \end{pmatrix}}^{\star} = \varphi_n(x)$$


Likewise if n is odd we have

$$\overset{\star}{\underbrace{x}}_{n} \mapsto I^{-1/2} \cdot \left(\underbrace{ \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \end{array} \\ p \end{array} \\ \vdots \end{array} \\ \vdots \end{array} }^{p} \\ \begin{array}{c} \end{array} \\ p \end{array} \right) = \varphi_{n}(x)$$

We will now show that $Sym(\mathcal{P})$ is indeed the planar algebra of $M_0 \otimes M_0^{op} \subset M_0 \boxtimes M_0^{op}$. First we need the following technical lemma, which will imply the compatibility of gluing for $Sym(\mathcal{P})$.

Lemma 4.4. Let T be a planar tangle which is partially labelled by q's. Beginning from the marked interval and moving clockwise, number the outgoing strings of each q by $1, \ldots, 6$. Assume that the

following condition is satisfied: for each q appearing in T, there is another one such that strings 1 and 2 of the first are connected to strings 4 and 3 of the second, respectively. Suppose that T contains a subtangle of the following form:



Let T' be the tangle obtained from T by removing this subtangle and replacing it with the following:


*тг	- 1 - 1
¦Ц	ЩĽ
(p)	(p)
İΠ	Π
<u>, I</u> L	_ []_'

Then in the planar algebra $\mathcal{C}_k(\mathcal{P} \otimes \mathcal{P}^{rev})$, we have $Z_T = Z_{T'}$.


Proof. We may assume without loss of generality that T is connected. Indeed, suppose there were some connected component consisting of internal discs and strings. If the subtangle in question is contained in this, then we may restrict to this connected component. Otherwise we may remove this component, which for fixed inputs will contribute the same multiplicative factor to both Z_T and $Z_{T'}$.

Label the upper and lower q's appearing in the diagram by q_t and q_b , respectively. Now consider following strings 1 and 2 of the q_b : by assumption we will arrive at strings 4 and 3 of another q, and so on. Following strings 1 and 2 of this new q we would arrive at strings 4 and 3 of another q, and so on. Eventually we must arrive back at strings 4 and 3 of q_b . There are two cases to consider, depending on whether this path meets q_t or not. First suppose that it does not. Then the winding number of this path around q_t is either 0 or 1. We may assume without loss of generality that it is equal to 0. Indeed, if it is equal to 1, then consider the path obtained by starting at strings 1 and 2 of the q_t and following the same procedure. Then this path does not meet q_b , and it is easy to see that the winding number around q_b must be equal to 0 by planarity. We may then rotate the picture by 180 degrees, reversing the roles of q_t and q_b .

We may now isotope the diagram to obtain the following picture:



Note that we are using the assumption that T is connected here, otherwise there could be something appearing in the region adjacent to the marked interval of q_b . By applying relation (4) of Proposition 4.1, we may modify T as follows (without effecting Z_T):




The result then follows from an application of Lemma 4.2 (or Proposition 4.1 (3) if there are only 2 q's appearing).

Now consider the second case, where the path constructed above contains q_t as well as q_b . Consider the part of the path which begins at string 1 of q_b and ends at string 4 of q_t . Suppose


STEPHEN CURRAN[†]

that we extend this downward to connect back to q_b to obtain a closed path. Then the winding number of this path around any point in region adjacent to the marked interval of q_b is either 0 or 1. As above, by reversing the roles of q_b and q_t if necessary, we may assume that it is equal to 1.

After isotoping, we may then arrange the diagram as follows:



where S is a subtangle of T. Apply relation (4) of Proposition 4.1 to obtain the following:



The result then follows from (5) of Proposition 4.1.

Remark 4.5. Note that the tangle T' obtained from T as above still satisfies the hypotheses of the Lemma (modulo an application of Proposition 4.1 (2) if necessary). So by iterating this procedure, we may make this replacement for every occurrence of the subdiagram which appeared in the statement of the lemma.

Note that if T is fully labelled by q's and satisfies the hypotheses of the Lemma, then by iterating this procedure and applying Proposition 4.1 we can compute the partition function Z_T . In particular, we see that in the (unshaded) planar algebra generated by q in the unshaded 2-cabling of $C_k(\mathcal{P} \otimes \mathcal{P}^{rev})$ we have the relation

which together with the rotational invariance of q and the loop parameter I determines the partition function.

Theorem 4.6. $Sym(\mathcal{P})$ is a spherical C^* -planar algebra, and the identification of $Sym(\mathcal{P})_n$ with $\mathcal{M}'_0 \cap \mathcal{M}_n$ is an isomorphism between $Sym(\mathcal{P})$ and the planar algebra of the inclusion $M_0 \otimes M_0^{op} \subset M_0 \boxtimes M_0^{op}$.

Proof. Since $C_k(\mathcal{P} \otimes \mathcal{P}^{rev})$ is a planar algebra, it is clear that the action Z_T^{Sym} depends only on the isotopy class of T. We must show that the action is compatible with the gluing of tangles. So suppose that we have two tangles T and S, such that the *i*-th input disc of T and the output disc of S both have the same number of marked points 2n. Now we have $Z_{T\circ_i S}^{Sym} = \sigma(T \circ_i S) Z_{f(T\circ_i S)}$ while $Z_T^{Sym} \circ_i Z_S^{Sym} = \sigma(T)\sigma(S) Z_{f(T)} \circ Z_{f(S)} = \sigma(T)\sigma(S) Z_{f(T)\circ_i f(S)}$. It is clear that $\sigma(T\circ_i S) = \sigma(T)\sigma(S)$, so it remains to show that $Z_{f(T)\circ_i f(S)} = Z_{f(T\circ_i S)}$.

Now in $f(T) \circ_i f(S)$ there are *n* copies of the diagram from the statement of Lemma 4.4 which are arranged around the former boundary of the *i*-th input disc of \widetilde{T} . Since $T \circ_i S$ is a shaded tangle, it is not hard to see that $f(T) \circ_i f(S)$ satisfies the conditions of that lemma, and so we may make the indicated replacements. After applying (2) of Proposition 4.1 if needed, the resulting tangle is then clearly isotopic to $f(T \circ_i S)$, and hence we have $Z_{f(T \circ_i S)} = Z_{f(T) \circ_i f(S)}$ as desired. This proves that $Sym(\mathcal{P})$ is a planar algebra.

The remaining properties of a spherical C^* -planar algebra follow easily from the corresponding properties of $\mathcal{C}_k(\mathcal{P} \otimes \mathcal{P}^{rev})$ (note that one has to be careful about the spin factors when checking sphericality).

Now in Example 4.3, we have shown that the identification of $Sym(\mathcal{P})_n$ with $\mathcal{M}'_0 \cap \mathcal{M}_n$ is compatible with the action of several classes of tangles. But it is a well-known result of Jones (see [13, Theorem 4.2.1]) that the planar algebra structure on $(\mathcal{M}'_0 \cap \mathcal{M}_n)_{n\geq 0}$ is uniquely determined by the action of these tangles. So it follows that this identification gives an isomorphism of planar algebras between $Sym(\mathcal{P})$ and $\mathcal{P}(\mathcal{M}_0 \subset \mathcal{M}_1)$, which completes the proof. \Box

Remark 4.7. We have worked with the inclusion $M_0 \otimes M_0^{op} \subset M_0 \boxtimes M_0^{op}$ instead of the symmetric enveloping inclusion $M_1 \otimes M_1^{op} \subset M_1 \boxtimes M_1^{op}$ for simplicity. However, we can now recover the planar algebra of the symmetric enveloping inclusion (or asymptotic inclusion) as follows. Since $M_0 \otimes M_0^{op} \subset M_0 \boxtimes M_0^{op}$ is isomorphic to the compression by $e_0 \otimes e_0^{op}$ of the inclusion $M_2 \otimes M_2^{op} \subset$ $M_2 \boxtimes M_2^{op}$, it follows that these inclusions have the same planar algebra. The latter subfactor is isomorphic to the symmetric enveloping inclusion for $M_1 \subset M_2$ by [1]. The planar algebra of $M_1 \subset M_2$ is \mathcal{P}^{op} , the dual planar algebra of \mathcal{P} in which the shadings are reversed [13]. As discussed by Popa [19], the planar algebra of the symmetric enveloping inclusion of a subfactor depends only on the planar algebra of that subfactor. By duality, it now follows that the planar algebra of the symmetric enveloping inclusion $M_1 \otimes M_1^{op} \subset M_1 \boxtimes M_1^{op}$ is isomorphic to $Sym(\mathcal{P}^{op})$.

5. FUSION RULES AND THE AFFINE CATEGORY

In this section we compute the fusion rules for the asymptotic inclusion, recovering some results of Ocneanu and Evans-Kawahigashi [4]. In particular, we show that the fusion rules for $\mathcal{M}_1 - \mathcal{M}_1$ bimodules are described by the affine category of \mathcal{P} . We remark that it has recently been shown by Das, Ghosh and Gupta [3] that the category of affine Hilbert representations is equivalent to the Drinfeld center of the fusion category associated to \mathcal{P} (see also [2]).

The affine category: We briefly recall the definition of the *affine category* of a planar algebra \mathcal{P} , for further details see [15], [6].

Definition 5.1. An affine (n, m)-tangle T is a planar tangle with outer box equal to $\{(x, y) \in \mathbb{R}^2 : \max(|x|, |y|) = 2\}$, a distinguished inner box equal to $\{(x, y) \in \mathbb{R}^2 : \max(|x|, |y|) = 1\}$, with marked points $\{(-2, \frac{2i}{n}) : -n \leq i < n\}$ (resp. $\{(-1, \frac{i}{m}) : -m \leq k < m\}$) on the boundary of the outer box (resp. distinguished inner box) and with distinguished intervals occuring at the top of the outer box and distinguished inner box.

Typically one must specify the shading near the boundary of the outer (resp. distinguished inner box) when n = 0 (resp. m = 0). Here we will only be concerned with the 'positive' part of the affine category, where these regions are assumed to be unshaded.


If \mathcal{P} is a planar algebra we say that an affine (n, m)-tangle T is \mathcal{P} -labelled if we have an assignment of elements of \mathcal{P} to each input box of T, except for the distinguished one. Note that T then determines a linear map $P_m \to P_n$ by assigning the element of P_m to the distinguished input box and applying Z_T .

Define the (positive) affine category $Aff^+(\mathcal{P})$ to have one object for each $k \geq 0$, and with morphisms equal to isotopy classes of \mathcal{P} -labelled affine tangles, where the isotopies are required to fix the outer and distinguished inner boxes. If T (resp. S) is an affine (n,k) (resp. (k,m)) tangle, the composition TS is the affine (n,m)-tangle obtained by scaling S by a factor of 1/2, composing the resulting diagram with T and then rescaling the resulting tangle. Let $F\mathcal{A}^+(\mathcal{P})$ be the linearization of this category, i.e. the objects are the same but the morphisms are vector spaces with bases given by isotopy classes of affine tangles.

STEPHEN CURRAN[†]

Remark 5.2. The distinction between the affine category and the *annular category* of [15] is that annular tangles are taken up to isotopies which are not required to fix the outer and distinguished inner boxes.

For $k \ge 0$ we have a natural map ψ_k from (isotopy classes of) labelled planar (n+m+2k)-tangles to (isotopy classes of) affine (n, m)-tangles given by:



It is easy to see that every isotopy class of affine (n, m)-tangles is in the range of ψ_k for k sufficiently large.


Define $\mathcal{R} \subset F\mathcal{A}^+(\mathcal{P})$ by

$$\mathcal{R} = \operatorname{span} \left\{ \sum \psi_k(T_i) : T_i \text{ are } \mathcal{P}\text{-labelled } (n+m+2k)\text{-tangles such that } \sum Z_{T_i} = 0 \right\}.$$

Definition 5.3. The (positive) affine algebroid $\mathcal{A}^+(\mathcal{P}) = {\mathcal{A}(\mathcal{P})_{n,m} : n, m \ge 0}$ is the quotient of $F\mathcal{A}^+(\mathcal{P})$ by \mathcal{R} .

In other words, $\mathcal{A}^+(\mathcal{P})$ is the quotient of the universal affine algebroid of \mathcal{P} by all relations which hold in a contractible disc. It is easy to see that composition of affine tangles passes to $\mathcal{A}^+(\mathcal{P})$. It is clear from the definitions that ψ_k gives a well-defined linear map $P_{n+m+2k} \to \mathcal{A}(\mathcal{P})_{n,m}$. Moreover, we have the following description of the kernel.

Lemma 5.4. The kernel of $\psi_k : P_{n+m+2k} \to \mathcal{A}(\mathcal{P})_{n,m}$ is spanned by elements of the form



for $x \in P_{2n+k}$ and $y \in P_{2k}$.

Proof. We prove this first for the universal planar algebra $\mathcal{P}(L)$ with labelling set L = P (cf. [13]). In this case P_{n+m+2k} is replaced by the vector space with basis given by isotopy classes of \mathcal{P} -labelled n+m+2k-tangles, and $\mathcal{A}(\mathcal{P})_{n,m}$ is replaced by the vector space with basis given by isotopy classes of \mathcal{P} -labelled affine (n, m)-tangles. Since ψ_k maps basis elements to basis elements, its kernel is spanned by elements of the form $T_1 - T_2$, where T_1, T_2 are \mathcal{P} -labelled planar n+m+2k tangles such that $\psi_k(T_1)$ is isotopic to $\psi_k(T_2)$. By a standard topological argument it follows that the kernel is spanned by elements of the form appearing in the statement of the lemma, where x is a labelled n+2k-tangle and y is a labelled 2k-tangle. Now taking the quotient by the kernel of the partition function $Z: \mathcal{P}(L) \to \mathcal{P}$, we see that the result holds for \mathcal{P} as well.

Note that by Proposition 2.6, if \mathcal{P} is finite-depth then ψ_k is surjective if $2k \ge depth(\mathcal{P})$. In particular, if \mathcal{P} is finite-depth then $\mathcal{A}(\mathcal{P})_{n,m}$ is finite-dimensional for every $n, m \ge 0$.

A module over $\mathcal{A}^+(\mathcal{P})$ is a graded vector space $V = (V_n)$ together with an action of affine tangles. So to each affine (n, m)-tangle there is an associated linear map $V_m \to V_n$, which is compatible with composition of affine tangles. In particular, $\mathcal{P}^+ = (P_n)_{n\geq 0}$ is a module over $\mathcal{A}^+(\mathcal{P})$. The *weight* of a module is the least value of n such that V_n is non-trivial. If each V_n is a finite-dimensional Hilbert space, and we have

$$\langle av, w \rangle = \langle v, a^*w \rangle$$

for all $v, w \in V$ and $a \in \mathcal{A}^+(\mathcal{P})$, then we say that V is a Hilbert module over $\mathcal{A}^+(\mathcal{P})$. For Hilbert modules the notions of *indecomposability* and *irreducibility* are equivalent, see [15], [6].

Affine representations and Ocneanu's tube algebra:

We will now assume that \mathcal{P} is finite-depth. Let $2k \geq depth(\mathcal{P})$. For each $v \in \Gamma_+$, fix a minimal projection t_v in the central component of P_{2k} corresponding to v. Recall that $\operatorname{tr}_{P_{2k}}(t_v) = \delta^{-2k}\mu_v$. Let $t = \sum t_v$, then Ocneanu's tube algebra is the compression

$$Tube(\mathcal{P}) = t(\mathcal{A}(\mathcal{P})_k)t = \bigoplus_{v,w \in \Gamma_+} Tube(\mathcal{P})_{v,w},$$

where we set $Tube(\mathcal{P})_{v,w} = t_v(\mathcal{A}(\mathcal{P})_k)t_w$.

Remark 5.5. Ocneanu defined the tube algebra [4] using Turaev-Viro topological quantum field theory, but it is not hard to see that our definition is equivalent.

For $v \in \Gamma_+$ and $n \ge 0$, define $H_n(v) \subset P_{n+k}$ to be the range of the tangle:



 $H_n(v)$ is a Hilbert space with inner product $\langle x, y \rangle_{H_n(v)} = \frac{\delta^{2k}}{\mu_v} \langle x, y \rangle_{P_{n+k}}$.

Observe that

$$\left(\bigoplus_{v,w\in\Gamma_+} Tube(\mathcal{P})_{v,w} \otimes (H_n(v) \otimes H_m(w)^*)\right)_{n,m}$$

>0

has a natural algebroid structure given as follows: the product of $a \otimes x_1 \otimes y_1$ in $Tube(\mathcal{P})_{v,w} \otimes (H_n(v) \otimes H_k(w)^*)$ and $b \otimes x_2 \otimes y_2$ in $Tube(\mathcal{P})_{w',z} \otimes (H_k(w') \otimes H_m(z)^*)$ is given by

$$(a \otimes x_1 \otimes y_1)(b \otimes x_2 \otimes y_2) = \delta_{w,w'} \langle y_1, x_2 \rangle_{H_k(w)} \cdot (ab \otimes x_1 \otimes y_2).$$

For $n, m \ge 0$ define $\phi_{n,m} : \bigoplus_{v,w \in \Gamma_+} Tube(\mathcal{P})_{v,w} \otimes (H_n(v) \otimes H_m(w)^*) \to \mathcal{A}(\mathcal{P})_{n,m}$ by

for $a \otimes x \otimes y^* \in Tube(\mathcal{P})_{v,w} \otimes (H_n(v) \otimes H_m(w)^*).$

Proposition 5.6. $\phi = (\phi_{n,m})$ gives an isomorphism of algebroids

$$\mathcal{A}^+(\mathcal{P}) \simeq \left(\bigoplus_{v,w\in\Gamma_+} Tube(\mathcal{P})_{v,w} \otimes (H_n(v) \otimes H_m(w)^*)\right)_{n,m}$$

Proof. That $(\phi_{n,m})$ is compatible with composition follows easily from the identity

$$\bullet t_v \bullet y^* \bullet x \bullet t_w \bullet = \delta_{v,w} \langle y, x \rangle_{H_n(v)} \cdot t_v,$$

which holds since t_v, t_w are minimal projections.

It remains to show that $\phi_{n,m}$ is a linear isomorphism for each n, m. Suppose we have an element

$$z = \sum_{v,w \in \Gamma_+} a_{v,w} \otimes x_v \otimes y_w^* \in \bigoplus_{v,w \in \Gamma_+} Tube(\mathcal{P})_{v,w} \otimes (H_n(v) \otimes H_m(w)^*).$$

Then we have

$$\phi_{n,m}((t_v \otimes t_v \otimes x_v^*)z(t_w \otimes y_w \otimes t_w)) = \phi_{k,k}(a_{v,w} \otimes |x_v|^2 \cdot t_v \otimes |y_w|^2 \cdot t_w) = |x_v|^2 |y_w|^2 \cdot a_{v,w}$$

By the above it follows that $\phi_{n,m}(z) = 0$ implies z = 0, so $\phi_{n,m}$ is injective. Surjectivity follows from Proposition 2.6 and the fact that $1_{P_{2k}}$ is contained in the linear span of $\{xt_vy : x, y \in P_{2k}, v \in \Gamma_+\}$.

We can now determine the irreducible Hilbert modules over $\mathcal{A}^+(\mathcal{P})$. It is clear that any minimal central projection of $Tube(\mathcal{P})$ must be contained in $Tube(\mathcal{P})_{v,v}$ for some $v \in \Gamma_+$. Suppose that p is such a projection, and let V be a finite-dimensional Hilbert space on which $p(Tube(\mathcal{P}))p$ acts irreducibly. Extend this action to $Tube(\mathcal{P})$ by letting 1 - p act by zero. For $n \ge 0$ define $V_n = V \otimes H_n(v)$. Then $\mathcal{V} = (V_n)_{n>0}$ has a natural Hilbert module structure over

$$\left(\bigoplus_{w,z\in\Gamma_+} Tube(\mathcal{P})_{w,z}\otimes (H_n(w)\otimes H_m(z)^*)\right)_{n,m}$$

We can use the isomorphism of Proposition 5.6 to make \mathcal{V} into a Hilbert module over $\mathcal{A}^+(\mathcal{P})$.

Theorem 5.7. $\mathcal{V} = (V_n)_{n \geq 0}$ is an irreducible Hilbert module over $\mathcal{A}^+(\mathcal{P})$, of weight d(*, v)/2. Moreover, any irreducible Hilbert module over $\mathcal{A}^+(\mathcal{P})$ is of this form.

Proof. For fixed $n \ge 0$, the tensor product of the actions of $Tube(\mathcal{P})$ on V and $H_n(v) \otimes H_n^*(v) \simeq \mathcal{B}(H_n(v))$ on $H_n(v)$ is clearly irreducible, and therefore the action of $\mathcal{A}(\mathcal{P})_n$ on $V_n = V \otimes H_n(v)$ is irreducible as well. It follows from [15] that $V = (V_n)$ is an irreducible Hilbert module over $\mathcal{A}^+(\mathcal{P})$. The weight of this module is the smallest value of n for which $H_n(v)$ is non-trivial, which is equal to d(*, v)/2.


Now by [15], any irreducible Hilbert module $\mathcal{V} = (V_n)$ over $\mathcal{A}^+(\mathcal{P})$ must have the property that V_n is an irreducible module over $\mathcal{A}(\mathcal{P})_n$ for each $n \geq 0$. Moreover, if $\mathcal{W} = (W_n)$ is another irreducible Hilbert module such that V_n and W_n are isomorphic $\mathcal{A}(\mathcal{P})_n$ -modules for some n, then \mathcal{V} and \mathcal{W} are isomorphic modules over $\mathcal{A}^+(\mathcal{P})$. Since the isomorphism classes of irreducible modules over the multimatrix algebra $\mathcal{A}(\mathcal{P})_n$ correspond to central projections, it follows from Proposition 5.6 that every irreducible Hilbert module over $\mathcal{A}^+(\mathcal{P})$ is of the form described above. \Box

The principal and dual principal graphs:


We will relate the lattice of higher relative commutants for $\mathcal{M}_0 \subset \mathcal{M}_1$ with the affine category, which will allow us to compute the principal and dual principal graphs. First we need to further analyze the map ψ_k defined above. We will assume that $2k \geq depth(\mathcal{P})$, so that ψ_k is surjective.

First we show that the kernel is determined by the projection $p = p^{(1)} \otimes p^{(2)}$ from Section 2. For this we need the following lemma.


Lemma 5.8. Let $x \in P_{2(n+k)}$ and $y \in P_{2k}$, then we have



Proof. By Proposition 2.6, it suffices to show that




for $x_1, x_2, z_1, z_2 \in P_{n+2k}$. By Lemma 2.1, the left hand side is equal to



Where the equality above follows from rotating the top and bottom components on the left hand side by 180 degrees and then applying Lemma 2.1 again. But if we applied the same procedure to the right hand side of the first equation it is clear that we would obtain the same diagram, so the result follows. \Box

Given $x \in P_{2(n+k)}$, define $p \cdot x$ by



Proposition 5.9. If $x \in P_{2(n+k)}$ then $\psi_k(x) = \psi_k(p \cdot x)$. Moreover, if $\psi_k(x) = 0$ then $p \cdot x = 0$.

Proof. The first statement follows immediately from Proposition 2.3 (4). The second follows from combining Lemmas 5.4 and 5.8. \Box

It follows that ψ_k restricts to a linear isomorphism of $\{p \cdot x : x \in P_{2(n+k)}\}$ onto $\mathcal{A}(\mathcal{P})_n$. We now want to pull back the multiplication from $\mathcal{A}(\mathcal{P})_n$. To do this it is convenient to modify the map

 ψ_k . So we define $\tilde{\psi}_k : P_{2(n+k)} \to \mathcal{A}(\mathcal{P})_n$ by

It is clear that $\tilde{\psi}_k$ has the same kernel as ψ_k , so $\tilde{\psi}_k$ is a linear isomorphism when restricted to $\{p \cdot x : x \in P_{2(n+k)}\}$. For *n* even we also define $\tilde{\psi}_k^{op} : P_{2(n+k)} \to \mathcal{A}(\mathcal{P})_n$ by

Note that we require n even because of the shading. It is clear that $\tilde{\psi}_k^{op}$ is also a linear isomorphism when restricted to $\{p \cdot x : x \in P_{2(n+k)}\}$.

Now for $x, y \in P_{2(n+k)}$ define


$$x \star_q y = \begin{pmatrix} & & \\ & &$$

Proposition 5.10. For $x, y \in P_{2(n+k)}$ we have

$$\tilde{\psi}_k(x \star_q y) = \tilde{\psi}_k(x) \cdot \tilde{\psi}_k(y)$$
$$\tilde{\psi}_k^{op}(y \star_q x) = \tilde{\psi}_k^{op}(x) \cdot \tilde{\psi}_k^{op}(y)$$

Proof. This follows from Proposition 2.7.

Corollary 5.11. We have an embeddings $\mathcal{A}(\mathcal{P})_n \hookrightarrow P_{2n+4k}$ and $\mathcal{A}(\mathcal{P})_n \hookrightarrow P_{2n+4k}^{op}$ given by



for $x \in P_{2(n+k)}$.

Proof. This follows from Corollary 2.9 and the previous proposition.

Theorem 5.12. For $n \ge 2$ we have an isomorphism of commuting squares:


$$\begin{bmatrix} \mathcal{M}'_0 \cap \mathcal{M}_{2n-1} & \subset & \mathcal{M}'_0 \cap \mathcal{M}_{2n} \\ \cup & & \cup \\ \mathcal{M}'_1 \cap \mathcal{M}_{2n-1} & \subset & \mathcal{M}'_1 \cap \mathcal{M}_{2n} \end{bmatrix} \simeq \begin{bmatrix} p_{n-1}(P_{4(n-1)k})p_{n-1} & \subset & p_n(P_{2nk} \otimes P_{2nk}^{op})p_n \\ \cup & & \cup \\ p_{n-2}(\mathcal{A}(\mathcal{P})_{2(n-2)k})p_{n-2} & \subset & p_{n-1}(P_{4(n-1)k}^{op})p_{n-1} \end{bmatrix}$$

Proof. This isomorphisms follow from Proposition 3.8 and Theorem 3.2. The compatibility of the inclusions can be easily checked using the description of $Sym(\mathcal{P})$ from the previous section, the details are left to the reader.

Remark 5.13. We also have an isomorphism of the dual squares

$$\begin{bmatrix} \mathcal{M}_0' \cap \mathcal{M}_{2n} & \subset & \mathcal{M}_0' \cap \mathcal{M}_{2n+1} \\ \cup & & \cup \\ \mathcal{M}_1' \cap \mathcal{M}_{2n} & \subset & \mathcal{M}_1' \cap \mathcal{M}_{2n+1} \end{bmatrix} \simeq \begin{bmatrix} p_n(P_{2nk} \otimes P_{2nk}^{op})p_n & \subset & p_n(P_{4nk})p_n \\ \cup & & \cup \\ p_{n-1}(P_{4(n-1)k}^{op})p_{n-1} & \subset & p_{n-1}(\mathcal{A}(\mathcal{P})_{2(n-1)k})p_{n-1} \end{bmatrix}$$


where the inclusion $P_{4(n-1)k}^{op} \subset \mathcal{A}(\mathcal{P})_{2(n-1)k}$ is given by:




We can now compute the principal and dual principal graphs by determining the Bratelli diagrams for the above inclusions.

Proposition 5.14. We have the following Bratelli diagrams.

(1) For $n \geq 2$ the Bratelli diagram of $P_{4(n-1)k} \subset P_{2nk} \otimes P_{2nk}^{op}$ is as follows. The even vertices are indexed by pairs of $M_0 - M_0$ bimodules (X_v, \overline{X}_w) for $v, w \in \Gamma_+$. The odd vertices are indexed by $M_0 - M_0$ bimodules $X_z, z \in \Gamma_+$. The number of edges between (X_v, \overline{X}_w) and X_z is equal to the multiplicity of X_z in $X_v \otimes \overline{X}_w$, i.e. the dimension of the range of the partially labelled tangle:



(2) For $n \geq 1$ the Bratelli diagram of $P_{4nk}^{op} \subset \mathcal{A}(\mathcal{P})_{2nk}$ is as follows. The odd vertices are indexed by $M_0 - M_0$ bimodules \overline{X}_z for $z \in \Gamma_+$. The even vertices are indexed by irreducible Hilbert modules over $\mathcal{A}^+(\mathcal{P})$, or equivalently by minimal central projections in Ocneanu's tube algebra Tube(\mathcal{P}). Given an irreducible Hilbert module V over $\mathcal{A}^+(\mathcal{P})$, let π_V be a minimal projection in the central component of Tube(\mathcal{P}) corresponding to V via Theorem 5.4. Then the number of edges between \overline{X}_z and V is equal to the dimension of the range of the following partially labelled affine tangle, viewed as a linear map $P_{4k} \to \mathcal{A}(\mathcal{P})_k$.



Proof. Recall that if $i : A \to B$ is a unital inclusion of multi-matrix algebras and $p \in A$ (resp. $q \in B$) is a minimal projection in the central component labelled v (resp. w) then the number of

STEPHEN CURRAN[†]

edges between v and w in the Bratelli diagram of the inclusion is equal to the dimension of the subspace $\{i(p)bq : b \in B\} \subset B$. The results then follow from the previous proposition and the remark which followed.

Remark 5.15. It follows from Theorem 5.12 that the principal (resp. dual principal) graph of $M_0 \otimes M_0^{op} \subset M_0 \boxtimes M_0^{op}$ is the connected component of (X_*, X_*) (resp. X_*) of the Bratelli diagram described in (1) (resp. (2)) above.

The vertices in the Bratelli diagrams above which are not in the principal or dual principal graphs do not correspond to bimodules arising from the Jones tower \mathcal{M}_n . However, these 'missing' bimodules are easily constructed. Indeed, recall that we have $\mathcal{M}_3 \simeq p(M_{2k} \boxtimes M_{2k}^{op})p$. This gives $M_{2k} \boxtimes M_{2k}^{op}$ a $\mathcal{M}_i - \mathcal{M}_j$ bimodule structure for i, j = 0, 1. By adapting the arguments above it can be proved that there is an isomorphism of commuting squares:

$$\begin{bmatrix} \operatorname{Hom}_{\mathcal{M}_{0},\mathcal{M}_{1}} \left(L^{2} \left(M_{2k} \boxtimes M_{2k}^{op} \right) \right) &\subset \operatorname{Hom}_{\mathcal{M}_{0},\mathcal{M}_{0}} \left(L^{2} \left(M_{2k} \boxtimes M_{2k}^{op} \right) \right) \\ \cup & \cup \\ \operatorname{Hom}_{\mathcal{M}_{1},\mathcal{M}_{1}} \left(L^{2} \left(M_{2k} \boxtimes M_{2k}^{op} \right) \right) &\subset \operatorname{Hom}_{\mathcal{M}_{1},\mathcal{M}_{0}} \left(L^{2} \left(M_{2k} \boxtimes M_{2k}^{op} \right) \right) \end{bmatrix} \simeq \begin{bmatrix} P_{8k} &\subset P_{6k} \otimes P_{6k}^{op} \\ \cup & \cup \\ \mathcal{A}(\mathcal{P})_{2k} &\subset P_{8k}^{op} \end{bmatrix}$$

It follows that the fusion graphs of irreducible bimodules which appear in $L^2(M_{2k} \boxtimes M_{2k})$ are precisely the Bratelli diagrams of Proposition 5.14.

Fusion rules for \mathcal{M}_1 - \mathcal{M}_1 bimodules:

It is well-known that the quantum double of a fusion category is always braided. In the subfactor setting, it was likewise shown by Ocneanu [18] and Evans-Kawahigashi [4] that the fusion category of bimodules arising from the asymptotic inclusion is braided (see also [10, 17]). Below we show how the fusion rules and braiding for $\mathcal{M}_1 - \mathcal{M}_1$ bimodules can be seen in our framework.

Define $v_n \in \mathcal{A}(\mathcal{P})_{2nk}$ by




For
$$x \in P_{2(n+k)}$$
 define x^{op} by

$\begin{bmatrix} \star & & \\ $
--

Observe that we have

$$\tilde{\psi}_k(x^{op}) = v_n^* \big(\tilde{\psi}_k(x) \big) v_n$$
$$\tilde{\psi}_k^{op}(x^{op}) = v_n^* \big(\tilde{\psi}_k^{op}(x) \big) v_n$$

Define $i_n : \mathcal{A}(\mathcal{P})_{2nk} \otimes \mathcal{A}(\mathcal{P})_{2nk} \hookrightarrow \mathcal{A}(\mathcal{P})_{4(n+1)k}$ by



We then have

$$v_{2(n+1)}^*\big(i_n\big(\tilde{\psi}_k(x)\otimes\tilde{\psi}_k^{op}(y^{op})\big)\big)v_{2(n+1)}=i_n\big(\tilde{\psi}_k(y)\otimes\tilde{\psi}_k^{op}(x^{op})\big),$$

$$u_n = v_{2(n+1)} \cdot i_n(v_n \otimes v_n)$$

then we have

$$u_n^*(i_n(x\otimes y))u_n = i_n(y\otimes x)$$

for any $x, y \in \mathcal{A}(\mathcal{P})_{2nk}$.

Recall that we have $L^2(\mathcal{M}_{2n}) \simeq L^2(\mathcal{M}_n) \otimes_{\mathcal{M}_0} L^2(\mathcal{M}_n)$ as $\mathcal{M}_1 - \mathcal{M}_1$ bimodules. Moreover, the inclusion $\operatorname{Hom}_{\mathcal{M}_1,\mathcal{M}_1}(L^2(\mathcal{M}_n)) \otimes \operatorname{Hom}_{\mathcal{M}_1,\mathcal{M}_1}(L^2(\mathcal{M}_n)) \hookrightarrow \operatorname{Hom}_{\mathcal{M}_1,\mathcal{M}_1}(L^2(\mathcal{M}_n))$ is given by the following tangle (the top-left corner is shaded):


$$x \otimes y \mapsto$$

Proposition 5.16. The following diagram is commutative:

Proof. This can be verified by computing the tangle above in the planar algebra $Sym(\mathcal{P})$, using the description from the previous section.

We are now able to recover the following description of the fusion rules for $\mathcal{M}_1 - \mathcal{M}_1$ -bimodules due to Ocneanu [18] and Evans-Kawahigashi [4].

Corollary 5.17. Let V, W, Z be irreducible $\mathcal{M}_1 - \mathcal{M}_1$ -bimodules (corresponding to irreducible Hilbert modules over $\mathcal{A}^+(\mathcal{P})$). Let π_V, π_W, π_Z be minimal projections in the central components of $Tube(\mathcal{P})$ corresponding to V, W, Z. Let $\tilde{\pi}_V, \tilde{\pi}_W \in P_{4k}$ be such that $\pi_V = \psi_k(\tilde{\pi}_V)$ and $\pi_W = \psi_k^{op}(\tilde{\pi}_W)$. Then the multiplicity of Z in $V \otimes_{\mathcal{M}_1} W$ is the dimension of the range of the following tangle, viewed as a linear map from P_{8k} to $\mathcal{A}(\mathcal{P})_{6k}$:



Moreover, the fusion category of $\mathcal{M}_1 - \mathcal{M}_1$ bimodules is braided.

Proof. For the first statement we compute the Bratelli diagram of the inclusion

$$\operatorname{Hom}_{\mathcal{M}_1,\mathcal{M}_1}(L^2(\mathcal{M}_n)) \otimes \operatorname{Hom}_{\mathcal{M}_1,\mathcal{M}_1}(L^2(\mathcal{M}_n)) \hookrightarrow \operatorname{Hom}_{\mathcal{M}_1,\mathcal{M}_1}(L^2(\mathcal{M}_n)),$$

using the description from the proof of Proposition 5.14. As discussed above, the unitary operator $p_{n-1}u_np_{n-1} \in \operatorname{Hom}_{\mathcal{M}_1,\mathcal{M}_1}(L^2(\mathcal{M}_{2n}))$ implements the braiding for this inclusion.

STEPHEN CURRAN[†]

References

- S. Curran, V. Jones, and D. Shlyakhtenko, On the symmetric enveloping algebra of planar algebra subfactors, arXiv:1105.1721 [math.OA], 2011.
- [2] P. Das, S. K. Ghosh, and V. P. Gupta, Affine modules and the Drinfeld center, arXiv:1010.0460 [math.QA], 2010.
- [3] P. Das, S. K. Ghosh, and V. P. Gupta, Drinfeld center of planar algebra, arXiv:1203.3958 [math.QA], 2012.
- [4] D. E. Evans and Y. Kawahigashi, On Ocneanu's theory of asymptotic inclusions for subfactors, topological quantum field theories and quantum doubles, Internat. J. Math., 6 (1995), 205–228.
- [5] M. Freedman, C. Nayak, K. Walker, and Z. Wang, On picture (2+1)-TQFTs, in Topology and physics, vol. 12 of Nankai Tracts Math., World Sci. Publ., Hackensack, NJ, 2008, 19–106.
- [6] S. K. Ghosh, Planar algebras: a category theoretic point of view, J. Algebra, 339 (2011), 27–54.
- [7] F. M. Goodman, P. de la Harpe, and V. F. R. Jones, Coxeter graphs and towers of algebras, vol. 14 of Mathematical Sciences Research Institute Publications, Springer-Verlag, New York, 1989.
- [8] A. Guionnet, V. Jones, and D. Shlyakhtenko, A semi-finite algebra associated to a subfactor planar algebra, J. Funct. Anal., 261 (2011), 1345–1360.
- [9] A. Guionnet, V. F. R. Jones, and D. Shlyakhtenko, Random matrices, free probability, planar algebras and subfactors, in Quanta of maths, vol. 11 of Clay Math. Proc., Amer. Math. Soc., Providence, RI, 2010, 201–239.
- [10] M. Izumi, The structure of sectors associated with Longo-Rehren inclusions. I. General theory, Comm. Math. Phys., 213 (2000), 127–179.
- [11] M. Izumi, The structure of sectors associated with Longo-Rehren inclusions. II. Examples, Rev. Math. Phys., 13 (2001), 603–674.
- [12] V. F. R. Jones, Index for subfactors, Invent. Math., 72 (1983), 1–25.
- [13] V. F. R. Jones, Planar algebras, arXiv:9909027 [math.QA], 1999.
- [14] V. F. R. Jones, The planar algebra of a bipartite graph, in Knots in Hellas '98 (Delphi), vol. 24 of Ser. Knots Everything, World Sci. Publ., River Edge, NJ, 2000, 94–117.
- [15] V. F. R. Jones, The annular structure of subfactors, in Essays on geometry and related topics, Vol. 1, 2, vol. 38 of Monogr. Enseign. Math., Enseignement Math., Geneva, 2001, 401–463.
- [16] R. Longo and K.-H. Rehren, Nets of subfactors, Rev. Math. Phys., 7 (1995), 567–597, Workshop on Algebraic Quantum Field Theory and Jones Theory (Berlin, 1994).
- [17] M. Müger, From subfactors to categories and topology. II. The quantum double of tensor categories and subfactors, J. Pure Appl. Algebra, 180 (2003), 159–219.
- [18] A. Ocneanu, Quantized groups, string algebras and Galois theory for algebras, in Operator algebras and applications, Vol. 2, vol. 136 of London Math. Soc. Lecture Note Ser., Cambridge Univ. Press, Cambridge, 1988, 119–172.
- [19] S. Popa, Symmetric enveloping algebras, amenability and AFD properties for subfactors, Math. Res. Lett., 1 (1994), 409–425.
- [20] S. Popa, An axiomatization of the lattice of higher relative commutants of a subfactor, Invent. Math., 120 (1995), 427–445.
- [21] S. Popa, Some properties of the symmetric enveloping algebra of a subfactor, with applications to amenability and property T, Doc. Math., 4 (1999), 665–744.
- [22] D. Shlyakhtenko, Free probability, planar algebras, subfactors and random matrices, in Proceedings of the International Congress of Mathematicians, (Hyderabad, India, 2010), vol. III, 1603–1623.

DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES, CA 90095. *E-mail address:* curransr@math.ucla.edu