On a Certain Subalgebra of $U_q(\mathfrak{sl}_2)$ Related to the Degenerate q-Onsager Algebra^{*}

Tomoya HATTAI † and Tatsuro ITO ‡

[†] Iida Highschool, 1-1, Nonoe, Suzu, Ishikawa 927-1213, Japan E-mail: tmyhtti@m2.ishikawa-c.ed.jp

[‡] School of Mathematical Sciences, Anhui University, 111 Jiulong Road, Hefei 230601, China E-mail: tito@staff.kanazawa-u.ac.jp

Received September 30, 2014, in final form January 15, 2015; Published online January 19, 2015 http://dx.doi.org/10.3842/SIGMA.2015.007

Abstract. In [Kyushu J. Math. 64 (2010), 81–144], it is discussed that a certain subalgebra of the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$ controls the second kind TD-algebra of type I (the degenerate q-Onsager algebra). The subalgebra, which we denote by $U'_q(\widehat{\mathfrak{sl}}_2)$, is generated by e_0^+ , e_1^\pm , $k_i^{\pm 1}$ (i = 0, 1) with e_0^- missing from the Chevalley generators e_i^\pm , $k_i^{\pm 1}$ (i = 0, 1) of $U_q(\widehat{\mathfrak{sl}}_2)$. In this paper, we determine the finite-dimensional irreducible representations of $U'_q(\widehat{\mathfrak{sl}}_2)$. Intertwiners are also determined.

Key words: degenerate *q*-Onsager algebra; quantum affine algebra; TD-algebra; augmented TD-algebra; TD-pair

2010 Mathematics Subject Classification: 17B37; 05E30

1 Introduction

Throughout this paper, the ground field is \mathbb{C} and q stands for a nonzero scalar that is not a root of unity. The symbols ε , ε^* stand for an integer chosen from $\{0,1\}$. Let $\mathcal{A}_q = \mathcal{A}_q^{(\varepsilon,\varepsilon^*)}$ denote the associative algebra with 1 generated by z, z^* subject to the defining relations [4]

(TD)
$$\begin{cases} \left[z, \left[z, \left[z, z^*\right]_q\right]_{q^{-1}}\right] = -\varepsilon \left(q^2 - q^{-2}\right)^2 [z, z^*], \\ \left[z^*, \left[z^*, \left[z^*, z\right]_q\right]_{q^{-1}}\right] = -\varepsilon^* \left(q^2 - q^{-2}\right)^2 [z^*, z], \end{cases}$$

where [X, Y] = XY - YX, $[X, Y]_q = qXY - q^{-1}YX$. This paper deals with a subalgebra of the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$ that is closely related to \mathcal{A}_q in the case of $(\varepsilon, \varepsilon^*) = (1, 0)$. If $(\varepsilon, \varepsilon^*) = (0, 0)$, \mathcal{A}_q is isomorphic to the positive part of $U_q(\widehat{\mathfrak{sl}}_2)$. If $(\varepsilon, \varepsilon^*) = (1, 1)$, \mathcal{A}_q is called the q-Onsager algebra. If $(\varepsilon, \varepsilon^*) = (1, 0)$, \mathcal{A}_q may well be called the *degenerate q-Onsager algebra*.

The algebra \mathcal{A}_q arises in the course of the classification of TD-pairs of type I, which is a critically important step in the study of representations of Terwilliger algebras for P- and Q-polynomial association schemes [3]. For this reason, \mathcal{A}_q is called the TD-algebra of type I. Precisely speaking, the TD-algebra of type I is standardized to be the algebra \mathcal{A}_q , where q is the main parameter for TD-pairs of type I; so $q^2 \neq \pm 1$ and q is allowed to be a root of unity. In our case where we assume q is not a root of unity, the classification of the TD-pairs of type I is equivalent to determining the finite-dimensional irreducible representations $\rho : \mathcal{A}_q \to \text{End}(V)$ with the property that $\rho(z)$, $\rho(z^*)$ are both diagonalizable. Such irreducible representations

^{*}This paper is a contribution to the Special Issue on New Directions in Lie Theory. The full collection is available at http://www.emis.de/journals/SIGMA/LieTheory2014.html

of \mathcal{A}_q are determined in [4] via embeddings of \mathcal{A}_q into the augmented TD-algebra \mathcal{T}_q . (In the case of $(\varepsilon, \varepsilon^*) = (1, 1)$, the diagonalizability condition of $\rho(z)$, $\rho(z^*)$ can be dropped, because it turns out that this condition always holds for every finite-dimensional irreducible representation ρ of the q-Onsager algebra \mathcal{A}_q .) \mathcal{T}_q is easier than \mathcal{A}_q to study representations for, and each finite-dimensional irreducible representation $\rho : \mathcal{A}_q \to \operatorname{End}(V)$ with $\rho(z), \rho(z^*)$ diagonalizable can be extended to a finite-dimensional irreducible representation of \mathcal{T}_q via a certain embedding of \mathcal{A}_q into \mathcal{T}_q .

The augmented TD-algebra $\mathcal{T}_q = \mathcal{T}_q^{(\varepsilon,\varepsilon^*)}$ is the associative algebra with 1 generated by $x, y, k^{\pm 1}$ subject to the defining relations

$$(TD)_0 \begin{cases} kk^{-1} = k^{-1}k = 1, \\ kxk^{-1} = q^2x, \\ kyk^{-1} = q^{-2}y, \end{cases}$$
(1)

and

$$(TD)_{1} \quad \begin{cases} \left[x, [x, [x, y]_{q}]_{q^{-1}}\right] = \delta\left(\varepsilon^{*}x^{2}k^{2} - \varepsilon k^{-2}x^{2}\right), \\ \left[y, [y, [y, x]_{q}]_{q^{-1}}\right] = \delta\left(-\varepsilon^{*}k^{2}y^{2} + \varepsilon y^{2}k^{-2}\right), \end{cases}$$
(2)

where $\delta = -(q-q^{-1})(q^2-q^{-2})(q^3-q^{-3})q^4$. The finite-dimensional irreducible representations of \mathcal{T}_q are determined in [4] via embeddings of \mathcal{T}_q into the $U_q(\mathfrak{sl}_2)$ -loop algebra $U_q(L(\mathfrak{sl}_2))$.

Let e_i^{\pm} , $k_i^{\pm 1}$ (i = 0, 1) be the Chevalley generators of $U_q(L(\mathfrak{sl}_2))$. So the defining relations of $U_q(L(\mathfrak{sl}_2))$ are

$$k_{0}k_{1} = k_{1}k_{0} = 1, \qquad k_{i}k_{i}^{-1} = k_{i}^{-1}k_{i} = 1, \qquad k_{i}e_{i}^{\pm}k_{i}^{-1} = q^{\pm 2}e_{i}^{\pm},$$

$$k_{i}e_{j}^{\pm}k_{i}^{-1} = q^{\mp 2}e_{j}^{\pm}, \quad i \neq j, \qquad [e_{i}^{+}, e_{i}^{-}] = \frac{k_{i} - k_{i}^{-1}}{q - q^{-1}}, \qquad [e_{i}^{+}, e_{j}^{-}] = 0, \quad i \neq j,$$

$$[e_{i}^{\pm}, [e_{i}^{\pm}, e_{j}^{\pm}]_{q}]_{q^{-1}}] = 0, \qquad i \neq j.$$
(3)

Note that if $k_0k_1 = k_1k_0 = 1$ is replaced by $k_0k_1 = k_1k_0$ in (3), we have the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$: $U_q(L(\mathfrak{sl}_2))$ is the quotient algebra of $U_q(\widehat{\mathfrak{sl}}_2)$ by the two-sided ideal generated by $k_0k_1 - 1$. For a nonzero scalar s, define the elements x(s), y(s), k(s) of $U_q(L(\mathfrak{sl}_2))$ by

$$\begin{aligned} x(s) &= -q^{-1} (q - q^{-1})^2 (se_0^+ + \varepsilon s^{-1} e_1^- k_1), \\ y(s) &= \varepsilon^* se_0^- k_0 + s^{-1} e_1^+, \\ k(s) &= sk_0. \end{aligned}$$
(4)

Then the mapping

$$\varphi_s: \ \mathcal{T}_q \to U_q(L(\mathfrak{sl}_2)), \qquad x, y, k \mapsto x(s), y(s), k(s), \tag{5}$$

gives an injective algebra homomorphism. If $(\varepsilon, \varepsilon^*) = (0, 0)$, the image $\varphi_s(\mathcal{T}_q)$ coincides with the Borel subalgebra generated by e_i^+ , $k_i^{\pm 1}$ (i = 0, 1). If $(\varepsilon, \varepsilon^*) = (1, 0)$, the image $\varphi_s(\mathcal{T}_q)$ is properly contained in the subalgebra generated by e_0^+ , e_1^\pm , $k_i^{\pm 1}$ (i = 0, 1) with e_0^- missing from the generators; we denote this subalgebra by $U'_q(L(\mathfrak{sl}_2))$. Through the natural homomorphism $U_q(\widehat{\mathfrak{sl}}_2) \to U_q(L(\mathfrak{sl}_2))$, pull back the subalgebra $U'_q(L(\mathfrak{sl}_2))$ and denote the pre-image by $U'_q(\widehat{\mathfrak{sl}}_2)$:

$$U_q'(\widehat{\mathfrak{sl}}_2) = \left\langle e_0^+, e_1^\pm, k_i^{\pm 1} \, | \, i = 0, 1 \right\rangle \subset U_q(\widehat{\mathfrak{sl}}_2).$$

In [4], it is shown that in the case of $(\varepsilon, \varepsilon^*) = (1, 0)$, all the finite-dimensional irreducible representations of \mathcal{T}_q are produced by tensor products of evaluation modules for $U'_q(L(\mathfrak{sl}_2))$ via the embedding φ_s of \mathcal{T}_q into $U'_q(L(\mathfrak{sl}_2))$. Using this fact and the Drinfel'd polynomials, we show in this paper that there are no other finite-dimensional irreducible representations of $U'_q(L(\mathfrak{sl}_2))$ and hence of $U'_q(\widehat{\mathfrak{sl}}_2)$ than those afforded by tensor products of evaluation modules, if we apply suitable automorphisms of $U'_q(L(\mathfrak{sl}_2))$, $U'_q(\widehat{\mathfrak{sl}}_2)$ to adjust the types of the representations to be (1, 1). Here we note that the evaluation parameters are allowed to be zero for $U'_q(L(\mathfrak{sl}_2))$, $U'_q(\widehat{\mathfrak{sl}}_2)$. Details will be discussed in Sections 2 and 3, where the isomorphism classes of finite-dimensional irreducible representations of $U'_q(\widehat{\mathfrak{sl}}_2)$ are also determined. In Section 4, intertwiners will be determined for finite-dimensional irreducible $U'_q(\widehat{\mathfrak{sl}}_2)$ modules.

In our approach, Drinfel'd polynomials are the key tool for the classification of finitedimensional irreducible representations of $U_q(\widehat{\mathfrak{sl}}_2)$, $U'_q(\widehat{\mathfrak{sl}}_2)$, although they are not the main subject of this paper. They are defined in [4], and the point is that they are directly attached to \mathcal{T}_q -modules, not to $U_q(\widehat{\mathfrak{sl}}_2)$ - or $U'_q(\widehat{\mathfrak{sl}}_2)$ -modules. (In the case of $(\varepsilon, \varepsilon^*) = (0, 0)$, they turn out to coincide with the original ones up to the reciprocal of the variable.) So in our approach to the case of $(\varepsilon, \varepsilon^*) = (0, 0)$, finite-dimensional irreducible representations are naturally classified firstly for the Borel subalgebra of $U_q(\widehat{\mathfrak{sl}}_2)$ and then for $U_q(\widehat{\mathfrak{sl}}_2)$ itself. This will be briefly demonstrated in Section 3 as a warm-up for the case of $(\varepsilon, \varepsilon^*) = (1, 0)$, thus giving another proof to the classification theorem of Chari–Pressley [2] and to the main theorems (Theorems 1.16 and 1.17) of [1].

We now review Drinfel'd polynomials for \mathcal{T}_q -modules [4, p. 119]. Let V be a finite-dimensional \mathcal{T}_q -module. We assume the following properties for V:

(D)₀: k is diagonalizable on V with
$$V = \bigoplus_{i=0}^{d} U_i$$
, $k|_{U_i} = sq^{2i-d}$, $0 \le i \le d$, for some nonzero constant s;
(D)₁: dim $U_0 = 1$.

By the relations $(TD)_0 : kk^{-1} = k^{-1}k = 1$, $kxk^{-1} = q^2x$, $kyk^{-1} = q^{-2}y$, it holds that $xU_i \subseteq U_{i+1}$, $yU_i \subseteq U_{i-1}$ $(0 \leq i \leq d)$, where $U_{-1} = U_{d+1} = 0$. So the one-dimensional subspace U_0 is invariant under $y^i x^i$ and we have the sequence $\{\sigma_i\}_{i=0}^{\infty}$ of eigenvalues σ_i of $y^i x^i$ on U_0 : $\sigma_i = y^i x^i|_{U_0}$. Notice that $\sigma_0 = 1$ and $\sigma_i = 0$, $d+1 \leq i$. The Drinfel'd polynomial $P_V(\lambda)$ of the \mathcal{T}_q -module V is defined by

$$P_V(\lambda) = \sum_{i=0}^d (-1)^i \frac{\sigma_i}{(q-q^{-1})^{2i}([i]!)^2} \prod_{j=i+1}^d \left(\lambda - \varepsilon s^{-2} q^{2(d-j)} - \varepsilon^* s^2 q^{-2(d-j)}\right),$$

where $[i] = [i]_q = (q^i - q^{-i})/(q - q^{-1})$ and $[i]! = [1][2] \cdots [i]$ with the understanding of [0]! = 1. Since $\sigma_0 = 1$, $P_V(\lambda)$ is a monic polynomial of degree d.

If V is an irreducible \mathcal{T}_q -module, it is known that V in fact satisfies the properties $(D)_0$, $(D)_1$ [4, Lemma 1.2, Theorem 1.8], and these properties provide a rather simple short proof for the 'injective' part of [4, Theorem 1.9], i.e., for the fact that the isomorphism class of the irreducible \mathcal{T}_q -module V is determined by the trio $(\{\sigma_i\}_{i=0}^{\infty}, s, d)$.

If V is a tensor product of evaluation modules for $U_q(L(\mathfrak{sl}_2))$ in the case of $(\varepsilon, \varepsilon^*) = (1, 1), (0, 0)$ or for $U'_q(L(\mathfrak{sl}_2))$ in the case of $(\varepsilon, \varepsilon^*) = (1, 0)$, we regard V as a \mathcal{T}_q -module via the embedding φ_s of (5). Then it is apparent that the \mathcal{T}_q -module V satisfies the properties $(D)_0, (D)_1$. Moreover it is known that a product formula holds for the Drinfel'd polynomial $P_V(\lambda)$ and it turns out that $P_V(\lambda)$ does not depend on the parameter s of the embedding φ_s [4, Theorem 5.2]. The 'surjective' part of [4, Theorem 1.9] follows from the structure of the zeros of the Drinfel'd polynomial for such a tensor product of evaluation modules regarded as a \mathcal{T}_q -module via the embedding φ_s .

2 Finite-dimensional irreducible representations of $U'_{q}(\mathfrak{sl}_{2})$

The subalgebra $U'_q(\widehat{\mathfrak{sl}}_2)$ of the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$ is generated by e_0^+ , e_1^\pm , $k_i^{\pm 1}$ (i = 0, 1), e_0^- missing from the generators, and has by the triangular decomposition of $U_q(\widehat{\mathfrak{sl}}_2)$ the defining relations

$$k_{0}k_{1} = k_{1}k_{0}, \qquad k_{i}k_{i}^{-1} = k_{i}^{-1}k_{i} = 1, \qquad k_{0}e_{0}^{+}k_{0}^{-1} = q^{2}e_{0}^{+}, \qquad k_{1}e_{1}^{\pm}k_{1}^{-1} = q^{\pm2}e_{1}^{\pm}, k_{1}e_{0}^{+}k_{1}^{-1} = q^{-2}e_{0}^{+}, \qquad k_{0}e_{1}^{\pm}k_{0}^{-1} = q^{\mp2}e_{1}^{\pm}, \qquad [e_{1}^{+}, e_{1}^{-}] = \frac{k_{1}-k_{1}^{-1}}{q-q^{-1}},$$

$$[e_{0}^{+}, e_{1}^{-}] = 0, \qquad \left[e_{i}^{+}, [e_{i}^{+}, e_{j}^{+}]_{q}\right]_{q^{-1}}\right] = 0, \qquad i \neq j.$$

$$(6)$$

Note that if $k_0k_1 = k_1k_0$ is replaced by $k_0k_1 = k_1k_0 = 1$ in (6), we have the defining relations for $U'_q(L(\mathfrak{sl}_2))$.

Let V be a finite-dimensional irreducible $U'_q(\widehat{\mathfrak{sl}}_2)$ -module. Let us first observe that the $U'_q(\widehat{\mathfrak{sl}}_2)$ module V is obtained from a $U'_q(L(\mathfrak{sl}_2))$ -module by applying an automorphism of $U'_q(\widehat{\mathfrak{sl}}_2)$ as follows. Since the element k_0k_1 belongs to the centre of $U'_q(\widehat{\mathfrak{sl}}_2)$, k_0k_1 acts on V as a scalar s by Schur's lemma. Since k_0k_1 is invertible, the scalar s is nonzero: $k_0k_1|_V = s \in \mathbb{C}^{\times}$. Observe that $U'_q(\widehat{\mathfrak{sl}}_2)$ admits an automorphism that sends k_0 to $s^{-1}k_0$ and preserves k_1 . Hence we may assume $k_0k_1|_V = 1$. Then we can regard V as a $U'_q(L(\mathfrak{sl}_2))$ -module.

Let V be a finite-dimensional irreducible $U'_q(L(\mathfrak{sl}_2))$ -module. For a scalar θ , set $V(\theta) = \{v \in V \mid k_0 v = \theta v\}$. So if $V(\theta) \neq 0$, θ is an eigenvalue of k_0 and $V(\theta)$ is the corresponding eigenspace of k_0 . For an eigenvalue θ and an eigenvector $v \in V(\theta)$, it holds that $e_0^+ v \in V(q^2\theta)$ by the relation $k_0 e_0^+ = q^2 e_0^+ k_0$ and $e_1^{\pm} v \in V(q^{\pm 2}\theta)$ by $k_0 e_1^{\pm} = q^{\pm 2} e_1^{\pm} k_0$. We have

$$e_0^+ V(\theta) \subseteq V(q^2\theta), \qquad e_1^\pm V(\theta) \subseteq V(q^{\pm 2}\theta).$$
 (7)

If dim V = 1, then $e_0^+ V = 0$, $e_1^\pm V = 0$ by (7) and $k_0|_V = \pm 1$ by $[e_1^+, e_1^-] = (k_1 - k_1^{-1})/(q - q^{-1}) = (k_0^{-1} - k_0)/(q - q^{-1})$. Such a $U'_q(L(\mathfrak{sl}_2))$ -module V is said to be trivial. Assume dim $V \ge 2$. Choose an eigenvalue θ of k_0 on V. Then $\sum_{i \in \mathbb{Z}} V(q^{\pm 2i}\theta)$ is invariant under the actions of the generators by (7), and so we have $V = \sum_{i \in \mathbb{Z}} V(q^{\pm 2i}\theta)$ by the irreducibility of the $U'_q(L(\mathfrak{sl}_2))$ -module V. Since V is finite-dimensional, there exists a positive integer d and a nonzero scalar s_0 such that the eigenspace decomposition of k_0 is

$$V = \bigoplus_{i=0}^{d} V(s_0 q^{2i-d}).$$
(8)

We want to show that $s_0 = \pm 1$ holds in (8).

Consider the subalgebra of $U'_q(L(\mathfrak{sl}_2))$ generated by e_1^{\pm} , $k_1^{\pm 1}$ and denote it by $\mathcal{U} : \mathcal{U} = \langle e_1^{\pm}, k_1^{\pm 1} \rangle$. Regard V as a \mathcal{U} -module. Since \mathcal{U} is isomorphic to the quantum algebra $U_q(\mathfrak{sl}_2)$, V is a direct sum of irreducible \mathcal{U} -modules, and for each irreducible \mathcal{U} -submodule W of V, the eigenvalues of $k_1 = k_0^{-1}$ on W are either $\{q^{2i-\ell} \mid 0 \leq i \leq \ell\}$ or $\{-q^{2i-\ell} \mid 0 \leq i \leq \ell\}$ for some nonnegative integer ℓ . This implies that (i) $s_0 = \pm q^m$ for some $m \in \mathbb{Z}$ and (ii) if θ is an eigenvalue of k_0 , so is θ^{-1} . It follows from (i) that $V = \bigoplus_{i=0}^d V(\pm q^{2i-d+m})$, and so by (ii), we obtain m = 0, i.e., $s_0 = \pm 1$.

Observe that $U'_q(L(\mathfrak{sl}_2))$ admits an automorphism that sends k_i to $-k_i$ (i = 0, 1) and e_1^+ to $-e_1^+$. Hence we may assume $s_0 = 1$ in (8). Note that in this case, k_1 has the eigenvalues $\{s_1q^{2i-d} \mid 0 \leq i \leq d\}$ with $s_1 = 1$. Such an irreducible module or the irreducible representation

afforded by such is said to be of type (1, 1), indicating $(s_0, s_1) = (1, 1)$. We conclude that the determination of finite-dimensional irreducible representations for $U'_q(\widehat{\mathfrak{sl}}_2)$ is, via automorphisms, reduced to that of type (1, 1) for $U'_q(L(\mathfrak{sl}_2))$.

In the rest of this section, we shall introduce evaluation modules for $U'_q(L(\mathfrak{sl}_2))$ and state our main theorem that every finite-dimensional irreducible representation of type (1, 1) of $U'_q(L(\mathfrak{sl}_2))$ is afforded by a tensor product of them. For $a \in \mathbb{C}$ and $\ell \in \mathbb{Z}_{\geq 0}$, let $V(\ell, a)$ denote the $(\ell + 1)$ dimensional vector space with a basis v_0, v_1, \ldots, v_ℓ . Using (6), it can be routinely verified that $U'_q(L(\mathfrak{sl}_2))$ acts on $V(\ell, a)$ by

$$k_0 v_i = q^{2i-\ell} v_i, \qquad k_1 v_i = q^{\ell-2i} v_i, \qquad e_0^+ v_i = aq[i+1] v_{i+1}, \\ e_1^+ v_i = [\ell-i+1] v_{i-1}, \qquad e_1^- v_i = [i+1] v_{i+1},$$

$$(9)$$

where $v_{-1} = v_{\ell+1} = 0$ and $[t] = [t]_q = (q^t - q^{-t})/(q - q^{-1})$. This $U'_q(L(\mathfrak{sl}_2))$ -module $V(\ell, a)$ is irreducible and called an *evaluation module*. The basis v_0, v_1, \ldots, v_ℓ of the $U'_q(L(\mathfrak{sl}_2))$ -module $V(\ell, a)$ is called a *standard basis*. The vector v_0 is called the *highest weight vector*. Note that the evaluation parameter a is allowed to be zero. Also note that if $\ell = 0$, $V(\ell, a)$ is the trivial module. We denote the evaluation module $V(\ell, 0)$ by $V(\ell)$, allowing $\ell = 0$, and use the notation $V(\ell, a)$ only for an evaluation module with $a \neq 0$ and $\ell \geq 1$.

The $U_q(\mathfrak{sl}_2)$ -loop algebra $U_q(L(\mathfrak{sl}_2))$ has the coproduct $\Delta : U_q(L(\mathfrak{sl}_2)) \to U_q(L(\mathfrak{sl}_2)) \otimes U_q(L(\mathfrak{sl}_2))$ defined by

$$\Delta(k_i^{\pm 1}) = k_i^{\pm 1} \otimes k_i^{\pm 1}, \qquad \Delta(e_i^+) = k_i \otimes e_i^+ + e_i^+ \otimes 1,$$

$$\Delta(e_i^- k_i) = k_i \otimes e_i^- k_i + e_i^- k_i \otimes 1.$$
(10)

The subalgebra $U'_q(L(\mathfrak{sl}_2))$ is closed under Δ . Thus given a set of evaluation modules $V(\ell)$, $V(\ell_i, a_i)$ $(1 \le i \le n)$ for $U'_q(L(\mathfrak{sl}_2))$, the tensor product

$$V(\ell) \otimes V(\ell_1, a_1) \otimes \dots \otimes V(\ell_n, a_n)$$
⁽¹¹⁾

becomes a $U'_q(L(\mathfrak{sl}_2))$ -module via Δ . Note that by the coassociativity of Δ , the way of putting parentheses in the tensor product of (11) does not affect the isomorphism class as a $U'_q(L(\mathfrak{sl}_2))$ module. Also note that if $\ell = 0$ in (11), then V(0) is the trivial module and the tensor product of (11) is isomorphic to $V(\ell_1, a_1) \otimes \cdots \otimes V(\ell_n, a_n)$ as $U'_q(L(\mathfrak{sl}_2))$ -modules. Finally we allow n = 0, in which case we understand that the tensor product of (11) means $V(\ell)$.

With the evaluation module $V(\ell, a)$, we associate the set $S(\ell, a)$ of scalars $aq^{-\ell+1}, aq^{-\ell+3}, \ldots, aq^{\ell-1}$:

$$S(\ell, a) = \{ aq^{2i-\ell+1} \, | \, 0 \le i \le \ell - 1 \}.$$

The set $S(\ell, a)$ is called a *q*-string of length ℓ . Two *q*-strings $S(\ell, a)$, $S(\ell', a')$ are said to be in general position if either

- (i) the union $S(\ell, a) \cup S(\ell', a')$ is not a q-string, or
- (ii) one of $S(\ell, a)$, $S(\ell', a')$ includes the other.

Below is the main theorem of this paper. It classifies the isomorphism classes of the finitedimensional irreducible $U'_q(L(\mathfrak{sl}_2))$ -modules of type (1, 1).

Theorem 1. The following (i), (ii), (ii), (iv) hold.

(i) A tensor product $V = V(\ell) \otimes V(\ell_1, a_1) \otimes \cdots \otimes V(\ell_n, a_n)$ of evaluation modules is irreducible as a $U'_q(L(\mathfrak{sl}_2))$ -module if and only if $S(\ell_i, a_i)$, $S(\ell_j, a_j)$ are in general position for all $i, j \in \{1, 2, ..., n\}$. In this case, V is of type (1, 1).

- (ii) Consider two tensor products $V = V(\ell) \otimes V(\ell_1, a_1) \otimes \cdots \otimes V(\ell_n, a_n)$, $V' = V(\ell') \otimes V(\ell'_1, a'_1) \otimes \cdots \otimes V(\ell'_m, a'_m)$ of evaluation modules and assume that they are both irreducible as a $U'_q(L(\mathfrak{sl}_2))$ -module. Then V, V' are isomorphic as $U'_q(L(\mathfrak{sl}_2))$ -modules if and only if $\ell = \ell'$, n = m and $(\ell_i, a_i) = (\ell'_i, a'_i)$ for all $i, 1 \leq i \leq n$, with a suitable reordering of the evaluation modules $V(\ell_1, a_1), \ldots, V(\ell_n, a_n)$.
- (iii) Every non-trivial finite-dimensional irreducible $U'_q(L(\mathfrak{sl}_2))$ -module of type (1,1) is isomorphic to some tensor product $V(\ell) \otimes V(\ell_1, a_1) \otimes \cdots \otimes V(\ell_n, a_n)$ of evaluation modules.
- (iv) If a tensor product $V = V(\ell) \otimes V(\ell_1, a_1) \otimes \cdots \otimes V(\ell_n, a_n)$ of evaluation modules is irreducible as a $U'_q(L(\mathfrak{sl}_2))$ -module, then any change of the orderings of the evaluation modules $V(\ell)$, $V(\ell_1, a_1), \ldots, V(\ell_n, a_n)$ for the tensor product does not change the isomorphism class of the $U'_q(L(\mathfrak{sl}_2))$ -module V.

3 Proof of Theorem 1(i), (ii), (iii)

Discard the evaluation module $V(\ell)$ from the statement of Theorem 1 and replace $U'_q(L(\mathfrak{sl}_2))$ by $U_q(L(\mathfrak{sl}_2))$ or by \mathcal{B} , where \mathcal{B} is the Borel subalgebra of $U_q(L(\mathfrak{sl}_2))$ generated by e_i^+ , $k_i^{\pm 1}$ (i = 0, 1). Then it precisely describes the classification of the isomorphism classes of finitedimensional irreducible modules of type (1, 1) for $U_q(L(\mathfrak{sl}_2))$ [2] or for \mathcal{B} [1]. There is a one-toone correspondence of finite-dimensional irreducible modules of type (1, 1) between $U_q(L(\mathfrak{sl}_2))$ and \mathcal{B} : every finite-dimensional irreducible $U_q(L(\mathfrak{sl}_2))$ -module of type (1, 1) is irreducible as a \mathcal{B} module and every finite-dimensional irreducible \mathcal{B} -module of type (1, 1) is uniquely extended to a $U_q(L(\mathfrak{sl}_2))$ -module. This sort of correspondence of finite-dimensional irreducible modules exists between $U'_q(L(\mathfrak{sl}_2))$ and \mathcal{T}_q via the embedding φ_s of (5), where \mathcal{T}_q is the augmented TDalgebra with $(\varepsilon, \varepsilon^*) = (1, 0)$. Namely, we shall show that (i) every finite-dimensional irreducible $U'_q(L(\mathfrak{sl}_2))$ -module of type (1, 1) is irreducible as a \mathcal{T}_q -module via certain embedding φ_s of (5), and (ii) every finite-dimensional irreducible \mathcal{T}_q -module is uniquely extended to a $U'_q(L(\mathfrak{sl}_2))$ module of type (1, 1) via the embedding φ_s of (5) with s uniquely determined. Since finitedimensional irreducible \mathcal{T}_q -modules are classified in [4], this gives a proof of Theorem 1.

Apart from the Drinfel'd polynomials, the key to our understanding of the correspondence is the following two lemmas about $U_q(\mathfrak{sl}_2)$ -modules. Let \mathcal{U} denote the quantum algebra $U_q(\mathfrak{sl}_2)$: \mathcal{U} is the associative algebra with 1 generated by X^{\pm} , $K^{\pm 1}$ subject to the defining relations

$$KK^{-1} = K^{-1}K = 1, \qquad KX^{\pm}K^{-1} = q^{\pm 2}X^{\pm}, \qquad [X^+, X^-] = \frac{K - K^{-1}}{q - q^{-1}}.$$
 (12)

Lemma 1 ([4, Lemma 7.5]). Let V be a finite-dimensional \mathcal{U} -module that has the following weight-space (K-eigenspace) decomposition:

$$V = \bigoplus_{i=0}^{d} U_i, \qquad K|_{U_i} = q^{2i-d}, \qquad 0 \le i \le d.$$

Let W be a subspace of V as a vector space. Assume that W is invariant under the actions of X^+ and K:

$$X^+W \subseteq W, \qquad KW \subseteq W.$$

If it holds that

$$\dim(W \cap U_i) = \dim(W \cap U_{d-i}), \qquad 0 \le i \le d,$$

then $X^-W \subseteq W$, i.e., W is a U-submodule.

Lemma 2. If V is a finite-dimensional \mathcal{U} -module, the action of X^- on V is uniquely determined by those of X^+ , $K^{\pm 1}$ on V.

Proof. The claim holds if V is irreducible as a \mathcal{U} -module. By the semi-simplicity of \mathcal{U} , it holds generally.

As a warm-up for the proof of Theorem 1, we shall demonstrate how to use these lemmas in the case of the corresponding theorem [2] for $U_q(L(\mathfrak{sl}_2))$. We want, and it is enough, to show part (iii) of the theorem for $U_q(L(\mathfrak{sl}_2))$ by using the classification of finite-dimensional irreducible \mathcal{B} -modules. This is because the parts (i), (ii), (iv) are well-known in advance of [2], while the finite-dimensional irreducible \mathcal{B} -modules are classified in [4] rather straightforward by the product formula of Drinfel'd polynomials without using the part (iii) in question.

Let V be a finite-dimensional irreducible $U_q(L(\mathfrak{sl}_2))$ -module of type (1, 1). Then V has the weight-space decomposition

$$V = \bigoplus_{i=0}^{d} U_i, \qquad k_0|_{U_i} = q^{2i-d}, \qquad 0 \le i \le d.$$

Regard V as a \mathcal{B} -module. Let W be a minimal \mathcal{B} -submodule of V. Note that W is irreducible as a \mathcal{B} -module. We want to show W = V, i.e., V is irreducible as a \mathcal{B} -module. Since the mapping $(e_0^+)^{d-2i}$: $U_i \to U_{d-i}$ is a bijection and $W \cap U_i$ is mapped into $W \cap U_{d-i}$ by $(e_0^+)^{d-2i}$, we have $\dim(W \cap U_i) \leq \dim(W \cap U_{d-i}), 0 \leq i \leq \lfloor d/2 \rfloor$. Similarly from the bijection $(e_1^+)^{d-2i}$: $U_{d-i} \to U_i$, we get $\dim(W \cap U_{d-i}) \leq \dim(W \cap U_i)$. Thus it holds that

 $\dim(W \cap U_i) = \dim(W \cap U_{d-i}), \qquad 0 \le i \le d.$

Consider the algebra homomorphism from \mathcal{U} to $U_q(L(\mathfrak{sl}_2))$ that sends X^+ , X^- , $K^{\pm 1}$ to e_0^+ , e_0^- , $k_0^{\pm 1}$, respectively. Regard V as a \mathcal{U} -module via this algebra homomorphism. Then $X^+W \subseteq W$, $KW \subseteq W$. Since $\dim(W \cap U_i) = \dim(W \cap U_{d-i})$, $0 \leq i \leq d$, we have by Lemma 1 that $X^-W \subseteq W$, i.e., $e_0^-W \subseteq W$. Similarly, Lemma 1 can be applied to the \mathcal{U} -module V via the algebra homomorphism from \mathcal{U} to $U_q(L(\mathfrak{sl}_2))$ that sends X^+ , X^- , $K^{\pm 1}$ to e_1^+ , e_1^- , $k_1^{\pm 1}$, respectively, in which case the weight-space decomposition of the \mathcal{U} -module V is $V = \bigoplus_{i=0}^{d} U_{d-i}$, $K|_{U_{d-i}} = q^{2i-d}$, $0 \leq i \leq d$. Consequently, we get $X^-W \subseteq W$, i.e., $e_1^-W \subseteq W$. Thus W is $U_q(L(\mathfrak{sl}_2))$ -invariant and we have W = V by the irreducibility of the $U_q(L(\mathfrak{sl}_2))$ -module V. We conclude that every finite-dimensional irreducible $U_q(L(\mathfrak{sl}_2))$ -module of type (1, 1) is irreducible as a \mathcal{B} -module.

Now consider the class of finite-dimensional irreducible \mathcal{B} -modules V, where V runs through all tensor products of evaluation modules that are irreducible as a $U_q(L(\mathfrak{sl}_2))$ -module:

$$V = V(\ell_1, a_1) \otimes \cdots \otimes V(\ell_n, a_n).$$

Then it turns out that the Drinfel'd polynomials $P_V(\lambda)$ of the irreducible \mathcal{B} -modules V exhaust all that are possible for finite-dimensional irreducible \mathcal{B} -modules of type (1,1), as shown in [4, Theorem 5.2] by the product formula

$$P_V(\lambda) = \prod_{i=1}^n P_{V(\ell_i, a_i)}(\lambda), \qquad P_{V(\ell_i, a_i)}(\lambda) = \prod_{\zeta \in S(\ell_i, a_i)} (\lambda + \zeta).$$

Since the Drinfel'd polynomial $P_V(\lambda)$ determines the isomorphism class of the \mathcal{B} -module V of type (1,1) [4, the injectivity part of Theorem 1.9'], there are no other finite-dimensional

irreducible \mathcal{B} -modules of type (1,1). This means that every finite-dimensional irreducible \mathcal{B} module of type (1,1) comes from some tensor product of evaluation modules for $U_a(L(\mathfrak{sl}_2))$.

Let V be a finite-dimensional irreducible $U_q(L(\mathfrak{sl}_2))$ -module of type (1,1). Then V is irreducible as a \mathcal{B} -module and so there exists an irreducible $U_q(L(\mathfrak{sl}_2))$ -module $V' = V(\ell_1, a_1) \otimes \cdots \otimes V(\ell_n, a_n)$ such that V, V' are isomorphic as \mathcal{B} -modules. By Lemma 2, V, V' are isomorphic as $U_q(L(\mathfrak{sl}_2))$ -modules. This completes the proof of part (iii) of the theorem for $U_q(L(\mathfrak{sl}_2))$.

The proof of Theorem 1 can be given by an argument very similar to the one we have seen above for the case of $U_q(L(\mathfrak{sl}_2))$. We prepare two more lemmas for the case of $U'_q(L(\mathfrak{sl}_2))$ to make the point clearer. Set $(\varepsilon, \varepsilon^*) = (1, 0)$ and let \mathcal{T}_q be the augmented TD-algebra defined by $(TD)_0$, $(TD)_1$ in (1), (2). For $s \in \mathbb{C}^{\times}$, let φ_s be the embedding of \mathcal{T}_q into $U'_q(L(\mathfrak{sl}_2))$ given by (4), (5).

Lemma 3. Let V_1 , V_2 be finite-dimensional irreducible $U'_q(L(\mathfrak{sl}_2))$ -modules. If V_1 , V_2 are isomorphic as $\varphi_s(\mathcal{T}_q)$ -modules for some $s \in \mathbb{C}^{\times}$, then V_1 , V_2 are isomorphic as $U'_q(L(\mathfrak{sl}_2))$ -modules.

Proof. By (4), $\varphi_s(\mathcal{T}_q)$ is generated by $se_0^+ + s^{-1}e_1^-k_1$, e_1^+ and $k_i^{\pm 1}$ (i = 0, 1). Since $\langle e_1^{\pm}, k_1^{\pm 1} \rangle$ is isomorphic to the quantum algebra $U_q(\mathfrak{sl}_2)$, the action of e_1^- on V_i , i = 1, 2, is uniquely determined by those of e_1^+ , $k_1^{\pm 1} \in \varphi_s(\mathcal{T}_q)$ by Lemma 2. Apparently the action of e_0^+ on V_i , i = 1, 2, is uniquely determined by those of $se_0^+ + s^{-1}e_1^-k_1$, e_1^- , k_1 , and hence by that of $\varphi_s(\mathcal{T}_q)$. So the action of $U'_q(L(\mathfrak{sl}_2))$ on V_i , i = 1, 2, is uniquely determined by that of $\varphi_s(\mathcal{T}_q)$.

Lemma 4. Let V be a finite-dimensional irreducible $U'_q(L(\mathfrak{sl}_2))$ -module of type (1,1). Then there exists a finite set Λ of nonzero scalars such that V is irreducible as a $\varphi_s(\mathcal{T}_q)$ -module for each $s \in \mathbb{C}^{\times} \setminus \Lambda$.

Proof. For $s \in \mathbb{C}^{\times}$, regard V as a $\varphi_s(\mathcal{T}_q)$ -module. Let W be a minimal $\varphi_s(\mathcal{T}_q)$ -submodule of V. It is enough to show that W = V holds if s avoids finitely many scalars. By (8) with $s_0 = 1$, the eigenspace decomposition of $k_1 = k_0^{-1}$ on V is $V = \bigoplus_{i=0}^{d} U_{d-i}, k_1|_{U_{d-i}} = q^{2i-d}, 0 \leq i \leq d$. The subalgebra $\langle e_1^{\pm}, k_1^{\pm 1} \rangle$ of $U'_q(L(\mathfrak{sl}_2))$ is isomorphic to the quantum algebra $\mathcal{U} = U_q(\mathfrak{sl}_2)$ in (12) via the correspondence of $e_1^{\pm}, k_1^{\pm 1}$ to $X^{\pm}, K^{\pm 1}$. The element $(e_1^+)^{d-2i}$ maps U_{d-i} onto U_i bijectively, $0 \leq i \leq [d/2]$. Also $(e_1^-k_1)^{d-2i}$ maps U_i onto U_{d-i} bijectively, $0 \leq i \leq [d/2]$.

The element $(e_1^+)^{d-2i}$ belongs to $\varphi_s(\mathcal{T}_q)$. So $(e_1^+)^{d-2i}W \subseteq W$. Since the mapping $(e_1^+)^{d-2i}$: $U_{d-i} \to U_i$ is a bijection, we have $\dim(W \cap U_{d-i}) \leq \dim(W \cap U_i), 0 \leq i \leq \lfloor d/2 \rfloor$.

The element $(e_1^-k_1)^{d-2i}$ does not belong to $\varphi_s(\mathcal{T}_q)$, but $(e_0^+ + s^{-2}e_1^-k_1)^{d-2i}$ does. By (7), $(e_0^+ + s^{-2}e_1^-k_1)^{d-2i}$ maps U_i to U_{d-i} , $0 \le i \le [d/2]$. We want to show it is a bijection if s avoids finitely many scalars. Identify U_{d-i} with U_i as vector spaces by the bijection $(e_1^+)^{d-2i}$ between them. Then it makes sense to consider the determinant of a linear map from U_i to U_{d-i} . Set $t = s^{-2}$ and expand $(e_0^+ + te_1^-k_1)^{d-2i}$ as

 $t^{d-2i}(e_1^-k_1)^{d-2i}$ + lower terms in t.

Each term of the expansion gives a linear map from U_i to U_{d-i} . So the determinant of $(e_0^+ + te_1^-k_1)^{d-2i}|_{U_i}$ equals

$$t^{(d-2i)\dim U_i} \det(e_1^{-}k_1)^{d-2i} \Big|_{U_i} + \text{lower terms in } t,$$
(13)

and this is a polynomial in t of degree $(d-2i) \dim U_i$, since $\det(e_1^-k_1)^{d-2i}|_{U_i} \neq 0$. Let Λ_i be the set of nonzero s such that $t = s^{-2}$ is a root of the polynomial in (13). Then if $s \in \mathbb{C}^{\times} \setminus \Lambda_i$, $(e_0^+ + s^{-2}e_1^-k_1)^{d-2i}$ maps U_i to U_{d-i} bijectively.

Set $\Lambda = \bigcup_{i=0}^{[d/2]} \Lambda_i$. Choose $s \in \mathbb{C}^{\times} \setminus \Lambda$. Then the mapping $(e_0^+ + s^{-2}e_1^-k_1)^{d-2i} : U_i \to U_{d-i}$ is a bijection for $0 \leq i \leq [d/2]$. Since $e_0^+ + s^{-2}e_1^-k_1$ belongs to $\varphi_s(\mathcal{T}_q)$, we have $(e_0^+ + s^{-2}e_1^-k_1)^{d-2i} = 0$.

 $s^{-2}e_1^-k_1)^{d-2i}W \subseteq W$ and so $\dim(W \cap U_i) \leq \dim(W \cap U_{d-i})$. Since we have already shown $\dim(W \cap U_{d-i}) \leq \dim(W \cap U_i)$, we obtain $\dim(W \cap U_i) = \dim(W \cap U_{d-i})$, $0 \leq i \leq \lfloor d/2 \rfloor$. Therefore by Lemma 1, we have $e_1^-W \subseteq W$. Since $(e_0^+ + s^{-2}e_1^-k_1)W \subseteq W$, the inclusion $e_0^+W \subseteq W$ follows from $e_1^-W \subseteq W$ and so W is $U'_q(L(\mathfrak{sl}_2))$ -invariant. Thus W = V holds by the irreducibility of V as a $U'_q(L(\mathfrak{sl}_2))$ -module.

Proof of Theorem 1. We use the classification of finite-dimensional irreducible \mathcal{T}_q -modules in the case of $(\varepsilon, \varepsilon^*) = (1, 0)$ [4, Theorem 1.18]:

- (i) A tensor product $V = V(\ell) \otimes V(\ell_1, a_1) \otimes \cdots \otimes V(\ell_n, a_n)$ of evaluation modules is irreducible as a $\varphi_s(\mathcal{T}_q)$ -module if and only if $-s^{-2} \notin S(\ell_i, a_i)$ for all $i \in \{1, \ldots, n\}$ and $S(\ell_i, a_i)$, $S(\ell_j, a_j)$ are in general position for all $i, j \in \{1, \ldots, n\}$.
- (ii) Consider two tensor products $V = V(\ell) \otimes V(\ell_1, a_1) \otimes \cdots \otimes V(\ell_n, a_n)$, $V' = V(\ell') \otimes V(\ell'_1, a'_1) \otimes \cdots \otimes V(\ell'_m, a'_m)$ of evaluation modules and assume that they are both irreducible as a $\varphi_s(\mathcal{T}_q)$ -module. Then V, V' are isomorphic as $\varphi_s(\mathcal{T}_q)$ -modules if and only if $\ell = \ell', n = m$ and $(\ell_i, a_i) = (\ell'_i, a'_i)$ for all $i \in \{1, \ldots, n\}$ with a suitable reordering of the evaluation modules $V(\ell_1, a_1), \ldots, V(\ell_n, a_n)$.
- (iii) Every finite-dimensional irreducible \mathcal{T}_q -module V, dim $V \ge 2$, is isomorphic to a \mathcal{T}_q -module $V' = V(\ell) \otimes V(\ell_1, a_1) \otimes \cdots \otimes V(\ell_n, a_n)$ on which \mathcal{T}_q acts via some embedding $\varphi_s : \mathcal{T}_q \to U'_q(L(\mathfrak{sl}_2))$.

Part (i) of Theorem 1 follows immediately from the part (i) above, due to Lemma 4. Part (ii) of Theorem 1 follows immediately from the part (ii) above, the 'if' part due to Lemma 3 (and Lemma 4) and the 'only if' part due to Lemma 4.

We want to show part (iii) of Theorem 1. Let V be a finite-dimensional irreducible $U'_q(L(\mathfrak{sl}_2))$ module of type (1, 1). By Lemma 4, there exists a nonzero scalar s such that V is irreducible as a $\varphi_s(\mathcal{T}_q)$ -module. By the part (iii) above, for the proof of which Drinfel'd polynomials play the key role, the \mathcal{T}_q -module V via φ_s is isomorphic to some \mathcal{T}_q -module $V' = V(\ell) \otimes V(\ell_1, a_1) \otimes$ $\cdots \otimes V(\ell_n, a_n)$ via some embedding $\varphi_{s'}$ of \mathcal{T}_q into $U'_q(L(\mathfrak{sl}_2))$. Since k_0 has the same eigenvalues on V, V', we have s = s' and so V, V' are isomorphic as $\varphi_s(\mathcal{T}_q)$ -modules. By Lemma 3, V, V' are isomorphic as $U'_q(L(\mathfrak{sl}_2))$ -modules. Part (iv) will be shown in the next section.

4 Intertwiners: Proof of Theorem 1(iv)

In this section, we show that for ℓ , $m \in \mathbb{Z}_{>0}$, $a \in \mathbb{C}^{\times}$, there exists an intertwiner between the $U'_q(L(\mathfrak{sl}_2))$ -modules $V(\ell, a) \otimes V(m)$, $V(m) \otimes V(\ell, a)$, i.e., a nonzero linear map R from $V(\ell, a) \otimes V(m)$ to $V(m) \otimes V(\ell, a)$ such that

$$R\Delta(\xi) = \Delta(\xi)R, \qquad \forall \xi \in U'_q(L(\mathfrak{sl}_2)), \tag{14}$$

where Δ is the coproduct from (10). If such an intertwiner R exists, then it is routinely concluded that $V(\ell, a) \otimes V(m)$ is isomorphic to $V(m) \otimes V(\ell, a)$ as $U'_q(L(\mathfrak{sl}_2))$ -modules and any other intertwiner is a scalar multiple of R, since $V(m) \otimes V(\ell, a)$ is irreducible as a $U'_q(L(\mathfrak{sl}_2))$ -module by Theorem 1.

Using the theory of Drinfel'd polynomials [4] for the augmented TD-algebra $\mathcal{T}_q = \mathcal{T}_q^{(\varepsilon,\varepsilon^*)}$ with $(\varepsilon,\varepsilon^*) = (1,0)$, we shall firstly show that $V(\ell,a) \otimes V(m)$ is isomorphic to $V(m) \otimes V(\ell,a)$ as $U'_q(L(\mathfrak{sl}_2))$ -modules. This proves Theorem 1(iv), since it is well-known [2, Theorem 4.11] that $V(\ell_i,a_i) \otimes V(\ell_j,a_j)$ and $V(\ell_j,a_j) \otimes V(\ell_i,a_i)$ are isomorphic as $U_q(\widehat{\mathfrak{sl}}_2)$ -modules, if $S(\ell_i,a_i)$ and $S(\ell_j,a_j)$ are in general position. Finally we shall construct an intertwiner explicitly.

Let us denote the $U'_{q}(L(\mathfrak{sl}_{2}))$ -modules $V(\ell, a) \otimes V(m), V(m) \otimes V(\ell, a)$ by V, V':

$$V = V(\ell, a) \otimes V(m), \qquad V' = V(m) \otimes V(\ell, a).$$

Recall we assume that the integers ℓ , m and the scalar a are nonzero. Let us denote a standard basis of the $U'_q(L(\mathfrak{sl}_2))$ -module $V(\ell, a)$ (resp. V(m)) by v_0, v_1, \ldots, v_ℓ (resp. v'_0, v'_1, \ldots, v'_m) in the sense of (9). Recall V(m) is an abbreviation of V(m, 0) and the action of $U'_q(L(\mathfrak{sl}_2))$ on V, V'are via the coproduct Δ of (10).

Let \mathcal{U} denote the subalgebra of $U'_q(L(\mathfrak{sl}_2))$ generated by e_1^{\pm} , k_1^{\pm} . The subalgebra \mathcal{U} is isomorphic to the quantum algebra $U_q(\mathfrak{sl}_2)$. Let V(n) denote the irreducible \mathcal{U} -module of dimension n+1: V(n) has a standard basis x_0, x_1, \ldots, x_n on which \mathcal{U} acts as

$$k_1 x_i = q^{n-2i} x_i, \qquad e_1^+ x_i = [n-i+1] x_{i-1}, \qquad e_1^- x_i = [i+1] x_{i+1},$$

where $[t] = [t]_q = (q^t - q^{-t})/(q - q^{-1}), x_{-1} = x_{n+1} = 0$. We call x_n (resp. x_0) the *lowest* (*highest*) weight vector: $k_1x_n = q^{-n}x_n, e_1^-x_n = 0$ ($k_1x_0 = q^nx_0, e_1^+x_0 = 0$). Note that $V(\ell, a)$ is isomorphic to $V(\ell)$ as \mathcal{U} -modules.

By the Clebsch–Gordan formula, $V = V(\ell, a) \otimes V(m)$ is decomposed into the direct sum of \mathcal{U} -submodules $\widetilde{V}(n)$, $|\ell - m| \leq n \leq \ell + m$, $n \equiv \ell + m \mod 2$, where $\widetilde{V}(n)$ is the unique irreducible \mathcal{U} -submodule of V isomorphic to V(n). With $n = \ell + m - 2\nu$, we have

$$V = V(\ell, a) \otimes V(m) = \bigoplus_{\nu=0}^{\min\{\ell, m\}} \widetilde{V}(\ell + m - 2\nu).$$
(15)

Let \widetilde{x}_n denote a lowest weight vector of the \mathcal{U} -module V(n). So

$$\Delta(k_1)\widetilde{x}_n = q^{-n}\widetilde{x}_n, \qquad \Delta(e_1^-)\widetilde{x}_n = 0.$$

Since V has a basis $\{v_{\ell-i} \otimes v'_{m-j} \mid 0 \le i \le \ell, 0 \le j \le m\}$ and k_1 acts on it by $\Delta(k_1)(v_{\ell-i} \otimes v'_{m-j}) = q^{-(\ell+m)+2(i+j)}v_{\ell-i} \otimes v'_{m-j}$, the lowest weight vector \widetilde{x}_n of $\widetilde{V}(n)$ can be expressed as

$$\widetilde{x}_n = \sum_{i+j=\nu} c_j v_{\ell-i} \otimes v'_{m-j}, \qquad n = \ell + m - 2\nu$$

Solving $\Delta(e_1^-)\tilde{x}_n = 0$ for the coefficients c_j , we obtain

$$\frac{c_j}{c_{j-1}} = -q^{m-2j+2} \frac{[\ell - \nu + j]}{[m-j+1]}$$

and so with a suitable choice of c_0

$$\widetilde{x}_n = \sum_{j=0}^{\nu} (-1)^j q^{j(m-j+1)} [\ell - \nu + j]! [m-j]! v_{\ell-\nu+j} \otimes v'_{m-j},$$
(16)

where $n = \ell + m - 2\nu$ and $[t]! = [t][t - 1] \cdots [1]$.

Lemma 5. $\Delta(e_0^+)\widetilde{x}_n = aq\widetilde{x}_{n+2}.$

Proof. By (10), we have $\Delta(e_0^+) = e_0^+ \otimes 1 + k_0 \otimes e_0^+$. By (9), the element e_0^+ vanishes on V(m) and acts on $V(\ell, a)$ as aqe_1^- . Since $e_1^- v_{\ell-\nu+j} = [\ell - (\nu - 1) + j]v_{\ell-(\nu-1)+j}$, the result follows from (16), using $v_{\ell+1} = 0$.

Corollary 1. Any nonzero $U'_q(L(\mathfrak{sl}_2))$ -submodule of $V(\ell, a) \otimes V(m)$ contains $\widetilde{x}_{\ell+m}$, the lowest weight vector of the \mathcal{U} -module $V(\ell, a) \otimes V(m)$.

We are ready to prove the following

Theorem 2. The $U'_q(L(\mathfrak{sl}_2))$ -modules $V(\ell, a) \otimes V(m)$, $V(m) \otimes V(\ell, a)$ are isomorphic for every $\ell, m \in \mathbb{Z}_{>0}, a \in \mathbb{C}^{\times}$.

Proof. Let $\mathcal{T}_q = \mathcal{T}_q^{(\varepsilon,\varepsilon^*)}$ be the augmented TD-algebra with $(\varepsilon,\varepsilon^*) = (1,0)$. Let $\varphi_s : \mathcal{T}_q \to U'_q(L(\mathfrak{sl}_2))$ denote the embedding of \mathcal{T}_q into $U'_q(L(\mathfrak{sl}_2))$ given in (5). By [4, Theorem 5.2], the Drinfel'd polynomial $P_V(\lambda)$ of the $\varphi_s(\mathcal{T}_q)$ -module $V = V(\ell, a) \otimes V(m)$ turns out to be

$$P_V(\lambda) = \lambda^m \prod_{i=0}^{\ell-1} \left(\lambda + aq^{2i-\ell+1}\right)$$

(Note that the parameter s of the embedding φ_s does not appear in $P_V(\lambda)$. So the polynomial $P_V(\lambda)$ can be called the Drinfel'd polynomial attached to the $U'_a(L(\mathfrak{sl}_2))$ -module V.)

Let W be a minimal $U'_q(L(\mathfrak{sl}_2))$ -submodule of $V = V(\ell, a) \otimes V(m)$; notice that we have shown the irreducibility of the $U'_q(L(\mathfrak{sl}_2))$ -module $V' = V(m) \otimes V(\ell, a)$ in Theorem 1 but not yet of $V = V(\ell, a) \otimes V(m)$. By Corollary 1, W contains the lowest and hence highest weight vectors of V. In particular, the irreducible $U'_q(L(\mathfrak{sl}_2))$ -module W is of type (1,1). By Lemma 4, there exists a finite set Λ of nonzero scalars such that W is irreducible as a $\varphi_s(\mathcal{T}_q)$ -module for any $s \in \mathbb{C}^{\times} \setminus \Lambda$. By the definition, the Drinfel'd polynomial $P_W(\lambda)$ of the irreducible $\varphi_s(\mathcal{T}_q)$ module W coincides with $P_V(\lambda)$:

$$P_W(\lambda) = P_V(\lambda).$$

By Theorem 1, $V' = V(m) \otimes V(\ell, a)$ is irreducible as a $U'_q(L(\mathfrak{sl}_2))$ -module. So by Lemma 4, there exists a finite set Λ' of nonzero scalars such that V' is irreducible as a $\varphi_s(\mathcal{T}_q)$ -module for any $s \in \mathbb{C}^{\times} \setminus \Lambda'$. By [4, Theorem 5.2], the Drinfel'd polynomial $P_{V'}(\lambda)$ of the irreducible $\varphi_s(\mathcal{T}_q)$ -module V' coincides with $P_V(\lambda)$:

$$P_{V'}(\lambda) = P_V(\lambda).$$

Both of the irreducible $\varphi_s(\mathcal{T}_q)$ -modules W, V' have type s, diameter $d = \ell + m$ and the Drinfel'd polynomial $P_V(\lambda)$. By [4, Theorem 1.9'], W and V' are isomorphic as $\varphi_s(\mathcal{T}_q)$ -modules. By Lemma 3, W and V' are isomorphic as $U'_q(L(\mathfrak{sl}_2))$ -modules. In particular, dim $W = \dim V'$. Since dim $V' = \dim V$, we have W = V, i.e., V and V' are isomorphic as $U'_q(L(\mathfrak{sl}_2))$ -modules.

Finally we want to construct an intertwiner R between the irreducible $U'_q(L(\mathfrak{sl}_2))$ -modules V, V'. Regard $V' = V(m) \otimes V(\ell, a)$ as a \mathcal{U} -module. By the Clebsch–Gordan formula, we have the direct sum decomposition

$$V' = V(m) \otimes V(\ell, a) = \bigoplus_{\nu=0}^{\min\{\ell, m\}} \widetilde{V}'(\ell + m - 2\nu),$$
(17)

where $\widetilde{V}'(n)$ is the unique irreducible \mathcal{U} -submodule of V' isomorphic to V(n), $n = \ell + m - 2\nu$. Let \widetilde{x}'_n be a lowest weight vector of the \mathcal{U} -module $\widetilde{V}'(n)$. By (16), we have

$$\widetilde{x}'_{n} = \sum_{j=0}^{\nu} (-1)^{j} q^{j(\ell-j+1)} [m-\nu+j]! [\ell-j]! v'_{m-\nu+j} \otimes v_{\ell-j}$$
(18)

up to a scalar multiple, where $n = \ell + m - 2\nu$. It can be easily checked as in Lemma 5 that the lowest weight vectors \widetilde{x}'_n , $n = \ell + m - 2\nu$, $0 \le \nu \le \min\{\ell, m\}$, are related by

$$(e_1^- \otimes 1)\widetilde{x}'_n = \widetilde{x}'_{n+2},\tag{19}$$

where $V' = V(m) \otimes V(\ell, a)$ is regarded as a $(\mathcal{U} \otimes \mathcal{U})$ -module in the natural way.

Lemma 6. $\Delta(e_0^+)\widetilde{x}'_n = -aq \cdot q^{n+2}\widetilde{x}'_{n+2}$

Proof. We have $\Delta(e_0^+)\widetilde{x}'_n = aq(k_1^{-1} \otimes e_1^-)\widetilde{x}'_n$, since $\Delta(e_0^+) = e_0^+ \otimes 1 + k_0 \otimes e_0^+$, and e_0^+ vanishes on V(m) and acts on $V(\ell, a)$ as aqe_1^- . Express $k_1^{-1} \otimes e_1^-$ as $k_1^{-1} \otimes e_1^- = (k_1^{-1} \otimes 1)(1 \otimes e_1^-) = (k_1^{-1} \otimes 1)(\Delta(e_1^-) - e_1^- \otimes k_1^{-1}) = (k_1^{-1} \otimes 1)\Delta(e_1^-) - k_1^{-1}e_1^- \otimes k_1^{-1} = (k_1^{-1} \otimes 1)\Delta(e_1^-) - q^2(e_1^- \otimes 1)\Delta(k_1^{-1})$. Since $\Delta(e_1^-)\widetilde{x}'_n = 0$, $\Delta(k_1^{-1})\widetilde{x}'_n = q^n\widetilde{x}'_n$, the result follows from (19).

There exists a unique linear map

$$R_n: V = V(\ell, a) \otimes V(m) \to V'(n)$$

that commutes with the action of \mathcal{U} and sends \tilde{x}_n to \tilde{x}'_n . The linear map R_n vanishes on $\tilde{V}(t)$ for $t \neq n$ and affords an isomorphism between $\tilde{V}(n)$ and V'(n) as \mathcal{U} -modules. If R is an intertwiner in the sense of (14), then R can be expressed as

$$R = \sum_{\nu=0}^{\min\{\ell,m\}} \alpha_{\nu} R_{\ell+m-2\nu},$$
(20)

regarding R as an intertwiner for the \mathcal{U} -modules V, V'. By (14), we have

$$R\Delta(e_0^+) = \Delta(e_0^+)R. \tag{21}$$

Apply (21) to the lowest weight vector \tilde{x}_n in (16). By Lemma 5, $\Delta(e_0^+)\tilde{x}_n = aq\tilde{x}_{n+2}$ and so with $n = \ell + m - 2\nu$, we have

$$R\Delta(e_0^+)\widetilde{x}_n = aq\alpha_{\nu-1}\widetilde{x}'_{n+2}.$$
(22)

On the other hand, $R\tilde{x}_n = \alpha_{\nu}\tilde{x}'_n$, $n = \ell + m - 2\nu$, and so by Lemma 6, we have

$$\Delta(e_0^+)R\widetilde{x}_n = -aq\alpha_\nu q^{n+2}\widetilde{x}'_{n+2}.$$
(23)

By (22), (23), we have $\alpha_{\nu}/\alpha_{\nu-1} = -q^{-n-2} = -q^{-\ell-m+2(\nu-1)}$ and so

$$\alpha_{\nu} = (-1)^{\nu} q^{-\nu(\ell+m-\nu+1)} \tag{24}$$

by choosing $\alpha_0 = 1$. An intertwiner exists by Theorem 2. If it exists, it has to be in the form of (20), (24). Thus we obtain the following.

Theorem 3. The linear map

. . .

$$R = \sum_{\nu=0}^{\min\{\ell,m\}} (-1)^{\nu} q^{-\nu(\ell+m-\nu+1)} R_{\ell+m-2\nu}$$

is an intertwiner between the $U'_q(L(\mathfrak{sl}_2))$ -modules $V(\ell, a) \otimes V(m)$, $V(m) \otimes V(\ell, a)$. Any other intertwiner is a scalar multiple of R.

Remark 1. Let R(a,b) be an intertwiner between the irreducible $U'_q(L(\mathfrak{sl}_2))$ -modules $V = V(\ell, a) \otimes V(m, b), V' = V(m, b) \otimes V(\ell, a)$, where $a \neq 0, b \neq 0$:

$$R(a,b): V = V(\ell,a) \otimes V(m,b) \to V' = V(m,b) \otimes V(\ell,a).$$

As in (20), we write

$$R(a,b) = \sum_{\nu=0}^{\min\{\ell,m\}} \alpha_{\nu} R_{\ell+m-2\nu}.$$

Recall R_n is the linear map from V to V' that commutes with the action of $\mathcal{U} = \langle e_1^{\pm}, k_1^{\pm} \rangle$ and sends \tilde{x}_n to \tilde{x}'_n , where $\tilde{x}_n, \tilde{x}'_n$ are the lowest weight vectors of $\tilde{V}(n), \tilde{V}'(n)$ from (15), (17) that are explicitly given by (16), (18) and satisfy $(e_1^- \otimes 1)\tilde{x}_n = \tilde{x}_{n+2}, (e_1^- \otimes 1)\tilde{x}'_n = \tilde{x}'_{n+2}$ as in (19). Since the $U'_q(L(\mathfrak{sl}_2))$ -modules V, V' can be extended to $U_q(\widehat{\mathfrak{sl}}_2)$ -modules, we have by [2, Theorem 5.4]

$$\alpha_{\nu} = \prod_{j=0}^{\nu-1} \frac{a - bq^{\ell+m-2j}}{b - aq^{\ell+m-2j}},\tag{25}$$

where we choose $\alpha_0 = 1$. Note that the denominator and the numerator of (25) are non-zero, since V, V' are assumed to be irreducible and so $S(\ell, a), S(m, b)$ are in general position.

The intertwiner R(a, b) with $a \neq 0$, $b \neq 0$ is derived from the universal *R*-matrix for the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$ [5]. If we put b = 0 in (25), then the spectral parameter u disappears, where $q^{2u} = a/b$, and we get (24). In this sense, the intertwiner R of Theorem 3 is related to the universal *R*-matrix for $U_q(\widehat{\mathfrak{sl}}_2)$, but we cannot expect that R comes from it directly, because the $U'_q(L(\mathfrak{sl}_2))$ -modules $V = V(\ell, a) \otimes V(m)$, $V' = V(m) \otimes V(\ell, a)$ cannot be extended to $U_q(\widehat{\mathfrak{sl}}_2)$ -modules. In order to derive both of the intertwiners R(a, b), R from a universal *R*-matrix directly, we need to construct it for the subalgebra $U'_q(\widehat{\mathfrak{sl}}_2)$ of $U_q(\widehat{\mathfrak{sl}}_2)$.

Acknowledgements

The research of the second author is supported by Science Foundation of Anhui University (Grant No. J10117700037).

References

- [1] Benkart G., Terwilliger P., Irreducible modules for the quantum affine algebra $U_q(\hat{sl}_2)$ and its Borel subalgebra, J. Algebra **282** (2004), 172–194, math.QA/0311152.
- [2] Chari V., Pressley A., Quantum affine algebras, Comm. Math. Phys. 142 (1991), 261–283.
- [3] Ito T., Tanabe K., Terwilliger P., Some algebra related to P- and Q-polynomial association schemes, in Codes and Association Schemes (Piscataway, NJ, 1999), DIMACS Ser. Discrete Math. Theoret. Comput. Sci., Vol. 56, Amer. Math. Soc., Providence, RI, 2001, 167–192, math.CO/0406556.
- [4] Ito T., Terwilliger P., The augmented tridiagonal algebra, Kyushu J. Math. 64 (2010), 81–144, arXiv:0904.2889.
- [5] Tolstoy V.N., Khoroshkin S.M., Universal *R*-matrix for quantized nontwisted affine Lie algebras, *Funct. Anal. Appl.* 26 (1992), 69–71.