

# Darboux transformations, finite reduction groups and related Yang-Baxter maps

S. Konstantinou-Rizos and A. V. Mikhailov

*Department of Applied Mathematics, University of Leeds, Leeds*

mmskr@leeds.ac.uk, A.V.Mikhailov@leeds.ac.uk

November 10, 2018

## Abstract

In this paper we construct Yang-Baxter (YB) maps using Darboux matrices which are invariant under the action of finite reduction groups. We present 6-dimensional YB maps corresponding to Darboux transformations for the Nonlinear Schrödinger (NLS) equation and the derivative Nonlinear Schrödinger (DNLS) equation. These YB maps can be restricted to 4-dimensional YB maps on invariant leaves. The former are completely integrable and they also have applications to a recent theory of maps preserving functions with symmetries [14]. We give a 6-dimensional YB-map corresponding to the Darboux transformation for a deformation of the DNLS equation. We also consider vector generalisations of the YB maps corresponding to the NLS and DNLS equation.

## 1 Introduction

The Yang-Baxter equation has a fundamental role in the theory of quantum and classical integrable systems. In particular, Yang-Baxter maps, namely the set theoretical solutions [12] of the Yang-Baxter equation have been of great interest for several researchers in the area of Mathematical Physics. They are related to several concepts of integrability as, for instance, the multidimensionally consistent equations [3, 4, 7, 27, 28, 29]. Especially, for those Yang-Baxter maps which admit Lax representation [34], there are corresponding hierarchies of commuting transfer maps which preserve the spectrum of their monodromy matrix [36, 37]. Therefore, the construction of Yang-Baxter maps is important.

There are several methods of construction of Yang-Baxter maps, coming from several fields of mathematics as the quantum group theory, representation theory, Lie group theory etc. Here, we construct Yang-Baxter maps using Darboux transformations for Lax operators of integrable partial differential equations. These Yang-Baxter maps can be reduced to completely itnegrable maps on invariant leaves.

The Yang-Baxter equation

$$Y^{12} \circ Y^{13} \circ Y^{23} = Y^{23} \circ Y^{13} \circ Y^{12}, \quad (1)$$

originates in the works of Yang [38] and Baxter [6]. Here  $Y^{ij}$  denotes the action of a linear operator  $Y : U \otimes U \rightarrow U \otimes U$  on the  $ij$  factor of the triple tensor product  $U \otimes U \otimes U$ , where

$U$  is a vector space. In this form, equation (1) is known in the literature as the *quantum YB equation*.

Drinfel'd in 1992 [12] proposed to replace  $U$  by an arbitrary set  $A$  and, therefore, the tensor product  $U \otimes U$  by the Cartesian product  $A \times A$ . In our paper  $A$  is an algebraic variety in  $K^N$ , where  $K$  is any field of zero characteristic, such as  $\mathbb{C}$  or  $\mathbb{Q}$ .

In [36] Veselov proposed the term *Yang Baxter map* for the set-theoretical solutions of the quantum YB equation. Specifically, we consider the map  $Y : A \times A \rightarrow A \times A$ ,

$$Y : (x, y) \mapsto (u(x, y), v(x, y)). \quad (2)$$

Furthermore, we define the functions  $Y^{i,j} : A \times A \times A \rightarrow A \times A \times A$  for  $i, j = 1, 2, 3$ ,  $i \neq j$ , which appear in equation (1), by the following relations

$$Y^{12}(x, y, z) = (u(x, y), v(x, y), z), \quad (3)$$

$$Y^{13}(x, y, z) = (u(x, z), y, v(x, z)), \quad (4)$$

$$Y^{23}(x, y, z) = (x, u(y, z), v(y, z)), \quad (5)$$

where  $x, y, z \in A$ . The variety  $A$ , in general, can be of any dimension. Thus, elements  $x \in A$  are points in  $K^N$ . The map  $Y^{ji}$ ,  $i < j$ , is defined as  $Y^{ij}$  where we swap  $u(k, l) \leftrightarrow v(l, k)$ ,  $k, l = x, y, z$ . For example,  $Y^{21}(x, y, z) = (v(y, x), u(y, x), z)$ .

The map (2) is a YB map, if it satisfies the YB equation (1). Moreover, it is called *reversible* if the composition of  $Y^{ij}$  and  $Y^{ji}$  is the identity map,

$$Y^{ij} \circ Y^{ji} = Id. \quad (6)$$

We use the term *parametric YB map* when  $u$  and  $v$  are attached with parameters  $a, b \in K^n$ , namely  $u = u(x, y; a, b)$  and  $v = v(x, y; a, b)$ , meaning that the following map

$$Y_{a,b} : (x, y; a, b) \mapsto (u(x, y; a, b), v(x, y; a, b)), \quad (7)$$

satisfies the *parametric YB equation*

$$Y_{a,b}^{12} \circ Y_{a,c}^{13} \circ Y_{b,c}^{23} = Y_{b,c}^{23} \circ Y_{a,c}^{13} \circ Y_{a,b}^{12}. \quad (8)$$

Following Suris and Veselov in [34], we call a *Lax matrix for a parametric YB map*, a matrix  $L = L(x; c; \lambda)$  depending on a variable  $x$ , a parameter  $c$  and a *spectral parameter*  $\lambda$ , such that the *Lax-equation*

$$L(u; a, \lambda)L(v; b, \lambda) = L(y; b, \lambda)L(x; a, \lambda), \quad \text{for any } \lambda \in K, \quad (9)$$

is satisfied due to the YB map. The above is also called a *refactorisation problem*.

It is obvious that the Lax-equation (9) does not always have a unique solution, which motivated Kouloukas and Papageorgiou in [20] to propose the term *strong Lax matrix* for a YB map. This is when the Lax-equation is equivalent to a YB map

$$(u, v) = Y_{a,b}(x, y). \quad (10)$$

Actually, the uniqueness of refactorisation (9) is a sufficient condition for the solutions of the Lax-equation to define a reversible YB map [18, 20, 37] of the form (10). In the opposite case, one may need to check if the obtained map satisfies the YB equation.

One of the most famous parametric YB maps is Adler's map [1]

$$(x_1, x_2) \longrightarrow (u, v) = \left( x_2 - \frac{a-b}{x_1+x_2}, x_1 + \frac{a-b}{x_1+x_2} \right), \quad (11)$$

which occurs from the 3-D consistent discrete potential KDV equation [26, 30]. In terms of Lax matrices, Adler's map (11) is obtained from the following strong Lax matrix [34, 37]

$$L(x; a, \lambda) = \begin{pmatrix} x & 1 \\ x^2 + a - \lambda & x \end{pmatrix}. \quad (12)$$

In [31, 32] a variety of YB maps is constructed using the symmetries of multi-field equations on quad graphs.

It follows from the structure of the Lax-equation (9) that we can extract *invariants* of the YB map, which we denote as  $I_i(x, y)$ . The invariants are useful if one is interested in the dynamics of such maps. In terms of dynamics, the most interesting maps are those which are not involutive. Although, involutive maps have also useful applications [15]. In all the cases presented in the next sections, our YB maps are not involutive.

In this paper we are interested in the integrability of the YB maps as finite discrete maps. The transfer dynamics of YB maps is discussed in [37].

Now, following [13, 35] we define integrability for YB maps.

**Definition 1.1.** *A  $2N$ -dimensional Yang-Baxter map,*

$$Y : (x_1, \dots, x_{2N}) \mapsto (u_1, \dots, u_{2N}), \quad u_i = u_i(x_1, \dots, x_{2N}), \quad i = 1, \dots, 2N,$$

*is said to be completely integrable or Liouville integrable if*

1. *there is a Poisson matrix  $[J]_{ij} = \{x_i, x_j\}$ , of rank  $2r$ , which is invariant under  $Y$ , namely the matrix  $[\tilde{J}]_{ij} = \{u_i, u_j\}$  has the same functional form with  $J$ ,*
2. *map  $Y$  has  $r$ -functionally independent invariants, which are in involution with respect to the corresponding Poisson bracket, i.e.  $\{I_i, I_j\} = 0$ ,  $i, j = 1, \dots, r$ ,  $i \neq j$ ,*
3. *there are  $k = 2N - 2r$  in the number Casimir functions,  $C_i$ ,  $i = 1, \dots, k$ , which are invariant under  $Y$ , namely  $C_i \circ Y = C_i$ .*

## 2 Organisation of the paper

In the next section we briefly give some introduction to the notion of the *reduction group* [24], *automorphic Lie algebras* [8, 9, 21, 22] and *Darboux transformations* to make this text self-contained. We state three cases of reduction groups which are representative for all the *finite reduction groups with degenerate orbits*[21, 22].

In Section 4 we use Darboux transformations presented in [17] to derive YB maps. In particular, we consider Darboux matrices for the NLS equation, the DNLS equation and for a deformation of the DNLS equation. For these Darboux matrices the refactorisation is not unique. Therefore, for the corresponding 6–dimensional YB maps which are derived from the refactorisation problem, in principle, one needs to check the YB property separately. Yet, the entries of these Darboux matrices obey certain differential equations which possess first integrals. There is a natural restriction of the Darboux map on the affine variety corresponding to a level set of these first integrals. These restrictions make the refactorisation unique and this guarantees that the induced 4–dimensional YB maps satisfy the YB equation and they are reversible [37]. We show that these YB maps have Poisson structure. However, the first integrals are not always very useful for the reduction because, in general, they are polynomial equations. In fact, in the case of the DNLS equation the corresponding 6-dimensional YB map cannot be reduced to a 4-dimensional one on the invariant leaves explicitly. However, after a change in the variables we obtain a reducible map. In the case of the deformation of the DNLS equation, we can only derive the 6–dimensional YB map and its implicit restriction on 4–dimensional invariant leaves.

In Section 5 we consider the vector generalisations of the Adler-Yamilov YB map and the YB map corresponding to the DNLS equation.

### 3 Automorphic Lie algebras and Darboux Transformations and reduction groups with degenerate orbits

The reduction group was first introduced in [24]. It is a discrete group of automorphisms of a Lax operator and its elements are simultaneous gauge transformations and fractional-linear transformations of the spectral parameter.

Automorphic Lie algebras were introduced in [21, 22] and studied in [8, 9, 21, 22]. These algebras constitute a subclass of infinite dimensional Lie algebras and their name is due to their construction which is very similar to the one for automorphic functions.

Darboux transformations and their relations to the theory of *integrable systems* have been extensively studied [23, 33]. Such transformations can be derived from Lax pairs as, for instance, in [33], or in a more systematic algebraic manner in [11, 17].

We are interested in Darboux transformations corresponding to Lax operators of the following form

$$\mathfrak{L} = \mathfrak{L}(\mathbf{p}(x); \lambda) = D_x + U(\mathbf{p}(x); \lambda), \quad (13)$$

where  $U$  belongs to an automorphic Lie algebra.

In the rest of the text we use “ $\mathfrak{L}$ ” for Lax operators and “ $L$ ” for Lax matrices of the refactorisation problem (9).

By Darboux transformations we understand maps

$$\mathfrak{L}(\mathbf{p}(x); \lambda) \rightarrow \tilde{\mathfrak{L}} := \mathfrak{L}(\tilde{\mathbf{p}}(x); \lambda) = M\mathfrak{L}M^{-1}, \quad (14)$$

$M$  is a matrix called the *Darboux matrix*. They map fundamental solutions,  $\Psi$ , of the equation  $\mathfrak{L}\Psi = 0$  to other fundamental solutions,  $\tilde{\Psi} = M\Psi$ , of the equation  $\tilde{\mathfrak{L}}\tilde{\Psi} = 0$ .

The structure of Lax operators has a natural Lie algebraic interpretation in terms of Kac-Moody algebras and automorphic Lie algebras [8, 9, 21, 22]. While a Kac-Moody algebra is associated with an automorphism of finite order, automorphic Lie algebras correspond to a finite group of automorphisms, namely the *reduction group* [24].

In [8, 9] it was proved that in the case of  $2 \times 2$  matrices, which we study in this paper, the essentially different reduction groups are the trivial group (with no reduction), the cyclic group  $\mathbb{Z}_2$  (leading to the Kac-Moody algebra  $A_1^1$ ) and the Klein group  $\mathbb{Z}_2 \times \mathbb{Z}_2$  [22, 25].

We shall present 4 and 6-dimensional YB maps for all the following cases. The trivial group associated with the NLS equation [39]

$$p_t = p_{xx} + 4p^2q, \quad q_t = -q_{xx} - 4pq^2. \quad (15)$$

The  $\mathbb{Z}_2$  group associated to the DNLS equation [16]

$$p_t = p_{xx} + 4(p^2q)_x, \quad q_t = -q_{xx} + 4(pq^2)_x. \quad (16)$$

and the  $\mathbb{Z}_2 \times \mathbb{Z}_2$  group associated to the deformation of the DNLS equation [25]

$$p_t = p_{xx} + 8(p^2q)_x - 4q_x, \quad q_t = -q_{xx} + 8(pq^2)_x - 4p_x. \quad (17)$$

The above mentioned groups are representative of all the *finite reduction groups* with *degenerate orbits*, namely orbits corresponding to the fixed points of the fractional linear transformations of the spectral parameter.

## 4 Derivation of YB maps

In [17] we used Darboux transformations to construct integrable systems of discrete equations, which have the multidimensional consistency property [3, 4, 7, 27, 28, 29]. The compatibility condition of Darboux transformations around the square is exactly the same with the Lax equation (9). Therefore, in this paper, we use Darboux transformations to construct YB maps.

We start with the well known example of the Darboux transformation for the nonlinear Schrödinger equation and construct its associated YB map.

### 4.1 The Nonlinear Schrödinger equation

In the case of NLS equation, the Lax operator is given by

$$\mathfrak{L}(p, q; \lambda) = D_x + \lambda U_1 + U_0, \quad \text{where } U_1 = \sigma_3 = \text{diag}(1, -1), \quad U_0 = \begin{pmatrix} 0 & 2p \\ 2q & 0 \end{pmatrix}. \quad (18)$$

The Darboux Transformation,  $M$ , of  $\mathfrak{L}$  is given by [17, 33]

$$M = \lambda \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} f & p \\ \tilde{q} & 1 \end{pmatrix}. \quad (19)$$

The entries of (19), according to definition (14), must satisfy the following system of equations

$$\partial_x f = 2(pq - \tilde{p}\tilde{q}), \quad \partial_x p = 2(pf - \tilde{p}), \quad \partial_x \tilde{q} = 2(q - \tilde{q}f),$$

which admits the following first integral

$$\partial_x(f - p\tilde{q}) = 0. \quad (20)$$

This integral implies that  $\partial_x \det M = 0$ .

In correspondence with (19), we define the matrix

$$M(\mathbf{x}, X; \lambda) = \lambda \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} X & x_1 \\ x_2 & 1 \end{pmatrix} \quad (21)$$

and substitute it into the Lax equation (9)

$$M(\mathbf{u}, U; \lambda)M(\mathbf{v}, V; \lambda) = M(\mathbf{y}, Y; \lambda)M(\mathbf{x}, X; \lambda), \quad (22)$$

to derive the following system of equations

$$\begin{aligned} v_1 &= x_1, \quad u_2 = y_2, \quad U + V = X + Y, \quad u_2v_1 = x_1y_2, \\ u_1 + Uv_1 &= y_1 + x_1Y, \quad u_1v_2 + UV = x_2y_1 + XY, \quad v_2 + u_2V = x_2 + Xy_2. \end{aligned}$$

The corresponding algebraic variety is a union of two six-dimensional components. The first one is obvious from the refactorisation problem (22), and it corresponds to the permutation map

$$\mathbf{x} \mapsto \mathbf{u} = \mathbf{y}, \quad \mathbf{y} \mapsto \mathbf{v} = \mathbf{x}, \quad X \mapsto U = Y, \quad Y \mapsto V = X,$$

which is a trivial YB map. The second one can be represented as a rational 6-dimensional non-involutive map of  $K^3 \times K^3 \rightarrow K^3 \times K^3$

$$\begin{aligned} x_1 \mapsto u_1 &= \frac{y_1 + x_1^2 x_2 - x_1 X + x_1 Y}{1 + x_1 y_2}, & y_1 \mapsto v_1 &= x_1, \\ x_2 \mapsto u_2 &= y_2, & y_2 \mapsto v_2 &= \frac{x_2 + y_1 y_2^2 + y_2 X - y_2 Y}{1 + x_1 y_2}, \\ X \mapsto U &= \frac{y_1 y_2 - x_1 x_2 + X + x_1 y_2 Y}{1 + x_1 y_2}, & Y \mapsto V &= \frac{x_1 x_2 - y_1 y_2 + x_1 y_2 X + Y}{1 + x_1 y_2}, \end{aligned} \quad (23)$$

which, one can easily check that, satisfies the YB equation.

It follows from (22) that the trace of  $M(\mathbf{y}, Y; \lambda)M(\mathbf{x}, X; \lambda)$ , is a polynomial in  $\lambda$  whose coefficients are

$$\text{Tr}(M(\mathbf{y}; b, \lambda)M(\mathbf{x}; a, \lambda)) = \lambda^2 + \lambda I_1(\mathbf{x}, \mathbf{y}, X, Y) + I_2(\mathbf{x}, \mathbf{y}, X, Y),$$

where

$$I_1(\mathbf{x}, \mathbf{y}, X, Y) = X + Y \quad \text{and} \quad I_2(\mathbf{x}, \mathbf{y}, X, Y) = x_2 y_1 + x_1 y_2 + XY, \quad (24)$$

and those are invariants for the YB map (23).

In the following section we show that the YB map (23) can be reduced to a 4– dimensional YB map which has Poisson structure.

#### 4.1.1 Restriction on invariant leaves: The Adler-Yamilov map

We now take into account the first integral (20) of the Darboux matrix (19). This first integral requires that the matrix entries  $f$ ,  $q$  and  $\tilde{q}$  are related as

$$f - p\tilde{q} = \alpha = \text{constant}.$$

Therefore, the entries of matrix (21) must obey the following equation

$$X - x_1x_2 = a, \tag{25}$$

respectively.

The matrix (21) now takes the form

$$M(\mathbf{x}; a, \lambda) = \lambda \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} a + x_1x_2 & x_1 \\ x_2 & 1 \end{pmatrix}. \tag{26}$$

In this case, the Lax equation (9),

$$M(\mathbf{u}; a, \lambda)M(\mathbf{v}; b, \lambda) = M(\mathbf{y}; b, \lambda)M(\mathbf{x}; a, \lambda), \tag{27}$$

has a unique solution  $\mathbf{u} = \mathbf{u}(\mathbf{x}, \mathbf{y})$ ,  $\mathbf{v} = \mathbf{v}(\mathbf{x}, \mathbf{y})$  which defines a map  $\mathbf{x} \rightarrow \mathbf{u}(\mathbf{x}, \mathbf{y})$ ,  $\mathbf{y} \rightarrow \mathbf{v}(\mathbf{x}, \mathbf{y})$ , given by

$$(\mathbf{x}, \mathbf{y}) \xrightarrow{Y_{a,b}} \left( y_1 - \frac{a-b}{1+x_1y_2}x_1, y_2, x_1, x_2 + \frac{a-b}{1+x_1y_2}y_2 \right). \tag{28}$$

Therefore, the above map is a reversible parametric YB map with strong Lax matrix (26). Moreover, it is not involutive.

The map (28) first appeared in the work of Adler Yamilov [5]. Its interpretation as a YB map was given in [19, 31].

From the trace of  $M(\mathbf{y}; b, \lambda)M(\mathbf{x}; a, \lambda)$  we obtain the following invariants for the map (28)

$$I_1(\mathbf{x}, \mathbf{y}) = x_1x_2 + y_1y_2 + a + b, \tag{29}$$

$$I_2(\mathbf{x}, \mathbf{y}) = (a + x_1x_2)(b + y_1y_2) + x_1y_2 + x_2y_1 + 1. \tag{30}$$

The constant terms in  $I_1, I_2$  can be omitted. It is easy to check that  $I_1, I_2$  are in involution with respect to invariant Poisson brackets defined as

$$\{x_1, x_2\} = \{y_1, y_2\} = 1, \quad \text{and all the rest} \quad \{x_i, y_j\} = 0, \tag{31}$$

and the corresponding Poisson matrix is invariant under the YB map (28). Therefore the map (28) is completely integrable.

The Adler-Yamilov map is a restriction of the YB map (23) on the invariant leaves

$$A_a = \{(x_1, x_2, X) \in \mathbb{R}^3; X = a + x_1x_2\}, \quad B_b = \{(y_1, y_2, Y) \in \mathbb{R}^3; Y = b + y_1y_2\}. \tag{32}$$

## 4.2 Derivative NLS equation: $\mathbb{Z}_2$ reduction

The Lax operator for the DNLS equation [10, 16] is given by

$$\mathfrak{L}(p, q; \lambda) = D_x + \lambda^2 U_2 + \lambda U_1, \quad \text{where} \quad U_2 = \sigma_3, \quad U_1 = \begin{pmatrix} 0 & 2p \\ 2q & 0 \end{pmatrix}, \quad (33)$$

and  $\sigma_3$  is a Pauli matrix. Operator  $\mathfrak{L}$  is invariant with respect to the following involution

$$\mathfrak{L}(\lambda) = \sigma_3 \mathfrak{L}(-\lambda) \sigma_3, \quad (34)$$

where  $\mathfrak{L}(\lambda) \equiv \mathfrak{L}(p, q; \lambda)$ . Involution (34) generates the so-called reduction group [24, 22] and it is isomorphic to  $\mathbb{Z}_2$ .

The Darboux matrix in this case is given by [17]

$$M := \lambda^2 \begin{pmatrix} f & 0 \\ 0 & 0 \end{pmatrix} + \lambda \begin{pmatrix} 0 & fp \\ f\tilde{q} & 0 \end{pmatrix} + \begin{pmatrix} c & 0 \\ 0 & 1 \end{pmatrix}, \quad (35)$$

whose entries  $p$ ,  $\tilde{q}$  and  $f$  obey the following system of equations

$$\partial_x p = 2p(\tilde{p}\tilde{q} - pq) - \frac{2}{f}(\tilde{p} - cp), \quad (36)$$

$$\partial_x \tilde{q} = 2\tilde{q}(\tilde{p}\tilde{q} - pq) - \frac{2}{f}(c\tilde{q} - q), \quad (37)$$

$$\partial_x f = 2f(pq - \tilde{p}\tilde{q}). \quad (38)$$

System (36)-(38) has a first integral which obliges the determinant of matrix (35) to be  $x$ -independent, and it is given by

$$\partial_x (f^2 p \tilde{q} - f) = 0. \quad (39)$$

Using the entries of (35) as variables, namely  $(p, \tilde{q}, f; c) \rightarrow (x_1, x_2, X; 1)$ , we define the matrix

$$M(\mathbf{x}, X; \lambda) = \lambda^2 \begin{pmatrix} X & 0 \\ 0 & 0 \end{pmatrix} + \lambda \begin{pmatrix} 0 & x_1 X \\ x_2 X & 0 \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (40)$$

The Lax equation implies the following equations

$$\begin{aligned} u_1 U + v_1 V &= x_1 X + y_1 Y, & u_2 U + v_2 V &= x_2 X + y_2 Y, \\ UV &= XY, & v_1 UV &= x_1 XY, & u_2 UV &= y_2 XY, & u_2 v_1 UV &= x_1 y_2 XY, \\ U + V + u_1 v_2 UV &= X + Y + x_2 y_1 XY. \end{aligned} \quad (41)$$

As in the case of nonlinear Schrödinger equation, the algebraic variety consists of two components. The first 6-dimensional component corresponds to the permutation map

$$\mathbf{x} \mapsto \mathbf{u} = \mathbf{y}, \quad \mathbf{y} \mapsto \mathbf{v} = \mathbf{x}, \quad X \mapsto U = Y, \quad Y \mapsto V = X, \quad (42)$$

and the second corresponds to the following 6-dimensional YB map

$$\begin{aligned} x_1 \mapsto u_1 &= f_1(\mathbf{x}, \mathbf{y}, X, Y), & y_1 \mapsto v_1 &= f_2(\pi \mathbf{y}, \pi \mathbf{x}, Y, X), \\ x_2 \mapsto u_2 &= f_2(\mathbf{x}, \mathbf{y}, X, Y), & y_2 \mapsto v_2 &= f_1(\pi \mathbf{y}, \pi \mathbf{x}, Y, X), \\ X \mapsto U &= f_3(\mathbf{x}, \mathbf{y}, X, Y), & Y \mapsto V &= f_3(\pi \mathbf{y}, \pi \mathbf{x}, Y, X), \end{aligned} \quad (43)$$



where  $\pi$  is the *permutation function*,  $\pi(x_1, x_2) = (x_2, x_1)$ ,  $\pi^2 = 1$  and  $f_1, f_2$  and  $f_3$  are given by

$$f_1(\mathbf{x}, \mathbf{y}, X, Y) = \frac{-1}{f_3(\mathbf{x}, \mathbf{y})} \frac{x_1 X + (y_1 - x_1) Y - x_1 x_2 y_1 X Y - x_1^2 x_2 X^2}{x_1 x_2 X + x_1 y_2 Y - 1}, \quad (44)$$

$$f_2(\mathbf{x}, \mathbf{y}, X, Y) = y_2, \quad (45)$$

$$f_3(\mathbf{x}, \mathbf{y}, X, Y) = \frac{x_1 x_2 X + x_1 y_2 Y - 1}{x_1 y_2 X + y_1 y_2 Y - 1} X. \quad (46)$$

One can verify that the above map is a non-involutive YB map. The invariants of this map are given by

$$I_1(\mathbf{x} \cdot \pi \mathbf{y}, X, Y) = XY \quad \text{and} \quad I_2(\mathbf{x}, \mathbf{y}, X, Y) = (\mathbf{x} \cdot \pi \mathbf{y}) XY + X + Y. \quad (47)$$

However, this map cannot be easily reduced on invariant leaves as in the case of NLS equation. This is due to the fact that the first integral of system (36)-(38) is

$$f - f^2 p \tilde{q} = k, \quad (48)$$

and therefore the relation between the entries  $x_1, x_2$  and  $X$  of (40) reads

$$X - X^2 x_1 x_2 = k. \quad (49)$$

The solution for  $X$  of the above equation can only be expressed in terms of square roots of  $x_1$  and  $x_2$ . Thus, in this case is difficult to solve the Lax equation, and if we do so the resulting map will not be presentable because of its length.

One option is to present the YB map implicitly. Specifically, the corresponding 4-dimensional map is given by

$$\begin{aligned} x_1 \mapsto u_1 &= f_1(\mathbf{x}, \mathbf{y}, X, Y), & y_1 \mapsto v_1 &= f_2(\pi \mathbf{y}, \pi \mathbf{x}, Y, X), \\ x_2 \mapsto u_2 &= f_2(\mathbf{x}, \mathbf{y}, X, Y), & y_2 \mapsto v_2 &= f_1(\pi \mathbf{y}, \pi \mathbf{x}, Y, X), \end{aligned} \quad (50)$$

where  $f_i, i = 1, 2$ , are given by (44)-(45) where  $X$  and  $Y$  are given by

$$X - X^2 x_1 x_2 = a, \quad Y - Y^2 y_1 y_2 = b. \quad (51)$$

Nevertheless, we can choose another parametrisation of matrix (35), in order to make the obtained YB map explicitly reducible on the invariant leaves.

#### 4.2.1 $\mathbb{Z}_2$ reduction: A reducible 6-dimensional YB map

Now, lets go back to the Darboux matrix (35) and replace  $(fp, f\tilde{q}, f; c) \rightarrow (x_1, x_2, X; 1)$ , namely

$$M(\mathbf{x}, X; \lambda) = \lambda^2 \begin{pmatrix} X & 0 \\ 0 & 0 \end{pmatrix} + \lambda \begin{pmatrix} 0 & x_1 \\ x_2 & 0 \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (52)$$

From the Lax equation we obtain the following equations

$$\begin{aligned} u_2 v_1 &= x_1 y_2, & u_2 V &= X y_2, & U v_1 &= x_1 Y, & UV &= XY \\ u_1 + v_1 &= x_1 + y_1, & U + u_1 v_2 + V &= X + x_2 y_1 + Y, & u_2 + v_2 &= x_2 + y_2. \end{aligned}$$

Now, the first 6-dimensional component of the algebraic variety corresponds to the trivial map (42) and the second component corresponds to a map of the form (43), with  $f_1, f_2$  and  $f_3$  now given by

$$f_1(\mathbf{x}, \mathbf{y}, X, Y) = \frac{(x_1 + y_1)X - x_1Y - x_1x_2(x_1 + y_1)}{X - x_1(x_2 + y_2)}, \quad (53)$$

$$f_2(\mathbf{x}, \mathbf{y}, X, Y) = \frac{X - x_1(x_2 + y_2)}{Y - y_2(x_1 + y_1)}y_2, \quad (54)$$

$$f_3(\mathbf{x}, \mathbf{y}, X, Y) = \frac{X - x_1(x_2 + y_2)}{Y - y_2(x_1 + y_1)}Y. \quad (55)$$

This map has the following invariants

$$I_1(\mathbf{x}, \mathbf{y}, X, Y) = XY, \quad I_2(\mathbf{x}, \mathbf{y}, X, Y) = \mathbf{x} \cdot \pi \mathbf{y} + X + Y, \quad (56)$$

$$I_3(\mathbf{x}, \mathbf{y}, X, Y) = x_1 + y_1, \quad I_4(\mathbf{x}, \mathbf{y}, X, Y) = x_2 + y_2. \quad (57)$$

#### 4.2.2 Restriction on invariant leaves

Now, taking into account the first integral

$$f - (fp)(f\tilde{q}) = k = \text{constant}, \quad (58)$$

the entries of matrix (52) must satisfy the following equation

$$X - x_1x_2 = k, \quad (59)$$

and thus matrix (52) takes the following form

$$M(\mathbf{x}; k; \lambda) = \lambda^2 \begin{pmatrix} k + x_1x_2 & 0 \\ 0 & 0 \end{pmatrix} + \lambda \begin{pmatrix} 0 & x_1 \\ x_2 & 0 \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (60)$$

The Lax-equation for matrix (60) is equivalent to the following

$$(\mathbf{x}, \mathbf{y}) \xrightarrow{Y_{a,b}} \left( y_1 + \frac{a-b}{a-x_1y_2}x_1, \frac{a-x_1y_2}{b-x_1y_2}y_2, \frac{b-x_1y_2}{a-x_1y_2}x_1, x_2 + \frac{b-a}{b-x_1y_2}y_2 \right), \quad (61)$$

and therefore the above map is a reversible parametric YB map. Moreover, it is not involutive.

The invariants of map (61) are given by

$$I_1(\mathbf{x}, \mathbf{y}) = (a + x_1x_2)(b + y_1y_2), \quad I_2(\mathbf{x}, \mathbf{y}) = (x_1 + y_1)(x_2 + y_2) + a + b. \quad (62)$$

The constant terms in  $I_1$  and  $I_2$  can be omitted. These are the invariants we retrieve from the trace of  $M(\mathbf{y}; b, \lambda)M(\mathbf{x}; a, \lambda)$ . However, the quantities  $x_1 + y_1$  and  $x_2 + y_2$  in  $I_2$  are invariants themselves. The Poisson bracket in this case is given by

$$\{x_1, x_2\} = \{y_1, y_2\} = \{x_2, y_1\} = \{y_2, x_1\} = 1, \quad \text{and all the rest } \{x_i, y_j\} = 0. \quad (63)$$

The rank of the Poisson matrix is 2,  $I_1$  is one invariant and  $I_2 = C_1C_2 + a + b$ , where  $C_1 = x_1 + y_1$  and  $C_2 = x_2 + y_2$  are Casimir functions. The latter are preserved by (61), namely  $C_i \circ Y_{a,b} = C_i$ ,  $i = 1, 2$ . Therefore, map (61) is completely integrable.

Map (61) is a restriction of the YB map (53)-(55) on the invariant leaves

$$A_a = \{(x_1, x_2, X) \in \mathbb{R}^3; X = a + x_1x_2\}, \quad B_b = \{(y_1, y_2, Y) \in \mathbb{R}^3; Y = b + y_1y_2\}. \quad (64)$$

Moreover, the map (61) can be expressed as a map of two variables on the symplectic leaf

$$x_1 + y_1 = c_1, \quad x_2 + y_2 = c_2. \quad (65)$$

### 4.3 A deformation of the DNLS equation: Dihedral Group

In the case of dihedral reduction group, the Lax operator is given by

$$\begin{aligned} \mathfrak{L}(p, q; \lambda) &= D_x + \lambda^2 U_2 + \lambda U_1 + \lambda^{-1} U_{-1} - \lambda^{-2} U_{-2}, \quad \text{where} \\ U_2 \equiv U_{-2} &= \sigma_3, \quad U_1 = \begin{pmatrix} 0 & 2p \\ 2q & 0 \end{pmatrix} \quad U_{-1} = \sigma_1 U_1 \sigma_1, \end{aligned} \quad (66)$$

and  $\sigma_1, \sigma_3$  are Pauli matrices. Here, the reduction group consists of the following set of transformations acting on the Lax operator (66) ,

$$\mathfrak{L}(\lambda) = \sigma_3 \mathfrak{L}(-\lambda) \sigma_3 \quad \text{and} \quad \mathfrak{L}(\lambda) = \sigma_1 \mathfrak{L}(\lambda^{-1}) \sigma_1, \quad (67)$$

and it is isomorphic to  $\mathbb{Z}_2 \times \mathbb{Z}_2 \cong D_2$ , [22].

In this case, the Darboux matrix is given by [17]

$$M = f \left( \begin{pmatrix} \lambda^2 & 0 \\ 0 & \lambda^{-2} \end{pmatrix} + \lambda \begin{pmatrix} 0 & p \\ \tilde{q} & 0 \end{pmatrix} + g \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{\lambda} \begin{pmatrix} 0 & \tilde{q} \\ p & 0 \end{pmatrix} \right), \quad (68)$$

where its entries obey the following equations

$$\partial_x p = 2((\tilde{p}\tilde{q} - pq)p + (p - \tilde{p})g + q - \tilde{q}), \quad (69)$$

$$\partial_x \tilde{q} = 2((\tilde{p}\tilde{q} - pq)\tilde{q} + p - \tilde{p} + (q - \tilde{q})g), \quad (70)$$

$$\partial_x g = 2((\tilde{p}\tilde{q} - pq)g + (p - \tilde{p})p + (q - \tilde{q})\tilde{q}), \quad (71)$$

$$\partial_x f = -2(\tilde{p}\tilde{q} - pq)f. \quad (72)$$

It can be shown that the above system of differential equations admits two first integrals,  $\partial_x \Phi_i = 0$ ,  $i = 1, 2$ , where

$$\Phi_1 := f^2(g - p\tilde{q}) \quad \text{and} \quad \Phi_2 := f^2(g^2 + 1 - p^2 - \tilde{q}^2). \quad (73)$$

It follows from (73), that the quantities  $f$  and  $fg$  can be expressed in terms of  $p$  and  $\tilde{q}$ , as solutions of quadratic equations. Then, the Darboux matrix depends only on two variables and we construct a 4-dimensional parametric YB map. Although, we have omitted this map because of its length.

In the next section we construct a 6-dimensional map from (68).

### 4.3.1 Dihedral group: A 6-dimensional YB map

We now consider the matrix  $N := fM$ , where  $M$  is given by (68), and we change  $(p, \tilde{q}, f^2) \rightarrow (x_1, x_2, X)$ . Then,

$$N(\mathbf{x}, X; c_1, \lambda) = \begin{pmatrix} \lambda^2 X + x_1 x_2 X + c_1 & \lambda x_1 X + \lambda^{-1} x_2 X \\ \lambda x_2 X + \lambda^{-1} x_1 X & \lambda^{-2} X + x_1 x_2 X + c_1 \end{pmatrix}, \quad (74)$$

where we have substituted the product  $f^2 g$  by

$$f^2 g = c_1 + x_1 x_2 X, \quad (75)$$

using the first integral,  $\Phi_1$ , in (73).

The Lax equation for the Darboux matrix (74) reads

$$N(\mathbf{u}, U; a, \lambda)N(\mathbf{v}, V; b, \lambda) = N(\mathbf{y}, Y; b, \lambda)N(\mathbf{x}, X; a, \lambda), \quad (76)$$

from where we obtain an algebraic system of equations, omitted because of its length.

The first 6-dimensional component of the corresponding algebraic variety corresponds to the trivial YB map

$$\mathbf{x} \mapsto \mathbf{u} = \mathbf{y}, \quad \mathbf{y} \mapsto \mathbf{v} = \mathbf{x}, \quad X \mapsto U = \frac{a}{b}Y, \quad Y \mapsto V = \frac{b}{a}X,$$

and the second component corresponds to the following map

$$\begin{aligned} x_1 \mapsto u_1 &= \frac{f(\mathbf{x}, \mathbf{y}, X, Y; a, b)}{g(\mathbf{x}, \mathbf{y}, X, Y; a, b)}, & y_1 \mapsto v_1 &= x_1 \\ x_2 \mapsto u_2 &= y_2, & y_2 \mapsto v_2 &= \frac{f(\pi\mathbf{y}, \pi\mathbf{x}, Y, X; b, a)}{g(\pi\mathbf{y}, \pi\mathbf{x}, Y, X; b, a)} \\ X \mapsto U &= \frac{g(\mathbf{x}, \mathbf{y}, X, Y; a, b)}{h(\mathbf{x}, \mathbf{y}, X, Y; a, b)}, & Y \mapsto V &= \frac{g(\pi\mathbf{y}, \pi\mathbf{x}, Y, X; b, a)}{h(\pi\mathbf{y}, \pi\mathbf{x}, Y, X; b, a)}, \end{aligned} \quad (77)$$

where  $f, g$  and  $h$  are given by

$$\begin{aligned} f(\mathbf{x}, \mathbf{y}, X, Y; a, b) &= a^2 b^2 x_1 X + a^2 b [x_2 - y_2 + 2x_1 x_2 y_1 + x_1^2 (y_2 - 3x_2)] XY + \\ & a^2 (y_2^2 - 1) [y_1 (1 + x_1^2) - x_1 (1 + y_1^2)] XY^2 - ab^2 (x_1^2 - 1) (y_2 - x_2) X^2 - \\ & ab (x_1^2 - 1) [x_2^2 (3x_1 - y_1) - x_1 - y_1 + 2y_2 (y_1 y_2 - x_1 x_2)] X^2 Y - \\ & a (x_1^2 - 1) (y_2^2 - 1) [y_2 (y_1^2 - 1) + x_2 (y_1^2 - 2x_1 y_1 + 1)] X^2 Y^2 + \\ & y_1 (x_1^2 - 1)^2 (x_2^2 - 1) (y_2^2 - 1) X^3 Y^2 + b (x_1^2 - 1)^2 (x_2^2 - 1) (y_2 - x_2) X^3 Y + \\ & a^3 b (y_1 - x_1) Y, \\ g(\mathbf{x}, \mathbf{y}, X, Y; a, b) &= a^2 b^2 X + 2a^2 b y_2 (y_1 - x_1) XY + a^2 (y_2^2 - 1) (x_1 - y_1)^2 XY^2 + \\ & 2ab (x_1^2 - 1) (1 - x_2 y_2) X^2 Y + 2a x_2 (x_1^2 - 1) (y_2^2 - 1) (x_1 - y_1) X^2 Y^2 + \\ & (x_1^2 - 1)^2 (x_2^2 - 1) (y_2^2 - 1) X^3 Y^2, \\ h(\mathbf{x}, \mathbf{y}, X, Y; a, b) &= a^2 b^2 - 2ab^2 x_1 (y_2 - x_2) X - 2ab (x_1 y_1 - 1) (y_2^2 - 1) XY \\ & b^2 (x_1^2 - 1) (x_2 - y_2)^2 X^2 - 2b y_1 (x_2 - y_2) (x_1^2 - 1) (y_2^2 - 1) X^2 Y + \\ & (x_1^2 - 1) (y_1^2 - 1) (y_2^2 - 1)^2 X^2 Y^2. \end{aligned} \quad (78)$$

It can be verified that this is a parametric YB map. From  $\text{Tr}(N(\mathbf{x}, X; \lambda)N(\mathbf{y}, Y; \lambda))$  we extract the following invariants for the above map

$$I_1(\mathbf{x}, \mathbf{y}, X, Y; a, b) = XY, \quad (79)$$

$$I_2(\mathbf{x}, \mathbf{y}, X, Y; a, b) = bX + aY + (x_1 + y_1)(x_2 + y_2)XY, \quad (80)$$

$$I_3(\mathbf{x}, \mathbf{y}, X, Y; a, b) = 2bx_1x_2X + 2ay_1y_2Y + 2(\mathbf{x} \cdot \mathbf{y} + x_1x_2y_1y_2)XY + 2ab. \quad (81)$$

### 4.3.2 Restriction to invariant leaves: An implicit map

Using the first integrals  $\Phi_1$  and  $\Phi_2$  we can reduce the above 6-dimensional map to a 4-dimensional YB map implicitly. In particular, from equations  $\Phi_1 = c_1$  and  $\Phi_2 = c_2$  one can obtain

$$(1 - x_1^2 - x_2^2 + x_1^2x_2^2)X^2 + (2x_1x_2 - c_2)X + 1 = 0, \quad (82)$$

where we have rescaled  $c_1 \rightarrow 1$ .

Therefore the 4-dimensional map is given by

$$(u_1, u_2, v_1, v_2) = \left( \frac{f(\mathbf{x}, \mathbf{y}, X, Y; a, b)}{g(\mathbf{x}, \mathbf{y}, X, Y; a, b)}, y_2, x_1, \frac{f(\pi\mathbf{y}, \pi\mathbf{x}, X, Y; b, a)}{g(\pi\mathbf{y}, \pi\mathbf{x}, X, Y; b, a)} \right), \quad (83)$$

where  $f$ ,  $g$  and  $h$  are given by the above relations and  $X$  and  $Y$  are given by

$$(1 - x_1^2 - x_2^2 + x_1^2x_2^2)X^2 + (2x_1x_2 - a)X + 1 = 0, \quad (84)$$

$$(1 - y_1^2 - y_2^2 + y_1^2y_2^2)Y^2 + (2y_1y_2 - b)Y + 1 = 0. \quad (85)$$

## 4.4 Dihedral group: A linearised YB map

We replace  $(f\tilde{q}, fp) \rightarrow (x_1, x_2)$  in the Darboux matrix (68) to become

$$M(\mathbf{x}; k, \lambda) = \begin{pmatrix} \lambda^2 f & 0 \\ 0 & \lambda^{-2} f \end{pmatrix} + \lambda \begin{pmatrix} 0 & x_1 \\ x_2 & 0 \end{pmatrix} + fg \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{\lambda} \begin{pmatrix} 0 & x_2 \\ x_1 & 0 \end{pmatrix}. \quad (86)$$

Then equations  $\Phi_1 = c_1$  and  $\Phi_2 = c_2$  imply that the quantities  $f$  and  $fg$  are given by

$$f = \frac{1}{2}\sqrt{1 + (x_1 + x_2)^2} + \frac{1}{2}\sqrt{k^2 + (x_1 - x_2)^2}, \quad (87)$$

$$fg = \frac{1}{2}\sqrt{1 + (x_1 + x_2)^2} - \frac{1}{2}\sqrt{k^2 + (x_1 - x_2)^2}, \quad (88)$$

where  $(c_1, c_2) \rightarrow (\frac{1-k^2}{4}, \frac{1+k^2}{2})$ .

The linear approximation to the YB map is given by

$$\begin{pmatrix} x_1 \\ x_2 \\ y_1 \\ y_2 \end{pmatrix} \xrightarrow{U_0} \begin{pmatrix} u_1 \\ u_2 \\ v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} \frac{(a-1)(a-b)}{(a+1)(a+b)} & \frac{a-b}{a+b} & \frac{2a}{a+b} & \frac{(a+1)(b-a)}{(b+1)(a+b)} \\ 0 & 0 & 0 & \frac{a+1}{b+1} \\ \frac{b+1}{a+1} & 0 & 0 & 0 \\ \frac{(a-b)(b+1)}{(a+1)(a+b)} & \frac{2b}{a+b} & \frac{b-a}{a+b} & \frac{(b-1)(b-a)}{(b+1)(a+b)} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ y_1 \\ y_2 \end{pmatrix} \quad (89)$$

which is a linear parametric YB map and it is not involutive.

## 5 $2N \times 2N$ -dimensional YB maps

In this section we consider the vector generalisations of the YB maps (28) and (61). We replace the variables,  $x_1$  and  $x_2$ , in the Lax matrices with  $N$ -vectors  $\mathbf{w}_1$  and  $\mathbf{w}_2^T$  to obtain  $2N \times 2N$  YB maps. In what follows we use the following notation for a  $n$ -vector  $\mathbf{w} = (w_1, \dots, w_n)$

$$\mathbf{w} = (\mathbf{w}_1, \mathbf{w}_2), \quad \text{where} \quad \mathbf{w}_1 = (w_1, \dots, w_N), \quad \mathbf{w}_2 = (w_{N+1}, \dots, w_{2N}) \quad (90)$$

and also

$$\langle u_i | := \mathbf{u}_i, \quad |w_i\rangle := \mathbf{w}_i^T \quad \text{and their dot product with} \quad \langle u_i, w_i\rangle. \quad (91)$$

### 5.1 NLS equation

Replacing the variables in (26) with  $N$ -vectors, namely

$$M(\mathbf{w}; a, \lambda) = \begin{pmatrix} \lambda + a + \langle w_1, w_2\rangle & \langle w_1 | \\ |w_2\rangle & I \end{pmatrix}, \quad (92)$$

we obtain a unique solution of the Lax-Equation given by the following  $2N \times 2N$  map

$$\begin{cases} \langle u_1 | = \langle y_1 | + f(z; a, b)\langle x_1 |, \\ \langle u_2 | = \langle y_2 |, \end{cases} \quad (93)$$

and

$$\begin{cases} \langle v_1 | = \langle x_1 |, \\ \langle v_2 | = \langle x_2 | + f(z; b, a)\langle y_2 |, \end{cases} \quad (94)$$

where  $f$  is given by

$$f(z; b, a) = \frac{b-a}{1+z}, \quad z := \langle x_1, y_2\rangle. \quad (95)$$

The above is a non-involutive parametric  $2N \times 2N$  YB map with strong Lax matrix given by (92). As a YB map it appears in [31], but it is originally introduced by Adler [2]. Moreover, one can construct the above  $2N \times 2N$  map for the  $N \times N$  Darboux matrix (92) by taking the limit of the solution of the refactorisation problem in [20].

Two invariants of this map are given by

$$I_1(\mathbf{x}, \mathbf{y}; a, b) = \langle x_1, x_2\rangle + \langle y_1, y_2\rangle, \quad (96)$$

$$I_2(\mathbf{x}, \mathbf{y}; a, b) = b\langle x_1, x_2\rangle + a\langle y_1, y_2\rangle + \langle x_1, y_2\rangle + \langle x_2, y_1\rangle + \langle x_1, x_2\rangle\langle y_1, y_2\rangle. \quad (97)$$

These are the invariants which are obtained from the trace of  $M(\mathbf{y}; b, \lambda)M(\mathbf{x}; a, \lambda)$  and they are not enough to prove Liouville integrability.

## 5.2 $\mathbb{Z}_2$ reduction

In the case of  $Z_2$  we consider, instead of (60), the following matrix

$$M(\mathbf{w}; a, \lambda) = \begin{pmatrix} \lambda^2(a + \langle w_1, w_2 \rangle) & \lambda \langle w_1 | \\ \lambda | w_2 \rangle & I \end{pmatrix}, \quad (98)$$

we obtain a unique solution for the Lax-Equation given by the following  $2N \times 2N$  map

$$\begin{cases} \langle u_1 | = \langle y_1 | + f(z; a, b) \langle x_1 |, \\ \langle u_2 | = g(z; a, b) \langle y_2 |, \end{cases} \quad (99)$$

and

$$\begin{cases} \langle v_1 | = g(z; b, a) \langle x_1 |, \\ \langle v_2 | = \langle x_2 | + f(z; b, a) \langle y_2 |, \end{cases} \quad (100)$$

where  $f$  and  $g$  are given by

$$f(z; a, b) = \frac{a-b}{a-z}, \quad g(z; a, b) = \frac{a-z}{b-z}, \quad z := \langle x_1, y_2 \rangle. \quad (101)$$

The above map is a non-involutive parametric  $2N \times 2N$  YB map with strong Lax matrix given by (98).

The invariants of the above map, coming from the trace of  $M(\mathbf{y}; b, \lambda)M(\mathbf{x}; a, \lambda)$ , are given by

$$I_1(\mathbf{x}, \mathbf{y}; a, b) = \langle x_1 + y_1, x_2 + y_2 \rangle, \quad (102)$$

$$I_2(\mathbf{x}, \mathbf{y}; a, b) = b \langle x_1, x_2 \rangle + a \langle y_1, y_2 \rangle + \langle x_1, x_2 \rangle \langle y_1, y_2 \rangle. \quad (103)$$

In fact, both vectors of the inner product in  $I_1$  are invariants. However, as in the vector generalisation of the Adler-Yamilov map, the invariants are not enough to claim Liouville integrability.

The Liouville integrability of the vector generalisations of the YB maps we consider in this section is an open problem.

## Acknowledgements

We would like to thank A. Veselov, T. Kouloukas, V. Papageorgiou, A. Dzhamay and G. Grahovski for the discussions and their comments, P. Xenitidis for helping to improve the text and special thanks to D. Tsubelis for making available his computing facilities. A.V.M. would like to acknowledge support from EPSRC (EP/I038675/1). S.K.R. would like to acknowledge William Right Smith scholarship and John E. Crowther scholarship.

## References

## References

- [1] ADLER V 1993 Recuttings of polygons *Funktsional. Anal. i Prilozhen.* **27** 79–82.
- [2] ADLER V 1994 Nonlinear superposition principle for the Jordan NLS equation *Phys. Lett. A* **190** 53–58.
- [3] ADLER V, BOBENKO A, AND SURIS Y 2003 Classification of integrable equations on quad-graphs. The consistency approach *Comm. Math. Phys.* **233** 513–543.
- [4] ADLER V, BOBENKO A, AND SURIS Y 2004 Geometry of Yang-Baxter maps: pencils of conics and quadrirational mappings *Comm. Anal. Geom.* **12** 967–1007.
- [5] ADLER V AND YAMILOV R 1994 Explicit auto-transformations of integrable chains *J. Phys. A* **27** 477–492.
- [6] BAXTER R 1972 Partition function of the eight-vertex lattice model *Ann. Physics* **70** 193–228.
- [7] BOBENKO A AND SURIS Y. 2002 Integrable systems on quad-graphs *Int. Math. Res.* **11** 573–611.
- [8] BURY R 2010 Automorphic Lie algebras, corresponding integrable systems and their soliton solutions *Ph.D. thesis, Un. of Leeds*.
- [9] BURY R AND MIKHAILOV A 2012 Automorphic Lie algebras and corresponding integrable systems. I. to be submitted
- [10] CAI H, LIU F, AND HUANG N 2003 Hamiltonian formalism of the derivative nonlinear Schrödinger equation *Commun. Theor. Phys.* **39** 181–188.
- [11] CIEŚLIŃSKI J 1995 An algebraic method to construct the Darboux matrix *J. Math. Phys.* **36** 5670–5706.
- [12] DRINFEL'D V 1992 On some unsolved problems in quantum group theory *Lecture Notes in Math.* **1510** 1–8.
- [13] FORDY A 2012 Integrable Poisson maps from cluster exchange relations *Lecture notes - SIDE summer school, Ningbo*.
- [14] FORDY A AND KASSOTAKIS A 2013 Integrable Maps which Preserve Functions with Symmetries *J. Phys. A* **46** 205201.
- [15] KASSOTAKIS P AND NIESZPORSKI M 2012 On non-multiaffine consistent-around-the-cube lattice equations *Phys. Lett. A* **376** 3135–3140.
- [16] KAUP D AND NEWELL A 1978 An exact solution for a derivative nonlinear Schrödinger equation *J. Mathematical Phys.* **19** 798–801.



- [17] KONSTANTINOU-RIZOS S, MIKHAILOV A, AND XENITIDIS P 2012 Reduction groups and integrable difference systems of NLS type *To be submitted*.
- [18] KOULOUKAS T 2010 Yang-Baxter maps, poisson structure and integrability *Ph.D thesis, Un. of Patras, Greece*.
- [19] KOULOUKAS T AND PAPAGEORGIU V 2009 Yang-Baxter maps with first-degree-polynomial  $2 \times 2$  Lax matrices *J. Phys. A* **42** 404012.
- [20] KOULOUKAS T AND PAPAGEORGIU V 2011 Poisson yang-baxter maps with binomial lax matrices *J. Math. Phys.* **52** 073502.
- [21] LOMBARDO S 2004 Reductions of integrable equations and automorphic Lie algebras *Ph.D. thesis, Un. of Leeds*.
- [22] LOMBARDO S AND MIKHAILOV A 2005 Reduction groups and automorphic Lie algebras *Comm. Math. Phys.* **258** 179–202.
- [23] MATVEEV V AND SALLE M 1991 Darboux transformations and solitons *Integrable systems in statistical mechanics, Springer series in nonlinear dynamics*.
- [24] MIKHAĬLOV A 1981 The reduction problem and the inverse scattering method *Physica D* **3** 73-117.
- [25] MIKHAĬLOV A, SHABAT A, AND YAMILOV R 1988 Extension of the module of invertible transformations. Classification of integrable systems *Comm. Math. Phys.* **115** 1–19.
- [26] NIJHOFF F AND CAPEL H 1995 The discrete Korteweg-de Vries equation *Acta Appl. Math.* **39** 133–158.
- [27] NIJHOFF F 2002 Lax pair for the Adler (lattice Krichever-Novikov) system *Phys. Lett. A* **297** 49–58.
- [28] NIJHOFF F 2010 Discrete systems and integrability *Academic lecture notes*.
- [29] NIJHOFF F AND WALKER A 2001 The discrete and continuous Painlevé VI hierarchy and the Garnier systems *Glasg. Math. J.* **43A** 109–123.
- [30] PAPAGEORGIU V, NIJHOFF F, AND CAPEL H 1990 Integrable mappings and nonlinear integrable lattice equations *Phys. Lett. A* **147** 106–114.
- [31] PAPAGEORGIU V AND TONGAS A 2007 Yang-Baxter maps and multi-field integrable lattice equations *J. Phys. A* **40** 12677–12690.
- [32] PAPAGEORGIU V, TONGAS A, AND VESELOV A 2006 Yang-Baxter maps and symmetries of integrable equations on quad-graphs *J. Math. Phys.* **47** 083502, 16.
- [33] ROGERS C AND SCHIEF W 2002 Bäcklund and Darboux transformations. Geometry and modern applications in soliton theory, *Cambridge texts in applied mathematics*.

- [34] SURIS Y AND VESELOV A 2003 Lax matrices for Yang-Baxter maps *J. Nonlinear Math. Phys.* **10** 223–230.
- [35] VESELOV A 1991 Integrable maps *Russ. Math. Surveys* **46** 1–51.
- [36] VESELOV A 2003 Yang-Baxter maps and integrable dynamics *Phys. Lett. A* **314** 214–221.
- [37] VESELOV A 2007 Yang-Baxter maps: dynamical point of view *J. Math. Soc. Japan* **17** 145–167.
- [38] YANG C 1967 Some exact results for the many-body problem in one dimension with repulsive delta-function interaction *Phys. Rev. Lett.* **19** 1312–1315.
- [39] ZAHAROV V AND ŠABAT A 1979 Integration of the nonlinear equations of mathematical physics by the method of the inverse scattering problem. II *Funktsional. Anal. i Prilozhen.* **13** 13–22.