On peaked solitary waves of the Degasperis - Procesi equation

Shijun Liao

State Key Laboratory of Ocean Engineering, Dept. of Mathematics School of Naval Architecture, Ocean and Civil Engineering Shanghai Jiao Tong University, Shanghai 200240, China

(Email address: sjliao@sjtu.edu.cn)

Abstract The Degasperis - Process (DP) equation describing the propagation of shallow water waves contains a physical parameter ω , and it is well-known that the DP equation admits solitary waves with a peaked crest when $\omega = 0$. In this article, we illustrate, for the first time, that the DP equation admits peaked solitary waves even when $\omega \neq 0$. This is helpful to enrich our knowledge and deepen our understandings about peaked solitary waves of the DP equation.

PACS Number: 47.35.Bb

Key Words Peaked solitary waves, discontinuity, Degasperis - Processi equation

1 Introduction

The solitary surface wave was first discovered by Russell [17] in 1834. Since then, many models for shallow water waves have been developed, such as the Boussinesq equation [3], the Korteweg & de Vries (KdV) equation [8] and the Benjamin-Bona-Mahony (BBM) equation [1], and so on. All of them admit solitary waves with smooth crest. In contrast to the KdV, Boussinesq and BBM equations, the celebrated Camassa-Holm (CH) equation [4]

$$u_t + 2\omega u_x - u_{xxt} + 3uu_x = 2u_x u_{xx} + uu_{xxx} \tag{1}$$

can model both phenomena of soliton interaction and wave breaking (see [5]), where u(x,t) denotes the wave elevation, x, t are the temporal and spatial variables, ω is a constant related to the critical shallow water wave speed, and the subscript denotes the partial differentiation, respectively. The CH equation (1) is integrable and bi-Hamiltonian. Especially, when $\omega = 0$, the CH equation (1) has the peaked solitary wave [4]

$$u(x,t) = c \exp(-|x - ct|)$$

Degasperis and Procesi [7] investigated a generalized equation

$$u_t + 2\omega u_x - u_{xxt} + (\beta + 1)uu_x = \beta u_x u_{xx} + uu_{xxx}$$

$$\tag{2}$$

and found that it is completely integrable only when $\beta = 2$, corresponding to the CH equation (1), and $\beta = 3$, corresponding to the so-called Degasperis - Procesi (DP) equation

$$u_t + 2\omega u_x - u_{xxt} + 4uu_x = 3u_x u_{xx} + uu_{xxx},\tag{3}$$

respectively. The DP equation (3) can model the propagation of shallow water waves with small amplitude and large wave-length. It is interesting that, like the CH equation (1), the DP equation (3) also admits solitary waves with a peaked crest when $\omega = 0$.

Currently, Liao [13] gained the closed-form solutions of the peaked solitary waves of the KdV equation, the modified KdV equation, the Boussinesq equation and the BBM equation. Besides, by means of the Mathematica package BVPh 1.0 that is based on the homotopy analysis method (HAM) [9–12], a powerful analytic method for highly nonlinear equations, Liao [14] found that the CH equation (1) also admits peaked solitary waves even in the case of $\omega \neq 0$. All of these indicate that most of mainstream models of shallow water waves admit peaked solitary waves.

In this article, using the HAM-based Mathematica package BVPh 1.0 for nonlinear boundary value problems in a similar way, we illustrate that the DP equation (3) also admits peaked solitary waves even when $\omega \neq 0$. In addition, we further illustrate that Eq. (2) for arbitrary constant β and the more generalized equation

$$u_t + 2\omega u_x - u_{xxt} + \alpha u u_x = \beta u_x u_{xx} + u u_{xxx} \tag{4}$$

for arbitrary constants α and β also admit peaked solitary waves even when $\omega \neq 0$, although they are not integrable in general cases.

2 Peaked solitary waves of Eq. (4) in the case of $\omega \neq 0$

Let us first consider the propagating solitary waves of the DP equation (3) with permanent form, corresponding to $\alpha = 4$ and $\beta = 3$ of Eq. (4). To avoid the repeat, Eq. (4) is used to describe the basic ideas of our analytic approach. Writing $\xi = x - c t$ and $w(\xi) = c u(x, t)$, the original equation (4) becomes

$$w''' - \left(1 - \frac{2\omega}{c}\right)w' + \alpha ww' = \beta w'w'' + ww''',\tag{5}$$

subject to the boundary conditions

$$w \to 0, w' \to 0, w'' \to 0, \text{ as } |\xi| \to +\infty,$$
 (6)

where the prime denotes the differentiation.

The corresponding linearized equation

$$w''' - \left(1 - \frac{2\omega}{c}\right)w' = 0\tag{7}$$

has the peaked solitary wave

$$w(\xi) = A \exp(-\mu|\xi|), \ \mu = \sqrt{1 - \frac{2\omega}{c}},$$
 (8)

provided $\omega < c/2$, where A = w(0). The solution of Eq. (5) can be expressed in the form

$$w(\xi) = \sum_{n=1}^{+\infty} a_n \exp(-n\mu|\xi|),$$

where a_n is a constant to be determined. Using the homotopy analysis method (HAM) [9–12], an analytic technique for highly nonlinear differential equations, we gain the series solution

$$w(\xi) = w_0(\xi) + \sum_{m=0}^{+\infty} w_m(\xi).$$
(9)

Here

$$w_0(\xi) = A \exp(-\mu|\xi|)$$

is the initial guess, w_m for $m \ge 1$ is governed by

$$\mathcal{L}[w_m(\xi) - \chi_m \ w_{m-1}(\xi)] = c_0 \ \delta_{m-1}(\xi), \tag{10}$$

subject to the boundary condition

$$w_m(0) = 0, \ w_m \to 0, \ \text{as } |\xi| \to +\infty,$$
(11)

where

$$\mathcal{L}f = f''' - \mu^2 f'$$

is an auxiliary linear operator, and

$$\delta_n = w_n''' - \mu^2 w_n' + \sum_{j=0}^n \left[\beta \ w_j \ w_{n-j}' - \alpha \ w_j' \ w_{n-j}'' - w_j \ w_{n-j}''' \right], \tag{12}$$

$$\chi_n = \begin{cases} 1 & \text{when } n > 1, \\ 0 & \text{otherwise.} \end{cases}$$
(13)

Note that $c_0 \neq 0$ is an auxiliary parameter, called the convergence-control parameter, which provides us a convenient way to guarantee the convergence of the approximation series. For details, please refer to Liao [9, 11, 12]. In fact, directly using the HAM-based Mathematica package BVPh 1.0 (see Part II of Liao's book [12]) for nonlinear boundary-value/eigenvalue problems, it is straightforward to gain high-order analytic approximations of (5) and (6). For details, please refer to the Appendix.

The accuracy of the *m*th-order approximation of $w(\xi)$ is defined by the averaged residual square of the governing equation (5) in the domain $\xi \in [0, a]$:

$$\mathcal{E}_m = \frac{1}{a} \int_0^a \left[\mathcal{N}\left(\sum_{j=0}^m w_j\right) \right]^2 d\xi, \tag{14}$$

Table 1: $u'(0_+)$ of the *m*th-order analytic approximations and the residual squares of the DP equation (3) in the case of $c = 1, \omega = 1/4$, and $A = \pm 1/10$ given by the HAM-based Mathematica package BVPh 1.0 with the convergence-control parameter $c_0 = -1$.

| | A = 1/10 | | A = -1/10 | |
|----|-------------|-----------------|-------------|-----------------|
| m | $u'(0_{+})$ | \mathcal{E}_m | $u'(0_{+})$ | \mathcal{E}_m |
| 2 | -0.065250 | 3.8 E-9 | 0.074675 | 4.8E-9 |
| 4 | -0.065110 | 2.0 E-12 | 0.074779 | 1.6E-12 |
| 6 | -0.065105 | 9.2 E-15 | 0.074782 | 5.9E-15 |
| 8 | -0.065105 | 2.4 E-17 | 0.074782 | 1.6E-17 |
| 10 | -0.065105 | 5.2 E-20 | 0.074782 | 3.5E-20 |

where

$$\mathcal{N}w = w''' - \mu^2 w' + \alpha \, ww' - \beta \, w'w'' - ww'''.$$

We choose a = 10 in this article, because the wave elevation decays exponentially.

For simplicity, we investigate the case c = 1 in this article. Without loss of generality, we first consider the case of $\omega = 1/4$ and $A = \pm 1/10$ for the DP equation (3). As shown in Table 1, the averaged residual squares of the 10th-order analytic approximations decrease to 5.2×10^{-20} in the case of A = 1/10 and to 3.5×10^{-20} in the case of A = -1/10, respectively. Besides, the corresponding values of $u'(0_+)$ (the limit is taken as $\xi \to 0$ from the right) quickly converge to -0.065105 and 0.074782, respectively. All of these clearly indicate that the series given by the Mathematica package BVPh 1.0 converge to the solution of the DP equation (3). The solutions for larger values of |A| can be gained in a similar way. All of these solitary waves have a peaked crest, as shown in Fig. 1. These clearly illustrate that, like the CH equation (1), the DP equation (3) also admits peaked solitary waves even when $\omega \neq 0$. Besides, like the CH equation (1), the phase speed of the peaked solitary waves of the DP equation (3) has nothing to do with the wave amplitude A. This is quite interesting.

As pointed out by Degasperis and Procesi [7], the generalized equation (2) is completely integrable only when $\beta = 2$ or $\beta = 3$, corresponding to the CH equation (1) or the DP equation (3), respectively. Does Eq. (2) admit peaked solitary waves when $\beta \neq 2$ and $\beta \neq 3$?

Without loss of generality, let us consider the case of $c = 1, \omega = 1/4, \beta = 5/2$ and $A = \pm 1/10$ for Eq. (2), corresponding to $\alpha = 7/2, \beta = 5/2$ for Eq. (4). Similarly, by means of the HAM-based Mathematica package BVPh 1.0, we gain the convergent approximations with $c_0 = -1$. As shown in Table 2, the averaged residual squares of the 10th-order analytic approximations decrease to 4.4×10^{-21} in the case of A = 1/10 and to 3.2×10^{-21} in the case of A = -1/10, respectively. Similarly, using the Mathematica package BVPh 1.0, we gain the convergent approximations for larger values of A, as shown in Fig. 2. Note that all of these solitary waves have a peaked



Figure 1: The peaked solitary waves u(x,t) of the DP equation (3) in the case of $\omega = 1/4$ and c = 1. Solid line: $A = \pm 3/10$; Dashed line: $A = \pm 1/5$; Dash-dotted line: $A = \pm 1/10$.

crest! Thus, the generalized equation (2) in the case of $\omega \neq 0$ also admits peaked solitary waves, even when $\beta \neq 2$ and $\beta \neq 3$ so that it is *not* integrable!

How about the more generalized equation (4)?

Without loss of generality, let us consider the generalized equation (4) in the case of $c = 1, \omega = 1/4, \alpha = 11/3, \beta = 5/2$ and $A = \pm 1/10$. Similarly, by means of the HAM-based Mathematica package BVPh 1.0, we gain the convergent approximations with $c_0 = -1$. As shown in Table 3, the averaged residual squares of the 10thorder analytic approximations decrease to 1.1×10^{-20} in the case of A = 1/10 and to 7.8×10^{-21} in the case of A = -1/10, respectively. Similarly, using the Mathematica package BVPh 1.0, we gain the convergent approximations for larger values of |A|, as shown in Fig. 3. It should be emphasized that all of these solitary waves have a peaked crest. These illustrate that the generalized equation (4) also admits peaked solitary waves for arbitrary constants α and β even when $\omega \neq 0$, although the corresponding equation is *not* integrable in general cases. Therefore, the existence of peaked solitary waves might be a common property of the generalized equation (4).

As pointed out by Constantin and Molinet [6], all of these peaked solitary waves should be understood mathematically as weak solutions of the generalized equation (4). However, physically, this kind of discontinuity of wave elevation widely appears in fluid mechanics, such as dam break [19] in hydrodynamics and shock waves in aerodynamics, which have clear physical meanings. Note also that such kind of discontinuous problems belong to the so-called Riemann problem [2,16,18,19], a classical field of fluid mechanics.

Table 2: $u'(0_+)$ of the *m*th-order analytic approximations and the residual squares of Eq. (2) in the case of $c = 1, \omega = 1/4, A = \pm 1/10$ and $\beta = 5/2$ given by the HAM-based Mathematica package BVPh 1.0 with the convergence-control parameter $c_0 = -1$.

| | A = 1/10 | | A = -1/10 | |
|----|-------------|-----------------|-----------|-----------------|
| m | $u'(0_{+})$ | \mathcal{E}_m | $u'(0_+)$ | \mathcal{E}_m |
| 2 | -0.066002 | 2.1 E-9 | 0.074251 | 2.6E-9 |
| 4 | -0.065904 | 5.7 E-13 | 0.074323 | 5.3E-13 |
| 6 | -0.065901 | 1.6 E-15 | 0.074325 | 1.1E-15 |
| 8 | -0.065901 | 3.0 E-18 | 0.074325 | 2.2E-18 |
| 10 | -0.065901 | 4.4 E-21 | 0.074325 | 3.2E-21 |

Table 3: $u'(0_+)$ of the *m*th-order analytic approximations and the residual squares of the generalized equation (4) in the case of $c = 1, \omega = 1/4, A = \pm 1/10, \alpha = 11/3$ and $\beta = 5/2$ given by the HAM-based Mathematica package BVPh 1.0 with the convergence-control parameter $c_0 = -1$.

| | A = 1/10 | | A = -1/10 | |
|----|-----------|-----------------|-------------|-----------------|
| m | $u'(0_+)$ | \mathcal{E}_m | $u'(0_{+})$ | \mathcal{E}_m |
| 2 | -0.065541 | 2.3 E-9 | 0.074576 | 2.9E-9 |
| 4 | -0.065428 | 9.8 E-13 | 0.074658 | 7.6E-13 |
| 6 | -0.065425 | 3.3 E-15 | 0.074660 | 2.2E-15 |
| 8 | -0.065425 | 6.6 E-18 | 0.074660 | 4.6E-18 |
| 10 | -0.065425 | 1.1 E-20 | 0.074660 | 7.8E-21 |

3 Concluding remarks

In this paper, using the HAM-based Mathematica package BVPh 1.0 as a useful tool, we illustrate that, like the CH equation (1), the DP equation (3) also admits peaked solitary waves even when $\omega \neq 0$. Furthermore, we illustrate in a similar way that the generalized equations (2) and (4) also admit peaked solitary waves even when $\omega \neq 0$, and the corresponding equation is *not* integrable! As pointed out by Liao [13], nearly all mainstream models of shallow water waves admit peaked solitary waves. This work further confirms this view-point. This is mainly because the fully nonlinear water wave equations might admit peaked solitary waves, as pointed out by Liao [15]. Therefore, mathematically speaking, peaked solitary waves widely exist and might be a common property of models for water waves, although it should be carefully investigated whether or not such kind of peaked solitary waves indeed could be regarded as weak solutions and besides have physical meanings.



Figure 2: The peaked solitary waves u(x,t) of Eq. (2) in the case of $\omega = 1/4$, c = 1 and $\beta = 5/2$, corresponding to $\alpha = 7/2$, $\beta = 5/2$ of Eq. (4). Solid line: $A = \pm 3/10$; Dashed line: $A = \pm 1/5$; Dash-dotted line: $A = \pm 1/10$.

Acknowledgement

This work is partly supported by the State Key Lab of Ocean Engineering (Approval No. GKZD010056-6) and the National Natural Science Foundation of China.



Figure 3: The peaked solitary waves u(x,t) of (4) in the case of $\omega = 1/4, c = 1$, $\alpha = 11/3$ and $\beta = 5/2$. Solid line: $A = \pm 3/10$; Dashed line: $A = \pm 1/5$; Dash-dotted line: $A = \pm 1/10$.

References

- Benjamin, B., Bona, J.L. and Mahony, J.J. Model equations for long waves in nonlinear dispersive systems. *Philos. Trans. Roy. Soc. London*, 272: 47-78 (1972)
- [2] Bernetti, R., Titarev, V.A. and Toro, E.F. Exact solution of the Riemann problem for the shallow water equations with discontinuous bottom geometry. J. Computational Phys. 227: 3212 – 3243 (2008)
- [3] Boussinesq, J. Théorie des ondes et des remous qui se propagent le long d'un canal rectangulaire horizontal, en communiquant au liquide contenu dans ce canal des vitesses sensiblement pareilles de la surface au fond. Journal de Math'ematiques Pures et Appliquées. Deuxiéme Série 17: 55 - 108 (1872).
- [4] Camassa, R. and Holm, D. D. An integrable shallow water equation with peaked solitons. *Phys. Rev. Lett.* **71**, 1661 - 1664 (1993).
- [5] Constantin, A. Existence of permanent and breaking waves for a shallow water equation: a geometric approach. Ann. Inst. Fourier, Grenoble. 50, No.2, 321 -362 (2000)
- [6] Constantin, A. and Molinet, L. Global weak solutions for a shallow water equation. Comm. Math. Phys. 211, 45 - 61 (2000)
- [7] Degasperis, A. and Procesi, M. Asymptotic integrability. Symmetry and Perturbation Theory, Edited by Degasperis, A. and Gaeta, G., World Scientific, New Jersey, 22 – 37 (1999).

- [8] Korteweg, D. J. and de Vries, G. On the Change of Form of Long Waves Advancing in a Rectangular Canal, and on a New Type of Long Stationary Waves'. *Phil. Mag.* 39, 422 - 443 (1895)
- [9] Liao, S.J. An explicit, totally analytic approximation of Blasius viscous flow problems. Int. J. Non-Linear Mechanics, 34 (4): 759 - 778 (1999).
- [10] Liao, S.J. A uniformly valid analytic solution of 2D viscous flow past a semiinfinite flat plate. J. Fluid Mechanics, 385: 101-128 (1999).
- [11] Liao, S.J. Beyond Perturbation Introduction to the Homotopy Analysis Method, Chapman Hall/CRC Press, Boca Raton, 2003.
- [12] Liao, S.J. Homotopy Analysis Method in Nonlinear Differential Equations, Springer & Higher Education Press, Heidelberg & Beijing, 2012.
- [13] Liao, S.J. Two kinds of peaked solitary waves of the KdV, BBM and Boussinesq equations, arXiv: 1203.3917, 2012.
- [14] Liao, S.J. On peaked solitary waves of Camassa-Holm equation, arXiv: 1204.4517, 2012.
- [15] Liao, S.J., On a new type of solitary surface waves in finite water depth, arXiv: 1204.3354, 2012.
- [16] Rosatti, R. and Begnudelli, L. The Riemann Problem for the one-dimensional, free-surface Shallow Water Equations with a bed step: Theoretical analysis and numerical simulations. J. Computational Phys. 229: 760 – 787 (2010)
- [17] Russell, J.S. Report on waves. Fourteenth Meeting of the British Association for the Advancement of Science (1844).
- [18] Wu, Y.Y. and Cheung, K.F. Explicit solution to the exact Riemann problem and application in nonlinear shallow-water equations. Int. J. Numer. Meth. Fluids, 57: 1649 – 1668 (2008)
- [19] Zoppou, C. and Roberts, S. Numerical solution of the two-dimensional unsteady dam break. Applied Mathematical Modelling, 24: 457 – 475 (2000)

Appendix

The use of HAM-based Mathematica package BVPh 1.0

The Mathematica package BVPh 1.0 for nonlinear boundary-value/eigenvalue problems is developed and issued by Liao [12] (Part II), which is based on the HAM and is free available online. Using the BVPh 1.0, it is straightforward to gain the analytic approximations of the peaked solitary waves of Eqs. (5) and (6) for given w(0) = A. Here, we briefly describe how to do it in case of $c = 1, \omega = 1/4, \alpha = 4, \beta = 3$ and A = 1/10, corresponding to the DP equation (3).

- 1. First, download the BVPh 1.0 (the code file is named by BVPh_1.0.txt) online (http://numericaltank.sjtu.edu.cn/BVPh.htm) and save it in a directory such as C:/math/DP as an example.
- 2. Then, run the computer algebra system Mathematica, and type the following command one by one:

```
SetDirectory["C:\math\DP"];
```

<<InputDP.txt

The file named InputDP.txt contains the following Mathematica commands and necessary definitions for BVPh 1.0:

```
(* Install the BVPh 1.0 *)
<<BVPh1_0.txt;
(* Define the physical and control parameters *)
            = 1;
TypeEQ
ApproxQ
            =
               0;
ErrReq
            = 10^{(-30)};
zRintegral = 10;
(* Define the governing equation *)
ALPHA = 4;
BETA = 3;
mu2 = 1-2*omega/c;
f[z_,u_,lambda_] := D[u,{z,3}]-mu2*D[u,z] \setminus
    + ALPHA*u*D[u,z] - BETA*D[u,z]*D[u,{z,2}] - u*D[u,{z,3}] ;
If[BETA == 2 && ALPHA == 3, Print["CH equation"]];
If[BETA == 3 && ALPHA == 4, Print["DP equation"]];
```

```
(* Define Boundary conditions *)
zR = Infinity;
OrderEQ = 3;
BC[1,z_,u_,lambda_] := Limit[u-A, z -> 0 ];
BC[2,z_u_lambda_] := Limit[u, z \rightarrow zR];
BC[3,z_,u_,lambda_] := Limit[D[u,z], z -> zR ];
(* Define initial guess *)
mu = Sqrt[mu2];
u[0] = A*Exp[-mu*z];
(* Define output term *)
output[z_,u_,k_]:= Print["output = ",D[u[k],z] /. z->0//N];
(* Defines the auxiliary linear operator *)
L[u_] := D[u, \{z, 3\}] - mu2 * D[u, z];
(* Print input and control parameters *)
PrintInput[u[z]];
(* Set convergence-control parameter c0 and physical parameters *)
c0 = -1;
  = 1/10;
Α
omega = 1/4;
    = 1;
С
Print[" c0 = ",c0, " omega = ",omega, " c = ", c, " A = ",A];
(* Gain up to 10th-order HAM approxiamtion *)
BVPh[1,10];
(* Get results in the whole domain *)
For [k=0, k \le 10, k++, W[k] = U[k] /. z > Abs[x]];
(* Show the 5th and 10th-order approximation *)
Plot[{W[5],W[10]},{x,-10,10},PlotRange->{Min[A,0],Max[A,0]}]
```