

ON REMOVABILITY PROPERTIES OF ψ -UNIFORM DOMAINS IN BANACH SPACES

M. HUANG, M. VUORINEN, AND X. WANG *

ABSTRACT. Suppose that E denotes a real Banach space with the dimension at least 2. The main aim of this paper is to show that a domain D in E is a ψ -uniform domain if and only if $D \setminus P$ is a ψ_1 -uniform domain, and D is a uniform domain if and only if $D \setminus P$ also is a uniform domain, whenever P is a closed countable subset of D satisfying a quasihyperbolic separation condition. This condition requires that the quasihyperbolic distance (w.r.t. D) between each pair of distinct points in P has a lower bound greater than or equal to $\frac{1}{2}$.

1. INTRODUCTION AND MAIN RESULTS

The quasihyperbolic metric of a domain in a metric space was introduced by F. W. Gehring and his students B. Palka and B. Osgood in the 1970's [2, 3] in the setup of the Euclidean space \mathbb{R}^n , $n \geq 2$. Since its first appearance, the quasihyperbolic metric has become an important tool in geometric function theory and in its generalizations to metric spaces and to Banach spaces [20]. For instance, Väisälä's theory of quasiconformal maps in Banach spaces relies on the quasihyperbolic metric. Yet, some basic questions of the quasihyperbolic geometry in Banach spaces and even in Euclidean spaces are open. Such a basic question as the convexity of quasihyperbolic balls has been studied in [7, 9, 13, 22] only recently.

In this paper, we study the classes of uniform domains [12] and the wider class of ψ -uniform domains [24] in Banach spaces and the stability of these classes of domains under the removal of a countable set of points. The motivation for this study stems from the discussions in [6, 18]. In these two papers, the removability questions were studied for the classes of uniform domains and John domains. For latest results in the case of ψ -uniform domains in \mathbb{R}^n see [8]. Our main result extends these results to Banach spaces. We begin with some basic definitions and the statements of our results. The proofs and necessary supplementary notation and terminology will be given thereafter.

Throughout the paper, we always assume that E denotes a real Banach space with the dimension at least 2. The norm of a vector z in E is written as $|z|$, and for each pair of points z_1, z_2 in E , the distance between them is denoted by $|z_1 - z_2|$ and

File: rem20130620.tex, printed: 2018-11-12, 17.22

2000 *Mathematics Subject Classification*. Primary: 30C65, 30F45; Secondary: 30C20.

Key words and phrases. Uniform domain, ψ -uniform domain, removability property, quasihyperbolic metric.

* Corresponding author.

The research was partly supported by NSF's of China (No. 11071063 and No. 11101138) and by the Academy of Finland, Project 2600066611.

the closed line segment with endpoints z_1 and z_2 by $[z_1, z_2]$. We always use $\mathbb{B}(x_0, r)$ to denote the open ball $\{x \in E : |x - x_0| < r\}$ centered at x_0 with radius $r > 0$. Similarly, for the closed balls and spheres, we employ the usual notations $\overline{\mathbb{B}}(x_0, r)$ and $\partial\mathbb{B}(x_0, r)$, respectively. When $x_0 = 0$, briefly, we denote $\mathbb{B}(x_0, r) = \mathbb{B}(r)$, in particular, $\mathbb{B} = \mathbb{B}(1)$. In \mathbb{R}^n , we specially denote $\mathbb{B}^n(x_0, r)$ the ball centered at x_0 with radius r , $\mathbb{B}^n(r)$ the ball centered at 0 with radius r , and \mathbb{B}^n the ball centered at 0 with radius 1. We adopt some basic terminology following closely [11, 16, 17, 18].

Definition 1. A domain D in E is said to be *c-uniform* if there exists a constant c with the property that each pair of points z_1, z_2 in D can be joined by a rectifiable arc γ in D satisfying

- (1) $\min_{j=1,2} \ell(\gamma[z_j, z]) \leq c d_D(z)$ for all $z \in \gamma$, and
- (2) $\ell(\gamma) \leq c |z_1 - z_2|$,

where $\ell(\gamma)$ denotes the arc length of γ , $\gamma[z_j, z]$ the part of γ between z_j and z , and $d_D(z)$ the distance from z to the boundary ∂D of D [12]. Also we say that γ is a *double c-cone arc*.

Definition 2. Let $\psi : [0, \infty) \rightarrow [0, \infty)$ be a homeomorphism. A domain D in E is called *ψ -uniform* if

$$k_D(z_1, z_2) \leq \psi\left(\frac{|z_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}}\right)$$

for all $z_1, z_2 \in D$ [24].

It is a basic fact that uniform domains are special cases of ψ -uniform domains. See Section 2.

Simple examples show, see Example 2.5, that removing a countable closed set E from a uniform domain D may yield a domain $D \setminus E$ which is not ψ -uniform for any ψ . This motivates us to study countable subsets of a domain satisfying the following quasihyperbolic separation condition.

We say that a countable closed set P in a domain $D \subset E$ satisfies a quasihyperbolic separation condition if

$$(1.1) \quad P = \{x_j \in D : j = 1, 2, \dots\} \text{ and } k_D(x_i, x_j) \geq \frac{1}{2}, \quad \forall i \neq j.$$

In what follows, for a given domain D , the symbol P stands for a fixed countable subset of D with this property.

The purpose of this paper is to study the following problem.

Problem 1. Suppose that D is a ψ -uniform (resp. c -uniform) domain in E and P is as in (1.1). Is it true that $G = D \setminus P$ is a ψ_1 -uniform (resp. c_1 -uniform) domain, where ψ_1 (resp. c_1) depends only on ψ (resp. c)?

We are now in a position to formulate our results.

Theorem 1. *A domain D in E is a ψ -uniform domain if and only if $G = D \setminus P$ is a ψ_1 -uniform domain, where ψ and ψ_1 are $(2^{12}, 1, 3, 2^7)$ -equivalent.*

Here for two homeomorphisms ψ and $\psi_1 : [0, \infty) \rightarrow [0, \infty)$, we say that they are (a_1, a_2, a_3, a_4) -equivalent if there are positive constants a_1, a_2, a_3 and a_4 such that $\psi_1(t) = a_1\psi(a_2t)$ and $\psi(t) = a_3\psi_1(a_4t)$ for $t \geq 0$. As a corollary of Theorem 1, we have

Corollary 1. *A domain D in E is a c -uniform domain if and only if $G = D \setminus P$ is a c_1 -uniform domain, where the constants c and c_1 depend only on each other.*

Remark 1. *It easily follows from the discussions in Section 4 that Theorem 1 and Corollary 1 still hold when we replace the constant $\frac{1}{2}$ in P in (1.1) by any positive constant κ .*

The proofs of Theorem 1 and Corollary 1 will be given in Section 4. We will prove several lemmas in Section 3, which will be used later on, and in Section 2, some preliminaries will be introduced.

2. PRELIMINARY RESULTS

2.1. Quasihyperbolic distance and neargeodesics.

The *quasihyperbolic length* of a rectifiable arc or a path α in the norm metric in D is the number:

$$\ell_{k_D}(\alpha) = \int_{\alpha} \frac{|dz|}{d_D(z)}.$$

Gehring and Palka [3] introduced the quasihyperbolic metric of a domain in \mathbb{R}^n and it has been recently used by many authors in the study of quasiconformal mappings [1, 5, 22, 24, 25] and related questions [4]. Many of the basic properties of this metric may be found in [2, 17, 18, 21, 25].

For each pair of points z_1, z_2 in D , the *distance ratio metric* $j_D(z_1, z_2)$ between z_1 and z_2 is defined by

$$j_D(z_1, z_2) = \log \left(1 + \frac{|z_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}} \right).$$

The *quasihyperbolic distance* $k_D(z_1, z_2)$ between z_1 and z_2 is defined in the usual way:

$$k_D(z_1, z_2) = \inf\{\ell_{k_D}(\alpha)\},$$

where the infimum is taken over all rectifiable arcs α joining z_1 to z_2 in D . For all z_1, z_2 in D , we have [21]

$$(2.1) \quad k_D(z_1, z_2) \geq \inf_{\alpha} \left\{ \log \left(1 + \frac{\ell(\alpha)}{\min\{d_D(z_1), d_D(z_2)\}} \right) \right\} \geq \left| \log \frac{d_D(z_2)}{d_D(z_1)} \right|,$$

where the infimum is taken over all rectifiable curves α in D connecting z_1 and z_2 . Since $\ell(\alpha) \geq |z_1 - z_2|$ in (2.1), for all z_1, z_2 in D , we have

$$(2.2) \quad k_D(z_1, z_2) \geq j_D(z_1, z_2).$$

Next, if $|z_1 - z_2| \leq d_D(z_1)$, then we have [17], [23, Lemma 2.11]

$$(2.3) \quad k_D(z_1, z_2) \leq \log \left(1 + \frac{|z_1 - z_2|}{d_D(z_1) - |z_1 - z_2|} \right) \leq \frac{|z_1 - z_2|}{d_D(z_1) - |z_1 - z_2|},$$

where the last inequality follows from the following elementary inequality

$$(2.4) \quad \frac{r}{1 - r/2} \leq \log \frac{1}{1 - r} \leq \frac{r}{1 - r} \quad \text{for } 0 \leq r < 1.$$

The following observation easily follows from (2.2) and Definition 2.

Proposition 1. *If D is ψ -uniform, then the homeomorphism ψ satisfies*

$$\psi(t) \geq \log(1 + t)$$

for $t > 0$.

In [18], Väisälä characterized uniform domains by the quasihyperbolic metric.

Theorem A. ([2, 18]) *For a domain D , the following are quantitatively equivalent:*

- (1) D is a c -uniform domain;
- (2) $k_D(z_1, z_2) \leq c' j_D(z_1, z_2)$ for all $z_1, z_2 \in D$;
- (3) $k_D(z_1, z_2) \leq c'_1 j_D(z_1, z_2) + d$ for all $z_1, z_2 \in D$.

In the case of domains in \mathbb{R}^n , the equivalence of items (1) and (3) in Theorem A is due to Gehring and Osgood [2] and the equivalence of items (2) and (3) due to Vuorinen [24]. By Theorem A, we see that uniformity implies ψ -uniformity.

Example 2.5. For $r > 0$ and $j = 10, 11, \dots$ let $E(r, j)$ be a finite set of points in $A = \mathbb{B}^n(\frac{21r}{20}) \setminus \mathbb{B}^n(r)$ such that for every point $z \in A$ we have $d(z, A) < r/(20j)$. Let $G(r, j) = \mathbb{R}^n \setminus E(r, j)$ and $a_j = \frac{11r}{10}e_1$, $b_j = \frac{9r}{10}e_1$. It is readily seen that

$$k_{G(r,j)}(a_j, b_j) \geq \frac{r/20}{r/(20j)} = j.$$

Let $E = \cup_{j=10}^{\infty} E(2^{-j}, j) \cup \{0\}$. Clearly, $\mathbb{B}^n \setminus E$ is not ψ -uniform for any ψ although \mathbb{B}^n is uniform.

Recall that an arc α from z_1 to z_2 is a *quasihyperbolic geodesic* if $\ell_{k_D}(\alpha) = k_D(z_1, z_2)$. Each subarc of a quasihyperbolic geodesic is obviously a quasihyperbolic geodesic. It is known that a quasihyperbolic geodesic between every pair of points in E exists if the dimension of E is finite, see [2, Lemma 1]. This is not true in Banach spaces [19, Example 2.9]. In order to remedy this shortage, Väisälä introduced the following concepts [18].

Definition 3. Let D be a domain in E and $\nu > 1$. An arc $\alpha \subset D$ is a ν -neargeodesic if $\ell_{k_D}(\alpha[x, y]) \leq \nu k_D(x, y)$ for all $x, y \in \alpha$.

Obviously, a ν -neargeodesic is a quasihyperbolic geodesic if and only if $\nu = 1$. The smoothness of geodesics has been studied recently in [14].

In [19], Väisälä proved the following property concerning the existence of neargeodesics in Banach spaces.

Theorem B. ([19, Theorem 3.3]) *Fix $\nu > 1$. Then for $\{z_1, z_2\} \subset D \subset E$ there is a ν -neargeodesic in D joining z_1 and z_2 .*

2.2. Quasiconvexity.

Definition 4. We say that an arc γ in $D \subset E$ is c -*quasiconvex* in the norm metric if it satisfies the condition

$$\ell(\gamma[z_1, z_2]) \leq c|z_1 - z_2|$$

for every z_1, z_2 in γ .

The following result is due to Schäffer [15].

Theorem C. ([15, 4.4]) *Suppose that S in E is a sphere, that T is a 2-dimensional linear subspace in E and that the intersection $S \cap T$ contains at least two points. For every pair $\{z_1, z_2\} \subset T \cap S$, if $\gamma \subset T \cap S$ is the minor arc or a half circle with the endpoints z_1 and z_2 , then γ is 2-quasiconvex.*

3. SEVERAL LEMMAS

We recall that D denotes a domain in E and $G = D \setminus P$, where $P \subset D$ is a countable set satisfying the quasihyperbolic separation condition (1.1). The aim of this section is to prove several lemmas on which the proofs of Theorem 1 and Corollary 1 will be based. Our first lemma concerns the number of the points of P , which are contained in the balls $\mathbb{B}(x, \lambda d_D(x))$ in D for varying values of λ .

Lemma 1. (1) *Let $\lambda_0 \equiv 1 - \exp(-1/4) = 0.22\dots$. If $\lambda \in (0, \lambda_0)$ and $x \in D$, then $\mathbb{B}(x, \lambda d_D(x))$ contains at most one point of P .*

(2) *If $\lambda \in (0, \lambda_0)$, $x \in G$, and $\frac{d_G(x)}{d_D(x)} < \lambda$, then $\mathbb{B}(x, \lambda d_D(x))$ contains exactly one point of P .*

(3) *If $\lambda \leq \frac{1}{16}$, $x \in G$, and $\frac{d_G(x)}{d_D(x)} < \lambda$, then $\mathbb{B}(x, \lambda d_D(x))$ contains exactly one point x_i in P which satisfies*

$$d_G(z) = |x_i - z|$$

for all $z \in \overline{\mathbb{B}}(x, \frac{1}{16}d_D(x))$.

Proof. (1) Let $x \in D$. It follows from (2.3) that

$$k_D(\mathbb{B}(x, \lambda d_D(x))) = \sup\{k_D(u, v) : u, v \in \mathbb{B}(x, \lambda d_D(x))\} \leq 2 \log \frac{1}{1 - \lambda} < \frac{1}{2}.$$

Then (1.1) implies that (1) holds.

(2) Let $x \in G$ with $\frac{d_G(x)}{d_D(x)} < \lambda$. Then there exists some i such that $|x - x_i| < \lambda d_D(x)$. Hence $x_i \in \mathbb{B}(x, \lambda d_D(x))$ and by (1) $\mathbb{B}(x, \lambda d_D(x))$ cannot contain points in $P \setminus \{x_i\}$.

We now prove (3). Because $\frac{1}{5} \in (0, \lambda_0)$, we see by (1) that $\mathbb{B}(x, \frac{1}{5}d_D(x))$ contains at most one point of P . By (2) there exists a unique point $x_i \in \mathbb{B}(x, \lambda d_D(x)) \cap P$. Then

$$d_G(x) = |x - x_i| < \lambda d_D(x),$$

because for $j \neq i$ we have

$$|x - x_j| \geq \frac{1}{5}d_D(x) - \lambda d_D(x) > \lambda d_D(x)$$

by the choice of λ . In the same way we see that for all $z \in \mathbb{B}(x, \frac{1}{15}d_D(x))$ we have for all $j \neq i$

$$|z - x_j| \geq |x - x_j| - |x - z| > \frac{1}{5}d_D(x) - \lambda d_D(x) = (\frac{1}{5} - \lambda)d_D(x)$$

whereas

$$|z - x_i| \leq |z - x| + |x - x_i| < \frac{1}{15}d_D(x) + \lambda d_D(x) = (\frac{1}{15} + \lambda)d_D(x).$$

Because $\frac{1}{5} - \lambda > \frac{1}{15} + \lambda$, we see that for all $z \in \overline{\mathbb{B}}(x, \frac{1}{16}d_D(x))$,

$$d_G(z) = |x_i - z|.$$

The proof is complete. \square

The constant λ_0 in Lemma 1 obviously depends on the constant $\frac{1}{2}$ in the quasihyperbolic separation condition (1.1). At the same time it is easy to see that a result similar to Lemma 1 also holds if the constant $\frac{1}{2}$ in the definition of the set P (1.1) is replaced by $\sigma \in (0, \frac{1}{2})$. Only the constants would change and e.g. λ_0 would be replaced by λ_σ , with

$$2 \log \frac{1}{1 - \lambda_\sigma} = \sigma.$$

Given a point w in a domain $U \subset \mathbb{R}^n$ it is clear that

$$k_U(x, y) \leq k_{U \setminus \{w\}}(x, y)$$

for all $x, y \in U \setminus \{w\}$. For points $x, y \in U \setminus \mathbb{B}(w, \theta d_U(w))$, $0 < \theta < 1$, we also have an opposite inequality as shown in [24, Lemma 2.53]. The next few lemmas deal with this situation for Banach spaces. We start by a comparison theorem for the metrics k_G and j_G .

Lemma 2. *For $0 < \mu \leq \frac{1}{32}$ and $w_1, w_2 \in G$, if $w_2 \in \overline{\mathbb{B}}(w_1, \mu d_D(w_1))$ and $\min\{d_G(w_1), d_G(w_2)\} \leq \frac{\mu}{2}d_D(w_1)$, then*

$$k_G(w_1, w_2) \leq \frac{13}{2}j_G(w_1, w_2).$$

Proof. Clearly, we have

$$(3.1) \quad \begin{aligned} \max\{d_G(w_1), d_G(w_2)\} &\leq \min\{d_G(w_1), d_G(w_2)\} + |w_1 - w_2| \\ &\leq \frac{3\mu}{2}d_D(w_1). \end{aligned}$$

By Lemma 1 (3), we see that there exists some point $x_i \in \mathbb{B}(w_1, \frac{3\mu}{2}d_D(w_1)) \cap P$ satisfying

$$|w_2 - x_i| = d_G(w_2) \quad \text{and} \quad |w_1 - x_i| = d_G(w_2).$$

Without loss of generality, we may assume

$$(3.2) \quad \min\{d_G(w_1), d_G(w_2)\} = d_G(w_2).$$

We use $w_{1,1}$ to denote the intersection point of the closed segment $[w_1, x_i]$ with the sphere $\mathbb{S}(x_i, \min\{d_G(w_1), d_G(w_2)\})$. It is possible that $w_{1,1} = w_1$. Let T denote a 2-dimensional linear subspace of E passing thorough the points w_1, w_2 and x_i , and ω_0 the circle $T \cap \mathbb{S}(x_i, \min\{d_G(w_1), d_G(w_2)\})$. Then $w_{1,1}$ and w_2 divide the circle ω_0 into two parts β_i and $\beta_{1,i}$. Without loss of generality, we may assume that $\ell(\beta_i) \leq \ell(\beta_{1,i})$. Then it follows from (3.2) that for each $z \in \beta_i$,

$$|z - w_1| \leq |w_1 - x_i| + |x_i - z| \leq 2\mu d_D(w_1),$$

whence

$$\beta_i \subset \overline{\mathbb{B}}(w_1, 2\mu d_D(w_1)),$$

and so Lemma 1 yields that for each $z \in \beta_i$,

$$d_G(z) = d_G(w_2),$$

which, together with (2.3) and Theorem C, shows that

$$\begin{aligned} k_G(w_1, w_2) &\leq k_G(w_1, w_{1,1}) + \ell_{k_G}(\beta_i) \\ &\leq \log\left(1 + \frac{|w_1 - w_{1,1}|}{d_G(w_2)}\right) + \frac{2|w_2 - w_{1,1}|}{d_G(w_2)} \\ &\leq \log\left(1 + \frac{|w_1 - w_{1,1}|}{d_G(w_2)}\right) + \frac{4}{\log 3} \log\left(1 + \frac{|w_2 - w_{1,1}|}{d_G(w_2)}\right) \\ &< \left(1 + \frac{6}{\log 3}\right) \log\left(1 + \frac{|w_1 - w_2|}{d_G(w_2)}\right) \\ &\leq \frac{13}{2} j_G(w_1, w_2), \end{aligned}$$

since $\frac{|w_2 - w_{1,1}|}{d_G(w_2)} \leq 4$ and $|w_2 - w_{1,1}| \leq 2|w_1 - w_2|$. Hence the proof follows. □

The next two results are related to k_G and k_D .

Lemma 3. *For $w_1 \in G$, suppose $d_G(w_1) = \frac{1}{128}d_D(w_1)$. If $w_2 \in \mathbb{S}(w_1, \frac{1}{32}d_D(w_1))$, then*

$$k_G(w_1, w_2) \leq 2^9 k_D(w_1, w_2).$$

Proof. It follows from Lemma 1 that there is a unique element x_i in the intersection $P \cap \overline{\mathbb{B}}(w_1, \frac{1}{64}d_D(w_1))$ such that

$$d_G(w_2) = |w_2 - x_i| \geq \frac{3}{128}d_D(w_1) = 3d_G(w_1).$$

Then Lemma 2, (2.1) and the Bernoulli inequality [25, (3.6)] imply

$$\begin{aligned}
k_G(w_1, w_2) &\leq \frac{13}{2} \log \left(1 + \frac{|w_1 - w_2|}{d_G(w_1)} \right) \\
&= \frac{13}{2} \log \left(1 + \frac{128|w_1 - w_2|}{d_D(w_1)} \right) \\
&< 2^9 \log \left(1 + \frac{|w_1 - w_2|}{d_D(w_1)} \right) \\
&\leq 2^9 k_D(w_1, w_2),
\end{aligned}$$

which shows that the lemma is true. \square

Lemma 4. *Let $w_1, w_2 \in G$ and let γ denote a 2-neargeodesic joining w_1 and w_2 in D . If $d_G(z) \geq \frac{1}{128}d_D(z)$ for each $z \in \gamma$, then*

$$k_G(w_1, w_2) \leq 2^8 k_D(w_1, w_2).$$

Proof. Obviously, we get

$$\begin{aligned}
k_G(w_1, w_2) &\leq \ell_{k_G}(\gamma[w_1, w_2]) = \int_{\gamma[w_1, w_2]} \frac{|dx|}{d_G(x)} \leq 128 \int_{\gamma[w_1, w_2]} \frac{|dx|}{d_D(x)} \\
&= 128 \ell_{k_D}(\gamma[w_1, w_2]) \leq 2^8 k_D(w_1, w_2).
\end{aligned}$$

Hence the proof is complete. \square

Our last lemma in this section is as follows.

Lemma 5. *Let $w_1, w_2 \in D$. If $|w_1 - w_2| \geq \frac{1}{c}d_D(w_1)$ ($c \geq 2$), then*

$$|w_1 - w_2| \geq \frac{1}{c+1}d_D(w_2).$$

Proof. Suppose on the contrary that

$$|w_1 - w_2| < \frac{1}{c+1}d_D(w_2).$$

Then

$$d_D(w_1) \geq d_D(w_2) - |w_1 - w_2| > \frac{c}{c+1}d_D(w_2),$$

which shows that

$$|w_1 - w_2| \geq \frac{1}{c}d_D(w_1) > \frac{1}{c+1}d_D(w_2).$$

This is the desired contradiction. \square

4. THE PROOFS OF THEOREM 1 AND COROLLARY 1

4.1. **The proof of Theorem 1.** We first prove the sufficiency. Suppose that G is a ψ_1 -uniform domain. Then we shall prove that for $z_1, z_2 \in D$,

$$(4.1) \quad k_D(z_1, z_2) \leq \psi\left(\frac{|z_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}}\right),$$

where $\psi(t) = 3\psi_1(2^7 t)$ for $t > 0$, which implies that D is a ψ -uniform domain. Without loss of generality, we assume that

$$\min\{d_G(z_1), d_G(z_2)\} = d_G(z_1).$$

We divide the proof into two cases.

4.1.1. *We first suppose that $|z_1 - z_2| \leq \frac{1}{2}d_D(z_1)$.*

Then it follows from (2.3) and Proposition 1 that

$$(4.2) \quad \begin{aligned} k_D(z_1, z_2) &\leq \log\left(1 + \frac{|z_1 - z_2|}{d_D(z_1) - |z_1 - z_2|}\right) \leq 2\log\left(1 + \frac{|z_1 - z_2|}{d_D(z_1)}\right) \\ &\leq 2\psi_1\left(\frac{|z_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}}\right). \end{aligned}$$

4.1.2. *We then suppose that $|z_1 - z_2| > \frac{1}{2}d_D(z_1)$.*

Hence by Lemma 5, we have

$$(4.3) \quad |z_1 - z_2| \geq \frac{1}{3}d_D(z_2).$$

In the following, we separate the rest discussions to two parts.

Case 1. $d_G(z_1) \geq \frac{1}{64}d_D(z_1)$.

Under this assumption, we have

$$(4.4) \quad \begin{aligned} k_D(z_1, z_2) &\leq k_G(z_1, z_2) \leq \psi_1\left(\frac{|z_1 - z_2|}{d_G(z_1)}\right) \leq \psi_1\left(\frac{64|z_1 - z_2|}{d_D(z_1)}\right) \\ &\leq \psi_1\left(\frac{64|z_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}}\right). \end{aligned}$$

Case 2. $d_G(z_1) < \frac{1}{64}d_D(z_1)$.

For a proof under this assumption, we let $u_1 \in \mathbb{S}(z_1, \frac{1}{32}d_D(z_1))$. Then by (2.3)

$$(4.5) \quad k_D(z_1, u_1) \leq \frac{32|z_1 - u_1|}{31d_D(z_1)} = \frac{1}{31} < \log\left(1 + \frac{|z_1 - z_2|}{d_D(z_1)}\right).$$

Moreover, we get

$$(4.6) \quad |u_1 - z_2| \leq |z_1 - z_2| + |z_1 - u_1| \leq \frac{17}{16}|z_1 - z_2|,$$

and it follows from Lemma 1 and the assumption “ $d_G(z_1) < \frac{1}{64}d_D(z_1)$ ” that there exists only one element x_i in $\overline{\mathbb{B}}(z_1, \frac{1}{64}d_D(z_1)) \cap P$ such that $d_G(u_1) = |u_1 - x_i|$ and so

$$(4.7) \quad d_D(u_1) > d_G(u_1) = |u_1 - x_i| \geq |u_1 - z_1| - |z_1 - x_i| \geq \frac{1}{64}d_D(z_1).$$

In this part, we again distinguish two possibilities.

Subcase 1. $d_G(z_2) \geq \frac{1}{120}d_D(z_2)$.

Then by (4.5), (4.6) and (4.7), we have

$$\begin{aligned}
(4.8) \quad k_D(z_1, z_2) &\leq k_D(z_1, u_1) + k_D(u_1, z_2) \\
&\leq \log \left(1 + \frac{|z_1 - z_2|}{d_D(z_1)} \right) + k_G(u_1, z_2) \\
&\leq j_D(z_1, z_2) + \psi_1 \left(\frac{|u_1 - z_2|}{\min\{d_G(u_1), d_G(z_2)\}} \right) \\
&\leq j_D(z_1, z_2) + \psi_1 \left(\frac{120|u_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}} \right) \\
&\leq j_D(z_1, z_2) + \psi_1 \left(\frac{2^7|z_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}} \right) \\
&< 2\psi_1 \left(\frac{2^7|z_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}} \right).
\end{aligned}$$

Subcase 2. $d_G(z_2) < \frac{1}{120}d_D(z_2)$.

Under this hypothesis, to get a homeomorphism ψ from ψ_1 , we take $u_2 \in \mathbb{S}(z_2, \frac{1}{32}d_D(z_2))$. It follows from Lemma 1 and (4.3) that there is only one element x_i in $\overline{\mathbb{B}}(z_2, \frac{1}{32}d_D(z_2)) \cap P$ such that $d_G(u_2) = |u_2 - x_i|$ and so

$$(4.9) \quad d_D(u_2) \geq d_G(u_2) = |u_2 - x_i| \geq |u_2 - z_2| - |z_2 - x_i| \geq \frac{11}{480}d_D(z_2)$$

and by (4.3), we have

$$(4.10) \quad |u_1 - u_2| \leq |z_1 - z_2| + |z_1 - u_1| + |u_2 - z_2| \leq \frac{37}{32}|z_1 - z_2|.$$

It follows from (4.3), (2.3) and (4.5) that

$$(4.11) \quad k_D(z_2, u_2) \leq \frac{32|z_2 - u_2|}{31d_D(z_2)} < \log \left(1 + \frac{|z_1 - z_2|}{d_D(z_2)} \right).$$

Then we infer from (4.5), (4.7), (4.9), (4.10), (4.11) and Proposition 1 that

$$\begin{aligned}
(4.12) \quad k_D(z_1, z_2) &\leq k_D(z_1, u_1) + k_G(u_1, u_2) + k_D(u_2, z_2) \\
&\leq 2j_D(z_1, z_2) + k_G(u_1, u_2) \\
&\leq 2j_D(z_1, z_2) + \psi_1 \left(\frac{|u_1 - u_2|}{\min\{d_G(u_1), d_G(u_2)\}} \right) \\
&\leq 2j_D(z_1, z_2) + \psi_1 \left(64 \frac{|u_1 - u_2|}{\min\{d_D(z_1), d_D(z_2)\}} \right) \\
&\leq 2j_D(z_1, z_2) + \psi_1 \left(74 \frac{|z_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}} \right) \\
&< 3\psi_1 \left(74 \frac{|z_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}} \right).
\end{aligned}$$

Since ψ_1 is increasing, by taking $\psi(t) = 3\psi_1(2^7 t)$ for $t \geq 0$, we easily see from the inequalities (4.2), (4.4), (4.8) and (4.12) that (4.1) holds.

Next we prove the necessity. Suppose that D is a ψ -uniform domain. Then we shall prove that for $z_1, z_2 \in G$,

$$(4.13) \quad k_G(z_1, z_2) \leq 2^{12} \psi \left(\frac{|z_1 - z_2|}{\min\{d_G(z_1), d_G(z_2)\}} \right),$$

which implies that G is a ψ_1 -uniform domain with $\psi_1 = 2^{12}\psi$.

Without loss of generality, we may assume that $\min\{d_G(z_1), d_G(z_2)\} = d_G(z_1)$. In the following, we consider the two cases where $d_G(z_1) \leq \frac{1}{64}d_D(z_1)$ and $d_G(z_1) > \frac{1}{64}d_D(z_1)$, respectively.

4.1.3. *We first suppose that $d_G(z_1) \leq \frac{1}{64}d_D(z_1)$.*

Let γ be a 2-neargeodesic joining z_1 and z_2 in D . The existence of γ follows from Theorem B. We separate the discussions in this case to two parts.

Case 3. $|z_1 - z_2| \leq \frac{1}{32}d_D(z_1)$.

Then by Lemma 2 and Proposition 1, we have

$$\textbf{Claim 1.} \quad k_G(z_1, z_2) \leq \frac{13}{2} \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right) \leq \frac{13}{2} \psi \left(\frac{|z_1 - z_2|}{d_G(z_1)} \right).$$

Case 4. $|z_1 - z_2| > \frac{1}{32}d_D(z_1)$.

Under this condition, Lemma 5 implies

$$(4.14) \quad |z_1 - z_2| \geq \frac{1}{33}d_D(z_2).$$

Obviously, there exists some point $v_1 \in \gamma \cap \mathbb{S}(z_1, \frac{1}{32}d_D(z_1))$ such that

$$\gamma[z_2, v_1] \subset D \setminus \mathbb{B}(z_1, \frac{1}{32}d_D(z_1)).$$

By Lemma 1, there exists some point $x_{i,1} \in \overline{\mathbb{B}}(z_1, \frac{1}{64}d_D(z_1)) \cap P$ such that

$$(4.15) \quad d_G(v_1) = |v_1 - x_{i,1}| \geq \frac{1}{64}d_D(z_1) \geq \frac{1}{66}d_D(v_1),$$

since $d_D(v_1) \leq d_D(z_1) + |z_1 - v_1| \leq \frac{33}{32}d_D(z_1)$.

It follows from Lemma 2 and (4.15) that

$$(4.16) \quad k_G(z_1, v_1) \leq \frac{13}{2} \log \left(1 + \frac{|z_1 - v_1|}{d_G(z_1)} \right) \leq \frac{13}{2} \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right).$$

In this part, we again distinguish two possibilities.

Subcase 3. $d_G(z) \geq \frac{1}{128}d_D(z)$ for each $z \in \gamma[v_1, z_2]$.

By Lemma 4, we know

$$\begin{aligned} k_G(v_1, z_2) &\leq 2^8 k_D(v_1, z_2) \leq 2^8 \ell_{k_D}(\gamma[v_1, z_2]) \leq 2^8 \ell_{k_D}(\gamma[z_1, z_2]) \\ &\leq 2^9 k_D(z_1, z_2) \leq 2^9 \psi \left(\frac{|z_1 - z_2|}{d_G(z_1)} \right), \end{aligned}$$

since D is ψ -uniform, and so the following inequality easily follows from (4.16).

Claim 2. $k_G(z_1, z_2) \leq k_G(z_1, v_1) + k_G(v_1, z_2) \leq 519\psi\left(\frac{|z_1 - z_2|}{d_G(z_1)}\right)$.

Subcase 4. There exists some point $z \in \gamma[v_1, z_2]$ such that $d_G(z) < \frac{1}{128}d_D(z)$.

Under this assumption, it follows from (4.15) that there exists some point y_1 which is the first point in γ along the direction from v_1 to z_2 such that

$$d_G(y_1) = \frac{1}{128}d_D(y_1).$$

Then Lemma 4 shows

$$(4.17) \quad \begin{aligned} k_G(v_1, y_1) &\leq 2^8 k_D(v_1, y_1) \leq 2^8 \ell_{k_D}(\gamma[z_1, z_2]) \\ &\leq 2^9 k_D(z_1, z_2) \leq 2^9 \psi\left(\frac{|z_1 - z_2|}{d_D(z_1)}\right). \end{aligned}$$

To get a homeomorphism ψ_1 from ψ in this possibility, we consider two cases.

Subsubcase 1. $|z_2 - y_1| \leq \frac{1}{32}d_D(y_1)$.

Then we see from Lemma 2 that

$$(4.18) \quad k_G(y_1, z_2) \leq \frac{13}{2} \log\left(1 + \frac{|y_1 - z_2|}{\min\{d_G(y_1), d_G(z_2)\}}\right).$$

Thus we have the following claim.

Claim 3. $k_G(z_1, z_2) \leq 564\psi\left(\frac{|z_1 - z_2|}{d_G(z_1)}\right)$.

We now prove this claim. Since

$$d_D(z_2) \geq d_D(y_1) - |z_2 - y_1| \geq \frac{31}{32}d_D(y_1) \geq 31|z_2 - y_1|,$$

we infer from (4.14) that

$$(4.19) \quad |z_1 - z_2| \geq \frac{31}{33}|y_1 - z_2|,$$

and by (4.18), we get

$$k_G(y_1, z_2) \leq \frac{13}{2} \log\left(1 + \frac{33|z_1 - z_2|}{31 \min\{d_G(y_1), d_G(z_2)\}}\right).$$

Let us leave the proof of Claim 3 for a moment and prove the following inequality

$$(4.20) \quad k_G(y_1, z_2) \leq 35 \log\left(1 + \frac{|z_1 - z_2|}{d_G(z_2)}\right).$$

To prove this estimate, obviously, we only need to consider the case $d_G(z_2) \geq d_G(y_1)$. Since

$$d_G(z_2) \leq d_G(y_1) + |z_2 - y_1| \quad \text{and} \quad |z_2 - y_1| \leq \frac{1}{32}d_D(y_1) = 4d_G(y_1),$$

we see from (4.18) and the Bernoulli inequality [25, (3.6)] that

$$k_G(y_1, z_2) \leq \frac{13}{2} \log\left(1 + \frac{|y_1 - z_2|}{d_G(y_1)}\right) \leq \frac{13}{2} \log\left(1 + \frac{5|y_1 - z_2|}{d_G(z_2)}\right) < 35 \log\left(1 + \frac{|z_1 - z_2|}{d_G(z_2)}\right).$$

Hence (4.20) is true.

Now, we come back to the proof of Claim 3. It follows from (4.16), (4.17) and (4.20) that

$$\begin{aligned}
 k_G(z_1, z_2) &\leq k_G(z_1, v_1) + k_G(v_1, y_1) + k_G(y_1, z_2) \\
 &\leq \frac{13}{2} \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right) + 2^9 \psi \left(\frac{|z_1 - z_2|}{d_G(z_1)} \right) + 35 \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_2)} \right) \\
 &\leq 42 \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right) + 2^9 \psi \left(\frac{|z_1 - z_2|}{d_G(z_1)} \right) \\
 &\leq 564 \psi \left(\frac{|z_1 - z_2|}{d_G(z_1)} \right),
 \end{aligned}$$

which shows that Claim 3 is true.

Subsubcase 2. $|z_2 - y_1| > \frac{1}{32}d_D(y_1)$.

Obviously, there exists some point $v_2 \in \gamma \cap \mathbb{S}(y_1, \frac{1}{32}d_D(y_1))$ such that

$$\gamma[z_2, v_2] \subset D \setminus \mathbb{B}(y_1, \frac{1}{32}d_D(y_1)).$$

By Lemma 1, we see that there exists some point $x_{i,2} \in P \cap \overline{\mathbb{B}}(y_1, \frac{1}{128}d_D(y_1))$ such that

$$\begin{aligned}
 (4.21) \quad d_G(v_2) &= |v_2 - x_{i,2}| \geq |v_2 - y_1| - |y_1 - x_{i,2}| \\
 &\geq \frac{3}{128}d_D(y_1) \geq \frac{1}{44}d_D(v_2),
 \end{aligned}$$

since $d_D(v_2) \leq d_D(y_1) + |y_1 - v_2| \leq \frac{33}{32}d_D(y_1)$.

Moreover, by Lemma 3, we have

$$(4.22) \quad k_G(y_1, v_2) \leq 2^9 k_D(y_1, v_2) \leq 2^{10} k_D(z_1, z_2).$$

If $d_G(z) \geq \frac{1}{128}d_D(z)$ for each $z \in \gamma[v_2, z_2]$, then Lemma 4 implies

$$(4.23) \quad k_G(v_2, z_2) \leq 2^8 k_D(v_2, z_2) \leq 2^9 k_D(z_1, z_2),$$

and thus we infer from (4.16), (4.17), (4.22), (4.23) that

$$k_G(z_1, z_2) \leq \frac{13}{2} \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right) + 3 \cdot 2^{10} k_D(z_1, z_2).$$

Hence we have the following estimate.

Claim 4. $k_G(z_1, z_2) \leq 2^{12} \psi \left(\frac{|z_1 - z_2|}{d_G(z_1)} \right)$.

For the remaining case, that is, there exists some point $z \in \gamma[v_2, z_2]$ such that $d_G(z) < \frac{1}{128}d_D(z)$, similar arguments as in Subcase 3 show that there exists some point $y_2 \in \gamma[v_2, z_2]$ satisfying

$$d_G(y_2) = \frac{1}{128}d_D(y_2)$$

and

$$(4.24) \quad k_G(v_2, y_2) \leq 2^8 k_D(v_2, y_2).$$

Now, if $|z_2 - y_2| \leq \frac{1}{32} d_D(y_2)$, then the similar reasoning as in the proof of (4.20) shows that

$$(4.25) \quad k_G(y_2, z_2) \leq 35 \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_2)} \right).$$

Then it follows from (4.16), (4.17), (4.22), (4.24) and (4.25) that

$$\begin{aligned} k_G(z_1, z_2) &\leq k_G(z_1, v_1) + k_G(v_1, y_1) + k_G(y_1, v_2) + k_G(v_2, y_2) + k_G(y_2, z_2) \\ &\leq 2^{10} k_D(z_1, z_2) + 42 \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right). \end{aligned}$$

Hence we reach the following estimate.

Claim 5. $k_G(z_1, z_2) \leq 2^{12} \psi \left(\frac{|z_1 - z_2|}{d_G(z_1)} \right)$.

We assume now that $|z_2 - y_2| > \frac{1}{32} d_D(y_2)$. Then there exists some point $v_3 \in \gamma[y_2, z_2] \cap \mathbb{S}(y_2, \frac{1}{32} d_D(y_2))$ such that

$$\gamma[z_2, v_3] \subset D \setminus \mathbb{B}(y_2, \frac{1}{32} d_D(y_2)),$$

and the similar reasoning as in the proof of (4.22) shows that

$$k_G(y_2, v_3) \leq 2^9 k_D(y_2, v_3) \leq 2^{10} k_D(z_1, z_2).$$

By repeating the procedure as above, we will reach a finite sequence of points in γ :

- (1) $\{z_1, v_1, y_1, v_2, y_2, \dots, v_t, z_2\}$ such that $d_G(z) \geq \frac{1}{128} d_D(z)$ for each $z \in \gamma[v_t, z_2]$;
or
- (2) $\{z_1, v_1, y_1, v_2, y_2, \dots, v_t, y_t, z_2\}$ such that $|z_2 - y_t| \leq \frac{1}{32} d_D(y_t)$.

It follows from (2.1) that

$$k_D(y_i, v_{i+1}) \geq \log \left(1 + \frac{|y_i - v_{i+1}|}{d_D(y_i)} \right) \geq \log \frac{33}{32}$$

for each $i \in \{1, \dots, t\}$, and we see that

$$t \leq \frac{k_D(z_1, z_2)}{\log \frac{33}{32}}.$$

For the former case, i.e., when the statement (1) as above holds, we have shown that

- (1) $k_G(z_1, v_1) \leq \frac{13}{2} \log \left(1 + \frac{|z_1 - v_1|}{d_G(z_1)} \right) \leq \frac{13}{2} \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right)$;
- (2) $k_G(v_i, y_i) \leq 2^8 k_D(v_i, y_i)$, where $i \in \{1, \dots, t-1\}$;
- (3) $k_G(y_i, v_{i+1}) \leq 2^9 k_D(y_i, v_{i+1})$, where $i \in \{1, \dots, t\}$; and
- (4) $k_G(v_t, z_2) \leq 2^8 k_D(v_t, z_2) \leq 2^9 k_D(z_1, z_2)$.

Hence we obtain

$$\begin{aligned}
 k_G(z_1, z_2) &\leq k_G(z_1, v_1) + \sum_{i=1}^{t-1} k_G(v_i, y_i) + \sum_{i=2}^t k_G(y_{i-1}, v_i) + k_G(v_t, z_2) \\
 &\leq \frac{13}{2} \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right) + 2^8 \sum_{i=1}^{t-1} k_D(v_i, y_i) + 2^9 \sum_{i=2}^t k_D(y_{i-1}, v_i) \\
 &\quad + 35 \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_2)} \right) \\
 &\leq \frac{13}{2} \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right) + 2^{10} k_D(z_1, z_2) + 35 \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_2)} \right),
 \end{aligned}$$

which shows

Claim 6. $k_G(z_1, z_2) \leq 2^{11} \psi \left(\frac{|z_1 - z_2|}{d_G(z_1)} \right)$.

For the latter case, i.e., the statement (2) as above holds, we also have shown that

- (1) $k_G(z_1, v_1) \leq \frac{13}{2} \log \left(1 + \frac{|z_1 - v_1|}{d_G(z_1)} \right) \leq \frac{13}{2} \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right)$;
- (2) $k_G(v_i, y_i) \leq 2^8 k_D(v_i, y_i)$, where $i \in \{1, \dots, t\}$;
- (3) $k_G(y_i, v_{i+1}) \leq 2^9 k_D(y_i, v_{i+1})$, where $i \in \{1, \dots, t-1\}$, and
- (4) $k_G(y_t, z_2) \leq 35 \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_2)} \right)$.

Hence we get

$$\begin{aligned}
 k_G(z_1, z_2) &\leq k_G(z_1, v_1) + \sum_{i=1}^t k_G(v_i, y_i) + \sum_{i=2}^t k_G(y_{i-1}, v_i) + k_G(y_t, z_2) \\
 &\leq \frac{13}{2} \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right) + 2^8 \sum_{i=1}^t k_D(v_i, y_i) + 2^9 \sum_{i=2}^t k_D(y_{i-1}, v_i) \\
 &\quad + 35 \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_2)} \right) \\
 &\leq \frac{13}{2} \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right) + 2^{10} k_D(z_1, z_2) + 35 \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_2)} \right),
 \end{aligned}$$

which implies

Claim 7. $k_G(z_1, z_2) \leq 2^{11} \psi \left(\frac{|z_1 - z_2|}{d_G(z_1)} \right)$.

By taking $\psi_1(t) = 2^{12} \psi(t)$ for $t \geq 0$, we see that in the case $d_G(z_1) \leq \frac{1}{64} d_D(z_1)$, (4.13) follows from Claims 1 ~ 7.

4.1.4. We then suppose that $d_G(z_1) > \frac{1}{64} d_D(z_1)$.

Also, we divide the discussions into two cases.

Case 5. $|z_1 - z_2| \leq \frac{1}{2} d_G(z_1)$.

By (2.3) we have

$$k_G(z_1, z_2) \leq \frac{2|z_1 - z_2|}{d_G(z_1)}.$$

Hence by Proposition 1, we have

Claim 8. $k_G(z_1, z_2) \leq 3 \log \left(1 + \frac{|z_1 - z_2|}{d_G(z_1)} \right) \leq 3\psi \left(\frac{|z_1 - z_2|}{d_G(z_1)} \right).$

Case 6. $|z_1 - z_2| > \frac{1}{2}d_G(z_1)$

Since $|z_1 - z_2| > \frac{1}{128}d_D(z_1)$, we know from Lemma 5 that

$$(4.26) \quad |z_1 - z_2| > \frac{1}{129}d_D(z_2).$$

Let β be a 2-neargeodesic joining z_1 and z_2 in D . We divide the rest discussions into two subcases.

Subcase 5. $d_G(z) \geq \frac{1}{64}d_D(z)$ for each $z \in \beta$.

In this case, the following inequality easily follows from Lemma 4.

Claim 9. $k_G(z_1, z_2) \leq 2^8 k_D(z_1, z_2) \leq 2^8 \psi \left(\frac{|z_1 - z_2|}{d_G(z_1)} \right).$

Subcase 6. There exists some point $z \in \beta$ such that $d_G(z) < \frac{1}{64}d_D(z)$.

Obviously, it follows from the assumption “ $d_G(z_1) > \frac{1}{64}d_D(z_1)$ ” that there exists point p_1 which is the first point in β along the direction from z_1 to z_2 such that

$$d_G(p_1) = \frac{1}{64}d_D(p_1).$$

Then Lemma 4 shows

$$(4.27) \quad k_G(z_1, p_1) \leq 2^8 k_D(z_1, p_1).$$

To get a homeomorphism ψ_1 from ψ , we consider the case where $|z_2 - p_1| \leq \frac{1}{32}d_D(p_1)$ and the case where $|z_2 - p_1| > \frac{1}{32}d_D(p_1)$, respectively.

Subsubcase 3. $|z_2 - p_1| \leq \frac{1}{32}d_D(p_1)$.

Under this condition, it follows from (4.26) that

$$(4.28) \quad |z_2 - p_1| \leq \frac{1}{31}d_D(z_2) \leq \frac{129}{31}|z_1 - z_2|$$

since $d_D(z_2) \geq d_D(p_1) - |z_2 - p_1| \geq \frac{31}{32}d_D(p_1)$.

By Lemma 1, we have

$$d_G(z_2) \leq \frac{1}{16}d_D(p_1) \leq 4d_G(p_1).$$

Then we know from Lemma 2 and (4.28) that

$$k_G(p_1, z_2) \leq \frac{13}{2} \log \left(1 + \frac{|z_2 - p_1|}{\min\{d_G(z_2), d_G(p_1)\}} \right) \leq 28 \log \left(1 + \frac{|z_2 - p_1|}{d_G(z_2)} \right),$$

which, together with (4.27), implies

Claim 10. $k_G(z_1, z_2) \leq 28\psi\left(\frac{|z_1 - z_2|}{d_G(z_1)}\right)$.

Subsubcase 4. $|z_2 - p_1| > \frac{1}{32}d_D(p_1)$.

Obviously, there exists some point $q_1 \in \beta[p_1, z_2]$ such that

$$\beta[q_1, z_2] \subset D \setminus \mathbb{B}(p_1, \frac{1}{32}d_D(p_1)).$$

By Lemma 1, we see that there exists some point $x_{i,3} \in P \cap \overline{\mathbb{B}}(p_1, \frac{1}{128}d_D(p_1))$ such that

$$(4.29) \quad \begin{aligned} d_G(q_1) &= |q_1 - x_{i,3}| \geq |q_1 - p_1| - |p_1 - x_{i,3}| \\ &\geq \frac{3}{128}d_D(p_1) \geq \frac{1}{44}d_D(q_1), \end{aligned}$$

since $d_D(q_1) \leq d_D(p_1) + |p_1 - q_1| \leq \frac{33}{32}d_D(p_1)$.

Then the similar reasoning as in Subsubcase 2 in Subsection 4.1.3 implies that we will get a finite sequence of points in β :

- (1) $\{z_1, p_1, q_1, \dots, p_s, z_2\}$ such that $d_G(z) \geq \frac{1}{128}d_D(z)$ for each $z \in \gamma[p_s, z_2]$; or
- (2) $\{z_1, p_1, q_1, \dots, p_s, q_s, z_2\}$ such that $|z_2 - q_s| \leq \frac{1}{32}d_D(q_s)$.

It follows from the similar arguments as in Claims 6 and 7 in Subsubsection 2 that

Claim 11. $k_G(z_1, z_2) \leq 2^{11}\psi\left(\frac{|z_1 - z_2|}{d_G(z_1)}\right)$.

Now we are in a position to conclude that the truth of (4.13) for the case $d_G(z_1) > \frac{1}{64}d_D(z_1)$ follows from Claims 8 ~ 11. Hence the proof of Theorem 1 is complete. \square

4.2. The proof of Corollary 1. First, we prove the sufficiency. Suppose $G = D \setminus P$ is a c_1 -uniform domain. Then Theorem A implies that there exists a constant c'_1 depending only on c_1 such that for all x and y in G ,

$$k_G(x, y) \leq c'_1 j_G(x, y).$$

By Theorem 1, we see that for all z_1 and z_2 in D ,

$$k_D(z_1, z_2) \leq 3c'_1 \log\left(1 + \frac{128|z_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}}\right) \leq 384c'_1 j_D(z_1, z_2),$$

which, together with Theorem A, shows that D is a c -uniform domain, where c depends only on c_1 .

Next, we prove the necessity. Suppose that D is a c -uniform domain. Then Theorem A implies that there exists a constant c' depending only on c such that for all x and y in D ,

$$k_D(x, y) \leq c' j_D(x, y).$$

By Theorem 1, we see that for all z_1 and z_2 in $G = D \setminus P$,

$$k_G(z_1, z_2) \leq 2^{12}c' j_G(z_1, z_2),$$

which, together with Theorem A, shows that G is a c_1 -uniform domain, where c_1 depends only on c . \square

Acknowledgement. This research was carried out when the first author was an academic visitor at the University of Turku and supported by the Academy of Finland grant of the second author with the Project number 2600066611. The authors are indebted to the referee for his/her valuable suggestions.

REFERENCES

1. L. V. AHLFORS, Quasiconformal reflections, *Acta Math.*, **109** (1963), 291–301.
2. F. W. GEHRING AND B. G. OSGOOD, Uniform domains and the quasi-hyperbolic metric, *J. Analyse Math.*, **36** (1979), 50–74.
3. F. W. GEHRING AND B. P. PALKA, Quasiconformally homogeneous domains, *J. Analyse Math.*, **30** (1976), 172–199.
4. P. HÄSTÖ, Z. IBRAGIMOV, D. MINDA, S. PONNUSAMY AND S. SAHOO, Isometries of some hyperbolic-type path metrics, and the hyperbolic medial axis. In the tradition of Ahlfors-Bers. IV, 63–74, *Contemp. Math.*, 432, Amer. Math. Soc., Providence, RI, 2007.
5. J. HEINONEN AND P. KOSKELA, Definitions of quasiconformality, *Invent. Math.*, **120** (1995), 61–79.
6. M. HUANG, S. PONNUSAMY AND X. WANG, Decomposition and removability properties of John domains, *Proc. Indian Acad. Sci. (Math. Sci.)*, **118** (2008), 357–370.
7. R. KLÉN, Local convexity properties of quasihyperbolic balls in punctured space, *J. Math. Anal. Appl.*, **342** (2008), 192–201.
8. R. KLÉN, Y. LI, S. K. SAHOO AND M. VUORINEN, On the stability of φ -uniform domains, *Monatshefte Math.*(to appear) arXiv: 0812.4369v5 [math.MG], 2012.
9. R. KLÉN, A. RASILA, AND J. TALPONEN, Quasihyperbolic geometry in euclidean and Banach spaces, Proc. ICM2010 Satellite Conf. International Workshop on Harmonic and Quasiconformal Mappings (HMQ2010), eds. D. Minda, S. Ponnusamy, N. Shanmugalingam, *J. Analysis*, **18** (2010), 261–278, arXiv: 1104.3745v1 [math.CV].
10. H. LINDÉN, Quasihyperbolic geodesics and uniformity in elementary domains. Dissertation, University of Helsinki, Helsinki, 2005. *Ann. Acad. Sci. Fenn. Math. Diss.* 146 (2005), 50 pp.
11. O. MARTIO, Definitions of uniform domains, *Ann. Acad. Sci. Fenn. Ser. A I Math.*, **5** (1980), 197–205.
12. O. MARTIO AND J. SARVAS, Injectivity theorems in plane and space. *Ann. Acad. Sci. Fenn. Ser. A I Math.* **4** (1979), no. 2, 383–401.
13. A. RASILA AND J. TALPONEN, Convexity properties of quasihyperbolic balls on Banach spaces, *Ann. Acad. Sci. Fenn.* **37** (2012), 215–228, arXiv:1007.3197v1 [math. CV].
14. A. RASILA AND J. TALPONEN, On Quasihyperbolic Geodesics in Banach Spaces, arXiv:1301.0900 [math.CV]
15. J. J. SCHÄFFER, Inner diameter, perimeter, and girth of spheres. *Math. Ann.*, **173**(1967), 59–82.
16. J. VÄISÄLÄ, Uniform domains, *Tohoku Math. J.*, **40** (1988), 101–118.
17. J. VÄISÄLÄ, Free quasiconformality in Banach spaces. I, *Ann. Acad. Sci. Fenn. Ser. A I Math.*, **15** (1990), 355–379.
18. J. VÄISÄLÄ, Free quasiconformality in Banach spaces. II, *Ann. Acad. Sci. Fenn. Ser. A I Math.*, **16** (1991), 255–310.
19. J. VÄISÄLÄ, Relatively and inner uniform domains, *Conformal Geom. Dyn.*, **2** (1998), 56–88.
20. J. VÄISÄLÄ, The free quasiworld, Freely quasiconformal and related maps in Banach spaces, *Quasiconformal geometry and dynamics (Lublin 1996)*, *Banach Center Publications*, Vol. 48, Polish Academy of Science, Warsaw, ed. by Bogdan Bojarski, Julian Lawrynowicz, Olli Martio, Matti Vuorinen and Józef Zajac, 1999, 55–118.
21. J. VÄISÄLÄ, Quasihyperbolic geodesics in convex domains, *Results Math.*, **48** (2005), 184–195.
22. J. VÄISÄLÄ, Quasihyperbolic geometry of planar domains. *Ann. Acad. Sci. Fenn. Math.* **34** (2009), 447–473.

23. M. VUORINEN, Capacity densities and angular limits of quasiregular mappings. *Trans. Amer. Math. Soc.* **263** (1981), 343–354.
24. M. VUORINEN, Conformal invariants and quasiregular mappings, *J. Analyse Math.*, **45** (1985), 69–115.
25. M. VUORINEN, Conformal geometry and quasiregular mappings (Monograph, 208 pp.), Lecture Notes in Math. Vol. 1319, Springer-Verlag, 1988.

M. HUANG, DEPARTMENT OF MATHEMATICS, HUNAN NORMAL UNIVERSITY, CHANGSHA, HUNAN 410081, PEOPLE'S REPUBLIC OF CHINA
E-mail address: mzhuang79@163.com

M. VUORINEN, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TURKU, 20014 TURKU, FINLAND
E-mail address: vuorinen@utu.fi

X. WANG, DEPARTMENT OF MATHEMATICS, HUNAN NORMAL UNIVERSITY, CHANGSHA, HUNAN 410081, PEOPLE'S REPUBLIC OF CHINA
E-mail address: xtwang@hunnu.edu.cn