

A method to determine algebraically integral Cayley digraphs on finite Abelian group

LI Fei *

(School of Statistics and Applied Mathematics,
Anhui University of Finance and Economics,
Bengbu, 233030, Anhui, P.R.China)

Abstract Researchers in the past have studied eigenvalues of Cayley digraphs or graphs. We are interested in characterizing Cayley digraphs on a finite Abelian group G whose eigenvalues are algebraic integers in a given number field K . And we succeed in finding a method to do so by proving Theorem 1. Also, the number of such Cayley digraphs is computed.

Keywords Cayley digraph, integral graph, algebraic integer, Abelian group

Mathematics Subject Classification 05C20

1 Introduction and Main Results

In this paper, the finite digraphs without loops and multiple edges are considered. Let G be a group and let S be a subset of G that does not contain the identity. Then the Cayley digraph (see [8]) $D(G, S)$ is the digraph with vertex set G and edge set $E(D(G, S)) = \{gh \mid hg^{-1} \in S\}$. If S is inverse-closed, then the definition reduces

* Partially supported by Foundation of NSSFC(No.13CTJ006). E-mail: cczxf@163.com

to that of a Cayley graph. When G is cyclic, $D(G, S)$ is called circulant digraphs. For the prerequisites for Cayley digraphs we refer to [5, 8].

A digraph is called algebraically integral over a number field K if its spectrum consists of some algebraic integers of K . Since 1974, Harary and Schwenk (see [9]) began to research integral graphs, which in fact are algebraically integral over the rational number field \mathbb{Q} . There are some literature studying integral Cayley graphs and the spectrum of Cayley digraphs(see e.g., [1][2][3][4][6][14]).

We call $D(G, S)$ Gaussian integral if it is algebraically integral over Gaussian field $\mathbb{Q}(i)$. There are several authors who have studied digraphs with Gaussian spectra(see[7][11][15]). And in our previous results, algebraically integral circulant digraphs was discussed completely(see [13]).

In this paper, we obtain the necessary and sufficient conditions for algebraically integral Cayley digraphs on finite Abelian groups. Also, the number of such algebraically integral Cayley digraphs is calculated.

Let n be the order of G . By the theorem of finite Abelian groups, G is isomorphic to the direct sum of some cyclic groups. In the following, we assume that $G = C_1 \oplus C_2 \oplus \cdots \oplus C_m$, where $C_i = \mathbb{Z}/n_i\mathbb{Z}$ (n_i not necessary be a power of a prime number)for some integer $n_i, 1 \leq i \leq m$. We assume without loss of generality the fixed number field K is contained in the n -th cyclotomic field $\mathbb{Q}(\zeta_n)$. The main results are stated as follows.

A. (see Theorem 1.) The Cayley digraph $D(G, S)$ is algebraically integral over K if and only if S is a union of some orbits Hg_i 's.

B. (see Proposition 3.) For $G = (\mathbb{Z}/n_1\mathbb{Z}) \oplus (\mathbb{Z}/n_2\mathbb{Z}) \oplus \cdots \oplus (\mathbb{Z}/n_m\mathbb{Z})$,

there are at most $2^{r(G,K)}$ Cayley digraphs algebraically integral over K .

Remark. Based on Theorem 1, we can get a method to determine integral and Gaussian integral Cayley digraphs on a finite Abelian group completely.

2 Proof of Main Theorem

Let $H = Gal(\mathbb{Q}(\zeta_n)/K)$ be the Galois group of $\mathbb{Q}(\zeta_n)$ over K . Since $H \subseteq Gal(\mathbb{Q}(\zeta_n)/\mathbb{Q}) \cong (\mathbb{Z}/n\mathbb{Z})^*$, for each $\sigma \in H$, there is an element $a \in (\mathbb{Z}/n\mathbb{Z})^*$ such that $\sigma(\zeta_n) = \zeta_n^a$. Notice that $n = n_1 n_2 \cdots n_m$. For each n_i , we define a group operation π_i of H on $\mathbb{Z}/n_i\mathbb{Z}$ by $\pi_i(\sigma, x) = \sigma(x) = ax \pmod{n_i}$ for $x \in \mathbb{Z}/n_i\mathbb{Z}$. Then we have a operation of H on G by $\sigma(x_1, x_2, \cdots, x_m) = (\pi_1(\sigma, x_1), \pi_2(\sigma, x_2), \cdots, \pi_m(\sigma, x_m))$ for $(x_1, x_2, \cdots, x_m) \in G$. Using the orbit decomposition formula, $G \setminus \{0\}$ is the disjoint union of the distinct orbits, and we can write $G \setminus \{0\} = \bigsqcup_{i \in I} Hg_i$, where I is some index set, and the g_i are elements of distinct orbits.

We obtain a necessary and sufficient condition for the Cayley digraph $D(G, S)$ to be algebraically integral as in the following theorem.

Theorem 1. The Cayley digraph $D(G, S)$ is algebraically integral over K if and only if S is a union of some orbits Hg_i 's.

In the following, we give the proof of Theorem 1 in the case of $m = 2$, i.e., G is a direct sum of two summands.

Let \widehat{G} denote the group of multiplicative homomorphisms from G to \mathbb{C}^* (see [10]), where \mathbb{C} is the complex field. Then we have isomorphism $G \cong \widehat{G}$. For each $(a, b) \in G$, there is an element $\chi_{(a,b)} \in \widehat{G}$ such that $\chi_{(a,b)}(s, t) = \zeta_{n_1}^{as} \times \zeta_{n_2}^{bt}$, $(s, t) \in G$.

The spectrum of Cayley digraph $D(G, S)$ is given by [3],

$$\text{spec}(D(G, S)) = \{\lambda_{(0,0)}, \lambda_{(0,1)}, \dots, \lambda_{(0,n_2-1)}, \lambda_{(1,0)}, \dots, \lambda_{(n_1-1,0)}, \dots, \lambda_{(n_1-1,n_2-1)}\},$$

where $\lambda_{(a,b)} = \sum_{(s,t) \in S} \chi_{(a,b)}(s, t) = \sum_{(s,t) \in S} \zeta_{n_1}^{as} \cdot \zeta_{n_2}^{bt}$.

Proposition 1. If S is a union of some orbits Hg_i 's, then $D(G, S)$ is algebraically integral over K .

Proof. For each orbit Hg_i , we have $\sigma Hg_i = \{\sigma(hg_i) \mid h \in H\} = Hg_i, \sigma \in H$.

So $\sigma S = S$, since S is a union of some orbits Hg_i 's. Thus

$$\begin{aligned} \sigma(\lambda_{(a,b)}) &= \sum_{(s,t) \in S} \sigma(\zeta_{n_1}^{as} \cdot \zeta_{n_2}^{bt}) = \sum_{(s,t) \in S} \sigma(\zeta_{n_1}^{as}) \cdot \sigma(\zeta_{n_2}^{bt}) \\ &= \sum_{(s,t) \in S} \zeta_{n_1}^{a\sigma(s)} \cdot \zeta_{n_2}^{b\sigma(t)} = \lambda_{(a,b)}, \end{aligned}$$

for every element $\sigma \in H$. By the Galois theory (see [12]) for finite Galois extensions, we get $\lambda_{(a,b)} \in K$. Notice that $\lambda_{(a,b)}$ are algebraic integers, hence $D(G, S)$ is algebraically integral over K . ■

In the other hand, we have the following proposition.

Proposition 2. If $D(G, S)$ is algebraically integral over K , then S is a union of some orbits Hg_i 's .

Before the proof, we need some preparations.

Let Γ be a $(n - 1)$ -order square matrix with the row and column index in $G \setminus \{(0, 0)\}$. Namely, we suppose $\Gamma = (\gamma_{(a,b)(\alpha,\beta)})$, where $\gamma_{(a,b)(\alpha,\beta)}$ is the entry of the (a, b) -th row and (α, β) -th column, with $(a, b), (\alpha, \beta) \in G \setminus \{(0, 0)\}$. And $\gamma_{(a,b)(\alpha,\beta)} = \chi_{(a,b)}(\alpha, \beta) = \zeta_{n_1}^{a\alpha} \times \zeta_{n_2}^{b\beta}$. For the matrix Γ , we have the following lemma.

Lemma 1. The matrix Γ is nonsingular.

Proof. Let χ be a nontrivial character of finite Abelian group G , we know that $\sum_{g \in G} \chi(g) = 0$. Then the sum of all entries in each row of Γ is -1 . Hence, to prove this lemma, it suffices to show the $n \times n$ block matrix $\Gamma' = \begin{pmatrix} 1 & 1 \\ 1 & \Gamma \end{pmatrix}$ is invertible.

Let F be the space of complex valued class function on G . All the elements of \widehat{G} form an orthonormal basis of F (see [10]). Suppose there exist n complex numbers $k_{(a,b)}$, $(a,b) \in G$ such that $\sum_{(a,b) \in G} k_{(a,b)} R_{(a,b)} = (0, 0, \dots, 0)$, where $R_{(a,b)}$ is the (a,b) -th row of the matrix Γ' . Then we have the class function $\sum_{(a,b) \in G} k_{(a,b)} \chi_{(a,b)} = 0$. So $k_{(a,b)} = 0$ for all $(a,b) \in G$, which shows that the row vectors of Γ' are linearly independent and so Γ' is invertible. ■

Let τ be such a $(n-1)$ -dimension column vector as

$$\tau = (v_{(0,1)}, v_{(0,2)}, \dots, v_{(0,n_2-1)}, v_{(1,0)}, \dots, v_{(n_1-1,0)}, \dots, v_{(n_1-1,n_2-1)})^T$$

with $v_{(a,b)} = 1$ for $(a,b) \in S$ and 0 otherwise. It is easy to see that

$$\Gamma\tau = (\lambda_{(0,1)}, \lambda_{(0,2)}, \dots, \lambda_{(0,n_2-1)}, \lambda_{(1,0)}, \dots, \lambda_{(n_1-1,0)}, \dots, \lambda_{(n_1-1,n_2-1)})^T.$$

Let τ_i be the $(n-1)$ -dimension column vector for the orbit Hg_i just as τ for S . We denote W the vector space $\{\omega \in K^{n-1} \mid \Gamma\omega \in K^{n-1}\}$ and $V \subset K^{n-1}$ the the vector space spanned by the vectors $\{\tau_i, i \in I\}$. We obtain the following lemma for W and V .

Lemma 2. W and V are the same vector space.

Proof. By Proposition 1, $\Gamma\tau_i \in K^{n-1}, i \in I$. So $V \subset W$. Let $\omega \in W$ and

$\omega = (\omega_{(0,1)}, \omega_{(0,2)}, \dots, \omega_{(0,n_2-1)}, \omega_{(1,0)}, \dots, \omega_{(n_1-1,0)}, \dots, \omega_{(n_1-1,n_2-1)})^T$, $u = \Gamma\omega = (u_{(0,1)}, u_{(0,2)}, \dots, u_{(0,n_2-1)}, u_{(1,0)}, \dots, u_{(n_1-1,0)}, \dots, u_{(n_1-1,n_2-1)})^T$. First, we show that $u_{(a,b)} = u_{(c,d)}$ if $(a, b), (c, d)$ in the same orbit Hg_i . Because $(a, b), (c, d) \in Hg_i$, there exist an element $\sigma \in H$ such that $\sigma(a, b) = (c, d)$, namely, $\sigma(a) = c(\text{mod } n_1)$ and $\sigma(b) = d(\text{mod } n_2)$. In fact,

$$\begin{aligned}
u_{(a,b)} &= \sigma(u_{(a,b)}) = \sigma\left(\sum_{(k,l) \in G \setminus \{0\}} \omega_{(k,l)} \chi_{(a,b)}(k, l)\right) = \sum_{(k,l) \in G \setminus \{0\}} \omega_{(k,l)} \sigma(\chi_{(a,b)}(k, l)) \\
&= \sum_{(k,l) \in G \setminus \{0\}} \omega_{(k,l)} \sigma(\zeta_{n_1}^{ak} \cdot \zeta_{n_2}^{bl}) = \sum_{(k,l) \in G \setminus \{0\}} \omega_{(k,l)} \zeta_{n_1}^{\sigma(a)k} \cdot \zeta_{n_2}^{\sigma(b)l} \\
&= \sum_{(k,l) \in G \setminus \{0\}} \omega_{(k,l)} \zeta_{n_1}^{ck} \cdot \zeta_{n_2}^{dl} = u_{(c,d)},
\end{aligned}$$

which implies that $\Gamma(W) \subset V$. Notice that the matrix Γ is nonsingular by Lemma 1 and $V \subset W$. Hence $\dim W = \dim V$, and $W = V$. ■

Now it comes to prove Proposition 2.

Proof of Proposition 2. Since $D(G, S)$ is algebraically integral over K , $\Gamma\tau \in K^{n-1}$. We have $\tau \in W$. By Lemma 2, $\tau \in V$ and $\tau = \sum_{i \in I} c_i \tau_i$ for some coefficients $c_i \in K$. By the construction of τ and τ_i 's, we conclude that S is the union of the Hg_i 's with $c_i = 1$. The proof is completed. ■

Remark Merging Proposition 1 and Proposition 2 together, we have Theorem 1 in the case of $m = 2$. By [3], $\text{spec}(D(G, S)) = \{\lambda_g \mid g = (g_1, g_2, \dots, g_m) \in G\}$, where $\lambda_g = \sum_{(s_1, s_2, \dots, s_m) \in S} \zeta_{n_1}^{g_1 s_1} \zeta_{n_1}^{g_2 s_2} \cdot \zeta_{n_2}^{g_m s_m}$. So far, we have given the ideas and methods to the proof of Theorem 1. It is not hard to find these ideas and methods can be applied to the general case. In another words, with the formula of spectrum, we can prove Theorem 1 for any $m \in \mathbb{Z}^+$ (\mathbb{Z}^+ , the set of all positive integers) in the same way as above. So we obtain the result of Theorem 1.

3 Calculating the number of algebraically integral Cayley digraphs

Let $G_{n_i} (1 \leq i \leq m)$ be the set of all the orbits of $\mathbb{Z}/n_i\mathbb{Z}$ under H . Denote P_G the collection of Cartesian product $\{P = p_1 \times p_2 \times \cdots \times p_m \mid P \neq 0, p_i \in G_{n_i}\}$. For a Cartesian product $P \in P_G$, choose one element $\rho \in P, \rho = (a_1, a_2, \cdots, a_m)$. Define $\mathbb{Q}(P)$ the cyclotomic field

$$\mathbb{Q}(\zeta_{n_1}^{a_1}, \zeta_{n_2}^{a_2}, \cdots, \zeta_{n_m}^{a_m}) = \mathbb{Q}(\zeta_{n_1}^{a_1}) \cdot \mathbb{Q}(\zeta_{n_2}^{a_2}) \cdots \mathbb{Q}(\zeta_{n_m}^{a_m}),$$

$[\mathbb{Q}(P) : \mathbb{Q}(P) \cap K]$ the dimension of $\mathbb{Q}(P)$ as vector space over $\mathbb{Q}(P) \cap K$ and $|P|$ the cardinal number of P . It is easy to see that $\mathbb{Q}(P)$ is well-defined. By Galois theory (see [12]) in number field, we have $|P| = \prod_{i=1}^m [\mathbb{Q}(\zeta_{n_i}^{a_i}) : \mathbb{Q}(\zeta_{n_i}^{a_i}) \cap K]$.

Let f be the map $Gal(\mathbb{Q}(P)/\mathbb{Q}(P) \cap K) \rightarrow \prod_{i=1}^m Gal(\mathbb{Q}(\zeta_{n_i}^{a_i})/\mathbb{Q}(\zeta_{n_i}^{a_i}) \cap K)$ by restriction, namely, $\sigma \mapsto (\sigma|_{\mathbb{Q}(\zeta_{n_1}^{a_1})}, \sigma|_{\mathbb{Q}(\zeta_{n_2}^{a_2})}, \cdots, \sigma|_{\mathbb{Q}(\zeta_{n_m}^{a_m})})$. It is easy to see that f is injective. So every orbit contained in the Cartesian product P has $[\mathbb{Q}(P) : \mathbb{Q}(P) \cap K]$ elements, which implies that P can be divided equally into $\frac{|P|}{[\mathbb{Q}(P) : \mathbb{Q}(P) \cap K]}$ orbits under H . Totally, we get the following Lemma.

Lemma 3. Under the operation of H , the Cartesian product P described as above is divided into $\frac{\prod_{i=1}^m [\mathbb{Q}(\zeta_{n_i}^{a_i}) : \mathbb{Q}(\zeta_{n_i}^{a_i}) \cap K]}{[\mathbb{Q}(P) : \mathbb{Q}(P) \cap K]}$ orbits.

Let $r(G, K) = \sum_{P \in P_G} \frac{\prod_{i=1}^m [\mathbb{Q}(\zeta_{n_i}^{a_i}) : \mathbb{Q}(\zeta_{n_i}^{a_i}) \cap K]}{[\mathbb{Q}(P) : \mathbb{Q}(P) \cap K]}$. By Lemma 3, $r(G, K)$ is the orbits number of group operation of H on G . So we obtain the following proposition by Theorem 1.

Proposition 3. For $G = (\mathbb{Z}/n_1\mathbb{Z}) \oplus (\mathbb{Z}/n_2\mathbb{Z}) \oplus \cdots \oplus (\mathbb{Z}/n_m\mathbb{Z})$, there are at most $2^{r(G, K)}$ Cayley digraphs algebraically integral over K .

References

1. RC. Alperin, BL. Peterson, Integral Sets and Cayley Graphs of Finite Groups. Electron. J. Combin, 2012,19: #P44.
2. J.Sander, T.Sander, The Exact Maximal Energy of Integral Circulant Graphs with Prime Power Order. Contributions to Discrete Mathematics, 2013, 8(2):19-40.
3. K. Babai, Spectra of Cayley Graphs. Journal of Combinatorial Theory, Series B, 1979, 27: 180-189.
4. K. Balinska, D.Cvetkovic, Z.Radosavljevic, S.Simic, D.Stevanovic, A Survey on Integral Graphs. Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat, 2002,13: 42-65.
5. N. Biggs, Algebraic Graph theory. Amsterdam: North-Holland, 1985.
6. W. G.Bridges, R. A.Mena, Rational G-matrices with rational eigenvalues. Journal of Combinatorial Theory, Series A, 1982,32(2): 264-280.
7. F. Esser, F. Harary, Digraphs with real and Gaussian spectra. Discrete Appl. Math, 1980, 2: 113-124.
8. C. Godsil, G. Royle, Algebraic Graph theory. New York: Springer-Verlag, 2001.
9. F. Harary, A. Schwenk, Which graphs have integral spectra, in: R. Bari, F. Harary (Eds.), Graphs and Combinatorics. Berlin: Springer-Verlag, 1974.

10. J. P. Serre, Linear Representations of Finite Groups. New York: Springer-Verlag, 1977.
11. Y. Xu, J. X. Meng, Gaussian integral circulant digraphs. Discrete Mathematics, 2011, 311: 45-50.
12. S. Lang, Algebra. 3rd edition, New York: Springer-verlag, 2002.
13. Fei Li, Circulant Digraphs Integral over Number Fields. Discrete Mathematics, 2013, 313: 821-823.
14. W. So, Integral circulant graphs, Discrete Mathematics, 2005, 306: 153-158.
15. Xiangdong Hou, On the G-Matrices with Entries and Eigenvalues in $\mathbb{Q}(i)$, Graphs and Combinatorics, 1992, 8:53-64.