

On 021-Avoiding Ascent Sequences

William Y.C. Chen

Center for Combinatorics, LPMC-TJKLC, Nankai University
Tianjin 300071, P.R. China, chen@nankai.edu.cn

Alvin Y.L. Dai

Center for Combinatorics, LPMC-TJKLC, Nankai University
Tianjin 300071, P.R. China, alvin@cfc.nankai.edu.cn

Theodore Dokos

Department of Mathematics, University of California, Los Angeles
Box 951555, Los Angeles, CA 90095-1555, USA, t.dokos@gmail.com

Tim Dwyer

Department of Mathematics, Dartmouth College,
Hanover, NH 03755, USA, jtimdwyer@gmail.com

Bruce E. Sagan

Department of Mathematics, Michigan State University,
East Lansing, MI 48824-1027, USA, sagan@math.msu.edu

Abstract

Ascent sequences were introduced by Bousquet-Mélou, Claesson, Dukes and Kitaev in their study of $(\mathbf{2} + \mathbf{2})$ -free posets. An ascent sequence of length n is a nonnegative integer sequence $x = x_1x_2 \dots x_n$ such that $x_1 = 0$ and $x_i \leq \text{asc}(x_1x_2 \dots x_{i-1}) + 1$ for all $1 < i \leq n$, where $\text{asc}(x_1x_2 \dots x_{i-1})$ is the number of ascents in the sequence $x_1x_2 \dots x_{i-1}$. We let \mathcal{A}_n stand for the set of such sequences and use $\mathcal{A}_n(p)$ for the subset of sequences avoiding a pattern p . Similarly, we let $S_n(\tau)$ be the set of τ -avoiding permutations in the symmetric group S_n . Duncan and Steingrímsson have shown that the ascent statistic has the same distribution over $\mathcal{A}_n(021)$ as over $S_n(132)$. Furthermore, they conjectured that the pair (asc, rlm) is equidistributed over $\mathcal{A}_n(021)$ and $S_n(132)$ where rlm is the right-to-left minima statistic. We prove this conjecture by constructing a bivariate-preserving bijection.

Keywords: 021-avoiding ascent sequence, 132-avoiding permutation, right-to-left minimum, number of ascents, bijection

AMS subject classification: 05A05, 05A19

1 Introduction

The objective of this note is to establish a bijection which leads to the equidistribution of the pair of statistics (asc, rlm) over 021-avoiding ascent sequences and over 132-avoiding

permutations. This confirms a conjecture posed by Duncan and Steingrímsson [5].

Let us give an overview of the notation and terminology. Let S_n denote the set of permutations of $[n]$, where $[n] = \{1, 2, \dots, n\}$. Given a permutation $\pi \in S_n$ and a permutation $\tau \in S_k$, we say that a subsequence $\pi_{i_1}\pi_{i_2}\dots\pi_{i_k}$, $1 \leq i_1 < i_2 < \dots < i_k \leq n$, of π is of *pattern* τ if it is order isomorphic to τ , that is, this subsequence has the same relative order as τ . If π does not contain any subsequence of pattern τ , then we say that π *avoids* τ , or π is τ -*avoiding*. We denote by $S_n(\tau)$ the set of τ -avoiding permutations in S_n . For example, the permutation 763894512 contains the subsequence 3952 of pattern 2431, but it is 1234-avoiding. Pattern avoiding permutations have been intensively studied in recent years from many points of view, see [1, 6, 7].

Ascent sequences were introduced by Bousquet-Mélou, Claesson, Dukes and Kitaev [2]. For a sequence $x = x_1x_2\dots x_n$ of nonnegative integers, we say that an index i ($1 \leq i < n$) is an *ascent* if $x_i < x_{i+1}$. We denote by $\text{asc}(x)$ or merely $\text{asc } x$ the number of ascents of x . A sequence $x = x_1x_2\dots x_n$ is called an *ascent sequence* if $x_1 = 0$ and

$$x_i \leq \text{asc}(x_1x_2\dots x_{i-1}) + 1$$

for all $1 < i \leq n$. For example, $x = 010122$ is an ascent sequence while $x = 010142$ is not since $x_5 = 4 > \text{asc}(0101) + 1 = 3$. We let \mathcal{A}_n denote the set of ascent sequences of length n . For an ascent sequence, a *pattern* is a word on a nonnegative integers $\{0, 1, \dots, k\}$, where each element i appears at least once. Containment and avoidance of patterns for ascent sequences are defined in the same way as for permutations. For example, the ascent sequence 01231234 has five occurrences of the pattern 001, namely, the subsequences 112, 113, 114, 223, 224, and the ascent sequence 01012203 is 021-avoiding. We denote by $\mathcal{A}_n(p)$ the set of ascent sequences of length n avoiding pattern p .

In addition to the ascent statistic, we will be interested in the number of right-to-left minima. A *right-to-left minimum* of any sequence x of nonnegative integers is an index i such that $x_i < x_j$ for all $j > i$. The number of right-to-left minima of x is denoted by $\text{rlm}(x) = \text{rlm } x$. For example, $\text{rlm}(010122) = 3$.

Ascent sequences are closely connected to $(\mathbf{2} + \mathbf{2})$ -free posets [2], upper-triangular matrices [4], Stoimenow's matchings [3], and the Catalan numbers C_n [5]. In particular a poset is called $(\mathbf{2} + \mathbf{2})$ -*free* if it does not contain an induced subposet which is isomorphic to the disjoint union of two 2-element chains. Bousquet-Mélou, Claesson, Dukes and Kitaev [2] found a bijection from $(\mathbf{2} + \mathbf{2})$ -free posets to ascent sequences which maps the number of levels of the poset to the number of ascents of the sequence. Dukes and Parviainen [4] established a bijection between ascent sequences and nonnegative upper-triangular matrices. Duncan and Steingrímsson [5] have shown that $\#\mathcal{A}_n(p) = C_n$ for any of the patterns $p = 101, 0101, \text{ or } 021$. It is well known that $\#S_n(132) = C_n$ and Duncan-Steingrímsson also showed that the ascent statistic is equidistributed over $\mathcal{A}_n(021)$ and $S_n(132)$. Furthermore, they proposed the following conjecture.

Conjecture 1.1. *The bivariate statistic (asc, rlm) has the same distribution over $\mathcal{A}_n(021)$ and $S_n(132)$.*

The objective of this note is to give a bijective proof of the above conjecture.

2 Proof of the conjecture

In order to construct our bijection, we will need the concept of the special maximum value of an ascent sequence $x = x_1x_2 \dots x_n$. The *special maximum value* of x is the largest integer M such that there is an index i with $x_i = M$ and

$$x_i = \text{asc}(x_1x_2 \dots x_{i-1}) + 1 \tag{2.1}$$

giving us equality in the defining relation for an ascent sequence. To illustrate the notion, if $x = 01013312434$ then $M = 3$. Note that, except for the zero sequence, M will always exist since the first 1 in any nonzero sequence satisfies (2.1). So we define $M = 0$ for a zero sequence.

Also define a *special maximum index* as an index i where x_i satisfies $x_i = M$ as well as condition (2.1). The first index i with $x_i = M$ is always a special maximum index. In fact, if $[i, j]$ is the largest interval of indices starting with the first special maximum index and satisfying $x_i = x_{i+1} = \dots = x_j = M$ then we claim that these are exactly the special maximum indices. To see this, first note that if $k \in [i, j]$ then k is a special maximum index because

$$x_k = x_i = \text{asc}(x_1x_2 \dots x_{i-1}) + 1 = \text{asc}(x_1x_2 \dots x_{k-1}) + 1$$

since there are no ascents between x_i and x_k . To see that no other index can be special maximum, suppose $x_k = M$ with $k \geq j + 2$. Now $x_j > x_{j+1}$ because if $x_j < x_{j+1}$ then the special maximum value would be at least $M + 1$. Thus there must be an ascent between x_{j+1} and x_k so that (2.1) is no longer an equality when $i = k$. Call the special maximum value *unique* if there is only one special maximum index and *repeated* otherwise.

Theorem 2.1. *The bivariate statistic (asc, rlm) has the same distribution over $\mathcal{A}_n(021)$ and $S_n(132)$.*

Proof. We will inductively build a bijection $\phi_n: \mathcal{A}_n(021) \rightarrow S_n(132)$ preserving the bivariate statistic. To do so, we will need decompositions of $\mathcal{A}_n(021)$ and $S_n(132)$ into pieces indexed by smaller subscripts. We will start on the ascent side.

A simple but important observation for what follows is that $p \in \mathcal{A}_n(021)$ if and only if the nonzero entries of p are weakly increasing. We will use this fact to construct a bijection $f = f_n$ between $\mathcal{A}_n(021)$ and the set of pairs

$$\bigcup_{i=1}^n \mathcal{A}_{i-1}(021) \times \mathcal{A}_{n-i}(021).$$

Consider $x \in \mathcal{A}_n(021)$ and suppose first that x has a repeated special maximum value M . Let k be any of the special maximum indices and define

$$f(x) = (\epsilon, z)$$

where ϵ is the empty sequence and z is x with x_k removed. For example, if $x = 01013300304$ then $z = 0101300304$. Clearly z still avoids 021 since its nonzero entries still increase and, since x 's special maximum value was repeated, z still has M as its special maximum value. Since the special maximum value does not change, one can construct an inverse map from $\mathcal{A}_0(021) \times \mathcal{A}_{n-1}(021)$ back to the elements of \mathcal{A}_n which have a repeated special maximum in the obvious way. Finally note that in this case

$$\text{asc } x = \text{asc } z \quad \text{and} \quad \text{rlm } x = \text{rlm } z.$$

Now suppose that x has a unique special maximum value $x_i = M$. Here we let

$$f(x) = (y, z)$$

where $y = x_1 \dots x_{i-1}$ and z is obtained from $x' = x_{i+1} \dots x_n$ by subtracting $M - 1$ from all the nonzero entries. To illustrate, if $x = 0101300304$ then $y = 0101$ and $z = 00102$. It is clear that $y \in \mathcal{A}_{i-1}(021)$. To show that $z \in \mathcal{A}_{n-i}(021)$, we first note that z still has weakly increasing nonzero elements and so avoids 012. We must also demonstrate that z is an ascent sequence. Since the defining condition for an ascent sequence is trivial for zero elements, we need only consider $z_r \neq 0$. But since we have subtracted the same amount from all nonzero entries of x' , the index r is an ascent of z if and only if the index $r+i$ is an ascent of x' . Also, since M was the special maximum value, we have $\text{asc}(x_1 \dots x_i) = M$, $x_i > x_{i+1}$ and $x_{r+i} \leq \text{asc}(x_1 x_2 \dots x_{r+i-1})$ for any $r \geq 1$. It follows that for any $z_r \neq 0$ we have

$$\begin{aligned} z_r &= x_{r+i} - M + 1 \\ &\leq \text{asc}(x_1 \dots x_{r+i-1}) - M + 1 \\ &= \text{asc}(x_1 \dots x_i) + \text{asc}(x_{i+1} \dots x_{r+i-1}) - M + 1 \\ &= \text{asc}(z_1 \dots z_{r-1}) + 1 \end{aligned}$$

which is what we wished to prove. Constructing the inverse of this part of the map is similar to what was done in the first case.

It will be useful to record what happens to our two statistics in the second case defining f . For the ascent statistic, we have everything in place from the previous paragraph and the fact that, by definition of M , $x_i > x_{i-1}$. Thus

$$\begin{aligned} \text{asc } x &= \text{asc}(x_1 \dots x_i) + \text{asc}(x_{i+1} \dots x_n) \\ &= \text{asc}(x_1 \dots x_{i-1}) + 1 + \text{asc}(x_{i+1} \dots x_n) \\ &= \text{asc } y + \text{asc } z + 1. \end{aligned}$$

In terms of right-to-left minima, we distinguish two subcases. If $i \leq n - 1$ then, since $x_{i+1} = 0$, the right-to-left minima of x must occur in the sequence $x_{i+1} \dots x_n$. Since the subtraction of $M - 1$ does not change the positions of these minima, we have

$$\text{rlm } x = \text{rlm } z.$$

On the other hand, if $i = n$ then $z = \epsilon$ and x_n is a right-to-left minimum, giving

$$\text{rlm } x = \text{rlm } y + 1.$$

We will now review the standard decomposition of $S_n(132)$ which gives a bijection g from this set to

$$\bigcup_{i=1}^n S_{i-1}(132) \times S_{n-i}(132).$$

If $\pi \in S_n(132)$ then we write $\pi = \pi_L n \pi_R$ where π_L, π_R are the elements to the left and right of n , respectively. Define the index i by $\pi_i = n$. Then it is well known that $\pi \in S_n(132)$ if and only if π_L, π_R avoid 132 and every element of π_L is bigger than every element of π_R . So we let

$$g(\pi) = (\rho, \sigma)$$

where $\rho \in S_{i-1}(132)$ and $\sigma \in S_{n-i}(132)$ are order isomorphic to π_L and π_R , respectively.

As with the ascent sequence decomposition, we have to consider what happens to our statistics in two separate cases. The first is when $\rho = \epsilon$, equivalently, $i = 1$. So $\pi = n \pi_R$ and so n makes no contribution either to the ascents or right-to-left maxima. It follows that

$$\text{asc } \pi = \text{asc } \sigma \quad \text{and} \quad \text{rlm } \pi = \text{rlm } \sigma$$

just as in the corresponding case for ascent sequences.

Now suppose $1 < i \leq n$. So there will be an ascent ending at n and all other ascents of π correspond to ascents of ρ or ascents of σ . It follows that

$$\text{asc } \pi = \text{asc } \rho + \text{asc } \sigma + 1.$$

For the right-to-left minima we again break into two subcases depending on whether the second component of our bijection is ϵ or not. If $\sigma \neq \epsilon$ then $i < n$ and the right-to-left minima of π are all in π_R because of the relative sizes of the elements of π_R and π_L . This gives

$$\text{rlm } \pi = \text{rlm } \sigma.$$

Now consider $\sigma = \epsilon$ so that $\pi = \pi_L n$. Thus n is a right-to-left minimum of π as is every right-to-left minimum of π_L . So in this subcase

$$\text{rlm } \pi = \text{rlm } \rho + 1.$$

Finally, we construct $\phi_n: \mathcal{A}_n(021) \rightarrow S_n(132)$ as follows. Start with $\phi_0(\epsilon) = \epsilon$. Assuming that ϕ_i has been defined for all $i < n$, we define ϕ_n to be the composition

$$\mathcal{A}_n(021) \xrightarrow{f} \bigcup_{i=1}^n \mathcal{A}_{i-1}(021) \times \mathcal{A}_{n-i}(021) \xrightarrow{h} \bigcup_{i=1}^n S_{i-1}(132) \times S_{n-i}(132) \xrightarrow{g^{-1}} S_n(132)$$

where the restriction of h to $\mathcal{A}_{i-1}(021) \times \mathcal{A}_{n-i}(021)$ is $\phi_{i-1} \times \phi_{n-i}$. It should be clear from the equations derived for asc and rlm when defining f and g that this bijection preserves the bivariate. \square

Acknowledgments. We wish to thank Einar Steingrímsson for helpful comments. This work was supported by the 973 Project, the PCSIRT Project of the Ministry of Education, and the National Science Foundation of China.

References

- [1] M. Bóna, *Combinatorics of Permutations*, CRC Press, 2004.
- [2] M. Bousquet-Mélou, A. Claesson, M. Dukes and S. Kitaev, $(2+2)$ -free posets, ascent sequence and pattern avoiding permutations, *J. Combin. Theory, Ser. A*, 117 (7) (2010), 884-909.
- [3] A. Claesson, M. Dukes and S. Kitaev, A direct encoding of Stoimenow's matchings as ascent sequences, *Australas. J. Combin.*, 49 (2011), 47-59.
- [4] M. Dukes and R. Parviainen, Ascent sequences and upper triangular matrices containing non-negative integers, *Electron. J. Combin.*, 17 (1) (2010), #R53.
- [5] P. Duncan and E. Steingrímsson, Pattern avoidance in ascent sequences, *Electron. J. Combin.*, 18 (2011), #P226.
- [6] C. Krattenthaler, Permutations with restricted patterns and Dyck paths, *Adv. Appl. Math.*, 27 (2001), 510-530.
- [7] R. Simion and F.W. Schmidt, Restricted permutations, *European J. Combin.*, 6 (1985), 383-406.