Monads of regular theories

Stanisław Szawiel, Marek Zawadowski

November 20, 2018

Abstract

We characterize the category of monads on Set and the category of Lawvere theories that are equivalent to the category of regular equational theories.

Contents

1	Introduction	1
2	Presentations of categories of algebras	4
3	Full operads	13
4	Regular Lawvere theories vs regular operads	16
5	Semi-analytic monads vs regular operads	23
6	Equational theories vs regular operads	27
7	Cartesian and weakly cartesian monads	29

1 Introduction

The category of algebras of a (finitary) equational theory can be equivalently described as either a category of models of a Lawvere theory or as a category of algebras of a finitary monad on the category *Set* or a category of algebras of a (generalized) operad, in which not only permutations but all functions between finite sets act on operations. In fact, the four categories of (finitary) equational theories, Lawvere theories, finitary monads on *Set* and (generalized) operads are equivalent. These equivalences induce a correspondence between various subcategories. In [SZ] we have given an intrinsic characterizations of equational theories and Lawvere theories that correspond to the analytic and polynomial monads on *Set*. This was achieved via correspondence with symmetric and rigid operads.

Recall that an equational theory is regular if it has a set of axioms in which each equation contains the same variable on both sides. Thus the theories of monoids and of sup-lattices are regular but the theory of groups is not. The category of regular equational theories and regular morphisms **RegET** was defined in [SZ] but the notion of a regular theory was discovered and studied in universal algebra (cf. [Pl]) and will turn fifty soon.

The main objective of this paper is to describe the categories of regular Lawvere theories **RegLT**, of semi-analytic monads **SanMnd**, and regular operads **RegOp** that correspond to the category of regular theories.

In [SZ] symmetric operads proved to be very useful in describing correspondences of this kind. Full and regular operads (cf. [Tr], Section 2) play the same role here. The category of full operads can be thought of as the category of monoids in the monoidal category $Set^{\mathbb{F}}$, where \mathbb{F} is the skeleton of the category of finite sets, with the substitution tensor. Thus a monoid in $Set^{\mathbb{F}}$ consists of operations that are equipped with not only a multiplication operator but also an action of the whole category \mathbb{F} . As the left Kan extension $Set^{\mathbb{F}} \to \mathbf{End}$ along the inclusion $\mathbb{F} \to Set$ is an equivalence of monoidal categories (**End** is the category of finitary endofunctors on Set) the category of full operads is equivalent to the category of finitary monads on Set. Regular operads can be thought of as monoids in the monoidal category $Set^{\$}$ with substitution tensor, where \$ is the skeleton of the category of finite sets and surjections. Thus in regular operads only surjections act on operations.

We identify the essential image of the left Kan extension $Set^{\mathbb{S}} \to \mathbf{End}$ along inclusion $\mathbb{S} \to Set$, as the category of semi-analytic functors and semi-analytic natural transformations **San**. Semi-analytic functors have similar characterizations as analytic ones; cf. Section 2. A finitary endofunctor on *Set* is semi-analytic iff it preserves pullbacks along monomorphisms. Semi-analytic functors also have presentations via series similar to the series that represent analytic functors. A natural transformation is semi-analytic iff the naturality squares for monomorphism are pullbacks. The category of semi-analytic functors **San** is a monoidal subcategory of **End** and the monoids there form the category of semi-analytic monads equivalent to the category of regular theories; cf. Section 5.

The category of regular Lawvere theories is defined very much in the spirit of the definition of the category of analytic Lawvere theories, cf. Section 2. A regular Lawvere theory is a Lawvere theory with nicely behaving isomorphisms and a factorization system consisting of the class of projections and the class of regular morphisms. Regular morphisms in a Lawvere theory are by definitions those morphisms that are right orthogonal to all projections. A regular interpretation of Lawvere theories is an interpretation of Lawvere theories that preserves regular morphisms. We show that the category of regular Lawvere theories is equivalent to the category of regular operads and hence also to the category of semi-analytic monads and to the category of regular equational theories; cf. Section 6.

The category of semi-analytic monads **SanMnd** contains subcategories of cartesian **CartMnd** and weakly cartesian **wCartMnd** monads (i.e. those that preserve and weakly preserve pullbacks, respectively). One may think that such categories of monads can correspond to some natural subcategories of the category of Lawvere theories and of the category of equational theories. In Section 7 we characterize the subcategories of $Set^{\$}$ that have as their essential images **wCartMnd** and **CartMnd**. As these characterizations are a bit technical we do not try to rephrase those conditions in terms of either Lawvere theories or equational theories.

In this way we shall describe the second level (level r) of the following diagram





The vertical lines denote adjoint equivalences. Thus up to equivalence there are only four categories in it, one on each level. One equivalent to the category of all finitary monads on Set, second equivalent to the category of all semi-analitic monads on Set, third equivalent to the category of all analytic monads on Set, and forth equivalent to the category of all polynomial monads on Set. These levels are denoted by letters f, r, a, and p, respectively. Thus all four columns of equational theories, Lawvere theories, monads and operads are equivalent. These columns are denoted by letters e, l, m and o, respectively. The vertical functors heading up, in all columns but the column of operads, are inclusions of subcategories. In the column of operads the functors heading up are more like free extensions of the actions. The lower functors are full embeddings and the upper are embeddings that are full on isomorphisms. The vertical functors heading down, the right adjoints to those heading up are monadic. All the squares in the diagram commute up to canonical isomorphisms. The notation concerning categories involved is displayed in the above

diagram.

The notation concerning functors is not on the diagram but it is meant to be systematic referring to levels and columns they 'connect'. The horizontal functors are denoted using letters from both columns they connect; the codomain by the script letter, the domain by its subscript, and the level is denoted by superscript. Thus the functor **AnMnd** \rightarrow **AnLT** will be denoted by \mathcal{L}_m^a . We usually drop superscripts when it does not lead to confusion. Thus we can write, for example, $\mathcal{E}_o = \mathcal{E}_o^p : \mathbf{RiOp} \rightarrow \mathbf{RiET}$. The vertical functors heading up are denoted by the script letter \mathcal{P} with superscript indicating the column and subscript indicating the level of the codomain. The vertical functors heading down are denoted by the script letter \mathcal{Q} with subscript and superscript as with those heading up. Thus we have, for example, functors $\mathcal{P} = \mathcal{P}^o = \mathcal{P}_a^o : \mathbf{RiOp} \rightarrow \mathbf{SOp}$ and $\mathcal{Q} = \mathcal{Q}_a = \mathcal{Q}_a^m : \mathbf{Mnd} \rightarrow \mathbf{AnMnd}$. We will also refer to various diagonal morphisms and then we need to extend the notation concerning vertical functors by specifying both the columns of the domain and the codomain. For example, we write $\mathcal{P}_f^{ol} : \mathbf{SOp} \rightarrow \mathbf{LT}$ to denote one such functor and its right adjoint will be denoted by $\mathcal{Q}_a^{lo} : \mathbf{LT} \rightarrow \mathbf{SOp}$. In principle this notation will leave the codomain not always uniquely specified but in practice it it sufficient, and in fact, usually much less is needed.

Notation

 $n = \{0, \ldots, n-1\}, [n] = \{0, \ldots, n\}, (n] = \{1, \ldots, n\}, \omega$ - denotes the set of natural numbers. The set X^n is interpreted as $X^{(n)}$ when convenient. The skeletal category equivalent to the category of finite sets Set_{fin} will be denoted by \mathbb{F} . We will be assuming that the objects of \mathbb{F} are sets (n) for $n \in \omega$. The subcategories of \mathbb{F} with the same objects as \mathbb{F} but having as objects bijection, surjections and injections will be denoted by \mathbb{B} , \mathbb{S} , \mathbb{I} , respectively. When S_n acts on a set A_n on the right and on the set B_n on the left, the set $A \otimes_n B$ is the usual tensor product of S_n -sets.

2 Presentations of categories of algebras

Equational theories

The category of regular theories, denoted by **RegET**, was introduced in [SZ]. It is a subcategory of all equational theories, denoted **ET**.

Operads

The symmetric operads provide yet another way of presenting models of an equational theory. This kind of presentation is usually very convenient, however the models defined by such operads are more specific than models of arbitrary equational theories. For example, if \mathcal{O} is a symmetric operad then the free algebra functor $Set \to Alg(\mathcal{O})$ preserves weak wide pullbacks. Below we extend the definition of an operad so that it captures all the equational theories but still keeps the operadic flavor. The main difference is that instead of having just symmetric groups acting on sets of operations we have actions of the morphisms of the whole skeleton of the category of finite sets \mathbb{F} . Symmetric operads can be thought of as monoids for the substitution tensor on the category $Set^{\mathbb{F}}$. By **End** we denote the category of finitary endofunctors of Set. It is a strict monoidal category with the tensor being composition. The substitution tensor on $Set^{\mathbb{F}}$ makes the equivalence of categories $Set^{\mathbb{F}} \to \mathbf{End}$, given by left Kan extension, a strong monoidal equivalence. This immediately shows that the category of \mathbb{F} -operads is equivalent to the category of finitary monads on Set. The definition below was first explicitly spelled out in [Tr].

A full operad (or \mathbb{F} -operad) \mathcal{O} consists a family of sets \mathcal{O}_n , for $n \in \omega$, a unit element $\iota \in \mathcal{O}_1$ for any $k, n, n_1, \ldots, n_k \in \omega$ with $n = \sum_{i=1}^k n_i$ a multiplication operation

$$*: \mathcal{O}_{n_1} \times \ldots \times \mathcal{O}_{n_k} \times \mathcal{O}_k \longrightarrow \mathcal{O}_n$$
$$(b_1, \ldots, b_k, a) \mapsto \langle b_1, \ldots, b_k \rangle * a$$

a left action of the morphisms on operations

$$: \mathbb{F}(n,m) \times \mathcal{O}_n \longrightarrow \mathcal{O}_m$$

for $n, m \in \omega$, such that the multiplication is associative with unit ι and compatible with the category action, i.e. for $a \in \mathcal{O}_n$

$$\langle a \rangle * \iota = a = \langle \iota, \dots, \iota \rangle * a; \quad 1_n \cdot a = a$$

for $a \in \mathcal{O}_n$, $b_i \in O_{k_i}$, $c_{ij} \in \mathcal{O}_{m_{ij}}$, for $i \in (n]$, $j \in (k_i]$ we have

$$\langle c_{1,1},\ldots,c_{1,k_1},\ldots,c_{n,1},\ldots,c_{n,k_n}\rangle * (\langle b_1,\ldots,b_n\rangle * a) =$$

$$= \langle \langle c_{1,1}, \dots, c_{1,k_1} \rangle * b_1, \dots, \langle c_{n,1}, \dots, c_{n,k_n} \rangle * b_n \rangle * a$$

and for $\phi \in \mathbb{F}(n,m)$, $a \in \mathcal{O}_n \ \psi_i \in \mathbb{F}(k_i, l_i)$, $b_i \in \mathcal{O}_{k_i}$ for $i \in (m]$, we have

$$\langle \psi_1 \cdot b_1, \dots, \psi_n \cdot b_n \rangle * (\phi \cdot a) = (\langle \psi_1 \dots, \psi_n \rangle \star \phi) \cdot (\langle b_{\phi(1)}, \dots, b_{\phi(n)} \rangle * a)$$

in \mathcal{O}_l , where

$$\langle \psi_1 \dots, \psi_n \rangle \star \phi : (k] = \sum_{j=1}^n (k_{\phi(j)}] \longrightarrow \sum_{i=1}^m (l_i] = (l]$$

is a function such that j-th summand of the domain $(k_{\phi(j)}]$ is sent to the $\phi(j)$ -th summand $(l_{\phi(j)}]$ of the codomain by the function $\psi_{\phi(j)}$, i.e.

$$\langle j, r \rangle \mapsto \langle \phi(j), \psi_{\phi(j)}(r) \rangle$$

for $j \in (n]$ and $r \in (k_{\phi(j)}]$. This definition refers to the obvious lexicographic order on both $(k] = \sum_{j=1}^{n} (k_{\phi(j)})$ and $(l] = \sum_{i=1}^{m} (l_i)$. It is an extension of the multiplication operation in the operad of symmetries.

$$\sum_{j=1}^{n} (k_{\phi(j)}] \xrightarrow{\langle \psi_1 \dots, \psi_n \rangle \star \phi} \sum_{i=1}^{m} (l_i]$$

$$\kappa_j \uparrow \qquad \qquad \uparrow \lambda_{\phi(j)}$$

$$(k_{\phi(j)}] \xrightarrow{\psi_{\phi(j)}} (l_{\phi(j)})$$

Remark The operation \star is defined on a family of function indexed by $k, l \in \omega$ between sets:

$$\coprod_{n,m\in\omega,\ \phi\in\mathbb{F}(n,m)}\ \coprod_{k_i,l_i\in\omega,\ k=\sum_{j=1}^n k_{\phi(j)},\ l=\sum_{i=1}^m l_i}\ \prod_{i=1}^m \mathbb{F}(k_i,l_i) \xrightarrow{} \mathbb{F}(k,l)$$

A morphism of full operads $f : \mathcal{O} \to \mathcal{O}'$ is a function that respects arities of operations, unit, compositions, and the actions of functions from \mathbb{F} .

The operation \star , when applied to morphisms in S, returns a morphism in S. Therefore these definitions make sense if we restrict morphisms to surjections i.e. morphisms in S^1 . In this way, we obtain the notion of a *regular operad*, a *morphism of regular operads* and the whole category of regular operads denoted **RegOp**.

We have functors

$$\mathbf{FOp} \xrightarrow{\mathcal{Q}_r^o} \mathbf{RegOp} \xrightarrow{\mathcal{Q}_a^o} \mathbf{SOp}$$

'restricting actions' along the inclusions $\mathbb{B} \to \mathbb{S} \to \mathbb{F}$. They have left adjoints:

$$\mathbf{FOp} \longleftarrow \mathcal{P}_{f}^{o} \qquad \mathbf{RegOp} \longleftarrow \mathcal{P}_{r}^{o} \qquad \mathbf{SOp}$$

We sketch the definitions of those functors below.

For a symmetric operad \mathcal{O} the regular operad $\mathcal{P}_r^o(\mathcal{O})$ has

$$\coprod_{k\in\omega} \mathbb{S}(k,n)\otimes_k \mathcal{O}_k$$

as the set of *n*-ary operations. The unit of $\mathcal{P}_r^o(\mathcal{O})$ is

$$[1,\iota] \in \mathbb{S}(1,1) \otimes_1 \mathcal{O}_1 \subset \prod_{k \in \omega} \mathbb{S}(k,1) \otimes_k \mathcal{O}_k$$

¹Clearly, these definitions make also sense if we restrict morphisms in \mathbb{F} to injections i.e. morphisms in \mathbb{I} . But we will study such operads elsewhere.

The action of the category $\[mathbb{S}\]$ in $\mathcal{P}_r^o(\mathcal{O})$

$$: \mathbb{S}(n,m) \times \coprod_{k \in \omega} \mathbb{S}(k,n) \otimes_k \mathcal{O}_k \longrightarrow \coprod_{k \in \omega} \mathbb{S}(k,n) \otimes_k \mathcal{O}_k$$

is

$$\phi \cdot [f,a] = [\phi \circ f,a]$$

where $\phi: (n] \to (m], f: (k] \to (n]$ are surjection and $a \in \mathcal{O}_k$. The composition in $\mathcal{P}_r^o(\mathcal{O})$

$$\prod_{l_1 \in \omega} \mathbb{S}(l_1, n_1) \otimes_{l_1} \mathcal{O}_{l_1} \times \ldots \times \prod_{l_m \in \omega} \mathbb{S}(l_m, n_m) \otimes_{l_m} \mathcal{O}_{l_m} \times \prod_{k \in \omega} \mathbb{S}(k, m) \otimes_k \mathcal{O}_k \longrightarrow \prod_{k \in \omega} \mathbb{S}(k, n) \otimes_k \mathcal{O}_k$$

is given by

$$\langle [\psi_1, b_1], \dots, [\psi_m, b_m], [\phi, b] \rangle \mapsto [\langle \psi_1, \dots, \psi_m \rangle \star \phi, \langle b_{\phi(1)}, \dots, b_{\phi(k)} \rangle * a]$$

where $\varphi \in S(k, m)$, $a \in \mathcal{O}_k$, $\sum_{i=1}^m n_i = n$, $\psi_i \in S(l_i, n_i)$, $b_i \in \mathcal{O}_{l_i}$, for $i \in (m]$. The definition of the functor \mathcal{P}_a^o on morphisms is left for the reader.

For a regular operad \mathcal{O} the full operad $\mathcal{P}_{f}^{o}(\mathcal{O})$ has

$$\coprod_{k\in\omega}\mathbb{I}(k,n)\otimes_k\mathcal{O}_k$$

as the set of *n*-ary operations. The action of the category \mathbb{F} in $\mathcal{P}_{f}^{o}(\mathcal{O})$

$$: \mathbb{F}(n,m) \times \prod_{k \in \omega} \mathbb{I}(k,n) \otimes_k \mathcal{O}_k \longrightarrow \prod_{k \in \omega} \mathbb{I}(k,n) \otimes_k \mathcal{O}_k$$

is

$$\phi \cdot [f, a] = [f', \phi' \cdot a]$$

where $\phi : (n] \to (m]$ is a function, $f : (k] \to (n]$ is an injection, $a \in \mathcal{O}_k$ and ϕ', f' is the epi-mono factorization of $\phi \circ f$:



The rest of the definition of $\mathcal{P}_{f}^{o}(\mathcal{O})$ is similar to the remaining part of the definition of $\mathcal{P}_{r}^{o}(\mathcal{O})$ given above.

Lawvere theories

The category of Lawvere theories will be denoted by **LT**, see [Lw], [KR], [SZ] for details. \mathbb{F}^{op} is the initial Lawvere theory with the obvious projections. Let **T** be any Lawvere theory. By $\pi : \mathbb{F}^{op} \to \mathbf{T}$ we denote the unique morphism from \mathbb{F}^{op} to **T**. The class of *projections* in **T** is the closure under isomorphisms of the image of the injections in \mathbb{F} under π . A morphism r in T is *regular* iff r is right orthogonal to all projection morphisms in **T**. By a factorization system we mean a factorization system in the sense of [FK], see [CJKP] sec 2.8.

Aut(n) is the set of automorphisms of n in **T**. Recall from [SZ], that a Lawvere theory has simple automorphisms if the canonical function

$$\rho_n: S_n \times Aut(1)^n \longrightarrow Aut(n)$$

such that

$$(\sigma, a_1, \ldots, a_n) \mapsto a_1 \times \ldots \times a_n \circ \pi_\sigma$$

is a bijection, for $n \in \omega$.

A Lawvere theory **T** is a *regular Lawvere theory* iff the projection morphisms and the regular morphisms form a factorization system and **T** has simple automorphisms. A *regular interpretation* of Lawvere theories is a morphism of Lawvere theories that preserves regular morphisms. Thus we have a non-full subcategory of **LT** of regular Lawvere theories with regular interpretations **RegLT**. The theory \mathbb{F}^{op} is regular.

We have inclusion functors

AnLT
$$\xrightarrow{\mathcal{P}_r^l}$$
 RegLT $\xrightarrow{\mathcal{P}_f^l}$ LT

Monads

We introduce here a notion of a semi-analytic monad that is broader then that of an analytic monad but it still retains some combinatorial flavor.

Recall that an analytic (endo)functor on Set can be defined by any of the following conditions

- 1. finitary functor preserving weak wild pullbacks;
- 2. Kan extension of a functor from \mathbb{B} to *Set*;
- 3. functor (isomorphic to one) having an analytic presentation $\sum_{n \in \omega} X^n \otimes_n A_n$, where the *n*-coefficient A_n is a (left) S_n -set for $n \in \omega$.

Similarly we shall define a *semi-analytic* (endo)functor on *Set* as a functor satisfying any of three equivalent conditions (see Proposition 2.1, Theorem 2.2)

- 1. finitary functor preserving pullbacks along monomorphisms;
- 2. Kan extension of a functor from \$ to Set along the inclusion functor $\mathbf{i}_{\$}:\$ \to Set$;
- 3. functor (isomorphic to one) having a semi-analytic presentation $\sum_{n \in \omega} \begin{bmatrix} X \\ n \end{bmatrix} \otimes_n A_n$, where the category S acts on coefficients A_n on the left (see below).

The first two characterizations of the semi-analytic functors are clear. We shall describe the third one and show that it is equivalent to the other two. For the time being, the first definition of a semi-analytic functor is the official one. A natural transformation $\phi : F \to G$ is *semi-cartesian* iff the naturality squares for monomorphisms are pullbacks. The category of semi-analytic functors with the semi-cartesian natural transformations will be denoted by **San**.

Examples

- 1. The functor $\mathcal{P}_{\leq n} : Set \longrightarrow Set$ associating to a set X the set of subsets of X with at most *n*-elements is not analytic, if n > 2, as it can be easily seen that it does not preserve weak pullbacks. However, it preserves pullbacks along monos and hence it is semi-analytic.
- 2. If U is a set, $n \in \omega$ then the functor $(-)_{\leq n}^U : Set \to Set$, associating to a set X the set of functions from U to X with an at most n-element image, is not analytic, if |U| > n > 2. Again it can be easily seen that it does not preserve weak pullbacks. However, it is semi-analytic.
- 3. We will see later that the functor part of any monad on *Set* that comes from a regular equational theory is semi-analytic.

For a set X and $n \in \omega$ the set

 $\left[\begin{array}{c} X\\n\end{array}\right]$

denotes the set of monomorphisms from (n] to X. If X has less than n elements, this set is empty and if X is a finite set, then it has $\binom{|X|}{n} \cdot k!$ elements. The notation is in analogy with the notation $\binom{|X|}{n}$

denoting the set of n-element subsets of X.

Clearly, $\begin{bmatrix} X \\ n \end{bmatrix}$ is not functorial in X on its own but if we build a series with such sets and coefficients that are related by surjections we do get a functor. To see this let

$$A: \mathbb{S} \to Set$$

be a functor, or equivalently a sequence of coefficient sets $\{A_n\}_{n \in \omega}$ on which the category S acts on the left. As S_n acts on $\begin{bmatrix} X\\n \end{bmatrix}$ on the right, by composition it makes sense to form a set

$$\left[\begin{array}{c}X\\n\end{array}\right]\otimes_n A_n$$

and a whole coproduct

$$\hat{A}(X) = \sum_{n \in \omega} \begin{bmatrix} X \\ n \end{bmatrix} \otimes_n A_n$$

The above formula is functorial in X, i.e. \hat{A} can be defined on morphisms as follows. Let $f: X \to Y$ be a function and $[\vec{x}, a]$ an element of $\begin{bmatrix} X \\ n \end{bmatrix} \otimes_n A_n$. We take the epi-mono factorization α, \vec{y} of $f \circ \vec{x}$

$$\begin{array}{ccc} (n] & & \overrightarrow{x} & X \\ \alpha & & & & \downarrow f \\ (m] & & & \overrightarrow{y} & Y \end{array}$$

and we put

$$\hat{A}(f)([\vec{x},a]) = [\vec{y}, A(\alpha)(a)]$$

which is an element in $\begin{bmatrix} Y \\ m \end{bmatrix} \otimes_m A_m$, i.e. in $\hat{A}(Y)$. As the factorization is unique up to an isomorphism, $\hat{A}(f)$ is well defined.

If $\tau : A \to B$ is a natural transformation of functors it induces a natural transformation of functors $\hat{\tau} : \hat{A} \longrightarrow \hat{B}$

as follows. For $[\vec{x}, a]$ in $\begin{bmatrix} X\\ n \end{bmatrix} \otimes_n A_n$ we put

$$\hat{\tau}([\vec{x}, a]) = [\vec{x}, \tau(a)]$$

Proposition 2.1. The functors (-): Set^S \longrightarrow End is well defined and it is isomorphic to the left Kan extension along the inclusion functor $\mathbf{i}_{\mathbf{s}} : \mathbf{S} \rightarrow Set$.

Proof. It is well known (see [CWM]) that the Kan extensions can be calculated with coends. Thus for $A: \mathbb{S} \to Set$ and a function $f: X \to Y$ we have

$$Lan_{\mathbf{i}_{\mathbf{S}}}(A)(X) = \int^{n \in \mathcal{S}} X^n \times A_n \xrightarrow{Lan_{\mathbf{i}_{\mathbf{S}}}(A)(f)} \int^{n \in \mathcal{S}} Y^n \times A_n = Lan_{\mathbf{i}_{\mathbf{S}}}(A)(Y)$$

 $\|\vec{x}, a\| \mapsto \|f \circ \vec{x}, a\|$

where $\|\vec{x}, a\|$ is the equivalence class of the equivalence relation \approx on $\sum_{n \in \omega} X^n \times A_n$ generated by the relation \sim such that

$$\langle \vec{y} \circ \phi, a \rangle \sim \langle \vec{y}, \phi \cdot a \rangle$$

for $\vec{y}: (m] \to X$, $\phi: (n] \to (m] \in \mathbb{S}$ and $a \in A_n$. Let us call an representant $\langle \vec{x}, a \rangle$ of a class $||\vec{x}, a||$ minimal iff \vec{x} is injective. Any element $\langle \vec{x}, a \rangle$ is ~-related to one of form $\langle \vec{y}, f \cdot a \rangle$ where f, \vec{y} is the surjective-injective factorization of \vec{x} , i.e. any element is ~-related to a minimal one. It is easy to see that any two minimal representatives of \approx -equivalence class are ~-related.

We define an isomorphism

$$\kappa_A : \hat{A} \longrightarrow Lan_{\mathbf{i}_{\mathbf{S}}}(A)$$

so that on a set X

$$(\kappa_A)_X : \hat{A}(X) \longrightarrow Lan_{\mathbf{i}_S}(A)(X)$$

we put

$$[\vec{x}, a] \mapsto \|\vec{x}, a\|$$

where $\vec{x}: (n] \to X$ and $a \in A_n$. Taking the minimal elements in the equivalence class is the inverse function. Thus $(\kappa_A)_X$ is a bijection. It is easy to see that κ_A defined this way is natural in X, i.e. it is a natural isomorphism.

For a natural transformation $\tau: A \to B$ in $Set^{\mathbb{S}}$ the square

$$\hat{A} \xrightarrow{\tau_A} Lan_{\mathbf{i}_{\mathbf{s}}}(A)$$

$$\hat{\tau} \downarrow \qquad \qquad \downarrow Lan_{\mathbf{i}_{\mathbf{s}}}(\tau)$$

$$\hat{B} \xrightarrow{\tau_B} Lan_{\mathbf{i}_{\mathbf{s}}}(B)$$

obviously commutes, as both compositions of an element $[\vec{x}, a] \in \hat{A}(X)$ are equal to $\|\vec{x}, \tau(a)\|$. Thus

$$\kappa: (\hat{-}) \longrightarrow Lan_{\mathbf{i}_{\mathrm{S}}}$$

is also a natural isomorphism, as required. \Box

Theorem 2.2. The functor (-): Set^S \longrightarrow End is faithful, full on isomorphisms, and its essential image is the category of semi-analytic functors San.

The theorem will be proved via a series of lemmas.

Lemma 2.3. The essential image of the functor (-): Set[§] \longrightarrow End is contained in the category San.

Proof. First we check that for $A \in Set^{\mathbb{S}}$, \hat{A} preserves pullbacks along monos. Let



be a pullback in Set. We need to show that the square

$$\begin{split} \sum_{n \in \omega} \begin{bmatrix} P \\ n \end{bmatrix} \otimes_n A_n & \xrightarrow{\hat{A}(f')} & \sum_{n \in \omega} \begin{bmatrix} Y \\ n \end{bmatrix} \otimes_n A_n \\ & \hat{A}(\beta') \\ & & & \\ \sum_{n \in \omega} \begin{bmatrix} Z \\ n \end{bmatrix} \otimes_n A_n & \xrightarrow{\hat{A}(f)} & \sum_{n \in \omega} \begin{bmatrix} X \\ n \end{bmatrix} \otimes_n A_n \end{split}$$

is also a pullback in Set. Let $[\vec{y}, a] \in \begin{bmatrix} Y \\ n \end{bmatrix} \otimes_n A_n$ and $[\vec{z}, b] \in \begin{bmatrix} Z \\ m \end{bmatrix} \otimes_m A_m$ be such that $\hat{A}(f)([\vec{z}, b]) = \hat{A}(\beta)([\vec{y}, a])$

Let $\alpha : (m] \to (n']$ and $\vec{z}' : (n] \to X$ be a surjection-injection factorization of the function $f \circ \vec{z} : (m] \to X$. Then there is a $\sigma \in S_n$ such that

$$\vec{z} \circ \sigma = \beta \circ \vec{y}, \quad A(\alpha)(b) = A(\sigma)(a)$$

We have

$$f \circ \vec{z} = \vec{z}' \circ \alpha = \beta \circ \vec{y} \circ \sigma^{-1} \circ \alpha$$

Hence there is a function $\vec{p}: (m] \to P$ as in the following diagram



such that

$$\beta'\circ \vec{p}=\vec{z}, \qquad f'\circ \vec{p}=\vec{y}\circ \sigma^{-1}\circ \alpha$$

The function \vec{p} is an injection since \vec{z} is. Thus $[\vec{p}, b]$ is an element of $\begin{bmatrix} P \\ m \end{bmatrix} \otimes_m A_m$. We have

$$\hat{A}(\beta')([\vec{p},b]) = [\beta' \circ \vec{p},b] = [\vec{z},b]$$

Moreover, on representatives we have

$$\langle f' \circ \vec{p}, b \rangle = \langle \vec{y} \circ \sigma^{-1} \circ \alpha, b \rangle \sim$$
$$\sim \langle \vec{y}, A(\sigma^{-1} \circ \alpha)(b) \rangle = \langle \vec{y}, A(\sigma^{-1})(A(\alpha)(b)) \rangle =$$
$$= \langle \vec{y}, A(\sigma^{-1})(A(\sigma)(a)) \rangle = \langle \vec{y}, a \rangle$$

and hence

$$\hat{A}(f')([\vec{p},b]) = [f' \circ \vec{p},b] = [\vec{y},a]$$

Thus \hat{A} preserves pullbacks along monos indeed.

Now let $\tau : A \to B$ be a morphism in $Set^{\mathbb{S}}$. We shall show that $\hat{\tau} : \hat{A} \to \hat{B}$ is semi-cartesian. Let $f : X \to Y$ be an injection in Set. We need to show that the square

$$\begin{array}{c|c}
\hat{A}(X) & \xrightarrow{\hat{\tau}_X} \hat{B}(X) \\
\hat{A}(f) & & & & \\
\hat{A}(Y) & & & & \\
\hat{\tau}_Y & \hat{B}(X)
\end{array}$$

is a pullback. Let $[\vec{y}, a] \in \hat{A}(Y)$ and $[\vec{x}, b] \in \hat{B}(X)$ be such that

$$[\vec{y}, \tau_n(a)] = \hat{\tau}_Y([\vec{y}, a]) = \hat{B}(f)([\vec{x}, b]) = [f \circ \vec{x}, b]$$

where $a \in A_n$. Note that as f is an injection, so is $f \circ \vec{x}$ and hence $\langle f \circ \vec{x}, b \rangle$ represents an element in $\hat{B}(Y)$. The above equality means that $b \in B_n$ and there is a $\sigma \in S_n$ such that

$$\vec{y} \circ \sigma = f \circ \vec{x}, \quad \tau_n(a) = B(\sigma)(b)$$

We have an element $[\vec{x} \circ \sigma^{-1}, a]$ such that

$$\hat{A}(f)([\vec{x}\circ\sigma^{-1},a]) = [f\circ\vec{x}\circ\sigma^{-1},a] = [\vec{y},a]$$

On representatives we have

$$\langle \vec{x}, \tau_n(a) \rangle \sim \langle \vec{x}, \hat{B}(\sigma^{-1}(\tau_n(a))) \rangle = \langle \vec{x}, b \rangle$$

and hence we have

$$\hat{\tau}_X([\vec{x} \circ \sigma^{-1}, a]) = [\vec{x}, \tau_n(a)] = [\vec{x}, b]$$

as well. This means that $\hat{\tau}$ is semi-cartesian indeed. \square

Lemma 2.4. The functor (-): Set[§] \longrightarrow End is faithful and full on semi-cartesian morphisms. In particular it is full on isomorphisms.

Proof. One can easily verify that if two natural transformations $\hat{\tau}, \hat{\tau}' : \hat{A} \to \hat{B}$ agree on elements of the form $[1_{(m]}, a]$ for $m \in \omega$ and $a \in \hat{A}(m]$ then natural transformation $\tau, \tau' : A \to B$ are equal. Thus (-) is faithful.

To show that (-) is full, let us fix two functors $A, B \in Set^{\mathbb{S}}$ and a semi-cartesian natural transformation $\psi : \hat{A} \to \hat{B}$. Let $m \in \omega$ and $a \in A_m$. We shall define $\tau_m(a)$. We have $\psi_{(m]}([1_{(m]}, a]) = [f, b] \in \hat{B}(m]$ for some injection $f : (k] \to (m]$ and $b \in B_k$. We claim that k = mand f is a bijection. Suppose to the contrary, that k < m. We have that $\hat{B}(f)([1_{(k]}, b]) = [f, b]$. Since f is an injection and ψ is semi-cartesian, the square

is a pullback and hence there is an element $[g,c] \in A(k]$ such that

$$\hat{A}(f)([g,c]) = [1_{(m]},a], \quad \psi_{(m]}([g,c]) = [1_{(k]},b]$$

The first equality implies that the proper mono $f \circ g$ is epi (as its codomain is (m] and its mono part is $1_{(m]}$). This is a contradiction. Hence f is a bijection and we can apply the functor B to it. We put

$$\tau_m(a) = B(f)(b)$$

Thus we have defined a transformation $\tau : A \to B$.

To show that τ is natural, let us fix a surjection $\beta : (m] \to (n]$ and $a \in A_m$. Then using definitions of \hat{A} , \hat{B} and the naturality of ψ we have

$$[1_{(n]}, B(f)(\tau_m(a))] = \hat{B}(f)([1_{(m]}, \tau_m(a)]) =$$
$$= \hat{B}(f) \circ \psi_{(m]}([1_{(m]}, a]) = \psi_{(n]} \circ \hat{A}(f)([1_{(m]}, a]) =$$
$$= \psi_{(n]}([1_{(n]}, \hat{A}(f)(a))]) = [1_{(n]}, \tau_n(\hat{A}(f)(a))]$$

But this means that

$$B(f)(\tau_m(a)) = \tau_n(\hat{A}(f)(a))$$

Since a and f were arbitrary, τ is natural.

Finally, we show that $\hat{\tau} = \psi$. Let us fix a set X and $[\vec{x} : (k] \to X, a \in A_k] \in \hat{A}(X)$. Note that $\hat{A}(\vec{x})([1_{(k]}, a]) = [\vec{x}, a]$. Using the naturality of ψ and $\hat{\tau}$ on \vec{x} and the definition of τ we have

$$\psi_X([\vec{x}, a]) = \psi_X \circ \hat{A}(\vec{x})([1_{(k]}, a]) =$$

$$= \hat{B}(\vec{x}) \circ \psi_{(k]}([1_{(k]}, a]) = \hat{B}(\vec{x})([1_{(k]}, \tau_k(a)]) =$$

$$= \hat{B}(\vec{x}) \circ \hat{\tau}_{(k]}(([1_{(k]}, a]) = \hat{\tau}_X \circ \hat{A}(\vec{x})(([1_{(k]}, a]) =$$

$$= \hat{\tau}_X([\vec{x}, a])$$

Since X and $[\vec{x}, a]$ were arbitrary $\hat{\tau} = \psi$. \Box

Lemma 2.5. Each semi-cartesian functor is in the essential image of $(\hat{-})$: Set^S \longrightarrow End.

Proof. Let us fix a semi-cartesian functor. We define a functor $A : \$ \to Set$ and a natural isomorphism $\tau : \hat{A} \to F$.

Put

$$A(n] = F(n] - \bigcup_{(m] \xrightarrow{f} (n] \in \mathbb{I}, \ m < n} F(f)(A(m])$$

Note that the sum over the empty index set is empty. For $\alpha : [n] \to [m]$ in \mathbb{S} we put

$$A(\alpha) = F(\alpha)_{\lceil A \rceil n} : A[n) \longrightarrow A[m)$$

We need to show that $A(\alpha)$ is well defined, i.e. for $a \in A(n]$ we have $F(\alpha) \in A(m]$. Clearly, $F(\alpha) \in F(m]$. Suppose to the contrary, that $F(\alpha)(a) \notin A(m]$. Then there is a proper mono $f: (k] \to (m]$ and $b \in A(k]$ such that $F(\alpha)(a) = F(f)(b)$. We form a pullback of a surjection α and a proper mono f

$$\begin{array}{c} (p] & \xrightarrow{\alpha'} & (k] \\ f' \downarrow & & \downarrow f \\ (n] & \xrightarrow{\alpha} & (m] \end{array}$$

Thus f' is again a proper mono. F preserves this pullback, i.e. the square

$$F(p) \xrightarrow{F(\alpha')} F(k)$$

$$F(f') \downarrow \qquad \qquad \downarrow F(f)$$

$$F(n) \xrightarrow{F(\alpha)} F(m)$$

is again a pullback. Thus there is an element $c \in F(p]$ such that

$$f(\alpha')(c) = b, \qquad f(f')(c) = a$$

The later equality means that $a \notin A(n]$ contrary to the supposition. Thus $A : \mathbb{S} \to Set$ is a well defined functor.

For a set X we define the function

$$\sum_{n \in \omega} \begin{bmatrix} X \\ n \end{bmatrix} \otimes_n A_n \xrightarrow{\tau_X} F(X)$$
$$\tau_X([\vec{x}, a]) = F(\vec{x})(a)$$

where $n \in \omega$, $\vec{x} : (n] \to X$ is an injection and $a \in A(n]$.

First we show that the transformation $\tau : \hat{A} \to F$ is natural. Let $f : X \to Y$ be a function. Then we have an epi/mono factorization of f', \vec{y} of the function $f \circ \vec{x}$

$$(n) \xrightarrow{\vec{x}} X$$

$$f' \downarrow \qquad \qquad \downarrow f$$

$$(m) \xrightarrow{\vec{y}} Y$$

Thus we have

by

$$F(f)\tau_X([\vec{x},a]) = F(f \circ \vec{x})(a) =$$
$$F(\vec{y} \circ f')(a) = F(\vec{y})(F(f')(a)) =$$
$$= \tau_Y(A(f')(a))$$

i.e. τ is natural.

It remains to show that τ is an isomorphism. Fix a set X. Let $x \in F(x)$. As F is finitary, there is an $n \in \omega$, $f: (n] \to X$ and $y \in F(n]$ such that F(f)(y) = x. Let α , g be an epi/mono factorization of f



Then $[g, A(\alpha)(y)] \in \begin{bmatrix} X \\ m \end{bmatrix} \otimes_m A_m$ and

$$\tau_X([g, A(\alpha)(y)]) = F(g)(A(\alpha)(y)) =$$
$$= F(g \circ \alpha)(y) = F(f)(y) = x$$

Thus τ_X is onto.

Now suppose that
$$[\vec{x}, a] \in \begin{bmatrix} X\\ n \end{bmatrix} \otimes_n A_n$$
 and $[\vec{y}, b] \in \begin{bmatrix} X\\ m \end{bmatrix} \otimes_m A_m$ and $\tau_X([\vec{x}, a]) = \tau_X([\vec{y}, b])$

We form a pullback of monos

$$(n] \xrightarrow{\vec{x}} X$$

$$f \downarrow \qquad \qquad \downarrow \vec{y}$$

$$(k] \xrightarrow{g} (m]$$

that F preserves, i.e. we have a pullback

$$A(n] \longrightarrow F(n] \xrightarrow{F(\vec{x})} F(X)$$

$$F(f) \downarrow \qquad \qquad \downarrow F(g) \xrightarrow{F(g)} F(m]$$

$$A(m]$$

Since

$$F(\vec{x})(a) = \tau_X([\vec{x}, a]) = \tau_X([\vec{y}, b]) = F(\vec{y})(b)$$

there is a $c \in F(k]$ such that

$$F(f)(c) = a, \quad F(g)(c) = b$$

As $a \in A(n]$ and $b \in B(m]$ we must have that n = k = m and both f and g are bijections. Put $\sigma = f \circ g^{-1}$. Then

$$\vec{x} \circ \sigma = \vec{y}, \quad a = A(\sigma)(b)$$

which means that $[\vec{x}, a] = [\vec{y}, b]$. This τ_X is an injection, as required. \Box

A monad (M, η, μ) on *Set* is a *semi-analytic monad* iff M is a semi-analytic functor and both η and μ are semi-cartesian natural transformations. The category of semi-analytic monads with the semi-cartesian morphisms will be denoted by **SanMnd**.

We have inclusion functors

AnMnd
$$\xrightarrow{\mathcal{P}_r^m}$$
 SanMnd $\xrightarrow{\mathcal{P}_f^m}$ Mnd

3 Full operads

In this section we describe the equivalence of the category of full operads **FOp** to categories of equational theories **ET**, Lawvere theories **LT** and monads **Mnd**. As we said, the categories **FOp** and **Mnd** are categories of monoids of two equivalent monoidal categories thus they are obviously equivalent. Therefore we will just define the equivalences of categories



with the domain **FOp** and their essential inverses leaving all the verifications to the reader. In [Tr] it was shown that the category of full operads is equivalent to the category of abstract clones. The latter category in known to be equivalent to **ET**.

Let us fix a morphism $h: \mathcal{O} \to \mathcal{O}'$ of **F**-operads. We denote by $\kappa_i^n: (1] \to (n]$ the function that sends 1 to *i*, where $n \in \omega$ and we denote π_i^n the action of κ_i^n on the unit $\iota \in \mathcal{O}_1$, i.e. $\pi_i^n = \kappa_i^n \cdot \iota$.

The functor \mathcal{E}_o

The equational theory $\mathcal{E}_o(\mathcal{O}) = (L, A)$ has as the set of operation $L = \coprod_{n \in \omega} \mathcal{O}_n$. For $f \in \mathcal{O}_k$, $f_i \in \mathcal{O}_{n_i}, i = 1, \ldots, k, n = \sum_{i=1}^k n_i$, and $\psi: (k] \to (n]$ we have the following axioms in A:

$$f(f_1(x_1, \dots, x_{n_1}), \dots, f_k(x_{n-n_k+1}, \dots, x_n)) = ((f_1, \dots, f_k) \star f)(x_1, \dots, x_n) : \vec{x}^n$$

$$f(x_{\psi(1)},\ldots,x_{\psi(k)})=(\psi\cdot f)(x_1,\ldots,x_n):\vec{x}^r$$

and

$$\iota(x_1) = x_1 : \vec{x}^1$$

The interpretation $\mathcal{E}_o(h)$: $\mathcal{E}_o(\mathcal{O}) \to \mathcal{E}_o(\mathcal{O}')$ sends function symbol $f \in \mathcal{O}_n$ to the term $h(f)(x_1,\ldots,x_n): \vec{x}^n$. This ends the definition of the functor \mathcal{E}_o .

The essential inverse of the functor \mathcal{E}_o

$$\mathcal{O}p_e: \mathbf{ET} \longrightarrow \mathbf{FOp}$$

is defined as follows. For an equational theory T = (L, A), the F-operad $\mathcal{Q} = \mathcal{O}p_e(T)$ has as *n*-ary operations \mathcal{Q}_n the terms in context \vec{x}^n modulo provability in T. The action

$$\cdot : \mathbb{F}(n,m) \times \mathcal{Q}_n \longrightarrow \mathcal{Q}_m$$

is defined as

$$\phi \cdot [t(x_1, \dots, x_n) : \vec{x}^n] = [t(x_{\phi(1)}, \dots, x_{\phi(n)}) : \vec{x}^m]$$

where $[t(x_1, \ldots, x_n) : \vec{x}^n]$ is an operation in a \mathcal{Q}_n and $\phi : (n] \to (m]$ is a function. The composition in \mathcal{Q}

$$\star:\mathcal{Q}_{n_1}\times\ldots\times\mathcal{Q}_{n_k}\times\mathcal{Q}_k\longrightarrow Q_n$$

where $n = \sum_{i=1}^{k} n_i$ is defined by substitution with α -conversion, i.e. for $[t_i(x_1, \ldots, x_{n_i}) : \vec{x}^{n_i}] \in \mathcal{Q}_{n_i}$ and $[s(x_1, \ldots, x_k) : \vec{x}^k] \in \mathcal{Q}_k$ we have

$$([t_1(x_1,\ldots,x_{n_1}):\vec{x}^{n_1}],\ldots,[t_1(x_1,\ldots,x_{n_k}):\vec{x}^{n_k}])\star[s(x_1,\ldots,x_k):\vec{x}^k] = [s(t_1(x_{\alpha_1(1)}),\ldots,x_{\alpha_1(n_1)})),\ldots,t_k(x_{\alpha_k(1)}),\ldots,x_{\alpha_k(n_k)})):\vec{x}^k]$$

where $\alpha_i : (n_i] \to (n]$, for i = 1, ..., k is the obvious embedding.

The functor \mathcal{L}_o

A morphism

$$\langle f_1,\ldots,f_k\rangle:n\longrightarrow k$$

in the Lawvere theory $\mathcal{L}_o(\mathcal{O})$ is a k-tuple such that $f_i \in \mathcal{O}$, for $i = 1, \ldots, k$. The identity on n is

$$\langle \pi_1^n, \dots, \pi_n^n \rangle : n \longrightarrow n$$

and the *i*-th projection from n is

$$\pi_i^n: n \longrightarrow 1$$

Recall that π_i^n is the value of the action of the function $\kappa_i^n : (1] \to (n]$ that picks *i* on the unit ι . The composition of morphisms

$$\langle g_1, \dots, g_n \rangle : m \longrightarrow n, \quad \langle f_1, \dots, f_k \rangle : n \longrightarrow k$$

is

$$\langle t_1,\ldots,t_k\rangle:m\longrightarrow k$$

where

$$t_i = \psi \cdot ((g_1, \dots g_n) * f_i)$$

for i = 1, ..., k, where $\psi : (n \cdot m] \to (m]$ is given by

$$\psi(r) = ((r-1) \operatorname{mod} m) + 1$$

for $r \in (n \cdot m]$. The interpretation $\mathcal{L}_o(h) : \mathcal{L}_o(\mathcal{O}) \to \mathcal{L}_o(\mathcal{O}')$ sends morphism $\langle f_1, \ldots, f_k \rangle : n \longrightarrow k$ to morphism $\langle h(f_1), \ldots, h(f_k) \rangle : n \longrightarrow k$. This ends the definition of the functor \mathcal{L}_o .

The essential inverse of the functor \mathcal{L}_o

$$\mathcal{O}p_l: \mathbf{LT} \longrightarrow \mathbf{FOp}$$

is defined as follows. For a Lawvere theory **T** the set of *n*-ary operation of the operad $Q = \mathcal{O}p_l(\mathbf{T})$ is $Q_n = \mathbf{T}(n, 1)$. The action in \mathcal{Q} is given by

$$\phi \cdot f = f \circ \pi_{\phi}$$

where $f \in \mathcal{Q}_n$, ϕ as above, and $\pi : \mathbb{F}^{op} \to \mathbf{T}$ is the unique morphism from the initial Lawvere theory. The composition in \mathcal{Q} is given by products and composition in \mathbf{T}

$$(f_1,\ldots,f_k)\star g = g\circ(f\times\ldots,\times f_k)$$

The functor \mathcal{M}_o

Let $f: X \to Y$ be a function. The functor part of the monad $\mathcal{M}_o(\mathcal{O})$ is defined via the coend formula

$$\mathcal{M}_o(\mathcal{O})(X) = \int^{n \in \mathbb{F}} X^n \times \mathcal{O}_r$$

Thus an element of this set is an equivalence class of pairs $\langle \vec{x}, t \rangle \in X^n \times O_n$. We put

$$\mathcal{M}_o(\mathcal{O})(f)([\vec{x},t]) = [f \circ \vec{x},t]$$

The unit of the monad

$$\eta_X^{\mathcal{O}}: X \to \mathcal{M}(\mathcal{O})(X)$$

is given by

 $x \mapsto [\tilde{x}, \iota]$

where $\tilde{x}: (1] \to X$ sends 1 to x. The multiplication

$$\mu_X^{\mathcal{O}}: \mathcal{M}_o^2(\mathcal{O})(X) = \int^{k, n_1, \dots, n_k \in \omega} X^n \times \mathcal{O}_{n_1} \times \dots \times \mathcal{O}_{n_k} \times \mathcal{O}_k \longrightarrow \mathcal{M}_o(\mathcal{O})(X)$$

where $n = \sum_{i=1}^{k} n_i$, is given by

$$[\vec{x}, s_1, \dots, s_k, t] \mapsto [\vec{x}, (s_1, \dots, s_k) * t]$$

The essential inverse of the functor \mathcal{M}_o

$$\mathcal{O}p_m:\mathbf{Mnd}\longrightarrow\mathbf{FOp}$$

is defined as follows. For a monad $T = (T, \eta, \mu)$ in **Mnd** the set of *n*-ary operations in the operad $\mathcal{Q} = Op_m(T)$ is $Q_n = T(n]$. The action on $t \in Q_n$ is

$$\phi \cdot t = T(\phi)(t)$$

Let $t_i \in T(n_i]$ and $s \in T(k]$. then we have corresponding morphisms

 $\lceil s \rceil : (1] \longrightarrow T(k], \quad \lceil t \rceil : (k] \longrightarrow T(n_1] + \dots t(n_k]$

'picking' s and the k terms t_1, \ldots, t_k . The action

 $(t_1,\ldots,t_k)\star s$

is the element 'picked' by the morphism

$$(1) \xrightarrow{\lceil s \rceil} T(k) \xrightarrow{T(\lceil t \rceil)} T(T(n_1] + \ldots + T(n_k]) \xrightarrow{T(\lceil T(\alpha_1), \ldots, T(\alpha_k)])} T^2(n] \xrightarrow{\mu} T(n]$$

in T(n], where functions α_i are as above.

4 Regular Lawvere theories vs regular operads

In this section we study the relations between Lawvere theories and regular operads. We shall describe the adjunction $\mathcal{Q}_r^{ol} \dashv \mathcal{P}_f^{lo}$ and the properties of the embedding \mathcal{P}_f^{lo} .



The functor $\mathcal{P}_f = \mathcal{P}_f^{ol}: \mathbf{RegOp}
ightarrow \mathbf{LT}$

Let \mathcal{O} be a regular operad; ι , \cdot , * denote the unit, action (of \$), and composition in \mathcal{O} , respectively. We define a Lawvere theory $\mathcal{P}_f(\mathcal{O})$ as follows. The set of objects of $\mathcal{P}_f(\mathcal{O})$ is the set of natural numbers ω . A morphism from n to m is an equivalence class of spans



such that $\phi: (r] \to (n]$ is a function called *amalgamation*, $f: (r] \to (m]$ is a monotone function called the *arity function* (as it determines the arities of the operations g_i), $r_i = |f^{-1}(i)|$ and we have $g_i \in \mathcal{O}_{r_i}$ for $i \in (m]$ and $r = \sum_{i=1}^m r_i$, moreover ϕ and f are jointly mono (or what comes out the same, ϕ is mono on the fibers of f). Two spans $\langle \phi, f, g_i \rangle_{i \in m}$ and $\langle \phi', f', g'_j \rangle_{j \in m'}$ are equivalent whenever f = f' and there are permutations $\sigma_i: (r_i] \to (r_i]$ for $i \in (m]$



such that

$$g_i = \sigma_i \cdot g'_i, \quad \phi \circ \sum_i \sigma_i = \phi'$$

In particular $|f^{-1}(i)| = r_i = |f'^{-1}(i)|$, for $i \in (m]$. By $\sum_i \sigma_i : r \to r$ we mean the permutation that is formed by placing permutations σ_i 'one after another'. Thus, it respects the fibers of f, i.e. $f \circ \sum_i \sigma_i = f'$. A morphism in the category $\mathcal{P}_f(\mathcal{O})$ is a class of spans modulo the above equivalence relation.

We could represent morphisms in $\mathcal{P}_f(\mathcal{O})$ as spans without the requirement that ϕ and f are jointly mono. But then the relation identifying the spans would be more complicated. Instead of permutations σ_i we would need to consider surjections. But this relation is not an equivalence relation and we would need to work with the equivalence relation generated by such a relation. However, it might happen, as with the compositions defined below, that an operation on spans naturally leads to a span whose amalgamation ϕ is not injective on fibers of f. In such a case we can *regularize* the span as follows. Let $\langle \phi, f, g_i \rangle_{i \in m}$ be a span as above but with ϕ not necessarily injective on the fibers of f. Let ϕ_i be the restriction of ϕ to the fiber $f^{-1}(i)$ for $i \in (m]$. Let $\phi_i = \phi'_i \circ \psi_i$ be an epi-mono factorization of ϕ_i and $g'_i = \psi_i \cdot g_i$, for $i \in (m]$. Then the *regularization* of the span $\langle \phi, f, g_i \rangle_{i \in m}$ is the span



where $r' = \sum_{i \in (m]} r'_i, r'_i = |dom(\phi'_i)|$, for $i \in (m], \phi' = (\phi'_1 + \ldots + \phi'_m) : r' \to n$. f' is the monotone map sending the elements in the domain of ϕ'_i to i, for $i \in (m]$.

The composition of morphisms $\langle \phi, f, g_i \rangle_{i \in (m]} : n \to m$ and $\langle \phi', f', g'_j \rangle_{j \in (k]} : m \to k$ to $\langle \phi'', f'', g''_j \rangle_{i \in (k]} : n \to k$ is defined in two steps as follows. In the diagram



the square is a pullback of f along ϕ' . The function \overline{f} is so chosen that it is monotone. We define a span by $f'' = f' \circ \overline{f}$, $\phi'' = \phi' \circ \overline{\phi}$, and $g''_j = g'_j * \langle g_{\phi'(l)} \rangle_{l \in f'^{-1}(j)}$. Finally, we take a regularization of this span to get a regular span that represents the composition. We leave for the reader the verification that the composition is a congruence with respect to the equivalence relation on spans.

The identity on n is the span



The *i*-projection $\pi_i^n : n \to 1$ on *i*-th coordinate is the span



where $i \in (n]$ and $\tilde{i}(1) = i$.

For a morphism of regular operads $h: \mathcal{O} \to \mathcal{O}'$, we define a functor

$$\mathcal{P}_f(h): \mathcal{P}_f(\mathcal{O}) \longrightarrow \mathcal{P}_f(\mathcal{O}')$$

so that for a morphism $\langle \phi, f, g_i \rangle_{i \in (m]} : n \to m$ in $\mathcal{P}_f(\mathcal{O})$, we have

$$\mathcal{P}_f(h)(\langle \phi, f, g_i \rangle_{i \in \{m\}}) = \langle \phi, f, h(g_i) \rangle_{i \in \{m\}} : n \to m$$

in $\mathcal{P}_f(\mathcal{O}')$. This ends the definition of the functor \mathcal{P}_f .

The functor $\mathcal{Q}_r = \mathcal{Q}_r^{lo} : \mathbf{LT} \longrightarrow \mathbf{RegOp}$

Let **T** be a Lawvere theory. Recall that $\pi : \mathbb{F}^{op} \to \mathbf{T}$ is the morphism from the initial Lawvere theory. The operad $\mathcal{Q}_r(\mathbf{T})$ consists of operations of **T**, i.e. morphisms to 1. In detail it can be described as follows. The set of *n*-ary operations $\mathcal{Q}_r(\mathbf{T})_n$ is the set of *n*-ary operations $\mathbf{T}(n, 1)$ of **T**, for $n \in \omega$. The action

$$\cdot : \mathbb{S}(n,m) \times \mathcal{Q}_r(\mathbf{T})_n \longrightarrow \mathcal{Q}_r(\mathbf{T})_m$$

is given, for $f \in \mathbf{T}(n, 1)$ and $\phi \in \mathbb{S}(n, m)$, by

$$\phi \cdot f = f \circ \pi_{\phi}$$

The identity of $\mathcal{Q}_r(\mathbf{T})$ is $\iota = 1_1 \in \mathbf{T}(1,1)$. The composition

$$*: \mathcal{Q}_r(\mathbf{T})_{n_1} \times \mathcal{Q}_r(\mathbf{T})_{n_k} \times \mathcal{Q}_r(\mathbf{T})_k \longrightarrow \mathcal{Q}_r(\mathbf{T})_n$$

is given, for $f \in \mathcal{Q}_r(\mathbf{T})_k$ and $f_i \in \mathcal{Q}_r(\mathbf{T})_{n_i}$, where $i \in (k]$, $n = \sum_{i \in k} n_i$, by

$$\langle f_1, \ldots, f_k \rangle * f = f \circ (f_1 \times \ldots, \times f_k)$$

where $f_1 \times \ldots, \times f_k$ is defined using the chosen projections in **T** and \circ is the composition in **T**. If $F : \mathbf{T} \to \mathbf{T}'$ is a morphism of Lawvere theories then the map of regular operads

$$\mathcal{Q}_r(F): \mathcal{Q}_r(\mathbf{T}) \to \mathcal{Q}_r(\mathbf{T}')$$

is defined, for $f \in \mathcal{Q}_r(\mathbf{T})_n$, by

$$\mathcal{Q}_r(F)(f) = F(f)$$

This ends the definition of the functor Q_r .

The adjunction $\mathcal{P}_f \dashv \mathcal{Q}_r$ and the properties of the functor \mathcal{P}_f

We note for the record

Proposition 4.1. The functors $\mathcal{P}_f : \operatorname{RegOp} \longrightarrow \operatorname{LT}$ and $\mathcal{Q}_r : \operatorname{LT} \to \operatorname{RegOp}$ are well defined. \Box

We have an easy

Lemma 4.2. Let \mathcal{O} be a regular operad and $n \in \omega$. An isomorphism on n in $P_r(\mathcal{O})$ has a representation by a span of the following form



where $\phi : (n] \to (n]$ is a bijection, $a_i \in \mathcal{O}_1$ is an invertible operation, i.e. there is $b_i \in \mathcal{O}_1$ such that $a_i * b_i = \iota = b_i * a_i$ for $i \in (n]$. It is the unique span in its equivalence class.

Proof. Suppose that we have two spans



representing two morphisms that compose to identity 1_n both ways. Then, since the displayed composition is 1_n , the functions ϕ , f' are surjections. Hence the functions ϕ' , f are surjections, as well. Having this it is easy to notice that ϕ sends elements in different fibers of f to different elements. Thus ϕ (and ϕ') must be in fact a bijection.

If we take a composition $\langle g_i \rangle_{i \in f'^{-1}(j)} * h_j$ and regularize it multiplying by a surjection, we will get ι . This implies that h_j cannot be a nullary operation and that the arity of g_i 's is at most one. Now, as the above spans represent morphisms that compose both ways to identity, it is easy to see that we get the required description. \Box

Proposition 4.3. We have an adjunction $\mathcal{P}_f \dashv \mathcal{Q}_r$. The functor \mathcal{P}_f is faithful.

Proof. We shall show that $\mathcal{P}_f \dashv \mathcal{Q}_r$. For a regular operad \mathcal{O} the unit is

$$\eta_{\mathcal{O}}: \mathcal{O} \longrightarrow \mathcal{Q}_r(\mathcal{P}_f(\mathcal{O}))$$
$$\mathcal{O}_n \ni g \mapsto \langle 1_n, !, g \rangle$$

For Lawvere theory \mathbf{T} the counit is

$$\varepsilon_{\mathbf{T}}: \mathcal{P}_f \mathcal{Q}_r(\mathbf{T}) \longrightarrow \mathbf{T}$$

$$\langle \phi, f, g_i \rangle_{i \in (m]} \mapsto (g_1 \times \ldots \times g_m) \circ \pi_{\phi}$$

We verify the triangular equalities. For $g \in \mathcal{Q}_r(\mathbf{T})_n = \mathbf{T}(n, 1)$ we have

$$\begin{aligned} \mathcal{Q}_r(\varepsilon_{\mathbf{T}}) \circ \eta_{\mathcal{Q}_r(\mathbf{T})}(g) &= \\ &= \mathcal{Q}_r(\varepsilon_{\mathbf{T}})(\langle \mathbf{1}_n, \mathbf{!}, g \rangle) = \\ &= g \circ \pi_{\mathbf{1}_n} = g \end{aligned}$$

For $\langle \phi, f, g_i \rangle_{i \in (m]} \in \mathcal{P}_f(\mathcal{O})$ we have

$$\varepsilon_{\mathcal{P}_{f}(\mathcal{O})} \circ \mathcal{P}_{f}(\eta_{\mathcal{O}})(\langle \phi, f, g_{i} \rangle_{i \in \{m\}}) =$$

$$= \varepsilon_{\mathcal{P}_{f}(\mathcal{O})}(\langle \phi, f, \langle 1_{r_{i}}, !, g_{i} \rangle_{i \in \{m\}}) =$$

$$= (\langle 1_{r_{1}}, !, g_{1} \rangle \times \ldots \times \langle 1_{r_{m}}, !, g_{m} \rangle) \circ \pi_{\phi} =$$

$$= \langle \phi, f, g_{i} \rangle_{i \in \{m\}}$$

As the unit $\eta_{\mathcal{O}}$ is mono, \mathcal{P}_f is faithful. \Box

Corollary 4.4. The triangle



commutes up to an isomorphism.

Proof. By Proposition 4.3 it is enough to show that triangle



commutes up to an isomorphism, where the functor $\mathcal{O}p_l^f$ is the left adjoint to \mathcal{L}_o^f and together they form an adjoint equivalence. The functor $\mathcal{O}p_l^f$ is defined as the functor \mathcal{Q}_r except the action involved is the action of the whole category \mathbb{F} rather than its subcategory \mathbb{S} . As the functor \mathcal{Q}_r^o forgets this additional part of the action the above diagram clearly commutes. \Box

Proposition 4.5. The functor $\mathcal{P}_f : \operatorname{RegOp} \to \operatorname{LT}$ is full on isomorphisms and its essential image is RegLT. In particular RegOp is equivalent to RegLT.

Proof. Recall that we have a unique morphism of Lawvere theories from the initial theory $\pi : \mathbb{F}^{op} \to \mathcal{P}_f(\mathcal{O})$. For a function $\phi : m \to n$, π_{ϕ} the morphism π_{ϕ} is represented by the span of form



The class of projection morphisms, is the closure under isomorphism of the class of morphisms $\{\pi_{\phi} : \phi \in \mathbb{I}\}$ in $\mathcal{P}_f(\mathcal{O})$. Using Lemma 4.2, it is easy to see that the projection morphisms in $\mathcal{P}_f(\mathcal{O})$ are (represented by) the spans of the form



where ϕ is an injection and a_i is an invertible unary operation.

The regular morphisms in $\mathcal{P}_f(\mathcal{O})$ are (represented by) the spans of the form



where ϕ is a bijection.

Clearly, both classes contain isomorphisms and are closed under composition. Any morphism $\langle \phi, f, g_i \rangle_{i \in \{m\}} : n \to m$ in $\mathcal{P}_f(\mathcal{O})$ has a projection-regular factorization as follows



Thus to show that projections and (what we have described as) regular morphisms form a factorization system it remains to show that projection morphisms are left orthogonal to the regular morphisms. Let



be a commutative square in $\mathcal{P}_f(\mathcal{O})$ with the left vertical morphism $\langle \phi, 1_r, a_i \rangle_{j \in [k]}$ a structural map and right vertical morphism $\langle 1_m, !, g \rangle$ a regular map. We have chosen the right bottom to be 1 to simplify notation but the general case is similar. The commutativity means that r = r' and there is a permutation $\sigma \in S_r$ such that

$$\psi = \phi \circ \phi' \circ \sigma$$

and

$$\langle a_{\phi'(1)}, \ldots, a_{\phi'(r)} \rangle * g' = \sigma \cdot (\langle h_1, \ldots, h_m \rangle * g)$$

Putting into the square a diagonal morphism $\langle \phi' \circ \sigma, f, h_i \rangle_{i \in [m]}$



where

$$\bar{h_i} = \langle a_{\phi' \circ \sigma(l)}^{-1} \rangle_{l \in f^{-1}(i)} * h_i$$

we see that the permutations 1_r and σ show that both triangles commute. Thus regular morphisms are indeed right orthogonal to the structural ones and $\mathcal{P}_f(\mathcal{O})$ is a regular Lawvere theory.

From the description of the functor $\mathcal{P}_f(h) : \mathcal{P}_f(\mathcal{O}) \to \mathcal{P}_f(\mathcal{O}')$ and the description of the structure of $\mathcal{P}_f(\mathcal{O})$, it is clear that $\mathcal{P}_f(h)$ sends the regular (projection) morphisms to the regular (projection) morphisms. Thus $\mathcal{P}_f(h)$ is a regular morphism of Lawvere theories.

Now let **T** be any Lawvere theory. As the class of regular morphisms in **T** is right orthogonal to a class of morphisms and it is closed under composition, finite products and isomorphisms. Moreover, for any surjection in $\psi : (n] \to (m]$ in **F** the image $\pi_{\psi} : m \to n$ in **T** is regular as it is orthogonal to all projection morphisms in **T**. Thus surjections in **S** act on all regular morphisms $f: n \to 1$ in **T** on the left

$$\cdot : \mathbb{S}(n,m) \times \mathbf{T}(n,1) \longrightarrow \mathbf{T}(m,1)$$

by

$$(\psi, f) \mapsto f \circ \pi_{\psi}$$

Hence the regular operations of any Lawvere theory \mathbf{T} form a regular operad. The unit is the identity morphism on 1. The composition $\langle f_1, \ldots, f_n \rangle * f$ is defined to be $f \circ (f_1 \times \ldots \times f_n)$. The action of S is defined as above. Let us denote this operad by \mathbf{T}^r . We have an inclusion morphism of regular operads $\mathbf{T}^r \to \mathcal{Q}_r(\mathbf{T})$. By adjunction we get a morphism of Lawvere theories

$$\xi_{\mathbf{T}}: \mathcal{P}_f(\mathbf{T}^r) \longrightarrow \mathbf{T}$$

Clearly, $\xi_{\mathbf{T}}$ is bijective on objects. If \mathbf{T} is regular then $\xi_{\mathbf{T}}$ is full (faithful) since the projection-regular factorization exists (is unique and $\pi : \mathbb{F}^{op} \to \mathbf{T}$ is faithful).

If $I: \mathbf{T} \to \mathbf{T}'$ is a regular interpretation between any Lawvere theories, then the diagram



commutes, where I^r is the obvious restriction of I to \mathbf{T}^r . Thus the essential image of \mathcal{P}_f is indeed the category of regular Lawvere theories and regular interpretations. An isomorphic interpretation of Lawvere theories is always regular. Therefore \mathcal{P}_f is full on isomorphisms. \Box

We have

Proposition 4.6. The functor $Q_r : LT \to RegOp$ is monadic.

Proof. We shall verify the assumption of the Beck monadicity theorem. By Proposition 4.3 Q_r has a left adjoint. It is easy to see that Q_r reflects isomorphisms. We shall verify that **LT** has and Q_r preserves Q_r -contractible coequalizers.

Let $I, I': \mathbf{T}' \to \mathbf{T}$ be a pair of interpretations between Lawvere theories so that

$$\mathcal{Q}_r(\mathbf{T}') \xrightarrow[]{\mathcal{Q}_r(I')}{\underbrace{\mathcal{Q}_r(I')}{r}} \mathcal{Q}_r(\mathbf{T}) \xrightarrow[]{q} \mathcal{O}$$

is a split coequalizer in **RegOp**. We define a Lawvere theory $\mathbf{T}_{\mathcal{O}}$ so that a morphism from n to m in $\mathbf{T}_{\mathcal{O}}$ is an m-tuple $\langle g_1, \ldots, g_m \rangle$ with $g_i \in \mathcal{O}_n$, for $i = 1, \ldots, m$. The compositions and the identities in $\mathbf{T}_{\mathcal{O}}$ are defined in the obvious way from the compositions and the unit in \mathcal{O} . The projections $\bar{\pi}_i^n$ in $\mathbf{T}_{\mathcal{O}}$ are the images of the projections π_i^n in \mathbf{T} , i.e. $\bar{\pi}_i^n = q(\pi_i^n)$.

The functor $\tilde{q}: \mathbf{T} \to \mathbf{T}_{\mathcal{O}}$ is defined, for $f: n \to m$ in \mathbf{T} , as

$$\tilde{q}(f) = \langle q(\pi_1^m \circ f), \dots, q(\pi_m^m \circ f) \rangle$$

First we verify, that $\mathbf{T}_{\mathcal{O}}$ has finite products. For this, it is enough to verify that $\langle f_1, \ldots, f_n \rangle * \bar{\pi}_i^n = f_i$, where * is the multiplication in the operad \mathcal{O} . The uniqueness of the morphism into the product is obvious from the construction. We have routine calculations

$$\langle f_1, \dots, f_n \rangle * \bar{\pi}_i^n =$$

$$q \circ s(\langle f_1, \dots, f_n \rangle * q(\pi_i^n)) =$$

$$\langle q \circ s(f_1), \dots, q \circ s(f_n) \rangle * (q \circ s \circ q(\pi_i^n)) =$$

$$\langle q \circ s(f_1), \dots, q \circ s(f_n) \rangle * (q \circ Op(I) \circ t(\pi_i^n)) =$$

$$\langle q \circ s(f_1), \dots, q \circ s(f_n) \rangle * (q \circ Op(I') \circ t(\pi_i^n)) =$$

$$\langle q \circ s(f_1), \dots, q \circ s(f_n) \rangle * (q(\pi_i^n)) =$$

$$q(\langle s(f_1), \dots, s(f_n) \rangle * \pi_i^n) =$$

$$q(s(f_i) = f_i$$

It is obvious that \tilde{q} is a morphism of Lawvere theories and that $Q_r(\tilde{q}) = q$. It remains to verify that \tilde{q} is a coequalizer in **LT**. Let $p: \mathbf{T} \to \mathbf{S}$ be a morphism in **LT** coequalizing I and I'.



The morphism $\mathcal{Q}_r(p)$ coequalizes $\mathcal{Q}_r(I)$ and $\mathcal{Q}_r(I')$ in **RegOp**. Thus there is a unique morphism k in **RegOp** making the triangle on the right

$$\mathcal{Q}_{r}(\mathbf{T}') \xrightarrow{\mathcal{Q}_{r}(I)} \mathcal{Q}_{r}(\mathbf{T}') \xrightarrow{q} \mathcal{O}$$

$$\mathcal{Q}_{r}(p) \xrightarrow{k} \mathcal{Q}_{r}(\mathbf{S})$$

commute. We define the functor \tilde{k} so that

$$\tilde{k}(\langle f_1,\ldots,f_n\rangle) = \langle k(f_1),\ldots,k(f_n)\rangle$$

for any morphism $\langle f_1, \ldots, f_n \rangle$ in $\mathbf{T}_{\mathcal{O}}$. The verification that \tilde{k} is the required unique functor is left for the reader. \Box

5 Semi-analytic monads vs regular operads

The main objective of this section is to show that the square

$$\begin{array}{c} \mathbf{FOp} & \xrightarrow{\mathcal{M}_{o}^{f}} & \mathbf{Mnd} \\ \mathcal{P}_{f}^{o} & & \uparrow \mathcal{P}_{f}^{m} \\ \mathbf{RegOp} & \xrightarrow{\mathcal{M}_{o}^{r}} & \mathbf{SanMnd} \end{array}$$
(3)

commutes up to isomorphism, with horizontal functors being equivalences of categories. The functor \mathcal{M}_{o}^{f} was defined in Section 3. \mathcal{P}_{f}^{m} is an inclusion. To this end we shall use Theorem 2.2 and some general considerations but the functor part

To this end we shall use Theorem 2.2 and some general considerations but the functor part of the monad $\mathcal{M}_o^r(\mathcal{O})$ will be described explicitly below. For a regular operad \mathcal{O} , the monad $\mathcal{M}_o^r(\mathcal{O})$ on a set X is defined as follows. The functor $\mathcal{M}_o^r(\mathcal{O})$ is the application of the functor $(-): Set^{\mathbb{S}} \longrightarrow End$ to \mathcal{O} considered as a functor $\mathcal{O}: \mathbb{S} \to Set$, see Section 2 for details.

In particular for a set X we have

$$\mathcal{M}_{o}^{r}(\mathcal{O})(X) = \sum_{n \in \omega} \begin{bmatrix} X \\ n \end{bmatrix} \otimes_{n} \mathcal{O}_{n}$$

In the set $\begin{bmatrix} X \\ n \end{bmatrix} \otimes_n \mathcal{O}_n$ we identify $\langle \vec{x} \circ \sigma, a \rangle$ with $\langle \vec{x}, \sigma \cdot a \rangle$ for $a \in \mathcal{O}_n, \vec{x} : (n] \to X$ and $\sigma \in S_n$.

Let $\gamma : \mathbb{S} \to \mathbb{F}$ be the inclusion functor. It induces the following diagram of categories and functors that we describe below

 γ^* is the functor of composing with γ . It has a left adjoint Lan_{γ} , the left Kan extension along γ . For $C \in Set^{\mathbb{S}}$ it is given by the coend formula

$$Lan_{\gamma}(C)(X) = \int^{n \in \mathbb{S}} X^n \times C(n)$$

The functor $i^{sa} : \mathbf{San} \to \mathbf{End}$ is just an inclusion. The equivalence

$$i_{\mathbb{S}}: Set^{\mathbb{S}} \longrightarrow \mathbf{San}$$

is defined by left Kan extension that may be given by the coend formula and a coproduct

$$i_{\mathbb{S}}(C)(X) = \int^{n \in \mathbb{S}} X^n \times C(n) = \sum_{n \in \omega} \begin{bmatrix} X \\ n \end{bmatrix} \otimes_n C(n)$$

where $C \in Set^{\$}$. Similarly, the equivalence

$$i_{\mathbb{F}}: Set^{\mathbb{F}} \longrightarrow \mathbf{End}$$

is defined by left Kan extension that is given by the coend formula

$$i_{\mathbb{F}}(B)(X) = \int^{n \in \mathbb{F}} X^n \times C(n)$$

where $B \in Set^{\mathbb{F}}$. The following calculation shows that the right hand square in the above diagram commutes:

$$i_{\mathbb{F}}(Lan_{\gamma}(C)(X) \cong \int^{m \in \mathbb{F}} X^m \times Lan_{\gamma}(C)(m] \cong$$
$$\cong \int^{m \in \mathbb{F}} X^m \times \sum_{n \in \omega} (m]^n \otimes_n C(n] \cong$$
$$\cong \sum_{n \in \omega} X^n \otimes_n C(n]$$

The functor $(-)^{sa}$, right adjoint to i^{sa} , is given by the formula

$$F^{sa}(X) = \sum_{n \in \omega} \begin{bmatrix} X \\ n \end{bmatrix} \otimes_n F(n)$$

for $F \in \mathbf{End}$. $(-)^{sa}$ associates to functors and natural transformations their 'semi-analytic parts'.

Note that both **San** and **End** are strict monoidal categories with tensor given by composition, and i^{sa} is a strict monoidal functor. Thus its right adjoint $(-)^{sa}$ has a unique lax monoidal structure making the adjunction $i^{sa} \dashv (-)^{sa}$ a monoidal adjunction. This in turn gives us a monoidal monad (\mathbb{W}, η, μ) on **San**.

We have (cf. [Z]) a 2-natural transformation \mathcal{U}

$$\operatorname{MonCat} \xrightarrow[|-|]{\operatorname{Mon}} \operatorname{Cat}$$

where **MonCat** is the 2-category of monoidal categories, lax monoidal functors, and monoidal transformations; **Mon** is the 2-functor associating the monoids objects to monoidal categories, |-| is the forgetful functor forgetting the monoidal structure, and \mathcal{U} is a 2-natural transformation whose component at a monoidal category M is the forgetful functor from monoids in M to the underlying category of $M: \mathcal{U}_M: \mathbf{Mon}(M) \to |M|$.

Applying \mathcal{U} to the monoidal adjunction and $i^{sa} \dashv (-)^{sa}$ and monoidal monad \mathbb{W} we get an adjunction between categories of monoids and a monad on **Mon(San**). The unnamed arrow in the above diagram is **Mon** (i^{sa}) . But the monoids in **End** and **San** are exactly monads and hence we get the left most adjunction $\mathcal{P}_f^m \dashv \mathcal{Q}_r^m$ together with the monad $(\overline{\mathbb{W}}, \overline{\eta}, \overline{\mu})$ on the category of semi-analytic monads.

On the other hand, on the categories $Set^{\mathbb{F}}$ and $Set^{\mathbb{S}}$ there are substitution tensors making $i_{\mathbb{F}}$ and $i_{\mathbb{S}}$ monoidal equivalences and γ^* and Lan_{γ} monoidal adjunctions. Thus we can apply the 2-functor **Mon** to this adjunction and obtain an adjunction $\mathbf{Mon}(Lan_{\gamma}) \dashv \mathbf{Mon}(\gamma^*)$ as in the diagram

The unnamed functor is $Mon((-)^{sa})$. But monoids in $Set^{\mathbb{F}}$ and $Set^{\mathbb{S}}$ are (equivalent to) the categories of full and regular operads, respectively. The verification that the right most square commutes serially is left for the reader. We obtain

Proposition 5.1. The square (3) of categories and functors commutes up to an isomorphism.

Proof. Both horizontal adjunctions in the square (3) are obtained from equivalent monoidal adjunctions. It remains to show that the identifications we obtained above are isomorphic to the functors \mathcal{M}^f_o and \mathcal{M}^r_o , respectively. This is left for the reader. \Box

There are free monads on finitary functors (cf. [Barr]) and free semi-analytic monads on semianalytic functors. The adjunctions $F \dashv U$ and $\widehat{F} \dashv \widehat{U}$ induce monads \mathbb{R} and $\widehat{\mathbb{R}}$, respectively. $\widehat{\mathbb{R}}$ is the finitary version of what is called 'the monad for all monads' in [Barr]. Putting this additional data to the above diagram and simplifying it at the same time we get a diagram



In the above diagram the square of the left adjoints commutes. Thus, the square of the right adjoint commutes as well. This shows in particular that the free monad on a semi-analytic functor is semi-analytic.

The monad \overline{W} is a lift of a monad W to the category of R-algebras **SanMnd** and, by [Be] we obtain

Theorem 5.2. The monad \mathbb{R} for regular monads distributes over the monad \mathbb{W} for finitary functors, *i.e.* we have a distributive law

$$\lambda: \mathbb{R}\mathbb{W} \longrightarrow \mathbb{W}\mathbb{R}$$

The category of algebras of the composed monad $W\mathbb{R}$ on **SanMnd** is equivalent to the category **Mnd** of all finitary monads on Set. \Box

Remark. We arrived at the above theorem with essentially no calculations. We give below explicit formulas how to calculate the values of some functors mentioned above and we shall also describe the coherence morphism on the monoidal monad W. The coherence morphism φ : $W \circ W \longrightarrow W(- \circ -)$ is a 'finite' description of the distributive law λ .

First we describe the adjunction $i^{sa} \dashv (-)^{sa}$. We shall drop the inclusion i^{sa} when possible. Let $A \in \mathbf{San}$ and $G \in \mathbf{End}$ and X be a set. The regular functor A is given by the (functor of) its coefficients. Its value at X is

$$A(X) = \sum_{n \in \omega} \begin{bmatrix} X \\ n \end{bmatrix} \otimes_n A_n$$

The value of G^{sa} at X is

$$G^{sa}(X) = \sum_{n \in \omega} \begin{bmatrix} X \\ n \end{bmatrix} \otimes_n G(n]$$

Thus

$$W(A)(X) = A^{sa}(X) = \sum_{n,m\in\omega} \begin{bmatrix} X\\ n \end{bmatrix} \otimes_n (n)^m \otimes_m A_m$$

The unit of the adjunction $i^{sa} \dashv (-)^{sa}$ at X

$$(\eta_A)_X : A(X) \longrightarrow A^{sa}(X)$$

is given by

$$[\vec{x}, a] \mapsto [\vec{x}, 1_n, a]$$

where $\vec{x}: (n] \to X$ and $a \in A_n$.

The counit of the adjunction at X

$$(\varepsilon_G)_X : \sum_{n \in \omega} \begin{bmatrix} X \\ n \end{bmatrix} \otimes_n G(n) \longrightarrow G(X)$$

is given by

$$[\vec{x}, t] \mapsto G(\vec{x})(t)$$

where $\vec{x}: (n] \to X$ is an injection and $t \in G(n]$.

The multiplication in the monad \mathbbm{W}

$$(\mu_A)_X : \sum_{n,m,k\in\omega} \begin{bmatrix} X\\n \end{bmatrix} \otimes_n (n]^m \otimes_m (m]^k \otimes_k A_k \longrightarrow \sum_{n,k\in\omega} \begin{bmatrix} X\\n \end{bmatrix} \otimes_n (n]^k \otimes_k A_k$$

is given by composition

$$[\vec{x}, g, f, a] \mapsto [\vec{x}, g \circ f, a]$$

where $\vec{x}: (n] \to X, g: (m] \to (n], f: (k] \to (m], and a \in A_k$. This ends the definition of the monad W.

Now we shall describe the monoidal structure on W. If B is another analytic functor, the n-coefficient of the composition $A \circ B$ is given by

$$(A \circ B)_n = \sum_{m, n_1, \dots, n_m \in \omega, \sum_{i=1}^m n_i = n} (S_n \times B_{n_1} \times \dots \times B_{n_m} \times A_m)_{/\sim_n}$$

where the equivalence relation \sim_n is such that for $\sigma \in S_n$, $\sigma_i \in S_{n_i}$, $\tau \in S_m$, $b_i \in B_i$, for $i \in (m]$ and $a \in A_m$ we have

$$\langle \sigma, \sigma_1 \cdot b_1, \dots, \sigma_m \cdot b_m, \tau \cdot a \rangle \sim_n \langle \sigma \circ (\sigma_1, \dots, \sigma_m) \star \tau, b_{\tau(1)}, \dots, b_{\tau(m)}, a \rangle$$

The *n*-th coefficient of $W(A) \circ W(A)$ is given by

$$(\mathbb{W}(A) \circ \mathbb{W}(A))_n = \sum_{m, n_i, k_i \in \omega, \sum_{i=1}^m n_i = n} (S_n \times (n_1]^{k_1} \otimes_{k_1} A_{k_1} \times \ldots \times (n_m]^{k_m} \otimes_{k_m} A_{k_m} \times A_m)_{/\sim_n}$$

and the *n*-coefficient of $W(A^2)$ is given by

$$(\mathbb{W}(A^2))_n = \sum_{m,k,k_i \in \omega, \sum_{i=1}^m k_i = k} (n]^k \times A_{k_1} \times \ldots \times A_{k_m} \times A_m$$

The coherence morphism φ for W at the *n*-th coefficient of the functor A is

$$\varphi_n : (\mathbb{W}(A) \circ \mathbb{W}(A))_n \longrightarrow (\mathbb{W}(A^2))_n$$

is given by

$$\langle \sigma, [\sigma_1, a_1] \dots [\sigma_m, a_m], \tau, a \rangle \mapsto \langle \sigma \circ (\sigma_1 \dots \sigma_m) \star \tau, a_{\tau(1)} \dots a_{\tau(m)}, a \rangle$$

Note that this map is well defined at the level of equivalence classes.

6 Equational theories vs regular operads

In this section we study the relations between regular equational theories and regular operads. We shall show that the square \mathcal{E}^{f}



(4)

commutes up to an isomorphism, with \mathcal{P}_{f}^{e} being an inclusion and both horizontal functors being equivalences of categories. \mathcal{P}_{f}^{o} was defined in Section 2 and \mathcal{E}_{o}^{f} was defined in Section 3. We shall define \mathcal{E}_{o}^{r} .

The functor $\mathcal{E}_{o}^{r} : \operatorname{RegOp} \to \operatorname{ET}$

Let \mathcal{O} be a regular operad. We define an equational theory $\mathcal{E}_o^r(\mathcal{O}) = (L, A)$. As the set of *n*-ary function symbols we put $L_n = \mathcal{O}_n$ for $n \in \omega$. The set of axioms A contains the following three kinds of equations in context:

- 1. *unit*: $\iota(x_1) = x_1 : \vec{x}^1$ where $\iota \in \mathcal{O}_1$ is the unit of the operad \mathcal{O} ;
- 2. action: $a(x_{\tau(1)}, \ldots, x_{\tau(m)}) = (\tau \cdot a)(x_1, \ldots, x_n) : \vec{x}^n$ for all $a \in \mathcal{O}_m$ and surjections $\tau : (m] \to (n];$
- 3. multiplication: $a(a_1(x_1, ..., x_{k_1}), ..., a_m(x_{k-k_m+1}, ..., x_k)) = ((a_1, ..., a_m) * a)(x_1, ... x_k) : \vec{x}^k$ where $a \in \mathcal{O}_m, a_i \in \mathcal{O}_{k_i}$ for $i \in 1, ..., m, k = \sum_{i=1}^m k_i$;

Clearly, all equations are regular and hence the theory $\mathcal{E}_{o}^{r}(\mathcal{O})$ is regular.

Suppose that $h: \mathcal{O} \to \mathcal{O}'$ is a morphism of regular operads. We define the interpretation

$$\mathcal{E}_{o}^{r}(h):\mathcal{E}_{o}^{r}(\mathcal{O})\longrightarrow\mathcal{E}_{o}^{r}(\mathcal{O}')$$

For $a \in \mathcal{O}_n$ we put

$$\mathcal{E}_o^r(h)(a) = h(a)(x_1, \dots, x_n) : \vec{x}^n$$

for $n \in \omega$. Clearly, $\mathcal{E}_o^r(h)$ is a regular interpretation.

Proposition 6.1. The square (4) commutes up to a natural isomorphism.

Proof. Let \mathcal{O} be a regular operad. We define an interpretation of equational theories

$$I_{\mathcal{O}}: \mathcal{E}^r_o(\mathcal{O}) \longrightarrow \mathcal{E}^f_o \mathcal{P}^o_f(\mathcal{O})$$

by

$$\mathcal{O}_n \ni a \mapsto [[1_n, a](x_1, \dots, x_n) : \vec{x}^n]$$

where $a \in \mathcal{O}_n$ and $[1_n, a]$ is an *n*-ary operation symbol of the theory $\mathcal{E}_o^f \mathcal{P}_f^o(\mathcal{O})$.

We need to verify that I is a well defined natural isomorphism. First we need to verify that $I_{\mathcal{O}}$ preserves axioms. The unit axiom is obvious. To prove the action axioms, we fix $a \in \mathcal{O}_m$ and $\tau : (m] \to (n]$ and we calculate using the theory $\mathcal{E}_o^f \mathcal{P}_f^o(\mathcal{O})$

$$I_{\mathcal{O}}(a(x_{\tau(1)}, \dots, x_{\tau(m)}) = [1_m, a](x_{\tau(1)}, \dots, x_{\tau(m)}) =$$
$$= \tau \cdot [1_m, a](x_1, \dots, x_n) = [1_m, \tau \cdot a](x_1, \dots, x_n) =$$
$$= I_{\mathcal{O}}(\tau \cdot a(x_1, \dots, x_n))$$

The calculations for the composition axioms are similar. The naturality of $I_{\mathcal{O}}$ is left for the reader.

We shall show that $I_{\mathcal{O}}$ is an isomorphism of theories. To this end we define an inverse interpretation

$$J_{\mathcal{O}}: \mathcal{E}_{o}^{f} \mathcal{P}_{f}^{o}(\mathcal{O}) \longrightarrow \mathcal{E}_{o}^{r}(\mathcal{O})$$

given by

$$\mathbb{I}(k,n) \otimes_k \mathcal{O}_k \ni [f,a] \mapsto a(x_{f(1)},\ldots,x_{f(k)}) : \vec{x}^n$$

Again we need to verify that $J_{\mathcal{O}}$ preserves the axioms and again we shall verify the action axiom only. Fix $\phi : [n) \to [m)$ and $[f, a] \in \mathbb{I}(k, n) \otimes \mathcal{O}_k$. Let $f' \circ \phi'$ be an epi-mono factorization in \mathbb{F} with $f' : [n') \to [m)$. Using the theory $\mathcal{E}_o^r(\mathcal{O})$, we have

$$J_{\mathcal{O}}((\phi \cdot [f, a])(x_1, \dots, x_n)) = [f', \phi' \cdot a](x_1, \dots, x_k) =$$

= $(\phi' \cdot a)(x_{f'(1),\dots,x_{f'(n')}}) = a(x_{f' \circ \phi'(1),\dots,x_{f' \circ \phi'(n')}}) =$
= $a(x_{\phi \circ f(1),\dots,x_{\phi \circ f(n')}}) = J_{\mathcal{O}}([f, a](x_{\phi(1),\dots,x_{\phi(n')}}))$

Finally, we need to verify that $I_{\mathcal{O}}$ and $J_{\mathcal{O}}$ are mutually inverse one to the other. The composition $J_{\mathcal{O}}I_{\mathcal{O}}$ sends operation $a \in \mathcal{O}_n$ to the term $a(x_1, \ldots, x_n) : \vec{x}^n$ so it is the identity. For an operation $[f, a] \in \mathbb{I}(k, n) \otimes_k \mathcal{O}_k$ in theory $\mathcal{E}^f_o \mathcal{P}^o_f(\mathcal{O})$ we have

$$I_{\mathcal{O}}J_{\mathcal{O}}([f,a]) = [1_n, a](x_{f(1)}, \dots, x_{f(n)}) : \vec{x}^n =$$
$$= f \cdot [1_n, a](x_1, \dots, x_k) : \vec{x}^k = [f, a](x_1, \dots, x_k) : \vec{x}^k$$

i.e. the composition $I_{\mathcal{O}}J_{\mathcal{O}}$ is the identity as well. \Box Finally, we have

Theorem 6.2. The functor $\mathcal{E}_o^r : \operatorname{RegOp} \longrightarrow \operatorname{RegET}$ is an equivalence of categories.

Proof. From Proposition 6.1 it follows that \mathcal{E}_o^r is faithful. To see that it is full consider a regular interpretation $I : \mathcal{E}_o^r(\mathcal{O}) \to \mathcal{E}_o^r(\mathcal{O}')$. We shall define a morphism of regular operads $h : \mathcal{O} \to \mathcal{O}'$ such that $\mathcal{E}_o^r(h) = I$. For $f \in \mathcal{O}_n$, $I(f)(\vec{x}^n) : \vec{x}^n$ is a regular term. But in theories in the image of \mathcal{E}_o^r every regular term is equal to a simple term, i.e. there is an $f' \in \mathcal{O}_n'$ such that $f'(\vec{x}^n) = I(f)(\vec{x}^n) : \vec{x}^n$ is a theorem in the theory $\mathcal{E}_o^r(\mathcal{O}')$. Thus we put h(f) = f'. It is left for the reader to verify that h has the required property.

To see that \mathcal{E}_o^r is essentially surjective let us fix a regular theory T = (L, A). Then the regular terms in T form a regular operad called T^{ro} . The unit is the term $x_1 : x_1$. The composition is defined by the substitution (making sure that we make the variable disjoint in different substituted term, via α -conversion). The action of a surjection $\phi : (n] \to (m]$ on a regular term $t(x_1, \ldots, x_n) : \vec{x}^n$ is again a regular term $t(x_{\phi(1)}, \ldots, x_{\phi(n)} : \vec{x}^n$. Again it is a matter of a routine verification that $\mathcal{E}_o^r(T^{ro}) \cong T$. \Box

Examples

- 1. Let $\mathbb{1}$ be the terminal equational theory. It has one constant, say e, and can be axiomatized by a single axiom: $v_1 = e : \vec{v}^1$. As a Lawvere theory it is the category that has exactly one morphism between any two objects. It is best seen at the level of Lawvere theories. Both theories $\mathbb{1}^a$ and the theory of commutative monoids are linear-regular and have exactly one analytic morphism $a : n \to 1$, for any n.
- 2. The functor $\mathcal{Q}_r^e : \mathbf{ET} \to \mathbf{RegET}$ is a right adjoint, it preserves the terminal object. Hence $\mathbb{1}^r$, the regular part $\mathbb{1}$, is the terminal regular theory. It is the theory of suplattices.

The embedding of the regular theories into all equational theories has a right adjoint $(-)^{sr}$, as well. The value of the functor $(-)^{sr}$ on the terminal equational theory 1 is the terminal regular theory, i.e. the theory of suplattices:

$$(x \lor y) \lor z = x \lor (y \lor z), \quad x \lor y = y \lor x, \quad x \lor \bot = x = \bot \lor x, \quad (x \lor x) = x$$

3. The theory 1 has a proper subcategory, in which $0 \not\cong 1$. It has no function symbols, and can be axiomatized by a single axiom: $v_1 = v^2 : \vec{v}^2$. The regular part of this theory is the theory of suplattices without a bottom element.

7 Cartesian and weakly cartesian monads

In this section we shall investigate two (strict monoidal) subcategories of **San** and their categories of monoids. The category of (weakly) cartesian functors and (weakly) cartesian natural transformations will be denoted by **Cart** (**wCart**). The corresponding categories of monoids: the category of (weakly) cartesian monads will be denoted by **CartMnd** (**wCartMnd**). Thus we have embeddings full on isomorphisms

$$\mathbf{Cart} \longrightarrow \mathbf{wCart} \longrightarrow \mathbf{San}$$

which are strict monoidal and induce embeddings of categories of monoids

$\mathbf{CartMnd} \longrightarrow \mathbf{wCartMnd} \longrightarrow \mathbf{SanMnd}$

The characterizations of the subcategories of equational theories **ET** and of Lawvere theories **LT** corresponding to **CartMnd** and **wCartMnd** are a bit technical and we are not going to describe it in detail here. Clearly, the objects are some regular theories satisfying additional conditions and similarly for morphisms. We shall content ourselves with a description of subcategories of $Set^{\mathbb{S}}$ whose essential images are **wCart** and **Cart**, respectively. Note, however, that if (T, η, μ) is a semi-analytic monad such that the functor part T is the left Kan extension of a functor $R : \mathbb{S} \to Set$ then R is the functor of all regular operations in the equational theory corresponding to the monad T. Thus our description will in fact provide a description of the equational theories corresponding to monads in **wCart** and **Cart**.

Remarks. If in a weak pullback square

$$P \xrightarrow{m} A$$

$$h \downarrow f$$

$$B \xrightarrow{g} C$$

the morphism m is mono then the square is a pullback. Since m is mono iff the square

$$P \xrightarrow{1_P} P$$

$$1_P \downarrow \qquad \qquad \downarrow m$$

$$P \xrightarrow{m} A$$

is a pullback, it follows that if a functor weakly preserves pullbacks it does preserve monos as well. Recall the description of functor $(-) : Set^{\mathbb{S}} \longrightarrow End$ from Section 2, the Kan extension along $i_{\mathbb{S}} : \mathbb{S} \to Set$. We begin with the following observation

Lemma 7.1. Let $A : \mathbb{S} \to Set$ be a functor, $f : X \to Y$ a function, $[\vec{x} : (n] \to X, a \in A_n], [\vec{x}' : (n'] \to X, a' \in A_{n'}] \in \hat{A}(X)$. If

$$\hat{A}([\vec{x},a]) = \hat{A}([\vec{x}',a'])$$

then there are $m \in \omega$, surjections $g: n \to m$, $g': n' \to m$, and an injection $\vec{y}: m \to Y$ as in the diagram

$$(n] \xrightarrow{\vec{x}} X$$

$$g | (n') \xrightarrow{\vec{x}'} | f$$

$$(m] \xrightarrow{\vec{y}} Y$$

such that $\vec{y} \circ g = f \circ \vec{x} \ \vec{y} \circ g' = f \circ \vec{x}'$, and A(g)(a) = A(g')(a'). In particular

$$\hat{A}([\vec{x},a]) = [\vec{y}, A(g)(a)] = [\vec{y}, A(g')(a')] = \hat{A}([\vec{x}',a'])$$

Proof. Exercise. \Box

The following two Propositions identify the subcategory of $Set^{\mathbb{S}}$ whose essential image in **End** is **wCart**.

Proposition 7.2. Let $A : \mathbb{S} \to Set$ be a functor. The functor $\hat{A} : Set \to Set$ weakly preserves pullbacks iff the functor A satisfies the following condition (WPB):

for any pullback of surjections in $\mathbb F$



and a pair of elements $a \in A(m]$, $b \in A(n]$ such that $A(f_1)(a) = A(g_1)(b)$ there is an injection $h: (q] \to (p]$ and an element $c \in A(q]$ such that $f_2 \circ h$ and $g_2 \circ h$ are surjections and

$$A(g_2 \circ h)(c) = a$$
, and $A(f_2 \circ h)(c) = b$

Proof. \Rightarrow First we verify that the condition (WPB) is necessary. Let us fix the square and elements a and b as in the above condition. Then we have elements

$$[1_{(m]}, a] \in \begin{bmatrix} (m] \\ m \end{bmatrix} \otimes_m A(m) \subseteq \sum_{k \in \omega} \begin{bmatrix} (m] \\ k \end{bmatrix} \otimes_k A(k) = \hat{A}(X)$$

and

$$[1_{(n]}, b] \in \begin{bmatrix} (n] \\ n \end{bmatrix} \otimes_n A(n) \subseteq \hat{A}(X)$$

such that

$$\hat{A}(f_1)([1_{(m]},a]) = \hat{A}(g_1)([1_{(n]},b])$$

As f_1 , g_1 are surjections this means that

$$A(f_1)(a) = A(g_1)(b)$$

As \hat{A} weakly preserves pullbacks there is an $[h, c] \in \hat{A}(p]$ such that

$$\hat{A}(f_2)([h,c]) = [1_{(n]}, b], \text{ and } \hat{A}(g_2)([h,c]) = [1_{(m]}, a]$$

Thus for some $h: (q] \to (p], c \in A(q)$ we have $[h, c] \in \begin{bmatrix} p \\ q \end{bmatrix} \otimes_q A(q)$ and using the diagram

$$\begin{array}{c} (q] \xrightarrow{g_3} (m'] \\ \downarrow \vec{x} \\ f_3 \\ \downarrow \\ f_2 \\ (n'] \xrightarrow{g_2} (m] \\ f_2 \\ \downarrow \\ (n'] \xrightarrow{g_1} (k] \end{array}$$

we have

$$[\vec{y}, A(f_3)(c)] = [1_{(n]}, b]$$
 and $[\vec{x}, A(g_3)(c)] = [1_{(m]}, a$

This means that n = n', \vec{y} is a bijection, $\vec{y} \circ f_3 = f_2 \circ h$ is a surjection and $b = A(\vec{y} \circ f_3) = A(f_2 \circ h)$. Similarly, we get that m = m', $g_2 \circ h$ is a surjection and $a = A(g_2 \circ h)$. Thus the condition (WPB) is necessary.

 \Leftarrow To show that the condition (WPB) is sufficient we suppose that a functor $A : \mathbb{S} \to Set$ satisfies (WPB) and we shall show that $\hat{A} : Set \to Set$ weakly preserves pullbacks.

Let us fix a pullback in *Set* and

$$\begin{array}{c|c} R & \xrightarrow{g_2} & Y \\ f_2 \downarrow & & \downarrow f_1 \\ Z & \xrightarrow{g_1} & X \end{array}$$
 (5)

and elements $[\vec{z},b]\in \hat{A}(Z)$ and $[\vec{y},a]\in \hat{A}(Y)$ such that

$$\hat{A}(f_1)([\vec{y}, a]) = \hat{A}(g_1)([\vec{z}, b])$$

i.e. in the diagram



we have

$$[\vec{y'}, A(f_3)(a)] = [\vec{z'}, A(g_3)(b)]$$

i.e. there is a permutation $\sigma \in S_k$ such that

$$\vec{y} \circ \sigma = \vec{z'}$$
 and $A(f_3)(a) = A(\sigma \circ g_3)(b)$

Still in the above diagram we can form a pullback in $\mathbb F$

$$\begin{array}{c} (q] \xrightarrow{g_4} (m] \\ f_4 \downarrow & \downarrow f_3 \\ (n] \xrightarrow{\sigma \circ g_3} (k] \end{array}$$

and we have a morphism $v:(q] \to R$ into the pullback (5) such that

$$f_2 \circ v = \vec{z} \circ f_4, \quad g_2 \circ v = \vec{y} \circ g_4$$

Thus by assumption there is an injection $h: (r] \to (q]$ and a $c \in A(q)$ such that both $f_4 \circ h$ and $g_4 \circ h$ are surjections and

$$A(g_4 \circ h)(c) = a, \quad A(f_4 \circ h)(c) = b$$

From diagram (6) we see that $f_4 \circ h$ is a surjection, \vec{z} is an injection and $f_2 \circ v \circ h = \vec{z} \circ f_4 \circ h$. Thus

$$\hat{A}(f_2)(\hat{A}(v)([h,c]) = [\vec{z}, A(f_4 \circ h)(c) = [\vec{z}, b]$$

Similarly

$$\hat{A}(g_2)(\hat{A}(v)([h,c]) = [\vec{y},a]$$

Thus $\hat{A}(v)([h,c]) \in \hat{A}(P)$ is the sought element in the weak pullback

$$[h,c] \in \hat{A}(q] \xrightarrow{\hat{A}(v)} \hat{A}(P) \xrightarrow{\hat{A}(g_2)} \hat{A}(Y)$$
$$\hat{A}(f_2) \downarrow \qquad \qquad \downarrow \hat{A}(f_1)$$
$$\hat{A}(Z) \xrightarrow{\hat{A}(g_1)} \hat{A}(X)$$

Proposition 7.3. Let $\tau : A \to B$ be a natural transformation. Then τ is weakly cartesian iff $\hat{\tau} : \hat{A} \to \hat{B}$ is weakly cartesian.

Proof. Assume that $\tau: A \to B$ is a weakly cartesian natural transformation, $f: X \to Y$ a function. We shall show that the square

(6)

$$\begin{array}{c|c} \hat{A}(X) & \xrightarrow{\hat{\tau}_X} & \hat{B}(X) \\ \hat{A}(f) & & & & & \\ \hat{A}(Y) & & & & \\ \hat{A}(Y) & \xrightarrow{\hat{\tau}_Y} & \hat{B}(Y) \end{array}$$

is a weak pullback. Let us fix elements

$$[\vec{y}, a] \in \begin{bmatrix} Y\\ n \end{bmatrix} \otimes_n A(n) \subseteq \hat{A}(Y), \text{ and } [\vec{x}, b] \in \begin{bmatrix} X\\ m \end{bmatrix} \otimes_m B(m) \subseteq \hat{B}(X)$$

such that

$$\hat{B}(f)([\vec{x},b]) = \hat{\tau}([\vec{y},a]) \ (= [\vec{y},\tau_n(a)])$$

By Lemma 7.1 we have a surjection f'

$$(m] \xrightarrow{\vec{x}} X$$

$$f' \downarrow \qquad \qquad \downarrow f$$

$$(n] \xrightarrow{\vec{y}} Y$$

such that $B(f')(b) = \tau_n(a)$. Since τ is weakly cartesian, there is a $c \in A(m)$ such that

$$\tau_m(c) = b$$
, and $A(f')(c) = a$

Then

and

$$[\vec{x},c] \in \left[\begin{array}{c} X\\m\end{array}\right] \otimes_m A(m)$$

and moreover

$$\hat{A}(f)([\vec{x},c]) = [\vec{y},A(f')(c)] = [\vec{y},a]$$

 $\hat{\tau}_X([\vec{x}, c]) = [\vec{x}, \tau_m(c)] = [\vec{x}, b]$

To prove the converse let us assume that $\hat{\tau} : \hat{A} \to \hat{B}$ is a weakly cartesian natural transformation and that $f : (m] \to (n]$ is a surjection in S. We need to show that the square

$$\begin{array}{c|c} A(m] & \xrightarrow{\tau_m} B(m] \\ A(f) & & & B(f) \\ A(n) & & & T_n \end{array} \\ A(n) & \xrightarrow{\tau_n} B(n] \end{array}$$

is weakly cartesian. Fix $a \in A(n)$ and $b \in B(m)$ such that $\tau_n(a) = B(f)(b)$. Consider a weak pullback in Set

$$\begin{array}{c} \hat{A}(m] \xrightarrow{\hat{\tau}_m} \hat{B}(m] \\ \hat{A}(f) \\ \hat{A}(n] \xrightarrow{\hat{\tau}_n} \hat{B}(n] \end{array}$$

We have elements

$$[1_{(n]}, a] \in \begin{bmatrix} (n] \\ n \end{bmatrix} \otimes_n A(n) \subseteq \hat{A}(n], \text{ and } [1_{(m]}, b] \in \begin{bmatrix} (m] \\ m \end{bmatrix} \otimes_m B(m) \subseteq \hat{B}(m)$$

such that

$$\hat{\tau}_n([1_{(n]},a]) = [1_{(n]},\tau_n(a)] = [1_{(n]},B(f)(b)] = \hat{B}(f)([1_{(m]},b])$$

Thus, by our assumption, for some $k \in \omega$ there is an $[\vec{x}, c] \in \begin{bmatrix} m \\ k \end{bmatrix} \otimes_k A(k)$ such that

$$\hat{A}(f)([\vec{x},c]) = [1_{(n]},a], \text{ and } \hat{\tau}_m([\vec{x},c]) = [1_{(m]},b]$$

Thus $(\vec{x}, \tau_m(c)) \sim (1_{(m]}, b)$ and we have that $k = m, \vec{x}$ is a bijection, $(\vec{x}, c) \sim (1_{(m]}, A(\vec{x}(c)))$. Hence we also have $\tau_m(A(\vec{x})(c)) = b$. Moreover as the square

$$(m] \xrightarrow{\vec{x}} (m]$$

$$f \circ \vec{x} \downarrow \qquad \qquad \downarrow f$$

$$(n] \xrightarrow{1_{(n]}} (n]$$

commutes, we have

$$A(f)(A(\vec{x})(c)) = A(f \circ \vec{x})(c) = a$$

Thus $A(\vec{x})(c)$ is the element sought for a and b. Since f, a and b were arbitrary, τ is weakly cartesian. \Box

The final two Propositions identify the subcategory of $Set^{\$}$ whose essential image in **End** is **Cart**.

Proposition 7.4. Let $A : \mathbb{S} \to Set$ be a functor. The functor $\hat{A} : Set \to Set$ preserves pullbacks iff the functor A satisfies the condition (WPB) from the Proposition 7.2, and additionally satisfies the following condition (PB):

suppose that the square

$$\begin{array}{c} (p] \xrightarrow{g_2} (m] \\ f_2 \downarrow & \downarrow \\ (n] \xrightarrow{g_1} (k] \end{array}$$

is a pullback of surjections in \mathbb{F} , $\vec{x} : (q] \to (p]$, $\vec{x}' : (q'] \to (p]$ are two injections, $c \in A(q]$, $c' \in A(q')$ are two elements so that the functions

$$f_2 \circ \vec{x}, \quad f_2 \circ \vec{x}', \quad g_2 \circ \vec{x}, \quad g_2 \circ \vec{x}',$$

are surjections and

$$A(f_2 \circ \vec{x})(c) = A(f_2 \circ \vec{x}')(c'), \quad A(g_2 \circ \vec{x})(c) = A(g_2 \circ \vec{x}')(c').$$

Then q = q' and there is $\sigma \in S_q$ such that

$$\vec{x}' \circ \sigma = \vec{c}, \quad A(\sigma)(c) = c'.$$

Proof. Assume that A satisfies (WPB) and (PB). Thus, by Proposition 7.3, \hat{A} weakly preserves pullbacks. Let

$$P \xrightarrow{g_2} Y$$

$$f_2 \downarrow \qquad \qquad \downarrow f_1$$

$$Z \xrightarrow{g_1} X$$

be a pullback in Set. We shall show that the square

$$\begin{array}{c}
\hat{A}(P) \xrightarrow{\hat{A}(g_2)} \hat{A}(Y) \\
\hat{A}(f_2) \\
\hat{A}(Z) \xrightarrow{\hat{A}(g_1)} \hat{A}(X)
\end{array}$$

is a pullback, i.e. it satisfies the uniqueness condition. So let

$$[h,a] \in \begin{bmatrix} P \\ q \end{bmatrix} \otimes_q A(q), \text{ and } [h',a'] \in \begin{bmatrix} P \\ q' \end{bmatrix} \otimes_{q'} A(q')$$

be such that

$$\hat{A}(f_2)([h,a]) = \hat{A}(f_2)([h',a']), \text{ and } \hat{A}(g_2)([h,a]) = \hat{A}(g_2)([h',a'])$$

This implies the equalities of images of functions

$$im(f_2 \circ h) = im(f_2 \circ h'), \ im(g_2 \circ h) = im(g_2 \circ h'), \ im(g_1 \circ f_2 \circ h) = im(f_2 \circ h')$$

All these sets are finite, say, having n, m, and k elements, respectively. Thus we can form a commuting diagram



where the back square is a pullback of surjections (in \mathbb{F}) and all arrows from the back to the front are injections (for simplicity: inclusions). By definition the functions

$$f_4 \circ h, f_4 \circ h', g_4 \circ h, f_4 \circ h'$$

are surjections. Since \hat{A} preserves monos we also have

$$A(f_4 \circ h)(a) = A(f_4 \circ h')(a'), \quad A(g_4 \circ h)(a) = A(g_4 \circ h')(a')$$

Thus from the condition (PB) it follows that q = q' and there is a $\sigma \in S_q$ such that $h' \circ \sigma = h$ and $A(\sigma)(a) = a'$, i.e. [h, a] = [h', a'], as required.

To prove the converse we assume now that \hat{A} preserves pullbacks and we fix a pullback

$$(r) \xrightarrow{g_2} (m)$$

$$f_2 \downarrow \qquad \qquad \downarrow f_1$$

$$(n) \xrightarrow{g_1} (k)$$

in \mathbb{F} of surjections. As \hat{A} weakly preserves pullbacks, the above pullback is sent by \hat{A} to a weak pullback in *Set*. Then, using Lemma 7.1 it is easy to see that the condition (PB) expresses the fact that \hat{A} sends the above square to a pullback in *Set*. \Box

Proposition 7.5. Let $\tau : A \to B$ be a natural transformation. Then τ is cartesian iff $\hat{\tau} : \hat{A} \to \hat{B}$ is cartesian.

Proof. First assume that $\tau : A \to B$ is a cartesian natural transformation. Fix a function $f : X \to Y$. By Proposition 7.3 the square

$$\begin{array}{c|c} \hat{A}(X) & \xrightarrow{\tau_X} & \hat{B}(X) \\ \hat{A}(f) & & & & \\ \hat{A}(Y) & & & & \\ \hat{A}(Y) & \xrightarrow{\tau_Y} & \hat{B}(Y) \end{array}$$

is a weak pullback. We shall show that it also satisfies uniqueness property. Let us fix elements

$$[\vec{x}, a] \in \begin{bmatrix} X\\ n \end{bmatrix} \otimes_n A(n) \subseteq \hat{A}(X), \text{ and } [\vec{x}', a'] \in \begin{bmatrix} X\\ n' \end{bmatrix} \otimes_{n'} A(n') \subseteq \hat{A}(X)$$

such that

$$\hat{\tau}([\vec{x},a]) = \hat{\tau}([\vec{x}',a']), \text{ and } \hat{A}(f)([\vec{x},a]) = \hat{A}(f)([\vec{x}',a'])$$
(7)

The first equality means that $[\vec{x}, \tau_n(a)] = [\vec{x}', \tau_n(a')]$. Hence n = n' and there is a $\sigma \in S_n$ such that

$$\vec{x}' \circ \sigma = \vec{x}$$
, and $\tau_n(a') = B(\sigma)(\tau_n(a)) = \tau_n(A(\sigma)(a))$

By Lemma 7.1 and the second equality in (7) there are surjections g and g' and an injection \vec{y} as in the diagram



such that

$$\vec{y} \circ g = f \circ \vec{x}, \quad \vec{y}' \circ g' = f \circ \vec{x}', \quad \text{and} \quad A(g')(a') = A(g)(a)$$

Then we have

$$\vec{y} \circ g = f \circ \vec{x} \circ \sigma = \vec{y} \circ g' \circ \sigma$$

As \vec{y} is mono $g = g' \circ \sigma$. Thus

$$A(g')(a') = A(g)(a) = A(g')(A(\sigma)(a))$$

As τ is a cartesian natural transformation we get, from the fact that the naturality square for g' is a pullback, that $a' = A(\sigma)(a)$. But this means that

$$(\vec{x}',a') \sim (\vec{x}',A(\sigma)(a)) \sim (\vec{x}' \circ \sigma,a) \sim (\vec{x},a)$$

i.e. $[\vec{x}', a'] = [\vec{x}, a]$ and $\hat{\tau}$ is cartesian.

To show the converse assume that $\hat{\tau}$ is a cartesian natural transformation and $f: (n] \to (m]$ is a surjection. We need to show that the weak pullback

$$\begin{array}{c|c} A(n) & \xrightarrow{\tau_n} & B(n) \\ A(f) & & \downarrow B(f) \\ A(m) & \xrightarrow{\tau_m} & B(m) \end{array}$$

satisfies the uniqueness condition, as well. Fix $a, a' \in A(n)$ such that

$$A(f)(a) = A(f)(a')$$
, and $\tau_n(a) = \tau_n(a')$

By assumption the square

$$\begin{array}{c} \hat{A}(n) \xrightarrow{\tau_{(n]}} \hat{B}(n) \\ \hat{A}(f) & & & \\ \hat{A}(m) \xrightarrow{\tau_{(m]}} \hat{B}(m) \end{array}$$

is a pullback. We have elements $[1_{(n]}, a], [1_{(n]}, a'] \in \hat{A}(n]$ such that

$$\tau_{(n]}([1_{(n]}, a]) = [1_{(n]}, \tau_n(a)]) = [1_{(n]}, \tau_n(a')]) = \tau_{(n]}([1_{(n]}, a'])$$

and

$$\hat{A}(f)([1_{(n]},a]) = [1_{(n]},A(f)(a)] = [1_{(n]},A(f)(a')] = \hat{A}([1_{(n]},a'])$$

Thus $[1_{(n]}, a] = [1_{(n]}, a']$ and a = a', i.e. τ is a cartesian natural transformation. \Box

References

- [Barr] M. Barr, Coequalizers and Free Triples, Math. Z. 116, pp. 307-322 (1970).
- [Be] J. Beck, *Distributive laws*, Lecture Notes in Mathematics, vol 80, pp. 119140, (1969).
- [CJKP] A. Carboni, G. Janelidze, G.M. Kelly, P. Paré, On Localization and Stabilization for Factorization Systems, Applied Categorical Structures, vol 5, (1997), pp. 1-58.
- [FK] P. Freyd, G. M. Kelly, Categories of continous functors I, J. Pure Appl. Algebra vol 2, (1972), pp. 169-191.
- [KR] A. Kock, G. E. Reyes, *Doctrines in Categorical Logic*, In: Handbook of Mathematical Logic, J. Barwise ed., Studies in Logic and The Found. of Math, vol 90, North-Holland, (1977), pp. 283-313.
- [Lw] F. W. Lawvere, Functorial Semantics of Algebraic Theories and Some Algebraic Problems in the context of Functorial Semantics of Algebraic Theories, Ph.D. thesis, Columbia University, 1963. Reprinted in Theory and Applications of Categories, No. 5 (2004) pp. 1-121.
- [CWM] S. MacLane, Categories for the Working Mathematician, Graduate Texts in Mathematics 5, Springer-Verlag, New York 1998.
- [Pl] J. Płonka On a method of construction of abstract algebras, Fundamenta Mathematice, LXI, (1967), pp. 191-198.
- [Tr] S. N. Tronin, Abstract Clones and Operads, Siberian Mathematical Journal, Vol. 43, No. 4, pp. 746-755, 2002
- [SZ] S. Szawiel, M. Zawadowski, *Theories of analytic monads*, arXiv:1204.2703v2 [math.CT].
- M. Zawadowski, Formal Theory of Monoidal Monads, J. Pure Appl. Algebra vol 216, (2012), pp. 1932-1942.

Instytut Matematyki, Uniwersytet Warszawski, ul. S.Banacha 2, 00-913 Warszawa, Poland szawiel@mimuw.edu.pl, zawado@mimuw.edu.pl