

# A SEMI-ORDINARY $p$ -STABILIZATION OF SIEGEL EISENSTEIN SERIES FOR SYMPLECTIC GROUPS AND ITS $p$ -ADIC INTERPOLATION

HISA-AKI KAWAMURA

ABSTRACT. Given a prime number  $p$ , we introduce a certain  $p$ -stabilization of holomorphic Siegel Eisenstein series for the symplectic group  $\mathrm{Sp}(2n)_{/\mathbb{Q}}$  such that the resulting forms satisfy the semi-ordinary condition at  $p$ , that is, the associated eigenvalue of a generalized Atkin  $U_p$ -operator is a  $p$ -adic unit. In addition, we derive an explicit formula for all Fourier coefficients of such  $p$ -stabilized Siegel Eisenstein series, and conclude their  $p$ -adic interpolation problems. This states the existence of a quite natural generalization of the ordinary  $\Lambda$ -adic Eisenstein series which have been constructed by Hida and Wiles for  $\mathrm{GL}(2)_{/\mathbb{Q}}$ .

## 1. INTRODUCTION

Given a positive integer  $M$ , a Dirichlet character  $\chi$  modulo  $M$  and an integer  $\kappa \geq 2$  with  $\chi(-1) = (-1)^\kappa$ , the classical Eisenstein series is defined as follows: For each  $z \in \mathfrak{H}_1 = \{z \in \mathbb{C} \mid \mathrm{Im} z > 0\}$ , put

$$E_{\kappa, \chi}(z) (= E_{\kappa, \chi}^{(1)}(z)) := \left\{ \frac{2G(\chi^{-1})(-2\pi\sqrt{-1})^\kappa}{M^\kappa(\kappa-1)!} \right\}^{-1} \sum_{(c,d) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \chi^{-1}(d) (cMz + d)^{-\kappa},$$

where  $G(\chi^{-1}) = \sum_{n=1}^M \chi^{-1}(n) \exp(2\pi\sqrt{-1}n/M)$ . As is well-known,  $E_{\kappa, \chi}$  gives rise to a holomorphic modular form of weight  $\kappa$  and nebentypus character  $\chi$  for the congruence subgroup  $\Gamma_0(M)$  of  $\mathrm{SL}(2, \mathbb{Z})$  unless  $\kappa = 2$  and  $\chi$  is trivial (or identical). If  $\chi$  is trivial (i.e.,  $M = 1$ ) or primitive (i.e., the conductor of  $\chi$  equals  $M > 1$ ), then  $E_{\kappa, \chi}$  possesses the Fourier expansion

$$E_{\kappa, \chi}(z) = \frac{L(1-\kappa, \chi)}{2} + \sum_{m=1}^{\infty} \sigma_{\kappa-1, \chi}(m) q^m,$$

where  $L(s, \chi) = \prod_{l: \text{prime}} (1 - \chi(l)l^{-s})^{-1}$ ,  $\sigma_{\kappa-1, \chi}(m) = \sum_{0 < d \mid m} \chi(d) d^{\kappa-1}$  and  $q = \exp(2\pi\sqrt{-1}z)$ . This implies that  $E_{\kappa, \chi}$  is a non-cuspidal Hecke eigenform such that the associated  $L$ -function  $L(s, E_{\kappa, \chi})$  is taken of the form

$$L(s, E_{\kappa, \chi}) = \zeta(s)L(s-\kappa+1, \chi).$$

Let  $p$  be a fixed prime number not dividing  $M$ , which we assume to be odd for simplicity. Put

$$(1) \quad E_{\kappa, \chi}^*(z) := E_{\kappa, \chi}(z) - \chi(p)p^{\kappa-1}E_{\kappa, \chi}(pz).$$

We easily see that  $E_{\kappa, \chi}^*$  is also a non-cuspidal Hecke eigenform of weight  $\kappa$  and nebentypus character  $\chi$  for  $\Gamma_0(Mp) (\subset \Gamma_0(M))$ , and its Fourier expansion is taken of the form

$$(2) \quad E_{\kappa, \chi}^*(z) = \frac{L^{\{p\}}(1-\kappa, \chi)}{2} + \sum_{m=1}^{\infty} \sigma_{\kappa-1, \chi}^{\{p\}}(m) q^m,$$

---

*Date:* August 20th, 2018.

*2010 Mathematics Subject Classification.* 11F46 (Primary); 11F30, 11F33 (Secondary).

*Key words and phrases.* Siegel modular forms, Siegel Eisenstein series,  $\Lambda$ -adic forms,  $p$ -adic analytic families.

where  $L^{\{p\}}(s, \chi) := (1 - \chi(p)p^{-s})L(s, \chi) = \prod_{l \neq p} (1 - \chi(l)l^{-s})^{-1}$  and

$$\sigma_{\kappa-1, \chi}^{\{p\}}(m) := \sum_{\substack{0 < d \mid m, \\ \gcd(d, p) = 1}} \chi(d) d^{\kappa-1}.$$

It turns out that for  $E_{\kappa, \chi}^*$ , all the Hecke eigenvalues outside  $p$  agree with  $E_{\kappa, \chi}$ , but the eigenvalue of Atkin's  $U_p$ -operator, which is converted from the  $p$ -th Hecke operator  $T_p$  (cf. [AL, Mi]), is  $\sigma_{\kappa-1, \chi}^{\{p\}}(p) = 1$ . Namely, we have

$$L(s, E_{\kappa, \chi}^*) = \zeta(s)L^{\{p\}}(s - \kappa + 1, \chi).$$

This type of normalization  $E_{\kappa, \chi} \mapsto E_{\kappa, \chi}^*$  given by eliminating the  $p$ -part of every Fourier coefficient or equivalently, eliminating the latter half of the Euler factor at  $p$  was firstly introduced by Serre [Se], and we refer to it as the *ordinary*<sup>1</sup>  $p$ -*stabilization*. Here we should mention that every Fourier coefficient of  $E_{\kappa, \chi}^*$  depends on the weight  $\kappa$   $p$ -adically. Indeed, it follows immediately from Fermat's little theorem that for each positive integer  $m$ , the function  $\kappa \mapsto \sigma_{\kappa-1, \chi}^{\{p\}}(m)$  can be extended to an analytic function defined on the ring of  $p$ -adic integers  $\mathbb{Z}_p$ . In addition, the constant term  $L^{\{p\}}(1 - \kappa, \chi)/2$  is also interpolated by the  $p$ -adic  $L$ -function in the sense of Kubota-Leopoldt or Deligne-Ribet [DR]. This fact turns out to be the following theorem due to Hida and Wiles:

**Fact 1.1** (cf. Proposition 7.1.1 in [H1]; Proposition 1.3.1 in [W]). *Suppose that a Dirichlet character  $\chi$  modulo  $M$  is either trivial or primitive, and  $M$  is not divisible by  $p$ . Let  $K = \mathbb{Q}_p(\chi)$  be the field obtained by adjoining all values of  $\chi$  to the  $p$ -adic number field  $\mathbb{Q}_p$  and  $\mathcal{L} = K[[X]]$  the power series ring in one variable  $X$  over  $K$ , respectively. For each integer  $a$  with  $0 \leq a < p - 1$ , there exists a formal Fourier expansion*

$$\mathcal{E}_{\chi\omega^a}(X) (= \mathcal{E}_{\chi\omega^a}^{(1)}(X)) = \sum_{m=0}^{\infty} \mathcal{A}_{\chi\omega^a}(m; X) q^m \in \mathcal{L}[[q]],$$

where  $\omega : \mathbb{Z}_p^\times \rightarrow \mu_{p-1} = \{x \in \mathbb{Z}_p^\times \mid x^{p-1} = 1\}$ <sup>2</sup> denotes the Teichmüller character, such that for each integer  $\kappa > 2$  with  $\chi(-1) = (-1)^\kappa$  and  $\kappa \equiv a \pmod{p-1}$ <sup>3</sup>, we have

$$\mathcal{E}_{\chi\omega^a}((1+p)^\kappa - 1) = E_{\kappa, \chi}^*.$$

Moreover, if  $\varepsilon : 1 + p\mathbb{Z}_p \rightarrow \overline{\mathbb{Q}_p}^\times$  is a character having exact order  $p^r$  for some integer  $r \geq 0$ <sup>4</sup>, then for each integer  $\kappa \geq 2$ , we have

$$\mathcal{E}_{\chi\omega^a}(\varepsilon(1+p)(1+p)^\kappa - 1) = E_{\kappa, \chi\omega^{a-\kappa\varepsilon}}$$

as long as  $\omega^{a-\kappa\varepsilon}$  is non-trivial.

Strictly speaking, it turns out that  $\mathcal{E}_{\chi\omega^a}(X) \in \Lambda[[q]]$ , where  $\Lambda = \mathcal{O}_K[[X]]$ , unless  $\chi\omega^a$  is trivial. However, we note that if  $\chi\omega^a$  is trivial, then  $\mathcal{E}_{\chi\omega^a}(X)$  (more precisely, the constant term  $\mathcal{A}_{\chi\omega^a}(0; X)$ ) has a simple pole at  $X = 0$  and  $\overline{\mathcal{E}}_{\chi\omega^a}(X) := X \cdot \mathcal{E}_{\chi\omega^a}(X) \in \Lambda[[q]]$ . Hence the above-mentioned fact implies that  $\mathcal{E}_{\chi\omega^a}(X)$  or  $\overline{\mathcal{E}}_{\chi\omega^a}(X)$  according as  $\chi\omega^a$  is non-trivial or trivial, is a  $\Lambda$ -adic form of level  $Mp^\infty$  with nebentypus character  $\chi\omega^a$ , which interpolates  $p$ -adic families of non-cuspidal ordinary Hecke eigenforms  $\{E_{\kappa, \chi}^*\}$  and  $\{E_{\kappa, \chi\omega^{a-\kappa\varepsilon}}\}$  or their constant multiples, given by varying the weight  $\kappa$

<sup>1</sup>In general, a Hecke eigenform  $f$  is said to be *ordinary* at  $p$  if the eigenvalue of  $U_p$  (or  $T_p$ ) is a  $p$ -adic unit.

<sup>2</sup>Since  $\mu_{p-1} \simeq (\mathbb{Z}/p\mathbb{Z})^\times$ ,  $\omega$  can be identified with a Dirichlet character modulo  $p$  in a natural way.

<sup>3</sup>The latter condition is regarded as if  $\omega^{a-\kappa}$  corresponds to the trivial Dirichlet character.

<sup>4</sup>Namely,  $\varepsilon$  can be also regarded as a primitive Dirichlet character of conductor  $p^{r+1}$ .

$p$ -adically analytically (cf. [W, H1]). In this context, we refer to it as the *ordinary  $\Lambda$ -adic Eisenstein series* of genus 1 and level  $Mp^\infty$  with character  $\chi\omega^a$ .

The aim of the present article is to formulate a similar statement in the case of Siegel modular forms, that is, automorphic forms on the symplectic group  $\mathrm{Sp}(2n)/\mathbb{Q}$  of an arbitrary genus  $n \geq 1$ . Let us explain how it goes briefly: Given a positive integer  $\kappa > n + 1$  and a Dirichlet character  $\chi$  modulo  $M$ , let  $E_{\kappa, \chi}^{(n)}$  be the classical (holomorphic) Siegel Eisenstein series of weight  $\kappa$  and nebentypus character  $\chi$  for  $\Gamma_0(M)^{(n)} \subset \mathrm{Sp}(2n, \mathbb{Z})$  to be described in the subsequent §2, whenever either

- (i)  $M = 1$  and thus,  $\chi$  is trivial or
- (ii)  $M > 1$  is odd,  $\chi$  is primitive and  $\chi^2$  is locally non-trivial at every prime  $l \mid M^5$ .

For any prime number  $p \nmid M$  (not necessarily odd), we define, in §4, a certain  $p$ -stabilization map  $E_{\kappa, \chi}^{(n)} \mapsto (E_{\kappa, \chi}^{(n)})^*$  by means of the action of a linear combination of  $(U_{p,n})^i$ 's for  $i = 0, 1, \dots, n$ , where  $U_{p,n}$  denotes a generalized Atkin  $U_p$ -operator to be defined in §3 below, so that the associated eigenvalue of  $U_{p,n}$  is 1. This formulation is natural from the viewpoint of which in the case where  $n = 1$ ,  $E_{\kappa, \chi}^* = (E_{\kappa, \chi}^{(1)})^*$  can be easily written in terms of Atkin's original operator  $U_p = U_{p,1}$  as

$$(1') \quad E_{\kappa, \chi}^* = E_{\kappa, \chi} \parallel_{\kappa} (U_p - \chi(p)p^{\kappa-1}).$$

Following in the tradition of Skinner-Urban [SU], in the case where  $n > 1$ , we call  $(E_{\kappa, \chi}^{(n)})^*$  *semi-ordinary* at  $p$ . Moreover, we derive an explicit formula for all Fourier coefficients of  $(E_{\kappa, \chi}^{(n)})^*$ , which can be regarded as a natural generalization of the equation (2) (cf. Theorem 4.2 below). As a consequence of the above issues, we first show in Theorem 4.4 below the existence of a formal Fourier expansion with coefficients in  $\mathcal{L} = \mathrm{Frac}(\Lambda)$  associated to  $\chi\omega^a$  with some nonnegative integer  $a$ , whose specialization at  $X = (1 + p)^\kappa - 1$  coincides with the semi-ordinary  $p$ -stabilized Siegel Eisenstein series  $(E_{\kappa, \chi}^{(n)})^*$  for any  $\kappa$  taken as above. Finally in §5, for a fixed odd prime number  $p$ , we show that after taking a suitable constant multiple, the above-mentioned formal Fourier expansion is indeed a  $\Lambda$ -adic form of genus  $n$  and level  $Mp^\infty$  (or tame level  $M$  in the sense of Taylor [Ta]) with character  $\chi\omega^a$ , which can be viewed as a satisfactory generalization of Fact 1.1 (cf. Theorem 5.4 below).

**Acknowledgements.** The author is deeply grateful to Professors S. Böcherer, H. Hida, T. Ikeda, H. Katsurada, A.A. Panchishkin, V. Pilloni, R. Schulze-Pillot and J. Tilouine for their valuable suggestions and comments. This research was partially supported by the JSPS Grant-in-Aid for Young Scientists (No.26800016).

*Notation.* We summarize here some notation we will use in the sequel. We denote by  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}$ , and  $\mathbb{C}$  the ring of integers, fields of rational numbers, real numbers and complex numbers, respectively. Let  $\overline{\mathbb{Q}}$  denote the algebraic closure of  $\mathbb{Q}$  sitting inside  $\mathbb{C}$ . We put  $e(x) = \exp(2\pi\sqrt{-1}x)$  for  $x \in \mathbb{C}$ . Given a prime number  $p$ , we denote by  $\mathbb{Q}_p$ ,  $\mathbb{Z}_p$  and  $\mathbb{Z}_p^\times$  the field of  $p$ -adic numbers, the ring of  $p$ -adic integers and the group of  $p$ -adic units, respectively. Hereinafter, given a prime number  $p$ , we fix an algebraic closure  $\overline{\mathbb{Q}}_p$  of  $\mathbb{Q}_p$  and an embedding  $\iota_p : \overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$  once for all. Let  $\mathrm{val}_p$  denote the  $p$ -adic valuation on  $\overline{\mathbb{Q}}_p$  normalized so that  $\mathrm{val}_p(p) = 1$ , and  $|\cdot|_p$  the corresponding norm on  $\overline{\mathbb{Q}}_p$ , respectively. Let  $e_p$  be the continuous additive character of  $\overline{\mathbb{Q}}_p$  such that  $e_p(x) = e(x)$  for all  $x \in \mathbb{Q}$ . Let  $\mathbb{C}_p$  be the completion of the normed space  $(\overline{\mathbb{Q}}_p, |\cdot|_p)$ . We also fix, once for all, an isomorphism

---

<sup>5</sup>Namely, if we factor  $\chi$  as  $\chi = \prod_{l \mid M} \chi_l$ , then for each prime  $l$ ,  $\chi_l$  is not a quadratic character.

$\widehat{\iota}_p : \mathbb{C} \xrightarrow{\sim} \mathbb{C}_p$  such that the diagram

$$\begin{array}{ccc} \mathbb{C} & \xrightarrow[\simeq]{\widehat{\iota}_p} & \mathbb{C}_p \\ \uparrow & & \uparrow \\ \overline{\mathbb{Q}} & \xrightarrow[\iota_p]{} & \overline{\mathbb{Q}}_p \end{array}$$

is commutative. Given a prime number  $p$ , put

$$\mathbf{p} = \begin{cases} 4 & \text{if } p = 2, \\ p & \text{otherwise.} \end{cases}$$

Let  $\mathbb{Z}_{p,\text{tor}}^\times$  be the torsion subgroup of  $\mathbb{Z}_p^\times$ , that is,  $\mathbb{Z}_{2,\text{tor}}^\times = \{\pm 1\}$  and  $\mathbb{Z}_{p,\text{tor}}^\times = \mu_{p-1}$  if  $p \neq 2$ . We define the Teichmüller character  $\omega : \mathbb{Z}_p^\times \rightarrow \mathbb{Z}_{p,\text{tor}}^\times$  by putting  $\omega(x) = \pm 1$  according as  $x \equiv \pm 1 \pmod{4\mathbb{Z}_2}$  if  $p = 2$ , and  $\omega(x) = \lim_{i \rightarrow \infty} x^{p^i}$  for  $x \in \mathbb{Z}_p^\times$  if  $p \neq 2$ . In addition, for each  $x \in \mathbb{Z}_p^\times$ , put  $\langle x \rangle := \omega(x)^{-1}x \in 1 + \mathbf{p}\mathbb{Z}_p$ . Then we have a canonical isomorphism

$$\begin{array}{ccc} \mathbb{Z}_p^\times & \xrightarrow{\sim} & \mathbb{Z}_{p,\text{tor}}^\times \times (1 + \mathbf{p}\mathbb{Z}_p) \\ x & \mapsto & (\omega(x), \langle x \rangle). \end{array}$$

We note that the maximal torsion-free subgroup  $1 + \mathbf{p}\mathbb{Z}_p$  of  $\mathbb{Z}_p^\times$  is topologically cyclic, that is,  $1 + \mathbf{p}\mathbb{Z}_p = (1 + \mathbf{p})^{\mathbb{Z}_p}$ . As already mentioned in §1, the Teichmüller character  $\omega$  gives rise to a Dirichlet character  $\omega : (\mathbb{Z}/\mathbf{p}\mathbb{Z})^\times \simeq \mathbb{Z}_{p,\text{tor}}^\times \hookrightarrow \mathbb{C}_p^\times \simeq \mathbb{C}^\times$ . Let  $\varepsilon : 1 + \mathbf{p}\mathbb{Z}_p \rightarrow \overline{\mathbb{Q}}_p^\times$  be a character of finite order. More precisely, if  $\varepsilon$  has exact order  $p^m$  for some nonnegative integer  $m$ , then  $\varepsilon$  optimally factors through

$$1 + \mathbf{p}\mathbb{Z}_p / (1 + \mathbf{p}\mathbb{Z}_p)^{p^m} \simeq (1 + \mathbf{p})^{\mathbb{Z}_p} / (1 + \mathbf{p})^{p^m \mathbb{Z}_p} \simeq \mathbb{Z}_p / p^m \mathbb{Z}_p \simeq \mathbb{Z} / p^m \mathbb{Z}.$$

Since  $(\mathbb{Z}/p^m \mathbf{p}\mathbb{Z})^\times \simeq (\mathbb{Z}/\mathbf{p}\mathbb{Z})^\times \times \mathbb{Z}/p^m \mathbb{Z}$ , we may naturally regard  $\varepsilon$  as a Dirichlet character of conductor  $p^m \mathbf{p}$ . By abuse of notation, we will often identify such  $p$ -adic characters  $\omega$  and  $\varepsilon$  with the corresponding Dirichlet characters whose conductor is a powers of  $p$  in the sequel.

Given a positive integer  $n$ , let  $\text{GSp}(2n)$  be the group of symplectic similitudes over  $\mathbb{Q}$ , that is,

$$\text{GSp}(2n) := \{ g \in \text{GL}(2n) \mid {}^t g J g = \nu(g) J \text{ for some } \nu(g) \in \mathbb{G}_m \},$$

where  $J = \begin{bmatrix} 0_n & -1_n \\ 1_n & 0_n \end{bmatrix}$  with the  $n \times n$  unit (resp. zero) matrix  $1_n$  (resp.  $0_n$ ), and  $\text{Sp}(2n)$  the derived group of  $\text{GSp}(2n)$  characterized by the exact sequence

$$1 \rightarrow \text{Sp}(2n) \rightarrow \text{GSp}(2n) \xrightarrow{\nu} \mathbb{G}_m \rightarrow 1,$$

respectively. Namely,  $\text{GSp}(2) = \text{GL}(2)$  and  $\text{Sp}(2) = \text{SL}(2)$  in this setting. We note that every real point  $g = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \text{GSp}(2n, \mathbb{R})$  with  $\nu(g) > 0$ , where  $A, B, C, D \in \text{Mat}_{n \times n}(\mathbb{R})$ , acts on the Siegel upper-half space

$$\mathfrak{H}_n := \{ Z = X + \sqrt{-1}Y \in \text{Mat}_{n \times n}(\mathbb{C}) \mid {}^t Z = Z, Y > 0 \text{ (positive-definite)} \}$$

of genus  $n$  via the linear transformation  $Z \mapsto g \langle Z \rangle = (AZ + B)(CZ + D)^{-1}$ . If  $F$  is a function on  $\mathfrak{H}_n$ , then for each  $\kappa \in \mathbb{Z}$ , we define the slash action of  $g$  on  $F$  by

$$(F \parallel_\kappa g)(Z) := \nu(g)^{n\kappa - n(n+1)/2} \det(CZ + D)^{-\kappa} F(g \langle Z \rangle).$$

For each positive integer  $N$ , we shall consider the following congruence subgroups of level  $N$  for the full-modular group  $\text{Sp}(2n, \mathbb{Z})$ :

$$\Gamma_0(N)^{(n)} \text{ (resp. } \Gamma_1(N)^{(n)} \text{)} := \left\{ \gamma \in \text{Sp}(2n, \mathbb{Z}) \mid \gamma \equiv \begin{bmatrix} * & * \\ 0_n & * \end{bmatrix} \left( \text{resp. } \begin{bmatrix} * & * \\ 0_n & 1_n \end{bmatrix} \right) \pmod{N} \right\}.$$

For each  $\kappa \in \mathbb{Z}$ , let us denote by  $\mathcal{M}_\kappa(\Gamma_1(N))^{(n)}$  the space of (*holomorphic*) *Siegel modular forms* of genus  $n$ , weight  $\kappa$  and level  $N$ , that is,  $\mathbb{C}$ -valued holomorphic functions  $F$  on  $\mathfrak{H}_n$  satisfying the following two conditions:

- (i)  $F \parallel_{\kappa} \gamma = F$  for any  $\gamma \in \Gamma_1(N)^{(n)}$ .  
 (ii) For each  $\gamma \in \mathrm{Sp}(2n, \mathbb{Z})$ , the function  $F \parallel_{\kappa} \gamma$  possesses a Fourier expansion of the form

$$(F \parallel_{\kappa} \gamma)(Z) = \sum_{T \in \mathrm{Sym}_n^*(\mathbb{Z})} A_{F, \gamma}(T) e(\mathrm{tr}(TZ)),$$

where  $\mathrm{Sym}_n^*(\mathbb{Z})$  is the set of all half-integral symmetric matrices of degree  $n$  over  $\mathbb{Z}$ , namely,

$$\mathrm{Sym}_n^*(\mathbb{Z}) := \{T = [t_{ij}] \in \mathrm{Sym}_n(\mathbb{Q}) \mid t_{ii}, 2t_{ij} \in \mathbb{Z} \ (1 \leq i < j \leq n)\},$$

and  $\mathrm{tr}(\ast)$  denotes the trace. Then it is satisfied that

$$A_{F, \gamma}(T) = 0 \text{ unless } T \geq 0 \text{ (positive-semidefinite)}$$

for all  $\gamma \in \mathrm{Sp}(2n, \mathbb{Z})$ .

A Siegel modular form  $F \in \mathcal{M}_{\kappa}(\Gamma_1(N))^{(n)}$  is said to be *cuspidal* if it is satisfied that  $A_{F, \gamma}(T) = 0$  unless  $T > 0$  for all  $\gamma \in \mathrm{Sp}(2n, \mathbb{Z})$ . We denote by  $\mathcal{S}_{\kappa}(\Gamma_1(N))^{(n)}$  the subspace of  $\mathcal{M}_{\kappa}(\Gamma_1(N))^{(n)}$  consisting of all cuspidal forms. Given a Dirichlet character  $\chi : (\mathbb{Z}/N\mathbb{Z})^{\times} \rightarrow \mathbb{C}^{\times}$ , we denote by  $\mathcal{M}_{\kappa}(\Gamma_0(N), \chi)^{(n)}$  (resp.  $\mathcal{S}_{\kappa}(\Gamma_0(N), \chi)^{(n)}$ ) the subspace of  $\mathcal{M}_{\kappa}(\Gamma_1(N))^{(n)}$  (resp.  $\mathcal{S}_{\kappa}(\Gamma_1(N))^{(n)}$ ) consisting of all forms  $F$  with nebentypus character  $\chi$ , that is,

$$F \parallel_{\kappa} \gamma = \chi(\det D)F \text{ for any } \gamma = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \Gamma_0(N)^{(n)}.$$

In particular, whenever  $\chi$  is trivial, we naturally write  $\mathcal{M}_{\kappa}(\Gamma_0(N))^{(n)} = \mathcal{M}_{\kappa}(\Gamma_0(N), \mathrm{triv})^{(n)}$  and  $\mathcal{S}_{\kappa}(\Gamma_0(N))^{(n)} = \mathcal{S}_{\kappa}(\Gamma_0(N), \mathrm{triv})^{(n)}$ , respectively.

For a given pair of  $Z = [z_{ij}] \in \mathfrak{H}_n$  and  $T = [t_{ij}] \in \mathrm{Sym}_n^*(\mathbb{Z})$ , put

$$\mathbf{q}^T := e(\mathrm{tr}(TZ)) = \prod_{i=1}^n q_{ii}^{t_{ii}} \prod_{i < j \leq n} q_{ij}^{2t_{ij}},$$

where  $q_{ij} = e(z_{ij})$  ( $1 \leq i \leq j \leq n$ ). Since  $\begin{bmatrix} 1_n & S \\ 0_n & 1_n \end{bmatrix} \in \Gamma_1(N)^{(n)} \subset \Gamma_0(N)^{(n)}$  for each  $S \in \mathrm{Sym}_n(\mathbb{Z})$ , we easily see that if  $F \in \mathcal{M}_{\kappa}(\Gamma_1(N))^{(n)}$  (or  $\mathcal{M}_{\kappa}(\Gamma_0(N), \chi)^{(n)}$ ), then  $F$  possesses a Fourier expansion of the form

$$F(Z) = \sum_{\substack{T \in \mathrm{Sym}_n^*(\mathbb{Z}), \\ T \geq 0}} A_F(T) \mathbf{q}^T,$$

which is regarded as belonging to the ring  $\mathbb{C}[q_{ij}^{\pm 1} \mid 1 \leq i < j \leq n][[q_{11}, \dots, q_{nn}]]$ . Given a ring  $R$ , we write

$$R[[\mathbf{q}]]^{(n)} := R[q_{ij}^{\pm 1} \mid 1 \leq i < j \leq n][[q_{11}, \dots, q_{nn}]],$$

in a similar fashion to the notation of the ring of formal  $q$ -expansions  $R[[q]]$ . In particular, if  $F \in \mathcal{M}_{\kappa}(\Gamma_1(N))^{(n)}$  is a Hecke eigenform (i.e., a simultaneous eigenfunction of all Hecke operators whose similitude is coprime to  $N$ ), then it is well-known that the field  $K_F$  obtained by adjoining all Fourier coefficients (or equivalently, all Hecke eigenvalues) of  $F$  to  $\mathbb{Q}$  is an algebraic number field. Thus, by virtue of the presence of  $\iota_p$  and  $\widehat{\iota}_p$ , we may regard  $F \in K_F[[\mathbf{q}]]^{(n)}$  as sitting inside  $\mathbb{C}[[\mathbf{q}]]^{(n)}$  and  $\mathbb{C}_p[[\mathbf{q}]]^{(n)}$  interchangeably. For further details on the basic theory of Siegel modular forms set out above, see [AZ] or [Fr]. In particular, a comprehensive introduction to the theory of elliptic modular forms and Hecke operators can be found in [Mi].

## 2. SIEGEL EISENSTEIN SERIES FOR SYMPLECTIC GROUPS

In this section, we review some elementary facts on the Siegel Eisenstein series defined for  $\mathrm{Sp}(2n)_{/\mathbb{Q}}$  of an arbitrary genus  $n \geq 1$ . In particular, we describe an explicit form of its Fourier

expansion according to some previous works of Shimura (e.g., [S1, S2]), which is the starting point for the subsequent arguments.

Let  $N$  be a positive integer and  $\chi : (\mathbb{Z}/N\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$  a Dirichlet character, respectively. As mentioned in §1, for simplicity, we restrict ourselves to either of the following cases:

- (i)  $N = 1$ , that is,  $\chi$  is trivial;
- (ii)  $N > 1$  is odd,  $\chi$  is primitive and  $\chi^2$  is locally non-trivial at every prime  $l \mid N$ .

Given a positive integer  $n$ , if  $\kappa$  is an integer with  $\kappa > n + 1$  and  $\chi(-1) = (-1)^\kappa$ , then the (*holomorphic*) *Siegel Eisenstein series* of genus  $n$ , weight  $\kappa$  and level  $N$  with nebentypus character  $\chi$  is defined as follows: For each  $Z \in \mathfrak{H}_n$ , put

$$E_{\kappa, \chi}^{(n)}(Z) := 2^{-[(n+1)/2]} L(1 - \kappa, \chi) \prod_{i=1}^{[n/2]} L(1 - 2\kappa + 2i, \chi^2) \\ \times \sum_{\gamma = \begin{bmatrix} * & * \\ C & D \end{bmatrix} \in (P_{2n} \cap \Gamma_0(N)^{(n)}) \setminus \Gamma_0(N)^{(n)}} \chi^{-1}(\det D) \det(CZ + D)^{-\kappa},$$

where  $L(s, \psi)$  denotes Dirichlet's  $L$ -function associated with  $\psi$ , and  $P_{2n}$  the Siegel parabolic subgroup of  $\mathrm{Sp}(2n)$  consisting of all matrices  $g = \begin{bmatrix} * & * \\ 0_n & * \end{bmatrix}$ , respectively.

Let  $r$  be a positive integer. For each rational prime  $l$ , let  $\mathrm{Sym}_r^*(\mathbb{Z}_l)$  denote the set of all half-integral symmetric matrices of degree  $r$  over  $\mathbb{Z}_l$ . Given a nondegenerate  $S \in \mathrm{Sym}_r^*(\mathbb{Z}_l)$ , we define a formal power series  $b_l(S; X)$  in  $X$  by

$$b_l(S; X) := \sum_{R \in \mathrm{Sym}_r(\mathbb{Q}_l) / \mathrm{Sym}_r(\mathbb{Z}_l)} e_l(\mathrm{tr}(SR)) X^{\mathrm{val}_l(\mu_R)},$$

where  $\mu_R = [\mathbb{Z}_l^r + \mathbb{Z}_l^r R : \mathbb{Z}_l^r]$ . Put  $\mathfrak{D}_S := 2^{2[r/2]} \det S$ . We note that if  $r$  is even, then  $(-1)^{r/2} \mathfrak{D}_S \equiv 0$  or  $1 \pmod{4}$ , and thus, we may decompose it into the form

$$(-1)^{r/2} \mathfrak{D}_S = \mathfrak{d}_S \mathfrak{f}_S^2,$$

where  $\mathfrak{d}_S$  is the fundamental discriminant of the quadratic field extension  $\mathbb{Q}_l(\{(-1)^{r/2} \mathfrak{D}_S\}^{1/2}) / \mathbb{Q}_l$  and  $\mathfrak{f}_S = \{(-1)^{r/2} \mathfrak{D}_S / \mathfrak{d}_S\}^{1/2} \in \mathbb{Z}_l$ . Let  $\xi_l : \mathbb{Q}_l^\times \rightarrow \{\pm 1, 0\}$  denote the character defined by

$$\xi_l(x) = \begin{cases} 1 & \text{if } \mathbb{Q}_l(x^{1/2}) = \mathbb{Q}_l, \\ -1 & \text{if } \mathbb{Q}_l(x^{1/2}) / \mathbb{Q}_l \text{ is unramified,} \\ 0 & \text{if } \mathbb{Q}_l(x^{1/2}) / \mathbb{Q}_l \text{ is ramified.} \end{cases}$$

As shown in [Kil, Fe, S2], for each nondegenerate  $S \in \mathrm{Sym}_r^*(\mathbb{Z}_l)$ , there exists a polynomial  $F_l(S; X) \in \mathbb{Z}[X]$  whose constant term is 1 such that  $b_l(S; X)$  is decomposed as follows:

$$(3) \quad b_l(S; X) = F_l(S; X) \times \begin{cases} \frac{(1-X) \prod_{i=1}^{r/2} (1-l^{2i} X^2)}{1 - \xi_l((-1)^{r/2} \det S) l^{r/2} X} & \text{if } r \text{ is even,} \\ (1-X) \prod_{i=1}^{(r-1)/2} (1-l^{2i} X^2) & \text{if } r \text{ is odd} \end{cases}$$

(cf. Proposition 3.6 in [S2]). We note that  $F_l(S; X)$  satisfies the functional equation

$$(4) \quad F_l(S; l^{-r-1} X^{-1}) = F_l(S; X) \times \begin{cases} (l^{r+1} X^2)^{-\mathrm{val}_l(\mathfrak{f}_S)} & \text{if } r \text{ is even,} \\ \eta_l(S) (l^{(r+1)/2} X)^{-\mathrm{val}_l(\mathfrak{D}_S)} & \text{if } r \text{ is odd,} \end{cases}$$

where

$$\eta_l(S) := h_l(S) (\det S, (-1)^{(r-1)/2} \det S)_l (-1, -1)_l^{(r^2-1)/8}$$

in terms of the Hasse invariant  $h_l(S)$  in the sense of Kitaoka [Ki1] and the Hilbert symbol  $(*, *)_l$  defined over  $\mathbb{Q}_l$  (cf. Theorem 3.2 in [K1]). Thus, it turns out that  $F_l(S; X)$  has degree  $2\text{val}_l(\mathfrak{f}_S)$  or  $\text{val}_l(\mathfrak{D}_S)$  according as  $r$  is even or odd. We easily see that  $F_l(uS; X) = F_l(S; X)$  for each  $u \in \mathbb{Z}_l^\times$ , and that if  $S, T \in \text{Sym}_r^*(\mathbb{Z}_l)$  are equivalent over  $\mathbb{Z}_l$ , that is,  $T = {}^tUSU$  for some  $U \in \text{GL}(r, \mathbb{Z}_l)$ , then  $F_l(S; X) = F_l(T; X)$ . For further details on the above-mentioned issues, see [K1].

**Lemma 2.1.** *Let  $n, \kappa, N$  and  $\chi$  be taken as above.*

- (I)  $E_{\kappa, \chi}^{(n)} \in \mathcal{M}_\kappa(\Gamma_0(N), \chi)^{(n)}$  and it is a Hecke eigenform, that is, a simultaneous eigenfunction of Hecke operators defined at least for all primes not dividing the level  $N$ .
- (II) Let us consider a Fourier expansion of  $E_{\kappa, \chi}^{(n)}$  taken of the form

$$E_{\kappa, \chi}^{(n)}(Z) = \sum_{\substack{T \in \text{Sym}_n^*(\mathbb{Z}), \\ T \geq 0}} A_{\kappa, \chi}(T) \mathbf{q}^T.$$

Then every coefficient  $A_{\kappa, \chi}(T)$ , which is invariant under  $T \mapsto {}^tUTU$  for  $U \in \text{GL}(n, \mathbb{Z})$ , is described as follows:

- (IIa) For  $T = 0_n \in \text{Sym}_n^*(\mathbb{Z})$ , we have

$$A_{\kappa, \chi}(T) = 2^{-[(n+1)/2]} L(1 - \kappa, \chi) \prod_{i=1}^{[n/2]} L(1 - 2\kappa + 2i, \chi^2).$$

Therefore  $E_{\kappa, \chi}^{(n)}$  is not cuspidal.

- (IIb) If  $T \in \text{Sym}_n^*(\mathbb{Z})$  is taken of the form

$$T = \left[ \begin{array}{c|c} T' & \\ \hline & 0_{n-r} \end{array} \right]$$

for some nondegenerate  $T' \in \text{Sym}_r^*(\mathbb{Z})$  with  $0 < r \leq n$  (i.e.,  $\text{rank } T = \text{rank } T' = r$ ), then

$$A_{\kappa, \chi}(T) = 2^{[(r+1)/2] - [(n+1)/2]} \prod_{i=[r/2]+1}^{[n/2]} L(1 - 2\kappa + 2i, \chi^2) \\ \times \begin{cases} L(1 - \kappa + r/2, \left(\frac{\mathfrak{d}_{T'}}{*}\right) \chi) \prod_{l|f_{T'}} F_l(T'; \chi(l) l^{\kappa-r-1}) & \text{if } r \text{ is even,} \\ \prod_{l|\mathfrak{D}_{T'}} F_l(T'; \chi(l) l^{\kappa-r-1}) & \text{if } r \text{ is odd,} \end{cases}$$

where  $\left(\frac{\mathfrak{d}}{*}\right)$  denotes the Kronecker symbol.

**Remark 2.2.** For the convenience in the sequel, we make the convention for  $r = 0$ , that  $\mathfrak{D}_S = \mathfrak{d}_S = \mathfrak{f}_S = 1$ , and  $F_l(S; X) = 1$  for all primes  $l$ . This enables us to regard (IIa) as (IIb) for  $r = 0$ .

*Proof.* The assertion (I) is well-known. The assertion (II) can be obtained by exploiting an idea of Shimura [S2] as follows: Let  $\mathbb{A}$  be the ring of adèles over  $\mathbb{Q}$ ,  $G_{2n}(\mathbb{A}) = \text{Sp}(2n, \mathbb{A})$ ,  $P_{2n}(\mathbb{A}) = M_{2n}(\mathbb{A})N_{2n}(\mathbb{A})$  a Levi decomposition of the Siegel parabolic subgroup, where

$$M_{2n} := \left\{ \left[ \begin{array}{cc} A & 0_n \\ 0_n & {}^tA^{-1} \end{array} \right] \mid A \in \text{GL}(n) \right\}, \quad N_{2n} := \left\{ \left[ \begin{array}{cc} 1_n & B \\ 0_n & 1_n \end{array} \right] \mid {}^tB = B \right\},$$

and  $\chi : \mathbb{A}^\times / \mathbb{Q}^\times \rightarrow \mathbb{C}^\times$  the unitary Hecke character corresponding to  $\chi$ , respectively. For each  $s \in \mathbb{C}$ , let  $\text{Ind}_{P_{2n}(\mathbb{A})}^{G_{2n}(\mathbb{A})}(\chi \cdot |*|_{\mathbb{A}}^s)$  denote the normalized smooth representation induced from the character of  $\text{GL}(n, \mathbb{A}) \simeq M_{2n}(\mathbb{A})$  defined by  $A \mapsto \chi(\det A) |\det A|_{\mathbb{A}}^s$ , where  $|*|_{\mathbb{A}}$  denotes the norm on  $\mathbb{A}$ .

Choosing a suitable section  $\varphi^{(s)} \in \text{Ind}_{P_{2n}(\mathbb{A})}^{G_{2n}(\mathbb{A})}(\chi \cdot |*|_{\mathbb{A}}^s)$ , we define the Eisenstein series  $\mathbf{E}(\varphi^{(s)})$  on  $G_{2n}(\mathbb{A})$  by

$$\mathbf{E}(\varphi^{(s)})(g) := \sum_{\gamma \in P_{2n} \backslash G_{2n}} \varphi^{(s)}(\gamma g),$$

which converges absolutely for  $\text{Re}(s) \gg 0$ , and  $\mathbf{E}(\varphi^{(\kappa-(n+1)/2)})$  evaluates

$$2^{[(n+1)/2]} L(1 - \kappa, \chi)^{-1} \prod_{i=1}^{[n/2]} L(1 - 2\kappa + 2i, \chi^2)^{-1} E_{\kappa, \chi}^{(n)}.$$

Thus, the desired equation can be obtained from an explicit formula for Fourier coefficients of  $\mathbf{E}(\varphi^{(s)})$ , more precisely, local Whittaker functions  $\text{Wh}_T(\varphi_v^{(s)})$  defined on  $G_{2n}(\mathbb{Q}_v)$  for all places  $v$  of  $\mathbb{Q}$ , where  $\mathbb{A} = \prod_v \mathbb{Q}_v$ ,  $\chi = \prod_v \chi_v$  and  $\varphi^{(s)} = \prod_v \varphi_v^{(s)}$ . Whenever  $v$  is archimedean (i.e.,  $v = \infty$ ) or non-archimedean at which  $\chi_v$  is unramified (i.e.,  $v$  is a prime  $l$  not dividing  $N$ ), it has been proved by Shimura (cf. Equations 4.34-35K in [S1] and Proposition 7.2 in [S2]). Whenever  $v$  is a non-archimedean place at which  $\chi_v$  is ramified and  $\chi_v^2$  is non-trivial, the local Whittaker function  $\text{Wh}_T(\varphi_v^{(s)})$  is described by Takemori [T2] in which the argument relies upon a functional equation of  $\text{Wh}_T(\varphi_v^{(s)})$  due to Ikeda [Ik] (generalizing [Sw]).  $\square$

### 3. GENERALIZED ATKIN $U_p$ -OPERATOR

In this section, we recall the theory of Atkin's  $U_p$ -operator and its generalization. For further details on the facts set out below, see, for instance, [AZ, B , Ta].

Let  $p$  be a prime number and  $N$  a positive integer, respectively. If  $p \mid N$ , the double coset

$$\Gamma_0(N)^{(1)} \begin{bmatrix} 1 & 0 \\ 0 & p \end{bmatrix} \Gamma_0(N)^{(1)} = \bigsqcup_{s=0}^{p-1} \Gamma_0(N)^{(1)} \begin{bmatrix} 1 & s \\ 0 & p \end{bmatrix}$$

induces the following operator  $U_p = U_{p,1}$  acting on  $\mathcal{M}_{\kappa}(\Gamma_0(N), \chi)^{(1)}$ : For each  $f \in \mathcal{M}_{\kappa}(\Gamma_0(N), \chi)^{(1)}$ , put

$$(f \parallel_{\kappa} U_p)(z) := \sum_{s=0}^{p-1} \left( f \parallel_{\kappa} \begin{bmatrix} 1 & s \\ 0 & p \end{bmatrix} \right) (z) = p^{-1} \sum_{s=0}^{p-1} f\left(\frac{z+s}{p}\right) \in \mathcal{M}_{\kappa}(\Gamma_0(N), \chi)^{(1)}.$$

We easily see that it is written in terms of Fourier expansions as

$$(5) \quad f(z) = \sum_{m=0}^{\infty} a(m) q^m \mapsto (f \parallel_{\kappa} U_p)(z) = \sum_{m=0}^{\infty} a(pm) q^m$$

and this is still valid even if  $p \nmid N$ , however it maps from  $\mathcal{M}_{\kappa}(\Gamma_0(N), \chi)^{(1)}$  to  $\mathcal{M}_{\kappa}(\Gamma_0(Np), \chi)^{(1)}$  in this case. We refer to the operator  $U_p$ , regardless of whether  $p \mid N$  or not, as Atkin's  $U_p$ -operator.

**Remark 3.1.** Obviously,  $U_p$  coincides with the usual Hecke operator  $T_p$  if  $p \mid N$ . However,  $U_p$  is slightly different from  $T_p$  in general:

$$\Gamma_0(N)^{(1)} \begin{bmatrix} 1 & 0 \\ 0 & p \end{bmatrix} \Gamma_0(N)^{(1)} = \bigsqcup_{s=0}^{p-1} \Gamma_0(N)^{(1)} \begin{bmatrix} 1 & s \\ 0 & p \end{bmatrix} \sqcup \Gamma_0(N)^{(1)} \begin{bmatrix} p & 0 \\ 0 & 1 \end{bmatrix}$$

if  $p \nmid N$ .

Similarly, if  $n > 1$  and  $p \mid N$ , the following  $n$  double-coset operators at  $p$  are relevant for Siegel modular forms of genus  $n$  and level  $N$ :

$$U_{p,i} := \begin{cases} \Gamma_0(N)^{(n)} \operatorname{diag}(\underbrace{1, \dots, 1}_i, \underbrace{p, \dots, p}_{n-i}, \underbrace{p^2, \dots, p^2}_i, \underbrace{p, \dots, p}_{n-i}) \Gamma_0(N)^{(n)} & \text{if } 1 \leq i \leq n-1, \\ \Gamma_0(N)^{(n)} \operatorname{diag}(\underbrace{1, \dots, 1}_n, \underbrace{p, \dots, p}_n) \Gamma_0(N)^{(n)} & \text{if } i = n. \end{cases}$$

We note that if  $N$  is divisible by  $p$ , these operators  $U_{p,1}, \dots, U_{p,n-1}$  and  $U_{p,n}$  generate the dilating Hecke algebra at  $p$  acting on  $\mathcal{M}_\kappa(\Gamma_0(N), \chi)^{(n)}$ . In particular, we are interested in the operator  $U_{p,n}$  which plays a central role among them. Namely, we define the operator  $U_{p,n}$  on  $\mathbb{C}_p[[\mathbf{q}]]^{(n)}$  by

$$(6) \quad F = \sum_{T \geq 0} A_F(T) \mathbf{q}^T \mapsto F \parallel_\kappa U_{p,n} = \sum_{T \geq 0} A_F(pT) \mathbf{q}^T.$$

Indeed, we easily see that if  $F \in \mathcal{M}_\kappa(\Gamma_0(N), \chi)^{(n)}$  with some positive integers  $\kappa, N$  and a Dirichlet character  $\chi$ , then

$$F \parallel_\kappa U_{p,n} \in \begin{cases} \mathcal{M}_\kappa(\Gamma_0(N), \chi)^{(n)} & \text{if } p \mid N, \\ \mathcal{M}_\kappa(\Gamma_0(Np), \chi)^{(n)} & \text{if } p \nmid N \end{cases}$$

and the action of  $U_{p,n}$  commutes with those of the Hecke operators defined outside  $Np$ .

#### 4. SEMI-ORDINARY $p$ -STABILIZATION OF SIEGEL EISENSTEIN SERIES

Let us fix a positive odd integer  $M$  and a prime number  $p$  not dividing  $M$  once for all. In this section, for a pair of positive integers  $n, \kappa$  and a Dirichlet character  $\chi$  modulo  $M$  taken as in §2, we introduce a certain  $p$ -stabilization of the Siegel Eisenstein series  $E_{\kappa, \chi}^{(n)} \in \mathcal{M}_\kappa(\Gamma_0(M), \chi)$  such that it can be viewed as a natural generalization of the ordinary  $p$ -stabilization  $E_{\kappa, \chi}^{(1)} \mapsto (E_{\kappa, \chi}^{(1)})^*$  (cf. Equations (1) and (1')). Moreover, we derive an explicit form of the associated Fourier expansion

$$(E_{\kappa, \chi}^{(n)})^*(Z) = \sum_{\substack{T \in \operatorname{Sym}_n^*(\mathbb{Z}), \\ T \geq 0}} A_{\kappa, \chi}^*(T) \mathbf{q}^T,$$

which is expressed in a similar fashion to (2), and conclude its  $p$ -adic interpolation problem.

To begin with, we introduce the following two polynomials in  $X$  and  $Y$ :

$$\begin{aligned} \mathcal{P}_p^{(n)}(X, Y) &:= (1 - p^n XY) \prod_{i=1}^{[n/2]} (1 - p^{2n-2i+1} X^2 Y), \\ \mathcal{R}_p^{(n)}(X, Y) &:= \prod_{j=1}^n (1 - p^{j(2n-j+1)/2} X^j Y). \end{aligned}$$

In addition, let us denote by  $\tilde{\mathcal{R}}_p^{(n)}(X, Y)$  the reflected polynomial of  $\mathcal{R}_p^{(n)}(X, Y)$  with respect to  $Y$ , that is,

$$(7) \quad \tilde{\mathcal{R}}_p^{(n)}(X, Y) := Y^n \mathcal{R}_p^{(n)}(X, Y^{-1}) = \prod_{j=1}^n (Y - p^{j(2n-j+1)/2} X^j).$$

**Remark 4.1.** Whenever  $n = 1$  and  $2$ , a straightforward calculation yields

$$\begin{cases} \mathcal{P}_p^{(1)}(X, Y) = \mathcal{R}_p^{(1)}(X, Y) = 1 - pXY, \\ \mathcal{P}_p^{(2)}(X, Y) = \mathcal{R}_p^{(2)}(X, Y) = (1 - p^2XY)(1 - p^3X^2Y). \end{cases}$$

We note that if  $n > 2$ , then  $\mathcal{P}_p^{(n)}(X, Y) \neq \mathcal{R}_p^{(n)}(X, Y)$ , however,  $\mathcal{P}_p^{(n)}(X, 1)$  divides  $\mathcal{R}_p^{(n)}(X, 1)$  in general. For readers' convenience, we reveal the origins of these two polynomials here: Obviously, the former  $\mathcal{P}_p^{(n)}(X, Y)$  is relevant to the local factor at  $p$  of the Fourier coefficient  $A_{\kappa, \chi}(0_n)$  described in Lemma 2.1 (IIa):

$$(8) \quad \mathcal{P}_p^{(n)}(\chi(p)p^{\kappa-n-1}, 1) = (1 - \chi(p)p^{\kappa-1}) \prod_{i=1}^{[n/2]} (1 - \chi^2(p)p^{2\kappa-2i-1}).$$

The latter  $\mathcal{R}_p^{(n)}(X, Y)$  has been introduced by Kitaoka [Ki2] and Böcherer-Sato [BS] to describe the denominator of the formal power series  $\sum_{m=0}^{\infty} F_p(p^m S; X)Y^m$  for each nondegenerate  $S \in \text{Sym}_n^*(\mathbb{Z}_p)$  (cf. Equation (11) below).

Now, we introduce a  $p$ -stabilization of the Siegel Eisenstein series  $E_{\kappa, \chi}^{(n)} \in \mathcal{M}_{\kappa}(\Gamma_0(M), \chi)^{(n)}$  in terms of a linear combination of  $(U_{p,n})^i = \underbrace{U_{p,n} \circ \cdots \circ U_{p,n}}_i$  for  $i = 0, 1, \dots, n$  as follows:

**Theorem 4.2.** *For a pair of  $n, \kappa$  and  $\chi$  taken as above, put*

$$(9) \quad (E_{\kappa, \chi}^{(n)})^* := \frac{\mathcal{P}_p^{(n)}(\chi(p)p^{\kappa-n-1}, 1)}{\mathcal{R}_p^{(n)}(\chi(p)p^{\kappa-n-1}, 1)} \cdot E_{\kappa, \chi}^{(n)} \parallel_{\kappa} \tilde{\mathcal{R}}_p^{(n)}(\chi(p)p^{\kappa-n-1}, U_{p,n}).$$

Then we have

- (I)  $(E_{\kappa, \chi}^{(n)})^* \in \mathcal{M}_{\kappa}(\Gamma_0(Mp), \chi)^{(n)}$  and it is a Hecke eigenform such that all the eigenvalues outside  $Mp$  agree with those of  $E_{\kappa, \chi}^{(n)} \in \mathcal{M}_{\kappa}(\Gamma_0(M), \chi)^{(n)}$ .
- (II) If  $0 \leq T \in \text{Sym}_n^*(\mathbb{Z})$  is taken of the form

$$T = \left[ \begin{array}{c|c} T' & \\ \hline & 0_{n-r} \end{array} \right]$$

for some nondegenerate  $T' \in \text{Sym}_r^*(\mathbb{Z})$  with  $0 \leq r \leq n$ , then the  $T$ -th Fourier coefficient of  $(E_{\kappa, \chi}^{(n)})^*$  is taken of the following form:

$$A_{\kappa, \chi}^*(T) = 2^{[(r+1)/2] - [(n+1)/2]} \prod_{i=[r/2]+1}^{[n/2]} L^{\{p\}}(1 - 2\kappa + 2i, \chi^2) \\ \times \begin{cases} L^{\{p\}}(1 - \kappa + r/2, \left(\frac{\partial_{T'}}{*}\right) \chi) \prod_{\substack{l|f_{T'} \\ l \neq p}} F_l(T'; \chi(l)l^{\kappa-r-1}) & \text{if } r \text{ is even,} \\ \prod_{\substack{l|\mathfrak{D}_{T'} \\ l \neq p}} F_l(T'; \chi(l)l^{\kappa-r-1}) & \text{if } r \text{ is odd.} \end{cases}$$

Therefore we have  $(E_{\kappa, \chi}^{(n)})^* \parallel_{\kappa} U_{p,n} = (E_{\kappa, \chi}^{(n)})^*$ .

The preceding Theorem 4.2 totally insists that the assignment  $E_{\kappa, \chi}^{(n)} \mapsto (E_{\kappa, \chi}^{(n)})^*$  can be regarded as a  $p$ -stabilization which generalizes the ordinary  $p$ -stabilization of  $E_{\kappa, \chi} = E_{\kappa, \chi}^{(1)}$  explained in §1. Whenever  $n = 2$ , Skinner-Urban [SU] has already dealt with a similar type of  $p$ -stabilization for some Siegel modular forms so that the associated eigenvalue of  $U_{p,2}$  is a  $p$ -adic unit. Accordingly, in the same context, we call  $(E_{\kappa, \chi}^{(n)})^*$  *semi-ordinary* at  $p$  if  $n \geq 2$ .

**Remark 4.3.** It should be mentioned that if  $n > 1$ ,  $(E_{\kappa, \chi}^{(n)})^*$  may *not* satisfy the ordinary condition at  $p$  in the sense of Hida (cf. [H2, H3]). This disparity is inevitable at least for Siegel Eisenstein series in general. For instance, in the case where  $M = 1$ , because of the shape of Satake parameters at  $p$ , there is no way to produce from  $E_{\kappa}^{(n)} = E_{\kappa, \text{triv}}^{(n)} \in \mathcal{M}_{\kappa}(\Gamma_0(1))^{(n)}$  to a Hecke eigenform of level  $p$  such that the associated eigenvalues of  $U_{p,1}, \dots, U_{p,n-1}$  and  $U_{p,n}$  are  $p$ -adic units simultaneously. However, as mentioned in [SU], it turns out that the semi-ordinary condition concerning only on the eigenvalue of  $U_{p,n}$  is sufficient to adapt Hida's ordinary theory with some modification. (See also [Pi, BPS].)

*Proof of Theorem 4.2.* The assertion (I) is obvious from Lemma 2.1 (I) and the properties of  $U_{p,n}$  explained in §3. Since  $U_{p,n}$  does not effect on the Fourier coefficient  $A_{\kappa, \chi}(0_n)$ , the assertion (II) for  $r = 0$  follows immediately from Lemma 2.1 (IIa), Equations (8) and (9). Hereinafter, we suppose that  $r > 0$ . It follows by the definition of  $\tilde{\mathcal{R}}_p^{(n)}(X, Y)$  (cf. Equation (7)) that

$$\tilde{\mathcal{R}}_p^{(n)}(X, Y) = \sum_{m=0}^n (-1)^m s_m \left( \{p^{j(2n-j+1)/2} X^j \mid 1 \leq j \leq n\} \right) Y^{n-m},$$

where  $s_m(\{X_1, \dots, X_n\})$  denotes the  $m$ -th elementary symmetric polynomial in  $X_1, \dots, X_n$ . Thus, by Lemma 2.1 (IIb) and Equation (9), we have

$$\begin{aligned} A_{\kappa, \chi}^*(T) &= \frac{\mathcal{P}_p^{(n)}(\chi(p) p^{\kappa-n-1}, 1)}{\mathcal{R}_p^{(n)}(\chi(p) p^{\kappa-n-1}, 1)} \cdot 2^{[(r+1)/2] - [(n+1)/2]} \prod_{i=[r/2]+1}^{[n/2]} L(1 - 2\kappa + 2i, \chi^2) \\ &\times \sum_{m=0}^n (-1)^m s_m \left( \{\chi^j(p) p^{j(2n-j+1)/2+j(\kappa-n-1)} \mid 1 \leq j \leq n\} \right) F_p(p^{n-m} T'; \chi(p) p^{\kappa-r-1}) \\ &\times \begin{cases} L(1 - \kappa + r/2, \binom{\mathfrak{D}_{T'}}{*}) \chi \prod_{\substack{l \mid \mathfrak{f}_{T'} \\ l \neq p}} F_l(T'; \chi(l) l^{\kappa-r-1}) & \text{if } r \text{ is even,} \\ \prod_{\substack{l \mid \mathfrak{D}_{T'} \\ l \neq p}} F_l(T'; \chi(l) l^{\kappa-r-1}) & \text{if } r \text{ is odd.} \end{cases} \end{aligned}$$

Here we note that

$$\begin{cases} \mathcal{R}_p^{(n)}(X, Y) = \mathcal{R}_p^{(r)}(p^{n-r} X, Y) \prod_{j=r+1}^n (1 - p^{j(2n-j+1)/2} X^j Y), \\ \mathcal{P}_p^{(n)}(X, 1) = \mathcal{P}_p^{(r)}(p^{n-r} X, 1) \prod_{i=[r/2]+1}^{[n/2]} (1 - p^{2n-2i+1} X^2). \end{cases}$$

Thus, to prove the assertion (II), it suffices to show that the following equation holds valid for each nondegenerate  $T' \in \text{Sym}_r^*(\mathbb{Z}_p)$  with  $0 < r \leq n$ :

$$\begin{aligned} (10) \quad &\sum_{m=0}^r (-1)^m s_m \left( \{p^{j(2r-j+1)/2} X^j \mid 1 \leq j \leq r\} \right) F_p(p^{r-m} T'; X) \\ &= \frac{\mathcal{R}_p^{(r)}(X, 1)}{\mathcal{P}_p^{(r)}(X, 1)} \cdot \begin{cases} (1 - \xi_p((-1)^{r/2} \det T') p^{r/2} X) & \text{if } r \text{ is even,} \\ 1 & \text{if } r \text{ is odd.} \end{cases} \end{aligned}$$

(Indeed, the preceding equation (10) yields

$$\begin{aligned} & \frac{\mathcal{P}_p^{(n)}(X, 1)}{\mathcal{R}_p^{(n)}(X, 1)} \sum_{m=0}^n (-1)^m s_m \left( \{p^{j(2n-j+1)/2} X^j \mid 1 \leq j \leq n\} \right) F_p(p^{n-m} T'; p^{n-r} X) \\ &= \prod_{i=[r/2]+1}^{[n/2]} (1 - p^{2n-2i+1} X^2) \times \begin{cases} (1 - \xi_p((-1)^{r/2} \det T') p^{n-r/2} X) & \text{if } r \text{ is even,} \\ 1 & \text{if } r \text{ is odd,} \end{cases} \end{aligned}$$

and hence, by evaluating this at  $X = p^{\kappa-n-1}$ , we obtain the desired equation. ) On the other hand, Theorem 1 in [Ki2] (resp. Theorem 6 in [BS]) states that for each nondegenerate  $T' \in \text{Sym}_r^*(\mathbb{Z}_p)$ , the equation

$$(11) \quad \sum_{m=0}^{\infty} F_p(p^m T'; X) Y^m = \frac{\mathcal{S}_p(T'; X, Y)}{(1-Y)\mathcal{R}_p^{(r)}(X, Y)}$$

holds for some polynomial  $\mathcal{S}_p(T'; X, Y) \in \mathbb{Z}[X, Y]$  if  $p \neq 2$  (resp.  $p = 2$ ). Since the preceding equation yields

$$\sum_{m=0}^r (-1)^m s_m \left( \{p^{j(2r-j+1)/2} X^j \mid 1 \leq j \leq r\} \right) F_p(p^{r-m} T'; X) = \mathcal{S}_p(T'; X, 1),$$

we may interpret Equation (10) as

$$(12) \quad \mathcal{S}_p(T'; X, 1) = \frac{\mathcal{R}_p^{(r)}(X, 1)}{\mathcal{P}_p^{(r)}(X, 1)} \times \begin{cases} (1 - \xi_p((-1)^{r/2} \det T') p^{r/2} X) & \text{if } r \text{ is even,} \\ 1 & \text{if } r \text{ is odd.} \end{cases}$$

Whenever  $r = 1$ , we easily see that  $F_p(t; X) = \sum_{i=0}^{\text{val}_p(t)} (pX)^i$  for each  $t \in \mathbb{Z}_p \setminus \{0\}$ . Thus, we have

$$F_p(pt; X) - pX F_p(t; X) = 1 = \frac{\mathcal{R}_p^{(1)}(X, 1)}{\mathcal{P}_p^{(1)}(X, 1)} \quad (\text{cf. Remark 4.1}).$$

Whenever  $r > 1$ , for a given  $T$ , let  $\mathfrak{i}(T)$  denote the least integer  $m$  such that  $p^m T^{-1} \in \text{Sym}_r^*(\mathbb{Z}_p)$ . It is known that if  $r = 2$ , then for each nondegenerate  $T \in \text{Sym}_2^*(\mathbb{Z}_p)$ , the polynomial  $F_p(T; X)$  admits the explicit form

$$F_p(T; X) = \sum_{i=0}^{\mathfrak{i}(T)} (p^2 X)^i \left\{ \sum_{j=0}^{\text{val}_p(f_T)-i} (p^3 X^2)^j - \xi_p(-\det T) pX \sum_{j=0}^{\text{val}_p(f_T)-i-1} (p^3 X^2)^j \right\}$$

(cf. [K1]). Thus a simple calculation yields that

$$\begin{aligned} & F_p(p^2 T; X) - (p^2 X + p^3 X^2) F_p(pT; X) + p^5 X^3 F_p(T; X) \\ &= 1 - \xi_p(-\det T) pX = \frac{\mathcal{R}_p^{(2)}(X, 1)}{\mathcal{P}_p^{(2)}(X, 1)} \times (1 - \xi_p(-\det T) pX), \end{aligned}$$

and hence, Equation (12) also holds for  $r = 2$ . Now, we suppose that  $r > 2$ . We note that every nondegenerate  $T \in \text{Sym}_r^*(\mathbb{Z}_p)$  is equivalent, over  $\mathbb{Z}_p$ , to a canonical form

$$T = \left[ \begin{array}{c|c} T_1 & \\ \hline & T_2 \end{array} \right]$$

for some  $T_1 \in \text{Sym}_2^*(\mathbb{Z}_p)$  and  $T_2 \in \text{Sym}_{r-2}^*(\mathbb{Z}_p) \cap \text{GL}(r-2, \mathbb{Q}_p)$ . It follows from Theorems 4.1 and 4.2 in [K1] that

$$\frac{\mathcal{S}_p(T; X, 1)}{1 - \xi_p((-1)^{r/2} \det T) p^{r/2} X} = \frac{\mathcal{S}_p(T_2; p^2 X, 1)}{1 - \xi_p((-1)^{r/2-1} \det T_2) p^{r/2+1} X} \times \frac{(1 - p^{(r-1)(r+2)/2} X^{r-1})(1 - p^{r(r+1)/2} X^r)}{1 - p^{r+1} X^2}$$

if  $r$  is even, and

$$\mathcal{S}_p(T; X, 1) = \mathcal{S}_p(T_2; p^2 X, 1) \cdot \frac{(1 - p^{(r-1)(r+2)/2} X^{r-1})(1 - p^{r(r+1)/2} X^r)}{1 - p^{r+2} X^2}$$

if  $r$  is odd. (See also [K2, §3]) Thus, it is proved by induction on  $r$  that Equation (12) (and thus, (10)) holds in general. This completes the proof.  $\square$

As a straightforward conclusion of Theorem 4.2 above, we have

**Theorem 4.4.** *Let  $\chi$  be a Dirichlet character modulo  $M$  as above and  $\omega^a$  a power of the Teichmüller character with  $0 \leq a < \varphi(\mathfrak{p})$ <sup>6</sup>, respectively. For each  $n \geq 1$ , there exists a formal Fourier expansion*

$$\mathcal{E}_{\chi\omega^a}^{(n)}(X) = \sum_{\substack{T \in \text{Sym}_n^*(\mathbb{Z}), \\ T \geq 0}} \mathcal{A}_{\chi\omega^a}(T; X) \mathbf{q}^T \in \mathcal{L}[[\mathbf{q}]]^{(n)},$$

where  $\mathcal{L}$  is the field of fractions of  $\Lambda = \mathbb{Z}_p[\chi][[X]]$ , such that for each positive integer  $\kappa > n+1$  with  $\chi(-1) = (-1)^\kappa$  and  $\kappa \equiv a \pmod{\varphi(\mathfrak{p})}$ , we have

$$\mathcal{E}_{\chi\omega^a}^{(n)}((1 + \mathfrak{p})^\kappa - 1) = (E_{\kappa, \chi}^{(n)})^* \in \mathcal{M}_\kappa(\Gamma_0(Mp), \chi)^{(n)}.$$

Moreover, put

$$\begin{aligned} \mathcal{B}^{(n)}(X) &:= \prod_{i=1}^{[n/2]} \{(1 + \mathfrak{p})^{-2i} (1 + X)^2 - 1\} \prod_{j=0}^{[n/2]} \{(1 + \mathfrak{p})^{-j} (1 + X) - 1\} \\ &= X \prod_{i=1}^{[n/2]} \{(1 + \mathfrak{p})^{-i} (1 + X) - 1\}^2 \cdot \{(1 + \mathfrak{p})^{-i} (1 + X) + 1\}. \end{aligned}$$

Then  $\overline{\mathcal{E}}_{\chi\omega^a}^{(n)}(X) := \mathcal{B}^{(n)}(X) \cdot \mathcal{E}_{\chi\omega^a}^{(n)}(X)$  belongs to  $\Lambda[[\mathbf{q}]]^{(n)}$ .

*Proof.* As is well-known by Deligne-Ribet [DR] (generalizing the previous work of Kubota-Leopoldt), associated to a quadratic Dirichlet character  $\xi$ , a Dirichlet character  $\chi$  taken as above, and a power of the Teichmüller character  $\omega^a$  with  $0 \leq a < \varphi(\mathfrak{p})$ , there exists  $\Phi(\xi\chi\omega^a; X) \in \Lambda$  such that for each positive integer  $k > 1$ , we have

$$\Phi(\xi\chi\omega^a; (1 + \mathfrak{p})^k - 1) = \begin{cases} ((1 + \mathfrak{p})^k - 1) \cdot L^{\{p\}}(1 - k, \omega^{-k}) & \text{if } \xi\chi\omega^a \text{ is trivial,} \\ L^{\{p\}}(1 - k, \xi\chi\omega^{a-k}) & \text{otherwise.} \end{cases}$$

Accordingly, put

$$\Psi(\xi\chi\omega^a; X) := \begin{cases} X^{-1} \Phi(\xi\chi\omega^a; X) & \text{if } \xi\chi\omega^a \text{ is trivial,} \\ \Phi(\xi\chi\omega^a; X) & \text{otherwise.} \end{cases}$$

We easily see that  $\Psi(\xi\chi\omega^a; (1 + \mathfrak{p})^k - 1) = L^{\{p\}}(1 - k, \xi\chi\omega^{a-k})$  for each  $k > 1$ . More generally, it turns out that if  $\varepsilon : 1 + \mathfrak{p}\mathbb{Z}_p \rightarrow \overline{\mathbb{Q}_p}^\times$  is a character of finite order, then

$$(13) \quad \Psi(\xi\chi\omega^a; \varepsilon(1 + \mathfrak{p})(1 + \mathfrak{p})^k - 1) = L^{\{p\}}(1 - k, \xi\chi\omega^{a-k}\varepsilon)$$

<sup>6</sup>Here  $\varphi$  denotes Euler's totient function.

for any  $k > 1$ . On the other hand, for each  $x \in 1 + \mathfrak{p}\mathbb{Z}_p$ , put  $s(x) := \log_p(x)/\log_p(1 + \mathfrak{p})$ , where  $\log_p$  is the  $p$ -adic logarithm function in the sense of Iwasawa, and thus we have  $s : 1 + \mathfrak{p}\mathbb{Z}_p \xrightarrow{\sim} \mathbb{Z}_p$ .

Then for each  $T = \left[ \begin{array}{c|c} T' & \\ \hline & 0_{n-r} \end{array} \right] \in \text{Sym}_n^*(\mathbb{Z})$  with  $T' \in \text{Sym}_r^*(\mathbb{Z}) \cap \text{GL}(r, \mathbb{Q})$  and  $0 \leq r \leq n$ , we define  $\mathcal{A}_{\chi\omega^a}(T; X) \in \mathcal{L}$  as follows:

$$(14) \quad \mathcal{A}_{\chi\omega^a}(T; X) = 2^{[(r+1)/2]-[(n+1)/2]} \prod_{i=[r/2]+1}^{[n/2]} \Psi(\chi\omega^{2a-2i}; (1 + \mathfrak{p})^{-2i}(1 + X)^2 - 1) \\ \times \begin{cases} \Psi\left(\left(\frac{\mathfrak{d}_{T'}}{*}\right) \chi\omega^{a-r/2}; (1 + \mathfrak{p})^{-r/2}(1 + X) - 1\right) \\ \quad \times \prod_{\substack{l|T', \\ l \neq p}} F_l(T'; \chi\omega^a(l) l^{-r-1}(1 + X)^{s((l))}) & \text{if } r \text{ is even,} \\ \prod_{\substack{l|\mathfrak{D}_{T'}, \\ l \neq p}} F_l(T'; \chi\omega^a(l) l^{-r-1}(1 + X)^{s((l))}) & \text{if } r \text{ is odd.} \end{cases}$$

It follows from Theorem 4.2 (II) that if  $\kappa > n + 1$  and further if  $\kappa \equiv a \pmod{\varphi(\mathfrak{p})}$  (or equivalently,  $\omega^{a-\kappa}$  is trivial), then

$$\mathcal{A}_{\chi\omega^a}(T; (1 + \mathfrak{p})^\kappa - 1) = A_{\kappa, \chi}^*(T).$$

Thus we obtain the former assertion. Moreover, it follows directly from Equation (14) that  $\mathcal{B}^{(n)}(X) \cdot \mathcal{A}_{\chi\omega^a}(T; X) \in \Lambda$  for all  $T$ . Therefore  $\overline{\mathcal{E}}_{\chi\omega^a}^{(n)}(X) \in \Lambda[[\mathfrak{q}]]^{(n)}$ . This completes the proof.  $\square$

## 5. SEMI-ORDINARY $\Lambda$ -ADIC SIEGEL EISENSTEIN SERIES

In this section, we fix an *odd* prime number  $p$ . We show that for a Dirichlet character  $\chi$  modulo a positive odd integer  $M$  with  $p \nmid M$  and an integer  $a$  with  $0 \leq a < p - 1$ , the formal Fourier expansion  $\mathcal{E}_{\chi\omega^a}^{(n)}(X)$  (resp.  $\overline{\mathcal{E}}_{\chi\omega^a}^{(n)}(X)$ ) with coefficients in  $\mathcal{L} = \mathbb{Q}_p(\chi)[[X]]$  (resp.  $\Lambda = \mathbb{Z}_p[\chi][[X]]$ ) defined in Theorem 4.4, give rise to classical Siegel modular forms via the specialization at  $X = \varepsilon(1 + p)(1 + p)^\kappa - 1$ , where  $\kappa$  is a positive integer sufficiently large and  $\varepsilon : 1 + p\mathbb{Z}_p \rightarrow \overline{\mathbb{Q}}_p^\times$  is a character of finite order.

First, from Lemma 2.1 (II), we deduce the following:

**Theorem 5.1.** *If  $\kappa$  is a positive integer with  $\kappa > n + 1$  and further if  $\varepsilon : 1 + p\mathbb{Z}_p \rightarrow \overline{\mathbb{Q}}_p^\times$  is a character having exact order  $p^m$  for some nonnegative integer  $m$ , then*

$$\mathcal{E}_{\chi\omega^a}^{(n)}(\varepsilon(1 + p)(1 + p)^\kappa - 1) = E_{\kappa, \chi\omega^{a-\kappa}\varepsilon}^{(n)} \in \mathcal{M}_\kappa(\Gamma_0(Mp^{m+1}), \chi\omega^{a-\kappa}\varepsilon)^{(n)}$$

as long as  $\omega^{2a-2\kappa}\varepsilon^2$  is non-trivial.

*Proof.* For each  $T = \left[ \begin{array}{c|c} T' & \\ \hline & 0_{n-r} \end{array} \right] \in \text{Sym}_n^*(\mathbb{Z})$  with  $T' \in \text{Sym}_r^*(\mathbb{Z}) \cap \text{GL}(r, \mathbb{Q})$  and  $0 \leq r \leq n$ , the equation (14) is specialized as

$$\mathcal{A}_{\chi\omega^a}(T; \varepsilon(1 + p)(1 + p)^\kappa - 1) = 2^{[(r+1)/2]-[(n+1)/2]} \prod_{i=[r/2]+1}^{[n/2]} L^{\{p\}}(1 - 2\kappa + 2i, \chi\omega^{2a-2\kappa}\varepsilon^2)$$

$$\times \begin{cases} L\{p\}(1 - \kappa + r/2, \left(\frac{\partial_{T'}}{*}\right) \chi\omega^{a-\kappa}\varepsilon) \\ \quad \times \prod_{\substack{l|f_{T'}, \\ l \neq p}} F_l(T'; \chi\omega^{a-\kappa}\varepsilon(l) l^{\kappa-r-1}) & \text{if } r \text{ is even,} \\ \prod_{\substack{l|\mathfrak{D}_{T'}, \\ l \neq p}} F_l(T'; \chi\omega^{a-\kappa}\varepsilon(l) l^{\kappa-r-1}) & \text{if } r \text{ is odd.} \end{cases}$$

Thus, if  $\omega^{2a-2\kappa}\varepsilon^2$  is non-trivial, then Lemma 2.1 (II) yields

$$\mathcal{A}_{\chi\omega^a}(T; \varepsilon(1+p)(1+p)^\kappa - 1) = A_{\kappa, \chi\omega^{a-\kappa}\varepsilon}(T).$$

This completes the proof.  $\square$

On the other hand, by exploiting the vertical control theorem for  $p$ -adic Siegel modular forms in the sense of Hida [H2], we may deduce a slight weaker version of the preceding theorem as follows:

**Theorem 5.2.** *If  $\kappa$  is a positive integer with  $\kappa > n(n+1)/2$ , and further if  $\varepsilon : 1 + p\mathbb{Z}_p \rightarrow \overline{\mathbb{Q}}_p^\times$  is a character having exact order  $p^m$  for some nonnegative integer  $m$ , then*

$$\mathcal{E}_{\chi\omega^a}^{(n)}(\varepsilon(1+p)(1+p)^\kappa - 1) \in \mathcal{M}_\kappa(\Gamma_0(Mp^{m+1}), \chi\omega^{a-\kappa}\varepsilon)^{(n)}$$

as long as  $\omega^{a-\kappa}\varepsilon$  is non-trivial.

*Proof.* It follows from Theorem 4.4 that  $\mathcal{E}_{\chi\omega^a}^{(n)}((1+p)^{\kappa'} - 1) \in \mathcal{M}_{\kappa'}(\Gamma_0(Mp), \chi)^{(n)}$  for infinitely many integers  $\kappa'$  with  $\kappa' > n+1$  and  $\kappa' \equiv a \pmod{p-1}$ . Thus, Théorème 1.1 in [Pi] (generalizing [H2] in more general settings) turns out that if an integer  $\kappa$  is sufficiently large, then the specialization of  $\mathcal{E}_{\chi\omega^a}^{(n)}(X)$  at  $X = \varepsilon(1+p)(1+p)^\kappa - 1$  gives rise to an overconvergent  $p$ -adic Siegel modular form of weight  $\kappa$ , tame level  $M$  and Iwahori level  $p^{m+1}$  with character  $\chi\omega^{a-\kappa}\varepsilon$  (for the precise definition, see [SU, Pi]). In addition, the explicit formula for Fourier coefficients of  $\mathcal{E}_{\chi\omega^a}^{(n)}(\varepsilon(1+p)(1+p)^\kappa - 1)$  also yields that it is an eigenform of  $U_{p,n}$  whose eigenvalue is 1. Therefore, if  $\kappa$  satisfies the condition  $\kappa > n(n+1)/2$  (cf. Hypothèse 4.5.1 in [BPS]), then the above-mentioned overconvergent form is indeed classical, that is,

$$\mathcal{E}_{\chi\omega^a}^{(n)}(\varepsilon(1+p)(1+p)^\kappa - 1) \in \mathcal{M}_\kappa(\Gamma_0(Mp^{m+1}), \chi\omega^{a-\kappa}\varepsilon)^{(n)}$$

(cf. Théorème 5.3.1 in [ibid]). Thus we obtain the assertion.  $\square$

**Remark 5.3.** On the contrary to Theorem 4.4, we may not deduce similar statements to Theorems 5.1 and 5.2 in the case where  $p = 2$ .

Now, as a summary of Theorems 4.4, 5.1 and 5.2 above, we have the following statement which can be viewed as a generalization of Fact 1.1:

**Theorem 5.4.** *Let  $\chi$  be a Dirichlet character modulo  $M$  taken as above. For each integer  $a$  with  $0 \leq a < p-1$ , there exists a formal Fourier expansion  $\overline{\mathcal{E}}_{\chi\omega^a}^{(n)}(X) \in \Lambda[[\mathbf{q}]]^{(n)}$  such that if  $\kappa > n(n+1)/2$  and further if  $\varepsilon : 1 + p\mathbb{Z}_p \rightarrow \overline{\mathbb{Q}}_p^\times$  is a character having exact order  $p^m$  with some  $m \geq 0$ , then*

$$\overline{\mathcal{E}}_{\chi\omega^a}^{(n)}(\varepsilon(1+p)(1+p)^\kappa - 1) \in \mathcal{M}_\kappa(\Gamma_0(Mp^{m+1}), \chi\omega^{a-\kappa}\varepsilon)^{(n)}$$

is a Hecke eigenform whose eigenvalue of  $U_{p,n}$  is 1. In particular,

$$\overline{\mathcal{E}}_{\chi\omega^a}^{(n)}(\varepsilon(1+p)(1+p)^\kappa - 1) = C_{\kappa, \varepsilon}^{(n)} \times \begin{cases} (E_{\kappa, \chi}^{(n)})^* & \text{if } \omega^{a-\kappa}\varepsilon \text{ is trivial,} \\ E_{\kappa, \chi\omega^{a-\kappa}\varepsilon}^{(n)} & \text{if } \omega^{2a-2\kappa}\varepsilon^2 \text{ is non-trivial,} \end{cases}$$

where  $C_{\kappa, \varepsilon}^{(n)} = \mathcal{B}^{(n)}(\varepsilon(1+p)(1+p)^\kappa - 1)$ .

## REFERENCES

- [AZ] A.N. Andrianov and V.G. Zhuravlev, *Modular forms and Hecke operators*, Transl. of Math. Monogr., vol.145, Amer. Math. Soc., Providence, RI, 1995.
- [AL] A.O.L. Atkin and J. Lehner, *Hecke operators on  $\Gamma_0(m)$* , Math. Ann. **185** (1970), 134–160.
- [BPS] S. Bijakowski, V. Pilloni and B. Stroh, *Classicit  de formes modulaires surconvergentes*, Ann. of Math. **183** (2016), no.3, 975–1014.
- [B ] S. B cherer, *On the Hecke operator  $U(p)$ . With an appendix by Ralf Schmidt*, J. Math. Kyoto Univ. **45** (2005), no.4, 807–829.
- [BS] S. B cherer and F. Sato, *Rationality of certain formal power series related to local densities*, Comment. Math. Univ. St.Paul. **36** (1987), no.1, 53–86.
- [CP] M. Courtieu and A.A. Panchishkin, *Non-Archimedean L-Functions and Arithmetical Siegel Modular Forms*, Lecture Notes in Math., vol.1471, Springer-Verlag, Berlin, 2004.
- [DR] P. Deligne and K.A. Ribet, *Values of abelian L-functions at negative integers over totally real fields*, Invent. Math. **59** (1980), 227–286.
- [Fe] P. Feit, *Poles and residues of Eisenstein series for symplectic and unitary groups*, Mem. Amer. Math. Soc. **61**, no.346, 1986.
- [Fr] E. Freitag, *Siegelsche Modulfunktionen*, Grundlehren Math. Wiss., vol.254, Springer-Verlag, Berlin, 1983.
- [H1] H. Hida, *Elementary theory of L-functions and Eisenstein series*, London Math. Soc. Student Text, vol.26, Cambridge University Press, Cambridge, 1993.
- [H2] H. Hida, *Control theorems for coherent sheaves on Shimura varieties of PEL-type*, J. Inst. Math. Jussieu **1** (2002), no.1, 1–76.
- [H3] H. Hida, *p-adic automorphic forms on Shimura varieties*, Springer Monographs in Math., Springer-Verlag, New York, 2004.
- [Ik] T. Ikeda, *On the functional equation of the Siegel series*, J. Number Theory **172** (2017), 44–62.
- [K1] H. Katsurada, *An explicit formula for Siegel series*, Amer. J. Math. **121** (1999), no.2, 415–452.
- [K2] H. Katsurada, *Euler factor of a certain Dirichlet series attached to Siegel Eisenstein series*, Abh. Math. Sem. Univ. Hamburg **71** (2001), 81–90.
- [Ki1] Y. Kitaoka, *Dirichlet series in the theory of quadratic forms*, Nagoya Math. J. **95** (1984), 73–84.
- [Ki2] Y. Kitaoka, *Local densities of quadratic forms and Fourier coefficients of Eisenstein series*, Nagoya Math. J. **103** (1986), 149–160.
- [Mi] T. Miyake, *Modular forms*, Springer Monographs in Math., Springer-Verlag, Berlin, 2006.
- [Pa] A.A. Panchishkin, *On the Siegel-Eisenstein measure and its applications*, Israel J. Math. **120** (2000), part B, 467–509.
- [Pi] V. Pilloni, *Sur la th orie de Hida pour le groupe  $\mathrm{GSp}_{2g}$* , Bull. Soc. Math. France **140** (2012), no.3, 335–400.
- [Se] J.-P. Serre, *Formes modulaires et fonctions z ta p-adiques, Modular functions of one variable, Vol. III* (Proc. Internat. Summer School, Univ. Antwerp, 1972), 191–268, Lecture Notes in Math., vol.350, Springer-Verlag, Berlin, 1973.
- [S1] G. Shimura, *Confluent hypergeometric functions on tube domains*, Math. Ann. **260** (1982), no.3, 269–302.
- [S2] G. Shimura, *Euler products and Fourier coefficients of automorphic forms on symplectic groups*, Invent. Math. **116** (1994), no.1–3, 531–576.
- [SU] C. Skinner and E. Urban, *Sur les d formations p-adiques de certaines repr sentations automorphes*, J. Inst. Math. Jussieu **5** (2006), no.4, 626–698.
- [Sw] W.J. Sweet Jr., *A computation of the gamma matrix of a family of p-adic zeta integrals*, J. Number Theory **55** (1995), no.2, 222–260.
- [T1] S. Takemori, *p-adic Siegel-Eisenstein series of degree two*, J. Number Theory **132** (2012), no.6, 1203–1264.
- [T2] S. Takemori, *Siegel Eisenstein series of degree n and  $\Lambda$ -adic Eisenstein series*, J. Number Theory **149** (2015), 105–138.
- [Ta] R.L. Taylor, *Congruences between modular forms*, Ph.D thesis, Princeton University, 1988.
- [W] A. Wiles, *On ordinary  $\lambda$ -adic representations associated to modular forms*, Invent. Math. **94** (1988), no.3, 529–573.

DEPARTMENT OF MATHEMATICS, HIROSHIMA UNIVERSITY, 1-7-1, KAGAMIYAMA, HIGASHI-HIROSHIMA, JAPAN  
 E-mail address: hisa@hiroshima-u.ac.jp