

# GEOMETRIC PROPERTIES OF SOME TOTALLY ORDERED COMPACT SETS

MOHAMMAD DAHER AND KHALIL SAADI

ABSTRACT. In this paper, we show that there are a totally ordered compact  $K$  separable, a Hausdorff topology  $\tau'$  on  $C(K)$  and two closed subspaces  $Y_1, Y_2$  of  $(C(K), \tau_p)$  such that  $(C(K), \tau')$  is not universally measurable,  $(C(K), \tau_p) = (Y_1, \tau_p) \oplus (Y_2, \tau_p)$ ,  $(Y_1, \tau_p)$  is isomorphic to  $(Y_2, \tau_p)$ ,  $(Y_j, \tau_p) = (Y_j, \tau')$ ,  $j \in \{1, 2\}$  and  $Bor(C(K), \tau') \otimes Bor(C(K), \tau') \neq Bor(C(K)) \times C(K), \tau' \otimes \tau'$ , this is the main result of this work.

We start this work to construct totally ordered non metrisable compact sets  $K_\mu(E)$  from a reference set  $E$  which is totally ordered, and from a positive Borel measure on  $E$  satisfying some reasonable assumptions.

At first, we introduce a totally ordered set  $E$ , which is equipped with the order topology  $\tau_0$ , and a non-negative Borel measure on  $(E, \tau_0)$ . To these two elements we associate a compact set  $K = K_\mu(E)$ , obtained as spectrum of a certain  $C^*$ -subalgebra of  $L^\infty(E, \mu)$ . We impose some natural properties on  $E$  and  $\mu$  that make the compact  $K$  not metrisable.

Afterward, we introduce a certain function  $h_{\mu, E} = h$  defined on a subset of  $K$  with values in  $E$  or in  $\overline{E} = E \cup \{-\infty, +\infty\}$ . We study the surjectivity and the continuity of the function  $h$ ; this function plays an important role for understanding the geometric properties of  $K$ .

We will see that  $E$  is homeomorphic to  $K$ , When  $E$  is a separable connected strongly Lindelöf space and if  $\mu(I) > 0$  for every nonempty interval of  $E$ .

We study in detail a particular case when  $E$  is a locally compact totally ordered group satisfying the following conditions:

$$\forall a, b \in E \text{ such that } a < b, \exists c \in E \text{ verifying } a < c < b. \quad (*)$$

$$a \leq b \iff a - b \leq 0, \quad a, b \in E. \quad (**)$$

---

2020 *Mathematics Subject Classification.* 54H10, 54F05; Secondary 54C35.

*Key words and phrases.* Totally ordered compact set, Lindelöf space, Borel set and mapping, Rosenthal compact set.

Our main result (theorem 6) will be shown, for this case and we will see that the compact  $K$  ( $K$  is a Rosenthal compact set) has the following property:

$Bor(C(K), \tau_p) \otimes Bor(C(K), \tau_p) = Bor(C(K)) \times C(K), \tau_p \otimes \tau_p)$  (with the continuous hypothesis) but  $Bor(K) \otimes Bor(K) \neq Bor(K \times K)$ .

Recall that if  $L$  is a metrisable compact set, then  $Bor(L) \otimes Bor(L) = Bor(L \times L)$ .

We will see in part 6 (lemma 14) that if  $L$  is a Rosenthal compact set, then  $(C(L), \tau_p)$  is universally measurable. In theorem 6 the Hausdorff topology  $\tau'$  on  $C(K)$  satisfying  $(Y_j, \tau_p) = (Y_j, \tau')$ ,  $j \in \{1, 2\}$  and  $(C(K), \tau_p) = (Y_1, \tau_p) \oplus (Y_2, \tau_p)$ ,  $(Y_1, \tau_p)$  but  $(C(K), \tau')$  is not universally measurable (here  $Y_1$  and  $Y_2$  are two closed subspaces of  $(C(K), \tau_p)$ ).

Let  $(E, \leq)$  be a totally ordered set. The order topology  $\tau_0$  on  $E$  is the topology generated by the intervals of the form  $\{x \in E; x < a\}$  or of the form  $\{x \in E; x > b\}$ ,  $a, b \in E$ . This topology is Hausdorff (if  $x < a < y$  in  $E$ ,  $V_x = \{t \in E; t < a\}$  and  $V_y = \{t \in E; t > a\}$  are disjoint neighborhoods of  $x$  and  $y$ ; if there is no  $a$  between  $x$  and  $y$ ,  $V_x = \{t \in E; t < y\}$  and  $V_y = \{t \in E; t > x\}$  do the job).

We denote by  $\tau_1$  (resp.  $\tau_2$ ) the topology on  $E$  generated by the intervals of the form  $\{x \in E; x > a\}$  or  $\{x \in E; x \leq b\}$  (resp. of the form  $\{x \in E; x < a\}$  or  $\{x \in E; x \geq b\}$ ),  $a, b \in E$ .

Let  $(X, \tau)$  be a topological space and  $X_1$  a subset of  $X$ .  $(X_1, \tau)$  (since unless is said otherwise) denotes the topological space formed by the relative open subsets (a relative open subset of  $X_1$  is of the form  $X_1 \cap V$ , where  $V$  is an open subset of  $(X, \tau)$ ).

## 1. CONSTRUCTION OF TOTALLY ORDERED COMPACT SETS

In this part we construct totally ordered compact sets  $K = K_\mu(E)$  non metrisables, in passing, we introduce the function  $h$ , this function help us to know the topological structure of  $K$ .

Let  $(E, \leq)$  be a totally ordered set. If  $E$  admits a maximum (resp. a minimum) we denote

$$+\infty = \max\{x \in E; x \in E\} \text{ (resp. } -\infty = \min\{x \in E; x \in E\}).$$

Let  $\mu$  be a positive Borel measure on  $(E, \tau_0)$ . One supposes that the measure  $\mu$  verifies the following condition:

$$\mu(\{u \in E; u > t\}) > 0, \quad \forall t \in E \setminus \{+\infty\}. \quad (1.1)$$

Let  $t \in E$  and let  $f^t$  be the characteristic function of  $\{u \in E; u > t\}$ . Denote by  $A$  the  $C^*$ -subalgebra of  $L^\infty(\mu)$  (with unit) generated by the

family  $\{f^t, t \in E\}$  (the scalars are complex scalars). Let  $K(E)$  be the set of characters on  $A$ . By [9], [24],  $K_\mu(E)$  is  $\sigma(A^*, A)$  compact and  $A = C(K_\mu(E))$ .

As  $f^s f^t = f^{s \vee t}$  when  $s, t \in E$ , the family  $(f^t)_{t \in E}$  is stable under product, and  $A$  is simply the closure in  $L^\infty(\mu)$  of the vector space generated by the constant function  $\mathbf{1}$  (the unit  $e$  of  $A$ ) and the family  $(f^t)_{t \in E}$ . As a consequence, a character  $\theta \in K$  is fully determined by its values on the  $f^t$ s: if  $\theta(f^t) = \eta(f^t)$  for all  $t \in E$ , then  $\theta = \eta$ .

If  $\theta \in K_\mu(E)$  is a character and  $\chi_F$  the characteristic function of a subset  $F \subset E$ , then  $\theta(\chi_F) = 0$  or  $1$  because  $(\chi_F)^2 = \chi_F$ . In particular,  $\theta(f^t) \in \{0, 1\}$  and furthermore, when  $s < t$  in  $E$ , we have  $\theta(f^s) \geq \theta(f^t)$  since  $f^s = f^t + \chi_{(s,t]}$ .

For every  $t \in E$  let

$$[t] = \left\{ t' \in E; \|f^t - f^{t'}\|_{L^\infty(E, \mu)} = 0 \right\} = \{t' \in E; \mu([t \wedge t', t \vee t']) = 0\}.$$

In the following, we suppose that

$$\{[t]; t \in E\} \text{ is uncountable.} \quad (1.2)$$

Since  $\|f^t - f^{t'}\|_{L^\infty(E, \mu)} \geq 1$  for  $[t] \neq [t']$ ,  $C(K_\mu(E)) = A$  is not a separable space, hence  $K_\mu(E)$  is not metrisable.

We define the function  $\psi_{\mu, E} : K_\mu(E) \times E \rightarrow \{0, 1\}$  by

$$\psi_{\mu, E}(\theta, t) = \theta(f^t), \quad \theta \in K, t \in E.$$

When we have that  $-\infty, +\infty \in E$ , the function  $\psi_{\mu, E}$  is defined on  $K_\mu(E) \times [E \setminus \{-\infty, +\infty\}]$ , with values in  $\{0, 1\}$  as was said above.

Let  $B_1^{\mu, E}$  be the subset of all characters  $\theta \in K$  such that the map  $(E, \tau_0) \ni t \mapsto \psi_{\mu, E}(\theta, t)$  is continuous; this means that the two sets  $\{t \in E; \psi_{\mu, E}(\theta, t) = \varepsilon\}$ , for  $\varepsilon = 0, 1$ , are both closed (and both open).

Let now  $\theta \in K_\mu(E) \setminus B_1^{\mu, E}$ ; there are two types of discontinuities.

*Case 1*,  $\{t \in E; \psi_{\mu, E}(\theta, t) = 0\}$  is not closed. Let  $h_{\mu, E}(\theta)$  be a point in the closure of  $\{t \in E; \psi_{\mu, E}(\theta, t) = 0\}$  in  $E$  such that  $\psi_{\mu, E}(\theta, h_{\mu, E}(\theta)) = 1$ .

**Lemma 1.** *The point  $h_{\mu, E}(\theta)$  is unique.*

*Proof.* Suppose that there exists  $h'(\theta) \in E$  in the closure of the set  $E_0(\theta) = \{t \in E; \psi_{\mu, E}(\theta, t) = 0\}$ , such that  $\psi_{\mu, E}(\theta, h'(\theta)) = 1$  and  $h_{\mu, E}(\theta) < h'(\theta)$ . The set  $\{u \in E; u < h'(\theta)\}$  is then an open neighborhood of  $h_{\mu, E}(\theta)$ , but  $h_{\mu, E}(\theta)$  is in the closure of  $E_0(\theta)$ , hence there is  $u_1 \in E_0(\theta)$  such that  $u_1 < h'(\theta)$ . This is impossible, because the map  $t \mapsto \psi_{\mu, E}(\theta, t)$  is a non-increasing function. Thus  $h_{\mu, E}(\theta)$  is unique. ■

*Case 2*,  $\{t \in E; \psi_{\mu,E}(\theta, t) = 1\}$  is not closed. We let now  $h_{\mu,E}(\theta)$  be the unique point in the closure of  $\{t \in E; \psi_{\mu,E}(\theta, t) = 1\}$  in  $E$  such that  $\psi_{\mu,E}(\theta, h_{\mu,E}(\theta)) = 0$ .

Let  $B_2^{\mu,E} = \{\theta \in K_\mu(E) \setminus B_1^{\mu,E}; \{t \in E; \psi_{\mu,E}(\theta, t) = 0\} \text{ is not closed}\}$  and let  $B_3^{\mu,E} = \{\theta \in K_\mu(E) \setminus B_1^{\mu,E}; \{t \in E; \psi_{\mu,E}(\theta, t) = 1\} \text{ is not closed}\}$ .

When the space  $E$  is fixed, we write  $K = K_\mu(E)$ ,  $h_{\mu,E} = h$ ,  $\psi = \psi_{\mu,E}$  and  $B_j^{\mu,E} = B_j$ ,  $j \in \{1, 2, 3\}$ .

**Lemma 2.** *For every  $\theta \in B_2 \cup B_3$ , we have*

$$h(\theta) = \sup\{t \in E; \psi(\theta, t) = 1\}. \quad (1.3)$$

*Proof.* Remember that the map  $t \mapsto \psi(\theta, t)$  is a non-increasing function; let  $E_j(\theta) = \{t \in E; \psi(\theta, t) = j\}$ , for  $j = 0, 1$ .

If  $\theta \in B_2$ , we have  $\psi(\theta, h(\theta)) = 1$  while  $h(\theta)$  is in the closure of  $E_0(\theta)$ ; when  $v > h(\theta)$ , the set  $\{u \in E; u < v\}$  is a neighborhood of  $h(\theta)$ , hence there is  $u_1 \in E$  such that  $u_1 < v$  and  $\psi(\theta, u_1) = 0$ , thus  $\psi(\theta, v) = 0$ : when  $\theta \in B_2$ , the point  $h(\theta)$  is actually the largest element of  $E_1(\theta)$ .

If  $\theta \in B_3$ , then  $\psi(\theta, h(\theta)) = 0$ ; if  $\psi(\theta, v) = 1$ , we have  $v < h(\theta)$ , showing already that  $h(\theta)$  is an upper bound of  $E_1(\theta)$ . If now we have  $a < h(\theta)$ , then  $V = \{t \in E; t > a\}$  is a neighborhood of  $h(\theta)$ , implying that there is  $v_1 \in E$  such that  $v_1 > a$  and  $\psi(\theta, v_1) = 1$ . This shows that  $a < h(\theta)$  cannot be an upper bound of  $E_1(\theta)$ . ■

By an argument similar to that of Lemma 2, we see that:

**Lemma 3.** *For every  $\theta \in B_2 \cup B_3$  :*

$$h(\theta) = \inf\{t \in E; \psi(\theta, t) = 0\}.$$

Note that  $B_2 \cap B_3 = \emptyset$ ,  $B_2 = \{\theta \in K \setminus B_1; \psi(\theta, h(\theta)) = 1\}$  and  $B_3 = \{\theta \in K \setminus B_1; \psi(\theta, h(\theta)) = 0\}$ .

**Remark 1.** *Assume that  $(E, \tau_0)$  is separable and that  $(E, \leq)$  satisfies the condition (\*). Then  $(E, \tau_0)$  has a countable basis.*

*Proof.* Indeed, let  $(a_n)_{n \in \mathbb{N}}$  be a dense sequence in  $(E, \tau_0)$ . For every  $m, n \in \mathbb{N}$ , consider  $V_{m,n} = \{x \in E; a_n > x > a_m\}$  and let  $M = \{(m, n) \in \mathbb{N}^2; V_{m,n} \neq \emptyset\}$ . It is easy to see that  $(V_{m,n})_{(m,n) \in M}$  forms a basis of  $(E, \tau_0)$ . ■

**Example 1.** *Let  $E$  be a separable uncountable abelian locally compact totally ordered group [12]. Assume that  $E$  satisfies the condition (\*) and that the order topology coincides with the topology of  $E$ . Choose  $\mu = m$  the Haar measure on  $E$ . We observe that every non empty open*

set in  $E$  is of strictly positive measure, and since  $\mu(\{t\}) = 0$  for every  $t \in E$ , the conditions (1.1) and (1.2) are satisfied. We shall study this example in detail in the last part.

**Example 2.** Let  $E = [0, 1]$  and  $\mu = m$  the Lebesgue measure, in corollary 12 we will see that  $K = K_m(E)$  is homeomorphic to the two-arrows space  $[0, 1] \times \{0, 1\}$ .

**Proposition 1.** The compact set  $K$  is totally ordered (i.e., the topology on  $K$  is defined by a total order relation).

*Proof.* We define on  $K$  the following relation, for all  $\theta_1, \theta_2 \in K$ :

$$\theta_1 \leq \theta_2 \Leftrightarrow \psi(\theta_1, t) = \theta_1(f^t) \leq \theta_2(f^t) = \psi(\theta_2, t), \quad \forall t \in E.$$

Let us show that  $(K, \leq)$  is totally ordered. Pick  $\theta_1, \theta_2 \in K$ ,  $\theta_1 \neq \theta_2$ . We may assume that there exists  $t_0 \in E$  such that  $\psi(\theta_1, t_0) = 0$  and  $\psi(\theta_2, t_0) = 1$ . Let  $t \in E$ . If  $t < t_0$ ,  $\psi(\theta_2, t) = 1$ , hence  $\psi(\theta_1, t) \leq \psi(\theta_2, t)$ . If  $t \geq t_0$ ,  $\psi(\theta_1, t) = 0$ , hence  $\psi(\theta_1, t) \leq \psi(\theta_2, t)$ , this implies that  $\theta_1 < \theta_2$ . It remains to show that the topology of  $K$  and the order topology are identical.

Since  $K$  is compact and  $(K, \tau_0)$  is Hausdorff (see the introduction), it is enough to show that the order topology is weaker than the topology of  $K$ . Given  $\theta \in K$ , let

$$Z = \{\alpha \in K; \alpha < \theta\}.$$

Suppose that  $Z$  is not empty and  $\alpha \in Z$ . Since  $\alpha, \theta$  are  $\{0, 1\}$ -valued on the family  $(f^t)_{t \in E}$  and  $\alpha < \theta$ , it is clear that there is  $t \in E$  such that  $0 = \alpha(f^t) < \theta(f^t) = 1$ , hence

$$Z = \cup_{t \in E} \{\alpha \in K; \psi(\alpha, t) = 0 \text{ and } \psi(\theta, t) = 1\}.$$

On the other hand,  $K \ni \alpha \mapsto \alpha(f^t)$  is continuous for the topology of  $K$ , namely, the  $\sigma(A^*, A)$  topology; for every  $t \in E$  the set

$$\{\alpha \in K; \psi(\alpha, t) = 0 \text{ and } \psi(\theta, t) = 1\}$$

is an open subset of  $K$ , hence  $Z$  is an open subset of  $(K, \sigma(A^*, A))$ . By the same proof,  $Z' = \{\alpha \in K; \alpha > \theta\}$  is an open subset of  $K$ . ■

**Definition 1.** Let  $K$  be a Hausdorff compact space. We say that  $C(K)$  admits an equivalent  $\tau_p$ -Kadec norm, if there exists an equivalent norm  $\rho$  on  $C(K)$  such that the strong topology and the  $\tau_p$  topology coincide on  $\{f \in C(K); \rho(f) = 1\}$ .

**Corollary 1.** The space  $C(K)$  admits an equivalent  $\tau_p$ -Kadec norm.

*Proof.* This is a consequence of Proposition 1, and of Theorem A in [10]. ■

When  $(X, \tau)$  is a topological space, we denote by  $\text{Bor}(X, \tau)$  the Borel  $\sigma$ -algebra of subsets of  $X$  that is generated by the class of open subsets of  $(X, \tau)$ .

**Corollary 2.** *We have  $\text{Bor}(C(K), \|\cdot\|) = \text{Bor}(C(K), \tau_p)$ .*

*Proof.* Corollary 1 shows us that there exists on  $C(K)$  an equivalent  $\tau_p$ -Kadec norm. By [13], it follows that  $\text{Bor}(C(K), \|\cdot\|) = \text{Bor}(C(K), \tau_p)$ . ■

## 2. SURJECTIVITY OF $h$

In this part, we study the surjectivity of  $h$ . We find sufficient conditions for  $h$  to be surjective.

On the set  $\{[t]; t \in E\}$  we define an order relation by

$$[t] < [u] \text{ if } t < u \text{ and } \mu([t, u]) > 0$$

(we know that  $[t] = [u]$  when  $\mu([t \wedge u, t \vee u]) = 0$ ). It is obvious that the set  $\{[t]; t \in E\}$  is totally ordered by this relation.

**Definition 2.** *A totally ordered set is said to be complete if every nonempty subset that has an upper bound, has a least upper bound.*

One sees then that every nonempty subset  $F$  that has a lower bound has a greatest lower bound, by applying the preceding definition to the set  $F'$  of lower bounds of  $F$ .

We denote by  $\theta'_E = \theta'$  the maximal element of  $K$  and by  $\theta''_E = \theta''$  the minimal element of  $K$ . Note that  $\psi(\theta', t) = 1$  and  $\psi(\theta'', t) = 0$ , for every  $t \in E \setminus \{-\infty, +\infty\}$ . It is clear that  $\theta', \theta'' \in B_1$ , the subset of  $K$  consisting of those  $\theta$  with  $t \mapsto \theta(f^t)$  continuous on  $(E, \tau_0)$ .

**Remark 2.** *Denote by  $h_j$  the restriction of  $h$  to  $B_j$ ,  $j \in \{2, 3\}$ ; then  $h_j : B_j \rightarrow E$  is an injective function, for  $j \in \{2, 3\}$ .*

*Proof.* Let us show for example that  $h_2$  is an injective function. Let  $\theta_1, \theta_2 \in B_2$  such that  $\theta_1 < \theta_2$ . There exists  $t \in E$  such that  $\psi(\theta_1, t) = 0$  and  $\psi(\theta_2, t) = 1$ , it follows that  $h_2(\theta_1) < t \leq h_2(\theta_2)$ , (by Lemma 2, we know that  $t \leq h_2(\theta_2)$ ) hence  $h_2(\theta_1) \neq h_2(\theta_2)$ . ■

**Remark 3.** Suppose that  $(E, \leq)$  is complete. Let  $\theta \in B_1 \setminus \{\theta', \theta''\}$ . It is clear that  $\{t \in E; \psi(\theta, t) = 1\}$  and  $\{u \in E; \psi(\theta, u) = 0\}$  are both nonempty, for otherwise  $\theta$  would coincide on every  $f^t$ ,  $t \in E$ , with either  $\theta'$  or  $\theta''$ , implying that  $\theta = \theta'$  or  $\theta = \theta''$ .

Suppose that  $(E, \leq)$  is complete and let  $\theta \in B_1 \setminus \{\theta', \theta''\}$ ; there exists  $u' \in E$  with  $\psi(\theta, u') = 0$ . Note that we have  $u' \geq u$  for every  $u \in E$  such that  $\psi(\theta, u) = 1$ , hence  $\sup\{t \in E; \psi(\theta, t) = 1\}$  exists by completeness. Let us extend to  $B_1 \setminus \{\theta', \theta''\}$  the definition of  $h$  by setting  $h(\theta) = \sup\{t \in E; \psi(\theta, t) = 1\}$  (now, this formula holds for every  $\theta \in K \setminus \{\theta', \theta''\}$ , see Lemma 2).

**Lemma 4.** Suppose that the order on  $E$  is complete. Then for every  $\theta \in B_1 \setminus \{\theta', \theta''\}$ , one has  $\psi(\theta, h(\theta)) = 1$ .

*Proof.* Let  $\theta \in B_1 \setminus \{\theta', \theta''\}$  and set  $E_1(\theta) = \{t \in E; \psi(\theta, t) = 1\}$ . It is easy to see that  $h(\theta) = \sup E_1(\theta)$  belongs to the closure of  $E_1(\theta)$  for the order topology  $\tau_0$ . But  $E_1(\theta) = \{t \in E; \theta(f^t) = 1\}$  is closed since  $\theta \in B_1$  and  $t \mapsto \theta(f^t)$  is continuous. It follows that  $h(\theta) \in E_1(\theta)$ . ■

Denote  $\overline{E} = E \cup \{-\infty, +\infty\}$  ( $\overline{E} = E$ , if  $-\infty, +\infty \in E$ ). It is obvious that  $\overline{E}$  is totally ordered. Denote by  $e$  the unit element of  $C(K)$ . When  $F$  is a subset of  $E$ , denote by  $\chi_F$  the characteristic function of  $F$ .

**Lemma 5.** Let  $t \in E \setminus \{-\infty\}$ , then there exists a character  $\theta_{t,E} \in K$  (we denote  $\theta_{t,E} = \theta_t$ , When  $E$  is fixed) such that for every  $u \in E$ , we have  $\theta_t(f^u) = 1$  and only if  $[u] \leq [t]$  (the value is 0 otherwise, of course).

*Proof.* We shall define a function  $\theta_t$  on  $A$  by setting first

$$\begin{cases} \theta_t(f^u) = 1 & \text{if } [u] \leq [t], \\ \theta_t(f^u) = 0 & \text{if } [u] > [t], \\ \theta_t(e) = 1, \end{cases} \quad (2.1)$$

then extending it to the  $\mathbb{C}$ -vector subspace  $V$  of  $L^\infty(\mu)$  generated by the family  $(f^t)_{t \in E}$  and the unit  $e$  of  $A$  (constantly equal to 1 on  $E$ ). As  $f^u = f^v$  in  $L^\infty(\mu)$  when  $[u] = [v]$ , every  $g \in V$  can be expressed as

$$g = \sum_{j=1}^n \alpha_j f^{t_j} + \alpha_0 e, \quad (2.2)$$

where  $\alpha_j \in \mathbb{C}$ ,  $t_j \in E$  are such that  $[t_i] \neq [t_j]$  for  $i \neq j$ , and  $f^{t_j} \neq e$ ,  $f^{t_j} \neq 0$  for all  $1 \leq j \leq n$  (as  $L^\infty$  classes; condition (1.1) implies that  $f^t \neq 0$  when  $t \neq +\infty$ ). Since  $(E, \leq)$  is totally ordered, we may assume that  $[t_1] < [t_2] < \dots < [t_n]$ ; since  $f^{t_1} \neq e$  in  $A \subset L^\infty(\mu)$  and

$e - f^{t_1} = \chi_{F_0}$ ,  $F_0 \subset E$ , we have  $\mu(F_0) > 0$ ; it follows that  $g = \alpha_0$  on  $F_0$  and  $|\alpha_0| \leq \|g\|_{L^\infty(\mu)}$ . Next,  $[t_1] < [t_2]$  implies that  $f^{t_1} - f^{t_2} = \chi_{F_1}$  with  $F_1 = (t_1, t_2]$ ,  $\mu(F_1) > 0$ , and  $g = \alpha_0 + \alpha_1$  on  $F_1$ ; it follows that  $|\alpha_0 + \alpha_1| \leq \|g\|_{L^\infty(\mu)}$ . We proceed in this way to  $f^{t_n} \neq 0$ , with the value  $\sum_{j=0}^n \alpha_j$  on the set  $F_n = \{t \in E; t > t_n\}$  of positive measure. Finally,

$$\sup_{0 \leq k \leq n} \left| \sum_{j=0}^k \alpha_j \right| \leq \|g\|_{L^\infty(\mu)}. \quad (2.3)$$

It follows from (2.3) that  $e, f^{s_1}, \dots, f^{s_m}$  are linearly independent in  $A$  when  $[s_i] \neq [s_j]$  for  $i \neq j$  and  $f^{s_j} \neq e$ ,  $f^{s_j} \neq 0$  for all  $1 \leq j \leq m$ . Indeed, assuming as we may that  $s_1 < s_2 < \dots < s_m$ , the equality

$$\sum_{j=1}^m \alpha_j f^{s_j} + \alpha_0 e = 0$$

implies  $\sum_{j=0}^k \alpha_j = 0$  for  $0 \leq k \leq m$ , hence  $\alpha_j = 0$  for  $j = 0, \dots, m$ .

It follows that we may extend  $\theta_t$  in a natural way as a linear form on  $V$  by setting  $\theta_t(g) = \sum_{j=1}^n \alpha_j \theta_t(f^{t_j}) + \alpha_0$  whenever  $g$  is expressed as in (2.2). Furthermore,  $\theta_t$  is a continuous function on  $V$  equipped with the  $L^\infty(\mu)$  norm: indeed, pick  $g = \sum_{j=1}^n \alpha_j f^{t_j} + \alpha_0 e$  in the previous form and assume  $t_1 < t_2 < \dots < t_n$ . We shall check that  $\theta_t(g)$  is equal to one of the sums  $\sum_{j=0}^k \alpha_j$ ,  $k \in \{0, \dots, n\}$ , depending on the position of  $t$ : if  $[t] < [t_1]$ , we have that  $\theta_t(f^{t_j}) = 0$  for  $1 \leq j \leq n$ , hence  $\theta_t(g) = \alpha_0$ ; if  $[t_k] \leq [t] < [t_{k+1}]$  and  $k \in \{1, \dots, n-1\}$ , we see that  $\theta_t(f^{t_j}) = 1$  when  $j \leq k$  and  $= 0$  otherwise, thus  $\theta_t(g) = \alpha_0 + \alpha_1 + \dots + \alpha_k$ ; finally, when  $[t] \geq [t_n]$ , we have  $\theta_t(f^{t_j}) = 1$  for every  $j$  and  $\theta_t(g) = \alpha_0 + \dots + \alpha_n$ . Therefore, we have by (2.3) that

$$|\theta_t(g)| \leq \|g\|_{L^\infty(\mu)}.$$

Since the set  $\{f^u; u \in E\} \cup \{e\}$  forms a total set in  $A$ ,  $\theta_t$  extends (in a unique way) to a continuous linear form on  $A = C(K)$ .

We check finally that  $\theta(t)$  is a character. First  $\theta_t(f^u f^v) = \theta_t(f^u) \theta_t(f^v) = \theta_t(f^u) \theta_t(f^v)$  because the values are 0 or 1. Let  $u, v$  in  $E$  with  $[u] < [v]$ ; we have  $f^u f^v = f^v$  and we can check that  $\theta_t(f^u f^v) = \theta_t(f^u) \theta_t(f^v)$  by considering the cases  $[t] < [u]$  (values  $u \mapsto 0, v \mapsto 0$ ),  $[u] \leq [t] < [v]$  (values  $u \mapsto 1, v \mapsto 0$ ), and  $[v] \leq [t]$  (values  $u \mapsto 1, v \mapsto 1$ ). This implies the result, by expanding products  $g_1 g_2$  with  $g_1, g_2$  expressed as in (2.2). It follows that  $\theta_t \in K$ . ■

Let us set  $h(\theta') = +\infty$  and  $h(\theta'') = -\infty$ .

**Lemma 6.** *Let  $t \in E \setminus \{-\infty\}$ . Then*

a)  $\sup\{u \in E; u \in [t]\} = b \in [t]$  and  $\{u \in E; u > b\}$  is not closed subset of  $E$ , if and only if there exists a character  $\theta_{t,E} \in B_2$  (we denote  $\theta_{t,E} = \theta_t$ , When  $E$  is fixed) such that  $h(\theta_t) = b \in [t]$ .

b)  $\inf\{u \in E; u \in [t]\} = a \in [t]$  and  $\{u \in E; u < a\}$  is not closed subset of  $E$ , if and only if there exists  $\theta'_{t,E} = \theta'_t \in B_3$  such that  $h(\theta'_t) = a \in [t]$ .

c) Assume  $(E, \leq)$  complete. Then  $\sup\{u \in E; u \in [t]\} = b \in [t]$  and  $\{u \in E; u > b\}$  is closed subset of  $E$ , if and only if there exists  $\theta''_{t,E} = \theta''_t \in B_1 \setminus \{\theta''\}$  such that  $h(\theta''_t) = b \in [t]$ .

*Proof.*

a). Assume that  $\sup\{u \in E; u \in [t]\} = b \in [t]$  and  $\{u \in E; u > b\}$  is not closed subset of  $E$ . Consider the character  $\theta_t$  given by Lemma 5. We show that  $\theta_t \in B_2^0$  and  $h(\theta_t) = b$ . Indeed, let  $u \in E$ . If  $u > b$ , then  $[u] > [b]$  because  $\sup\{v \in E; v \in [t]\} = b$ . If  $[u] > [b]$ , it is obvious that  $u > b$ . We conclude that  $u > b$  if and only if  $[u] > [b]$ , or if and only if  $[u] > [t]$ , since  $[b] = [t]$ .

Therefore, by the formula (2.1), we have  $\psi(\theta_t, u) = 0$  if and only if  $u > b$ . Thus  $\{u \in E; \psi(\theta_t, u) = 0\} = \{u \in E; u > b\}$  is not closed subset of  $E$  by the hypothesis. We conclude that  $\theta_t \in B_2$ . Finally by Lemma 2,

$$h(\theta_t) = \sup\{u \in E; \psi(\theta_t, u) = 1\} = \sup\{u \in E; u \leq b\} = b.$$

Conversely, assume that there is  $\theta_t \in B_2$  such that  $h(\theta_t) = b \in [t]$ . By Lemma 2, we know that  $\psi(\theta_t, v) = 1$  when  $v < b$  and  $\psi(\theta_t, v) = 0$  when  $v > b$ . Thus the set  $E_0(\theta_t) = \{u \in E; \psi(\theta_t, u) = 0\}$  contains  $\{u \in E; u > b\}$  and is contained in  $\{u \in E; u \geq b\}$ ; but  $b \notin E_0(\theta_t)$ , otherwise  $E_0(\theta_t)$  would be closed, contrary to the assumption  $\theta_t \in B_2$ . It follows therefore that  $E_0(\theta_t) = \{u \in E; u > b\}$  is not closed subset of  $E$ . It remains to show that  $\sup\{u \in E; u \in [t]\} = b$ .

Let  $u \in [t] = [b]$ . We have then  $\psi(\theta_t, u) = \psi(\theta_t, b) = 1$ , and also  $u \leq h(\theta_t) = b$  by Lemma 2. It follows that  $b = \sup\{u \in E; u \in [t]\}$ . ■

b). The proof of b) is similar to that of a), replacing in Lemma 5 the formula

$$\begin{cases} \theta_t(f^u) = 1 & \text{if } [u] \leq [t], \\ \theta_t(f^u) = 0 & \text{if } [u] > [t], \\ \theta_t(e) = 1, \end{cases}$$

by the formula

$$\begin{cases} \theta'_t(f^u) = 1 & \text{if } [u] < [t], \\ \theta'_t(f^u) = 0 & \text{if } [u] \geq [t], \\ \theta'_t(e) = 1. \end{cases} \quad (2.4)$$

and replacing Lemma 2 by Lemma 3. ■

c). Suppose that  $\sup\{u \in E; u \in [t]\} = b \in [t]$  and  $\{u \in E; u > b\}$  is a closed subset

*Case 1,  $t = +\infty$  (if  $-\infty, +\infty \in E$ ):* Note that by (1.1)  $[t] = \{+\infty\}$ . Consider  $\theta_t'' = \theta'$ , it is clear that  $\theta_t'' \in B_1$  and  $h(\theta_t''(t)) = b = +\infty$ .

*Case 2,  $t < +\infty$ :* First note that  $b \neq -\infty$ . Consider the character  $\theta_t$  given by Lemma 5. Let us show that  $\theta_t \in B_1$ . Note that  $u > b$  if and only if  $[u] > [b]$ , because  $\sup\{v \in E; v \in [t]\} = b$ . It implies that  $\{u \in E; \psi(\theta_t, u) = 0\} = \{u \in E; [u] > [b]\} = \{u \in E; u > b\}$ ; this is a closed subset of  $E$  by hypothesis, as well as  $\{u \in E; u \leq b\}$  (obviously), and the latter set is  $\{u \in E; \psi(\theta_t, u) = 1\}$ ; it follows that  $\theta_t \in B_1$ . By Lemma 4,

$$h(\theta_t) = \sup\{u \in E; \psi(\theta_t, u) = 1\} = \sup\{u \in E; u \leq b\} = b.$$

We put  $\theta_t'' = \theta_t$ . It is obvious that  $\theta_t'' \neq \theta''$ .

Conversely, assume that there exists  $\theta_t'' \in B_1 \setminus \{\theta''\}$  such that  $h(\theta_t'') = b \in [t]$ .

*Case 1,  $\theta_t'' = \theta'$ :* We note that  $b = +\infty$ , then  $\{t \in E; t > b\} = \emptyset$ , which is closed subset of  $E$ . On the other hand  $\sup\{v \in E; v \in [t]\} = b = +\infty$ .

*Case 2,  $\theta_t'' \neq \theta'$ :* Since  $\psi(\theta_t'', h(\theta_t'')) = 1$  by Lemma 4, it follows that  $\{u \in E; \psi(\theta_t'', u) = 0\} = [\psi(\theta_t'', \cdot)]^{-1}\{0\} = \{u \in E; u > b\}$  is a closed subset of  $E$ , because  $\theta_t'' \in B_1$ . An argument similar to that of a) shows that  $\sup\{u \in E; u \in [t]\} = b$ . ■

**Corollary 3.** *Assume that the space  $(E, \leq)$  is complete and that we have  $\mu([t \wedge t', t \vee t']) > 0$  when  $t \neq t'$ . Then  $h|_{B_1 \cup B_2} : B_1 \cup B_2 \rightarrow \overline{E}$  is onto  $(-\infty, +\infty \notin E)$ .*

*Proof.* Note that the hypothesis implies that  $[t] = \{t\}$  for every  $t \in E$ . Let  $a \in E$ . If  $\{x \in E; x > a\}$  is not closed subset of  $E$ , by Lemma 6-a), there exists  $\theta \in B_2$  such that  $h(\theta) = a$ . If  $\{x \in E; x > a\}$  is closed subset of  $E$ , by Lemma 6-c), there exists  $\theta \in B_1$  such that  $h(\theta) = a$ . The points  $-\infty, +\infty$  are images of  $\theta', \theta''$ . Thus  $h|_{B_1 \cup B_2}$  is surjective. ■

**Proposition 2.** *Assume  $E$  satisfies (\*) and  $\mu([t \wedge t', t \vee t']) > 0$  for  $t \neq t' \in E$ . Then  $h_j : B_j \rightarrow E \setminus \{-\infty, +\infty\}$  is onto,  $j \in \{2, 3\}$ .*

*Proof.* It suffices to show by Lemma 6 that the sets  $\{u \in E; u > t\}$  and  $\{u \in E; u < t\}$  are not closed subsets of  $E$ , for every point  $t \in E \setminus \{-\infty, +\infty\}$ . Indeed, let  $a, b \in E$  be such that  $t \in ]a, b[$ . Since  $E$  satisfies the condition (\*), there exists  $c \in E$  such that  $a < c < t < b$ , so  $c \in \{u \in E; u < t\}$ , thus  $t$  is in the closure of  $\{u \in E; u < t\}$ . We

conclude that  $\{u \in E; u < t\}$  is not closed subset in  $E$ . In the same way we can see that  $\{u \in E; u > t\}$  is not a closed subset of  $E$ . ■

### 3. CONTINUITY OF $h$

In this part, we show that the function  $h$  is continuous on the subset of  $K$  where  $h$  is defined.

Recall that  $h(\theta') = +\infty$  and  $h(\theta'') = -\infty$ . Let  $K_1 = (K \setminus B_1) \cup \{\theta', \theta''\}$ . We denote again by  $h$  the restriction of  $h$  to  $K_1$ .

**Proposition 3.** *Assume that  $-\infty, +\infty \notin E$ . Then the map  $h : (K_1, \tau_0) \rightarrow (\overline{E}, \tau_0)$  is continuous.*

*Proof.* The topology of  $\overline{E}$  is generated by the intervals of the form  $\{u \in \overline{E}; u < \alpha\}$  or of the form  $\{u \in \overline{E}; u > \alpha\}$ ,  $\alpha \in E$ . Let  $\alpha \in E$  and let  $\theta_0$  belong to the set  $W = \{\theta \in K_1; h(\theta) < \alpha\}$ . Let us show that there exists an open neighborhood of  $\theta_0$  contained in  $W$ . Note that  $\theta_0 \neq \theta'$ .

By Lemma 3, we know when  $\theta_0 \neq \theta''$  that  $\psi(\theta_0, \alpha) = 0$  since  $h(\theta_0) < \alpha$  (and it is also true for  $\theta_0 = \theta''$ ), and we know that if  $\psi(\theta, \alpha) = 0$ , then  $h(\theta) \leq \alpha$ . The set

$$U = \{\theta \in K_1; \psi(\theta, \alpha) = 0\} = K_1 \cap \{\theta \in K; \theta(f^\alpha) = 0\}$$

is open in  $K_1$ , and  $U$  is contained in  $\{\theta \in K_1; h(\theta) \leq \alpha\}$ . Consider  $V = U \setminus \{\theta \in K_1; h(\theta) = \alpha\}$ . Note that  $V$  is an open set in  $K_1$ , because  $\{\theta \in K_1; h(\theta) = \alpha\}$  is finite by Remark 2, thus  $V$  is an open neighborhood of  $\theta_0$  contained in  $W$ .

Let  $\alpha \in E$  and let  $\theta_0$  belong to  $W' = \{\theta \in K_1; h(\theta) > \alpha\}$ . Using now Lemma 4, we can show in a similar way that the set

$$V' = \{\theta \in K_1; \psi(\theta, \alpha) = 1\} \setminus \{\theta \in K_1; h(\theta) = \alpha\}$$

is an open neighborhood of  $\theta_0$  and  $V' \subset W'$ . ■

**Remark 4.** *The space  $(E, \tau_0)$  is connected if and only if  $(E, \leq)$  is a complete and  $E$  satisfies the condition  $(*)$  [14, rem. (d), p. 58].*

**Corollary 4.** *Assume that  $E$  is connected and  $-\infty, +\infty \notin E$ . Then  $h : K \rightarrow (\overline{E}, \tau_0)$  is continuous.*

*Proof.* It suffices to see that  $B_1 = \{\theta', \theta''\}$ , because  $E$  is connected. ■

By an argument similar to that of Proposition 3 and its corollary, we can show the following result:

**Proposition 4.** *Assume that  $-\infty, +\infty \in E$ . Then  $h : (K_1, \tau_0) \rightarrow (E, \tau_0)$  is continuous.*

**Corollary 5.** *Assume that  $E$  is connected and  $-\infty, +\infty \in E$ . Then  $h : K \rightarrow (E, \tau_0)$  is continuous.*

**Proposition 5.** *Assume that  $(E, \leq)$  is complete and  $-\infty, +\infty \notin E$ . Then  $h : K \rightarrow \overline{E}$  is continuous.*

*Proof.* Let  $\alpha \in E$  and let  $\theta_0 \in W = h^{-1}(\{t \in \overline{E}; t < \alpha\}) = \{\theta \in K; h(\theta) < \alpha\}$ . We shall find an open neighborhood of  $\theta_0$  in  $W$ .

*Case 1,  $\theta_0 = \theta''$ :* Consider  $t_0 \in E$  such that  $t_0 < \alpha$  and let  $V = \{\theta \in K; \psi(\theta, t_0) = 0\}$ . Note that  $\theta'' \in V$ . It is obvious that  $V \subset W$ .

*Case 2,  $\theta_0 \in B_1 \setminus \{\theta''\}$ :* The set

$$\{u \in E; u < \alpha\} \cap \{t \in E; \psi(\theta_0, t) = 1\}$$

is an open neighborhood of  $h(\theta_0)$ , and because  $\psi(\theta_0, h(\theta_0)) = 1$  by Lemma 4, it follows that there exists  $a, b \in E$  ( $a < b$ ) such that  $h(\theta_0) \in ]a, b[ \subset \{u \in E; u < \alpha\} \cap \{t \in E; \psi(\theta_0, t) = 1\}$ . Denote  $V = \{\theta \in K; \psi(\theta, b) = 0\}$ ; the set  $V$  is an open neighborhood of  $\theta_0$ , because  $h(\theta_0) < b$  and  $\psi(\theta_0, h(\theta_0)) = 1$ . It remains to show that  $V \subset W$ . Indeed, let  $\theta \in V = \{\theta \in K; \psi(\theta, b) = 0\}$  (hence  $h(\theta) \leq b$ ). Let us show that  $h(\theta) < b$ . Assume that  $h(\theta) = b$ . Since  $h(\theta_0) < h(\theta) = b$ ,  $\theta_0 < \theta$ , there exists  $\gamma \in E$  such that  $\psi(\theta_0, \gamma) = 0$  and  $\psi(\theta, \gamma) = 1$ , this implies that  $a < h(\theta_0) < \gamma < h(\theta) = b$  ( $\gamma < h(\theta)$  because  $\psi(\theta, b) = 0$  and  $\psi(\theta, \gamma) = 1$ ). We deduce that  $\gamma \in ]a, b[ \subset \{u \in E; u < \alpha\} \cap \{t \in E; \psi(\theta_0, t) = 1\}$ , which is impossible (note that  $\psi(\theta_0, \gamma) = 0$ ), thus  $h(\theta) < b$ .

Now let us show that  $b \leq \alpha$ . Assume that  $b > \alpha$ . Since  $a < h(\theta_0) < \alpha$ ,  $\alpha \in ]a, b[$ , hence  $\alpha \in \{u \in E; u < \alpha\}$ , this is impossible. It follows that  $h(\theta) < b \leq \alpha$ , *i.e.*  $\theta \in W = \{\varphi \in K; h(\varphi) < \alpha\}$ . We treat the case  $\theta_0 \in B_2$  and the case  $\theta_0 \in B_3$ , as in Proposition 3. ■

By an argument similar to that of Proposition 5 we show:

**Proposition 6.** *Assume that  $(E, \leq)$  is complete and  $-\infty, +\infty \in E$ . Then  $h : K \rightarrow E$  is continuous.*

We define the measure  $\mu'$  on  $E$  by  $\mu'(\{t\}) = 1$ , for every  $t \in E$  ( $\mu'$  is the *counting measure*). Let  $A'$  be the  $C^*$ -subalgebra generated by  $\{f^t, t \in E\}$  in  $L^\infty(E, \mu') = \ell^\infty(E)$  and let  $K'_\mu(E) = K'$  be the set of characters on  $A'$ . Denote

$$h'(\theta) = \sup\{t \in E; \psi'(\theta, t) = 1\}$$

where  $\psi'(\theta, t) = \theta(f^t)$ ,  $\theta \in K'$ ,  $t \in E \setminus \{-\infty, +\infty\}$ .

4. GEOMETRIC PROPERTIES OF THE COMPACT SETS  $K$ 

In this part, we shall see that the geometry of  $K$  is related to the geometry of  $E$ .

Recall that the lexicographic order on  $E \times \{0, 1\}$  is defined in the following way: for all  $(r, i), (s, j) \in E \times \{0, 1\}$ ,

$(r, i) < (s, j)$  if and only if  $r < s$ , or  $r = s$  and  $i < j$ .

Note that if  $E$  is totally ordered, then  $E \times \{0, 1\}$  has the same property. Recall that  $\tau_1$  (resp.  $\tau_2$ ) denotes the topology on a totally ordered set generated by semi-open intervals  $(a, b]$  (resp.  $[a, b)$ ). We need the following classical lemmas:

**Lemma 7.** *Let  $E'$  be a subset of  $E$ . Then  $(E', \tau_2)$  is homeomorphic to  $(E' \times \{1\}, \tau_0)$ , viewed as topological subspace of  $(E \times \{0, 1\}, \tau_0)$  equipped with the order topology  $\tau_0$  of the lexicographic order.*

**Lemma 8.** *Assume that  $(E, \tau_0)$  is a strongly Lindelöf space. Then:*

- 1)  $(E, \tau_j)$  is a strongly Lindelöf space,  $j \in \{1, 2\}$ .
- 2)  $\text{Bor}(E, \tau_0) = \text{Bor}(E, \tau_j)$ ,  $j \in \{1, 2\}$ .
- 3) If  $(E, \tau_0)$  is separable, then  $(E, \tau_j)$  is separable,  $j \in \{1, 2\}$ .
- 4) If every subset of  $(E, \tau_0)$  is separable, then every subset of  $(E, \tau_j)$  is separable,  $j \in \{1, 2\}$ .

**Remark 5.** *Let  $(X, \tau)$  be a strongly Lindelöf space and let  $Z$  be a subspace of  $(X, \tau)$ . Then  $(Z, \tau)$  is a strongly Lindelöf space.*

*Proof of Lemma 7.* Consider  $F : (E', \tau_2) \rightarrow (E' \times \{1\}, \tau_0)$  the map defined by  $F(t) = (t, 1)$ ,  $t \in E'$ . We show that  $F$  is a homeomorphism. Let  $(a, j_0) \in E \times \{0, 1\}$  and let  $V = \{(t, 1) \in E' \times \{1\}; (t, 1) < (a, j_0)\}$  be an open subset of  $(E' \times \{1\}, \tau_0)$ . We have

$$F^{-1}(V) = \{t \in E'; (t, 1) < (a, j_0)\} = \{t \in E'; t < a\} \in (E', \tau_2).$$

Let  $(a, j_0) \in E \times \{0, 1\}$  and let  $W = \{(t, 1) \in E' \times \{1\}; (t, 1) > (a, j_0)\}$ .

*Case 1,  $j_0 = 1$ :* It is obvious that  $F^{-1}(W) = \{t \in E'; t > a\} \in (E', \tau_2)$ .

*Case 2,  $j_0 = 0$ :* We observe that  $F^{-1}(W) = \{t \in E'; t \geq a\} \in (E', \tau_2)$ .

We conclude that if  $V$  is an open subset of  $(E' \times \{1\}, \tau_0)$ , then  $F^{-1}(V)$  is an open set of  $(E', \tau_2)$  and conversely. ■

**Remark 6.** *By an argument similar to that of Lemma 7, we show that  $(E', \tau_1)$  is homeomorphic to  $(E' \times \{0\}, \tau_0)$ .*

*Proof of Lemma 8.*

1). We shall prove Lemma 8 for  $j = 1$ , the case  $j = 2$  can be shown by a similar argument. The argument that allows us to show 1) is similar to [14, p. 58, 59].

Indeed, let  $\{]a_i, b_i]; a_i < b_i, i \in I\}$  be a family of  $\tau_1$ -open intervals in  $E$ , let  $U = \bigcup_{i \in I} ]a_i, b_i]$  and  $V = \bigcup_{i \in I'} ]a_i, b_i[$ . We will show that there exists a countable subset  $I'$  of  $I$  such that  $U = \bigcup_{i \in I'} ]a_i, b_i]$ . Denote by  $J$  the subset of  $I$  consisting of those  $j$  such that  $b_j \notin V$ .

Let us show that  $\{b_j; j \in J\}$  is countable. Let  $b_j \neq b_{j'}, j, j' \in J$ . Assume that there exists  $t \in ]a_j, b_j[ \cap ]a_{j'}, b_{j'}[$ . If  $b_j < b_{j'}$ , then  $a_{j'} < t < b_j < b_{j'}$ , it follows that  $b_j \in ]a_{j'}, b_{j'}[ \subset V$ , which is impossible, hence  $]a_j, b_j[ \cap ]a_{j'}, b_{j'}[ = \emptyset$ , if  $b_j \neq b_{j'}$ . By hypothesis  $(E, \tau_0)$  is a strongly Lindelöf space, then there exists a countable subset  $H$  of  $J$ , such that  $\bigcup_{i \in J} ]a_i, b_i[ = \bigcup_{i \in H} ]a_i, b_i[$ . On the other hand, if  $b_j \neq b_{j'}, j, j' \in J$ , then  $]a_j, b_j[ \cap ]a_{j'}, b_{j'}[$  is empty; we conclude that  $\{b_j; j \in J\} = \{b_j; j \in H\}$ , *i.e.*  $\{b_j; j \in J\}$  is countable.

Since  $(E, \tau_0)$  is a strongly Lindelöf space, there exists a countable subset  $I_1$  of  $I$ , such that  $V = \bigcup_{i \in I_1} ]a_i, b_i[$ . Let  $I_2 = I_1 \cup H$ ; this set is a countable subset of  $I$ . Let

$$W = \bigcup_{i \in I_2} ]a_i, b_i].$$

We check that  $W = U$ , this will end the proof. It is clear that  $W \supset V$  since  $I_2 \supset I_1$ . Furthermore, if  $t \in U \setminus V$ , then  $t$  must be an endpoint  $b_i$  of some  $(a_i, b_i]$ , for an  $i \in I$ . This means that  $i \in J$  and  $b_i = t$  belongs to the family  $\{b_j; j \in H\}$ , hence to  $\bigcup_{j \in H} ]a_j, b_j] \subset W$ . The space  $(E, \tau_1)$  is therefore a strongly Lindelöf space. ■

2). It is clear by 1) that  $\text{Bor}(E, \tau_0) = \text{Bor}(E, \tau_1)$ . ■

3). Assume  $(E, \tau_0)$  is separable. Let  $(x_n)_{n \geq 0}$  be a dense sequence in  $(E, \tau_0)$  and  $M = \{b \in E; \exists a_b \in E \text{ such that } ]a_b, b] = \{b\}\}$ .

*Step 1:* Let us show that  $M$  is countable.

By 1),  $(E, \tau_1)$  is a strongly Lindelöf space, it follows that  $\bigcup_{b \in M} ]a_b, b] = \bigcup_{b \in M} \{b\} = M$  is countable. ■

*Step 2:* Let us show that  $M_2 = \{x_n; n \in \mathbb{N}\} \cup M$  is dense in  $(E, \tau_1)$ .

Consider  $]a, b]$  a nonempty open interval of  $(E, \tau_1)$ . If  $]a, b] = \{b\}$ , then  $b \in M$  and  $]a, b] \cap M_2 \neq \emptyset$ . If  $]a, b[ \neq \emptyset$ , then there exists  $x_n \in ]a, b[$ , *i.e.*  $x_n \in ]a, b] \cap M_2$ . We conclude that  $M_2$  is dense in  $(E, \tau_1)$ . ■

4). We need only observe that clearly, any subset  $E'$  of a strongly Lindelöf space  $E$  is strongly Lindelöf for the induced topology, and then apply 3) to  $E'$ . ■

**Remark 7.** *Suppose that  $(E, \tau_0)$  is a strongly Lindelöf space. Then  $E \times \{0, 1\}$  has the same property.*

*Proof.* By Lemma 8, the space  $(E, \tau_j)$  is a strongly Lindelöf space, when  $j \in \{1, 2\}$ . Using Lemma 7 and Remark 6, one obtains that  $(E \times \{0\}, \tau_0)$  and  $(E \times \{1\}, \tau_0)$  are strongly Lindelöf spaces, therefore  $E \times \{0, 1\} = E \times \{0\} \cup E \times \{1\}$  has the same property. ■

**Remark 8.** *Suppose that  $(E, \tau_0)$  is a separable strongly Lindelöf space (resp. every subset of  $(E, \tau_0)$  is separable). Then  $E \times \{0, 1\}$  is separable (resp. every subset of  $E \times \{0, 1\}$  is separable).*

*Proof.* By Lemma 8,  $(E, \tau_j)$  is separable,  $j \in \{1, 2\}$  (resp. every subset of  $(E, \tau_j)$  is separable,  $j \in \{1, 2\}$ ). On the other hand, by Lemma 7 and Remark 6,  $(E, \tau_1)$  is homeomorphic to  $E \times \{0\}$  and  $(E, \tau_2)$  is homeomorphic to  $E \times \{1\}$ , it follows that  $E \times \{0\}, E \times \{1\}$  are separables (resp. every subset of  $E \times \{0\}$  or of  $E \times \{1\}$  is separable), hence  $E \times \{0, 1\} = E \times \{0\} \cup E \times \{1\}$  is separable (resp. every subset of  $E \times \{0, 1\} = E \times \{0\} \cup E \times \{1\}$  is separable). ■

**Proposition 7.** *i) Assume that every subset of  $(E, \tau_0)$  is separable. Then  $B_1$  is a strongly Lindelöf space.*

*ii) Assume that  $(E, \tau_0)$  is separable and that  $(E, \leq)$  satisfies the condition (\*). Then  $B_1$  has a countable basis.*

*Proof.*

*i).* Let  $B$  be a subset of  $E$  and let  $(u_n)_{n \geq 0}$  be a dense sequence in  $(B, \tau_0)$ . For every  $u \in B$  denote  $V_u = \{\theta \in B_1; \psi(\theta, u) = 1\}$ . Let us show that  $\bigcup_{u \in B} V_u = \bigcup_{n \in \mathbb{N}} V_{u_n}$ .

Let  $\theta \in \bigcup_{u \in B} V_u$ . There exists  $u \in B$  such that  $\theta \in V_u$ , hence  $u \in H_\theta = \{t \in B; \psi(\theta, t) = 1\}$ . Thus there exists  $n \in \mathbb{N}$  such that  $u_n \in H_\theta$ , this implies that  $\theta \in V_{u_n}$ . It follows that  $\bigcup_{u \in B} V_u = \bigcup_{n \in \mathbb{N}} V_{u_n}$ .

If  $V'_u = \{\theta \in B_1; \psi(\theta, u) = 0\}$ ,  $u \in B$ , by a similar argument, we show that  $\bigcup_{u \in B} V'_u = \bigcup_{n \in \mathbb{N}} V'_{u_n}$ . For every  $u, v \in B$  let

$$W_{u,v} = \{\theta \in B_1; \psi(\theta, u) = 1 \text{ and } \psi(\theta, v) = 0\}.$$

Show that  $\bigcup_{u,v \in B} W_{u,v} = \bigcup_{m,n \in \mathbb{N}} W_{u_m, u_n}$ .

Let  $\theta \in W_{u,v}$ . Note that  $u \in H_\theta = \{t \in B; \psi(\theta, t) = 1\}$  and  $v \in H'_\theta = \{t \in B; \psi(\theta, t) = 0\}$ , hence there  $u_m \in H_\theta$  and  $u_n \in H'_\theta$ . It is obvious that  $\theta \in W_{u_m, u_n}$ , this implies that  $\bigcup_{u,v \in B} W_{u,v} = \bigcup_{m,n \in \mathbb{N}} W_{u_m, u_n}$ .

■

ii). Let  $(a_n)_{n \geq 0}$  be a dense sequence in  $(E, \tau_0)$ . For every  $m, n \in \mathbb{N}$  consider the sets  $W_{m,n} = \{\theta \in B_1; \psi(\theta, a_n) = 0 \text{ and } \psi(\theta, a_m) = 1\}$ ,  $V_m = \{\theta \in B_1; \psi(\theta, a_m) = 1\}$ ,  $V'_n = \{\theta \in B_1; \psi(\theta, a_n) = 0\}$ , and  $D = \{(m, n) \in \mathbb{N}^2; W_{m,n} \neq \emptyset\}$ . Let us show that

$$\{W_{m,n}; (m, n) \in D\} \cup \{V_m; m \in \mathbb{N}\} \cup \{V'_n; n \in \mathbb{N}\}$$

forms a basis of  $B_1$ .

In the proof of Proposition 2, we proved that  $u$  is in the closure of  $\{x \in E; x < u\}$ . Let now  $t, u \in E$  and

$$W_{t,u} = \{\theta \in B_1; \psi(\theta, u) = 0 \text{ and } \psi(\theta, t) = 1\} \neq \emptyset.$$

Pick  $\varphi \in W_{t,u}$ . The set  $\{x \in E; \psi(\varphi, x) = 0\}$  is an open neighborhood of  $u$ , hence  $\{x \in E; \psi(\varphi, x) = 0\} \cap \{x \in E; x < u\}$  is a nonempty open subset of  $E$ , hence there exists  $n \in \mathbb{N}$  such that  $\psi(\varphi, a_n) = 0$  and  $a_n < u$ . By a similar argument, one shows that there exists  $m \in \mathbb{N}$  such that  $\psi(\varphi, a_m) = 1$  and  $a_m > t$ . This yields that  $\varphi \in W_{m,n} \subset W_{t,u}$ .

By a reasoning similar to the previous one, we show that for every  $\varphi \in \{\theta \in B_1; \psi(\theta, u) = 0\}$  (resp.  $\varphi \in \{\theta \in B_1; \psi(\theta, t) = 1\}$ ), there exists  $n \in \mathbb{N}$  such that  $\varphi \in V'_n \subset \{\theta \in B_1; \psi(\theta, u) = 0\}$  (resp. there exists  $m \in \mathbb{N}$  such that  $\varphi \in V_m \subset \{\theta \in B_1; \psi(\theta, t) = 1\}$ ). ■

**Proposition 8.** *Assume that  $(E, \tau_0)$  is a strongly Lindelöf space. Then*

I)  $(B_j, \tau_0)$  is a strongly Lindelöf space ( $(B_j, \tau_0)$  viewed as topological subspace of  $(K, \tau_0)$ ),  $j \in \{1, 2\}$ .

II) *Assume that every subset of  $(E, \tau_0)$  is separable. Then every subset of  $(B_j, \tau_0)$  is separable,  $j \in \{1, 2\}$ .*

*Proof.* We shall prove I) and II) for  $j = 3$ , the proof for  $j = 2$  is similar. I).

*Step 1:* Let us show that the map  $h_3 : (B_3, \tau_0) \rightarrow (E, \tau_1)$  is continuous.

Since  $h_3 : (B_3, \tau_0) \rightarrow (E, \tau_0)$  is continuous (by Proposition 3), it suffices to show that  $(h_3)^{-1}(V)$  is an open subset of  $B_3$ , when  $V = \{t \in E; t \leq \alpha\}$ ,  $\alpha \in E$ . Let  $\alpha \in E$  and let  $V = \{t \in E; t \leq \alpha\}$ . Consider  $\theta_0 \in (h_3)^{-1}(V)$  and  $V_1 = \{\theta \in B_3; \psi(\theta, h_3(\theta_0)) = 0\}$ . It is clear that  $V_1$  is an open neighborhood of  $\theta_0$ . On the other hand, if  $\theta \in V_1$ ,  $h(\theta) \leq h(\theta_0) \leq \alpha$ , hence  $\theta \in (h_3)^{-1}(V)$ . This implies that  $V_1 \subset (h_3)^{-1}(V)$ . Thus  $(h_3)^{-1}(V)$  is an open subset of  $(B_3, \tau_0)$ . ■

*Step 2:* Let us show that the map  $h_3 : (B_3, \tau_0) \rightarrow (h_3(B_3), \tau_1)$  is a homeomorphism.

Consider  $t \in E$  and  $V_t = \{\theta \in B_3; \psi(\theta, t) = 1\}$ . By definition of  $B_3$ ,  $V_t = \{\theta \in B_3; h_3(\theta) > t\}$ , it follows that  $h_3(V_t) = \{u \in h_3(B_3); u > t\}$  is an open subset of  $(h_3(B_3), \tau_1)$ . In the same way, if we let  $W_t =$

$\{\theta \in B_3; \psi(\theta, t) = 0\} = \{\theta \in B_3; h_3(\theta) \leq t\}$ , we see that  $h_3(W_t) = \{u \in h_3(B_3); u \leq t\}$  is an open set of  $(h_3(B_3), \tau_1)$ . Thus  $h_3$  is a homeomorphism.

By Lemma 8 and Remark 5  $(h_3(B_3), \tau_1)$  is a strongly Lindelöf space, hence  $(B_3, \tau_0)$  has the same property. ■

II). By Lemma 8, every subset of  $(h_3(B_3), \tau_1)$  is separable, using step 2, one obtains that every subset of  $(B_3, \tau_0)$  is separable. ■

**Corollary 6.** *Assume that  $(E, \tau_0)$  is a strongly Lindelöf space,  $(E, \leq)$  is complete and every subset of  $(E, \tau_0)$  is separable. Then  $K$  is separable.*

*Proof.* First let us show that  $(B_1, \tau_0)$  is separable. By an argument similar to that of Proposition 8-II), one shows that the map  $h_1 : (B_1, \tau_0) \rightarrow (h(B_1), \tau_2)$ ,  $\theta \rightarrow h(\theta)$  is a homeomorphism. By Lemma 8,  $(h(B_1), \tau_2)$  is separable, hence  $(B_1, \tau_0)$  is separable. On the other hand, by Proposition 8,  $(B_j, \tau_0)$  is separable,  $j = 0, 1$ . Since  $K = B_1 \cup B_2 \cup B_3$ ,  $K$  is separable. ■

**Corollary 7.** *Assume that  $(E, \tau_0)$  is a separable space and satisfies the condition (\*). Then  $K$  is a separable strongly Lindelöf space.*

*Proof.* By Proposition 7,  $B_1$  has a countable basis. On the other hand by Remark 1,  $(E, \tau_0)$  has a countable basis, this implies that  $(E, \tau_0)$  is a strongly Lindelöf space and every subset of  $(E, \tau_0)$  is separable, therefore by Proposition 8,  $B_2, B_3$  are separable strongly Lindelöf spaces. Thus  $K = B_1 \cup B_2 \cup B_3$  is a separable strongly Lindelöf space. ■

**Remark 9.** *Let  $(L, <)$  non-empty totally ordered set; assume that the order topology of  $L$  is compact and separable. By [19] every subset of  $L$  is separable and the order topology of  $L$  is strongly Lindelöf.*

**Corollary 8.** *Assume that  $(E, \tau_0)$  is separable and connected. Then  $K$  is separable space.*

Let  $P$  be a Polish space. We denote by  $B_1(P)$  the space of first class functions (with real values or complex values) on  $P$ .

We denote by  $\tau_p$  the topology of pointwise convergence on  $B_1(P)$ .

**Remark 10.** *Let  $(E, \tau_0)$  be a separable strongly Lindelöf space. Suppose that the map  $h_j : B_j \rightarrow E \setminus \{-\infty, +\infty\}$  is onto, for  $j \in \{2, 3\}$ . Then  $K \setminus B_1$  is separable.*

*Proof.* Note first that  $(E \setminus \{-\infty, +\infty\}, \tau_0)$  is a separable strongly Lindelöf space. By Lemma 8,  $(E \setminus \{-\infty, +\infty\}, \tau_1)$  is a separable space. On the other hand by step 2 of Proposition 8, the map  $h_3 : (B_3, \tau_0) \rightarrow$

$(E \setminus \{-\infty, +\infty\}, \tau_1)$  is a homeomorphism, therefore  $(B_3, \tau_0)$  is separable. By a similar argument, one shows that  $(B_2, \tau_0)$  is separable.

As  $K - B_1 = B_2 \cup B_3$ ,  $K - B_1$  is separable. ■

**Definition 3.** Let  $L$  be a compact Hausdorff space. We say that  $L$  is a Rosenthal compact set, if there exists a Polish space  $P$  such that  $L$  continuously embedded in  $(B_1(P), \tau_p)$  [8].

**Definition 4.** Let  $X$  be a Hausdorff space.  $X$  is said to be an analytic set, if  $X$  is a continuous image of a Polish space [21, chap. II, p. 96].

**Proposition 9.** Suppose that  $(E, \tau_0)$  is an analytic set and suppose that  $-\infty, +\infty \notin E$ . Then  $K$  is a Rosenthal compact set.

*Proof.* Since  $(E, \tau_0)$  is analytic, there exists a Polish space  $P$  and a surjective continuous map  $\phi : P \rightarrow (E, \tau_0)$ . Let  $\theta \in K$ . Consider the map  $\sigma_\theta : P \rightarrow \{0, 1\}$ ,  $x \in P \rightarrow \psi(\theta, \phi(x))$ . Note at first that if  $\theta \in B_1$ ,  $\sigma_\theta$  is a continuous function. Let  $\theta \in B_2$ . Prove that  $\sigma_\theta$  is a first class function.

It is clear that

$$(\sigma_\theta)^{-1}\{1\} = \{x \in P; \psi(\theta, \phi(x)) = 1\} = \{x \in P; \phi(x) \leq h(\theta)\}$$

which is a closed subset of  $P$ . On the other hand

$$(\sigma_\theta)^{-1}\{0\} = \{x \in P; \psi(\theta, \phi(x)) = 0\} = \{x \in P; \phi(x) > h(\theta)\}$$

is an open subset of  $P$ . Therefore the sets  $(\sigma_\theta)^{-1}\{1\}$ ,  $(\sigma_\theta)^{-1}\{0\}$  are  $G_\delta$  subsets of  $P$ . It follows that  $\sigma_\theta$  is a first class function. If  $\theta \in B_3$ , by a similar argument, one shows that  $\sigma_\theta$  is a first class function. One defines the map  $\sigma : K \rightarrow B_1(P)$ , by  $\sigma(\theta) = \sigma_\theta$ ,  $\theta \in K$ . It is obvious that  $\sigma$  is into and continuous, hence  $K$  is a Rosenthal compact set. ■

**Proposition 10.** Suppose that  $(E, \leq)$  is complete and  $(E, \tau_0)$  is a strongly Lindelöf space. Then  $K$  is a strongly Lindelöf space.

*Proof.* We can suppose that  $-\infty, +\infty \notin E$ . Denote  $K_2 = B_1 \cup B_2 - \{\theta', \theta''\}$  and consider the map  $h|_{K_2} : (K_2, \tau_0) \rightarrow (h(K_2), \tau_2)$  (note that  $\sup\{t; \psi(\theta, t) = 1\}$  exists for all  $\theta \in K_2$  by Lemma 2 and Remark 3). By hypothesis,  $(E, \tau_0)$  is a strongly Lindelöf space, hence by Lemma 8,  $(E, \tau_2)$  has the same property, therefore  $(h(K_2), \tau_2)$  is a strongly Lindelöf space. Thus  $(K_2, \tau_0)$  is a strongly Lindelöf space, because  $h|_{K_2}$  is a homeomorphism (By an argument similar to that of Proposition 8-I)). Observe finally that  $K = K_2 \cup B_3 \cup \{\theta', \theta''\}$ ,  $K_2$  is a strongly Lindelöf space and  $B_3$  is a strongly Lindelöf space by Proposition 8, hence  $K$  is a strongly Lindelöf space. ■

Let  $L$  be a Hausdorff compact space. Denote  $M^+(L)$  the Radon probability measures space on  $L$ .

For every Banach space  $X$ , we denote by  $B_X$  the closed unit ball of  $X$ .

**Definition 5.** [2, chap. IX.8] *A topological space  $(X, \tau)$  is said to be a completely regular space, if  $(X, \tau)$  can be embedded in a compact Hausdorff space.*

It is clear that every topological subspace of a completely regular space is a completely regular space.

Recall that  $\mu'$  is the counting measure defined on  $E$  and  $h'(\theta) = \sup\{t \in E; \psi'(\theta, t) = 1\}$ ,  $\theta \in K' \setminus B'_1$ , where  $\psi'(\theta, t) = \theta(f^t)$ ,  $t \in E$ . Denote by  $B'_1$  the set of  $\theta \in K'$  such that the map  $t \in E \rightarrow \psi'(\theta, t) \in \{0, 1\}$  is continuous,  $B'_2 = \{\theta \in K' \setminus B'_1; \psi'(\theta, h'(\theta)) = 1\}$  and by  $B'_3 = \{\theta \in K' \setminus B'_1; \psi'(\theta, h'(\theta)) = 0\}$ .

**Remark 11.** *Suppose that  $(E, \leq)$  is complete ( $-\infty, +\infty \notin E$ ). Then  $(E, \tau_j)$  is a completely regular,  $j \in \{1, 2\}$ .*

*Proof.* We shall show this remark for  $j = 2$ . Consider  $\mu = \mu'$ . By Corollary 3,  $h'(K_2) = E$ . On the other hand, in the proof of Proposition 10, we have shown that  $h' : B'_2 \cup B'_1 \setminus \{\theta', \theta''\} = K_2 \rightarrow (E, \tau_2)$  is a homeomorphism. Thus  $(E, \tau_2)$  is a completely regular space. ■

Let

$$E_1 = \{a \in E; \{x \in E; x > a\} \text{ is not closed subset of } E\}$$

and

$$E_2 = \{a \in E; \{x \in E; x < a\} \text{ is not closed subset of } E\}.$$

**Remark 12.** *The space  $(E_2, \tau_1)$  is completely regular space.*

*Proof.* Consider  $\mu = \mu'$ . By Lemma 6-b), the restriction of  $h'$  (in Remark 11) to  $B'_3$  is onto on  $E_2$  and by step 2 of Proposition 8, this map is homeomorphism. On the other hand,  $B'_3$  is a completely regular space, hence  $(E_2, \tau_1)$  has the same property. ■

**Remark 13.** *Suppose that  $(E, \leq)$  satisfies the condition (\*). Since the point  $a$  is in the closure of the set  $\{x \in E; x < a\}$ , for every  $a \in E$ , the set  $\{x \in E; x < a\}$  is not closed. We conclude that  $E = E_2$ .*

**Proposition 11.** *Suppose that  $E$  is a Polish space and  $\mu(\{a\}) = 0$  for every  $a \in E$  (i. e.,  $\mu$  is diffuse on the Borel  $\sigma$ -algebra of  $E$ ). Then for every  $\xi \in C(K)^*$  the map  $t \in (E, \tau_0) \rightarrow \xi(f^t)$  is a first-class function.*

*Proof.* It suffices to show Proposition 11 when  $\xi \in B_{C(K)^*}$ .

*Step 1:* Let us show that for every  $g \in L^1(E, \mu)$  the map  $t \in E \rightarrow (g, f^t) = \int_E g(x)f^t(x)d\mu(x)$  is continuous.

Let  $g \in L^1(E, \mu)$  and let  $(t_n)_{n \geq 0}$  be a sequence in  $E$  such that  $t_n \rightarrow t$  in  $(E, \tau_0)$ . Consider  $y \in E$  such that  $y \neq t$ . We shall show that  $f^{t_n}(y) \rightarrow_{n \rightarrow +\infty} f^t(y)$ .

We show that there does not exist two subsequences  $(t_{m_k})_{k \geq 0}$  and  $(t_{n_k})_{k \geq 0}$  such that  $f^{t_{m_k}}(y) = 1$  and  $f^{t_{n_k}}(y) = 0$  for every  $k \in \mathbb{N}$ . Assume there are such subsequences. We have then  $t_{n_k} \geq y > t_{m_k}$  for every  $k \in \mathbb{N}$ , by passing to the limit we obtain  $y = t$ , which is impossible.

Thus we distinguish two cases:

*Case 1*, there exists  $n_0 \in \mathbb{N}$  such that  $f^{t_n}(y) = 1$  for every  $n \geq n_0$ : Note that  $y > t_n, \forall n \geq n_0$ , hence  $y \geq t$ . Since  $y \neq t, y > t$ , i.e.  $f^{t_n}(y) \rightarrow_{n \rightarrow +\infty} f^t(y)$ .

*Case 2*, there exists  $n_0$  such that  $f^{t_n}(y) = 0$  for every  $n \geq n_0$ : This implies that  $y \leq t_n$  for every  $n \geq n_0$ , by passing to the limit, we obtain  $y \leq t$ . Thus  $f^{t_n}(y) \xrightarrow[n \rightarrow \infty]{} f^t(y) = 0$ .

By hypothesis on  $\mu, (g, f^{t_n}) = \int_{E \setminus \{t\}} g(y) f^{t_n}(y) d\mu(y)$ . By Lebesgue's dominated convergence theorem, we have

$$\begin{aligned} (g, f^{t_n}) &= \int_{E \setminus \{t\}} g(y) f^{t_n}(y) d\mu(y) \\ &\xrightarrow[n \rightarrow \infty]{} \int_{E \setminus \{t\}} g(y) f^t(y) d\mu(y) \\ &= (g, f^t). \blacksquare \end{aligned}$$

*Step 2*: Let  $\xi \in M^+(K)$ . Let us show that the map  $t \in E \rightarrow \xi(f^t) \in \mathbb{R}$  is a first class function.

By Hahn-Banach theorem there exists  $\xi' \in B_{L^\infty(\mu)^*}$  which extends  $\xi$  with  $\|\xi\| = \|\xi'\|$ . Since  $B_{L^1(\mu)}$  is  $w^*$ -dense in  $B_{(L^\infty(\mu))^*} = B_{(L^1(\mu))^{**}}$ , there is a net  $(g_\alpha)_{\alpha \in I}$  in the unit ball of  $L^1(\mu)$  such that  $g_\alpha \xrightarrow{u} \xi'$  for the topology  $w^*$  in  $[L^\infty(\mu)]^*$ . There exist  $\xi_1, \xi_2 \in M^+(K)$  such that  $g_\alpha^+ \rightarrow \xi_1 \in M^+(K)$  and  $g_\alpha^- \rightarrow \xi_2 \in M^+(K), \sigma[C(K)^*, C(K)]$ , hence  $\xi = \xi_1 - \xi_2$ , because  $g_\alpha = g_\alpha^+ - g_\alpha^-$ , for all  $\alpha$  and  $M^+(K)$  is  $\sigma(C(K)^*, C(K))$  compact. By using Proposition 9,  $K$  is a Rosenthal compact set. On the other hand, by [8],  $M^+(K)$  is a Rosenthal compact set, and by [3]  $M^+(K)$  is angelic. Therefore, there exist two subsequences  $(g_{j_k}^+)_{k \geq 0}, (g_{i_k}^-)_{k \geq 0}$ , such that  $g_{j_k}^+ \xrightarrow[k \rightarrow +\infty]{} \xi_1$  and  $g_{i_k}^- \xrightarrow[k \rightarrow +\infty]{} \xi_2$ , for the topology  $w^*$ , i.e.  $h_k = g_{j_k}^+ - g_{i_k}^- \xrightarrow[k \rightarrow +\infty]{} \xi_1 - \xi_2 = \xi$  for the topology  $w^*$ . Thus,  $(h_k, f^t) \xrightarrow[k \rightarrow +\infty]{} \xi(f^t)$  for every  $t \in E$ .

Finally, by step 1, the map  $t \in E \rightarrow (h_k, f^t)$  is continuous for every  $k$ , which implies that  $t \in E \rightarrow \xi(f^t)$  is a first class function. ■

If  $\xi$  in the closed unit ball of  $[C(K)]^*$ , it suffices to see that there exists  $u^+, u^-, v^+, v^- \in M^+(K)$  such that  $\xi = (u^+ - u^-) + i(v^+ - v^-)$ , by using step 2, we obtain that the map  $t \in E \rightarrow \xi(f^t)$  is a first class function. ■

Denote  $J = [E \cup \{+\infty\}] \times \{1\} \cup [E \cup \{-\infty\}] \times \{0\}$  ( $-\infty, +\infty \notin E$ ). As  $J$  is a subset of  $\overline{E} \times \{0, 1\}$  and  $\overline{E} \times \{0, 1\}$  is totally ordered (for the lexicographic order),  $J$  is totally ordered. We denote again by  $\tau_0$  the order topology of  $J$ .

Let  $J_1$  be a subset of  $J$ . Recall that the notation  $(J_1, \tau_0)$  means that the topology  $\tau_0$  on  $J_1$  is the topology formed of relative open subsets of  $J_1$  in  $(J, \tau_0)$ . Let

$$J_1 = [h(B_2 \cup B_1 \setminus \{\theta''\}) \times \{1\}] \cup [h(B_3 \cup \{\theta''\}) \times \{0\}].$$

In following Theorem  $J_1$  is equipped by the relative topology  $\tau_0$ .

**Theorem 1.** *Assume that  $(E, \leq)$  is complete and  $-\infty, +\infty \notin E$ . Then  $K$  is topological homeomorphic and order isomorphic to  $J_1$ .*

*Proof.*

One defines  $T : K \rightarrow (J_1, \tau_0)$ , by

$$T(\theta) = \begin{cases} (h(\theta), 1), & \text{if } \theta \in B_2 \cup (B_1 \setminus \{\theta''\}) \\ (h(\theta), 0), & \text{if } \theta \in B_3 \cup \{\theta''\} \end{cases}, \quad \theta \in K.$$

Let us show that  $T$  is a homeomorphis. Since the restriction of  $h$  to  $B_j$  is injective,  $j \in \{1, 2, 3\}$ ,  $T$  is injective. So suffices to show that  $T$  is continuous, because  $K$  is a Hausdorff compact space.

Let  $(\alpha, \beta) \in J$  and let  $V = \{(x, y) \in J_1; (x, y) < (\alpha, \beta)\}$  be an open subset of  $J_1$ . We prove that  $T^{-1}(V)$  is an open subset of  $K$ . We distinguish two cases:

*Case 1,  $\beta = 0$ :* It is obvious that  $T^{-1}(V) = (h)^{-1}(\{t \in \overline{E}; t < \alpha\})$  which is an open subset of  $K$ , by Proposition 5.

*Case 2,  $\beta = 1$ :* Let  $\theta \in T^{-1}(V)$ . We shall find an open neighborhood  $C$  of  $\theta$  in  $T^{-1}(V)$ . We distinguish again two cases:

*Case 2-a,  $h(\theta) < \alpha$ :* Put  $C = h^{-1}(\{t \in \overline{E}; t < \alpha\})$  (note that  $\theta \in C$ ). if  $\varphi \in C$ ,  $T(\varphi) = (h(\varphi), j) < (\alpha, 1)$ , ( $j \in \{0, 1\}$ ), hence  $C \subset T^{-1}(V)$ . On the other hand, by Proposition 5,  $C$  is an open subset of  $K$ .

*Case 2-b,  $h(\theta) = \alpha > -\infty$ :* Note that  $\theta \neq \theta'$ , because  $\theta \in T^{-1}(V)$ . Denote  $C = \{\varphi \in K; \psi(\varphi, h(\theta)) = 0\}$ .  $C$  is an open subset of  $K$ .

Let us show that  $C$  contains  $\theta$ .

Since  $\theta \in T^{-1}(V)$ ,  $T(\theta) = (h(\theta), j) < (h(\theta), 1)$ , hence  $j = 0$ . By definition of  $T$   $\psi(\theta, h(\theta)) = 0$ , i.e.  $\theta \in C$ .

It remains to show that  $C \subset T^{-1}(V)$ . Let in fact  $\varphi \in C$ ; this means that  $\psi(\varphi, h(\theta)) = 0$ , we conclude that  $h(\varphi) \leq h(\theta)$ , if  $h(\varphi) = h(\theta) = \alpha$ , then  $\psi(\varphi, h(\varphi)) = \psi(\varphi, h(\theta)) = 0$ . It follows that  $T(\varphi) = (h(\varphi), 0) < (\alpha, 1)$ . Thus  $\varphi \in T^{-1}(V)$ .

If  $h(\varphi) < h(\theta) = \alpha$ , then  $T(\varphi) = (h(\varphi), j) < (\alpha, 1)$  ( $j \in \{0, 1\}$ ), hence  $\varphi \in T^{-1}(V)$ . Therefore  $C \subset T^{-1}(V)$ .

Let now  $(\alpha, \beta) \in J$  and let  $V = \{(x, y) \in J_1; (x, y) > (\alpha, \beta)\}$  be an open subset of  $J_1$ . Show that  $T^{-1}(V)$  is an open subset of  $K$ .

*Case 1,  $\beta = 1$ :* By Proposition 5,  $T^{-1}(V) = h^{-1}(\{t \in \overline{E}; t > \alpha\})$  is an open subset of  $K$ .

*Case 2,  $\beta = 0$ :* Consider  $\theta \in T^{-1}(V)$ . We distinguish two cases:

*Case 2-c,  $h(\theta) > \alpha$ :* The set  $C = h^{-1}(\{t \in \overline{E}; t > \alpha\})$  is an open subset of  $K$ , by Proposition 5 which contains  $\theta$ . The condition  $h(\varphi) > \alpha$  implies that  $T(\varphi) = (h(\varphi), j) > (\alpha, 0)$ , hence  $C \subset T^{-1}(V)$ .

*Case 2-d,  $h(\theta) = \alpha < +\infty$ :* Note that  $\theta \neq \theta''$ , because  $\theta \in T^{-1}(V)$ . Choose  $C = \{\varphi \in K; \psi(\varphi, h(\theta)) = 1\}$ . The hypothesis  $\theta \in T^{-1}(V)$  implies that  $T(\theta) = (h(\theta), j) > (h(\theta), 0)$ , hence  $j = 1$ , this means that  $\psi(\theta, h(\theta)) = 1$ . It follows that  $C$  contains  $\theta$ . Show that  $C \subset T^{-1}(V)$ . Let  $\varphi \in C$ . It is clear that  $h(\varphi) \geq h(\theta)$ , if  $h(\theta) = h(\varphi)$ , then  $\psi(\varphi, h(\varphi)) = \psi(\varphi, h(\theta)) = 1$ , hence  $T(\varphi) = ((h(\varphi), 1) > (\alpha, 0)$ , i.e.  $\varphi \in T^{-1}(V)$ . Finally if  $h(\theta) < h(\varphi)$ , then  $T(\varphi) = (h(\varphi), j) > (h(\theta), 0) = (\alpha, 0)$ , this implies that  $\varphi \in T^{-1}(V)$ . Thus  $C \subset T^{-1}(V)$ .

Let us show that  $T$  is an order isomorphism.

Let  $\theta, \varphi \in K$  such that  $\theta < \varphi$ . There exists  $t \in E$  such that  $\psi(\theta, t) = 0$  and  $\psi(\varphi, t) = 1$ . This implies that

$$h(\varphi) \geq t \geq h(\theta). \quad (4.1)$$

Obviously if  $h(\theta) < h(\varphi)$ , then  $T(\theta) = (h(\theta), j_1) < (h(\varphi), j_2) = T(\varphi)$ . If  $h(\theta) = h(\varphi)$ , by (4.1)  $h(\theta) = h(\varphi) = t$ , hence  $\theta \in B_3^0$  and  $\varphi \in B_2$ , this implies that  $T(\theta) = (h(\theta), 0) < (h(\varphi), 1) = T(\varphi)$ .

Conversely, suppose that  $T(\theta) < T(\varphi)$ . If  $h(\theta) < h(\varphi)$ , then  $\theta < \varphi$ , if  $h(\theta) = h(\varphi)$ , since  $T(\theta) = (h(\theta), j_1) < T(\varphi) = (h(\varphi), j_2)$ ,  $j_1 = 0$  and  $j_2 = 1$ . Thus  $\theta \in B_3$  and  $\varphi \in B_2$ , hence  $\theta < \varphi$ . ■

**Corollary 9.** *Suppose that  $E$  is connected and  $\mu([t \wedge t', t \vee t']) > 0$ , for  $t \neq t'$ . Then  $K$  is topological homeomorphic and order isomorphic to*

$$J = ([E \cup \{+\infty\}] \times \{1\} \cup [E \cup \{-\infty\}] \times \{0\}, \tau_0).$$

*Proof.* Since  $E$  is connected, by Remark 4,  $(E, \leq)$  is complete. On the other hand, by Proposition 2,  $h_j$  is surjective onto  $E \setminus \{-\infty, +\infty\}$ ,

$j \in \{2, 3\}$ , it follows that  $T$  (defined in Theorem 6) is surjective (note that  $B_1 = \{\theta', \theta''\}$ ). ■

**Corollary 10.** *Let  $E$  be an abelian group ( $-\infty, +\infty \notin E$ ). Assume that  $E$  separable and connected. Then  $K$  is topological homeomorphic and order isomorphic to*

$$([E \cup \{+\infty\}] \times \{1\} \cup [E \cup \{-\infty\}] \times \{0\}, \tau_0).$$

*Proof.* The hypothesis that  $E$  is connected implies that  $(E, \leq)$  is complete. On the other hand,  $E$  satisfies the condition  $(*)$ , the example (1) shows that the measure  $\mu = m$  verifies the hypothesis of Corollary 9, hence  $K$  is isomorphic to  $([E \cup \{+\infty\}] \times \{1\} \cup [E \cup \{-\infty\}] \times \{0\}, \tau_0)$ . ■

**Lemma 9.** *Assume that  $(E, \leq)$  is complete ( $E$  is a separable strongly Lindelöf space). Then:*

- 1)  $J = ([E \cup \{+\infty\}] \times \{1\}) \cup [(E \cup \{-\infty\}) \times \{0\}]$ ,  $\tau_0$  is compact.
- 2)  $(J, \tau_0)$  is a separable space.

*Proof.*

1). Consider  $\mu = \mu'$  the counting measure. By using Theorem 1, we obtain that the set  $([E \cup \{+\infty\}] \times \{1\}) \cup [(h'(B'_3) \cup \{-\infty\}) \times \{0\}]$  is compact in  $(J, \tau_0)$ , because this space is homeomorphic to  $K'$ . By a similar argument (by changing the statement of Theorem 1) we may show that the set  $([h'(B'_2) \cup \{+\infty\}] \times \{1\}) \cup [(E \cup \{-\infty\}) \times \{0\}]$  is compact in  $(J, \tau_0)$ , thus

$$\begin{aligned} & ([E \cup \{+\infty\}] \times \{1\}) \cup [(h'(B'_3) \cup \{-\infty\}) \times \{0\}] \\ & \cup [(h'(B'_2) \cup \{+\infty\}) \times \{1\}] \cup [(E \cup \{-\infty\}) \times \{0\}] = J \end{aligned}$$

is compact. ■

2). Since  $E$  is a separable strongly Lindelöf space,  $(\overline{E}, \tau_0)$  has the same property. By Lemma 8,  $(\overline{E}, \tau_2)$  is a separable strongly Lindelöf space. On the other hand Lemma 7 shows us that  $(\overline{E} \times \{1\}, \tau_0)$  is a separable subspace of  $(\overline{E} \times \{0, 1\}, \tau_0)$ . Note now that  $[E \cup \{+\infty\}] \times \{1\} = \overline{E} \times \{1\} \setminus \{(-\infty, 1)\}$ , hence  $([E \cup \{+\infty\}] \times \{1\}, \tau_0)$  is separable. By a similar argument, one shows that  $([E \cup \{-\infty\}] \times \{0\}, \tau_0)$  is a separable of  $\overline{E} \times \{0, 1\}$ . It follows that  $([E \cup \{+\infty\}] \times \{1\}) \cup [(E \cup \{-\infty\}) \times \{0\}] = J$  is a separable space of  $\overline{E} \times \{0, 1\}$ . Let  $i : (J, \tilde{\tau}_0) \rightarrow (J, \tau_0)$  be the identity map of  $J$  ( $(J, \tilde{\tau}_0)$ , where is a topological subspace of  $\overline{E} \times \{0, 1\}$ ). It is clear that  $i$  is continuous, thus  $(J, \tau_0)$  is separable. ■

By an argument similar to that of Theorem 1, one shows the following theorem

**Theorem 2.** *Suppose that  $(E, \leq)$  is complete and  $-\infty, +\infty \in E$ . Then  $K$  is topological homeomorphic and order isomorphic to*

$$([(h(B_2 \cup B_1 \setminus \{\theta''\}) \times \{1\}) \cup [(h(B_3 \cup \{\theta''\}) \times \{0\}), \tau_0).$$

By an argument similar to that of Corollary 9, one shows the following corollary:

**Corollary 11.** *Suppose that  $E$  is connected, that  $-\infty, +\infty \in E$  and that  $\mu([t \wedge t', t \vee t']) > 0$ , for  $t \neq t'$ . Then  $K$  is topological homeomorphic and order isomorphic to  $([E \setminus \{-\infty\}] \times \{1\} \cup [E \setminus \{+\infty\}] \times \{0\}, \tau_0)$ .*

**Theorem 3.**

**Remark 14.** *Let  $(E, \tau_0)$  be a separable strongly Lindelöf space. Suppose that  $(E, \leq)$  is complete and  $-\infty, +\infty \in E$ . Then  $K$  is a separable..*

#### 5. CONSTRUCTION OF A TOTALLY ORDERED COMPACT SET HOMEOMORPHIC TO $K_\mu(E)$

This part is devoted to find sufficient conditions for  $K$  to be isomorphic to  $E$ .

Recall that  $E_1 = \{t \in E; \{x \in E; x > t\} \text{ is not closed in } E\}$ , that  $E_2 = \{t \in E; \{x \in E; x < t\} \text{ is not closed in } E\}$ ,  $h(\theta') = +\infty$  and  $h(\theta'') = -\infty$ .

**Theorem 4.** *Assume that  $(E, \leq)$  is complete, that  $-\infty, +\infty \in E$ ,  $E \setminus \{+\infty, -\infty\} = E_1 \cup E_2$  and for every  $t \in E_1 \cup E_2$ , there exists  $b \in E_1$  and  $a \in E_2$  such that  $[t] = \{a, b\}$ ,  $b > a$ . Then the map  $h : K \rightarrow E$  is a homeomorphism.*

*Proof.*

*Step 1:* Show that  $h : K \rightarrow E$  is onto.

Let  $b \in E_1$ . By hypothesis, there exists  $a \in E_2$  such that  $[b] = \{b, a\}$  and  $a < b$ . By Lemma 6, there is  $\theta_b \in B_2$  such that  $h(\theta_b) = b$ . In the same way, for every  $a' \in E_2$ , there is  $\theta'_{a'} \in B_3$  such that  $h(\theta'_{a'}) = a'$ . ■

*Step 2:* Let us show that the map  $h : K \rightarrow E$  is into.

Suppose there exist  $\theta_1, \theta_2 \in K$  such that  $h(\theta_1) = h(\theta_2) = t_0$ . Since the restriction of  $h$  to  $B_1 \cup B_2$  (resp. to  $B_3$ ) is injective, by step 1 of Proposition 10 (resp. by Remark 2)), it is enough to treat the following cases:

*Case 1,  $\theta_1 \in B_2$  and  $\theta_2 \in B_3$ :*

*Case 1-a,  $t_0 \in E_1$ :* By hypothesis, there exists  $a \in E_2$  such that  $[t_0] = \{a, b\}$  and  $b = t_0 > a$ . On the other hand, since  $a \in [t_0] = [b]$

and  $\theta_2 \in B_3$ ,  $\psi(\theta_2, a) = \psi(\theta_2, b) = \psi(\theta_2, h(\theta_2)) = 0$ , this implies that  $a \geq h(\theta_2) = b$ , it is impossible because  $b = t_0 > a$ .

*Case 1-b*,  $t_0 = a \in E_2$ : By hypothesis, there exists  $b \in E_1$  such that  $[t_0] = \{a, b\}$  and  $t = t_0 < b$ . Observe that  $\psi(\theta_1, b) = \psi(\theta_1, a) = \psi(\theta_1, h(\theta_1)) = 1$ . Thus  $b \leq h(\theta_1) = t_0$ , which is impossible.

*Case 2*,  $\theta_1 \in B_1 \setminus \{\theta', \theta''\}$  and  $\theta_2 \in B_3$ :

*Case 2-c*,  $t_0 = b \in E_1$ : By hypothesis, there exists  $a \in E_2$  such that  $[t_0] = \{a, b\}$  and  $a < b = t_0$ . Note that  $\psi(\theta_2, a) = \psi(\theta_2, b) = 0$ , hence  $a \geq h(\theta_2) = t_0 = b$ , this is impossible.

*Case 2-d*,  $t_0 = a \in E_2$ : There exists  $b \in E_1$  such that  $[t_0] = \{a, b\}$  and  $t_0 = a < b$ . In this case, we have  $\psi(\theta_1, b) = \psi(\theta_1, a) = 1$ , by lemma 4, hence  $b \leq h(\theta_1) = a$ , which is impossible.

Finally, by Proposition 6,  $h$  is continuous, therefore  $h$  is a homeomorphism. ■

**Remark 15.** *Under the hypothesis of Theorem 4, we have  $B_1 = \{\theta', \theta''\}$ .*

*Proof.* Suppose that  $\theta \in B_1 \setminus \{\theta', \theta''\}$ . Note that  $h(\theta) = t_0 \in E \setminus \{-\infty, +\infty\}$ . In the proof of Theorem 4-step 1, we showed that there exists  $\varphi \in B_2 \cup B_3$  such that  $h(\varphi) = h(\theta) = t_0$ . We deduce that  $\theta = \varphi$ , which is impossible. ■

**Remark 16.** *Under the hypothesis of Theorem 4, by step 1, one notes that if  $\theta \in B_2^0$  (resp.  $\theta \in B_3$ )  $h(\theta) \in E_1$  (resp.  $h(\theta) \in E_2$ ).*

Consider  $E = [0, 1] \times \{0, 1\}$  equipped with the lexicographic order and  $\Phi : [0, 1] \rightarrow [0, 1] \times \{0, 1\}$  the map defined by  $\Phi(t) = (t, 1)$ ,  $t \in [0, 1]$ .  $\Phi$  is Borel. Indeed, let  $F : ([0, 1], \tau_2) \rightarrow ([0, 1] \times \{1\}, \tau_0)$  the map defined by  $F(t) = (t, 1)$ . In the proof of Lemma 7 we showed that  $F$  is a homeomorphism. On the other hand by Lemma 8,  $\text{Bor}([0, 1], \tau_2) = \text{Bor}([0, 1], \tau_0)$ , hence  $\Phi$  is Borel.

Put  $\mu = \Phi(m_1)$ , where  $m_1$  the Lebesgue measure on  $[0, 1]$ .

**Corollary 12.**  *$h : K \rightarrow (E = [0, 1] \times \{0, 1\}, \mu)$  is a homeomorphism.*

*Proof.* Put  $E_1 = [0, 1] \times \{1\} \setminus \{(1, 1)\}$  and  $E_2 = [0, 1] \times \{0\} \setminus \{(0, 0)\}$ . Let  $b = (t_0, 1) \in E_1$ . Suppose that  $b_1 = (t_1, j_1) \in E \times \{0, 1\}$  and  $t_0 > t_1$ ,  $j_1 \in \{0, 1\}$ . One has

$$\begin{aligned} \mu([b_1, b]) &= \mu(\{(t, j); (t_1, j_1) < (t, j) \leq (t_0, 1)\}) \\ &= m_1(\{t \in E; (t_1, j_1) < (t, 1) \leq (t_0, 1)\}) \\ &\geq m_1(\{t \in E; t_1 < t < t_0\}) > 0. \end{aligned}$$

If  $b_1 = (t_1, j_1)$  and  $t_0 < t_1$ , by a similar argument, one shows that  $\mu([b, b_1]) > 0$ .

Consider  $a = (t_0, 0)$ . It is clear that  $\mu(]a, b]) = 0$ , hence  $[b] = \{a, b\}$  (by above).

It remains to show that  $V = \{(t, j) \in E \times \{0, 1\}; ((t, j) > (t_0, 1))\}$  is not closed in  $E$  (by a similar argument, one shows that  $V = \{(t, j) \in E \times \{0, 1\}; ((t, j) < (t_0, 0))\}$  is not closed in  $E$ ).

Let  $(t_n)_{n \geq 0}$  be a sequence in  $]0, 1[$  such that for every  $n \in \mathbb{N}$   $t_n > t_0$  and  $t_n \xrightarrow{n \rightarrow +\infty} t_0$ . Since  $(t_n, 1) \in V$  for every  $n \in \mathbb{N}$ , it is enough to show that  $(t_n, 1) \xrightarrow{n \rightarrow +\infty} (t_0, 1) \notin V$ .

Let  $W = \{(u, j) \in E \times \{0, 1\}; (u, j) < (u_0, j_0)\}$  be an open neighborhood of  $(t_0, 1)$ . Since  $t_0 < u_0$  (because  $(t_0, 1) \in W$ ), there exists  $n_0 \in \mathbb{N}$  such that  $t_n < u_0$  for every  $n \geq n_0$ , it follows that  $(t_n, 1) \in W$ .

Consider  $W = \{(u, j) \in E \times \{0, 1\}; (u, j) > (u_0, j_0)\}$  an open neighborhood of  $(t_0, 1)$ . Obviously  $t_0 \geq u_0$ , hence  $t_n > t_0 \geq u_0$ . It follows that  $(t_n, 1) \in W$ , for every  $n$ . By Theorem 4,  $h$  is a homeomorphism. ■

In the following of this part, one supposes that  $(E, \tau_0)$  is a connected strongly Lindelöf space,  $-\infty, +\infty \in E$  and  $[t] = \{t\}$  for all  $t \in E$ . One defines the map  $\Pi_1 : E \rightarrow K$  by

$$\Pi_1(t) = \begin{cases} \theta_t, & \text{if } t \in ]-\infty, +\infty[ \\ \theta', & \text{if } t = +\infty \\ \theta'', & \text{if } t = -\infty. \end{cases}$$

By Lemma 6,  $\theta_t \in B_2$ , and  $h(\theta_t) = t, \forall t \in E \setminus \{-\infty, +\infty\}$ .

**Lemma 10.** *The map  $\Pi_1$  is Borel.*

*Proof.* Consider  $V_a = \{\theta \in K; \psi(\theta, a) = 1\}$  and  $V'_a = \{\theta \in K; \psi(\theta, a) = 0\}$ . It suffices to show  $(\Pi_1)^{-1}(V_a)$  and  $(\Pi_1)^{-1}(V'_a)$  are Borel subsets of  $E$  for every  $a \in E \setminus \{-\infty, +\infty\}$ , because by Proposition 10  $K$  is a strongly Lindelöf space (note that  $(E, \tau_0)$  is connected implies that  $(E, \leq)$  is complete and satisfies the condition (\*) (cf. [14, rem. (d), p. 58]).

Pick  $a \in E \setminus \{-\infty, +\infty\}$ . One has

$$\begin{aligned} (\Pi_1)^{-1}(V_a) &= \{t \in E \setminus \{-\infty, +\infty\}; \psi(\theta_t, a) = 1\} \cup \{+\infty\} = \\ \{t \in E \setminus \{-\infty, +\infty\}; a \leq h(\theta_t)\} \cup \{+\infty\} &= \{t \in E \setminus \{-\infty, +\infty\}; a \leq t\} \cup \{+\infty\}. \end{aligned}$$

Thus  $(\Pi_1)^{-1}(V_a)$  is Borel. By a similar argument, one shows that  $(\Pi_1)^{-1}(V'_a)$  is Borel. ■

Put  $\mu_1 = \Pi_1(\mu)$ . Note that  $\mu_1$  is a measure on  $K$ .

**Lemma 11.** *Let  $\theta_0 \in B_2$  and  $\theta_1 \in B_3$ . Then:*

I) The sets  $\{\varphi \in K; \varphi > \theta_0\}$ ,  $\{\varphi \in K; \varphi < \theta_1\}$  are not closed subsets in  $K$ .

II) For every  $\varphi_0 \in B_2$ , there exists  $\varphi_1 \in B_3$  such that  $[\varphi_0] = \{\varphi_0, \varphi_1\}$  and  $h(\varphi_0) = h(\varphi_1)$ .

*Proof.*

I). We shall show that the set  $\{\varphi \in K; \varphi > \theta_0\}$  is not closed, the proof is similar for  $\{\varphi \in K; \varphi < \theta_1\}$ .

*Step 1:* Let  $\varphi \in K$  such that  $\varphi > \theta_0$ . Show that  $h(\varphi) > h(\theta_0)$ .

Suppose that  $h(\varphi) = h(\theta_0)$ . Since the restriction of  $h$  to  $B_j$  is injective,  $j \in \{2, 3\}$ ,  $\varphi \in B_3$ , this is impossible because  $\psi(\varphi, h(\varphi)) = 0$ ,  $\psi(\theta_0, h(\theta_0)) = 1$  and  $\varphi > \theta_0$ . It follows that  $h(\varphi) > h(\theta_0)$ . ■

*Step 2:* Show that  $\{\varphi \in K; \varphi > \theta_0\}$  is not closed in  $K$ .

By the step 1,  $\{\varphi \in K; \varphi > \theta_0\} = \{\varphi \in K; h(\varphi) > h(\theta_0)\}$ . Suppose that this subset is closed in  $K$ . We shall show that the set  $\{t \in E; t > h(\theta_0)\}$  is closed, in this case the proof will be finished because  $E$  is connected, hence  $\{t \in E; t > h(\theta_0)\}$  is not closed. ■

By Proposition 6, the map  $h : K \rightarrow E$  is continuous and the set  $\{\varphi \in K; h(\varphi) > h(\theta_0)\}$  is a compact subset of  $K$ , it follows that  $h[\{\varphi \in K; h(\varphi) > h(\theta_0)\}] = \{h(\varphi); h(\varphi) > h(\theta_0), \varphi \in K\}$  is compact in  $E$ . On the other hand, by Proposition 2,  $h$  is surjective, this implies that  $\{h(\varphi); h(\varphi) > h(\theta_0)\} = \{t \in E; t > h(\theta_0)\}$  is compact, hence it is closed in  $E$ . ■

II). Let  $\varphi_0 \in B_2$ . By Lemma 6, there exists  $\varphi_1 \in B_3$  such that  $h(\varphi_0) = h(\varphi_1)$ , it is clear that  $\varphi_0 > \varphi_1$  (note that  $\psi(\varphi_0, h(\varphi_0)) = 1$  and  $\psi(\varphi_1, h(\varphi_1)) = 0$ ). On the other hand

$$\begin{aligned} \mu_1([\varphi_1, \varphi_0]) &= \mu(\{t \in E; \varphi_1 < \Pi_1(t) \leq \varphi_0\}) \\ &\leq \mu(\{t \in E; h(\varphi_1) \leq h(\Pi_1(t)) \leq h(\varphi_0)\}) \\ &= \mu(\{t \in E; h(\varphi_1) \leq t \leq h(\varphi_0)\}) = 0. \end{aligned}$$

Thus  $\varphi_1 \in [\varphi_0]$ .

Let  $\varphi \in K \setminus \{\theta', \theta'', \varphi_0, \varphi_1\}$ . Let us show that  $\varphi \notin [\varphi_0]$ .

*Case 1,  $\varphi > \varphi_0$ :* Since the restriction of  $h$  to  $B_j$  is injective,  $j = 2, 3$ ,  $h(\varphi) > h(\varphi_0)$ . Observe that

$$\begin{aligned} \mu_1([\varphi_0, \varphi]) &= \mu(\{t \in E; \varphi_0 < \Pi_1(t) \leq \varphi\}) \\ &\geq \mu(\{t \in E; h(\varphi_0) < h(\Pi_1(t)) < h(\varphi)\}) \quad (5.1) \\ &= \mu(\{t \in E; h(\varphi_0) < t < h(\varphi)\}) > 0, \end{aligned}$$

(because  $[t] = \{t\}$ , for all  $t \in E$ ) hence  $\varphi \notin [\varphi_0]$ .

*Case 2,  $\varphi < \varphi_0$ :* Let us to show that  $h(\varphi) < h_1(\varphi_0)$ . Assume that  $h(\varphi) = h_1(\varphi_0)$ , hence  $h(\varphi) = h(\varphi_0) = h(\varphi_1)$ , since the restriction of  $h$

to  $B_j$  is injective,  $j = 2, 3$  and  $\varphi \in B_0 \cup B_1$ , either  $\varphi = \varphi_0$  or  $\varphi = \varphi_1$ , which is impossible. Thus  $h(\varphi) < h(\varphi_0)$ . By a similar argument, one shows that  $\mu([\varphi, \varphi_0]) > 0$ . We deduce that  $[\varphi_0] = \{\varphi_0, \varphi_1\}$ . ■

Recall that  $\psi_{\mu_1, K}(\eta, \varphi) = f^\varphi(\eta)$ , where  $f^\varphi$  is the characteristic function of  $\{\theta \in K; \theta > \varphi\}$ . Let  $A_1$  be the  $C^*$ -subalgebra (with unit) generated by the family  $\{f^\varphi, \varphi \in K\}$  in  $L^\infty(\mu_1)$ . Note that  $A_1 = C(K_{\mu_1}(K))$ . By lemmas 2 and 4,  $h_K(\eta) = \sup\{\varphi \in K; \psi_{\mu_1, K}(\eta, \varphi) = 1\}$  exists for every  $\eta \in K(K)$  ( $\theta'_K = \max(K_{\mu_1}(K))$  and  $\theta''_K = \min(K_{\mu_1}(K))$ ).

**Corollary 13.** *The map  $h_{\mu_1, K} : K_{\mu_1}(K) \rightarrow K_\mu(E) = K$  is a homeomorphism.*

*Proof.* Consider the reference set  $\tilde{E} = K$ ,  $E_1 = B_2^{\mu, E} = B_2$ ,  $E_2 = B_3^{\mu, E} = B_3$ . By Lemma 11 and Theorem 4 we obtain that  $h_{\mu_1, K} : K_{\mu_1}(K) \rightarrow K$  is a homeomorphism (note that  $\mu_1$  satisfies the condition (1.1)). ■

Let  $\Pi_2 : K \rightarrow K_{\mu_1}(K)$  the map defined by

$$\Pi_2(\varphi) = \begin{cases} \theta_{\varphi, K} \in B_2^{\mu_1, K}, & \text{if } \varphi \in E_1 = B_2^{\mu, E} = B_2 \\ \theta'_{\varphi, K} \in B_3^{\mu_1, K}, & \text{if } \varphi \in E_2 = B_3^{\mu, E} = B_3 \\ \Pi_2(\theta'_E) = \theta'_K \text{ and } \Pi_2(\theta''_E) = \theta''_K, & \end{cases}$$

Where  $\theta_{\varphi, K}, \theta'_{\varphi, K}$  the elements founded by lemma 6. Note that  $\Pi_2(B_j) = B_j^{\mu_1, K}$ ,  $j \in \{2, 3\}$ .

**Lemma 12.** *The map  $\Pi_2 : K \rightarrow K_{\mu_1}(K)$  is Borel.*

*Proof.* By Proposition 10,  $K_{\mu_1}(K)$  is a strongly Lindelöf space, it suffices then to show that  $(\Pi_2)^{-1}(V_\theta), (\Pi_2)^{-1}(V'_\theta)$  are Borel subsets of  $K$ , where  $V_\theta = \{\eta \in K_{\mu_1}(K); \psi_{\mu_1, K}(\eta, \theta) = 1\}$  and  $V'_\theta = \{\eta \in K_{\mu_1}(K); \psi_{\mu_1, K}(\eta, \theta) = 0\}$ . We can suppose that  $\theta \in B_2$ . We have

$$\begin{aligned} (\Pi_2)^{-1}(V_\theta) &= \{\varphi \in B_2; \psi_{\mu_1, K}(\theta_{\varphi, K}, \theta) = 1\} \\ &\quad \cup \{\varphi \in B_3; \psi_{\mu_1, K}(\theta'_{\varphi, K}, \theta) = 1\} \cup \{\theta'\} \\ &= \{\varphi \in B_2; \theta \leq h_{\mu_1, K}(\theta_{\varphi, K})\} \\ &\quad \cup \{\varphi \in B_3; \theta < h_{\mu_1, K}(\theta'_{\varphi, K})\} \cup \{\theta'\} \\ &= \{\varphi \in B_2; \theta \leq \varphi\} \cup \{\varphi \in B_3; \theta < \varphi\} \cup \{\theta'\} \\ &= \{\varphi \in K; \theta \leq \varphi\}. \end{aligned}$$

Thus  $(\Pi_2)^{-1}(V_\theta)$  is a Borel subset of  $K$ . By a similar argument, one shows that  $(\Pi_2)^{-1}(V'_\theta)$  is a Borel subset of  $K$ . ■

Let  $\mu_2 = \Pi_2(\mu_1)$ .

**Lemma 13.** *Suppose that  $\mu([t \wedge t', t \vee t']) > 0$ , for every  $t \neq t'$ . Let  $\eta_0 \in B_2^K$  and  $\eta_1 \in B_3^K$ . Then:*

I) *The sets  $\{\eta \in K_{\mu_1}(K); \eta > \eta_0\}$ ,  $\{\eta \in K_{\mu_1}(K); \eta < \eta_1\}$  are not closed subsets of  $K_{\mu_1}(K)$ .*

II) *For every  $\lambda_0 \in B_2^{\mu_1, K}$ , there exists  $\lambda_1 \in B_3^{\mu_1, K}$  such that  $[\lambda_0] = \{\lambda_0, \lambda_1\}$  and  $h(h_{\mu_1, K}(\lambda_0)) = h(h_{\mu_1, K}(\lambda_1))$ .*

*Proof.*

I). Let us show that  $\{\eta \in K_{\mu_1}(K); \eta > \eta_0\}$  is not closed. Suppose that there exists  $\eta \in K_{\mu_1}(K)$  such that  $\eta > \eta_0$ . There exists  $\theta \in K$  such that  $\psi_{\mu_1, K}(\eta, \theta) = f^\theta(\eta) = 1$  and  $\psi_{\mu_1, K}(\eta_0, \theta) = f^\theta(\eta_0) = 0$ . Since  $\eta_0 \in B_2^{\mu_1, K}$ ,  $h_{\mu_1, K}(\eta) \geq \theta > h_{\mu_1, K}(\eta_0)$ . It follows that  $h_K(\eta) > h_K(\eta_0)$ . We conclude that  $\{\eta \in K(K); \eta > \eta_0\} = \{\eta \in K(K); h_{\mu_1, K}(\eta) > h_{\mu_1, K}(\eta_0)\}$ .

Assume now that  $\{\eta \in K_{\mu_1}(K); \eta > \eta_0\} = \{\eta \in K_{\mu_1}(K); h_{\mu_1, K}(\eta) > h_{\mu_1, K}(\eta_0)\}$  is closed. The map  $h_{\mu_1, K} : K_{\mu_1}(K) \rightarrow K$  is a homeomorphism by Corollary 13, it follows that

$h_{\mu_1, K}[\{\eta \in K_{\mu_1}(K); h_{\mu_1, K}(\eta) > h_{\mu_1, K}(\eta_0)\}] = \{h_{\mu_1, K}(\eta); h_K(\eta) > h_{\mu_1, K}(\eta_0), \eta \in K_{\mu_1}(K)\}$  is a compact subset of  $K$ . Since  $h_{\mu_1, K}$  is onto, the set

$\{h_{\mu_1, K}(\eta) \in K; h_{\mu_1, K}(\eta) > h_{\mu_1, K}(\eta_0), \eta \in K_{\mu_1}(K)\} = \{\varphi \in K; \varphi > h_{\mu_1, K}(\eta_0)\}$

is closed in  $K$ . On the other hand, by Remark 16,  $h_{\mu_1, K}(\eta_0) \in B_2$ , using Lemma 11-I) we obtain that  $\{\varphi \in K; \varphi > h_{\mu_1, K}(\eta_0)\}$  is not closed, which is impossible. Thus  $\{\eta \in K_{\mu_1}(K); \eta > \eta_0\}$  is not closed. By a similar argument, one shows that the set  $\{\eta \in K_{\mu_1}(K); \eta < \eta_1\}$  is not closed. ■

II). Let  $\lambda_0 \in B_2^{\mu_1, K}$ . Since  $h_3 : B_3 \rightarrow E \setminus \{-\infty, +\infty\}$  is onto, there exists  $\theta_1 \in B_3$  such that

$$h(\theta_1) = h(h_{\mu_1, K}(\lambda_0)). \quad (5.2)$$

On the other hand, by Corollary 13 and Remark 16, there exists  $\lambda_1 \in B_3^{\mu_1, K}$  such that  $h_{\mu_1, K}(\lambda_1) = \theta_1$ , this implies by 5.2 that

$$h(h_{\mu_1, K}(\lambda_0)) = h(h_{\mu_1, K}(\lambda_1)). \quad (5.3)$$

We shall prove that  $\lambda_1 < \lambda_0$ .

Suppose that  $\lambda_1 \geq \lambda_0$ . It follows that  $h_{\mu_1, K}(\lambda_1) = \theta_1 \geq h_{\mu_1, K}(\lambda_0)$ . Since  $\theta_1 \in B_3$ ,  $\psi(\theta_1, h(\theta_1)) = 0$ , hence  $\psi_{\mu_1, K}(h_{\mu_1, K}(\lambda_0), h(\theta_1)) = 0$ , this implies that  $h(h_{\mu_1, K}(\lambda_0)) < h(\theta_1)$  because by Corollary 13 and Remark 16,  $h_{\mu_1, K}(\lambda_0) \in B_2$ , it is impossible by (5.2) and (5.3).

Thus  $\lambda_1 < \lambda_0$ .

Let us show that  $\lambda_1 \in [\lambda_0]$ .

Observe that

$$\begin{aligned}
\mu_2([\lambda_0, \lambda_1]) &= \mu_2(\{\eta \in K_{\mu_1}(K); \lambda_0 < \eta \leq \lambda_1\}) \\
&= \mu_1(\{\theta \in K; \lambda_0 < \Pi_2(\theta) \leq \lambda_1\}) \\
&\leq \mu_1(\{\theta \in K; h_{\mu_1, K}(\lambda_0) \leq h_{\mu_1, K}(\Pi_2(\theta)) \leq h_{\mu_1, K}(\lambda_1)\}) \\
&= \mu_1(\{\theta \in K; h_{\mu_1, K}(\lambda_0) \leq \theta \leq h_{\mu_1, K}(\lambda_1)\}) \\
&= \mu(\{t \in E; h_{\mu_1, K}(\lambda_0) \leq \theta_t \leq h_{\mu_1, K}(\lambda_1)\}) \\
&\leq \mu(\{t \in E; h(h_{\mu_1, K}(\lambda_0)) \leq h(\theta_t) \leq h(h_{\mu_1, K}(\lambda_1))\}) \\
&= \mu(\{t \in E; h(h_{\mu_1, K}(\lambda_0)) \leq t \leq h(h_{\mu_1, K}(\lambda_1))\}) = 0.
\end{aligned}$$

We conclude that  $\lambda_0 \in [\lambda_1]$ . Let us show that  $\eta \in K_{\mu_1}(K) \setminus \{\theta'_K, \theta''_K, \lambda_0, \lambda_1\}$ . We show that  $\eta \notin [\lambda_0]$ .

*Case 1:  $\eta > \lambda_0$ .*

*Step 1:* We shall show that  $h(h_{\mu_1, K}(\eta)) > h(h_{\mu_1, K}(\lambda_1)) = h(h_{\mu_1, K}(\lambda_0))$ .

Indeed, suppose that  $h(h_{\mu_1, K}(\eta)) = h(h_{\mu_1, K}(\lambda_1)) = h(h_{\mu_1, K}(\lambda_0))$ . By Corollary 13 and Remark 16,  $h_{\mu_1, K}(\lambda_0) \in B_2$  and  $h_{\mu_1, K}(\lambda_1) \in B_3$ . We distinguish two cases:

*Case 1-a,  $\eta \in B_2^{\mu_1, K}$ :* By Corollary 13 and Remark 16,  $h_{\mu_1, K}(\eta) \in B_2$ . On the other hand, the restriction of  $h$  to  $B_2$  is injective, hence  $h_{\mu_1, K}(\eta) = h_{\mu_1, K}(\lambda_0)$ . This is impossible because the restriction of  $h_{\mu_1, K}$  to  $B_2^{\mu_1, K}$  is injective by Remark 2.

*Case 1-b,  $\eta \in B_3^{\mu_1, K}$ :* Since  $h_{\mu_1, K}(\eta) \in B_3$  and the restriction of  $h$  to  $B_3$  is injective,  $h_{\mu_1, K}(\eta) = h_{\mu_1, K}(\lambda_1)$ , this implies that  $\eta = \lambda_1$ , this is impossible (we have shown previously that  $\lambda_0 > \lambda_1$ ). We conclude that  $h(h_{\mu_1, K}(\eta)) > h(h_{\mu_1, K}(\lambda_1)) = h(h_{\mu_1, K}(\lambda_0))$ . ■

*Step 2:* Let us show that  $\mu_2([\lambda_0, \eta]) > 0$ .

We have

$$\begin{aligned}
\mu_2([\lambda_0, \eta]) &= \mu_2(\{\omega \in K_{\mu_1}(K); \lambda_0 < \omega < \eta\}) \\
&\geq \mu_2(\{\omega \in K_{\mu_1}(K); h_K(\lambda_0) < h_{\mu_1, K}(\omega) < h_{\mu_1, K}(\eta)\}) \\
&= \mu_1(\{\theta \in K; h_{\mu_1, K}(\lambda_0) < h_{\mu_1, K}(\Pi_2(\theta)) < h_{\mu_1, K}(\eta)\}) \\
&= \mu_1(\{\theta \in K_{\mu_1}(K); h_{\mu_1, K}(\lambda_0) < \theta < h_{\mu_1, K}(\eta)\}) \\
&\geq \mu_1(\{\theta \in K_{\mu_1}(K); h_K(h_{\mu_1, K}(\lambda_0)) < h(\theta) < h(h_{\mu_1, K}(\eta))\}) \\
&= \mu(\{t \in E; h(h_{\mu_1, K}(\lambda_0)) < h(\theta_t) \leq h(h_{\mu_1, K}(\eta))\}) \\
&= \mu(\{t \in E; h(h_{\mu_1, K}(\lambda_0)) < t < h(h_{\mu_1, K}(\eta))\}) > 0. \blacksquare
\end{aligned}$$

Thus  $\eta \notin [\lambda_0]$ .

*Case 2,  $\eta < \lambda_0$ :* By a similar argument, one shows that  $\eta \notin [\lambda_0]$ . It follows that  $[\lambda_0] = \{\lambda_0, \lambda_1\}$ . ■

Let  $A_2$  be the  $C^*$ -subalgebra which is generated by the family  $\{f^\eta; \eta \in K_{\mu_1}(K)\}$  (where  $f^\eta$  the characteristic function of  $\{\xi \in K_{\mu_1}(K); \xi > \eta\}$ ) in  $L^\infty(K_{\mu_1}(K), \mu_2)$ , ( $A_2 = C(K_{\mu_2}(K_{\mu_1}(K)))$ ).

Recall that  $B_1^{\mu_2, K_{\mu_1}(K)}$  is the set of  $\xi \in K_{\mu_2}(K_{\mu_1}(K))$  such that the map  $\eta \in (K_{\mu_1}(K)) \setminus \{\theta'_K, \theta''_K\} \rightarrow \psi_{\mu_2, K_{\mu_1}(K)}(\xi, \eta) = \xi(f^\eta)$  is continuous and  $B_j^{\mu_2, K_{\mu_1}(K)} = \left\{ \xi \in K_{\mu_2}(K_{\mu_1}(K)) \setminus B_1^{\mu_2, K_{\mu_1}(K)}; \psi_{\mu_1, K}(\xi, h_{K_{\mu_1}(K)}(\xi)) = 3 - j \right\}$ ,  $j = 2, 3$ , where  $h_{\mu_2, K_{\mu_1}(K)}(\xi) = \sup \left\{ \eta \in K_{\mu_1}(K); \psi_{\mu_2, K_{\mu_1}(K)}(\xi, \eta) = 1 \right\}$ .

**Corollary 14.** *The map  $h_{\mu_2, K_{\mu_1}(K)} : K_{\mu_2}(K_{\mu_1}(K)) \rightarrow K_{\mu_1}(K)$  is a homeomorphism.*

*Proof.* Consider the reference set  $\tilde{E} = K_{\mu_1}(K)$ ,  $(E)_1 = B_2^{\mu_2, K_{\mu_1}(K)}$ ,  $E_2 = B_3^{\mu_2, K_{\mu_1}(K)}$ . By Theorem 4 and Lemma 13,  $h_{\mu_2, K_{\mu_1}(K)} : K_{\mu_2}(K_{\mu_1}(K)) \rightarrow K_{\mu_1}(K)$  is a homeomorphism. ■

## 6. GENERAL PROPERTIES OF $(C(L), \tau_p)$ WHEN L IS A ROSENTHAL COMPACT SET

In this part, we show that the function  $\psi : (K \times (E, \tau_0)) \rightarrow \{0, 1\}$  is not Borel under some assumptions. Let  $L$  be Hausdorff compact. In [27] one shows under the Martin's Axiom that  $(C(L), \tau_p)$  is universally measurable. During this part, we show that  $(C(L), \tau_p)$  is universally measurable without Martin's Axioms, if  $L$  is a Rosenthal compact set. Finally, we show that if  $K$  is separable, then  $(C(K, \tau_p)$  is not measure-compact.

**Definition 6.** *Let  $(X, \tau)$  be a Hausdorff space. We say that  $(X, \tau)$  is a Radon space [21, p. 117], if for every positive finite Borel measure on  $(E, \tau)$  is a Radon measure.*

**Definition 7.** *Let  $(X, \tau)$  be a Hausdorff space. A positive finite Borel measure  $\nu$  on  $(X, \tau)$  is said to be normal if for every net  $(O_\alpha)$  of open sets is increasing to open set  $O$ ,  $\lim_\alpha \nu(O_\alpha) = \nu(O)$ .*

**Remark 17.** *Let  $(X, \tau)$  be a strongly Lindelöf space. Then every positive Borel measure on  $(X, \tau)$  is normal.*

**Remark 18.** *Let  $(X, \tau)$  be a Hausdorff topological space and let  $X_1 \subset X$ . Then  $\text{Bor}(X_1, \tau) = \{C \cap X_1; C \in \text{Bor}(X, \tau)\}$ .*

**Remark 19.** *Let  $L$  be a compact strongly Lindelöf space. Then  $K$  is a Radon space.*

*Proof.* As  $L$  is a strongly Lindelöf space, by Remark 17, every positive finite Borel measure on  $K$  is normal. On the other hand,  $K$  is universally measurable, by [22, th. 3.2] every normal positive finite Borel measure on  $K$  is a Radon measure, hence  $L$  is a Radon-space. ■

**Proposition 12.** *Let  $(E, \tau_0)$  be a strongly Lindelöf space such that  $-\infty, +\infty \notin E$ ; let  $E'$  be a subset of  $E$  such that  $(E', \tau_1)$  be completely regular space. Assume that there exists a measure  $\nu$  strictly positive on  $(E', \tau_0)$  such that  $\nu$  is diffuse on  $E$ . Then  $(E', \tau_1)$  is not universally measurable space.*

*Proof.* By Lemma 8,  $\text{Bor}(E, \tau) = \text{Bor}(E, \tau_1)$ . This implies by Remark 18 that  $\text{Bor}(E', \tau_0) = \text{Bor}(E', \tau_1)$ . Hence, there exists  $V \in \text{Bor}(E', \tau_1)$  such that  $0 < \nu(V) < +\infty$ . One defines the measure  $\nu_1$  on  $E'$  by,  $\nu_1(U) = \nu(U \cap V)$ ,  $U \in \text{Bor}(E', \tau_1)$ .  $\nu_1$  is a finite Borel measure on  $(E', \tau_1)$ . On the other hand, by Lemma 8,  $(E, \tau_1)$  is a strongly Lindelöf space, hence  $(E', \tau_1)$  has the same property by Remark 5, we conclude by Remark 17 that  $\nu_1$  is normal. Suppose now that  $(E', \tau_1)$  is universally measurable. By [22, th. 3.2],  $\nu_1$  is a Radon measure. It follows that there exists a compact set  $L_0$  of  $(E', \tau_1)$  such that  $\nu_1(L_0) > 0$ . Put  $x_0 = \max \{t; t \in L_0\}$ . By hypothesis on  $\nu$ , we have  $\nu_1(L_0 - \{x_0\}) > 0$ , then there exists a compact set  $L_1$  of  $L_0 - \{x_0\}$  satisfying  $\nu_1(L_1) > 0$ . Denote  $x_1 = \max \{t; t \in L_1\}$ . By induction one constructs a strictly decreasing sequence  $(x_n)_{n \geq 0}$  of  $L_0$ . Let  $\mathcal{U}$  be a non trivial ultrafilter on  $\mathbb{N}$ . Since  $L_0$  is compact, there exists  $a \in L_0$  such that  $x_n \xrightarrow{\mathcal{U}} a$  in  $(E', \tau_1)$ . Let  $O = \{t \in E'; t \leq a\}$ .  $O$  is an open neighborhood of  $a$ , this implies that  $\{n \in \mathbb{N}; x_n \in O\} = \{n \in \mathbb{N}; x_n \leq a\} \in \mathcal{U}$ . It is impossible because  $x_k \geq a$ , for every  $k \in \mathbb{N}$  and the sequence  $(x_n)_{n \geq 0}$  is strictly decreasing. ■

By Remark 11 and Proposition 12, one has the following corollary:

**Corollary 15.** *Let  $(E, \tau_0)$  be a strongly Lindelöf space. Assume that  $\nu$  strictly positive on  $(E, \tau_0)$  and diffuse on  $E$ ,  $(E, \leq)$  is complete and  $-\infty, +\infty \notin E$ . Then  $(E, \tau_1)$  is not universally measurable.*

Recall that

$$E_2 = \{a \in E; \{x \in E; x < a\} \text{ is not a closed subset of } E\}.$$

**Corollary 16.** *Assume that  $(E, \tau_0)$  is a strongly Lindelöf space, that  $-\infty, +\infty \notin E$ , that  $E_2 = E$ , the measure  $\mu$  is diffuse on  $E$  and that  $\mu(\{t \wedge t', t \vee t'\}) > 0$  for  $t \neq t'$ . Then  $B_3$  is not universally measurable.*

*Proof.* By Remark 12,  $(E, \tau_1)$  is completely regular. On the other hand Proposition 12 shows us that  $(E, \tau_1)$  is not universally measurable and by step 2 of Proposition 8-I),  $B_3$  is homeomorphic to  $(h(B_3), \tau_1)$ . But  $h(B_3) = E$  by Lemma 6, we conclude that  $B_3$  is not universally measurable. ■

**Remark 20.** *In Corollary 16, we can replace the hypothesis  $E_2 = E$ , by the hypothesis  $h_3 : B_3 \rightarrow E$  is onto.*

**Corollary 17.** *Let  $(E, \tau_0)$  be a strongly Lindelöf space, and suppose that  $-\infty, +\infty \notin E$ . Assume that there exists a diffuse measure  $\nu$  on  $E$  and there exists a compact set  $L_0$  of  $(E, \tau_0)$  verifying  $0 < \nu(L_0) < +\infty$ . Then  $(E, \tau_1)$  does not embed in any analytic set.*

*Proof.* Consider the identity map  $i : (E, \tau_0) \rightarrow (E, \tau_1)$ . By Lemma 8,  $i$  is Borel. Suppose now that there exists an analytic set  $Y$  such that  $(E, \tau_1)$  is a topological subspace of  $Y$ . Consider the measure  $\nu_1$  defined on  $L_0$  by  $\nu_1(B) = \nu(B)$ ,  $B \in \text{Bor}(L_0)$ . Since  $(L_0, \tau_0)$  is a strongly Lindelöf space,  $\nu_1$  is normal. On the other hand  $(L_0, \tau_0)$  is universally measurable, by [22, th.3.2]  $\nu_1$  is a Radon measure. By [23, th. 14], there is a compact subset  $L_1$  of  $L_0$  of measure strictly positive such that the restriction of  $i$  to  $L_1$  is continuous (with values in  $Y$ ). It follows that  $i(L_1) = L_1$  is compact in  $(E, \tau_1)$  and  $\nu_1(L_1) > 0$ . using Proposition 12 for  $E' = L_1$ , we obtain that  $L_1$  is not universally measurable, which is impossible. ■

**Corollary 18.** *Let  $(E, \tau_0)$  be a strongly Lindelöf such that  $-\infty, +\infty \notin E$ . Suppose that  $h_3$  is onto and the hypothesis on  $\nu$  in corollary 17 are satisfied for  $\nu = \mu$ . Then  $B_3$  does not embed in any analytic set.*

*Proof.* By Proposition 8-I)  $B_3$  is homeomorphic to  $(E, \tau_1)$ . For getting the corollary it suffices to use Corollary 17.

**Proposition 13.** *Suppose that  $(E, \tau_0)$  is separable, satisfying the condition (\*), and the measure  $\mu$  is diffuse on  $E$ ,  $h_3 : B_3 \rightarrow E$  is surjective,  $-\infty, +\infty \notin E$  and  $\mu(\{t \wedge t', t \vee t'\}) > 0$  for  $t \neq t'$ . Then the map  $\psi : (K \times (E, \tau_0)) \rightarrow \{0, 1\}$  is not Borel.*

*Proof.*

*Step 1:* Let us show that  $\text{Bor}(K) \otimes \text{Bor}(E) = \text{Bor}(K \times E)$ .

It is obvious that  $\text{Bor}(K) \otimes \text{Bor}(E) \subset \text{Bor}(K \times E)$ . We show the reverse inclusion. Let  $(O_k)_{k \geq 0}$  be a countable basis of  $(E, \tau_0)$  (a such basis exists by Remark 1) and let  $V = \bigcup_{i \in I} U_i \times W_i$ , where the  $U_i$  are open subsets of  $K$  and the  $W_i$  are open subsets of  $E$ . For every  $i \in I$ ,

there exists a subset  $M_i$  of  $\mathbb{N}$  such that  $W_i = \bigcup_{k \in M_i} O_k$ , hence  $V = \bigcup_{k \in \mathbb{N}} \left[ \left( \bigcup_{i \in Y_k} U_i \right) \times O_k \right]$ , where  $Y_k = \{i; k \in M_i\}$ . Since  $\bigcup_{i \in Y_k} U_i$  is an open subset of  $K$ , then

$$V \in \text{Bor}(K) \otimes \text{Bor}(E).$$

It follows that  $\text{Bor}(K \times E) \subset \text{Bor}(K) \otimes \text{Bor}(E)$ . ■

*Step 2:* Let us show that every Borel subset of  $K$  is universally measurable.

By Corollary 7,  $K$  is a strongly Lindelöf space, using Remark 19, one obtains that  $K$  is a Radon space. We conclude that every Borel subset of  $K$  is relatively universally measurable in  $K$ , by [22, th. 3.2] every Borel subset of  $K$  is universally measurable. ■

Suppose now that  $\psi$  is Borel. Consider the map  $\sigma : K \setminus B_1 \rightarrow K \times E$  defined by  $\sigma(\theta) = (\theta, h(\theta))$ ,  $\theta \in K \setminus B_1$ . This map  $\sigma$  is Borel because  $\text{Bor}(K) \otimes \text{Bor}(E) = \text{Bor}(K \times E)$  and  $h$  is continuous by Proposition 3. It follows that  $\psi \circ \sigma$  is Borel, hence  $(\psi \circ \sigma)^{-1}\{0\} = B_3$  is Borel subset of  $K$ . By the step 2,  $B_3$  is universally measurable, this is impossible by Remark 20. ■

Let  $(Y, \tau)$  be a Hausdorff topological space. We denote by  $\text{Bair}(Y)$  the  $\sigma$ -algebra generated by the continuous functions on  $Y$  and by  $P_\sigma(Y)$  the set of probability measures on  $(Y, \text{Bair}(Y))$ .

**Definition 8.** ([16]-[17]-[28]) *Let  $X$  be a completely regular space:*

a)  *$X$  is measure-compact, if for every measure  $\nu \in P_\sigma(X)$  and every net  $(f_\alpha)_{\alpha \in I}$  of bounded continuous decreasing functions on  $(X, \tau)$  such that  $f_\alpha \rightarrow 0$ ,  $\lim_{\alpha} \int_X f_\alpha d\nu = 0$ .*

b)  *$X$  is strongly measure-compact, if for every measure  $\nu \in P_\sigma(X)$  and every  $\varepsilon > 0$ , there is a compact  $L$  of  $X$  such that  $\nu^*(L) > 1 - \varepsilon$ , where*

$$\nu^*(L) = \inf \{ \nu(H); H \in \text{Bair}(X) \text{ and } L \subset H \}.$$

Note that if  $X$  is strongly measure-compact, then  $X$  is measure-compact.

**Remark 21.** *Suppose that  $(E, \leq)$  satisfies the condition (\*). Then  $(E, \tau_0)$  and  $(E, \tau_1)$  are regular spaces.*

*Proof.* Let us show for example that  $(E, \tau_1)$  is regular. Indeed, let  $V = ]a, b]$  an open neighborhood of  $x$ . Since  $(E, \leq)$  verifies the condition (\*), there is  $c \in E$  such that  $a < c < x \leq b$ , hence  $[c, b] = \overline{]c, b]} \subset V$  and  $]c, b]$  is an open neighborhood of  $x$ . We can use the same argument if  $V = ]a, b[$ ,  $a < b$ . ■

**Proposition 14.** *Let  $(E, \tau_0)$  be a separable strongly Lindelöf space satisfying the condition (\*). Suppose there is a probability measure  $\nu$  on  $(E, \tau_0)$ , diffuse on  $E$  and  $-\infty, +\infty \notin E$ . Then  $(E, \tau_1)$  is not strongly measure-compact.*

*Proof.* Since  $E$  satisfies (\*),  $E = E_2$ . By Remark 12,  $(E, \tau_1)$  is completely regular.

*Step 1:* Show that every open subset of  $(E, \tau_1)$  is in  $\text{Bair}(E, \tau_1)$ .

Pick  $a \in E$ . The set  $\{t \in E; t \leq a\}$  is open and closed in  $(E, \tau_1)$ , hence the characteristic function of  $\{t \in E; t \leq a\}$  is continuous on  $(E, \tau_1)$ . It follows that  $\{t \in E; t \leq a\} \in \text{Bair}(E, \tau_1)$ .

Since  $(E, \tau_0)$  is strongly Lindelöf and regular (by Remark 21), by [21, chap. II, prop. 1],  $\{a\}$  is  $G_\delta$  of  $(E, \tau_0)$ . Thus there exists a sequence of open subsets  $(O_n)_{n \geq 0}$  of  $(E, \tau_0)$  such that  $\{a\} = \bigcap_{n \geq 0} O_n$ . For every  $n \geq 0$ , there is an open interval  $I_n$  of  $O_n$  such that  $a \in I_n \subset O_n$ , this implies that there exists  $a_n \in E$  such that  $]a_n, a[ \subset I_n \subset O_n$ . Therefore  $\{a\} = \bigcap_{n \geq 0} ]a_n, a[$ .

On the other hand,  $]a_n, a[ = \{t \in E; t \leq a\} - \{t \in E; t \leq a_n\} \in \text{Bair}(E, \tau_1)$ . It follows that  $\{a\} \in \text{Bair}(E, \tau_1)$ . Thus for every  $a \in E$ ,  $\{t \in E; t < a\} = \{t \in E; t \leq a\} - \{a\} \in \text{Bair}(E, \tau_1)$ .

Since  $(E, \tau_1)$  is strongly Lindelöf, by Lemma 8, every open subset of  $(E, \tau_1)$  is in  $\text{Bair}(E, \tau_1)$ . ■

*Step 2:* Let  $L$  be a compact subset of  $(E, \tau_1)$ . Let us show that  $\nu(L) = 0$ .

Suppose  $\nu(L) > 0$ . By Proposition 12,  $L$  is not universally measurable, which is impossible. ■

*Step 3:* Let us show that  $(E, \tau_1)$  is not strongly measure-compact.

Note first by step 1 that  $\text{Bor}(E, \tau_1) = \text{Bair}(E, \tau_1)$ , hence  $\nu \in P_\sigma(E, \tau_1)$ .

Suppose now that  $(E, \tau_1)$  is a strongly measure-compact. Then there exists a compact  $L$  subset of  $(E, \tau_1)$  such that  $(\nu)^*(L) > 0$ . By step 1,  $L \in \text{Bair}(E, \tau_1)$ , this implies that  $\nu(L) = \nu^*(L) > 0$ , this is impossible by step 2. ■

Note that if  $X$  a Lindelöf space, then  $X$  is measure-compact [28]. In [16]-[17] one shows that there exists a measure-compact space which is not strongly measure-compact. The following remark gives another type of example.

**Remark 22.** *There exists a Lindelöf space  $Y$ , separable paracompact which is not strongly measure-compact.*

*Proof.* It suffices to apply Proposition 14 to  $E = ]0, 1[$ ,  $\nu$  is the Lebesgue measure, hence  $Y = (]0, 1[, \tau_1)$  is not strongly measure-compact. ■

Let  $L$  be a Hausdorff compact space. In [27] one shows under the Martin's Axiom that  $(C(L), \tau_p)$  is universally measurable. In the following Lemma, one shows that  $(C(L), \tau_p)$  is universally measurable without Martin's Axioms if  $L$  is a Rosenthal compact set.

**Proposition 15.** *Let  $L$  be a Rosenthal compact set. Then  $(C(L), \tau_p)$  is universally measurable.*

*Proof.* We shall show Lemma 15 if  $C(L)$  is the real continuous functions space. The proof is similar when one replaces  $\mathbb{R}$  by  $\mathbb{C}$ . By [22, th. 3.2] it suffices to show that every normal probability measure on  $(C(L), \tau_p)$  is a Radon measure. Let  $\nu$  a normal probability measure on  $(C(L), \tau_p)$ .

Denote, for every  $n \in \mathbb{N}^*$ ,  $B_n = \{g \in C(L); \|g\|_{C(L)} \leq n\}$  and  $\nu_n$  the measure defined by  $\nu_n(C) = \nu(C \cap B_n)$ ,  $C \in \text{Bor}(C(L), \tau_p)$ . Since  $\nu_n(C) \rightarrow_{n \rightarrow +\infty} \nu(C)$  and  $B_n$  is  $\tau_p$  closed, it is enough to show that  $\nu_n$  is a Radon measure for all  $n \geq 1$ .

*Step 1:* Let us show that for every  $n \geq 1$ ,  $\nu_n$  is a normal measure.

Let  $n \in \mathbb{N}^*$  and let  $(F_\alpha)$  be a net of closed subsets of  $(C(L), \tau_p)$  which is decreasing to a closed set  $F$  of  $(C(L), \tau_p)$ . Since  $B_n$  is  $\tau_p$ -closed, the net  $(F_\alpha \cap B_n)$  is decreasing to  $B_n \cap F$  and  $\nu$  is normal, it follows that  $\lim_{\alpha} \nu_n(F_\alpha) = \lim_{\alpha} \nu(F_\alpha \cap B_n) = \nu(F \cap B_n) = \nu((\bigcap_{i \in I} F_i) \cap B_n) = \nu_n(F)$ .

■

*Step 2:* Pick  $n \in \mathbb{N}$ . Let us show that  $\nu_n$  is a Radon measure.

We can suppose that  $\text{supp}(\nu_n)$  is included in the closed unit ball of  $(C(L), \|\cdot\|)$ . Denote  $V$  the vector subspace spanned by  $L$  in the dual of  $(C(L), \|\cdot\|)$ . For every  $v \in V$  let  $H_v$  the hyperplane defined by  $\nu$  in  $C(L)$ , i.e.

$$H_v = \{f \in C(L); (f, v) = v(f) = 0\}.$$

Consider the set  $I'$  formed of all elements  $v$  of  $V$  such that  $\nu_n(H_v) = 1$  and  $E_{\nu_n}$  is the intersection of  $H_v$  when  $v \in I'$  (note that  $0 \in I'$ ).

Let us show that  $\nu_n(E_{\nu_n}) = 1$ . Let  $\tilde{I} = \{S; S \text{ is a finite subset of } I'\}$  and  $H_S = \bigcap_{v \in S} H_v$ ,  $S \in \tilde{I}$ . It is clear that  $(H_S)_{S \in \tilde{I}}$  is a net of decreasing closed subsets of  $(C(L), \tau_p)$ ; since  $\nu_n$  is normal,

$$\nu_n(E_{\nu_n}) = \nu_n\left(\bigcap_{v \in I'} H_v\right) = \nu_n\left(\bigcap_{S \in \tilde{I}} H_S\right) = \inf_{S \in \tilde{I}} \nu_n(H_S) = 1,$$

because  $\nu_n(H_S) = 1$ , for every  $S \in \tilde{I}$ . For  $y \in L$  put

$$[y] = \{x \in L; (f, y) = (f, x), \forall f \in E_{\nu_n}\}$$

and  $L_{\nu_n} = \{[y]; y \in L\}$  ( $L_{\nu_n}$  as a topological subspace of  $[-1, +1]^{E_{\nu_n}}$ , where  $[y](f) = (f, y)$ ,  $f \in E_{\nu_n}$ ). We shall show that  $L_{\nu_n}$  is compact.

It is enough to show that  $L_{\nu_n}$  is closed in  $[-1, +1]^{E_{\nu_n}}$ . Let  $([y_\alpha])_{\alpha \in I}$  be a net in  $L_{\nu_n}$  such that  $[y_\alpha] \rightarrow z \in [-1, +1]^{E_{\nu_n}}$ . Since  $L$  is compact, there exists  $y \in L$  such that  $y_\alpha \rightarrow y \in L$ , this implies that  $[y_\alpha](f) = (f, y_\alpha) \rightarrow (f, y) = [y](f)$  for every  $f \in E_{\nu_n}$ . On the other hand,  $[y_\alpha](f) = (f, y_\alpha) \rightarrow z(f)$ , it follows that  $z(f) = [y](f)$  for every  $f \in E_{\nu_n}$ . We conclude that  $z = [y]$ .

*Step 2-a):* Suppose that  $[y] = [y']$ ,  $\nu_n$ -almost-everywhere,  $(y, y' \in L)$  and show that  $[y] = [y']$  everywhere.

Indeed, for almost all  $f \in C(L)$ ,  $(y, f) = (y', f)$ . Thus  $(y - y', f) = 0$  for almost all  $f \in C(L)$ , i.e. the hyperplane  $H_{y-y'}$  is of measure one, hence  $H_{y-y'}$  contains  $E_{\nu_n}$ . We deduce that for every  $f \in E_{\nu_n}$ ,  $(f, y) = (f, y')$ . ■

We define the map  $\Sigma : L \rightarrow L_{\nu_n}$ , by  $\Sigma(y) = [y]$ ,  $y \in L$  (it is clear that  $\Sigma$  is continuous) and the map  $U : L_{\nu_n} \rightarrow L^1(E_{\nu_n}, \nu_n)$  by  $U([x])(f) = f(x)$ ,  $x \in L$ ,  $f \in E_{\nu_n}$ . Note that  $U$  is into by step 3.

*Step 2-b):* Show that  $U$  is continuous. For that let  $F$  be a closed subset of  $L^1(E_{\nu_n}, \nu_n)$  and let  $[x]$  be a point in the closure of  $U^{-1}(F)$ . There exists a net  $([x_\alpha])_{\alpha \in I}$  in  $U^{-1}(F)$  such that  $[x_\alpha] \rightarrow [x]$ . Since  $L$  is compact, there exists  $x' \in L$  such that  $x_\alpha \rightarrow x'$  in  $L$ , by continuity of  $\Sigma$ ,  $[x_\alpha] \rightarrow [x']$  in  $L_{\nu_n}$ . It follows that  $[x'] = [x]$ .

On the other hand, by [3]  $L$  is angelic, hence there is a countable subsequence  $(x_{i_k})_{k \geq 0}$  such that  $x_{i_k} \rightarrow_{k \rightarrow +\infty} x'$ . By the continuity of  $\Sigma$  we have  $[x_{i_k}] \rightarrow_{k \rightarrow +\infty} [x'] = [x]$  in  $L_{\nu_n}$ . By dominated convergence theorem,  $U([x_{i_n}]) \rightarrow_{n \rightarrow +\infty} U([x])$  in  $L^1(E_{\nu_n}, \nu_n)$ . Since  $U([x_{i_k}]) \in F$ , for every  $k \in \mathbb{N}$ ,  $U([x]) \in F$ . It follows that  $U$  is continuous. ■

Thus  $L_{\nu_n}$  is metrisable, because  $U$  is a homeomorphism on its range. Therefore,  $(C(L_{\nu_n}), \|\cdot\|)$  is a Polish space, and  $(C(L_{\nu_n}), \tau_p)$  is defined by a separating family of seminorms, by [2, TG. 9, 68-6, th. 14]  $\text{Bor}(C(L_{\nu_n}), \|\cdot\|) = \text{Bor}(C(L_{\nu_n}), \tau_p)$ . Since  $(C(L_{\nu_n}), \|\cdot\|)$  is a Radon space [21, chap. II, th. 9] and  $(E_{\nu_n}, \tau_p)$  is a closed subspace of  $(C(L_{\nu_n}), \tau_p)$ ,  $(E_{\nu_n}, \tau_p)$  is a Radon space. We deduce that  $\nu_n$  is a Radon measure. ■

For every locally topological vector space  $X$ , we denote by  $\text{Cyl}(X)$  the  $\sigma$ -algebra generated by continuous linear forms on  $X$ .

Recall that  $(X, weak)$  coincides with  $\text{Cyl}(X)$  [6, th. 2.3].

**Lemma 14.** *Let  $L$  be a Rosenthal compact set. Then  $(C(L), \tau_p)$  is measure-compact if and only if  $(C(L), weak)$  is measure-compact.*

*Proof.* Show first that  $\text{Cyl}((C(L), weak)) = \text{Cyl}((C(L), \tau_p))$ . (It follows by [6, th. 2.3] that

$$\text{Bair}(C(L), weak) = \text{Bair}(C(L), \tau_p). \quad (6.1)$$

Indeed, it is clear that  $\text{Cyl}((C(L), \tau_p)) \subset \text{Cyl}((C(L), \text{weak}))$ . Let us show the converse inclusion. Let  $y^* \in M^+(L)$ . By [8],  $M^+(L)$  is a Rosenthal compact set, hence by [3] it is angelic, we deduce that there exists a sequence  $(y_n^*)_{n \geq 0}$  of finite support on  $L$  such that  $y_n^* \xrightarrow{n \rightarrow +\infty} y^*$  for the  $w^*$ -topology. For each  $n \in \mathbb{N}$ ,  $y_n^*$  is measurable with respect to  $\text{Cyl}((C(L), \tau_p))$ , this implies that  $y^*$  is measurable with respect to  $\text{Cyl}((C(L), \tau_p))$ . If  $y^* \in B_{(C(L))^*}$ , there exists  $\mu_1, \mu_2, \mu_3, \mu_4 \in M^+(L)$  such that  $y^* = \mu_1 - \mu_2 - i(\mu_3 - \mu_4)$ , hence  $y^*$  is measurable with respect to  $\text{Cyl}((C(L), \tau_p))$ , thus  $\text{Cyl}((C(L), \text{weak})) \subset \text{Cyl}((C(L), \tau_p))$ .

Suppose now that  $(C(L), \tau_p)$  is measure-compact. Let  $\nu$  be a measure defined on  $\text{Cyl}((C(L), \text{weak}))$ . By [25, th. 2],  $\nu$  extends to a normal measure  $\tilde{\nu}$  on  $(C(L), \tau_p)$ . On the other hand, by Proposition 15  $(C(L), \tau_p)$  is universally measurable, hence  $\tilde{\nu}$  is a Radon measure on  $(C(L), \tau_p)$ . Let  $\varepsilon > 0$  and let  $C \in \text{Bor}(C(L), \tau_p)$ . There exists a  $\tau_p$ -compact set  $H \subset C$  such that  $\tilde{\nu}(H) > \tilde{\nu}(C) - \varepsilon$ . For every integer  $n \in \mathbb{N}^*$ , consider  $B_n = \{g \in C(L); \|g\|_{C(L)} \leq n\}$ . Since  $\tilde{\nu}(H \cap B_n) \xrightarrow{n \rightarrow +\infty} \tilde{\nu}(H)$ , there is  $n_0 \in \mathbb{N}^*$  such that  $\tilde{\nu}(H \cap B_{n_0}) > \tilde{\nu}(C) - \varepsilon$ .

Note that  $H \cap B_{n_0}$  is an uniformly bounded  $\tau_p$ -compact subset of  $H$ , hence by [11]  $H \cap B_{n_0}$  is a weakly compact subset. By [21, chap. I, th. 16], one obtains that  $\tilde{\nu}$  extends to a Radon-measure  $\nu'$  on  $(C(L), \text{weak})$ . In particular  $\nu$  extends to a normal measure on  $(C(L), \text{weak})$ . Note that, if  $f$  is a bounded continuous function on  $(C(L), \text{weak})$ , then we have

$$\int_{C(L)} f(x) d\tilde{\nu}(x) = \int_{C(L)} f(x) d\nu'(x) = \int_{C(L)} f(x) d\nu(x).$$

Let now  $(f_\alpha)_{\alpha \in I}$  be a decreasing net of bounded continuous functions on  $(C(L), \text{weak})$  such that  $f_\alpha \rightarrow 0$  for the pointwise topology. By [25, th. 2]  $\int_{C(L)} f_\alpha(x) d\nu(x) = \int_{C(L)} f_\alpha(x) d\tilde{\nu}(x) \rightarrow 0$ . It follows that  $(C(L), \text{weak})$  is measure-compact.

Conversely, suppose that  $(C(L), \text{weak})$  is measure-compact. Since, by above  $\text{Bair}(C(L), \text{weak}) = \text{Bair}(C(L), \tau_p)$ , it follows that  $(C(L), \tau_p)$  is measure compact. ■

**Proposition 16.** *Suppose that  $K$  is separable. Then  $(C(K), \tau_p)$  is not measure-compact.*

*Proof.* It is clear that for every  $\theta \in K = B_1 \cup B_2 \cup B_3$ , the map  $t \rightarrow \psi(\theta, t)$  is measurable.

By an argument similar to that of Proposition 11, we show that for every  $\xi \in C(K)^*$ , the map  $t \in E \rightarrow \xi(f^t)$  is measurable.

Suppose now that  $(C(K), \tau_p)$  is measure-compact. Lemma 14 shows us that  $(C(K), weak)$  is measure-compact. Since the map  $t \rightarrow f_t$  is scalarly measurable and  $(C(K), weak)$  is measure-compact, by Theorem of [6], there exists a strongly measurable function  $g$  with values in  $C(K)$  such that for every  $\xi \in C(K)^*$   $\xi(f^t) = \xi(g(t))$ , for almost all  $t \in E$ .

Let  $(\theta_n)_{n \geq 0}$  be a dense sequence in  $K$ . For every  $n \in \mathbb{N}$  there exists a measurable subset  $D_n$  of  $E$  such that  $\mu(D_n) = 0$  and  $\theta_n(f^t) = \theta_n(g(t))$ , for all  $t \in (D_n)^c$ . Denote  $D = \bigcup_{n \geq 0} D_n$ . Since  $\theta_n(f^t) = \theta_n(g(t))$ , for all  $n \in \mathbb{N}$  and all  $t \in D^c$ ,  $\theta(f^t) = \theta(g(t))$ , for all  $\theta \in K$  and all  $t \in D^c$ . It follows that  $f^t = g(t)$ , for almost all  $t \in E$ , this is impossible, because  $\|f_t - f_{t'}\|_{L^\infty(E, \mu)} \geq 1$ , if  $t \neq t'$  and  $g$  has almost values in a separable subspace of  $C(K)$ . Thus  $(C(K), \tau_p)$  is not measure-compact. ■

7. THE CASE OF A SEPARABLE CONNECTED ABELIAN GROUP  
 VERIFYING THE CONDITION (\*\*)

In this part,  $E$  is a separable connected abelian group verifying the condition (\*\*) and  $E$  is locally compact.

By Remark 1,  $E$  has a countable basis. As  $E$  is locally compact, by [21, chap. II, th. 6],  $E$  is a Polish space.

Note that  $-\infty, +\infty \notin E$ . For  $g \in C(K) \subset L^\infty(E, \mu)$ ,  $\theta \in K$  and  $t \in E$ , one defines  $g_t \in C(K)$  by  $g_t(u) = g(u - t)$ , for almost all  $u \in E$  and  $\theta_t \in K$  by  $\theta_t(r) = \theta(r_t)$ ,  $r \in C(K)$ .

Let  $f$  be the characteristic function of  $\{t \in E; t > 0\}$ .

**Lemma 15.** *For every  $\theta \in B_2$ ,  $h(\theta_u) = h(\theta) - u$ , for all  $u \in E$ .*

*Proof.* Since  $E$  satisfies the condition (\*\*),  $f_t = f^t$ , for every  $t \in E$ . Let  $\theta \in B_2$  and let  $t, u \in E$  such that  $\psi(\theta_u, t) = 1$ . Hence we have  $\psi(\theta_u, t) = \theta_u(f_t) = \theta[(f_t)_u] = \theta(f_{t+u}) = \psi(\theta, t + u) = 1$ . It follows that  $t + u \leq h(\theta)$ . The condition (\*\*) implies that  $t \leq h(\theta) - u$ . By Lemma 2, one has  $h(\theta_u) = \sup \{t; \psi(\theta_u, t) = 1\} \leq h(\theta) - u$ .

Conversely, let  $t \in E$  such that  $\psi(\theta, t) = 1$ . For every  $u \in E$ , we have  $\psi(\theta_u, t - u) = \theta_u(f_{t-u}) = \theta((f_{t-u})_u) = \theta(f_t) = \psi(\theta, t) = 1$ , hence  $h(\theta_u) \geq t - u$ , since  $E$  satisfies the condition (\*\*),  $h(\theta_u) + u \geq t$ . By Lemma 2, we obtain that  $h(\theta_u) + u \geq h(\theta)$ , i.e.  $h(\theta_u) \geq h(\theta) - u$ , we deduce that  $h(\theta_u) = h(\theta) - u$ . ■

**Remark 23.** *In the previous lemma, one can replace  $B_2$  by  $B_3$  (using Lemma 3).*

**Remark 24.**  $B_2$  is not empty.

*Proof.* Indeed, suppose that  $B_2 = \emptyset$ . Proposition 17 tells us that  $B_3 = \emptyset$ , hence  $K = B_1$  which is metrisable (by Proposition 7-ii), It is impossible, because  $(E, \mu)$  satisfies the condition (1.2). ■

**Lemma 16.** For every  $\theta \in K \setminus B_1$ :

$$\sup \{-t \in E; \psi(\theta, t) = 0\} = -\inf \{t \in E; \psi(\theta, t) = 0\}.$$

*Proof.* Consider  $\theta \in K \setminus B_1$ . Put

$$\beta = -\inf \{t \in E; \psi(\theta, t) = 0\}.$$

Let  $t \in E$  such that  $\psi(\theta, t) = 0$ . By definition of  $-\beta$  we have  $-\beta \leq t$ . The condition (\*\*\*) implies that  $-\beta + (\beta - t) \leq t + (\beta - t)$ , hence  $-\beta \leq \beta$ , to complete the proof, it suffices to show that for every  $z \in E$  such that  $-\beta \leq z$ , and if  $\psi(\theta, t) = 0$ , then  $\beta \leq z$ . Let a such  $z \in E$ . Since  $E$  satisfies the condition (\*\*),  $t \geq -z$ , for every  $t \in E$  such that  $\psi(\theta, t) = 0$ . We deduce that  $\inf \{t \in E; \psi(\theta, t) = 0\} \geq -z$ , i.e.  $\beta \leq z$ . ■

For every  $\theta \in K$ , one defines  $\overset{\vee}{\theta} \in K$ , by  $\overset{\vee}{\theta}(g) = \theta(\overset{\vee}{g})$ , where  $\overset{\vee}{g}(x) = g(-x)$ , for almost  $x \in E$ ,  $g \in C(K) \subset L^\infty(E, m)$ .

**Proposition 17.** Let  $\theta \in K \setminus B_1$ . Then:

- i)  $h(\theta) = -h(\overset{\vee}{\theta})$ .
- ii)  $\varphi \in B_2$ , if and only if  $\overset{\vee}{\varphi} \in B_3$ .
- iii)  $(\overset{\vee}{\theta}') = \theta''$ .

*Proof.*

i). The measure  $\mu = m$  is invariant by translation, then  $\mu(\{t\}) = 0$ , for every  $t \in E$ . Thus we have

$$\begin{aligned} & \mu\left(\{y \in E; (f_{-t}^{\vee})(y) = 0\}\right) \\ &= \mu(\{y \in E; f_{-t}(-y) = 0\}) \\ &= \mu(\{y \in E; f(t - y) = 0\}) = \mu(\{y \in E; y \geq t\}) \\ &= \mu(\{y \in E; y > t\}) = \mu(\{y \in E; f_t(y) = 1\}) \\ &= \mu(\{y \in E; e(y) - f_t(y) = 0\}). \end{aligned}$$

By a similar argument we show that

$$\mu(\{y \in E; (f_{-t}^{\vee})(y) = 1\}) = \mu(\{y \in E; e(y) - f_t(y) = 1\}).$$

Thus for every  $t \in E$

$$(f_{-t}^{\vee}) = e - f_t. \tag{7.1}$$

Let  $\theta \in K \setminus B_1$  and  $t \in E$ . By (7.1),

$$\psi(\overset{\vee}{\theta}, -t) = \overset{\vee}{\theta}(f_{-t}) = \theta(f_{-t}) = \theta(e) - \psi(\theta, t) = 1 - \psi(\theta, t),$$

i. e.

$$\psi(\overset{\vee}{\theta}, -t) = 0 \text{ if and only if } \psi(\theta, t) = 1. \quad (7.2)$$

Using (7.2) and Lemma 2, one obtains

$$\begin{aligned} h(\theta) &= \sup\{t \in E; \psi(\theta, t) = 1\} = \sup\{t \in E; \psi(\overset{\vee}{\theta}, -t) = 0\} \\ &= \sup\{-t \in E; \psi(\overset{\vee}{\theta}, t) = 0\}. \end{aligned}$$

On the other hand, by Lemma 16 and Lemma 3

$$\begin{aligned} &\sup\{-t \in E; \psi(\overset{\vee}{\theta}, t) = 0\} \\ &= -\inf\{t \in E; \psi(\overset{\vee}{\theta}, t) = 0\} = -h(\overset{\vee}{\theta}). \end{aligned}$$

Therefore  $h(\theta) = -h(\overset{\vee}{\theta})$ . ■

ii). By (7.2) and (i),  $\psi(\varphi, h(\varphi)) = 1$  if and only if  $\psi(\overset{\vee}{\varphi}, h(\overset{\vee}{\varphi})) = \psi(\overset{\vee}{\varphi}, -h(\varphi)) = 0$ . Thus  $\varphi \in B_2$  if and only if  $\overset{\vee}{\varphi} \in B_3$ . ■

iii). Obviously by (7.1). ■

**Lemma 17.** *Let  $\theta \in B_j$  and  $t \in E$ . Then  $\theta_t \in B_j$ ,  $j \in \{2, 3\}$ .*

*Proof.* Suppose that  $\theta \in B_2$ . By Lemma 15 we have  $\psi(\theta_t, h(\theta_t)) = \psi(\theta_t, h(\theta) - t) = \psi(\theta, h(\theta)) = 1$ , hence  $\theta_t \in B_2$ . By a similar argument, one shows that  $\theta_t \in B_3$ , if  $\theta \in B_3$ . ■

**Proposition 18.** *For  $\theta \in B_2$ , the map  $U'_\theta : E \rightarrow K \times K$ ,  $t \rightarrow (\theta_t, (\overset{\vee}{\theta})_t)$  is not Borel.*

*Proof.* Suppose that there exists  $\theta \in B_2$  such that  $U'_\theta$  is a Borel function. Choose a subset  $H$  of  $E$  which is not in the Borel  $\sigma$ -algebra of  $E$ . We define the open subset  $V$  of  $K \times K$  by

$$\begin{aligned} V &= \bigcup_{u \in H} \left[ \{\varphi \in K; \psi(\varphi, -u + h(\theta)) = 1\} \right. \\ &\quad \left. \times \{\varphi \in K; \psi(\varphi, -u + h(\overset{\vee}{\theta})) = 0\} \right]. \end{aligned}$$

By Lemma 17, for every  $t \in E$ ,  $\theta_t \in B_2$  and  $(\check{\theta})_t \in B_3$  (because  $\check{\theta} \in B_3$ , by Proposition 17). Thus we have

$$\begin{aligned}
& (U'_\theta)^{-1}(V) \\
&= \bigcup_{u \in H} \{t \in E; \psi(\theta_t, -u + h(\theta)) = 1 \text{ and } \psi((\check{\theta})_t, -u + h(\check{\theta})) = 0\} \\
&= \bigcup_{u \in H} \{t \in E; h(\theta_t) \geq -u + h(\theta) \text{ and } h((\check{\theta})_t) \leq -u + h(\check{\theta})\} \\
&= \bigcup_{u \in H} \{t \in E; h(\theta) - t \geq -u + h(\theta) \text{ and } h(\check{\theta}) - t \leq -u + h(\check{\theta})\} \\
&= \bigcup_{u \in H} \{t \in E; -u \leq -t \leq -u\} = \bigcup_{u \in H} \{u\} = H,
\end{aligned}$$

because  $h(\theta_t) = h(\theta) - t$  and  $h((\check{\theta})_t) = h(\check{\theta}) - t$ , for every  $t \in E$ , by Lemma 15. This implies that  $H$  is a Borel subset of  $E$ , which is impossible. ■

**Corollary 19.** *The Borel  $\sigma$ -algebra of the product space  $K \times K$  strictly contains  $\text{Bor}(K) \otimes \text{Bor}(K)$ .*

*Proof.* Consider the maps  $\delta_1 : t \in E \rightarrow \theta_t \in B_2 \subset K$ ,  $\delta_2 : t \in E \rightarrow (\check{\theta})_t \in B_3 \subset K$ .

Let us show that  $\delta_1, \delta_2$  are Borel functions. Let  $u, t \in E$ . It is clear that the subsets of the form

$$\begin{aligned}
& (\delta_1)^{-1} [\{\varphi \in K; \psi(\varphi, u) = 0\}], \\
& (\delta_1)^{-1} [\{\varphi \in K; \psi(\varphi, t) = 1\}], \\
& (\delta_2)^{-1} [\{\varphi \in K; \psi(\varphi, u) = 0\}], \\
& (\delta_2)^{-1} [\{\varphi \in K; \psi(\varphi, t) = 1\}],
\end{aligned} \tag{7.3}$$

are Borel subsets of  $E$  (note for example that

$$\begin{aligned}
& (\delta_1)^{-1} [\{\varphi \in K; \psi(\varphi, u) = 0\}] = \{t \in E; \psi(\theta_t, u) = 0\} \\
& = \{t \in E; h(\theta_t) = h(\theta) - t < u\} = \{t \in E; t > h(\theta) - u\}.
\end{aligned}$$

On the other hand by Proposition 10,  $K$  is a strongly Lindelöf space, hence  $\delta_1$  and  $\delta_2$  are Borel functions, we deduce that for every  $S \in \text{Bor}(K) \otimes \text{Bor}(K)$ ,  $(U'_\theta)^{-1}(S) \in \text{Bor}(E)$ .

Suppose now that  $\text{Bor}(K \times K) = \text{Bor}(K) \otimes \text{Bor}(K)$ . By above,  $U'_\theta$  is a Borel function, this is impossible by Proposition 18. ■

L. Lindenstrauss and C. Stegall [15] showed that there exists a Banach space  $X$  which does not contain  $\ell^\infty$  isomorphically and a map

$\sigma : \{0, 1\}^{\mathbb{N}} \rightarrow X$  scalarly measurable, but  $\sigma$  is not weakly equivalent to a measurable function. In the following proposition, one gives another type of examples. In the fact, the considered function (in Proposition 19) is locally integrable in the sense of Riemann.

**Proposition 19.** *There exists a Banach space  $X$  and a function  $g : E \rightarrow X$  such that*

- I)  $X$  does not contain  $\ell^\infty$  isomorphically.
- II)  $g$  is scalarly measurable.
- III)  $g$  cannot be weakly equivalent to a strongly measurable function.
- IV)  $g$  is locally integrable in the sense of Riemann.

*Proof.*

I). Consider  $X = C(K)$ . Since for all  $\xi \in B_{C(K)^*}$  there exist  $\xi_1, \xi_2, \xi_3, \xi_4 \in M^+(K)$  such that  $\xi = \xi_1 - \xi_2 + i(\xi_3 - \xi_4)$ , by [3]-[8]  $B_{C(K)^*}$  is angelic, hence  $X$  does not contain  $\ell^\infty$  isomorphically, because by [18] the closed unit ball of  $(\ell^\infty)^*$  is not angelic. ■

II). Let  $g : E \rightarrow (C(K) \subset L^\infty(E))$  the function defined by  $g(t) = f_t$  for almost all  $t \in E$ . By Proposition 11,  $g$  is scalarly measurable. ■

III). In the proof of Proposition 16 we saw that  $g$  cannot be weakly equivalent to a strongly measurable function. ■

IV). Let  $a, b \in E$  such that  $a < b$ . Note that

$$\int_a^b \chi_{[t, +\infty[}(\omega) dt = \begin{cases} (a \vee \omega) - a, & \text{si } \omega \leq b \\ b - a, & \text{si } \omega > b. \end{cases}$$

Consider  $t_0 = a, t_1, \dots, t_n = b \in [a, b]$  such that  $t_0 < t_1 < \dots < t_n$ . For  $t_{k(\omega)} \leq \omega < t_{k(\omega)+1}$ , we have  $\sum_{k=1}^n \chi_{[t_k, +\infty[}(\omega)(t_k - t_{k-1}) = \sum_{k=1}^{k(\omega)} \chi_{[t_k, +\infty[}(\omega)(t_k - t_{k-1}) = \sum_{k=1}^{k(\omega)} (t_k - t_{k-1}) = t_{k(\omega)} - a$ . One deduces that

$$\sum_{k=1}^n \chi_{[t_k, +\infty[}(\omega)(t_k - t_{k-1}) = \begin{cases} t_{k(\omega)} - a, & \text{si } t_{k(\omega)} \leq \omega < t_{k(\omega)+1} \\ 0, & \text{si } \omega < a \\ \sum_{k=1}^n (t_k - t_{k-1}) = b - a, & \text{si } \omega > b. \end{cases}$$

Thus

$$\begin{aligned} & \left\| \sum_{k=1}^n \chi_{[t_k, +\infty[}(\cdot)(t_k - t_{k-1}) - \int_a^b \chi_{[t, +\infty[}(\cdot) dt \right\|_{L^\infty(E)} \\ & \leq \sup_{\omega \in [a, b]} |\omega - t_{k(\omega)}| \leq \sup_{1 \leq k \leq n} |t_k - t_{k-1}|. \end{aligned}$$

It follows that  $g$  is integrable in the sense of Riemann on  $[a, b]$ . ■

Let  $M$  be the subset of  $C(K)$  formed by even functions with values in  $\{0, 1, 2\}$  ( $g$  is even function on  $K$  if  $g = \check{g}$ ).

**Theorem 5.** *There exists a Hausdorff topology  $\tau'$  on  $C(K)$ , a  $C^*$ -subalgebra  $Y_1$  of  $C(K)$  and a Banach subspace  $Y_2$  of  $C(K)$  such that*

- 1)  $\tau_p$  is finer than  $\tau'$ .
- 2)  $(C(K), \tau')$  is not universally measurable.
- 3)  $Y_1$  and  $Y_2$  are  $\tau_p$  closed and  $(C(K), \tau_p) = (Y_1, \tau_p) \oplus (Y_2, \tau_p)$ .
- 4)  $(Y_1, \tau_p)$  is isomorphic to  $(Y_2, \tau_p)$ .
- 5)  $(Y_j, \tau_p) = (Y_j, \tau')$ ,  $j \in \{1, 2\}$ .
- 6)  $(Y_1, \tau_p)$  contains a closed discrete uncountable subset which is contain in  $M$ .
- 7)  $\text{Bor}(C(K), \tau') \otimes \text{Bor}(C(K), \tau') \neq \text{Bor}(C(K) \times C(K), \tau' \times \tau')$ .
- 8) The projection  $P : (C(K), \tau') \rightarrow (Y_1, \tau')$  is not a Borel function.
- 9) The set  $\{H : E \rightarrow (C(K), \tau'); H \text{ is a Borel function}\}$  is not a vector subspace of  $C(K)^E$ .

*Proof.*

1).

*Step 1:* Let  $\theta \in B_2$ . Let us show that  $\{\theta_t; t \in E\} = B_2$ .

First observe that  $B_2 \neq \emptyset$  by Remark 24. Let  $\varphi \in B_2$ . There exists  $t \in E$  such that  $h(\varphi) = h(\theta_t)$  (because  $h(\theta_t) = h(\theta) - t$  by Lemma 15). By Lemma 17,  $\theta_t \in B_2$ . Since  $h_2 : B_2^0 \rightarrow E$  is onto, by Remark 2,  $\varphi = \theta_t$ . ■

*Step 2 :* Show that there exists a sequence in  $B_2$  which is dense in  $K$ .

Since  $h_2$  is onto, there is a sequence  $(\theta_n)_{n \geq 0}$  in  $B_2$  such that  $(h(\theta_n))_{n \geq 0}$  is dense in  $E$ .

We will show that  $(\theta_n)_{n \geq 0}$  is dense in  $K$ . For that, let  $t \in E$ . Denote  $V_t = \{\theta \in K; \psi(\theta, t) = 1\}$ . Choose  $\theta \in K \setminus B_1$  such that  $t < h(\theta)$ , there exists  $\theta_m \in B_2$  satisfying  $t < h(\theta_m)$ , because the sequence  $(h(\theta_n))_{n \geq 0}$  is dense in  $E$ . Since  $\psi(\theta_m, h(\theta_m)) = 1$  and  $t < h(\theta_m)$ ,  $\theta_m \in V_t$ . Let  $u \in E$  and let  $U_u = \{\theta \in K; \psi(\theta, u) = 0\}$ . Choose  $\theta \in K \setminus B_1$  such that  $u > h(\theta)$ , there is  $\theta_{m'} \in B_2$  satisfying  $u > h(\theta_{m'})$ . It follows that  $\psi(\theta'_{m'}, u) = 0$ , because  $\psi(\theta'_{m'}, h(\theta'_{m'})) = 1$ . Thus  $\theta_{m'} \in U_u$ .

Let  $t, u \in E$  and let

$$\theta_0 \in W_{t,u} = \{\theta \in K; \psi(\theta, t) = 1 \text{ and } \psi(\theta, u) = 0\}.$$

We note that  $t < u$ . Since  $E$  satisfies the condition  $(*)$  and  $h$  is onto, there exists  $\theta_1 \in K \setminus B_1$  such that  $t < h(\theta_1) < u$ . We deduce that there exists  $\theta_n \in B_2$  verifying  $t < h(\theta_n) < u$ . Thus  $\theta_n \in W_{t,u}$ . ■

Let  $\tau'$  be the topology defined on  $C(K)$  by the condition that  $u_\alpha \rightarrow u$  in  $(C(K), \tau')$ , if and only if  $u_\alpha(\theta) \rightarrow u(\theta)$  for all  $\theta \in B_2 \cup B_1$ . Therefore  $\tau_p$  is finer than  $\tau'$ , and by step 2  $\tau'$  is Hausdorff.

2). Put  $D = \{f_t; t \in E\} \subset (C(K), \tau')$ .

*Step 1:* Show that  $(D, \tau')$  is a strongly Lindelöf space.

Pick  $\theta \in B_2$  and define the map  $\pi : (D, \tau') \rightarrow B_2$ , by  $\pi(f_t) = \theta_t$ ,  $t \in E$ . It suffices to show that  $\pi$  is a homeomorphism, because  $B_2$  is strongly Lindelöf.

Let us first show that  $\pi$  is injective. For that, let  $u, v \in E$  such that  $\theta_u = \theta_v$ . By Lemma 15,  $h(\theta) - u = h(\theta) - v$ , hence  $u = v$ . Let  $u \in E$ . Put  $V_u = \{\theta \in B_2; \psi(\theta, u) = 0\}$  and  $W_u = \{\theta \in B_2; \psi(\theta, u) = 1\}$ . Thus we have  $\pi^{-1}(V_u) = \{f_t; \psi(\theta_t, u) = 0\} = \{f_t; \theta_t(f_u) = 0\} = \{f_t; \theta_u(f_t) = 0\}$  is an open subset of  $(D, \tau')$ . In the same way, we have  $\pi^{-1}(W_u) = \{f_t; \theta_u(f_t) = 1\}$  is an open subset of  $(D, \tau')$ . We deduce that  $\pi$  is continuous.

We shall show that  $\pi^{-1}$  is continuous. Indeed, let  $t_1, t_2 \in E$ . Denote  $W_1 = \{f_t; f_t(\theta_{t_1}) = 1\}$  and  $W_2 = \{f_t; f_t(\theta_{t_2}) = 0\}$ . It is enough to show that  $\pi(W_j)$  is an open subset of  $B_2$ ,  $j \in \{1, 2\}$ , because the open sets of the previous form generated the topology  $\tau'$ .

Note that  $\pi(W_1) = \{\theta_t; f_t(\theta_{t_1}) = 1\} = \{\theta_t; f_{t_1}(\theta_t) = 1\}$  and  $\pi(W_2) = \{\theta_t; f_t(\theta_{t_2}) = 0\} = \{\theta_t; f_{t_2}(\theta_t) = 0\}$ . We deduce that  $\pi(W_1), \pi(W_2)$  are open subsets of  $B_2$ .

Thus  $\pi$  is a homeomorphism. ■

Let  $F_1$  be the closure of  $D$  in  $(C(K), \tau')$  and let  $\sigma_1 : (E, \tau_0) \rightarrow F_1 \subset (C(K), \tau')$  the map defined by  $\sigma_1(t) = f_t$ ,  $t \in E$ . We will show that  $\sigma_1$  is Borel.

Indeed, let  $O$  be an open subset of  $(D, \tau')$ . There exists a sequence  $(O_i)_{i \in I}$  of  $D$  such that  $O = \bigcup_{i \in I} O_i$  and each  $O_i$  is a finite intersection of open subsets, under the form  $W_1, W_2$ , where  $W_1 = \{f_t; f_t(\theta_{t_1}) = 1\}$ ,  $W_2 = \{f_t; f_t(\theta_{t_2}) = 0\}$ ,  $t_1, t_2 \in E$ . Note that for every  $\theta \in B_2 \cup B_1$  the map  $t \in E \rightarrow f_t(\theta) = \psi(\theta, t)$  is Borel, hence  $(\sigma_1^{-1})(O_i)$  is Borel, for all  $i \in I$ . By step 1,  $(D, \tau')$  is strongly Lindelöf, one concludes that there is a countable subset  $I_1$  of  $I$  such that  $O = \bigcup_{i \in I_1} O_i$ . It follows that  $(\sigma_1)^{-1}(O)$  is a Borel subset of  $(E, \tau_0)$ . Finally if  $O'$  is an open subset of  $F_1$ , then  $(\sigma_1)^{-1}(O') = (\sigma_1)^{-1}(O' \cap D)$  and  $O = O' \cap D$  is an open subset of  $(D, \tau')$ .

*Step 2:* Show that every compact of  $F_1$  is metrisable.

Note that if  $g \in F_1$ ,  $g(\theta) \in \{0, 1\}$ .

Let  $L$  be a compact subset of  $F_1$  and let  $(\theta_n)_{n \geq 0}$  be a sequence in  $B_2$  which is dense in  $K$  (a such sequence exists by step 2 of 1). Define the

map  $\xi : L \rightarrow \{0, 1\}^{\mathbb{N}}$  by  $\xi(g) = (g(\theta_n))_{n \geq 0}$ ,  $g \in L$ . It is clear that  $\xi$  is continuous and injective, this implies that  $L$  is metrisable. ■

Observe now that the topology  $\tau'$  is defined by a family of semi-norms which separates the points of  $C(K)$ , hence  $(C(K), \tau')$  is completely regular space. Since every subspace of a completely regular space is completely regular,  $(F_1, \tau')$  is completely regular.

Suppose that  $(C(K), \tau')$  is universally measurable.

*Step 3:* Show that  $F_1$  is universally measurable.

By [22, th. 3.2], it is enough to show that every normal probability measure on  $F_1$  is a Radon measure. Let  $\nu$  be a normal probability measure on  $F_1$ . Consider  $\nu'$  the measure defined on  $(C(K), \tau')$  by  $\nu'(B) = \nu(B \cap F_1)$ , where  $B \in \text{Bor}(C(K), \tau')$ . Let us show that  $\nu'$  is a normal measure on  $(C(K), \tau')$ .

Let  $i : F_1 \rightarrow (C(K), \tau')$  be the canonical injection. Since  $i$  is continuous and  $\nu$  is normal, by [22],  $i(\nu) = \nu'$  is a normal measure. On the other hand  $(C(K), \tau')$  is universally measurable and  $\nu'$  is normal, hence by [22, th. 3.2],  $\nu'$  is a Radon measure. Now we let  $B'$  be a Borel subset of  $F_1$ . Note that  $B'$  is a Borel subset of  $(C(K), \tau')$ .

Thus we have

$$\begin{aligned} \nu(B') &= \nu'(B') = \sup\{\nu'(L'); L' \text{ compact in } B'\} \\ &= \sup\{\nu(L'); L' \text{ compact in } B'\}. \end{aligned}$$

Therefore  $\nu$  is a Radon measure. It follows that  $F_1$  is universally measurable. ■

Let  $L$  be a compact subset of  $(E, \tau_0)$  of strictly positive measure. We can suppose that  $\mu(L) = 1$ . Consider the map  $\eta_1 : L \rightarrow F_1$ , the restriction of  $\sigma_1$  to  $L$  and  $\nu_1 = \eta_1(\mu_L)$ , where  $\mu_L$  the measure defined on  $L$  by

$$\mu_L(C) = \mu(C) = m(C), \quad C \in \text{Bor}(L) \subset \text{Bor}(E).$$

*Step 4:* Show that  $\nu_1$  is a normal measure on  $F_1$ .

Let  $(U_\alpha)$  be a net of open subsets in  $F_1$  which is increasing to an open subset of  $F_1$ . Observe that for every Borel subset  $B$  of  $F_1$ ,  $\nu_1(B) = \mu(\{t \in L; f_t \in B\}) = \mu(\{t \in L; f_t \in B \cap D\})$ .

By step 1 of 2),  $(D, \tau')$  is strongly Lindelöf, hence there exists a countable subset  $I_1$  of  $I$  such that  $\bigcup_{i \in I} U_i \cap D = \bigcup_{i \in I_1} U_i \cap D$ . Thus

$$\begin{aligned} \nu_1\left(\bigcup_{i \in I} U_i\right) &= \mu\left(\left\{t \in L; f_t \in \left(\bigcup_{i \in I} U_i\right) \cap D\right\}\right) \\ &= \mu\left(\left\{t \in L; f_t \in \bigcup_{i \in I} U_i \cap D\right\}\right) = \mu\left(\left\{t \in L; f_t \in \bigcup_{i \in I_1} U_i \cap D\right\}\right) \\ &= \mu\left(\left\{t \in L; f_t \in \left(\bigcup_{i \in I_1} U_i\right) \cap D\right\}\right) \\ &= \nu_1\left(\bigcup_{i \in I_1} U_i\right) = \sup_{i \in I_1} \nu_1(U_i) \leq \sup_{i \in I} \nu_1(U_i). \end{aligned}$$

This implies that  $\nu_1$  is a normal measure on  $F_1$ . ■

By step 3 of 2),  $F_1$  is universally measurable, since  $\nu_1$  is a normal probability measure on  $F_1$ , by [22, th. 3.2],  $\nu_1$  is a Radon measure, we deduce that there exists a compact subset  $G_1$  of  $F_1$  such that  $\nu_1(G_1) > \frac{1}{2}$ . We denote by  $\eta_2$  the restriction of  $\sigma_1$  to  $(\eta_1)^{-1}(G_1) \subset L$ . the map  $\eta_2$  is Borel with values in  $G_1$  and  $m((\eta_1)^{-1}(G_1)) = \nu_1(G_1) > \frac{1}{2}$ . The step 2 of 2) shows us that  $G_1$  is metrisable. By [5], there exists a compact subset  $L_1$  of  $((\eta_1)^{-1}(G_1), \tau_0)$  such that  $\mu(L_1) > \frac{1}{3}$  and the restriction of  $\sigma_1$  to  $L_1$  is continuous.

Since  $\mu$  is diffuse on  $E$ , by an argument similar to that of Proposition 12, one constructs a strictly decreasing sequence  $(t_n)_{n \geq 0}$  in  $L_1$ . There exists  $t_0 \in L_1$  such that  $t_n \xrightarrow{n \rightarrow +\infty} t_0$  (modulo subsequence). On the other hand, the restriction of  $\sigma_1$  to  $L_1$  is continuous, it follows that  $f_{t_n} \xrightarrow{n \rightarrow +\infty} f_{t_0}$  in  $(C(K), \tau')$ . For  $s = h(\theta) - t_0$ , we have  $f_{t_n}(\theta_s) \xrightarrow{n \rightarrow +\infty} f_{t_0}(\theta_s) = \psi(\theta_s, t_0) = \psi(\theta, h(\theta)) = 1$ . Thus for  $n$  large enough,  $\psi(\theta_s, t_n) = 1$ , hence  $t_n \leq h(\theta_s) = h(\theta) - s = t_0$ . For seeing a contradiction, it suffices to note that the sequence  $(t_n)_{n \geq 0}$  is strictly decreasing. ■

3). Let  $Y_1$  be the vector subspace formed by even functions in  $C(K) \subset L^\infty(E)$  and  $Y_2$  be the vector space formed by odd functions in  $C(K) \subset L^\infty(E)$ . Observe that if  $g \in C(K)$ ,  $\check{g} \in C(K)$ , because  $\check{f}_t = e - f_{-t}, \forall t \in E$ .

Let us show that  $Y_1$  is  $\tau_p$  closed in  $C(K)$ . For that let  $(r_\alpha)_{\alpha \in I}$  be a net in  $Y_1$  such that  $r_\alpha \rightarrow r$  in  $(C(K), \tau_p)$ . We must prove that  $r \in Y_1$ .

For  $\alpha \in I$  and  $\theta \in K$ , we have  $r_\alpha(\check{\theta}) = \check{r}_\alpha(\theta) = r_\alpha(\theta)$ , by passing to the limit we obtain that  $\check{r}(\theta) = r(\theta)$ , hence  $\check{r} = r \in Y_1$ . By a similar argument, we show that  $Y_2$  is  $\tau_p$  closed in  $C(K)$ .

On the other hand, for  $g \in C(K)$ ,  $g = (g + \overset{\vee}{g})/2 \oplus (g - \overset{\vee}{g})/2$ , this implies that  $(C(K), \tau_p) = (Y_1, \tau_p) \oplus (Y_2, \tau_p)$  (note that the map  $g \in C(K) \rightarrow \overset{\vee}{g} \in C(K)$  is continuous). ■

4). Define the operator  $U : (Y_1, \tau_p) \rightarrow (Y_2, \tau_p)$ , by  $U(g) = 2gf - g$ ,  $g \in Y_1$ . Let us show that  $U$  has values in  $Y_2$ . For that let  $t \in G$ . Observe that

$$(2gf - g)(-t) = 2g(t)f(-t) - g(t),$$

but  $f(-t) = e - f(t)$ , hence

$$\begin{aligned} (2gf - g)(-t) &= 2g(t)(e - f(t) - g(t)) \\ &= -[2g(t)f(t) - g(t)]. \end{aligned}$$

Since  $2(2gf - g)f - (2gf - g) = g$ ,  $U^{-1}(g) = 2gf - g$ ,  $g \in Y_2$ . We observe that  $U, U^{-1}$  are continuous. Thus  $U$  is an isomorphism. ■

5). Show that  $(Y_1, \tau') = (Y_1, \tau_p)$ . For  $\theta \in B_2 \cup B_1$  and  $g \in Y_1$  we have  $g(\overset{\vee}{\theta}) = (\overset{\vee}{g})(\theta) = g(\theta)$ . Since  $\{\overset{\vee}{\theta}; \theta \in B_2 \cup B_1\} = B_3 \cup B_1$  by Proposition 17, it follows that  $(Y_1, \tau') = (Y_1, \tau_p)$ .

Let us show that  $(Y_2, \tau') = (Y_2, \tau_p)$ . Let  $(g_\alpha)_{\alpha \in I}$  be a net in  $Y_2$  and let  $g \in Y_2$  such that  $g_\alpha(\theta) \rightarrow g(\theta)$ , for every  $\theta \in B_2 \cup B_1$ . We must prove that  $g_\alpha \rightarrow g$  in  $(Y_2, \tau_p)$ . Note that  $U^{-1}(g_\alpha)(\theta) = (2g_\alpha \times f - g_\alpha)(\theta)$  and  $U^{-1}(g)(\theta) = (2g \times f - g)(\theta)$ , hence  $U^{-1}(g_\alpha)(\theta) \rightarrow U^{-1}(g)(\theta)$ , for every  $\theta \in B_2 \cup B_1$ . Since  $(Y_1, \tau') = (Y_1, \tau_p)$ ,  $U^{-1}(g_\alpha)(\cdot) \rightarrow U^{-1}(g)(\cdot)$  in  $(Y_1, \tau_p)$ , we deduce that  $g_\alpha \rightarrow g$  in  $(Y_2, \tau_p)$ , because  $U$  is a homeomorphism. Thus  $(Y_2, \tau') = (Y_2, \tau_p)$ . ■

6). For every  $t \in E$  denote  $w_t = f_t + (\overset{\vee}{f_t}) = f_t + 1 - f_{-t}$  and  $V = \{\omega_t; t \in E\}$ . We shall show that  $(V, \tau_p)$  is discrete and closed in  $(Y_1, \tau_p)$ . Indeed, let  $t \in E$ ; choose  $\theta_1 \in B_2$  and  $\theta_2 \in B_3$  such that  $h(\theta_1) = h(\theta_2) = t$ . Denote

$$W = \left\{ \omega_s; |\omega_s(\theta_j) - \omega_t(\theta_j)| < \frac{1}{2}, j \in \{1, 2\} \right\}.$$

We observe that  $W$  is an open subset of  $(V, \tau_p)$ .

Let us show that  $W = \{\omega_t\}$ .

*Case 1,  $t < 0$ :* Let  $\omega_s \in W$ . Note that  $\omega_t(\theta_1) = 1 + 1 - 0 = 2$  and  $\omega_t(\theta_2) = 0 + 1 - 0 = 1$ , hence  $\omega_s(\theta_1) = 2$  and  $\omega_s(\theta_2) = 1$ , this implies that  $\psi(\theta_1, s) = 1$ ,  $\psi(\theta_1, -s) = 0$  and  $\psi(\theta_2, s) = \psi(\theta_2, -s)$ .

The case  $\psi(\theta_2, s) = \psi(\theta_2, -s) = 1$  is excluded, because it will lead that  $t > 0$ , which is impossible. We have then  $\psi(\theta_1, s) = 1$  and  $\psi(\theta_2, s) = \psi(\theta_2, -s) = 0$ . It follows that  $s \leq h(\theta_1) = t \leq s$ . It means that  $t = s$ .

*Case 2,  $t > 0$ :* Let  $\omega_s \in W$ . Note that  $\omega_t(\theta_1) = 1 + 1 - 1 = 1$  and  $\omega_t(\theta_2) = 0 + 1 - 1 = 0$ , hence  $\omega_s(\theta_1) = \psi(\theta_1, s) + 1 - \psi(\theta_1, -s) = 1$  and  $\omega_s(\theta_2) = \psi(\theta_2, s) + 1 - \psi(\theta_2, -s) = 0$ . Therefore  $\psi(\theta_1, s) = \psi(\theta_1, -s)$  and  $\psi(\theta_2, s) = 0$  (note that  $\psi(\theta_2, -s) = 1$ ). The case  $\psi(\theta_1, s) = \psi(\theta_1, -s) = 0$  leads that  $t < 0$ , which is impossible, hence  $\psi(\theta_1, s) = 1$  and  $\psi(\theta_2, s) = 0$ . It follows that  $s = t$ .

*Case 3,  $t = 0$ :* Let  $\omega_s \in W$ . It is clear that  $\omega_0(\theta_1) = \omega_0(\theta_2) = 1$ , which implies that  $\psi(\theta_1, s) = \psi(\theta_1, -s)$  and  $\psi(\theta_2, s) = \psi(\theta_2, -s)$ . Note that the case  $\psi(\theta_1, s) = \psi(\theta_1, -s) = 0$  and the case  $\psi(\theta_2, s) = \psi(\theta_2, -s) = 1$  are excluded. Thus  $\psi(\theta_1, s) = 1$  and  $\psi(\theta_2, s) = 0$ . This implies that  $s = 0$ . Therefore  $(V, \tau_p)$  is a discrete space in  $(Y_1, \tau_p)$ .

It remains to show that  $V$  is a closed subset in  $(Y_1, \tau_p)$ .

Let  $(\omega_{t_\alpha})_{\alpha \in I}$  be a net in  $V$  such that  $\omega_{t_\alpha} \rightarrow g \in Y_1$ . Choose a fixed point  $\theta$  in  $B_2$ . The sequence  $(\theta_{t_\alpha})_{\alpha \in I}$  is a net in  $K$ , hence, there exists  $\theta_0 \in K$  such that  $\theta_{t_\alpha} \rightarrow \theta_0 \in K$  (with respect to an ultrafilter  $\mathcal{U}$  on  $I$ ).

*Case 1,  $t_\alpha \rightarrow +\infty$ :* Let  $\varphi \in K \setminus \{\theta', \theta''\}$ . There exist  $v \in E^+$  and  $u \in E^-$  such that  $-u > v$ ,  $\psi(\varphi, u) = 1$  and  $\psi(\varphi, v) = 0$ . Put  $W = \{t \in E; t > v \text{ and } -t < u\}$ . We observe that  $S = \{\alpha \in I; t_\alpha \in W\}$  belongs to  $\mathcal{U}$ , hence for all  $\alpha \in S$ ,  $f_{t_\alpha}(\varphi) = \psi(\varphi, t_\alpha) = 0$  and  $f_{-t_\alpha}(\varphi) = \psi(\varphi, -t_\alpha) = 1$ , it follows that  $\omega_{t_\alpha}(\varphi) = 0 + 1 - 1 = 0$ , for every  $\alpha \in S$ . We conclude that  $g(\varphi) = 0$ . It is obvious that  $\omega_{t_\alpha}(\theta') = \omega_{t_\alpha}(\theta'') = 1$ , which implies that  $g(\theta') = g(\theta'') = 1$ . Therefore, we have  $\{\theta', \theta''\} = g^{-1}(\{1\})$ , hence this subset is closed and open, because  $g$  has values in  $\{0, 1, 2\}$  (note that  $K = \{\theta', \theta''\} \cup g^{-1}(\{0, 2\})$ ). Since  $\{\theta'\} = \{\theta', \theta''\} \setminus \{\theta''\}$ ,  $\{\theta'\}$  is an open subset, this means that there is  $t_0 \in E$  such that  $\{\theta'\} = \{\theta \in K; \psi(\theta, t_0) = 1\}$ , this is impossible, because the set  $\{\theta \in K; \psi(\theta, t_0) = 1\}$  contains the set  $\{\theta \in K; h(\theta) < t_0\}$ .

*Case 1,  $t_\alpha \rightarrow -\infty$ :* By an argument similar to the previous case, we show that this case can be excluded, because  $g(\varphi) \in \{0, 1, 2\}$ , for every  $\varphi \in K$ .

Since the cases 1 and 2 are excluded and  $E$  is a locally compact, there exists  $t_0 \in E$  such that  $t_\alpha \rightarrow t_0 \in E$ . By Lemma 15,  $h(\theta_{t_\alpha}) = h(\theta) - t_\alpha$ , which implies that  $t_\alpha \rightarrow h(\theta) - h(\theta_0) = t_0$  (because  $h$  is continuous by Proposition 5).

Show that  $g = \omega_{t_0}$ . Pick  $\varphi \in K \setminus B_1$ .

*Case 1,  $t_0 > 0$ .*

*Case 1-a),  $-t_0 < h(\varphi) < t_0$ :* Let  $W = \{t \in E; -t < h(\varphi) < t\}$ .  $W$  is an open neighborhood of  $t_0$ , hence

$$S = \{\alpha \in I; -t_\alpha < h(\varphi) < t_\alpha\} \in \mathcal{U}.$$

For every  $\alpha \in S$  we have  $f_{t_\alpha}(\varphi) = \psi(\varphi, t_\alpha) = \psi(\varphi, t_0) = 0$  and  $\psi(\varphi, -t_\alpha) = \psi(\varphi, -t_0) = 1$ , this implies that  $\omega_{t_\alpha}(\varphi) = \omega_{t_0}(\varphi) = 0 + 1 - 1 = 0$  for all  $\alpha \in S$ . Thus  $\omega_{t_\alpha}(\varphi) \rightarrow \omega_{t_0}(\varphi) = g(\varphi)$ .

*Case 1-b),  $h(\varphi) < -t_0$ .* Put

$$W = \{t \in E; h(\varphi) < -t < 0\}.$$

$W$  is an open neighborhood of  $t_0$ , hence  $S = \{\alpha \in I; h(\varphi) < -t_\alpha < 0\}$  belongs to  $\mathcal{U}$ . It is clear that  $\omega_{t_\alpha}(\varphi) = \omega_{t_0}(\varphi) = 0 + 1 - 0 = 1$  (because  $t_\alpha > -t_\alpha$  and  $\psi(\varphi, -t_\alpha) = 0$ , hence  $\psi(\varphi, t_\alpha) = 0$ ). We deduce that  $\omega_t(\varphi) \rightarrow \omega_{t_0}(\varphi) = 1 = g(\varphi)$ .

*Case 1-c),  $h(\varphi) > t_0$ :* By an argument similar to case 1-b), one shows that  $\omega_{t_\alpha}(\varphi) \rightarrow \omega_{t_0}(\varphi) = g(\varphi)$ .

*Case 2,  $t_0 < 0$ :* The case  $t_0 < h(\varphi) < -t_0$ ,  $h(\varphi) < -t_0$  and the case  $h(\varphi) > t_0$  will be treated by arguments similar to the previous cases.

*Case 3),  $h(\varphi) = t_0$ ,  $t_0 \in E$ :* Without losing the generality, we can suppose that  $\varphi \in B_2$ . In the proof of 1), we showed that  $B_2$  is dense in  $K$ . Hence there exists a sequence  $(\varphi_n)_{n \geq 0}$  in  $B_2$  such that  $\varphi_n \rightarrow_{n \rightarrow +\infty} \varphi$ . We can choose the sequence  $(\varphi_n)_{n \geq 0}$  in  $B_2 \setminus \{\varphi\}$ , because  $K$  does not contain isolated points. Let us show that for  $n$  large enough  $t_0 < h(\varphi_n)$  (note that the restriction of  $h$  to  $B_2$  is injective, hence  $h(\varphi_n) \neq t_0$ ).

Indeed, suppose that for every  $n \in \mathbb{N}$ , there exists  $m_n \in \mathbb{N}$  such that  $t_0 > h(\varphi_{m_n})$ . Since  $\psi(\varphi_{m_n}, h(\varphi_{m_n})) = 1$ ,  $\psi(\varphi_{m_n}, t_0) = 0$ . By passing to the limit, we obtain that  $\psi(\varphi, t_0) = \psi(\varphi, h(\varphi)) = 0$ , which is impossible, hence there exists  $n_0 \in \mathbb{N}$  such that  $t_0 < h(\varphi_n)$  for all  $n \geq n_0$ .

Applying the case 1-c), to  $\varphi = \varphi_n$ , we obtain  $\omega_{t_\alpha}(\varphi_n) \rightarrow \omega_{t_0}(\varphi_n) = g(\varphi_n)$  for every  $n \geq n_0$ . It follows that  $\omega_{t_0}(\varphi) = g(\varphi)$ , because  $g$  is continuous.

*Case 4,  $h(\varphi) > t_0 = 0$ :* Put

$$W = \{t \in E; h(\varphi) > t \text{ and } -h(\varphi) < t\}.$$

$W$  is an open neighborhood of  $t_0 = 0$ , hence

$$S = \{\alpha \in I; h(\varphi) > t_\alpha \text{ and } -h(\varphi) < t_\alpha\} \in \mathcal{U}.$$

If  $\alpha \in S$ ,  $\omega_{t_\alpha}(\varphi) = \omega_0(\varphi) = 1 + 1 - 1 = 1$ . We conclude that  $\omega_{t_\alpha}(\varphi) = 1 \rightarrow \omega_0(\varphi) = 1 = g(\varphi)$ .

The case  $h(\varphi) < t_0 = 0$  is treated by a similar argument. If  $\varphi \in B_1$ , it is clear that  $\omega_{t_\alpha}(\varphi) \rightarrow \omega_t(\varphi)$ . Thus  $V$  is a closed subset in  $(Y_1, \tau_p)$ . ■

7). We will show that the map  $\sigma_2 : t \in E \rightarrow (f_t) \in (C(K), \tau')$  is Borel.

Note that for every  $t \in E$ ,  $(f_t) = e - f_{-t}$ . Consider the map  $\Delta : E \rightarrow E$  defined by  $\Delta(t) = -t$ ,  $t \in E$ ,  $\Delta$  is a Borel function. Thus  $\sigma_2 = e - \sigma_1 \circ \Delta$ ,  $\sigma_2$  is a Borel function, because in the proof of 2) we showed that  $\sigma_1 : t \in E \rightarrow f_t$  is Borel.

Consider the map  $\phi_1 : t \in E \rightarrow (f_t, (f_t)) \in C(K) \times C(K)$ . It is obvious that if  $W \in \text{Bor}(C(K), \tau') \otimes \text{Bor}(C(K), \tau')$ , then  $\phi_1^{-1}(W)$  is a Borel subset of  $E$ .

Suppose now that  $\text{Bor}(C(K), \tau') \otimes \text{Bor}(C(K), \tau') = \text{Bor}(C(K) \times C(K), \tau' \times \tau')$ . Therefore the map  $\phi_1 : E \rightarrow (C(K) \times C(K), \tau' \times \tau')$ ,  $t \rightarrow (f_t, (f_t))$  is Borel.

Define the map  $\phi_2 : (C(K) \times C(K), \tau' \times \tau') \rightarrow (C(K), \tau')$ , by  $\phi_2(g, u) = g + u$ ,  $(g, u) \in C(K) \times C(K)$ . Observe that  $\phi_2$  is continuous, hence it is Borel. It follows that  $\phi_2 \circ \phi_1$  is Borel. We deduce that the map  $t \in E \rightarrow w_t = f_t + (f_t) \in (Y_1, \tau') = (Y_1, \tau_p)$  is Borel. But  $\text{Bor}(C(K), \tau_p) = \text{Bor}(C(K), \|\cdot\|)$  by Corollary 2, hence  $\text{Bor}(Y_1, \tau_p) = \text{Bor}(Y_1, \|\cdot\|)$ , which implies that the map  $t \in (E, \tau_0) \rightarrow \omega_t \in Y_1$  is strongly measurable, this is impossible, because in the proof of 6), we saw that  $(V, \tau_p)$  is uncountable discrete space, this means that the map  $t \in E \rightarrow \omega_t$  cannot be almost everywhere valued in a separable subspace of  $C(K)$ . ■

8). Suppose that the projection  $P : (C(K), \tau') \rightarrow (Y_1, \tau') = (Y, \tau_p)$  is Borel. Since the map  $\sigma_1 : t \in E \rightarrow f_t \in (C(K), \tau')$  is Borel, the map  $t \in E \rightarrow P \circ \sigma_1(t) = P f_t = w_t/2 \in (C(K), \tau')$  is Borel, but in the proof of 7) we saw that this is impossible. Thus  $P$  is not a Borel function. ■

9). Put  $H_1 = \sigma_1 : t \in E \rightarrow f_t \in (C(K), \tau')$  and  $H_2 = \sigma_2 : t \in E \rightarrow (f_t) \in (C(K), \tau')$ . We proved previously that  $H_1, H_2$  are Borel and in the end of the proof of 8) we proved that  $H_1 + H_2$  is not Borel. ■

**Definition 9.** Let  $(X, \tau)$  be a topological Hausdorff space.  $(X, \tau)$  is called a  $K$ -analytic set [4] if  $(X, \tau)$  is a continuous image of a set belonging to the family  $K_{\sigma\delta}$ .

By [20, th. 2.7.1], every  $K$ -analytic set is a Lindelöf space. On the other hand if  $L$  is separable and compact such that  $(C(L), \tau_p)$  is a Lindelöf space, then  $L$  is metrisable [1] (under  $MA + \neg CH$ ).

**Corollary 20.**  $(C(K), \tau')$  is not  $K$ -analytic.

*Proof.* Suppose that  $(C(K), \tau')$  is  $K$ -analytic. By [4]  $(C(K), \tau')$  is universally measurable, which is impossible by Theorem 5 2). ■

**Corollary 21.** *There exists two subspaces  $Z_1, Z_2$  of  $(C(K), \tau_p)$ ,  $\tau_p$ -closed such that  $Z_1 \cap Z_2 = \{0\}$ ,  $(Z_j, \tau')$  is universally measurable,  $j \in \{1, 2\}$  and  $(Z_1, \tau')$  is isomorphic to  $(Z_2, \tau')$ , but  $(Z_1 \oplus Z_2, \tau')$  is not universally measurable.*

*Proof.* Let  $Z_j = Y_j$ ,  $j \in \{1, 2\}$ . By Theorem 5, one has  $(Z_j, \tau') = (Z_j, \tau_p)$ ,  $Z_j$  is  $\tau_p$  closed in  $(C(K), \tau_p)$ ,  $j \in \{1, 2\}$  and  $(Z_1, \tau')$  is isomorphic to  $(Z_2, \tau')$ . On the other hand, Lemma 15 shows us that  $(C(K), \tau_p)$  is universally measurable, hence  $(Z_j, \tau') = (Z_j, \tau_p)$  is universally measurable (because a closed subset of an universally measurable set is universally measurable). Since  $(Z_1 \oplus Z_2, \tau') = (C(K), \tau')$ , by Theorem 5 2),  $(Z_1 \oplus Z_2, \tau')$  is not universally measurable. ■

**Remark 25.** *By Corollary 2,  $\text{Bor}(C(K), \tau_p) \otimes \text{Bor}(C(K), \tau_p)$  is equal to  $\text{Bor}(C(K), \|\cdot\|) \otimes \text{Bor}(C(K), \|\cdot\|)$ .*

On the other hand by [26, th. 3] (with the continuous hypothesis) one has

$$\text{Bor}(C(K), \|\cdot\|) \otimes \text{Bor}(C(K), \|\cdot\|) = \text{Bor}(C(K) \times C(K), \|\cdot\|).$$

Thus  $\text{Bor}(C(K) \times C(K), \tau_p \times \tau_p) \subset \text{Bor}(C(K) \times C(K), \|\cdot\|) = \text{Bor}(C(K), \|\cdot\|) \otimes \text{Bor}(C(K), \|\cdot\|) = \text{Bor}(C(K), \tau_p) \otimes \text{Bor}(C(K), \tau_p)$ . But  $\text{Bor}(C(K), \tau_p) \otimes \text{Bor}(C(K), \tau_p) \subset \text{Bor}(C(K) \times C(K), \tau_p \times \tau_p)$ , one deduces that

$$\text{Bor}(C(K), \tau_p) \otimes \text{Bor}(C(K), \tau_p) = \text{Bor}(C(K) \times C(K), \tau_p \times \tau_p).$$

**Acknowledgement.** I would like to thank Professors Bernard Maurey and Gilles Godefroy for the time they devoted to me during the preparation of this work.

#### REFERENCE

- [1] **Arkhangel'skiĭ A. V.** *Topological functions spaces*, Kluwer Acad. Publ. Dordrecht-Boston London, 1992.
- [2] **Bourbaki N.** *Éléments de mathématiques, topologie générale, chapitre 9, utilisation des nombres réels en topologie générale*, Hermann, Paris, 1959.
- [3] **Bourgain J., Fremlin D. H. and Talagrand M.** *Pointwise compact sets of Baire measurable functions*, Amer. J. Math., 1998, 100(4):845-886.
- [4] **Bressler D. W. and Sion M.** *The current theory of analytic sets*, Canad. J. Math., 1964, 16:207-230.
- [5] **Fremlin D. H.** *Measurable functions and almost continuous functions*, Manus Math, 1981, 33:387-405.
- [6] **Edgar G. A.** *Measurability in Banach spaces*, Indiana University J Math, 1977, 26(4):663-680.
- [7] **Ferrando J. C.** *Two new properties of the space  $C_p(X)$* , Top Appl, 2007, 154:1799-1803.

- [8] **Godefroy G.** *Compacts de Rosenthal*, Pac Jour of Math, 1980, 91:293-306.
- [9] **Godement R.** *Sur la théorie des représentations unitaires*, Ann Math, 1951, 53:68-124.
- [10] **Haydon R G., Jayne J E., Namioka I. and Rogers C A.** *Continuous functions on totally ordered spaces that are compact in their order topologies*, J Funct Anal, 2000, 178(1):23-63.
- [11] **Grothendieck A.** *Critères de compacité dans les espaces fonctionnels généraux*, Amer J. Math. 168-186, 1952.
- [12] **Hewitt E. and Ross K A.** *Abstract Harmonic Analysis I*, Springer-Verlag, Berlin-Heidelberg-New-York, 1963.
- [13] **Jayne J E., Namioka I. and Rogers C A.** *Continuous functions on product of compact Hausdorff spaces*, Mathematika, 1997:323-330.
- [14] **Kelley J L.** *General topology*, D. Van Nostrand Company I N C, 1955.
- [15] **Lindenstrauss J. and Stegall C.** *Exemples of separable spaces which do not contain  $\ell_1$  and whose duals are non separable*, Stud Math, 1975-1976, 54:81-105.
- [16] **Moran W.** *Measures and mappings on topological spaces*, Proceedings of the London Math Soc (3), 19:493-508.
- [17] **Moran W.** *The additivity of measures on completely regular spaces*, J London Math Soc, 1968, 43:633-639.
- [18] **Odell E. and Rosenthal H P.** *A double dual characterization of separable Banach spaces containing  $\ell^1$* , Israël J Math, 1975, 20, 375-384.
- [19] **Ostaszewski A J.** *A characterization of compact separable ordered spaces*, J Lond Math Soc, II Ser 7, 1974, 758-769.
- [20] **Rogers C A. and Jayne J E.** *K-analytic sets*, a chapter in *Analytic sets*, Acad Press, 1980:1-181.
- [21] **Schwartz L.** *Radon measures on arbitrary topological spaces and cylindrical measures*, Oxford University Press, (1973).
- [22] **Schwartz L.** *Certaines propriétés des mesures sur les espaces de Banach*, Séminaire Maurey-Schwartz, 1975-1976, Exposé 23.
- [23] **Schwartz L. and others.** *Radon measures*, Oxford U P Oxford, 1973.
- [24] **Stone M H.** *A general theory of spectra*, Proc Not Acad Sci U S A, 1940, 26:280-283.
- [25] **Sunyach Ch.** *Une caractérisation des espaces universellement mesurables*, C R Acad Sci, 1969, 268:864-866.
- [26] **Talagrand M.** *Est-ce que  $\ell^\infty$  est un espace mesurable*, Bull Sci Math, II<sup>e</sup> série, 1979, 103:255-258.
- [27] **Talagrand M.** *Solution d'un problème de A. Ionescu-Tulcea*, C R Acad Sci Paris, Sér A, 1976, 283:975-978.
- [28] **Varadarajan V S.** *Measures on topological spaces*, Mat. Sb. (N.S), 55(97), (Russian); English translation: Amer Math Soc Transl, 1965, (2) 48:161-228.

*Email address:* daher.mohammad@ymail.com, kh\_saadi@yahoo.fr