

# How large should whales be?

Aaron Clauset<sup>1,2,3,\*</sup>

<sup>1</sup>*Department of Computer Science, University of Colorado, Boulder CO, 80309 USA*

<sup>2</sup>*BioFrontiers Institute, University of Colorado, Boulder CO, 80303 USA*

<sup>3</sup>*Santa Fe Institute, Santa Fe NM, 87501 USA*

The observed body masses for all extant cetacean species, including the 175,000,000 g Blue Whale, are predicted, with no tunable parameters, by a macroevolutionary tradeoff between short-term selective advantages and long-term extinction risks from increased species body size, unfolding in a pelagic environment. Dramatic size differences between terrestrial and aquatic mammals are thus the result of increased convective heat loss in water, which induces a minimum viable size orders of magnitude larger than the 2 g limit induced by air for terrestrial mammals. The large minimum size would have presented a high macroevolutionary barrier to mammals becoming fully aquatic and explains the historical timing of mammals' invasion of aquatic habits.

Keywords: body mass distribution, cetaceans, macroevolution, cladogenesis

Cetaceans include the largest animals ever to live, including the Blue Whale (*Balaenoptera musculus*), which is nearly 30 times larger than an African elephant and twice as large as the largest sauropod. However, the evolutionary and ecological reasons for their enormous sizes or the possibility of still larger animals remains unclear. A deeper understanding of the mechanisms shaping cetacean sizes would shed light on the interaction of macroecological [7] and macroevolutionary processes [44], on long-term trends in species mass [3, 4], e.g., Cope's rule, the empirically observed tendency for species masses to increase within a lineage over evolutionary time [2, 43], and may inform conservation efforts.

Many major animal clades, including mammals, birds, fish and insects, seem to exhibit a canonical pattern in the distribution of species masses [1, 10, 24, 43]. For example, the most common size of a terrestrial mammal is roughly 40 g (common Pacific Rat, *Rattus exulans*). Both larger and smaller species are much less common, but asymmetrically so: the largest species, like the extinct Imperial Mammoth (*Mammuthus imperator*,  $10^7$  g), are orders of magnitude larger, while the smallest, like Remy's Pygmy Shrew (*Suncus remyi*, 2 g), are only a little smaller (Fig. 1).

Both the precise shape and the origins of its ubiquity have long been a topic of ecological interest. Recently, this characteristic pattern was shown to be a long-term consequence when a minimum viable body size, e.g., from physiological or thermoregulatory limits, constrains a tradeoff between short-term selective advantages [7] and long-term extinction risks from increased species size [10, 11] (Fig. 2a). Other studies have suggested that the pattern is caused by competition about a taxon-specific energetically optimal body size [8, 26, 39]; however, evidence of Cope's rule—descendant species tend to be larger than their ancestors—and the fact that most species are not close to their group's predicted optimal

size (among other reasons [24]) suggest that this theory may be flawed.

Here, we consider the body masses of cetaceans, which also exhibit the canonical right-skewed pattern (Fig. 1): the median size (356 kg, *Tursiops truncatus*) is close to the smallest (37.5 kg, *Pontoporia blainvillei*) but far from the largest (175,000 kg). This fact is notable because Cetacea is a modest-sized clade (77 extant species) within Mammalia and claims about canonical size distributions have primarily been made about much more speciose and higher-level taxonomic groupings. Furthermore, Cetacea is the largest and most diverse aquatic mammal clade (Sirenia has only five extant species). Cetacean sizes thus provide a unique test of the macroevolutionary tradeoff hypothesis. They allow us to test whether the same short-term-long-term tradeoff that explains the sizes of terrestrial mammals also explain the sizes of whales, and to investigate the impact of an aquatic environment on mammalian body size evolution.




FIG. 1: Terrestrial (includes semi-aquatic) [40] and fully aquatic mammal species mass distributions. Both show the canonical asymmetric pattern: the median size is flanked by a short left-tail down to a minimum viable size and a long right-tail out to a few extremely large species.

\*Electronic address: aaron.clauset@colorado.edu

We show that the tradeoff mechanism is universal for mammals. Using a cladogenetic diffusion model estimated from fossil and extant terrestrial mammal data, but with a minimum viable size appropriate for aquatic endotherms, we produce a statistically accurate, zero-parameter prediction of extant cetacean masses. This strong agreement implies that a single macroevolutionary tradeoff explains both aquatic and terrestrial mammal sizes. We close with a brief discussion of the ecological and evolutionary theoretical implications.

## I. NEUTRAL MODEL FOR CETACEAN SIZES

Following Clauset and Erwin [10], we model the tradeoff hypothesis as constrained cladogenetic diffusion.

First, a species of mass  $M$  produces descendant species with masses  $\lambda M$  (Fig. 2b), where  $\lambda$  is a random variable representing short-term selective effects on size [29, 43]. With each  $\lambda$  drawn independently, short-term selections on body sizes are uncorrelated across the clade and the clade's size distribution evolves according to a diffusion process. The trajectory of any particular lineage follows a kind of random walk. When the average size change between ancestors and descendants within a lineage is biased toward larger sizes (Cope's rule), we have  $\langle \ln \lambda \rangle > 0$  [2]. (Size variation between speciation events need not be modeled separately because its impact may be absorbed into  $\lambda$ .)

Second, species may not take any size and thus the diffusion process is constrained. On the upper end, the probability of species extinction rises gently with increasing size [25], e.g., due to larger energetic requirements, smaller species abundance and longer generational times. The net effect is a soft upper limit. Given a particular extinction risk curve, the number and size of very large species is determined by the total number of species in the clade, which sets the rate at which smaller-sized lineages migrate into the larger and more risky size ranges [48].

On the lower end, endothermy imposes a minimum viable mass—a hard lower limit—that prohibits evolution toward ever smaller sizes. For terrestrial mammals and birds, this thermoregulatory minimum size is known to occur at  $M_{\min} = 2 \text{ g}$  [33, 50], below which a species' convective heat loss in air is too high to maintain its internal temperature.

To extract precise predictions for the distribution of species sizes, we formalize this model mathematically. Following Clauset and Redner [11],  $c(x, t)$  denotes the density (fraction) of species having mass  $x = \ln M$  at time  $t$ . Under mild assumptions, the value  $c(x, t)$  obeys the convection-diffusion-reaction equation in the continuum limit:

$$\frac{\partial c}{\partial t} + v \frac{\partial c}{\partial x} = D \frac{\partial^2 c}{\partial x^2} + (k - A - Bx)c, \quad (1)$$

where  $v = \langle \ln \lambda \rangle$  is the bias or average change in size from ancestor to descendent,  $D = \langle (\ln \lambda)^2 \rangle$  is the diffusion




FIG. 2: (A) The characteristic distribution of species body sizes, observed in most major animal groups. Macroevolutionary tradeoffs between short-term selective advantages and long-term extinction risks, constrained by a minimum viable size  $M_{\min}$ , produce the distribution's long right-tail. (B) Schematic illustrating the cladogenetic diffusion model of species body-size evolution: a descendant species' mass is related to its ancestor's size  $M$  by a random multiplicative factor  $\lambda$ . Species become extinct with a probability that grows slowly with  $M$ .

coefficient or the variance in size change, and  $k - A$  is the size-independent (background) net speciation rate, which sets the absolute scale of the mass frequencies.

The aforementioned size constraints ensure convergence on a steady state distribution. To solve for its shape, we change variables  $\mu = v/D$ ,  $\alpha = (k - A)/D$ , and  $\beta = B/D$ , and require that the distribution go to zero at  $x = x_{\min}$ . It can then be shown that the steady-state distribution of sizes  $x$  is

$$c(x) \propto e^{\mu x/2} \text{Ai} \left[ \beta^{1/3}(x - x_{\min}) + z_0 \right], \quad (2)$$

where  $\text{Ai}[\cdot]$  is the Airy function and  $z_0 = -2.3381 \dots$  is the location of its first zero [11, 12].

In this way, the predicted shape of the species size distribution is fully determined by three model parameters:  $\mu$ , the normalized strength of Cope's rule,  $\beta$ , the normalized size-dependence of extinction risk, and  $x_{\min}$ , the logarithm of the minimum viable body size.

Estimates for  $\mu$  and  $\beta$  for terrestrial mammals have previously been derived from fossil and extant data. The resulting size distribution accurately reproduces both the extant sizes of terrestrial mammals [10] and their expansion during the late Cretaceous and early Paleogene [11, 51]. Removing either the size-dependence of extinction risk or the minimum viable size produces unrealistic predictions [10].

However, pelagic environments impose distinct physiological, ecological and evolutionary challenges for endothermic mammals, which are not reflected in the terrestrial model. One critical difference is the greater convective heat loss in water, which raises the minimum size of a competent aquatic endotherm. Thermoregulatory calculations and empirical data both estimate roughly  $M_{\min} = 7 \text{ kg}$  for extant cetaceans [15], about 3500 times larger than the minimum size in air.

## II. TESTS OF THE MODEL PREDICTIONS

To test the tradeoff hypothesis for cetaceans, we use the terrestrial mammal model, but we shift  $M_{\min}$  to its pelagic value. That is, we use a model that successfully explains the sizes of terrestrial mammal species but we change the environmental constraint imposed by  $M_{\min}$  so that the tradeoff unfolds in a pelagic environment. This produces an *ex ante* prediction  $\Pr(M)$  for extant cetacean species sizes. Notably, the prediction has no tunable parameters by which to improve its agreement with observed sizes. This property makes its accuracy a strong test of the tradeoff's universality.

Previous analyses of terrestrial mammal data yielded  $\hat{\mu} \approx 0.2$ , a slight tendency toward larger sizes within a lineage (Cope's rule), and  $\hat{\beta} \approx 0.08$ , a weak tendency for extinction to increase with body size [11, 12].

To test the model's accuracy, we constructed a novel body size data set for all 77 extant cetacean species, from 183 empirical size estimates [5, 6, 9, 13, 14, 16–23, 27, 30–32, 34–38, 40–42, 45–47, 49]. Only plausibly independent, scientifically derived estimates were included. Mass ranges were converted to point estimates by taking their midpoint, unless a mean value was also provided. Subsequently, the mean value of all point estimates for a given species was used; this yielded an average of 2.4 measurements per species. Tables I and II give the mass estimates, primary source(s) and data curation comments.

Figure 3 compares the predicted and empirical distributions. We determined the prediction's statistical plausibility relative to the empirical sizes via a standard two-tailed Kolmogorov-Smirnov test, evaluated numerically. The resultant value  $p_{ks} = 0.16 \pm 0.01$  exceeds all conventional thresholds for rejecting the null hypothesis, indicating that extant cetacean sizes are statistically indistinguishable from the predicted distribution. To simulate statistical uncertainty in  $\mu$  and  $\beta$ , we estimated  $p_{ks}$  via Monte Carlo by adding a small amount of Normally distributed noise to the terrestrial parameter val-




FIG. 3: (A) *Ex ante* predicted cetacean sizes, from a cladogenetic model fitted to terrestrial mammals but with a pelagic  $M_{\min}$  (see text), and empirical sizes of 77 extant cetacean species, as complementary cumulative distributions and as (B) smoothed probability densities.

ues. This yields a slightly lower but still non-significant  $p_{ks} = 0.07 \pm 0.03$ , which does not alter our conclusion. Thus, the macroevolutionary tradeoff fully explains the observed sizes of cetacean species as a group.

As an additional test, we consider whether the size of the largest cetacean species would be considered a statistical outlier. The probability of observing at least one species with size at least as large as the Blue Whale at  $M_* = 1.75 \times 10^8 \text{ g}$  was computed as  $p(M_*) = 1 - F(M_*)^n$  where  $F(M_*) = \int_{M_{\min}}^{M_*} \Pr(M) dM$  is the portion of the predicted distribution below  $M_*$  and  $n$  is the number of iid observations (extant species) drawn from  $\Pr(M)$ . Taking fixed parameters yields  $p = 0.91$ ; simulating statistical uncertainty via Monte Carlo (as above) yields  $p = 0.88 \pm 0.03$ , which is consistent with the fixed-parameter result.

Thus, the enormous size of the Blue Whale is not a statistical outlier relative to the predicted distribution, implying that the tradeoff mechanism alone is sufficient explanation of the Blue Whale's size. We note, however, that a species somewhat larger than the Blue Whale would also not be statistically unlikely, although no such species is known to have existed.

Finally, the value of  $\beta$  used here was estimated by fitting the model to extant terrestrial mammal data. Estimating  $\beta$  directly from the extant cetacean species size distribution only improves the fit of the model and thus cannot change our conclusions. Doing so yields  $\hat{\beta}_{\text{cete}} \approx 0.097$ , which is close to the terrestrial mammals value and supports our universality claim for the underlying processes shaping mammal evolution.

### III. DISCUSSION

The accuracy of the model's *ex ante* prediction confirms that terrestrial and aquatic mammal sizes are shaped by a single macroevolutionary tradeoff between short-term advantages—better resource fluctuation tolerance, thermoregulation and predator avoidance [7]—and long-term extinction risks—larger energetic requirements, smaller species abundance and longer generational times—of increased size, constrained by endothermic requirements at the lower end. Thus, cetaceans exhibit large body sizes because greater convective heat loss in water raises the minimum viable size, which shifts the canonical distribution upward and pushes its tail into size ranges inaccessible to terrestrial mammals.

This large minimum size was a significant barrier for early terrestrial mammals, preventing their invasion of pelagic niches prior to roughly 60 Ma. Before this time, there were few or no mammal species with sufficiently large masses [28], and therefore no mammals capable of living as competent aquatic endotherms. Only after the terrestrial mammal size distribution expanded in the late Cretaceous and early Paleogene [11, 51] was such a transition possible, and mammals successfully invaded the oceans almost immediately after these conditions were met.

The general pattern of extant cetacean body sizes is compactly explained by only the normalized strength of Cope's rule  $\mu$  and the normalized size-dependence of extinction risks  $\beta$ , which take universal values for all mammals, terrestrial and aquatic. Given the first archaeocete's size, species counts over geological time and the model diffusion rate, the model would predict when a species of a given size should first have appeared.

The success of the constrained cladogenetic diffusion model and the tradeoff hypothesis in predicting, without tunable parameters, the sizes of extant cetaceans further

suggests that the energetically optimal body size explanation of the species size distribution is flawed. This, in turn, suggests that the empirically observed pattern of size changes among insular species [26, 39], and their hypothesized explanation [8], should be revisited.

The short-term-long-term tradeoff dynamic described here is a neutral mechanism for species size evolution within a clade, as it omits explicit mechanisms for ecologically important processes like species interaction, geography, climate, etc. Deviations from the predictions, like the slight over-abundance of cetaceans near 800 kg and under-abundance near 100,000 kg (Fig. 3a), may indicate previously unrecognized non-neutral evolutionary or ecological processes. Similarly, to the extent that they can be measured in empirical data, changes in model parameter values over time may indicate broad-scale, non-stationary processes like climate change, e.g., in the value on  $x_{\min}$  or the magnitude of  $\mu$ , or clade-level ecological competition, as between mammals and dinosaurs prior to the K-Pg event.

Finally, our assumptions of size-related macroevolutionary tradeoffs are entirely general, but it remains unknown whether they hold for other major clades, including aquatic tetrapod groups like ichthyosaurs, plesiosaurs and turtles, or dinosaurs, fish and foraminifera, or whether it holds for other small taxonomic groups. A broad examination of minimum viable sizes and size-dependent extinction risks across groups and across geologic time may better elucidate the role of species size in major evolutionary transitions and ecological theory.

### Acknowledgments

The author thanks Doug Erwin and Sid Redner for helpful conversations. This work was supported in part by the Santa Fe Institute.

- 
- [1] Allen, C. R., A. S. Garmestani, T. D. Havlicek, P. A. Marquet, G. D. Peterson, C. Restrepo, C. A. Stow, and B. E. Weeks (2006). Patterns in body mass distributions: sifting among alternative hypotheses. *Ecology Letters* 9, 630–643.
  - [2] Alroy, J. (1998). Cope's rule and the dynamics of body mass evolution in North American fossil mammals. *Science* 280, 731–734.
  - [3] Alroy, J. (2000a). New methods for quantifying macroevolutionary patterns and processes. *Paleobiology* 26, 707–733.
  - [4] Alroy, J. (2000b). Understanding the dynamics of trends within evolving lineages. *Paleobiology* 26, 319–329.
  - [5] Best, R. C. and V. M. F. da Silva (1993). *Inia geoffrensis*. *Mammalian Species* 426, 1–8.
  - [6] Borsa, P. (2006). Marine mammal strandings in the new caledonia region, southwest pacific. *Comptes Rendus Biologies* 329, 277–288.
  - [7] Brown, J. H. (1995). *Macroecology*. Chicago: University of Chicago Press.
  - [8] Brown, J. H., P. A. Marquet, and M. L. Taper (1996). Darwinian fitness and reproductive power: reply to kozlowski. *American Naturalist* 147, 1092–1097.
  - [9] Clapham, P. J. and J. G. Mead (1999). Megaptera novaeangliae. *Mammalian Species* 604, 1–9.
  - [10] Clauset, A. and D. H. Erwin (2008). The evolution and distribution of species body size. *Science* 321, 399–401.
  - [11] Clauset, A. and S. Redner (2009). Evolutionary model of species body mass diversification. *Physical Review Letters* 102, 038103.
  - [12] Clauset, A., D. J. Schwab, and S. Redner (2009). How many species have mass  $M$ ? *American Naturalist* 173, 256–263.
  - [13] Cranford, T. W. (1999). The sperm whale's nose: Sexual selection on a grand scale? *Marine Mammal Science* 15, 1133–1157.
  - [14] Culik, B. M. (2004). *Review of Small Cetaceans*. Bonn, Germany: Conservation of Migratory Species of Wild

- Animals Secretariat. Marine Mammal Action Plan / Regional Seas Report and Studies no. 177.
- [15] Downhower, J. F. and L. S. Blumer (1988). Calculating just how small a whale can be. *Nature* 335, 675.
- [16] Jefferson, T. A. and N. B. Barros (1997). Peponocephala electra. *Mammalian Species* 553, 1–6.
- [17] Jefferson, T. A. and B. E. Curry (2003). Stenella clymene. *Mammalian Species* 726, 1–5.
- [18] Jefferson, T. A. and S. K. Hung (2004). Neophocaena phocaenoides. *Mammalian Species* 746, 1–12.
- [19] Jefferson, T. A. and L. Karczmarski (2001). Sousa chinensis. *Mammalian Species* 655, 1–9.
- [20] Jefferson, T. A. and S. Leatherwood (1994a). Lagenodelphis hosei. *Mammalian Species* 470, 1–5.
- [21] Jefferson, T. A. and S. Leatherwood (1994b). Right whale dolphins *Lissodelphis borealis* (peale, 1848) and *Lissodelphis peronii* (lacépède, 1804). *Handbook of Marine Mammals* 5, 335–362.
- [22] Jefferson, T. A., S. Leatherwood, and M. A. Webber (1993). *FAO species identification guide. Marine mammals of the world*. Food and Agriculture Organization of the United Nations.
- [23] Jefferson, T. A. and M. W. Newcomer (1993). *Lissodelphis borealis*. *Mammalian Species* 425, 1–6.
- [24] Kozłowski, J. and A. T. Gawelczyk (2002). Why are species' body size distributions usually skewed to the right? *Functional Ecology* 16, 419–432.
- [25] Liow, L. H., M. Fortelius, E. Bingham, K. Lintulaakso, H. Mannila, L. Flynn, and N. C. Stenseth (2008). Higher origination and extinction rates in larger mammals. *Proc. Natl. Acad. Sci. (USA)* 105, 6097–6102.
- [26] Lomolino, M. (1985). Body size of mammals on islands: the island rule re-examined. *American Naturalist* 125, 310–316.
- [27] Long, C. A. (1968). An analysis of patterns of variation in some representative Mammalia. Part I. a review of estimates of variability in selected measurements. *Transactions of the Kansas Academy of Science (1903–)* 71, 201–227.
- [28] Luo, Z.-X. (2007). Transformation and diversification in early mammal evolution. *Nature* 450, 1011–1019.
- [29] McShea, D. W. (1994). Mechanisms of large-scale evolutionary trends. *Evolution* 48, 1747–1763.
- [30] Mead, J. G., W. A. Walker, W. J. Houck, and Smithsonian Institution (1982). *Biological Observations on Mesoplodon carlhubbsi (Cetacea, Ziphiidae)*. Smithsonian contributions to zoology. Smithsonian Institution Press.
- [31] Nagorsen, D. (1985). Kogia simus. *Mammalian Species* 239, 1–6.
- [32] Newcomer, M. W., T. A. Jefferson, and R. L. Brownell, Jr. (1996). *Lissodelphis peronii*. *Mammalian Species* 531, 1–5.
- [33] Pearson, O. P. (1948). Metabolism of small mammals, with remarks on the lower limit of mammalian size. *Science* 108, 44.
- [34] Perrin, W. F. (1998). *Stenella longirostris*. *Mammalian Species* 599, 1–7.
- [35] Perrin, W. F. (2001). *Stenella attenuata*. *Mammalian Species* 683, 1–8.
- [36] Perrin, W. F. (2002). *Stenella frontalis*. *Mammalian Species* 702, 1–6.
- [37] Perrin, W. F., G. E. Zubitsova, and A. A. Kuz'min (2004). *Partial Catalog of Cetacean Osteological Specimens in Russian Museums*. NOAA Technical Memorandum NMFS. NOM-TM-NMFS-SWFSC-364.
- [38] Reeves, R. R. and S. Tracey (1980). Monodon monoceros. *Mammalian Species* 127, 1–7.
- [39] Sebens, K. P. (1987). The ecology of indeterminate growth in animals. *Annual Review of Ecology and Systematics* 18, 371–407.
- [40] Smith, F. A., S. K. Lyons, S. K. M. Ernest, K. E. Jones, D. M. Kaufman, T. Dayan, P. A. Marquet, J. H. Brown, and J. P. Haskell (2003). Body mass of Late Quaternary mammals. *Ecology* 84, 3403. MOM Version 3.6.1.
- [41] Stacey, P. J. and P. W. Arnold (1999). Orcaella brevirostris. *Mammalian Species* 616, 1–8.
- [42] Stacey, P. J., S. Leatherwood, and R. W. Baird (1994). *Pseudorca crassidens*. *Mammalian Species* 456, 1–6.
- [43] Stanley, S. M. (1973). An explanation for Cope's Rule. *Evolution* 27, 1–26.
- [44] Stanley, S. M. (1975). A theory of evolution above the species level. *Proc. Natl. Acad. Sci. (USA)* 72, 646–650.
- [45] Stewart, B. E. and R. E. A. Stewart (1989). *Delphinapterus leucas*. *Mammalian Species* 336, 1–8.
- [46] Uhen, M. D., R. E. Fordyce, and L. G. Barnes (1998a). Mysticeti. In C. M. Janis, G. F. Gunnell, and M. D. Uhen (Eds.), *Evolution of Tertiary Mammals of North American Volume 2: Small Mammals, Xenarthrans, and Marine Mammals*, pp. 607–628. Cambridge University Press.
- [47] Uhen, M. D., R. E. Fordyce, and L. G. Barnes (1998b). Odontoceti. In C. M. Janis, G. F. Gunnell, and M. D. Uhen (Eds.), *Evolution of Tertiary Mammals of North American Volume 2: Small Mammals, Xenarthrans, and Marine Mammals*, pp. 566–606. Cambridge University Press.
- [48] Van Valkenburgh, B., X. Wang, and J. Damuth (2004). Cope's rule, hypercarnivory, and extinction in North American canids. *Science* 306, 101–104.
- [49] Van Waerebeek, K., L. Barnett, A. Camara, A. Cham, M. Diallo, A. Djiba, A. O. Jallow, E. Ndiaye, A. O. S. O. Bilal, and I. L. Bamy (2004). Distribution, status, and biology of the atlantic humpback dolphin, *Sousa teuszii* (kenthal, 1892). *Aquatic Mammals* 30, 56–83.
- [50] West, G. B., W. H. Woodruff, and J. H. Brown (2002). Allometric scaling of metabolic rate from molecules and mitochondria to cells and mammals. *Proc. Natl. Acad. Sci. (USA)* 99, 2473–2478.
- [51] Wilson, G. P., A. R. Evans, I. J. Corfe, P. D. Smits, M. Fortelius, and J. Jernvall (2012). Adaptive radiation of multituberculate mammals before the extinction of dinosaurs. *Nature* 483, 457–460.

| group      | family          | species                     | mass (kg) | primary source (reference)  | curation comments  |
|------------|-----------------|-----------------------------|-----------|---|--|
| Mysticeti  | Balaenidae      | Balaena mysticetus          | 100000    | Smith etal (2003)   |  |
| Mysticeti  | Balaenidae      | Balaena mysticetus          | 87500     | Jefferson Leatherwood Webber (1993)   |  |
| Mysticeti  | Balaenidae      | Eubalaena australis         | 23000     | Smith etal (2003)   |  |
| Mysticeti  | Balaenidae      | Eubalaena australis         | 100000    | Jefferson Leatherwood Webber (1993)   |  |
| Mysticeti  | Balaenidae      | Eubalaena glacialis         | 23000     | Smith etal (2003)   |  |
| Mysticeti  | Balaenidae      | Eubalaena glacialis         | 90000     | Jefferson Leatherwood Webber (1993)   |  |
| Mysticeti  | Balaenopteridae | Balaenoptera acutorostrata  | 10000     | Smith etal (2003)   |  |
| Mysticeti  | Balaenopteridae | Balaenoptera acutorostrata  | 14000     | Jefferson Leatherwood Webber (1993), Perrin Zubitsova Kuzmin (2004)                                 |  |
| Mysticeti  | Balaenopteridae | Balaenoptera borealis       | 20000     | Smith etal (2003)   |  |
| Mysticeti  | Balaenopteridae | Balaenoptera borealis       | 30000     | Jefferson Leatherwood Webber (1993), Long 1968  |  |
| Mysticeti  | Balaenopteridae | Balaenoptera edeni          | 20000     | Smith etal (2003)   |  |
| Mysticeti  | Balaenopteridae | Balaenoptera edeni          | 22500     | Jefferson Leatherwood Webber (1993)   |  |
| Mysticeti  | Balaenopteridae | Balaenoptera musculus       | 190000    | Smith etal (2003)   |  |
| Mysticeti  | Balaenopteridae | Balaenoptera musculus       | 160000    | Jefferson Leatherwood Webber (1993), Morton ed 1997   |  |
| Mysticeti  | Balaenopteridae | Balaenoptera physalus       | 70000     | Smith etal (2003)   |  |
| Mysticeti  | Balaenopteridae | Balaenoptera physalus       | 75000     | Jefferson Leatherwood Webber (1993), Uhen Fordyce Barnes 1998 inJanisGunnellUhen                    |  |
| Mysticeti  | Balaenopteridae | Megaptera novaeangliae      | 30000     | Smith etal (2003)   |  |
| Mysticeti  | Balaenopteridae | Megaptera novaeangliae      | 35000     | Jefferson Leatherwood Webber (1993), Clapham Mead 1999, Uhen Fordyce Barnes 1998 inJanisGunnellUhen |  |
| Mysticeti  | Eschrichtiidae  | Eschrichtius robustus       | 28500     | Smith etal (2003)   |  |
| Mysticeti  | Eschrichtiidae  | Eschrichtius robustus       | 35000     | Jefferson Leatherwood Webber (1993), Uhen Fordyce Barnes 1998 inJanisGunnellUhen                    |  |
| Mysticeti  | Neobalaenidae   | Caperea marginata           | 3200      | Jefferson Leatherwood Webber (1993), Perrin Zubitsova Kuzmin (2004)                                 | estimate by Smith etal 2003 gives a mass 10x as large, so we omit Smith etal |
| Odontoceti | Delphinidae     | Cephalorhynchus commersonii | 72.4      | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Cephalorhynchus commersonii | 76        | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Cephalorhynchus commersonii | 86        | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Cephalorhynchus eutropia    | 45        | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Cephalorhynchus eutropia    | 63        | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Cephalorhynchus eutropia    | 60        | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Cephalorhynchus heavisidii  | 40        | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Cephalorhynchus heavisidii  | 65        | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Cephalorhynchus hectori     | 50        | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Cephalorhynchus hectori     | 57        | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Delphinus delphis           | 80        | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Delphinus delphis           | 135       | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Delphinus delphis           | 200       | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Feresa attenuata            | 170       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Feresa attenuata            | 225       | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Globicephala macrorhynchus  | 726       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Globicephala macrorhynchus  | 3600      | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Globicephala melas          | 800       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Globicephala melas          | 2000      | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Globicephala melas          | 1600      | Perrin Zubitsova Kuzmin (2004)  |  |
| Odontoceti | Delphinidae     | Grampus griseus             | 387.5     | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Grampus griseus             | 400       | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Lagenodelphis hosei         | 164       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Lagenodelphis hosei         | 210       | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Lagenodelphis hosei         | 210       | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Lagenodelphis hosei         | 209       | Jefferson Leatherwood 1994  |  |
| Odontoceti | Delphinidae     | Lagenorhynchus acutus       | 182       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Lagenorhynchus acutus       | 208.5     | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Lagenorhynchus acutus       | 205       | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Lagenorhynchus albirostris  | 180       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Lagenorhynchus albirostris  | 265       | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Lagenorhynchus australis    | 120       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Lagenorhynchus australis    | 115       | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Lagenorhynchus australis    | 115       | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Lagenorhynchus cruciger     | 110       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Lagenorhynchus obliquidens  | 120       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Lagenorhynchus obliquidens  | 180       | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Lagenorhynchus obliquidens  | 82.5      | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Lagenorhynchus obscurus     | 127.5     | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Lagenorhynchus obscurus     | 60        | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Lagenorhynchus obscurus     | 100       | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Lissodelphis borealis       | 113       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Lissodelphis borealis       | 115       | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Lissodelphis borealis       | 116       | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Lissodelphis borealis       | 113       | Jefferson Newcomer (1993)   |  |
| Odontoceti | Delphinidae     | Lissodelphis peronii        | 116       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Lissodelphis peronii        | 116       | Jefferson Leatherwood Webber (1993)   | Culik 2004 repeats this measurement, so we omit Culik                        |
| Odontoceti | Delphinidae     | Lissodelphis peronii        | 116       | Newcomer Jefferson Brownell 1996  |  |
| Odontoceti | Delphinidae     | Orcaella brevirostris       | 190       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Orcaella brevirostris       | 122.5     | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Orcaella brevirostris       | 123.5     | Stacey Arnold 1999  |  |
| Odontoceti | Delphinidae     | Orcinus orca                | 4300      | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Orcinus orca                | 8750      | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Orcinus orca                | 4685      | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Orcinus orca                | 7050      | Perrin Zubitsova Kuzmin (2004)  | reported mass mean of 2 specimens  |
| Odontoceti | Delphinidae     | Peponocephala electra       | 208       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Peponocephala electra       | 275       | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Peponocephala electra       | 228       | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Peponocephala electra       | 208       | Jefferson Barros 1997   |  |
| Odontoceti | Delphinidae     | Pseudorca crassidens        | 1360      | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Pseudorca crassidens        | 2000      | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Pseudorca crassidens        | 1360      | Stacey Leatherwood 1994   |  |
| Odontoceti | Delphinidae     | Sotalia fluviatilis         | 44        | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Sotalia fluviatilis         | 40        | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Sousa chinensis             | 265       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Sousa chinensis             | 284       | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Sousa chinensis             | 215       | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Sousa chinensis             | 265       | Jefferson Karczmarski (2001)  |  |
| Odontoceti | Delphinidae     | Sousa teuszii               | 100       | Smith etal (2003)   |  |
| Odontoceti | Delphinidae     | Sousa teuszii               | 284       | Jefferson Leatherwood Webber (1993)   |  |
| Odontoceti | Delphinidae     | Sousa teuszii               | 215       | Culik (2004)  |  |
| Odontoceti | Delphinidae     | Sousa teuszii               | 166       | Waerebeek etal (2004)   |  |

TABLE I: Cetacean size estimates (part 1).

| group      | family        | species                           | mass (kg) | primary source (reference)   | curation comments  |
|------------|---------------|-----------------------------------|-----------|--|--|
| Odontoceti | Delphinidae   | <i>Stenella attenuata</i>         | 120       | Jefferson Leatherwood Webber (1993)  | estimate from Smith etal 2003 is less than 1/2 of the values reported elsewhere, so we omit Smith etal |
| Odontoceti | Delphinidae   | <i>Stenella attenuata</i>         | 119       | Culik (2004)   |  |
| Odontoceti | Delphinidae   | <i>Stenella attenuata</i>         | 119       | Perrin (2001)  |  |
| Odontoceti | Delphinidae   | <i>Stenella clymene</i>           | 68        | Smith etal (2003)  |  |
| Odontoceti | Delphinidae   | <i>Stenella clymene</i>           | 85        | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Delphinidae   | <i>Stenella clymene</i>           | 80        | Culik (2004)   |  |
| Odontoceti | Delphinidae   | <i>Stenella clymene</i>           | 80        | Jefferson Curry (2003)   |  |
| Odontoceti | Delphinidae   | <i>Stenella coeruleoalba</i>      | 135.9     | Smith etal (2003)  |  |
| Odontoceti | Delphinidae   | <i>Stenella coeruleoalba</i>      | 156       | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Delphinidae   | <i>Stenella coeruleoalba</i>      | 156       | Culik (2004)   |  |
| Odontoceti | Delphinidae   | <i>Stenella frontalis</i>         | 110       | Smith etal (2003)  |  |
| Odontoceti | Delphinidae   | <i>Stenella frontalis</i>         | 143       | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Delphinidae   | <i>Stenella frontalis</i>         | 143       | Culik (2004)   |  |
| Odontoceti | Delphinidae   | <i>Stenella frontalis</i>         | 140       | Perrin (2002)  |  |
| Odontoceti | Delphinidae   | <i>Stenella longirostris</i>      | 50.5      | Smith etal (2003)  |  |
| Odontoceti | Delphinidae   | <i>Stenella longirostris</i>      | 77        | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Delphinidae   | <i>Stenella longirostris</i>      | 50.5      | Perrin 1998  | Culik 2004 repeats measurement of Perrin 1998, so we omit Culik  |
| Odontoceti | Delphinidae   | <i>Steno bredanensis</i>          | 130       | Smith etal (2003)  |  |
| Odontoceti | Delphinidae   | <i>Steno bredanensis</i>          | 150       | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Delphinidae   | <i>Steno bredanensis</i>          | 155       | Culik (2004)   |  |
| Odontoceti | Delphinidae   | <i>Tursiops truncatus</i>         | 175       | Smith etal (2003)  |  |
| Odontoceti | Delphinidae   | <i>Tursiops truncatus</i>         | 650       | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Delphinidae   | <i>Tursiops truncatus</i>         | 242       | Culik (2004)   |  |
| Odontoceti | Monodontidae  | <i>Delphinapterus leucas</i>      | 1360      | Smith etal (2003)  |  |
| Odontoceti | Monodontidae  | <i>Delphinapterus leucas</i>      | 1500      | Steward Steward (1989), Uhen Fordyce Barnes (1998) inJanisGunnellUhen                            |  |
| Odontoceti | Monodontidae  | <i>Delphinapterus leucas</i>      | 1500      | Culik (2004)   |  |
| Odontoceti | Monodontidae  | <i>Delphinapterus leucas</i>      | 1600      | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Monodontidae  | <i>Monodon monoceros</i>          | 900       | Smith etal (2003)  |  |
| Odontoceti | Monodontidae  | <i>Monodon monoceros</i>          | 1600      | Jefferson Leatherwood Webber (1993), Reeves Tracey (1980)  |  |
| Odontoceti | Monodontidae  | <i>Monodon monoceros</i>          | 1300      | Culik (2004)   |  |
| Odontoceti | Phocoenidae   | <i>Australophocaena dioptrica</i> | 65        | Smith etal (2003)  |  |
| Odontoceti | Phocoenidae   | <i>Neophocaena phocaenoides</i>   | 32.5      | Smith etal (2003)  |  |
| Odontoceti | Phocoenidae   | <i>Neophocaena phocaenoides</i>   | 85        | Culik (2004)   |  |
| Odontoceti | Phocoenidae   | <i>Neophocaena phocaenoides</i>   | 71.8      | Jefferson Hung (2004)  |  |
| Odontoceti | Phocoenidae   | <i>Phocoena phocoena</i>          | 52.5      | Smith etal (2003)  |  |
| Odontoceti | Phocoenidae   | <i>Phocoena phocoena</i>          | 57.5      | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Phocoenidae   | <i>Phocoena phocoena</i>          | 55        | Culik (2004)   |  |
| Odontoceti | Phocoenidae   | <i>Phocoena sinus</i>             | 42.5      | Smith etal (2003)  |  |
| Odontoceti | Phocoenidae   | <i>Phocoena spinipinnis</i>       | 60        | Smith etal (2003)  |  |
| Odontoceti | Phocoenidae   | <i>Phocoena spinipinnis</i>       | 85        | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Phocoenidae   | <i>Phocoenoides dalli</i>         | 102.5     | Smith etal (2003)  |  |
| Odontoceti | Phocoenidae   | <i>Phocoenoides dalli</i>         | 200       | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Phocoenidae   | <i>Phocoenoides dalli</i>         | 200       | Culik (2004)   |  |
| Odontoceti | Phocoenidae   | <i>Phocoenoides dalli</i>         | 200       | Jefferson (1988)   |  |
| Odontoceti | Physeteridae  | <i>Kogia breviceps</i>            | 431.5     | Culik (2004), Borsa (2006), Uhen Fordyce Barnes (1998) inJanisGunnellUhen                        |  |
| Odontoceti | Physeteridae  | <i>Kogia breviceps</i>            | 450       | Culik (2004)   |  |
| Odontoceti | Physeteridae  | <i>Kogia breviceps</i>            | 400       | Borsa (2006), Jefferson Leatherwood Webber (1993), Uhen Fordyce Barnes (1998) inJanisGunnellUhen |  |
| Odontoceti | Physeteridae  | <i>Kogia simus</i>                | 183.1     | Smith etal (2003)  |  |
| Odontoceti | Physeteridae  | <i>Kogia simus</i>                | 270       | Nagorsen 1985  |  |
| Odontoceti | Physeteridae  | <i>Kogia simus</i>                | 270       | Culik (2004)   | mass quoted as 2702 kg, but this is too big by an order of magnitude. Assumed to be 270.2kg            |
| Odontoceti | Physeteridae  | <i>Kogia simus</i>                | 210       | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Physeteridae  | <i>Physeter catodon</i>           | 14025     | Smith etal (2003)  |  |
| Odontoceti | Physeteridae  | <i>Physeter catodon</i>           | 57000     | Jefferson Leatherwood Webber (1993), Cranford (1999)   |  |
| Odontoceti | Platanistidae | <i>Inia geoffrensis</i>           | 129.25    | Smith etal (2003)  |  |
| Odontoceti | Platanistidae | <i>Inia geoffrensis</i>           | 160       | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Platanistidae | <i>Inia geoffrensis</i>           | 167.5     | Culik (2004)   |  |
| Odontoceti | Platanistidae | <i>Inia geoffrensis</i>           | 129.25    | Best Silva (1993)  |  |
| Odontoceti | Platanistidae | <i>Lipotes vexillifer</i>         | 187.5     | Jefferson Leatherwood Webber (1993)  | mass estimate from Smith etal 2003 is less than 1/2 estimated range here, so we omit Smith etal        |
| Odontoceti | Platanistidae | <i>Platanista gangetica</i>       | 115       | Smith etal (2003)  |  |
| Odontoceti | Platanistidae | <i>Platanista gangetica</i>       | 108       | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Platanistidae | <i>Platanista minor</i>           | 83.9146   | Smith etal (2003)  |  |
| Odontoceti | Platanistidae | <i>Pontoporia blainvillei</i>     | 40.5      | Smith etal (2003)  |  |
| Odontoceti | Platanistidae | <i>Pontoporia blainvillei</i>     | 34        | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Ziphiidae     | <i>Berardius arnuxii</i>          | 7000      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Berardius bairdii</i>          | 11380     | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Berardius bairdii</i>          | 12000     | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Ziphiidae     | <i>Hyperoodon ampullatus</i>      | 5800      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Hyperoodon planifrons</i>      | 3000      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Indopacetus pacificus</i>      | 2200      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon bidens</i>          | 3400      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon bowdoini</i>        | 2600      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon carlhubbsi</i>      | 1400      | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon carlhubbsi</i>      | 3400      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon carlhubbsi</i>      | 500       | Mean Walker Houck 1982   |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon densirostris</i>    | 2300      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon densirostris</i>    | 1033      | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon europaeus</i>       | 5600      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon europaeus</i>       | 1200      | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon ginkgodens</i>      | 1500      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon grayi</i>           | 2900      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon grayi</i>           | 1100      | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon hectori</i>         | 1000      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon layardii</i>        | 1500      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon mirus</i>           | 2100      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon mirus</i>           | 1400      | Jefferson Leatherwood Webber (1993)  |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon mirus</i>           | 1400      | Culik (2004)   |  |
| Odontoceti | Ziphiidae     | <i>Mesoplodon stejnegeri</i>      | 4800      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Tasmacetus shepherdii</i>      | 2500      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Ziphius cavirostris</i>        | 4775      | Smith etal (2003)  |  |
| Odontoceti | Ziphiidae     | <i>Ziphius cavirostris</i>        | 3000      | Jefferson Leatherwood Webber (1993)  |  |

TABLE II: Cetacean size estimates (part 2).